1 **Three-dimensional genome architecture connects chromatin structure and**

2 **function in a major wheat pathogen**

4 Ivona Glavincheska[1] and Cecile Lorrain[1*]

5 [1] Plant Pathology Group, Institute of Integrative Biology, ETH Zürich, Zürich, Switzerland

6 *Corresponding author: cecile.lorrain@usys.ethz.ch

ABSTRACT

Three-dimensional (3D) genome organization plays a central role in gene regulation and genome stability,

yet its functional relevance in fungal pathogens remains largely uncharacterized. Here, we present a high-

resolution Hi-C analysis of the wheat pathogen *Zymoseptoria tritici*, revealing a hierarchical genome

organization. The genome adopts a Rabl-like conformation characterized by centromere clustering and

the spatial segregation of accessory chromosomes, which remain tethered to core pericentromeres,

supporting their mitotic stability. At finer resolution, we identify homotypic interactions among B

compartments and self-interacting domains demarcated by distinct histone modifications and insulator-

like sequence motifs at their boundaries. A subset of highly insulated domains enriched in

heterochromatin and transposable elements engages in strong inter-domain interactions. During wheat

infection, several self-interacting domains exhibit coordinated transcriptional activity, linking spatial

genome folding to dynamic gene regulation. These findings uncover multi-scale principles of nuclear

organization in a major fungal plant pathogen and establish a framework for investigating 3D genome

function in filamentous fungi.

INTRODUCTION

The spatial organization of eukaryotic genomes is fundamental to cellular function, underpinning chromosomal dynamics, genome integrity, and the regulation of gene expression[1–3]. The physical arrangement of the chromatin in the nuclear space reflects a highly ordered and dynamic hierarchical architecture. Advances in chromosome conformation capture integrated with high-throughput sequencing (Hi-C) have significantly transformed our understanding of genome organization. In addition, Hi-C, combined with polymer physics models, can be used to differentiate between chromatin states and describe chromosomal conformation[4]. The comparison of 3D genome organization across the tree of life highlights both shared features and distinct aspects of genome organization between organisms, raising the question of whether common principles govern genome architecture[5].

In metazoan models, genome conformation studies have revealed a highly hierarchical chromatin organization[5]. At the largest scale, chromosomes are organized into distinct conformations within the nucleus, generally categorized into two main types: chromosome territory-like and Rabl-like organizations[2]. Chromosome territory-like organization, primarily observed in mammals, is characterized by minimal inter-chromosomal interactions, with each chromosome occupying a discrete nuclear region during interphase[2]. In contrast, the Rabl-like configuration involves a polarized arrangement where centromeres cluster at one end of the nucleus and telomeres align at the opposite end[2]. The Rabl chromosome configuration is also observed during mitosis[6]. Beyond these large-scale arrangements, epigenetically distinct compartments interact preferentially to further structure genome architecture[3,5,7]. In metazoans, A compartments correspond to transcriptionally active, euchromatic regions enriched in gene activity, while B compartments are associated with transcriptionally repressed, heterochromatic regions. This bipartite compartmentalization highlights the functional partitioning of the genome into regions of activity and repression. At a finer resolution, chromatin is organized into topologically

associating domains (TADs), self-interacting regions within which DNA sequences interact more

frequently than with sequences outside the domain[8]. The function of TADs remains debated, but they

have been described as playing an important role in regulating gene expression, partly by modulating the

interactions between regulatory elements and their target genes[9–11]. TAD boundaries act as insulators

that mediate chromatin looping to ensure proper genomic interactions and are enriched in architectural

proteins, such as the cohesin complex and the DNA-binding protein CCCTC-binding factor (CTCF), as well

as regions containing highly transcribed housekeeping genes[9]. The disruption of genome folding can

disrupt interactions between genes and regulatory elements, as observed in the enhancer hijacking of

proto-oncogenes[10,11]. In contrast to metazoan systems, other organisms exhibit 3D genome organization

patterns that, although reminiscent of features such as loops and compartments observed in mammals,

arise through distinct mechanisms. For instance, plants' compartment-like structures and Polycomb-

associated domains have been identified in gene-rich regions[12–15]. However, these domains are not

necessarily conserved across plant species and are not strongly associated with gene co-regulation as seen

in animals[15]. Understanding the establishment and regulatory impact of 3D genome organization is

essential. Yet, despite significant progress in metazoan models, these processes remain poorly

understood in many other eukaryotes, particularly in fungi.

The fungal kingdom, comprising millions of species, has significant ecological and economic impacts[16]. The

deployment of Hi-C sequencing has reshaped our understanding of fungal genome organization[17]. While

Hi-C has been widely applied to generate chromosome-level assemblies, high-resolution datasets remain

scarce for most fungal species, limiting our insights into their nuclear architecture[17]. Fungal genomes

generally exhibit a Rabl-like organization characterized by centromere clustering, as demonstrated in

yeast and filamentous fungi[17,18]. However, chromatin compartmentalization in fungi appears to be less

defined compared to that in metazoans. For instance, *Rhizophagus irregularis* is one of the few fungal

species with described A/B compartments[19]. TAD-like structures have also been identified in various fungi,

resembling metazoan TADs but with a notably smaller size. Although Hi-C matrices have revealed the

presence of TAD-like structures in fungal genomes, boundary-associated proteins potentially responsible

for domain insulation have yet to be identified. In the budding yeast *Saccharomyces cerevisiae*, interacting

domains of 5-10 kb depend on nucleosome positioning that create rigid chromatin segments that can

function as domain boundaries independently of architectural proteins[20]. In *S. cerevisiae*, these

interaction domains or TAD-like structures are implicated in critical genome functions such as

recombination repression, genome evolution, and the temporal organization of replication origins during

mitosis[20]. In the filamentous pathogen *Verticilium dahliae,* TAD-like structures exhibited strong internal

chromatin interactions and limited interactions beyond their boundaries, indicating roles in regulating

gene expression and maintaining chromosomal organization[21]. In *Neurospora crassa*, TAD-like structures

consist of weakly insulated euchromatic regions flanked by constitutive heterochromatin borders[22,23]. The

diverse chromatin organization observed across fungal species underscores the need for high-resolution

Hi-C to elucidate the fundamental principles of nuclear architecture and its influence on genome function.

The fungal wheat pathogen *Zymoseptoria tritici* emerged alongside the domestication of wheat in the

Fertile Crescent[24]. The genome of *Z. tritici* is compartmentalized into thirteen core and eight accessory

chromosomes[25,26]. The core chromosomes, which are gene-rich and transposable element (TE)-poor,

contrast with the gene-poor and TE-rich accessory chromosomes, an ancestral feature shared across the

Zymoseptoria genus[26]. The accessory chromosomes exhibit a high degree of presence/absence

polymorphism in *Z. tritici* populations[27], and some chromosomes appear to be more prone to

destabilization than others, as indicated by targeted loss experiments[28]. The genomic partitioning of *Z.*

tritici is further reflected by distinct epigenetic signatures, particularly in the presence of specific histone

modifications in accessory chromosomes[29]. The epigenetic landscape of *Z. tritici* exhibits the distinction

of euchromatin, constitutive and facultative heterochromatin, cytosine methylation (5mC) regions, small

acrocentric centromeres, and telomeric repeats, underlining the complexity of *Z. tritici*'s chromatin

organization[26,30–35]. Pathogenicity-related genes in *Z. tritici* are enriched within TE-rich regions of the

genome, which likely facilitates rapid evolution through genome rearrangements and epigenetic

regulation[35,36]. These genes are predominantly repressed in the absence of the host and maintained in a

transcriptionally silent state by heterochromatin marks[36]. Upon host colonization, chromatin remodeling

events activate the expression of pathogenicity-related genes[36]. Despite these detailed insights into the

genomic and epigenomic features of *Z. tritici*, the spatial organization of its genome and how these regions

fold and interact in three-dimensional space remains unexplored. This study aims to bridge this gap by

integrating high-resolution Hi-C to explore the 3D architecture of *Z. tritici*'s genome. To examine the

association between the epigenetic landscape and chromatin folding, we investigated the interplay

among genomic, epigenomic, and transcriptomic features, shedding light on the structural and regulatory

mechanisms that govern the genome of this major wheat pathogen.

RESULTS

The Rabl-like genome of *Z. tritici* reveals distinct conformation patterns between core and accessory
chromosomes
108
To investigate the three-dimensional (3D) genome organization of *Z. tritici* IPO323, we performed Hi-C on

two independent replicates (Supplementary Fig. S1A). Quality control analysis confirmed reproducibility

between replicates, with a Pearson correlation of 0.87 and 77–81% valid read pairs per replicate

(Supplementary Fig. S1B). We combined the two datasets to generate a high-resolution contact map

containing ~19.2 million deduplicated valid read pairs (Supplementary Table S1), allowing us to resolve

chromatin interactions at a minimal bin size of 3 kb. At this resolution, 80% of bins in the core

chromosomes were covered by at least 1,000 contacts (Fig. 1A; Supplementary Table S2). At 10 kb

resolution, all chromosomes except chromosome 21 had at least 1,000 contacts in 90% of bins

(Supplementary Table S2). As expected, the Hi-C contact matrix was dominated by local interactions,

reflected by a strong signal along the diagonal of the genome-wide contact map (Fig. 1A). The strongest

interchromosomal contacts were enriched at centromeres and pericentromeric regions, consistent with

a Rabl-like conformation (observed-to-expected contact > 3.5; Fig. 1A, 1B; Supplementary Fig. S2-S3).

Strong interchromosomal interactions were predominantly enriched in regions marked by constitutive

heterochromatin (H3K9me3; Supplementary Fig. S3; Supplementary Table S3). Notably, 32.6% of the

strong interchromosomal interactions (observed-to-expected contact > 3.5) occurred between two

H3K9me3-enriched blocks, revealing a heterochromatin network spanning multiple chromosomes

(Supplementary Fig. S3C).

Accessory chromosomes preferentially anchored to the pericentromeric regions of core chromosomes,

with 78.6% of core-accessory interactions spanning the full length of accessory chromosomes but

restricted to the pericentromeric regions of core chromosomes (observed-to-expected contact > 3.5; Fig.

1B; Supplementary Fig. S3D). This spatial organization suggests that entire accessory chromosomes

primarily interact with core chromosomes at their centromeres and short arms while remaining

segregated from the long core chromosome arms (Fig. 1B). As accessory chromosome stability is variable,

we examined the potential relationship between chromosome loss and core-accessory pericentromeric

interaction strength[28]. We performed a log-transformed Poisson regression, which revealed a significant

negative association between interaction strength normalized for accessory chromosome size and the

loss frequency of accessory chromosomes in IPO323 after β-tubulin inhibition treatment[28] (p = 5.33e-05

for the intercept; p = 0.0025 for normalized interaction strength; Figure S4). This negative correlation

suggests a link between core-accessory interactions and the stability of accessory chromosome

transmission. Overall, we uncovered the Rabl-like structure of the *Z. tritici* IPO323 genome characterized

by interchromosomal interactions between (peri)centromeric regions and a heterochromatin-enriched

interaction network that possibly contributes to the spatial organization of the genome and maintains the

nuclear positioning of accessory chromosomes.

To determine whether chromatin conformation of core and accessory chromosomes is reflected by

distinct contact frequency decay values, we estimated the frequency decay exponent (α) as a function of

genomic separation and contact frequency for each chromosome. Here, the core chromosomes exhibited

an average decay exponent of α = -1.22 (standard deviation (SD) = 0.1), with three distinct scaling regimes

depending on genomic separation, indicating a hierarchical structure of the chromatin (Fig. 1C;

Supplementary Table S4). At short distances (< 70 kb), contact frequency decayed more gradually than

the global average with α = -0.71 (SD = 0.01). Interactions between 70 kb and 1 Mb followed a more

heterogeneous and steeper decline with α = -1.43 (SD = 0.09). At distances beyond 1 Mb, contact

frequency declined at a slower rate relative to intermediate distances (α = -1; SD = 0.09; Fig. 1C;

Supplementary Table S4). In contrast, accessory chromosomes displayed a more uniform decay of contact

frequency, with a global exponent of α = -0.67 (SD = 0.08; Fig. 1C; Supplementary Table S4). The contact

frequency decay was steepest at distances within 70 kb (α = -0.71; SD=0.18), and followed a shallower

decline at distances > 70 kb (α = -0.62; SD=0.01). This pattern suggests accessory chromosomes function

as unitary compartments and lack multiple self-interacting domains (Fig. 1C; Supplementary Table S4).

Together, these findings highlight a distinction in chromatin folding between core and accessory

chromosomes: core chromosomes display hierarchical folding and distance-dependent structural

transitions indicating the presence of self-interacting domains, whereas accessory chromosomes maintain

a more uniform and potentially less constrained conformation.

B chromatin compartments in *Z. tritici* engage in homotypic interactions at the intrachromosomal level

To further investigate the segregation of *Z. tritici* chromatin, we inferred compartmentalization strength based on principal component analyses (PCA) of intrachromosomal contact matrices at 10 kb and 3 kb resolution bins (Fig. 2; Supplementary Fig. S4). At 10 kb bins, eigenvector values segregated the chromosome (sub)telomeres, supporting inter-arm interactions. At 3 kb, however, a finer distinction between interaction patterns was present with negative eigenvector values associated with H3K9me3 enrichment regions (Supplementary Fig. S5). Compartmentalization strength at 3 kb bins was assessed by comparing interaction frequencies between regions with similar (positive-positive or negative-negative) and opposing (positive-negative) eigenvector values (Fig. 2A; Supplementary Fig. S5, S6). Regions with negative eigenvectors exhibited stronger interactions (interaction strength = 2) than positive eigenvector regions (interaction strength = 1; Fig. 2A; Supplementary Table S5). These negative eigenvector regions were enriched with heterochromatin-associated histone modifications, particularly H3K9me2 and H3K9me3, while depleted in euchromatin-associated H3K4me2, consistent with the characteristics of canonical B-compartments seen in metazoans (Fig. 2B, Supplementary Fig. S6). The B-compartment blocks had an average size of ~60 kb, accounting for 14.74 Mb (~36%) of the genome, and are characterized by low gene density and a high density of TEs (Fig. 2C-D, Supplementary Table S6). Notably, we observed a significant enrichment of retrotransposons, particularly LINEs and LTR elements from the *Copia* superfamily (Fig. 2C-E, Supplementary Table S7).

By contrast, positive eigenvector regions were associated with characteristic features of A-compartments. The A-compartment blocks had an average length of 100 kb, spanning a total of 24.35 Mb, of which 80.3% are covered by genes (9,996 genes), while TEs made up only 4.6% of the A-compartment blocks (Fig. 2C). These regions exhibited significantly higher gene density and lower TE density than B-compartment blocks (Fig. 2D). Gene ontology (GO) enrichment analysis revealed an overrepresentation of functions related to translation, ribosome function, RNA binding, and intracellular non-membrane-bounded organelles (FDR-

adjusted *p* < 0.01), suggesting a functional link to essential cellular processes such as protein biosynthesis,

RNA stability, modification, and transport (Supplementary Table S8-S9). Additionally, A-compartments

were enriched in non-autonomous miniature inverted-repeat elements (MITEs; Fig. 2E). Consistent with

these observations, genes and TEs within A-compartment blocks exhibited significantly higher *in vitro*

expression than those in B-compartment blocks (Fig. 2E), indicating that transcriptional regulation of both

genes and TEs is dependent on chromatin compartmentalization. The identification of intrachromosomal

A/B compartment blocks demonstrates that chromatin organization in *Z. tritici* goes beyond the global

distinction between core and accessory chromosomes.

The presence of TAD-like organization in *Z. tritici* reveals boundaries enriched in facultative

heterochromatin and specific DNA binding motifs

To investigate the presence of TAD-like structures in the *Z. tritici* IPO323 genome, we analyzed insulation

scores calculated from a 3 kb resolution Hi-C contact matrix. We identified 513 TADs and 533 TAD

boundaries in *Z. tritici* IPO323, with an average TAD length of approximately 70 kb (Supplementary Table

S10-12). The distribution of TADs varied between core and accessory chromosomes, with 92% of TADs

(473 TADs) located on core chromosomes and only 8% (40 TADs) on accessory chromosomes

(Supplementary Table S12). On average, TADs on core chromosomes contained 25 genes and 6 TEs,

whereas TADs on accessory chromosomes contained an average of 14 genes and 17 TEs per TAD

(Supplementary Table S12).

Since TAD boundaries act as insulators between TADs, we identified them by detecting local insulation

minima and further examined whether TAD boundaries exhibited specific epigenomic and genomic

features (Fig. 3A, 3B). We observed that TAD boundaries were depleted of the constitutive

heterochromatin histone modifications H3K9me2 and H3K9me3 but enriched for facultative

heterochromatin modifications H3K36me3, H4K20me1, and H4K20me3 (Fig. 3B, Supplementary Fig. S7).

In accessory chromosomes, TAD boundaries were also enriched for accessory chromosome-associated

modifications, H3K27me2 and H3K27me3 (Supplementary Fig. S7). In addition, TAD boundaries exhibited

a lower density of TEs, reduced RIP-like signatures, and a higher density of genes than TADs (Fig. 3B,

Supplementary Fig. S7). The active transcription mark H3K4me2, however, did not show a pattern of

enrichment at TAD boundaries (Supplementary Fig. S7). These findings indicate that TAD boundaries are

typically depleted in constitutive heterochromatin and enriched in facultative heterochromatin.

To investigate if TAD boundaries could be associated with housekeeping and core gene functions, such as

transcription factors and regulators, we first investigated the features of the 613 genes overlapping >50%

with TAD boundaries, including conservation across 19 *Z. tritici* strains[37], functional annotation

enrichment, and expression *in vitro* (Supplementary Table S11). We detected only 69 core genes with

orthologs in all 19 *Z. tritici* reference genomes (Fisher's test, p-value = 0.76; Supplementary Tables S13-

S15). Similarly, we did not find enrichment of specific Gene Ontology (GO) terms or functional domains

(Supplementary Tables S13-S15), nor did genes at boundaries exhibit differences in *in vitro* expression

levels compared to TAD-associated genes (Supplementary Fig. S8). Finally, we searched whether specific

DNA binding motifs were present at TAD boundaries and identified a significant enrichment of 35 known

and novel DNA binding motifs, grouped in 9 clusters (Supplementary Table S16). Interestingly, one of the

strongest matches showed similarity to the binding motif recognized by the mammalian KRAB zinc finger

protein, ZNF263. ZNF263 was recently shown to function as an insulating factor at TAD boundaries[38].

IPO323 has a single annotated KRAB zinc finger protein (ZtIPO323_082310), but this protein exhibits low

sequence similarity in its zinc finger domains with the human ZNF263 protein (Supplementary Table S17).

These results suggest that TAD boundaries in *Z. tritici* exhibit distinct epigenetic features and may

incorporate DNA binding motifs that are potentially recognized by insulator proteins.

Highly insulated TADs define chromosomal conformation, while local TAD organization regulates transcriptional dynamics during plant infection

The analysis of the intrachromosomal contact matrix revealed heterogeneity in contact patterns across TADs, with certain TADs exhibiting strong insulation to their proximal neighbors, indicative of a compartmentalized chromatin environment (Fig. 3C-D). These TADs were characterized by elevated insulation scores and flanked by lower insulation regions, suggesting stronger local boundaries. To further investigate, we defined these highly insulated TADs (HI-TADs) by comparing the maximum insulation score within each TAD to the scores of its boundaries (Fig. 3D). TADs with a sum of boundary-to-maximum insulation scores above the genome-wide median were classified as HI-TADs. We identified 105 HI-TADs across the genome, representing 20% of the total TADs (Supplementary Table S18). Analysis of strongest interactions (observed-to-expected intrachromosomal contacts > 1.75) revealed significantly higher strong intrachromosomal interactions between HI-TADs compared to R-TAD/R-TAD and HI-TAD/R-TAD interactions (Mann-Whitney $p < 0.05$; Supplementary Fig. S9). HI-TADs displayed a unique epigenetic profile, with strong enrichment for heterochromatin-associated marks H3K9me2 and H3K9me3 and depletion of all other histone modifications (Fig. 3C–D). Of the 105 HI-TADs identified, 64% (68) were associated with B-compartments. HI-TADs exhibited higher Repeat-Induced Point Mutation (RIP) indices, increased TE density, and lower gene density than regular TADs. In addition, we observed that R-TADs have significantly higher gene expression levels *in vitro* than HI-TADs (Supplementary Fig. S10). HI-TADs were also enriched in effector genes, implying that heterochromatin dynamics within these regions may be restructured to enable effector gene activation during host infection (Fisher's exact p-value < 0.01; odds ratio = 1.54; Supplementary Tables S19). Taken together, these results suggest that HI-TADs represent interspersed heterochromatic islands that may contain genes relevant for pathogenicity.

To investigate whether TADs function as co-regulatory domains in *Z. tritici*, we analyzed transcriptomic

data from axenic culture and wheat infection. Using a linear model, we investigated whether genes and

TEs within the same TAD were differentially coexpressed. We classified a TAD as co-regulated if at least

10 genes and TEs exhibited coordinated transcriptional changes (Supplementary Tables S20-23). The TAD

organization had a significant impact on gene expression dynamics between *in vitro* and *in planta*

conditions, with the model outperforming the null expectation in explaining transcriptional variation at

different infection stages and accounting for between 5%-9% of the adjusted variance in log fold-change

(likelihood ratio test p < 0.01; Supplementary Table S24). In addition, we performed further linear

regression analyses using TAD membership, gene GC content, and distance to the nearest TE as predictors

to determine whether other factors linked to gene expression can further explain transcriptional variation.

Adding both gene GC content and distance to nearest TE to the linear regression model explained an

additional 3-6% of the variance (Supplementary Table S24). However, in the absence of TAD membership

as a predictor variable gene GC content and TE distance only explained <0.1% of the transcriptional

variance. We identified 46 co-regulated TADs (8.9% of all TADs) with significant transcriptional

coordination during at least one stage of infection (Supplementary Fig. S11; Supplementary Table S25).

Interestingly, different TADs showed stage-specific co-expression patterns, with 23 co-regulated TADs

identified at only a single infection stage, reflecting localized chromatin dynamics between biotrophic and

necrotrophic phases of infection (Supplementary Fig. S11). We identified 19 co-regulated TADs in the

accessory chromosomes and six localized to the arm of chromosome 7, which display accessory

chromosome features, including H3K27me3 enrichment and sparse gene content (Supplementary Fig.

S12). Another third of co-regulated TADs are classified as HI-TADs (16 TADs). Collectively, these results

highlight that specific TADs act as functional modules of transcriptional regulation in *Z. tritici*, with regions

associated with heterochromatin, low gene density, and accessory features showing a greater

dependency on domain organization for coordinated gene expression.

DISCUSSION

Our study provides the first comprehensive characterization of the hierarchical 3D chromatin organization in the major fungal wheat pathogen *Z. tritici*. The genome of *Z. tritici* adopts a Rabl-like nuclear conformation, consistent with other fungal genomes[19,21,23,39,40] but with distinct folding patterns between core and accessory chromosomes. Accessory chromosomes are anchored to the pericentromeric regions of core chromosomes, with interaction strength correlating with stability, indicating that pericentromeric association may reinforce their nuclear positioning and stability. We detected hierarchical folding within core chromosomes, while accessory chromosomes displayed a more uniform organization. Beyond this global architecture, we identified intrachromosomal compartments, with B compartments engaging in homotypic interactions and enriched for retrotransposons. Furthermore, we detected self-interacting TAD-like structures with boundaries enriched in active chromatin marks and putative insulator motifs that resemble the binding sites of a mammalian architectural protein. Additionally, we defined highly insulated TADs that form an interaction network of heterochromatic islands and harbor genes related to pathogenicity. A subset of TADs showed transcriptional co-regulation patterns that varied across infection stages, indicating a regulatory role for TAD organization in infection stage-specific gene expression. Our results establish a multilayered view of 3D genome organization in *Z. tritici*, revealing chromatin conformation as a key component of genome function.

The core-accessory genome compartmentalization of *Z. tritici* extends beyond linear genome characteristics and is mirrored in the three-dimensional organization of the nucleus. Within this nuclear context, accessory chromosomes are preferentially associated with the pericentromeric regions of core chromosomes while remaining excluded from interactions with core chromosome arms. Interestingly, the accessory chromosomes harboring the weakest interactions with core chromosomes, specifically their

pericentromeric regions, were also the ones most frequently lost following treatment with carbendazim,

a fungicide that disrupts β-tubulin polymerization and impairs mitotic spindle assembly[28,41]. This

observation suggests that robust pericentromeric tethering may play a stabilizing role for accessory

chromosomes during mitosis. Similar phenomena have been observed in *S. cerevisiae*, where

microtubule-disrupting agents reduce centromere clustering, implying that centromeres are bundled at

the spindle pole body during interphase[42,43]. These parallels suggest that centromeric architecture in *Z.*

tritici may serve a comparable stabilizing function. Further supporting this hypothesis, accessory

chromosomes with stronger pericentromeric associations, such as chromosome 19, exhibited greater

stability under carbendazim treatment[28]. This chromosome also shows high prevalence in natural

populations of *Z. tritici*[27], aligning spatial stability with evolutionary persistence. In contrast, despite the

general presence/absence polymorphism trend among accessory chromosomes, those most frequently

lost under experimental conditions were not necessarily the least prevalent in global populations[27,34]. One

possible explanation for this discrepancy is that the strength of core-accessory chromosome interactions

may be strain-specific. It is likely that in different natural strains of *Z. tritici*, the core-accessory interactions

may vary, affecting the stability and presence of accessory chromosomes in a context-dependent manner.

Additionally, evolutionary pressures and local adaptation may influence the retention or loss of these

chromosomes, contributing to the observed diversity in the presence of accessory chromosomes across

strains [28]. Further research, including multiple-strain Hi-C analyses, could provide insights into how core-

accessory chromosome interactions have shaped the genomic architecture of *Z. tritici*.

In addition to the global Rabl-like organization of *Z. tritici*, we identified distinct intrachromosomal

compartmentalization by the presence of B compartments enriched in retrotransposons and constitutive

heterochromatin marks. These B compartments exhibit strong homotypic interactions, forming a spatially

distinct subnetwork largely insulated from the euchromatic A compartments, showing minimal inter-

compartment connectivity. Unlike the well-defined A/B compartmentalization observed in the human

genome[3] or the checkerboard interaction patterns reported in *R. irregularis*[19], *Z. tritici* displays a genome

topology that is dominated by internal B–B interactions. Compartmentalization inferred through principal

component analysis of Hi-C contact matrices revealed that B compartments are small (~60 kb), with a

similar size and positioning of TADs identified in our study. Specifically, these B compartments co-localize

with highly insulated TADs that form an interlinked network of heterochromatic regions. This suggests

that *Z. tritici* chromatin exhibit a heterochromatin-driven architecture, where B compartments and highly

insulated TADs act as spatial reservoirs for transcriptionally repressed, TE-rich genomic regions. This

compartmentalization likely reinforces silencing by sequestering these regions away from the

transcriptionally active A compartments. Such an arrangement parallels chromatin architectures observed

in other fungi[19,21,39]. For example, in *N. crassa*, heterochromatic domains marked by H3K9me3 aggregate

into discrete chromatin regions that contribute to genome stability[22,44]. In *Z. tritici*, the co-occurrence of

small B compartments with TAD-like structures supports a fine-scale, heterochromatin-driven genome

topology that enables both regulatory insulation and structural organization.

We identified self-interacting TAD-like domains with specific epigenetic characterization of their

boundaries, supporting the presence of chromatin-interacting domains as a universal genomic feature[45].

Z. tritici displays TAD-like domains of ~70 kb in size, similar to the average TAD size described in other

fungi[21,39], but far smaller than metazoan TADs, which typically span over megabases[45]. The detection of

TAD-like structures in *Z. tritici* was further supported by analysis of contact probability decay slopes

consistent with expectations from polymer physics modeling. These patterns, showing elevated

interaction frequencies at sub-100 kb distances, are consistent with the presence of interaction domains[4].

Unlike TADs described in metazoans, we counted only a subset of co-regulated TADs, suggesting that

structural features of the genome coordinate local transcriptional regulation during host-pathogen

interactions, similar to other plant-associated fungi[21,39]. In metazoans, the concept of TADs is also strongly

linked to the active loop extrusion creation process, which is dependent on cohesin and the boundary-

associated protein CTCF, or other insulator proteins [46]. However, self-interacting domains across the tree

of life were described in the absence of insulator proteins, as well as cohesin[47,48]. For instance,

chromosomal interaction domains (CIDs) in prokaryotes structure the genome at the gene operon level,

with insulated boundaries associated with highly expressed genes[49–51]. In *Z. tritici*, TAD boundaries are

gene-dense, TE-poor, and enriched in facultative heterochromatin histone modifications, consistent with

boundaries described in *V. dahliae*[21]. In *N. crassa*, interspersed heterochromatin blocks have been

proposed to act as domain boundaries in place of insulator proteins[18,22]. However, our findings in *Z. tritici*

suggest a more nuanced organization, as we observed multiple TADs between heterochromatic blocks,

with well-defined boundaries marked by distinct epigenetic profiles that are similar to those in *V.*

dahliae[21]. Furthermore, we observed a significant enrichment of zinc finger transcription factor DNA-

binding motifs at *Z. tritici* domain boundaries, notably those associated with the vertebrate insulator

protein ZNF263[52]. In *V. dahliae*, boundary-enriched GAAG-motifs and TATA-motifs associate with a

decrease in chromatin accessibility[21]. Interestingly, the *Z. tritici* IPO323 genome contains a single

annotated gene (ZtIPO323_082310) that shares a similar domain architecture to ZNF263, suggesting that

insulator-like proteins may contribute to domain boundary formation in fungi.

Globally, TAD-like structures exhibit low insulation scores in *Z. tritici*, consistent with previous

observations in other fungi[21]. However, we identified a subset of highly insulated domains (HI-TADs) that

coincide with TE-rich, heterochromatic regions. Similar patterns have been described in *V. dahliae* and

Epichloë festucae, where accessory genomic regions (AGRs) form structurally distinct and functionally

specialized chromatin domains with stronger local interactions compared to surrounding regions[21,39]. In

Z. tritici, these HI-TADs also engage more frequently in strong interactions with one another compared to

R-TADs, which seems reminiscent of the spatial network formed by the "TAD cliques" observed in the

human genome, where clusters of heterochromatic domains co-localize in 3D space[53]. Comparable

chromatin bundling mechanisms have also been described in *C. elegans*, where H3K9me3 promotes

chromosome compaction via HP1-mediated heterochromatin interactions[54]. In *N. crassa*, interspersed

heterochromatin blocks are described to function as boundaries, structuring TAD-like domains. Targeted

removal of a heterochromatic block induces localized architectural changes with minimal effects on

fitness[44]. The heterochromatin interaction network in *Z. tritici* shares conceptual similarities with the

KNOT Engaged Elements (KNOTs) described in *Arabidopsis thaliana*[55]. KNOTs are heterochromatic islands

within euchromatin that interact both within and between chromosomes. Notably, TEs preferentially

insert into these regions, and under heat stress, KNOT-associated TEs are derepressed, demonstrating

that these structures are dynamic and functional in TE regulation[56]. Previous studies in *Z. tritici* have

shown that wheat infection is accompanied by a global activation of TEs and epigenetic changes[36]. We

found that HI-TADs are enriched for effector genes, and some of these TADs are co-regulated *in planta,*

suggesting potential derepression of HI-TADs during infection. Although the mechanisms and dynamics of

chromatin folding during host colonization remain largely uncharacterized in fungi, the presence of

specialized 3D genome organization represents an evolutionary strategy to compartmentalize potentially

disruptive genomic elements while preserving their accessibility for regulated expression under specific

developmental or stress-related conditions.

CONCLUSION

The fundamental principles of genome folding, such as compartmentalization and domain organization, are broadly conserved across eukaryotes, even if the canonical definitions developed in metazoan systems do not always apply uniformly across the tree of life[45]. In the fungus *Z. tritici*, we uncovered a complex, multilayered chromatin architecture characterized by centromere clustering, spatial segregation between core and accessory chromosomes, a long-range heterochromatin network, and TAD-like domain structures. Together, these features likely contribute to this species' remarkable genome plasticity, facilitating regulatory flexibility, adaptive potential, and pathogenicity. By generating a high-resolution map of *Z. tritici*'s 3D genome, we lay the groundwork for mechanistic investigations into how chromatin structure governs gene expression, transposable element dynamics, and genome evolution in filamentous fungi. More broadly, this work advances the emerging field of comparative 3D genomics in non-model species, opening new avenues for exploring genome architecture as a driver of fungal adaptation and plant-pathogen interactions.

METHODS

Sample preparation and Hi-C sequencing

The reference strain *Z. tritici* IPO323[25] was preserved in 50% glycerol at -80°C. Pre-cultures were prepared with inoculations of ~100uL of glycerol stock spore suspension onto solid YMS medium (4 g/L yeast extract, 4 g/L malt extract, 4 g/L sucrose, and 12 g/L of agar) and grown for 4 days at 18°C. Spores were scraped from plates and re-suspended into PBS buffer (8g/L NaCl, 1.44g/L Na_2HPO_4, 0.2g/L KCl, 0.24g/L KH_2PO_4) for sample preparation for epigenomics sequencing as described in Möller et al. 2023[31]. Spores were fixed with formaldehyde according to PhaseGenomics instructions (Seattle, Washington, USA). Hi-C

experiments were performed with the Phase Genomics Proximo Hi-C Kit (Fungal), which combines four restriction enzymes to ensure homogeneous digestion at AT-rich genomic regions. The resulting libraries were sequenced at Biomarker Technologies GmbH (BMKgene, Münster, Germany). The sequencing was performed with Illumina NovaSeq X, paired-end 150bp reads. Sample preparation, Hi-C experiment, and sequencing were performed for two independent replicates.

Hi-C data processing, binning, and visualization

Each Hi-C read-pairs was filtered and trimmed using Trimmomatic v0.39[57] with the following settings ILLUMINACLIP:${adapters}:2:30:10 LEADING:15 TRAILING:15 SLIDINGWINDOW:5:15 MINLEN:50. We used HiCUP-0.9.2[58] to evaluate the quality of each Hi-C dataset. Reads were mapped against the reference genome assembly of *Z. tritici* IPO323[25] using bwa-mem2[59]. The construction and analysis of Hi-C interaction matrices were performed using the HiCExplorer suite of tools[60]. Each replicate interaction matrix was created using *hicBuildMatrix*, by aligning the paired reads to the reference genome digested *in silico* with the four restriction enzymes DpnII (GATC), DdeI (CTNAG), HinfI (GANTC), and MseI (TTAA). Contact matrices were individually corrected using the iterative correction and eigenvector decomposition (ICE) method via the *hicCorrectMatrix* function. The corrected contact matrices from the two biological replicates were combined to generate Hi-C matrices at 1 kb, 3 kb, 10 kb, and 50 kb. The Hi-C contact maps were generated with *hicPlotMatrix*. The most significant pairwise interactions were extracted by filtering the observed-over-expected values above 3.5 for inter- and 1.75 for intra-chromosomal interactions. The interaction link plots were generated using Circos-0.69[61]. The HiCExplorer .h5 contact matrix files were converted into .cool format with *hicConvertFormat* for compatibility with the *cooltools* and *cooler* packages for visualization[62]. All scripts for the analyses and visualization are available at https://github.com/lorraincecile/Hi-C_Ztritici_IPO323 and 10.5281/zenodo.15395556.

Estimation of chromatin folding patterns with contact probability decay

To characterize chromatin folding behavior, we computed the decay of contact frequency as a function of genomic distance, a method grounded in polymer physics models of chromosome organization[4]. These models typically distinguish between different chromatin configurations based on the decay exponent (α) of the contact probability P(s), where *s* represents genomic separation. Contact decay slopes provide insights into the folding state: an exponent of ~1.5 corresponds to an equilibrium globule, indicative of tightly knotted, inaccessible chromatin; values between 1 and 2 reflect a fractal globule, a compact yet unknotted state; and exponents < 1 are characteristic of a tension globule, normally associated with finer scale self-interacting domain units[63]. We used the *cooltools.expected_cis*[62] function to compute the average contact probability, P(s), separately for core and accessory chromosomes. Genomic distances and P(s) values were transformed to log10 scales, and a linear regression was fitted to the resulting log–log data to estimate the decay exponent (α), which serves as a proxy for the underlying chromatin folding state.

Statistical Analysis of Chromosome Loss and Interaction Strength

To investigate the relationship between chromosome loss and interaction strength in *Z. tritici* IPO323, we extracted the average number of core-accessory chromosome strong interactions (observed vs. expected > 3.5), normalized by the size of the corresponding accessory chromosome. This normalization yielded the interaction strength per accessory chromosome. We compared this measure to the frequency of accessory chromosome loss following beta-tubulin inhibition, described in Habig et al. (2017)[28]. A log-transformed Poisson regression model was used to assess the impact of normalized interaction strength on accessory chromosome loss. The model was fitted using the glm function in R (version 4.4.1), with chromosome loss modeled as a Poisson-distributed response variable. The coefficients were tested for statistical significance using z-tests, with p-values calculated for both the intercept and the normalized

interaction strength variable. Model diagnostics, including overdispersion testing using the *dispersiontest*

function from the AER package[64], confirmed that the Poisson distribution was an appropriate fit. The fitted

model was then used to predict chromosome loss, and the observed versus predicted values were

visualized with ggplot2[65].

A/B compartmentalization analysis

Compartmentalization analysis used the matrix at 3 and 10 kb resolution with functions from the cooltools

package[62]. We calculated PCA eigenvectors for intra- and inter-chromosomal contacts as *expected_cis* and

expected_trans, respectively. GC content was used as a phasing track to orient the eigenvector values.

The first three eigenvector values were manually checked, and the first eigenvector was determined to

have the best compatibility with the interaction pattern of the contact maps. We used the 2.5% and 97.5%

percentile eigenvector values as the minimum and maximum range to calculate the compartmentalization

strength. The compartmentalization plot was generated using the *saddleplot* function[62].

Topology-associated domains (TAD) calling

We established 3 kb as the smallest bin size at which at least 1000 reads covered 80% of bins for each

chromosome[66], and therefore the smallest scale at which we can reliably call TAD-like structures

(Supplementary Table S2). TAD predictions were then generated using *hicFindTADs* for HiCExplorer[60] with

default parameters using the contact matrix at 3kb bin size. This function calculates the insulation score

by evaluating interaction frequencies between adjacent bins. TAD boundaries were identified at

significant local minima in physical interactions, where large differences between adjacent bins yield

negative insulation scores, indicating "strong" insulation[60]. TADs were classified into "regular" (R-TADs)

and "high insulation" (HI-TADs) TADs by calculating the difference between each TAD's maximum

insulation score and the minimum insulation scores of its adjacent TADs. TADs with a score difference

within the 20th quantile, insulation scores above the median, and flanked by two neighboring TADs with

negative insulation scores were considered HI-TADs.

Analysis of genomic features

Gene annotations with associated GO terms, PFAM and effector predictions were sourced from Lapalu et

al. (2023)[67] and accessible here (https://bioinfo.bioger.inrae.fr/portal/genome-portal/12/). Centromere

detections and annotations were sourced from Schotanus et al. (2015)[29]. Core genes were identified as

described in Minana-Posada et al. (2024)[68] as shared across 19 *Z. tritici* strains[37]. GO term enrichment

analysis was performed using *topGO*'s *runGOEnrichment* function[69]. GO terms with no gene associations

in the query lists or supported by only a single gene across the genome were excluded. False discovery

rate (FDR) correction was applied to all *p*-values. Enrichment analyses of effectors, core genes, and PFAM

domains within gene sets were performed using Fisher's exact tests, with FDR correction applied to *p*-

values. TE annotations were obtained from the manually curated library of Baril and Croll (2023)[70]. TAD

boundary motif identification and enrichment were performed with XSTREME with default parameters

and compared to JASPAR database motifs[71] using Tomtom from the MEME suite tools[72]. We used the RIP

composite index to identify RIP-like signatures in 1 kb sliding windows across the genome as described in

Lorrain et al. (2021)[35]. A positive RIP composite index value reflects the presence of RIP-like mutations.

The distribution of genes, TEs and RIP-like signatures across TADs and their boundaries was analyzed using

the EnrichedHeatmap package in R[73].

Gene and TE expression analyses

RNA-seq data for *Z. tritici* IPO323 during *in planta* infection, comprising four distinct infection stages, was

obtained from Haueisen et al. (2018)[74], while *in vitro* RNA-seq data was sourced from Möller et al.

(2023)[31]. After trimming low-quality reads, the resulting high-quality reads were aligned to the IPO323

reference genome using STAR[75], with parameters configured to allow the inclusion of multimapping reads.

To capture the expression of genes and TEs, multimapper reads were reassigned using TEtranscript[76],

which ensures precise allocation of reads to the transcript sequences[67]

(https://bioinfo.bioger.inrae.fr/portal/genome-portal/12/) and TE library of *Z. tritici* IPO323[70]. Differential

expression analysis was performed using DESeq2[77], comparing expression levels across the biotrophic

phases (stages A and B), necrotrophic phases (stages C and D)[74], and *in vitro* growth conditions.

Characterization of epigenetic profiles

Histone modifications, centromere detection, and cytosine DNA methylation (5mC) level enrichments

across the *Z. tritici* IPO323 genome were assessed using publicly available chromatin immunoprecipitation

followed by sequencing (ChIP-seq) and whole-genome bisulfite sequencing (WGBS) datasets. ChIP-seq

datasets for CenH3 and H3K9me2 were obtained from Schotanus et al. (2015)[29], while datasets for

H3K36me3, H4K20me3, H4K20me1, H3K27me3, H3K27me2, and H3K4me2 were sourced from Möller et

al. (2023)[31]. Reads were preprocessed using Trimmomatic v0.39[57], with parameters

ILLUMINACLIP:${adapters}:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:30 and

ILLUMINACLIP:${adapters}:2:30:10:2:keepBothReads HEADCROP:5 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:30 for the single and paired end samples respectively. The filtered reads

were mapped to the IPO323 reference genome using Bowtie2 with default settings[78]. To account for the

repetitive nature of TEs in the genome, multimapping reads were subsequently mapped using Allo[79], an

approach tailored to improve mapping accuracy in repeat-rich regions. Reads with a mapping quality score

below 30 and PCR duplicates identified by Picard v2.26.2[80] were excluded from further analysis. To

visualize enrichment levels across the genome, histone enrichment profiles were generated as bigWig and

bedGraph files in bins of 10 bp, 3 kb, and 10 kb.

Bisulfite sequencing data was retrieved from Möller et al. (2021)[32] for cytosine DNA methylation analysis.

Cultures were inoculated directly from glycerol stocks into liquid YMS medium and grown for 5 days. The

Illumina HiSeq3000 paired-end raw reads were filtered using Trimmomatic[57] parameters HEADCROP:10

CROP:140 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:50 and subsequently mapped and

deduplicated using Bismark[81]. The methylated sites in all contexts (CG, CHG, CHH) were extracted using

the function *bismark_methylation_extractor,* and filtered to remove sites not supported by at least four

reads and more than 50% of the mapped reads. The weighted methylation average was calculated for 10

bp, 3 kb and 10 kb windows. The distribution of histone marks and 5mC levels across TADs and their

boundaries was analyzed using the EnrichedHeatmap package in R[73].

TAD co-expression analysis

To investigate the co-regulation of genes and TEs within TADs, we constructed a linear model in Python

as described in Winter et al. (2018)[39]. The model utilized the log2-fold change in gene expression between

in planta and *in vitro* stages as the response variable, with TAD membership as the main predictor. We

developed nested models by incorporating additional genomic features known to affect gene expression,

including GC content and distance to the nearest TE. The nested models' performance was evaluated

using the Akaike Information Criterion (AIC) and likelihood ratio tests. We defined TADs as co-regulated

TADs with more than 10 co-expressed genes and TEs.

Code availability

537   All custom scripts generated for data analyses and visualization are available at

538   https://github.com/lorraincecile/Hi-C_Ztritici_IPO323. The Hi-C contact matrices and TAD annotations as

539   well as source data for figures and tables presented in this study are available at

540   10.5281/zenodo.15395556.

Data availability

The Hi-C sequencing data is publicly available from the NCBI Gene Expression Omnibus (GEO) under the accession GSE296663. *In planta*[74] and *in vitro*[31] RNA-seq datasets are available at the GEO accession number GSE106136 and the accession number NCBI BioProjects PRJNA902413, respectively. The ChIP-seq datasets for CenH3 and H3K9me2[29] are accessible through the SRA ID SRP059394. The ChIP-seq datasets for H3K36me3, H4K20me3, H4K20me1, H3K27me3, H3K27me2, and H3K4me2[31] are available at the SRA ID SRP411885. Whole-genome bisulfite sequencing[32] data are available at SRA under BioProject ID PRJNA614493.

Acknowledgments

Ivona Glavincheska and Cecile Lorrain are funded by a SNSF Ambizione grant (PZ00P3_209022). We thank Prof. Bruce McDonald and ETH Zurich for funding the Hi-C sequencing.

Authors contributions

IG and CL designed the experiments. IG performed data acquisition. IG and CL performed the data analysis and interpretation. CL conceptualized the study. IG and CL wrote and revised the manuscript.

REFERENCES

1. Sexton, T. & Cavalli, G. The Role of Chromosome Domains in Shaping the Functional Genome. *Cell* **160**, 1049–1059 (2015).
2. Hoencamp, C. *et al.* 3D genomics across the tree of life reveals condensin II as a determinant of architecture type. *Science* **372**, 984–989 (2021).
3. Lieberman-Aiden, E. *et al.* Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome. *Science* **326**, 289–293 (2009).

4. Esposito, A. *et al.* Polymer physics reveals a combinatorial code linking 3D chromatin

architecture to 1D chromatin states. *Cell Reports* **38**, 110601 (2022).

5. Szalay, M.-F., Majchrzycka, B., Jerković, I., Cavalli, G. & Ibrahim, D. M. Evolution and

function of chromatin domains across the tree of life. *Nat Struct Mol Biol* **31**, 1824–1837

(2024).

6. Cremer, T. *et al.* Rabl's model of the interphase chromosome arrangement tested in Chinise

hamster cells by premature chromosome condensation and laser-UV-microbeam

experiments. *Hum Genet* **60**, 46–56 (1982).

7. Álvarez-González, L. *et al.* Principles of 3D chromosome folding and evolutionary genome

reshuffling in mammals. *Cell Reports* **41**, 111839 (2022).

8. Dixon, J. R. *et al.* Topological Domains in Mammalian Genomes Identified by Analysis of

Chromatin Interactions. *Nature* **485**, 376 (2012).

9. Long, H. S. *et al.* Making sense of the linear genome, gene function and TADs. *Epigenetics

& Chromatin* **15**, 4 (2022).

10. Flavahan, W. A. *et al.* Insulator dysfunction and oncogene activation in IDH mutant gliomas.

Nature **529**, 110–114 (2016).

11. Valton, A.-L. & Dekker, J. TAD disruption as oncogenic driver. *Current Opinion in Genetics

& Development* **36**, 34–40 (2016).

12. Dong, P. *et al.* 3D Chromatin Architecture of Large Plant Genomes Determined by Local

A/B Compartments. *Mol Plant* **10**, 1497–1509 (2017).

13. Li, E. *et al.* Long-range interactions between proximal and distal regulatory regions in maize.

Nat Commun **10**, 2633 (2019).

14. Ricci, W. A. *et al.* Widespread long-range cis-regulatory elements in the maize genome. *Nat

Plants* **5**, 1237–1249 (2019).

15. Dong, P., Tu, X., Liang, Z., Kang, B.-H. & Zhong, S. Plant and animal chromatin three-

dimensional organization: similar structures but different functions. *J Exp Bot* **71**, 5119–

5128 (2020).

16. Case, N. T. *et al.* Fungal impacts on Earth's ecosystems. *Nature* **638**, 49–57 (2025).

17. Belton, J.-M. *et al.* Hi-C: A comprehensive technique to capture the conformation of

genomes. *Methods* **58**, 10.1016/j.ymeth.2012.05.001 (2012).

18. Torres, D. E., Reckard, A. T., Klocko, A. D. & Seidl, M. F. Nuclear genome organization in

fungi: from gene folding to Rabl chromosomes. *FEMS Microbiology Reviews* **47**, fuad021

(2023).

19. Yildirir, G. *et al.* Long reads and Hi-C sequencing illuminate the two-compartment genome

of the model arbuscular mycorrhizal symbiont Rhizophagus irregularis. *New Phytologist*

233, 1097–1107 (2022).

20. Schalbetter, S. A., Fudenberg, G., Baxter, J., Pollard, K. S. & Neale, M. J. Principles of

meiotic chromosome assembly revealed in S. cerevisiae. *Nat Commun* **10**, 4795 (2019).

21. Torres, D. E. *et al.* Implications of the three-dimensional chromatin organization for genome

evolution in a fungal plant pathogen. *Nat Commun* **15**, 1701 (2024).

22. Galazka, J. M. *et al.* Neurospora chromosomes are organized by blocks of importin alpha-

dependent heterochromatin that are largely independent of H3K9me3. *Genome Res.* **26**,

1069–1080 (2016).

23. Rodriguez, S. *et al.* The genome organization of Neurospora crassa at high resolution

uncovers principles of fungal chromosome topology. *G3 Genes|Genomes|Genetics* **12**,

jkac053 (2022).

24. Stukenbrock, E. H., Banke, S., Javan-Nikkhah, M. & McDonald, B. A. Origin and

Domestication of the Fungal Wheat Pathogen Mycosphaerella graminicola via Sympatric

Speciation. *Molecular Biology and Evolution* **24**, 398–411 (2007).

25. Goodwin, S. B. *et al.* Finished Genome of the Fungal Wheat Pathogen Mycosphaerella

graminicola Reveals Dispensome Structure, Chromosome Plasticity, and Stealth

Pathogenesis. *PLOS Genetics* **7**, e1002070 (2011).

26. Feurtey, A. *et al.* Genome compartmentalization predates species divergence in the plant

pathogen genus Zymoseptoria. *BMC Genomics* **21**, 588 (2020).

27. Feurtey, A. *et al.* A thousand-genome panel retraces the global spread and adaptation of a

major fungal crop pathogen. *Nat Commun* **14**, 1059 (2023).

28. Habig, M., Quade, J. & Stukenbrock, E. H. Forward Genetics Approach Reveals Host

Genotype-Dependent Importance of Accessory Chromosomes in the Fungal Wheat

Pathogen Zymoseptoria tritici. *mBio* **8**, e01919-17 (2017).

29. Schotanus, K. *et al.* Histone modifications rather than the novel regional centromeres of

Zymoseptoria tritici distinguish core and accessory chromosomes. *Epigenetics & Chromatin*

8, 41 (2015).

30. Habig, M., Lorrain, C., Feurtey, A., Komluski, J. & Stukenbrock, E. H. Epigenetic

modifications affect the rate of spontaneous mutations in a pathogenic fungus. *Nat Commun*

12, 5869 (2021).

31. Möller, M., Ridenour, J. B., Wright, D. F., Martin, F. A. & Freitag, M. H4K20me3 is important

for Ash1-mediated H3K36me3 and transcriptional silencing in facultative heterochromatin in

a fungal pathogen. *PLoS Genet* **19**, e1010945 (2023).

32. Möller, M. *et al.* Recent loss of the Dim2 DNA methyltransferase decreases mutation rate in

repeats and changes evolutionary trajectory in a fungal pathogen. *PLOS Genetics* **17**,

e1009448 (2021).

33. Möller, M. *et al.* Destabilization of chromosome structure by histone H3 lysine 27

methylation. *PLOS Genetics* **15**, e1008093 (2019).

34. Möller, M., Habig, M., Freitag, M. & Stukenbrock, E. H. Extraordinary Genome Instability

and Widespread Chromosome Rearrangements During Vegetative Growth. *Genetics* **210**,

517–529 (2018).

35. Lorrain, C., Feurtey, A., Möller, M., Haueisen, J. & Stukenbrock, E. Dynamics of

transposable elements in recently diverged fungal pathogens: lineage-specific transposable

element content and efficiency of genome defenses. *G3 Genes|Genomes|Genetics* **11**,

jkab068 (2021).

36. Meile, L. *et al.* Chromatin Dynamics Contribute to the Spatiotemporal Expression Pattern of

Virulence Genes in a Fungal Plant Pathogen. *mBio* **11**, 10.1128/mbio.02343-20 (2020).

37. Badet, T., Oggenfuss, U., Abraham, L., McDonald, B. A. & Croll, D. A 19-isolate reference-

quality global pangenome for the fungal wheat pathogen Zymoseptoria tritici. *BMC Biol* **18**,

12 (2020).

38. Ortabozkoyun, H. *et al.* Members of an array of zinc-finger proteins specify distinct *Hox*

chromatin boundaries. *Molecular Cell* **84**, 3406-3422.e6 (2024).

39. Winter, D. J. *et al.* Repeat elements organise 3D genome structure and mediate

transcription in the filamentous fungus Epichloë festucae. *PLoS Genet* **14**, e1007467

(2018).

40. Shao, W. *et al.* The jet-like chromatin structure defines active secondary metabolism in

fungi. *Nucleic Acids Research* **52**, 4906–4921 (2024).

41. Davidse, L. C. & Flach, W. Differential binding of methyl benzimidazol-2-yl carbamate to

fungal tubulin as a mechanism of resistance to this antimitotic agent in mutant strains of

Aspergillus nidulans. *J Cell Biol* **72**, 174–193 (1977).

42. Goto, B., Okazaki, K. & Niwa, O. Cytoplasmic microtubular system implicated in de novo

formation of a Rabl-like orientation of chromosomes in fission yeast. *J Cell Sci* **114**, 2427–

2435 (2001).

43. Jin, Q. W., Fuchs, J. & Loidl, J. Centromere clustering is a major determinant of yeast

interphase nuclear organization. *J Cell Sci* **113 (Pt 11)**, 1903–1912 (2000).

44. Reckard, A. T. *et al.* A Constitutive Heterochromatic Region Shapes Genome Organization

and Impacts Gene Expression in Neurospora crassa. *bioRxiv* 2024.06.07.597955 (2024)

doi:10.1101/2024.06.07.597955.

45. Szalay, M.-F., Majchrzycka, B., Jerković, I., Cavalli, G. & Ibrahim, D. M. Evolution and

function of chromatin domains across the tree of life. *Nat Struct Mol Biol* 1–14 (2024)

doi:10.1038/s41594-024-01427-y.

46. Davidson, I. F. *et al.* CTCF is a DNA-tension-dependent barrier to cohesin-mediated loop

extrusion. *Nature* **616**, 822–827 (2023).

47. Oberbeckmann, E., Quililan, K., Cramer, P. & Oudelaar, A. M. In vitro reconstitution of

chromatin domains shows a role for nucleosome positioning in 3D genome organization.

Nat Genet **56**, 483–492 (2024).

48. Wing Shun Li *et al.* Mature chromatin packing domains persist after RAD21 depletion in 3D.

Science Advances (2025) doi:10.1126/sciadv.adp0855.

49. Trussart, M. *et al.* Defined chromosome structure in the genome-reduced bacterium

Mycoplasma pneumoniae. *Nat Commun* **8**, 14665 (2017).

50. Le, T. B. K., Imakaev, M. V., Mirny, L. A. & Laub, M. T. High-resolution mapping of the

spatial organization of a bacterial chromosome. *Science* **342**, 731–734 (2013).

51. Marbouty, M. *et al.* Metagenomic chromosome conformation capture (meta3C) unveils the

diversity of chromosome organization in microorganisms. *Elife* **3**, e03318 (2014).

52. Ortabozkoyun, H. *et al.* Members of an array of zinc-finger proteins specify distinct *Hox*

chromatin boundaries. *Molecular Cell* **84**, 3406-3422.e6 (2024).

53. Liyakat Ali, T. M., Brunet, A., Collas, P. & Paulsen, J. TAD cliques predict key features of

chromatin organization. *BMC Genomics* **22**, 499 (2021).

54. Qian Bian *et al.* Histone H3K9 methylation promotes formation of genome compartments in

Caenorhabditis elegans via chromosome compaction and perinuclear anchoring.

Proceedings of the National Academy of Sciences of the United States of America **117**,

11459–11470 (2020).

55. Grob, S., Schmid, M. W. & Grossniklaus, U. Hi-C Analysis in *Arabidopsis* Identifies the

KNOT, a Structure with Similarities to the *flamenco* Locus of *Drosophila*. *Molecular Cell* **55**,

678–693 (2014).

56. Sun, L. *et al.* Heat stress-induced transposon activation correlates with 3D chromatin

organization rearrangement in Arabidopsis. *Nat Commun* **11**, 1886 (2020).

57. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* **30**, 2114–2120 (2014).

58. Wingett, S. *et al.* HiCUP: pipeline for mapping and processing Hi-C data. *F1000Res* **4**, 1310

(2015).

59. Vasimuddin, Md., Misra, S., Li, H. & Aluru, S. Efficient Architecture-Aware Acceleration of

BWA-MEM for Multicore Systems. in *2019 IEEE International Parallel and Distributed*

Processing Symposium (IPDPS) 314–324 (2019). doi:10.1109/IPDPS.2019.00041.

60. Ramírez, F. *et al.* High-resolution TADs reveal DNA sequences underlying genome

organization in flies. *Nature Communications* **9**, 189 (2018).

61. Krzywinski, M. I. *et al.* Circos: An information aesthetic for comparative genomics. *Genome*

Res. (2009) doi:10.1101/gr.092759.109.

62. Open2C *et al.* Cooltools: Enabling high-resolution Hi-C analysis in Python. *PLOS*

Computational Biology **20**, e1012067 (2024).

63. Sanborn, A. L. *et al.* Chromatin extrusion explains key features of loop and domain

formation in wild-type and engineered genomes. *Proc Natl Acad Sci U S A* **112**, E6456-

6465 (2015).

64. Kleiber, C. & Zeileis, A. Diagnostics and Alternative Methods of Regression. in *Applied*

Econometrics with R (eds. Kleiber, C. & Zeileis, A.) 1–27 (Springer, New York, NY, 2008).

doi:10.1007/978-0-387-77318-6_4.

65. Wickham, H. *Ggplot2*. (Springer International Publishing, Cham, 2016). doi:10.1007/978-3-

319-24277-4.

66. Rao, S. S. P. *et al.* A 3D map of the human genome at kilobase resolution reveals principles

of chromatin looping. *Cell* **159**, 1665–1680 (2014).

67. Lapalu, N. *et al. Improved Gene Annotation of the Fungal Wheat Pathogen* Zymoseptoria

Tritici *Based on Combined Iso-Seq and RNA-Seq Evidence*.

http://biorxiv.org/lookup/doi/10.1101/2023.04.26.537486 (2023)

doi:10.1101/2023.04.26.537486.

68. Miñana-Posada, S., Lorrain, C., McDonald, B. A. & Feurtey, A. Thermal adaptation in

worldwide collections of a major fungal pathogen. 2024.09.12.612681 Preprint at

https://doi.org/10.1101/2024.09.12.612681 (2024).

69. topGO. *Bioconductor* http://bioconductor.org/packages/topGO/ (2024).

70. Baril, T. & Croll, D. A pangenome-guided manually curated library of transposable elements

for Zymoseptoria tritici. *BMC Research Notes* **16**, 335 (2023).

71. Sandelin, A., Alkema, W., Engström, P., Wasserman, W. W. & Lenhard, B. JASPAR: an

open-access database for eukaryotic transcription factor binding profiles. *Nucleic Acids*

Research **32**, D91–D94 (2004).

72. MEME Suite: tools for motif discovery and searching | Nucleic Acids Research | Oxford

Academic. https://academic.oup.com/nar/article/37/suppl_2/W202/1135092.

73. Gu, Z., Eils, R., Schlesner, M. & Ishaque, N. EnrichedHeatmap: an R/Bioconductor package

for comprehensive visualization of genomic signal associations. *BMC Genomics* **19**, 234

(2018).

74. Haueisen, J. *et al.* Highly flexible infection programs in a specialized wheat pathogen. *Ecol*

Evol **9**, 275–294 (2018).

75. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

76. Jin, Y., Tam, O. H., Paniagua, E. & Hammell, M. TEtranscripts: a package for including

transposable elements in differential expression analysis of RNA-seq datasets.

Bioinformatics **31**, 3593–3599 (2015).

77. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

78. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,

357–359 (2012).

79. Morrissey, A., Shi, J., James, D. Q. & Mahony, S. Accurate allocation of multimapped reads

enables regulatory element analysis at repeats. *Genome Res* **34**, 937–951 (2024).

80. Picard Tools - By Broad Institute. https://broadinstitute.github.io/picard/.

81. Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-

Seq applications. *Bioinformatics* **27**, 1571–1572 (2011).

82. Minana-Posada, S., Feurtey, A., Alassimone, J., McDonald, B. A. & Lorrain, C. Responses

to temperature shocks in Zymoseptoria tritici reveal specific transcriptional reprogramming

and novel candidate genes for thermal adaptation. 2024.12.16.628617 Preprint at

https://doi.org/10.1101/2024.12.16.628617 (2024).

LEGENDS

Fig. 1: The Rabl-like three-dimensional structure of *Zymoseptoria tritici* is compartmentalized. a)

Heatmap of genome-wide normalized Hi-C contact map at 10kb resolution. Contact numbers per 10kb

bins are represented in a log scale. Chromosomes are represented by ideograms with purple marking the

centromeres[29]. b) Strongest inter-chromosomal interactions with an observed-to-expected value >3.5 at

10 kb bin resolution. Core chromosomes are represented in light blue. Accessory chromosomes are

represented in dark blue. Centromeres are marked by triangles. H3K9me3 enrichment is shown in the

orange histogram. Links represent inter-chromosomal pericentromere-pericentromere interactions

(purple), accessory-core chromosome (green), or any other interactions (grey). c) Contact probability (top)

and contact frequency decay slope (bottom) against genomic distance for core (green) and accessory

(orange) chromosomes.

Fig. 2: Homotypic interactions and epigenomic features of A/B compartments in the *Zymoseptoria tritici*

genome. a) Saddle plot representing genome-wide intrachromosomal Hi-C interactions at 3 kb resolution.

Homotypic B-B interactions are represented in the top-left, while A-A interactions are observed in the

bottom-right. b) Enrichment of histone modifications across A and B compartments, highlighting distinct

chromatin states. c) Genomic composition of A and B compartments, showing proportions of genes

(green), transposable elements (orange), and intergenic regions (grey). d) Density of genes (green) and

transposable elements (orange) across A and B compartments. e) Distribution of transposable element

families within A and B compartments. f) *In vitro* Expression levels of genes (green) and transposable

elements (orange) in A and B compartments.

Fig. 3: The genome of *Zymoseptoria tritici* forms topology-associated domain-like (TAD-like) structures

with specific genomic and epigenomic signatures. a) Close-up of Hi-C contact map showing local

interaction frequencies at 3 kb bins resolution and detection of self-interacting TAD-like structures.

Insulation score was calculated with a genome-wide sliding window approach. Predicted TAD-like domains

are represented by black triangles, and boundaries are indicated by red arrows. b) Heatmap of chromatin

insulation scores, histone modifications (H3K36me3 and H3K9me3), gene, and TE density enrichment

centered on domain boundaries, with 35 kb upstream and downstream regions shown per row. The top

panel shows the average profile across all boundaries depicted in the heatmap. c) Close-up example of

TAD-like annotations on chromosome 4 of *Z. tritici* IPO323. From top to bottom: strongest intra-

chromosomal links (observed/expected > 1.75) across chromosome 4; chromosome 4 Hi-C contact map

with TAD annotated by triangles; insulation scores within chromosome 4; histone modifications for

euchromatin H3K4me2, heterochromatin H3K9me2, H3K9me3, and facultative heterochromatin

H3K27me2, H3K27me3, H3K36me3, H3K20me1, H4K20me3. d) Heatmap of chromatin insulation scores,

histone modifications, 5-methylcytosine DNA methylation (5mC), repeat-induced point mutation (RIP)

composite index frequency, genes, and TEs density centered on TAD-like domains, with 35 kb upstream

and downstream regions shown per row. The top panel shows the average profile across all TADs depicted

in the heatmap. Highly-insulated TADs (HI-TADs) are depicted by solid lines, and "regular-TADs" (R-TADs)

are depicted by dashed lines.

Supplementary Fig. S1: Quality assessment of the Hi-C libraries replicates. a) Pearson correlation

between Hi-C contact map replicates in *Zymoseptoria tritici* IPO323 (Pearson correlation = 0.87). b) Hi-C

libraries quality filtering with the HiCUP-0.9.2[58] pipeline.

Supplementary Fig. S2: *Zymoseptoria tritici* IPO323 Hi-C contact map 10kb bins normalized per genomic

distance as observed/expected interactions.

Supplementary Fig. S3: Strongest interactions in *Zymoseptoria tritici* IPO323. Links represent the

strongest interactions with observed-to-expected values >3.5 at 10 kb bin resolution for a) all strong

interactions, b) inter-chromosomal pericentromere-pericentromere interactions, c) accessory-core

chromosome interactions with inter-chromosomal pericentromere-pericentromere interactions (purple),

accessory-core chromosome (green), H3K9me3-H3K9me3 interactions (orange) or any other interactions

(grey), d) H3K9me3-H3K9me3 interactions.

Supplementary Fig. S4: Relationship between chromosomal interaction strength and chromosome loss

in *Zymoseptoria tritici* strain IPO323, after carbendazim treatment[28]. Scatterplot showing the association

between log-transformed normalized Hi-C interaction strength (x-axis) and the frequency of chromosome

loss (y-axis) across individual chromosomes. A dashed grey line indicates the predicted linear relationship.

The model indicates a significant inverse relationship between interaction strength and chromosome loss

(Poisson GLM, $p < 0.05$).

Supplementary Fig. S5: Interchromosomal compartmentalization in *Zymoseptoria tritici* IPO323. a) Self-

interacting compartments were identified from Hi-C contact matrix at 10 kb resolution. Principal

component analysis (PCA) of interchromosomal contact frequencies was used to extract the first

eigenvector (E1), with positive and negative values annotated in yellow and purple, respectively. The

eigenvector track is shown above each panel. The Hi-C contact map with 10 kb resolution is shown for all

chromosomes, shown by ideograms on the side, with centromeres represented in purple. b) Saddle plot

representing genome-wide Hi-C interactions at 10 kb resolution. Homotypic interactions between regions

associated with a negative E1 are represented in the top-left, while positive E1 region interactions are

observed in the bottom-right.

Supplementary Fig. S6: Intrachromosomal A/B compartments in *Zymoseptoria tritici* IPO323. For each

chromosome, A/B compartments were identified from Hi-C contact matrices at 3 kb resolution. Principal

component analysis (PCA) of intrachromosomal contact frequencies was used to extract the first

eigenvector (E1), with positive and negative values corresponding to A (yellow) and B (purple)

compartments, respectively. The eigenvector track is shown above each panel. Enrichment of H3K9me3

per 10 kb window is displayed as a histogram below the eigenvector track. Hi-C contact maps are shown

for individual chromosomes. Centromeres are represented by yellow stars.

Supplementary Fig. S7: TAD-like boundaries characteristics in core and accessory chromosomes of

***Zymoseptoria tritici* IPO323.** Heatmaps of chromatin insulation scores, enrichment in histone

modifications (H3K4me2, H3K9me2, H3K9me3, H3K27me2, H3K27me3, H3K36me3, H4K20me1,

H4K20me3), 5-methylcytosine DNA methylation (5mC), repeat-induced point mutation (RIP) composite

index frequency, genes, and TEs density centered on domain boundaries, with 35 kb upstream and

downstream regions shown per row. The top panel shows the average profile across all boundaries

depicted in the heatmap.

Supplementary Fig. S8: *In vitro* Expression levels of genes in topology-associated domains and

boundaries. Genes are represented in green and transposable elements in orange. Statistical significance

of differences between TADs and boundaries was assessed using a two-sided Mann-Whitney U test.

Supplementary Fig. S9: Strongest interactions between topology-associated domains (TADs).

Distribution of observed-to-expected contact ratios > 1.75 from the 10 kb resolution interaction matrix,

highlighting the strongest intrachromosomal interactions classified as either highly insulated TADs (HI-

TADs) or regular TADs (R-TADs). Statistical significance of differences between groups was assessed using

a two-sided Mann-Whitney U test.

Supplementary Fig. S10: *In vitro* Expression levels of genes in regular topology-associated domains (R-

TADs) and highly insulated topology-associated domains (HI-TADs). Genes are represented in green and

transposable elements in orange. Statistical significance of differences between HI-TADs and R-TADs was

assessed using a two-sided Mann-Whitney U test.

Supplementary Fig. S11: Co-regulated topology-associated domains (TADs) during wheat infection

stages. Effect sizes from linear regression models quantified the contribution of each topologically

associating domain (TAD) to differential gene expression between in vitro growth on YMA medium and

various in planta infection stages (stage A: early biotrophic, stage B: late biotrophic, stage C: early

necrotrophic, and stage D: late necrotrophic). Each point represents the mean effect size per TAD,

accompanied by a 95% confidence interval. TADs with statistically significant effects (i.e., confidence

intervals not overlapping zero) are highlighted in color and annotated with their chromosome and TAD

identifiers. Values reflect normalized means derived from two independent biological replicates [74].

Supplementary Fig. S12: Overlap between TAD categories. Venn diagram illustrating the overlap among

three categories of topology-associated domains (TADs): co-regulated TADs (defined as containing ≥10

genes or transposable elements with coordinated expression), highly insulated TADs (HI-TADs; identified

by elevated insulation scores), and accessory chromosome TADs (located on accessory chromosomes).

Supplementary Table S1: Quality assessment of Hi-C sequencing data.

Supplementary Table S2: Hi-C matrix contact coverage per chromosome.

Supplementary Table S3: Strongest interchromosomal interactions. Hi-C interactions were normalized

in observed vs expected (o/e) contacts.

Supplementary Table S4: Contact frequency decay exponent per chromosome.

Supplementary Table S5: A/B compartments from PCA eigenvector analysis.

Supplementary Table S6: Gene/TE content of A/B compartments.

Supplementary Table S7: TE content of A/B compartments.

Supplementary Table S8: Gene ontology enrichment in A compartment.

Supplementary Table S9: Gene ontology enrichment in B compartment.

Supplementary Table S10: Topology-associated domains (TADs) annotation in the *Z. tritici* IPO323

genome.

Supplementary Table S11: TAD boundaries annotations in the *Z. tritici* IPO323 genome.

Supplementary Table S12: TADs summary metrics in core and accessory chromosomes.

Supplementary Table S13: Core orthologous genes enrichment in TAD boundaries. The core orthologous

genes are present in 19 reference genomes of *Z. tritici* [37,82].

Supplementary Table S14: PFAM domains enrichment in TAD boundaries.

Supplementary Table S15: Gene ontology enrichment in TAD boundaries.

Supplementary Table S16: Motifs enrichment in TAD boundaries.

Supplementary Table S17: ZNF263 BLASTp searches in *Zymoseptoria tritici* IPO323 proteome

Supplementary Table S18: Annotation of highly insulated TADs (HI-TADs) and regular TADs (R-TADs).

Supplementary Table S19: Effector gene enrichment in highly insulated TADs (HI-TADs)

Supplementary Table S20: Gene/TE expression analysis *in planta* stage A (early biotrophic phase)

versus *in vitro*.

Supplementary Table S21: Gene/TE expression analysis *in planta* stage B (late biotrophic phase) versus

***in vitro*.**

Supplementary Table S22: Gene/TE expression analysis *in planta* stage C (early necrotrophic phase)

versus *in vitro*.

Supplementary Table S23: Gene/TE expression analysis *in planta* stage D (late necrotrophic phase)

versus *in vitro*.

Supplementary Table S24: Table S23: Linear models for TAD co-expression analyses.

Supplementary Table S25: Summary metrics of co-regulated TADs.

a

10 kb Bin Reads (log1p)

1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13

Accessory chromosomes

c

Core chromosomes
Accessory chromosomes

Contact probability

$\sim s^{-0.75}$
$\sim s^{-1.5}$
$\sim s^{-1}$

Local slope

Genomic distance (bp)

b

Chr11, Chr10, Chr9, Chr8, Chr7, Chr6, Chr5, Chr4, Chr3, Chr2, Chr1, Chr12, Chr13, 14, 15, 16, 17, 19, 20, 21

H3K9me3

△ Centromeres
Pericentromere-pericentromere links
Accessory-core chromosome links
Other links

a

average observed/expected contact frequency

B+B

A+A

b

log2fc

	H3K4me2	H3K9me2	H3K9me3	H3K27me2	H3K27me3	H3K36me3	H4K20me1	H4K20me3
A	0.44	-0.08	-0.28	-0.02	-0.10	0.11	0.17	0.13
B	-1.29	1.35	1.58	0.09	0.41	-0.88	-0.65	-0.51

c

Gene annotation   TE annotation   No annotation

A: 80.3% 4.6% 15.1%

B: 39.2% 43.9% 16.9%

Total Length (Mb)

d

log2(count + 1) per 10 kb window

Gene Density   TE Density

***   ***

A   B   A   B

e

Percentage of all TEs in compartment

Compartment
A
B

DIRS · DNA transposons · DNA/Crypton · DNA/MULE-MuDR · DNA/Maverick · DNA/PIF-Harbinger · DNA/PiggyBac · DNA/TcMar · DNA/hAT · LINEs · LTR · LTR/Copia · MITEs · RC/Helitron · Retrotransposons · SINEs · Unclassified transposons

f

log2(RPKM+1)

Gene Expression   TE Expression

***   ***

A   B   A   B