An abundant merozoite surface protein of *Plasmodium falciparum* modulates

susceptibility to inhibitory antibodies

Isabelle G. Henshall[1], Jill Chmielewski[1], Dimuthu Angage[2], Ornella Romeo[1], Keng Heng Lai[1,

3], Kaitlin R. Turland[1, 3], Nicki Badii[2], Michael Foley[2, 4], Robin F. Anders[2], James Beeson[5, 6,7]

and Danny W Wilson[1, 3, 5, *].

[1]Research Centre for Infectious Diseases, School of Biological Sciences, The University of

Adelaide, Australia 5005

[2]Department of Biochemistry and Chemistry, La Trobe Institute for Molecular Sciences, La

Trobe University, Victoria, Australia 3086

[3]Institute for Photonics and Advanced Sensing (IPAS), University of Adelaide, 5005, SA,

Australia

[4]AdAlta, 15/2 Park Drive, Bundoora, Victoria, Australia 3083

[5]Burnet Institute, 85 Commercial Road, Melbourne 3004, Victoria, Australia

[6]Department of Infectious Diseases, University of Melbourne, Victoria, Australia

[7] School of Translational Medicine and Department of Microbiology, Monash University, Victoria,

Australia

*Corresponding Author: danny.wilson@adelaide.edu.au

Abstract

Malaria merozoite surface proteins (MSPs), are thought to have important roles in red blood

cell (RBC) invasion and their exposure on the parasite surface makes them attractive vaccine

candidates. However, their role in invasion has not been directly demonstrated and their

biological functions are unknown. One of the most abundant proteins is *Pf*MSP2, which is

likely an ancestral protein that has been maintained in the *Plasmodium falciparum* lineage and

is a focus of vaccine development, whose function remains unknown. Using CRISPR-Cas9

gene-editing, we removed *Pf*MSP2 from two different *P. falciparum* lines with no impact on

parasite replication or phenotype *in vitro*, demonstrating that it is not essential for RBC

invasion. However, loss of *Pf*MSP2 led to increased inhibitory potency of antibodies targeting

other merozoite proteins involved in invasion, particularly *Pf*AMA1. In a solid-phase model,

increasing concentrations of *Pf*MSP2 protein reduced binding of different antibodies against

*Pf*AMA1 in a dose dependent manner. These data suggest that *Pf*MSP2 can modulate the

susceptibility of merozoites to protective inhibitory antibodies. The results of this study change

our understanding of the potential functions of *Pf*MSP2 and establishes a new concept in

malaria where a surface protein can reduce the protective efficacy of antibodies targeting a

different antigen. These findings have important implications for understanding malaria

immunity and informing vaccine development.

Introduction

Plasmodium falciparum malaria remains a global health challenge with over half a million

deaths annually (World Health Organization, 2024). Worryingly, increasing parasite resistance

to frontline antimalarials and mosquito insecticide resistance are decreasing the effectiveness

of key control measures (World Health Organization, 2024). Development of the RTS,S and

R21 vaccines targeting the *P. falciparum* circumsporozoite protein is a step forward for malaria

vaccine development, however further improvements in vaccine efficacy and longevity are

needed to protect those most at risk – children under 5 years – and enable malaria elimination

(Beeson et al., 2019; Datoo et al., 2024; RTSS Clinical Trials Partnership, 2015).

Understanding the function of vaccine targets remains important for prioritisation and

optimisation of candidates in the development pipeline. Merozoites, the parasite stage that

invades host red blood cells (RBCs), have long been targeted for vaccine development since

successful blocking of merozoite invasion would prevent the repeated cycles of blood-stage

parasite multiplication and associated disease. While much work has been done to understand

the processes and steps required for merozoite invasion, only a few merozoite proteins have

successfully had functions defined. To date, most vaccines based on merozoite antigens have

demonstrated limited efficacy (Beeson et al., 2016) at least in part due to the extensive

polymorphisms exhibited by many of these antigens.

The surface of the merozoite is covered in a fibril coat which is comprised of multiple proteins,

broadly known as merozoite surface proteins (MSPs). These proteins have been proposed to

mediate weak initial merozoite RBC contact (Bannister et al., 1986; Beeson et al., 2016;

Cowman et al., 2017; Weiss et al., 2015) and in *P. falciparum* are dominated by GPI-anchored

proteins, the most abundant being *Pf*MSP1 and 2 (Gilson et al., 2006). *Pf*MSP1 has been

suggested to have a role in merozoite attachment to the RBC (Li et al., 2004), but recent

evidence instead suggests that it has a role in merozoite rupture from schizonts and cellular

interaction studies using optical tweezers do not support a substantial role for this protein in

RBC binding (Das et al., 2015; Kals et al., 2024), although it may have another function in

invasion. These recent insights into *Pf*MSP1 function, the best studied MSP, highlight what

little is known of MSPs and their roles in parasite survival.

P. falciparum merozoite surface protein 2 (*Pf*MSP2), a protein proposed to be essential for

parasite growth (Sanders et al., 2006), has been of long-term interest as a vaccine candidate.

*Pf*MSP2 is a ~28 kDa intrinsically disordered protein that has conserved N- and C- termini,

along with a central variable region that defines two main allelic groups, 3D7-like and FC27-

like (Adda et al., 2012). While *Pf*MSP2 was theorised to have a mechanical role in the early

steps of invasion (Anders et al., 2010), there is minimal supporting evidence for this, in part

due to the lack of tools to study *Pf*MSP2 function. Studies with recombinant proteins suggest

that it is likely *Pf*MSP2 interacts with lipids and forms complexes with itself on the merozoite

surface (Adda et al., 2009; Lu et al., 2019; Zhang et al., 2012). *Pf*MSP2 peptides have also

been reported to bind the RBC and inhibit *P. falciparum* growth, which would support *Pf*MSP2

mediating merozoite-RBC interactions (Ocampo et al., 2003). Unlike the majority of the

merozoite surface coat proteins, *Pf*MSP2 is not present in other malaria parasites that infect

humans and has been postulated to have evolved specifically in the *Laverania* subgenus of

Plasmodium, which includes *P. falciparum* (Black et al., 2002).

*Pf*MSP2 has been trialled in a vaccine (Combination B) combined with fragments of *Pf*MSP1

and *Pf*RESA. A Phase 1/2b trial of this vaccine showed efficacy in reducing the parasite burden

in children, which was specific to the *Pf*MSP2 3D7-like allele used in the vaccine formulation

(Genton et al., 2003). Vaccine-induced antibodies from clinical trials or experiments in animals

have little or no inhibitory activity in standard growth inhibition assays (Boyle et al., 2015;

McCarthy et al., 2011), but did show functional activity through the recruitment of

complement, promoting opsonic phagocytosis and antibody-dependent cellular inhibition

(Boyle et al., 2015; Feng et al., 2018; McCarthy et al., 2011). Naturally acquired antibodies are

also known to target *Pf*MSP2 and have been linked to protection (Fowkes et al., 2010; Osier et

al., 2010; Stanisic et al., 2009; Zerebinski et al., 2024). Studies have highlighted that it is the

Fc-mediated functional activities of these naturally acquired *Pf*MSP2 antibodies, as opposed

to direct blocking of *Pf*MSP2 protein function, that correlates best with protection (Boyle et

al., 2015, 2014; Osier et al., 2014; Reiling et al., 2019). While *Pf*MSP2 is a key target of naturally acquired malaria immunity, little is known about the function of *Pf*MSP2, hindering its advancement as a potential vaccine candidate.

Here we take a reverse genetics approach to assess the function of *Pf*MSP2, quantifying the impacts of *Pf*MSP2 deletion on parasite invasion and phenotype. Given our findings that deletion of MSP2 did not impact invasion *in vitro*, we tested the alternative hypothesis that merozoite surface proteins may function to modulate susceptibility of merozoites to inhibitory antibodies.

Results

Avian malaria MSP2-like proteins are structurally similar to *Laverania* MSP2s and show that MSP2 did not arise exclusively in this clade.

Genome annotations of *pfmsp2* show it lies between the merozoite surface protein *pfmsp5* (PF3D7_0206900) and *pfmsp*4 (PF3D7_0207000), and the conversed purine biosynthesis enzyme *adenylosuccinate lyase* (ADSL, PF3D7_0206700) (Figure 1A) (PlasmoDB; (Aurrecoechea et al., 2009)), with this arrangement previously thought to be restricted to the *Laverania* clade which includes parasites that infect great apes and *P. falciparum* (Black et al., 2002; Ferreira et al., 2015). As described by Escalante et al. (2022), we found the genomes of the two annotated avian *Plasmodium* spp. available on PlasmoDB, *P. relictum* and *P. gallinaceum*, to also have a putative gene between ADSL and MSP5 with structural similarities to *Laverania* MSP2s. The reduced length of the *P. relictum* (PRELSG_0415300) and *P. gallinaceum* (PGAL8A_00017700) MSP2-like proteins compared to those of *Laverania* parasites is accounted for by the absence of any extended central domains of the two avian malaria parasites (Supplementary Figure 1).

It is estimated that ~10 million years of evolution separate the avian malaria parasites *P. relictum* and *P. gallinaceum* from the *Laverania* (Böhme et al., 2018). Therefore, we explored whether similarities in amino acid sequence and predicted conformation would support a shared lineage and functional properties of MSP2 between *P. relictum* and *P. gallinaceum*

MSP2s with *Laverania* MSP2 sequences. The primary translation product of *Pf*MSP2 has

hydrophobic N- and C-terminal sequences that correspond to secretion signals (SP) and GPI-

attachment signals, respectively, which are absent from the mature polypeptide on the

merozoite surface. The putative SP (Supplementary Figure 1, Table 1) has an amino acid

similarity (BLSM62) of >89% across the *Laverania* and avian malaria parasites, whereas, the

C-terminal GPI signal sequences of ~24 residues shows >79% amino acid similarity and shared

hydrophobicity across the *Laverania* and avian malarias. Both *P. relictum* (Asp, Ala, Ser: based

on big-PI Predictor (Eisenhaber et al., 1999)) and *P. gallinaceum* (Asp, Gly Ala) MSP2s have

a sequence of three amino acids with short side chains immediately prior to this C-terminal

hydrophobic region which, although different to the conserved sequence seen with the

Laverania (Ser, Asn, Ile/Met (Smythe et al., 1991)), conforms to an amino acid sequence

expected for a GPI signal (Gilson et al., 2006) (Supplementary Figure 1). Typically, N-terminal

of a GPI cleavage site lies an unstructured linker region of up to 10 amino acids rich in polar

residues, as seen for *Laverania* MSP2s (polar residues: 50% 2 species to 70% 4 species). For

the *Laverania*, this region has almost no Acidic amino acids (Acidic residues: 0% 7 species to

10% 3 species) (Supplementary Figure 1, Table 1). In contrast, this region has reduced levels

of polar, but is enriched in acidic amino acids, for both *P. relictum* (20% polar, 40% acidic)

and *P. gallinaceum* (10% polar, 30% acidic). Although there are some differences in the N and

C-terminal regions between *P. relictum* and *P. gallinaceum* and the *Laverania* MSP2s, these

signal sequences are broadly conserved in properties which suggests a conserved sub-cellular

localisation.

Immediately downstream of the N-terminal SP lies the 24 amino acid N-terminal conserved

region (Supplementary Figure 2). This region of MSP2 has a propensity for alpha-helical

structure and fibril formation (Yang et al., 2010). Amino acid similarity through the N-terminal

conserved region is 78% across the *Laverania* and avian parasites, resulting in similar

proportions of hydrophobic, polar uncharged, basic and acidic amino acid across this region

(Table 1). Following the N-terminal conserved regions lies the dimorphic central variable

repeat region in *P. falciparum* MSP2s that are grouped as 3D7 or FC27-like amino acid

sequence structures. A defining feature of the 3D7-like MSP2 family are 4 or 8-mer tandem

repeats consisting of small amino acids (for 3D7: Gly, Gly, Ser, Ala) (MacRaild et al., 2015;

Smythe et al., 1991). The FC27-like MSP2 family also has repeats of 12 or 32 amino acids in

length, but these do not resemble those of the 3D7-like family. We found that the non-*P.*

falciparum Laverania and avian malaria MSP2 sequences compared consisted of >75% small

amino acids (some of: Gly, Ser, Ala, Thr) generally ordered in repeats downstream of the N-

terminal conserved region (Supplementary Figure 3). The number of repeats (4-repeats = 5

examples to 8 repeats = *P. relictum*), the length of the repeats (3-amino acids = 5 examples to

7-amino acids = *P. praefalciparum*) and whether the repeats were highly conserved or had

various levels of degeneracy (conserved = 4 examples to partially or highly degenerate = 10

examples) differed between and within species (Table 1, Supplementary Figure 3). A broader

search of an additional nine *Laverania* MSP2s identified from the NCBI experimental database

did not reveal any repeat or central variable regions that resembled FC27-like MSP2s (data not

shown). This comparison highlights the surprising finding that a region of known dimorphic

variability in *P. falciparum* MSP2 is likely to retain functional constraints that favour small

amino acid rich repeats immediately downstream of the N-terminal conserved region. Broadly

speaking, this groups the compared *Laverania* and avian malaria MSP2s as having a 3D7-like

repeat structure.

Upstream of the C-terminal GPI signal sequence lies the 50 amino acid residue C-terminal

conserved region (Supplementary Figure 4). This region contains two cysteine residues in all

Laverania MSP2s, which are thought to form an intramolecular disulphide bond (Zhang et al.,

2008). Two cysteine residues are present in the C-terminus of *P. gallinaceum*, but not *P.*

relictum MSP2 which has a single cysteine (Supplementary Figure 4). *Laverania* MSP2

similarity through the 50 residue sequence that makes up the C-terminal conserved region was

80%, with the hydrophobic residues making up between 20% (*P. billcollinsi*) and 34% (*P.*

adleri) of the amino acid content (Table 1). Amino acid similarity dropped to 62% when all

available C-terminal conserved domain residues were compared between *Laverania* and the

avian malarias due to the limited amino acid sequence similarity upstream of the cysteine

residue/s and apparent insertions of 20 (*P. relictum*) and 26 (*P. gallinaceum*) amino acids

downstream of the first Cysteine (Supplementary Figure 4). However, the percentage of

hydrophobic residues for *P. relictum* (33%) and *P. gallinaceum* (34%) was broadly in line with

that seen for the *Laverania*.

Extensive studies on recombinant proteins of the *P. falciparum* 3D7 and FC27 variants of

*Pf*MSP2 have shown this merozoite surface protein to be intrinsically disordered. Given the

evidence for conservation of the primary structure of MSP2s from distantly related malaria

parasites, we used AlphaFold2 predictions (Jumper et al., 2021) to examine whether there were

likely to be also conformational similarities between *Laverania* MSP2s and those of *P. relictum*

and *P. gallinaceum*. The MSP2s in the database of AlphaFold2-predicted structures retain their

N- and C-terminal signal sequences, but as these are both absent from mature MSP2 on the
merozoite surface they were deleted from the MSP2 sequences used for the AlphaFold2
predictions reported here. The predicted structures of the two avian parasite MSP2s, and other
Laverania MSP2s, are very similar to that of *Pf*3D7 MSP2, with most of the polypetide chains
lacking predicted secondary structure and, from the colour coding, have a predicted local
distance test (pLDDT) consistent with that of an intrinsically disordered protein over the
majority of the proteins length (Figure 1B). In contrast to the rest of the polypeptide, some α-
helical structure was predicted in the conserved 20-residue N-terminal region of all the
Laverania and avian MSP2s. NMR studies with recombinant *Pf*3D7 and *Pf*FC27 MSP2
showed this conserved N-terminal sequence to have a propensity for forming an α-helix that
was stabilized in the presence of lipid mimetics (MacRaild et al., 2012). The low pLDDT
values are consistent with this region in the other *Laverania* and *P. gallinaceum* MSP2 having
some propensity for α-helix formation, whereas higher pLDDT values suggest the possible
presence of a more stable α-helix in this region of *P. relictum* MSP2.

Having established that MSP2 is present across different *Plasmodium* lineages sharing similar
predicted amino acid compositions and structural features, the phylogenetic relationship
between annotated *Laverania* and putative avian malaria parasite MSP2s was examined. A
phylogenetic tree was generated based on alignments of representative sequences from each
species. MSP2 sequences from species belonging to the *Laverania* subgenus all grouped
together with two smaller clusters, corresponding to the known *Laverania* clade A and clade
B species (Figure 1C). As expected, *P. relictum* and *P. gallinaceum* MSP2 also group together.
The phylogenetic tree generated based on MSP2 sequences mirrors the proposed evolution of
Plasmodium spp. (Figure 1C). As postulated by Escalante et al. (2022) , these data suggest that
MSP2 was likely present in the ancestral *Plasmodium* parasite prior to the split of the
mammalian infecting *Plasmodium* spp. from those that infect birds and lizards. Given the
shared amino acid sequence and predicted conformational similarities, including the high levels
of intrinsically disordered sequence outside of the N- and C-terminal regions, the *P. relictum*
and *P. gallinaceum* MSP2-like proteins appear likely to contain enough properties to
potentially be functionally equivalent to *Laverania* MSP2s. This conservation of MSP2-like
proteins across different species suggests it has an important function.

Figure 1. Presence and structure of MSP2 across different *Plasmodium* species. **(A)** Schematic of the gene arrangement surrounding *msp2* on *P. falciparum* chromosome 2. **(B)** AlphaFold2 structural predictions of example *Laverania* (*P. falciparum* (3D7), *P. reichenowi*, *P. billcollinsi*, *P. adleri*) and the two avian (*P. relictum*, *P. gallinaceum*) malaria species. The N-terminal signal peptide and C-terminal GPI anchor sequences were removed before the proteins structure was predicted. The most N-terminal amino acid is indicated. Colours represent the predicted local distance difference test (pLDDT) scores, with dark blue representing very high confidence (>90%), light blue high confidence (90 to >70), yellow low confidence (70 to 50) and orange very low confidence (<50). the Modelled structure of the N-terminal helical region is provided as an inset for *P. falciparum* 3D7 MSP2 in isolation, and also in alignment with the helical domain for *P. falciparum* 3D7 (green for all) for the other examples of *Laverania* and avian MSPs (other colours). **(C)** Maximum likelihood tree showing relationship of MSP2 protein sequences found in different *P. falciparum* isolates and other *Plasmodium* species. Grouping of sequences from *P. falciparum* isolates into two main allele types can be seen as well as the separation of the *Laverania* and avian malaria species into their expected groups. Tree robustness tested by bootstrap.

Loss of *Pf*MSP2 does not noticeably impact growth or invasion of different *P. falciparum* lines *in vitro*.

Given previous unsuccessful attempts to disrupt *pfmsp2* (Sanders et al., 2006), and its high abundance on the merozoite surface (Gilson et al., 2006), *Pf*MSP2 has been traditionally viewed as an essential *P. falciparum* protein with an essential function in merozoite invasion. We employed CRISPR-Cas9 gene editing to determine whether the reported inability to knock-out *pfmsp2 in vitro* was a result of this protein being essential for parasite survival or because of the lower efficiency of the previously used gene-editing system (Sanders et al., 2006). Unexpectedly, we confirmed successful disruption of *pfmsp2* by replacing the coding sequence between 132 bp and 819 bp of the gene with a hDHFR drug selection cassette in the 3D7 *P. falciparum* laboratory-adapted line (Figure 2A and B), resulting in *Pf*3D7 ΔMSP2 parasites. We confirmed that *Pf*MSP2 was no longer expressed in *Pf*3D7 ΔMSP2 parasites by western blot, which detected a MSP2 band around 50 kDa in *Pf*3D7 WT parasite material but not in *Pf*3D7 ΔMSP2 parasites (Figure 2C). Similarly, immunofluorescence microscopy using anti-*Pf*MSP2 rabbit polyclonal antibodies showed the expected surface localisation of MSP2 on

259   *Pf*3D7 WT merozoites but not on *Pf*3D7 ΔMSP2 parasites (Figure 2D). We next assessed

260   whether knock-out of *Pf*MSP2 impacted on parasite growth over one or two cycles of

261   development *in vitro* and found that deletion of *Pf*3D7 MSP2 did not cause any significant

262   change in growth (Figure 2E and F).

Figure 2. *Pf*MSP2 is not essential for growth *in vitro* of *Pf*3D7 blood stage parasites. **(A-B)**

Schematic and agarose gel showing integration of knockout construct (band A+C) in the

msp2 gene locus and absence of the *msp2* sequence from the *Pf*3D7 ΔMSP2 line (band A+B).

(C) Western Blot of late schizont protein extracts confirms no *Pf*MSP2 is expressed in the

*Pf*3D7 ΔMSP2 line. *Pf*MSP2 detected by anti-*Pf*MSP2 2F2 3D7 mAb with *Pf*Aldolase

serving as loading control. Representative image shown. **(D)** Distribution of key merozoite

surface proteins in the presence or absence of *Pf*MSP2 was visualised by

immunofluorescence. *Pf*MSP2 (red for top two rows, purple in bottom two rows), the nucleus

stained by DAPI (blue in merge) and *Pf*AMA1 (purple, top two rows) or *Pf*MSP1 19 (red,

bottom two rows), and the coloured merge of the proceeding panels. Scale bar = 0.7 μm.

Representative images shown from a minimum of 10 schizonts imaged per condition. **(E-F)**

Growth of *Pf*3D7 WT compared to *Pf*3D7 ΔMSP2 *P. falciparum* parasites, measured as fold

increase in parasitaemia, over one (48 hrs) or two (96 hrs) cycles in either standard (still- (E))

or shaking (F) conditions. Parasitaemia was determined by flow cytometry at the start and

end to calculate fold increase. Graph displays mean ± S.D. of three independent experiments

performed with technical triplicates. Significance determined by unpaired t-test with $p < 0.05$

deemed significant.

As *Pf*MSP2 has been considered, to this point, as essential for *P. falciparum* growth *in vitro*

(Sanders et al., 2006), we investigated whether *Pf*MSP2 could also be removed from *Pf*Dd2,

an isolate of *P. falciparum* that differs from 3D7 in geographical origin, RBC receptor usage

and allelic type of *pfmsp2*. Using CRISPR-Cas9 we successfully knocked-out *msp2* in *Pf*Dd2

(Figure 3 A and B) and confirmed the absence of the protein in *Pf*Dd2 ΔMSP2 parasites using

western blot (Figure 3C). As we found for *Pf*3D7, deletion of *Pf*Dd2 *msp2* did not result in a

significant impact on parasite growth over one or two cycles of parasite replication (Figure

3D).

Given the abundance of *Pf*MSP2 on the merozoite surface, we postulated that loss of this

protein may impact on the timing of merozoite invasion of the RBC, which is not reflected in

reduced growth in *in vitro* cultures. Previous studies using live-cell microscopy have reported

that merozoites first contact the RBC and form weak initial interactions. The merozoite then

reorientates to bring apical organelles into position to release their contents onto the RBC,

which then leads to formation of the tight junction, merozoite invasion through the tight-

junction, formation of the parasitophorous vacuole and resealing of the RBC membrane post-

entry (Weiss et al., 2015). Given its surface localization, *Pf*MSP2 has been speculated to act in

the early phase of merozoite attachment. To assess whether knock-out of *pfmsp2* impacted on

the progression of invasion, we used live-cell microscopy of *Pf*Dd2 ΔMSP2 parasites and

compared the timing and key steps of invasion to *Pf*Dd2 WT parasites *in vitro*. This analysis

showed that *Pf*Dd2 ΔMSP2 knock-out parasites had no detectable alteration in the time the

parasite takes to attach to the RBC, the length or strength of RBC deformation as the merozoite

begins entry or for the time it took for invasion to be completed compared to *Pf*Dd2 WT (Figure

3 E-H). Together these data show *Pf*MSP2 is not essential for blood-stage replication *in vitro*

in two *P. falciparum* laboratory isolates from different geographical regions and knock-out of

*Pf*MSP2 does not seem to impact parasite growth or merozoite invasion *in vitro*. However, its

conservation across diverse *Plasmodium* spp. suggests it does play an important function or is

advantageous for parasite survival.

Figure 3. *Pf*Dd2 does not require MSP2 for asexual growth *in vitro*. **(A-B)** Successful integration of KO construct (schematic in A) into the *msp2* gene locus of *Pf*Dd2 ΔMSP2 was confirmed by PCR of genomic DNA (primers A+B amplify WT locus, primers D+E amplify integrated KO construct). **(C)** Loss of *Pf*MSP2 expression was demonstrated by western blot of schizont protein extract with *Pf*MSP2 detected by anti-*Pf*MSP2 FC27 and anti-*Pf*EXP2 as loading control. Representative image shown. **(D)** Growth of *Pf*Dd2 WT *P. falciparum* parasites and *Pf*Dd2 ΔMSP2 over one (48 hrs) or two (96 hrs) cycles. Parasitaemia was determined by flow cytometry at the start and end to calculate fold increase. Graph displays mean ± S.D. of three independent experiments performed with technical triplicates. **(E-H)** Key parameters of merozoite invasion were measured for both *Pf*Dd2 WT and *Pf*Dd2 ΔMSP2 parasites that had successfully invaded a RBC using live cell imaging of merozoite invasion. Time to merozoite attachment to RBCs **(E)**, length **(F)** and strength **(G)** of RBC deformation, and time to complete merozoite invasion **(H)** were measured by live microscopy. A minimum of 30 invading merozoites for each line were captured and assessed. Significance determined by unpaired t-test with $p < 0.05$ deemed significant.

Impact of MSP2 gene deletion on gene expression and invasion pathway usage.

Because of the close proximity of the *pfmsp2*, 4 and 5 genes on chromosome 2, similarities in their structure (e.g. GPI-anchor intrinsically disordered protein) and shared merozoite surface localization, we used qPCR to assess whether knock-out of *Pf*3D7 MSP2 impacted the expression levels of *Pf*3D7 MSP4 or MSP5. We found no change in MSP4 and MSP5 expression levels between *Pf*3D7 ΔMSP2 parasites and *Pf*3D7 WT parasites (Figure 4A). As expected, *Pf*MSP2 was not detected in the *Pf*3D7 ΔMSP2 parasites.

Secreted *P. falciparum* antigens belonging to the erythrocyte binding antigen (EBA) and reticulocyte binding homologue (Rh) families are known to facilitate host-cell invasion through specific receptors on the RBC surface. Reliance on specific EBA or Rh proteins, or their cognate host-cell receptor, define different invasion pathways that can be crudely described by the sensitivity of merozoite invasion to cleavage of RBC surface proteins with trypsin, chymotrypsin and neuraminidase (Baum et al., 2005; Duraisingh et al., 2003; Orlandi et al., 1992).When we assessed whether *Pf*MSP2 knock-out impacted merozoite invasion pathway

usage, we found that *Pf*3D7 ΔMSP2 parasites grew equally as well as *Pf*3D7 WT parasites in

all enzyme-treated RBCs (Figure 4B).

To investigate whether *Pf*MSP2 knock-out caused any changes in gene-expression of proteins

that may have a role in merozoite invasion and possibly compensate for loss of *Pf*MSP2, we

undertook RNA sequencing and differential gene expression analysis of *Pf*3D7 ΔMSP2 and

*Pf*3D7 WT parasites. As expected, the *Pf*MSP2 transcript, which was targeted for removal by

our CRISPR-Cas9 strategy, was not detected in the *Pf*3D7 ΔMSP2 parasites between 132 bp

and 819 bp (Supplementary Figure 5). After removal of genes belonging to variant surface

antigen families from the data, eight proteins were found to have a log2fold expression increase

above 1; none of these proteins are annotated as a merozoite surface or secreted protein linked

to merozoite invasion (Figure 4C, Table 2). Five of these eight upregulated proteins are of

unknown function and expressed at schizont stages, and only one (*Pf*3D7_0909100) has a

transmembrane domain that could possibly anchor it to a membrane surface as the GPI anchor

does for *Pf*MSP2 (Table 2). Three proteins were found to have a log2fold expression of <-1,

indicating a significant reduction in expression with *Pf*MSP2 knock-out, with *Pf*MSP2 being

one of these and the other two predicted to be exported proteins (Figure 4C, Table 2). While

these data do not exclude any of the *Plasmodium* proteins with unknown function that are

upregulated in *Pf*3D7 ΔMSP2 parasites as having a role that compensates for the loss of

*Pf*MSP2, only one is predicted to have a transmembrane domain that would bind it to a

membrane and no protein identified as up or down-regulated with *Pf*MSP2 knock-out has

previously been linked to merozoite invasion. Therefore, *Pf*MSP2 knock-out does appear to

have led to changes in gene expression, but does not appear to have a significant impact on the

regulation of known merozoite surface or secreted protein gene-expression that would suggest

loss of *Pf*MSP2 is compensated for by another protein.

Figure 4. The impact of the loss of *Pf*MSP2 on expression of known merozoite invasion genes
and invasion pathway utilization. **(A)** Impact of *Pf*MSP2 KO on transcription during schizonts
was assessed by qPCR for genes located in proximity to *Pfmsp2* on chromosome 2. Changes
in expression between *Pf*3D7 WT and *Pf*3D7 ΔMSP2 parasites was determined by qPCR

relative to *pfaldolase* expression with *pfsub1* serving as a schizont stage control. Graph displays mean ± S.D. of three independent RNA harvests. **(B)** Selective enzymatic cleavage of key RBC receptors showed no difference in invasion preference between *Pf*3D7MSP2 WT and *Pf*3D7 ΔMSP2 parasites. Parasitaemia was determined by flow cytometry and compared to growth in non-treated control RBCs. Graph displays mean ± S.D. of three independent experiments. **(C)** Log2(fold change) for differentially expressed genes, including multigene families, between the transcriptome of *Pf*3D7 WT and *Pf*3D7 ΔMSP2 schizonts. Black lines differentiate genes with a log2 (fold change) > 0.5 and < -0.5 with adjusted p-value < 0.05. Four independent schizont RNA harvests were performed and analysed. Significance determined by t-test with p< 0.05 deemed significant.

Impact of *Pf*MSP2 disruption on antibodies and inhibitors targeting secreted surface exposed antigens

Merozoites are key targets of antibodies during malaria infection with antibodies either able to directly block protein function and thus merozoite invasion, or recruit effectors such as complement leading to the destruction of the merozoite (Beeson et al., 2016). Given the prominence of *Pf*MSP2 on the merozoite surface we asked the question; how does the loss of *Pf*MSP2 affect antibody efficacy against merozoites?

*Pf*EBAs and *Pf*Rhs moderate early steps in invasion with a degree of redundancy between these proteins. Several EBAs and Rhs can be targeted with invasion inhibitory antibodies and are targets of acquired human antibodies (Persson et al., 2013, 2008; Richards et al., 2013). We found no significant difference in the ability of purified rabbit immunoglobulin raised against single (*Pf*Rh2b) or combinations of *Pf*EBAs and *Pf*Rhs (*Pf*EBA175/*Pf*Rh2b, *Pf*EBA175/*Pf*Rh4, *Pf*EBA175/*Pf*Rh2b/*Pf*Rh4) to inhibit growth of *Pf*3D7 WT or *Pf*3D7 ΔMSP2 parasites (Figure 5A). Similarly, knock-out of MSP2 in *Pf*Dd2 did not significantly change the parasites sensitivity to rabbit antibodies raised against isogeneic W2mef EBA175 (Figure 5B). Broadly speaking, these results mirror what was seen with selective cleavage of the RBC ligands of *Pf*EBAs and *Pf*Rhs using enzyme treatment of RBCs (Figure 4B), with no evidence that loss of *Pf*MSP2 changed either the importance of these secreted merozoite antigens during invasion or their sensitivity to antibodies targeting different antigens.

The interaction of *Pf*Rh5 with RBC surface receptor basigin is a critical, non-redundant interaction for *P. falciparum* merozoite invasion and a known target of direct invasion-

inhibitory antibodies (Alanine et al., 2019; Crosnier et al., 2011; Douglas et al., 2011). PfRh5

was the initial member of the PCRCR complex to be described, with subsequent studies

demonstrating that antibodies against other members of this complex could also block

merozoite invasion of the RBC (Scally et al., 2022). Given the importance of these secreted

antigens to current vaccine development efforts, we next investigated whether loss of PfMSP2

impacted on the parasite's sensitivity to antibodies targeting these antigens. There was no

difference in sensitivity between Pf3D7 WT and Pf3D7 ΔMSP2 parasites when treated with

invasion-inhibitory rabbit antibodies targeting PfRh5 (Figure 5C) or nanobodies targeting

PCRCR complex component PfPTRAMP (H8) (Figure 5D). Nanobodies targeting the PCRCR

complex component CSS (D2) showed a slight, but non-significant increase in invasion-

inhibitory activity in the absence of PfMSP2 (1.4-fold increase; IC_{50} value for Pf3D7 WT is 57

μg/mL; IC_{50} value for Pf3D7 ΔMSP2 is 42 μg/mL; p=0.3; Figure 5E). However, the Pf3D7

ΔMSP2 specific increase in invasion-inhibitory activity was amplified with the addition of the

human Fc domain to the anti-CSS nanobody (D2-Fc) to be 3-fold greater for Pf3D7 ΔMSP2

parasites (IC_{50} Pf3D7 WT 138 μg/mL; IC_{50} Pf3D7 ΔMSP2 46 μg/mL; p=0.0001; Figure 5F).

The glycosaminoglycan heparin is a specific inhibitor of invasion and has been observed to

bind to the 42 kDa C-terminal region of PfMSP1 the most abundant GPI-anchored protein on

the merozoite surface, marking this as a possible mechanism for its invasion inhibitory activity

(Boyle et al., 2010a). However, the highly charged state of heparin means that it could block

invasion through binding additional proteins. When tested against Pf3D7 WT and Pf3D7

ΔMSP2 parasites, we observed a small increase in inhibitory potency of heparin for parasites

lacking PfMSP2 (1.4-fold; IC_{50} Pf3D7 WT 0.65 IU/mL; IC_{50} Pf3D7 ΔMSP2 0.47 IU/mL;

p=0.0003; Figure 5G). We next tested rabbit antibodies raised against the C-terminal 19 kDa

fragment of PfMSP1. We found a small but non-significant trend towards increased inhibition

of growth in parasites that lacked PfMSP2 across repeat experiments (Figure 5H). These data

suggest that loss of PfMSP2 can impact on the invasion-inhibitory potency of some antibodies

that target secreted or surface antigens.

Figure 5. Impact of *Pf*MSP2 removal on efficacy of antibodies targeting other merozoite

surface-exposed antigens. Changes in antibody efficacy in the absence of *Pf*MSP2 was

assessed by measuring changes in antibody invasion inhibition and subsequent growth

compared to growth in the absence of antibody for both *P. falciparum* WT and ΔMSP2

parasites over 2 cycles. **(A)** Rabbit (Rb) IgG raised against merozoite antigens of the *Pf*3D7

EBA/Rh family. **(B)** Rabbit sera raised against *Pf*Dd2 EBA175. **(C)** Rabbit IgG raised against

*Pf*3D7 Rh5. **(D)** Nanobody (nAb) to *Pf*3D7 PTRAMP. **(E-F)** Nanobody and Fc-tagged

nanobody to *Pf*3D7 CSS. **(G)** The invasion inhibitory glycosaminoglycan heparin. **(H)** Rabbit

sera raised against *Pf*3D7 MSP1-19 (different vaccinated rabbit sera identified by numbers).

Graph displays mean ± S.D. of three different experiments. Significance was determined by

unpaired t-test when only a single concentration point was tested and for IC_{50} comparisons an

extra Sum-of-Squares F Test (best-fit $LogIC_{50}$) was performed with p< 0.05 deemed

significant.

Loss of *Pf*MSP2 consistently potentiates AMA1 invasion inhibitory antibodies

Subsequent to *Pf*Rh5 -basigin binding in the steps of merozoite invasion is the formation of the

tight-junction between *Pf*AMA1, secreted from the micronemes onto the merozoite surface,

and the rhoptry neck protein *Pf*RON2, which is inserted into the RBC membrane (Lamarque

et al., 2011; Srinivasan et al., 2011). In lines where *Pf*MSP2 was disrupted, we found a

consistent potentiation of antibodies that had anti-*Pf*AMA1 invasion-inhibitory activity.

Invasion-inhibitory polyclonal rabbit serum antibodies and purified immunoglobulin raised

against *Pf*AMA1 of *Pf*3D7 and *Pf*W2mef (isogenic to *Pf*Dd2), which express different alleles

of both *Pf*MSP2 and *Pf*AMA1, were found to be more inhibitory with *Pf*MSP2 knock-out

(Figure 6A and B). The polyclonal antibody with the greatest potentiation of inhibitory activity

was an IgG preparation purified from a rabbit antiserum raised against *Pf*W2mef AMA1. This

had a 3-fold increased potency against *Pf*Dd2 ΔMSP2 compared to *Pf*Dd2 WT (IC_{50} *Pf*Dd2

WT 0.015 mg/mL; IC_{50} *Pf*Dd2 ΔMSP2 0.005 mg/mL; p<0.0001) over two cycles of growth

(Figure 6B, C).

The mouse 1F9 (binds to the RON2 binding pocket and polymorphic loop 1d (Coley et al.,

2007)) and rat 4G2 (binds to a conserved epitope in the ectodomain (Kocken et al., 1998))

monoclonal antibodies have been shown to target different regions of *Pf*AMA1 and exhibit

selective inhibition against Pf3D7 AMA1. Here we found potentiation of invasion inhibitory activity with Pf3D7 MSP2 knock-out for both 1F9 (2.1-fold; IC$_{50}$ Pf3D7 WT 0.27 mg/mL; IC$_{50}$ Pf3D7 ΔMSP2 0.13 mg/mL; p<0.0001; Figure 6D) and 4G2 (3.2-fold; IC$_{50}$ Pf3D7 WT 1.15 mg/mL; IC$_{50}$ Pf3D7 ΔMSP2 0.36 mg/mL; p<0.0001; Figure 6E).

We also tested the invasion inhibitory effect of i-bodies, which are smaller single-domain antibody-like molecules inspired from the shark variable new antigen receptor (V$_{NAR}$) (Griffiths et al., 2018, 2016). When we tested the i-body WD34 (Angage et al., 2024) which binds a highly conserved epitope that includes the PfRON2-binding pocket on PfAMA1, we observed a small potentiation of PfAMA1 specific activity with knock-out of PfMSP2 in Pf3D7 (1.3-fold; IC$_{50}$ Pf3D7 WT 0.012 mg/mL; IC$_{50}$ Pf3D7 ΔMSP2 0.009 mg/mL; p=0.08 Figure 6F). Given WD34 was also found to be inhibitory to growth of W2mef parasites (Angage et al., 2024) we also tested its activity against our PfDd2 ΔMSP2 parasites and found potentiation of invasion inhibitory activity with loss of MSP2 in PfDd2 parasites (2-fold; IC$_{50}$ PfDd2 WT 0.016 mg/mL; IC$_{50}$ PfDd2 ΔMSP2 0.008 mg/mL; p=0.004 Figure 6G). As was observed for the anti-CSS nanobody, addition of a human Fc domain to WD34 amplified the inhibitory phenotype, with WD34 Fc having a 2.3-fold greater inhibitory activity against Pf3D7 ΔMSP2 than against Pf3D7 WT (IC$_{50}$ Pf3D7 WT 0.1 mg/mL; IC$_{50}$ Pf3D7 ΔMSP2 0.04 mg/mL; p=0.0004; Figure 6H). A second i-body, WD33 (Angage et al., 2024), which binds AMA1 between domain II and domain III, had very limited invasion inhibitory activity against Pf3D7 parasites and did not show improved potency with knock-out of Pf3D7 MSP2 (0.9-fold; IC$_{50}$ Pf3D7 WT 1.02 mg/mL; IC$_{50}$ Pf3D7 ΔMSP2 1.1 mg/mL; p=0.8; Figure 6I). Although the limited inhibitory activity for WD33 tagged with the Fc receptor prevented us determining an IC$_{50}$ at the concentrations feasible to test, there was no increased inhibition of Pf3D7 ΔMSP2 parasites compared to the low level seen with Pf3D7 WT parasites (data not shown). These data suggest that the potentiation of PfAMA1 targeted antibody inhibition with MSP2 knock-out is PfAMA1 epitope dependent.

To account for the differences in i-body sizes, we compared the WD34 and WD34 Fc i-bodies on a molar basis and found that they had similar activities against Pf3D7 MSP2 WT parasites (IC$_{50}$ 0.94 μM and 1 μM respectively) indicating that the Fc-tag itself did not modify epitope binding properties. Rather, it is the absence of PfMSP2 that results in increased inhibition of AMA1 function by WD34, and this is further potentiated by antibody size.

In light of these results, we hypothesised that removal of *Pf*MSP2 may improve antibody access

to other antigens that are targets of inhibitory antibodies. To assess whether this was the case,

we first tested whether increased *Pf*AMA1 polyclonal antibody binding to the surface of

merozoites could be observed for *Pf*3D7 ΔMSP2 compared to *Pf*3D7 MSP2 WT parasites using

a fluorescently tagged WD34 i-body (WD34-mCherry) for quantitative immunofluorescence

assays of late stage schizonts labelled with a single incubation and wash step. We observed

WD34-mCherry to have a significantly higher mean fluorescence intensity for *Pf*3D7 ΔMSP2

compared to *Pf*3D7 WT (Figure 7A), supporting that the *Pf*AMA1 i-bodies may have better

access in the absence of *Pf*MSP2. As a control, we tested labelling using a fluorescently tagged

WD33 i-body which binds to AMA1 (Angage et al., 2024), but we found did not have improved

inhibitory activity in the absence of MSP2 (Figure 6I). Mirroring the results of the growth

assay, there was no difference observed for WD33-eGFP binding fluorescence for *Pf*3D7

ΔMSP2 compared to *Pf*3D7 WT parasites (Figure 7B).

We next explored whether the addition of recombinant *Pf*MSP2 could impact on the inhibitory

antibody binding on-rate against substrate-bound *Pf*AMA1 recombinant protein using Surface

Plasmon Resonance (SPR). For the anti-*Pf*AMA1 targeting mAb 4G2, there was a non-

significant trend toward a lower on-rate in the presence of recombinant *Pf*MSP2 (Figure 7C).

For the i-body WD34-Fc, there was minimal difference in detectable *Pf*AMA1 binding to

*Pf*AMA1 in the presence or absence of *Pf*MSP2 (Figure 7C). Using a similar principle of

testing antibody binding to substrate-bound *Pf*AMA1 protein in the presence, or absence of

increasing concentrations of *Pf*MSP2, we assessed changes in levels of anti-*Pf*AMA1 antibody

binding using an enzyme linked immunosorbent assay (ELISA). With increasing

concentrations of *Pf*MSP2 added to wells coated with AMA1, we found decreased binding of

anti-*Pf*AMA1 mAb 4G2 and i-body WD34-Fc by ELISA (Figure 7D). Interestingly, *Pf*MSP2

concentrations ≤0.31 μg/mL, an antigen concentration below the level that could be detected

by an MSP2 mAb, were enough to impact on the binding of *Pf*AMA1 antibodies. By looking

at the impacts of removing *Pf*MSP2 from the merozoite surface, or adding this protein in to

recombinant assays, these data provide additional support that inhibitory antibodies targeting

*Pf*AMA1 bind at a higher rate in the absence of *Pf*MSP2.

Figure 6. Absence of *Pf*MSP2 from the merozoite surface impacts invasion inhibition by

*Pf*AMA1 antibodies. *Pf*3D7 (**A, D-F, H, I**) and *Pf*Dd2 (**B, C, G**) express different *Pf*MSP2

alleles and different *Pf*AMA1 alleles, yet both showed altered anti-AMA1 antibody growth

inhibition in the absence of *Pf*MSP2. The effect was seen with serum (**A-B**; different

vaccinated rabbit (Rb) sera identified by numbers), purified rabbit and mouse monoclonal

(mAb) antibodies **(C-E)** and i-bodies (ibA) **(F-J)**. Final parasitaemia was determined by flow

cytometry and compared to control. Graph displays mean ± S.D. of three or four independent

experiments. Significance was determined by unpaired t-test when only a single concentration

point was tested and for IC_{50} comparisons an extra Sum-of-Squares F Test (best-fit $LogIC_{50}$)

was performed with $p < 0.05$ deemed significant.

Figure 7. Quantitative fluorescence microscopy to assess whether differential binding may explain the increased potency of anti-*Pf*AMA1 invasion-inhibitory antibodies in the absence of *Pf*MSP2. Fluorescence intensity of fluorescently tagged anti-*Pf*AMA1 i-body (ibA) WD34-mCherry **(A)** and WD33-eGFP **(B)** for both *Pf*3D7 WT and *Pf*3D7 ΔMSP2, with a representative image for i-body WD34-mCherry. Nucleus in blue, i-body signal in red. Scale bar = 4 μm. Two independent experiments were performed with significance determined by unpaired t-test with p< 0.05 deemed significant. The lower overall mCherry signal required a higher antibody concentration (240 ng/mL) to have a comparable intensity measure to the eGFP tagged antibody (120 ng/mL) for *Pf*3D7 WT merozoites. **(C)** Read-out of the Surface plasmon resonance (SPR) antibody on-rate (association constant) for anti-*Pf*AMA1 mAb 4G2

and i-body WD34-Fc binding to *Pf*AMA1 in the presence or absence of *Pf*MSP2 protein.

Data represents the mean of 3 experiments with significance determined by unpaired t-test

with p< 0.05 deemed significant. **(D)** ELISA based assessment of the anti-*Pf*AMA1 mAb

4G2 and i-body WD34-Fc antibody binding levels to recombinant *Pf*3D7 AMA1 in the

presence or absence of recombinant *Pf*3D7 MSP2. PBS control demonstrates background

fluorescence. Dashed orange line provides a guide for peak absorbance levels. Anti-*Pf*MSP2

mAb shows increasing concentrations of *Pf*MSP2 protein results in decreased binding of

mAb 4G2 and i-body WD34-Fc. Data represents the mean of 3 experiments and error bars

are ±S.D.

Discussion

Despite decades of research and interest, the function of most merozoite surface proteins

remains unclear and targeting merozoite antigens in vaccine development has not achieved

high efficacy in clinical trials. Additionally, it has generally proven difficult to generate highly

inhibitory antibodies through vaccination with merozoite antigens. *Pf*MSP2 is one of the most

abundant proteins on the merozoite surface, has shown promise as a vaccine candidate and is

an established target of protective cytophilic antibodies (Boyle et al., 2015; Genton et al., 2002;

McCarthy et al., 2011; Osier et al., 2014, 2010). However, its function remains unclear. Our

findings here suggest that *Pf*MSP2 does not have an essential role in invasion *in vitro*, but can

modulate the sensitivity of merozoites to inhibitory antibodies targeting other antigens.

Despite its abundance on the merozoite surface and previous work suggesting a role in RBC

invasion, we found merozoites invade normally in the absence of this protein after gene

deletion. This does not exclude a role for *Pf*MSP2 *in vivo* where there are additional pressures,

such as immune-effector mechanisms and flow dynamics, on merozoite invasion. However,

given we have knocked-out *Pf*MSP2 from two different *P. falciparum* isolates, our findings do

not currently support a major role for *Pf*MSP2 in the mechanics of merozoite invasion. Thus,

it appears that the function of the two most abundant proteins on the merozoite surface,

*Pf*MSP1 (Das et al., 2015; Kals et al., 2024) and *Pf*MSP2, are not directly linked to merozoite

binding to the RBC and subsequent invasion. Deletion of MSP2 did lead to significant changes

in the expression of several genes, but there were no proteins with a clear link to merozoite

invasion and the biological relevance of these changes is not currently understood and requires

future investigation.

Both rabbit polyclonal and mouse monoclonal anti-*Pf*AMA1 antibodies tested were

consistently more inhibitory in the absence of *Pf*MSP2. Of the two *Pf*AMA1 targeting i-bodies

tested (Angage et al., 2024), the most potent WD34 was also more potent in the absence of

*Pf*MSP2, with this activity increasing significantly with the ~5.6-fold increase in size that

occurred upon addition of a mouse Fc region. Similarly, activity of a nanobody targeting the

PCRCR complex protein CSS showed increased potency in the absence of *Pf*MSP2, and again

this was increased further with antibody enlargement through addition of an Fc region. These

data suggest that *Pf*MSP2 may act to dampen the inhibitory activity of antibodies targeting

other antigens on the surface of the merozoite. Such immune dampening (called

conformational masking) by intrinsically disordered proteins, or protein regions, of host-cell

invasion related proteins has recently been demonstrated by modelling and gene-editing studies

in the viral pathogens HIV and HCV, respectively (Li et al., 2024; Stejskal et al., 2022). These

studies, along with our demonstration of increase antibody inhibitory efficacy with loss of the

intrinsically disordered *Pf*MSP2, support that this may be a widespread approach used by

pathogens to protect important protein functional sites from inhibitory antibodies.

A role in immune evasion has previously been suggested for *Pf*MSP2 and speculatively an

alternate explanation for the sensitisation observed is that this protein may shield key *Pf*AMA1

epitopes from antibody inhibition. For the *P. falciparum* 3D7 line, we found that *Pf*MSP2

knock-out led to higher *Pf*AMA1 binding as measured by increased fluorescence intensity for

the mCherry tagged i-body WD34 in the absence of *Pf*MSP2. Using a solid-phase model and

titrating recombinant *Pf*MSP2, we found that increasing concentrations of *Pf*MSP2 reduced

binding of an anti-*Pf*AMA1 mAb and i-body, even at concentrations where *Pf*MSP2 was not

detectable itself by antibodies. These data support that the presence of *Pf*MSP2 may reduce

antibody access (masking) to important invasion proteins exposed on the merozoites surface.

However, we acknowledge that the antibody-binding assays used (immunofluorescence assays,

SPR, ELISA) might not be sensitive enough to observe all potential changes in binding,

temporal changes when binding epitopes are exposed or differences in avidity that could lead

to increased antibody potency during the course of parasite invasion and growth, and these

factors may also contribute. Future studies of the merozoite surface and interactions between

proteins may help decipher in full how *Pf*MSP2 is protecting essential merozoite antigens from

invasion inhibitory antibody activity and reveal strategies for enhanced vaccine design.

Our observation that loss of *Pf*MSP2 potentiates the invasion-inhibitory activity of antibodies targeting other antigens opens an additional avenue for vaccine development, such as targeting *Pf*MSP2 to interfere with its potential role in immune evasion and thus potentiating antibodies against other merozoite antigens. Additionally, our results suggest that vaccines based on *Pf*AMA1 may need to be designed to specifically avoid the effect of *Pf*MSP2 modulating inhibitory antibodies, such as focussing on specific epitopes or structural features. To date, vaccines based on AMA1 have failed to achieve substantial efficacy (Sagara et al., 2009; Sheehy et al., 2012; Thera et al., 2011). We can speculate that one possible protective mechanism of action of the Combination B vaccine (Genton et al., 2002; McCarthy et al., 2011), which combined *Pf*MSP2, *Pf*MSP1 and *Pf*RESA, could be that the inhibitory activity of vaccine-induced antibodies raised against *Pf*MSP2 were then able to potentiate the activity of antibodies targeting other antigens, potentially *Pf*MSP1 which was also a target of the vaccine. Further investigation using the parasite lines developed in this study and a wider panel of antibodies could shed more light on this potentially novel mechanism of vaccine derived antibody efficacy.

The results of this study and work done by Escalante et al. (2022) demonstrate that MSP2-like sequences are found in the avian malarias *P. galinaceum* and *P. relictum*, which differs from the fields previous understanding that *Pf*MSP2 arose in the *Laverania* lineage of malaria parasites. This observation indicates that MSP2 was likely present in the ancestral malaria parasite before being lost in primate and rodent *Plasmodium* spp. Further insights into the evolution and loss of *Pf*MSP2 are likely to be found with additional whole genome sequences of bird and lizard malaria species. The presence of *msp2*-like sequences in some avian *Plasmodium* species and *Laverania* suggest a retention of *msp2* sequences in the *Laverania* which indicates an important role for MSP2 for these parasites. Indeed, our amino acid and structural prediction level comparison demonstrates that much of the protein properties conserved in the *Laverania* MSP2s are also recognisable, and in many cases conserved, in the avian malaria MSP2s, despite 10 million years of evolutionary divergence. This suggests that having an MSP2-like protein is beneficial for these parasites with very different hosts. However, CRISPR-Cas9 gene editing used in this work has shown that, in contrast to previous work, *Pf*MSP2 is not essential for *P. falciparum* blood stage parasite growth *in vitro*. Our findings raise the possibility that the retention of *Pf*MSP2 in the *Laverania*, and by association potentially the avian malaria parasites, may be linked to its apparent capacity to modulate the impact of antibodies against other merozoite surface-exposed antigens, rather than through a

direct mechanistic role in host-cell invasion. If MSP2 was to fulfill such a role in the *Laverania*

and avian malaria parasites, it raises the question of whether a similar protective system exists

in other *Plasmodium* spp. and what the key protein/s are that fulfil this role in these parasites.

The *Pf*3D7 MSP2 and *Pf*Dd2 MSP2 lines developed in this study will provide useful reagents

for investigating the potential of targeting a single, or both, *Pf*MSP2 allelic types with vaccines

that elicit different immune responses. These parasites could also be used to explore whether

vaccine-induced antibodies against other merozoite antigens could also be potentiated if any

immune evasion mechanism provided by *Pf*MSP2 is removed. This may be achievable with a

combination vaccine or through targeting an epitope of another merozoite vaccine candidate

that is not protected by *Pf*MSP2. Such information, investigated using the *Pf*MSP2 knock-out

parasites developed in this study, could be useful in prioritising vaccine candidates, specific

epitopes and combinations that target the merozoite.

Conclusion

Advancements in gene-editing techniques in *P. falciparum* have allowed us to show for the

first time, and contrary to previous work, that *Pf*MSP2 is not essential for *P. falciparum* growth

in vitro. Instead, we present a new concept that MSP2 can modulate the activity of invasion-

inhibitory antibodies and may have evolved for this purpose. It is possible that other merozoite

surface proteins may have similar roles and should be investigated in future studies. Our

observation that loss of *Pf*MSP2 potentiates the inhibitory activity of antibodies targeting other

merozoite surface exposed proteins reveals a new avenue to explore in vaccine development

by blocking or bypassing this function to improve the activity of vaccines targeting these

antigens.

Materials and Methods

Bioinformatic analysis of *Pf*MSP2, *Pf*MSP4 and *Pf*MSP5 like proteins in *Plasmodium* spp.

The PlasmoDB database (Aurrecoechea et al., 2009) was used to identify in the available

Plasmodium genomes the region between Adenylosuccinate lyase (PF3D7_0206700) and a

conserved protein of unknown function (PF3D7_0207100) where *Pf*MSP2, *Pf*MSP4 and

*Pf*MSP5 are localised in *P. falciparum*. All annotated MSP2, MSP4 and MSP5 protein

sequences were downloaded. Translated sequences of unannotated genes and putative

annotations in this region were also downloaded and aligned using the MUSCLE alignment

tool MEGA11: Molecular Evolutionary Genetics Analysis version 11 (Tamura et al., 2021) or

the Geneious Prime 2019.2.1 MUSCLE alignment tool (Biomatters Ltd.) and determined to be

MSP2, MSP4 or MSP5 based on sequence similarity in the conserved regions of these genes.

A BLAST (NCBI) search with the N-terminal and C-terminal conserved regions of MSP2 was

also performed to capture any sequences missed by the above approach. A maximum likelihood

phylogenetic tree of identified MSP2 sequences was generated using the default maximum

likelihood settings of MEGA11 with 500 bootstraps and the *P. gallinaceum* MSP2 as an

outgroup.

Modelling of MSP structure across Plasmodium spp. using Alpha-fold

In order to obtain models of the 3-dimensional protein structures of MSP2 for *P. falciparum*,

P. reichenowi, *P. billcollinsi*, *P. adleri*, *P. relictum* and *P. gallinaceum*, we first removed the

predicted N-terminal signal sequences and C-terminal GPI anchor sequence. The resulting

sequences were submitted to AlphaFold2 (Jumper et al., 2021). The resulting models were

subsequently analysed and visualised using PyMOL 2.4.0 (Schrödinger, USA).

Culture and synchronisation of *P. falciparum*

Plasmodium falciparum 3D7 and Dd2 was cultured in RPMI-HEPES media (Gibco) containing

0.5% Albumax ll (Gibco), 25 μM Gentamicin (Gibco), 367 μM hypoxanthine (Sigma Aldrich),

2 mM L-Glutamax (Gibco), 0.17% sodium bicarbonate (Sigma Aldrich) and O positive human

red blood cells (Lifeblood, Australia) at 3% parasitemia, 3% haematocrit. Cultures were grown

at 37°C under 1% oxygen, 5% CO_2 (BOC gas), in a humidified chamber. For gene-edited lines,

WR99210 (Jakobus Pharmaceuticals) was added to culture media. Sorbitol lysis (5%) (Sigma

Aldrich) to select for ring-stage parasites, Percoll Plus (Sigma Aldrich) selection of late-stage

parasites and heparin treatment (Pfizer) (Boyle et al., 2010b) were used to maintain

synchronicity of parasite cultures.

Generation of *Pf*MSP2 knockout line using CRISPR Cas9

Briefly, a flank region incorporating the 5' region upstream of *Pf*MSP2, the first 44 amino

acids of the protein, and a flank of the region immediately 3' of *Pf*MSP2 was inserted into the

pCC1 plasmid by restriction enzyme cloning. A 20 bp guide sequence was designed using EuPaGDT (Peng and Tarleton, 2015), annealed and inserted in the pUF_Cas9 plasmid by infusion cloning (Takara Bio). pUF_Cas9 guide plasmid (20 μg) and pCC1 *Pf*MSP2 repair plasmid (60 μg) was transfected directly into *P. falciparum* schizonts. Early schizonts were purified using a 70% percoll gradient and treated for 4 hrs with compound 1 (Taylor et al., 2010) at 2 μM. After incubation treated schizonts were washed twice and left shaking for 20 minutes before pelleting and resuspension in Complete Cytomix (plasmids plus 0.895% KCl, 0.0017% $CaCl_2$, 0.076% EGTA, 0.102% $MgCl_2$, 0.0871% K_2HPO_4, 0.068% KH_2PO_4, 0.708% HEPES). Schizonts were electroporated in 0.2 cm cuvettes (Biorad) at 800 V and 25 μF. Electroporated schizonts were mixed with warmed media and fresh RBCs and shaken to promote invasion for 20 minutes after which they were placed in a culture dish. Transfected parasites were then treated with WR99210 to select for integration of the hDHFR drug selection cassette and MSP2 knock-out.

To confirm integration gDNA was collected by saponin lysis of schizont stage parasite culture and gDNA extracted using the PureLink Genomic DNA Mini Kit (Invitrogen) as per manufacturer's instructions. Integration of the *Pf*MSP2 knockout construct into a portion of parasites was confirmed by PCR (Supplementary Table 1). Once confirmed cultures were cycled on and off WR99210 (Jakobus Pharmaceuticals) and 1 μM 5FC (Sigma Aldrich) to select for integrated parasites no longer expressing the guide plasmid. Limited dilution cloning was then performed to obtain parasite clones which had integrated the knockout construct.

Western blotting to detect *Pf*MSP2

Synchronised schizonts were harvested and the RBCs lysed by saponin. Briefly ~10 mL schizont (38-44 hrs) culture was lysed on ice for 10 minutes with 0.15% w/v saponin before pelleting by centrifugation and washing once in 0.075% w/v saponin followed by three washes in PBS containing protease inhibitors (CØmplete, Roche). Saponin treated schizont pellets were treated with DNAseI (Qiagen) for 10 minutes at room temperature before resuspension in reducing sample buffer (0.125 M Tris-HCL, 4% SDS, 20% glycerol, 10% beta-mercaptoethanol and 0.002% bromophenol blue). Proteins were separated by size using SDS-PAGE 4-12% Bis-Tris Gels (Bolt, Invitrogen) at 110V for 80 minutes before transfer to a nitrocellulose membrane (iBlot, Invitrogen) at 20V for 7 minutes. Blots were blocked from 1 hr to overnight in 3% skim milk PBS (Sigma Aldrich) before incubation with primary

antibodies (1/75000 mouse 2F2 anti-*Pf*MSP2 3D7 (Supplementary Table 2; (Adda et al.,
2012)) and 1/75000 rabbit anti-*Plasmodium* aldolase (Abcam) or 1/75000 rabbit anti-FC27 and
1/10000 mouse anti- EXP2 (a gift of Paul Gilson, Burnet Institute, Melbourne) for 1- 2 hrs.
IRDye 800CW goat anti-mouse (1/4000, LI-COR Biosciences) or IRDye 680RD goat anti-
rabbit (1/4000, LI-COR Biosciences) secondary antibodies were used for detection on the
Odyssey Infrared imaging system (LI-COR Biosciences). Quantification was performed using
Image Studio Lite 5.2.5 (LI-COR Biosciences).

Quantification of parasite expansion rate and invasion inhibition

Potential growth defects resulting from gene editing was assessed by flow cytometry. The
initial parasitemia of cultures was determined by flow cytometry and then measured again after
either 48 hrs or 96 hrs of growth. To determine parasitemia by flow cytometry Ethidium
Bromide (10 μg/mL BioRad) was added to 50 μL of 1% haematocrit culture for 30 mins to
allow for staining of parasite DNA. The stain was washed off and wells resuspended in 200μL
of PBS. Data was collected on a BD Accuri C6 Plus Flow Cytometer (BD Biosciences) and
analysed on FlowJo (FlowJo LLC). Briefly, forward scatter and side scatter was used to
determine the RBC population. From this population infected RBCs were gated on as highly
fluorescent in the PE channel. Final parasitemia was compared back to initial parasitemia to
determine the fold increase in parasitemia for each line.

To examine the invasion inhibitory effect of antibodies (Supplementary Table 2), 5 μL of
potential inhibitor was mixed with 45 μL of 0.2% schizont parasitemia at 1% haematocrit for
96 hrs before end parasitemia was determined by flow cytometry. Data is displayed as %
Growth of media only controls for each line.

To examine differences in RBC receptor preference RBCs were treated with different enzymes
that cleave different residues of RBC receptors which is used as a marker for which RBC
receptor/s are preferred by *P. falciparum* (Duraisingh et al., 2003). Infected RBCs at 1% ring
stage parasitemia were washed three times in RPMI-HEPES to remove excess protein. Packed
RBCs (20 μL) at 1% parasitemia were then resuspended with 5X RBC volume of pre-warmed
enzymes at desired concentration (Supplementary Table 3). RBC-enzyme mix was incubated
at 37°C for 45 mins with rotation to keep RBCs resuspended. Following incubation, RBCs
were washed three times and added to fresh complete media at 1% haematocrit in a U- bottom
96 well plate in technical duplicate. Plates were incubated for 72 hrs, sufficient time for *P.
falciparum* to complete one full cycle of growth and reach trophozoite stage again. Final

parasitemia was determined by flow cytometry, as detailed above, and compared to non-treated

controls. IC_{50} calculations data was log transformed then a nonliner regression log-(inhibitor)-

versus-response curve was calculated with Extra Sum-of-Squares F Test (best-fit $LogIC_{50}$)

used to compared IC_{50}s between conditions.

Immunofluorescence assay of Wildtype and *Pf*MSP2 KO parasites

Schizonts ~38 hrs old were treated with E64 (Sigma-Aldrich) for 5 hrs to allow development

of very mature schizonts desired for imaging. After E64 treatment, parasite cultures were fixed

in 4% v/v paraformaledehyde (PFA, Sigma-Aldrich), 0.0075% v/v glutaraldehyde (pH 7.5,

Electron Microscopy Sciences) solution for 30 min at room temperature shaking gently. Fixed

parasites were washed in 1X PBS and then resuspended in PBS at 1% haematocrit. Coverslips

(#1.5H high-precision coverslips, Carl Zeiss, Oberkochen, Germany) were coated in 0.01%

Poly-L-lysine (Sigma Aldrich) for 1 hr at room temperature, washed in miliQ water before the

fixed parasite culture was allowed to adhere for 1 hr. Cells were permeabilised by 0.1% Triton

X-100 PBS for 10 mins before blocking for 1 hr to overnight in 3% BSA 0.05% Tween 20

PBS. Primary antibodies were diluted in 1% BSA 0.05% Tween 20 PBS and incubated for 2

hrs. Coverslips were washed in 0.05% Tween 20 PBS three times before incubation with the

secondary antibody (1/500 goat anti-chicken/mouse/rabbit Alexa Fluor coupled secondary

antibodies -488 nm, 594 nm, 647 nm; Life Technologies) for 1 hr. After the secondary antibody

was washed off coverslips underwent a secondary fix in 4% v/v PFA, 0.0075% v/v

glutaraldehyde for 5 mins. The fixative was washed off and the coverslips dehydrated with

sequential 3 min ethanol treatment (70%, 90% and 100%). Coverslips were allowed to dry and

mounted with ProLong Gold antifade solution (refractive index 1.4) which contains 4', 6-

diamidino-2phenylindole, dihydrochloride (DAPI) (ThermoFisher Scientific) and allowed to

set overnight. Imaging was performed on the Zeiss LSM 800 using the Airyscan super-

resolution mode (Carl Zeiss, Obekochen, Germany).

Live cell microscopic analysis of merozoite RBC invasion

Highly synchronised *Pf*Dd2 and *Pf*Dd2 ΔMSP2 parasites at schizont stages were adjusted to

0.25% Haematocrit in complete culture medium. A volume of 200 μL was added to a well of

an iBidi 15 μ-Slide 8 well, glass bottom slide (iBidi 80827) and the slide was promptly returned

to sit on a prewarmed water block within a gassed box and left to incubate at 37°C to allow

iRBCs to settle. The slide was then transported directly to a prewarmed Nikon TiE microscope

with an environmental chamber heated to 37°C and gas mixture of 1% O_2, 5% CO_2 and 94%

nitrogen. Filming was conducted using either a 60X water or 100X oil objective. Differential

interference contrast (DIC) imaging was captured at 3 – 3.5 Volts with a camera exposure of

60 – 180 milliseconds.

Captured footage was processed using the Nikon analysis software (NIS-Elements AR

Analysis version 5.21.01) and analysis of the schizont rupture and merozoite invasion events

was performed manually. The merozoites which invaded successfully were tracked and

attachment time, reorientation time (if obvious), deformation start and end times, the

deformation score (Weiss et al., 2015; Wilson et al., 2015), invasion initiation and completion

times and echinocytosis start times were recorded.

RNA extraction

Highly synchronised cultures were pelleted, resuspended in TriZol (Invitrogen) and incubated

for 5 minutes at 37°C. Chloroform (Sigma Aldrich) was added, mixed and then spun at 12,000g

for 30 minutes at 4°C. Supernatant was collected and mixed with an equal volume of 70%

Ethanol and processed using the RNeasy Kit (Qiagen) to extract purified RNA. RNA was

DNAse l (Qiagen) treated for 30 mins at room temperature and cleaned up on RNeasy columns

according to the manufacturers protocol. Absence of gDNA was checked by qPCR of gDNA

with primers (*Pf*SUB1) and DNAse I treatment repeated if necessary.

qPCR of gene expression in schizonts

DNAse treated RNA was added to 40 mM dNTPs (Qiagen), 0.4 mg/mL random hexamer

(Qiagen) and incubated at 65°C for 5 mins. After incubation 5x Superscript Buffer (Invitrogen),

100 mM DTT (Invitrogen), 40 U/µL RNAseOUT (Invitrogen) and 200 µ/mL Superscript lll

Reverse Transcriptase (Invitrogen) was added and cycled 25°C 5 mins, 50°C for 60 mins and

70°C for 15 mins. Primers (Supplementary Table 4) were designed to detect *Pf*MSP2, *Pf*MSP4,

*Pf*MSP5, *Pf*SUB1 and Fructose Bisphosphate Aldolase as a reference gene (Salanti et al.,

2003). qPCR was performed on cDNA from WT and knockout parasites with PowerUp SYBR

Green Master Mix (Applied Biosystems) and cycling parameters; 95°C for 10 mins followed

by 40 cycles 95°C for 15 mins, 60°C for 14 mins before a dissociation step at 95°C for 2 mins

followed by 60°C for 2 mins with a 2% ramp to 95°C for 2 mins on the QuantStudio 7 Flex

System (ThermoFisher). qPCR was performed in triplicate for each of the three independent

experiments. Change in gene expression in knockout lines compared to WT was quantified by

the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

Differential gene expression in *Pf*MSP2 KO compared to WT schizonts

RNA sequencing of the schizont stage transcriptome of *Pf*3D7 MSP2 KO and *Pf*3D7 WT was

performed using the Illumina total RNA with Ribo Zero plus for library preparation and

sequenced on the NovaSeq 6000 platform with 2x150 bp paired end reads (Supplementary

Table 5). Reads were aligned to the *Pf*3D7 reference genome (PlasmoDB v61) using STAR

aligner (Dobin et al., 2013). Reads were called using Rsubread FeatureCounts (Liao et al.,

2014) with reads counted by CDS and summarised by gene. Differential gene analysis was

performed with DESeq2 (Love et al., 2014) and visualized in R studio using ggplots2

(Wickham, 2016).

Quantitative Immunofluorescence assay

Synchronous *Pf*3D7 MSP2 KO and *Pf*3D7 WT parasite cultures were treated with ML10

(Ressurreição et al., 2020) to allow mature schizonts to develop. Thin blood smears were then

made and fixed in 100%, -20°C methanol for 5 mins. Blocking of smears was performed in 1%

BSA PBS, then fluorophore conjugated WD34 (240 ng/mL) and WD33 (120 ng/mL) i-bodies

(Angage et al., 2024) in 1% BSA PBS were added and incubated at room temperature for 1 hr.

Smears were then washed, ethanol dehydrated and a coverslip mounted (ProLong Diamond

Antifade mountant with DAPI (refractive index 1.47, ThermoFisher Scientific)) and allowed

to cure for 48 hrs at room temperature. Fluorescence images were acquired using an Olympus

FV3000 confocal microscope equipped with a 100x oil objective (NA 1.4). Quantification of

parasite associated fluorescence was performed using a modified image analysis pipeline

(Liffner et al., 2020) implemented in Imaris™ (9.9 version) where non-specific background

fluorescence was removed using rolling ball background subtraction and a thresholding

algorithm applied to the image to generate binary masks, separating objects exceeding the

threshold (putative parasites) from the background. Binary masks were further refined to

exclude objects smaller or larger than a defined size range (< 3 μm and > 15 μm),

corresponding to the expected size of AMA i-body stained schizonts. Following segmentation

the surface area containing antibody signal were calculated and these values compared between

*Pf*3D7 MSP2 KO and *Pf*3D7 WT parasites.

Surface Plasmon Resonance (SPR) Assay

Surface plasmon resonance (SPR) was employed to determine the association rate constant

(K_a) of AMA1-antibody interactions in the presence of MSP2 using the Biacore T200 system

(Cytiva). Flow cells 1, 2, and 3 were activated for 14 minutes using a 1:1 mixture of 0.1 M N-

hydroxysuccinimide (NHS) and 0.4 M N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide

hydrochloride (EDC) at a flow rate of 5 μL/min. Flow cell one was immobilised with bovine

serum albumin (BSA) to serve as a reference surface. AMA1 was immobilised at a

concentration of 80 μg/mL in 10 mM sodium acetate (pH 4.5) to achieve a surface density of

1000 response units (RU) on flow cells Two and three. MSP2 was co-immobilised in flow

cell three to a surface density of 2000 RU. All surfaces were subsequently blocked with a 7-

minute injection of 1 M ethanolamine (pH 8.0). Purified antibodies were prepared in PBS

containing 0.005% Tween-20 (pH 7.4) and injected over the flow cells at a flow rate of 60

μL/min at 25°C. Single-cycle kinetics, using a bivalent analyte model, was applied to analyse

the interactions.

Enzyme-Linked Immunosorbent Assay (ELISA)

ELISAs to evaluate the effect of MSP2 on the binding of AMA1-specific antibodies were

carried out in a volume of 100 μL per well, either at room temperature for 1 hr or overnight

at 4°C. All washing steps consisted of three washes with PBS containing 0.1% Tween-20

(PBS-T). Nunc Maxisorp plates (Thermo Fisher Scientific, USA) were coated with AMA1

at a concentration of 0.8 μg/mL, followed by washing and subsequent coating with

increasing concentrations of MSP2. Wells were then blocked with 5% (w/v) skim milk in

PBS and then incubated for 1 hr at room temperature with the anti-AMA1 antibodies mouse

Fc-conjugated WD34 i-body or mouse monoclonal antibody (mAb) 4G2, or with anti-MSP2

mAb 9D8, at a concentration of 2.5 μg/mL. The primary antibody was washed off and the

HRP-conjugated secondary antibody (1:5000, Sigma Aldrich) was then applied for 1 hr at

room temperature before washing. For signal development to detect bound primary

antibodies, wells were incubated with 1-Step™ Ultra TMB-ELISA Substrate Solution

(Thermo Fisher Scientific, USA) according to the manufacturers instructions. The reaction

was stopped by adding 1 M sulfuric acid, and the absorbance at 450 nm was measured using a microplate reader.

Statistical Analysis

Three independent experiments were performed for all experiments in technical duplicate, unless otherwise stated. Data was graphed and statistical tests performed in Prism GraphPad v9 or R.

Acknowledgements

The authors would like to thank Alan Cowman, Stephen Scally and Jennifer Thompson (WEHI, Melbourne) for provision of PCRCR complex antibodies, Paul Gilson (Burnet Institute, Melb) for EXP2 antibodies, Dr Chris MacRaild (Monash University) for helpful discussion, Dr Sonja Frölich (the University of Adelaide) for assistance on quantitative immunofluorescence assays and Adelaide Microscopy for training and use of microscopy instruments. We thank Lifeblood, Australia, for provision of RBCs. DWW was supported by a Hospital Research Foundation (C-MCF-52-2019) and Australian Research Council (FT240100420) Future Fellowship. DWW and JGB by a Hospital Research Foundation Collaborative Grant (S-03-EOI-2021). JGB was supported by the National Health and Medical Research Council of Australia (Research Fellowship and Investigator Grant to JGB, Australian Centre for Research Excellence in Malaria Research). IGH and JC were supported by an Australian Research Council RTP scholarship. The Burnet Institute is supported by the NHMRC for Independent Research Institutes Infrastructure Support Scheme and the Victorian State Government Operational Infrastructure Support.

Author Contributions

Conceived and designed the experiments: DWW, IGH, JC, JGB, DA, MF, RA. Performed the experiments and phylogenetic analysis: IGH, JC, KHL, OR, DA, KRT, NB. Analysed the data: IGH, DWW, JC, KHL, OR, DA, MF. Provided key discussions on PfMSP2 and/or key reagents: RA, MF, DA. Paper writing: DWW, JC, IGH, JGB with critical input from RA, MF,

932   DA, OR, KRT and based on the PhD thesis of IGH.

Competing Interests

935   M.F. is the founding chief scientist and a shareholder in AdAlta Ltd., and R.F.A. is also a
936   shareholder in AdAlta. The other authors have declared no competing interests.

References

Adda, C.G., MacRaild, C.A., Reiling, L., Wycherley, K., Boyle, M.J., Kienzle, V., Masendycz, P., Foley, M.,
Beeson, J.G., Norton, R.S., Anders, R.F., 2012. Antigenic characterization of an intrinsically
unstructured protein, Plasmodium falciparum merozoite surface protein 2. Infect Immun 80,
4177–4185. https://doi.org/10.1128/IAI.00665-12

Adda, C.G., Murphy, V.J., Sunde, M., Waddington, L.J., Schloegel, J., Talbo, G.H., Vingas, K., Kienzle, V.,
Masciantonio, R., Howlett, G.J., Hodder, A.N., Foley, M., Anders, R.F., 2009. Plasmodium
falciparum merozoite surface protein 2 is unstructured and forms amyloid-like fibrils. Mol
Biochem Parasitol 166, 159–171. https://doi.org/10.1016/j.molbiopara.2009.03.012

Alanine, D.G.W., Quinkert, D., Kumarasingha, R., Mehmood, S., Donnellan, F.R., Minkah, N.K.,
Dadonaite, B., Diouf, A., Galaway, F., Silk, S.E., Jamwal, A., Marshall, J.M., Miura, K., Foquet, L.,
Elias, S.C., Labbé, G.M., Douglas, A.D., Jin, J., Payne, R.O., Illingworth, J.J., Pattinson, D.J., Pulido,
D., Williams, B.G., de Jongh, W.A., Wright, G.J., Kappe, S.H.I., Robinson, C. V, Long, C.A., Crabb,
B.S., Gilson, P.R., Higgins, M.K., Draper, S.J., 2019. Human Antibodies that Slow Erythrocyte
Invasion Potentiate Malaria-Neutralizing Antibodies. Cell 178, 216-228.e21.
https://doi.org/10.1016/j.cell.2019.05.025

Anders, R.F., Adda, C.G., Foley, M., Norton, R.S., 2010. Recombinant protein vaccines against the
asexual blood-stages of Plasmodium falciparum. Hum Vaccin 6, 39–53.
https://doi.org/10.4161/hv.6.1.10712

Angage, D., Chmielewski, J., Maddumage, J.C., Hesping, E., Caiazzo, S., Lai, K.H., Yeoh, L.M., Menassa,
J., Opi, D.H., Cairns, C., Puthalakath, H., Beeson, J.G., Kvansakul, M., Boddey, J.A., Wilson, D.W.,
Anders, R.F., Foley, M., 2024. A broadly cross-reactive i-body to AMA1 potently inhibits blood
and liver stages of Plasmodium parasites. Nat Commun 15, 7206.
https://doi.org/10.1038/s41467-024-50770-7

Aurrecoechea, C., Brestelli, J., Brunk, B.P., Dommer, J., Fischer, S., Gajria, B., Gao, X., Gingle, A., Grant,
G., Harb, O.S., Heiges, M., Innamorato, F., Iodice, J., Kissinger, J.C., Kraemer, E., Li, W., Miller, J.A.,
Nayak, V., Pennington, C., Pinney, D.F., Roos, D.S., Ross, C., Stoeckert, J.C.J., Treatman, C., Wang,
H., 2009. PlasmoDB: a functional genomic database for malaria parasites. Nucleic Acids Res 37,
D539–D543. https://doi.org/10.1093/nar/gkn814

Bannister, L.H., Mitchell, G.H., Butcher, G.A., Gennis, E.D., Cohen, S., 1986. Structure and development
of the surface coat of erythrocytic merozoites of Plasmodium knowlesi. Cell Tissue Res 245, 281–
290.

Baum, J., Maier, A.G., Good, R.T., Simpson, K.M., Cowman, A.F., 2005. Invasion by P. falciparum
Merozoites Suggests a Hierarchy of Molecular Interactions. PLoS Pathog 1, e37.

Beeson, J.G., Drew, D.R., Boyle, M.J., Feng, G., Fowkes, F.J.I.I., Richards, J.S., 2016. Merozoite surface
proteins in red blood cell invasion, immunity and vaccines against malaria. FEMS Microbiol Rev
40, 343–372. https://doi.org/10.1093/femsre/fuw001

Beeson, J.G., Kurtovic, L., Dobaño, C., Opi, D.H., Chan, J.-A., Feng, G., Good, M.F., Reiling, L., Boyle,
M.J., 2019. Challenges and strategies for developing efficacious and long-lasting
malaria vaccines. Sci Transl Med 11. https://doi.org/10.1126/scitranslmed.aau1458

Black, C.G., Barnwell, J.W., Huber, C.S., Galinski, M.R., Coppel, R.L., 2002. The Plasmodium vivax
homologues of merozoite surface proteins 4 and 5 from Plasmodium falciparum are expressed
at different locations in the merozoite. Mol Biochem Parasitol 120, 215–224.
https://doi.org/https://doi.org/10.1016/S0166-6851(01)00458-3

Böhme, U., Otto, T.D., Cotton, J.A., Steinbiss, S., Sanders, M., Oyola, S.O., Nicot, A., Gandon, S., Patra,
K.P., Herd, C., Bushell, E., Modrzynska, K.K., Billker, O., Vinetz, J.M., Rivero, A., Newbold, C.I.,
Berriman, M., 2018. Complete avian malaria parasite genomes reveal features associated with
lineage-specific evolution in birds and mammals. Genome Res 28, 547–560.
https://doi.org/10.1101/gr.218123.116

Boyle, M.J., Langer, C., Chan, J.-A., Hodder, A.N., Coppel, R.L., Anders, R.F., Beeson, J.G., 2014.
Sequential Processing of Merozoite Surface Proteins during and after Erythrocyte Invasion by
Plasmodium falciparum. Infect Immun 82, 924–936. https://doi.org/10.1128/IAI.00866-13

Boyle, M.J., Reiling, L., Feng, G., Langer, C., Osier, F.H., Aspeling-Jones, H., Cheng, Y.S., Stubbs, J.,
Tetteh, K.K.A., Conway, D.J., McCarthy, J.S., Muller, I., Marsh, K., Anders, R.F., Beeson, J.G., 2015.
Human antibodies fix complement to inhibit plasmodium falciparum invasion of erythrocytes

and are associated with protection against malaria. Immunity 42, 580–590.
https://doi.org/10.1016/j.immuni.2015.02.012

Boyle, M.J., Richards, J.S., Gilson, P.R., Chai, W., Beeson, J.G., 2010a. Interactions with heparin-like
molecules during erythrocyte invasion by Plasmodium falciparum merozoites. Blood 115, 4559–
4568. https://doi.org/10.1182/blood-2009-09-243725

Boyle, M.J., Wilson, D.W., Richards, J.S., Riglar, D.T., Tetteh, K.K.A., Conway, D.J., Ralph, S.A., Baum, J.,
Beeson, J.G., 2010b. Isolation of viable Plasmodium falciparum merozoites to define erythrocyte
invasion events and advance vaccine and drug development. Proc Natl Acad Sci U S A 107,
14378–14383. https://doi.org/10.1073/pnas.1009198107

Coley, A.M., Gupta, A., Murphy, V.J., Bai, T., Kim, H., Anders, R.F., Foley, M., Batchelor, A.H., 2007.
Structure of the Malaria Antigen AMA1 in Complex with a Growth-Inhibitory Antibody. PLoS
Pathog 3, e138.

Cowman, A.F., Tonkin, C.J., Tham, W.-H., Duraisingh, M.T., 2017. The Molecular Basis of Erythrocyte
Invasion by Malaria Parasites. Cell Host Microbe 22, 232–245.
https://doi.org/10.1016/j.chom.2017.07.003

Crosnier, C., Bustamante, L.Y., Bartholdson, S.J., Bei, A.K., Theron, M., Uchikawa, M., Mboup, S., Ndir,
O., Kwiatkowski, D.P., Duraisingh, M.T., Rayner, J.C., Wright, G.J., 2011. Basigin is a receptor
essential for erythrocyte invasion by Plasmodium falciparum. Nature 480, 534–537.
https://doi.org/10.1038/nature10606

Das, S., Hertrich, N., Perrin, A.J., Withers-Martinez, C., Collins, C.R., Jones, M.L., Watermeyer, J.M.,
Fobes, E.T., Martin, S.R., Saibil, H.R., Wright, G.J., Treeck, M., Epp, C., Blackman, M.J., 2015.
Processing of Plasmodium falciparum Merozoite Surface Protein MSP1 Activates a Spectrin-
Binding Function Enabling Parasite Egress from RBCs. Cell Host Microbe 18, 433–444.
https://doi.org/10.1016/j.chom.2015.09.007

Datoo, M.S., Dicko, A., Tinto, H., Ouédraogo, J.B., Hamaluba, M., Olotu, A., Beaumont, E., Ramos
Lopez, F., Natama, H.M., Weston, S., Chemba, M., Compaore, Y.D., Issiaka, D., Salou, D., Some,
A.M., Omenda, S., Lawrie, A., Bejon, P., Rao, H., Chandramohan, D., Roberts, R., Bharati, S.,
Stockdale, L., Gairola, S., Greenwood, B.M., Ewer, K.J., Bradley, J., Kulkarni, P.S., Shaligram, U.,
Hill, A.V.S., Mahamar, A., Sanogo, K., Sidibe, Y., Diarra, K., Samassekou, M., Attaher, O., Tapily,
A., Diallo, M., Dicko, O.M., Kaya, M., Maguiraga, S.O., Sankare, Y., Yalcouye, H., Diarra, S.,
Niambele, S.M., Thera, I., Sagara, I., Sylla, M., Dolo, A., Misidai, N., Simando, S., Msami, H., Juma,
O., Gutapaka, N., Paul, R., Mswata, S., Sasamalo, I., Johaness, K., Sultan, M., Alexander, A.,

Kimaro, I., Lwanga, K., Mtungwe, M., Khamis, K., Rugarabam, L., Kalinga, W., Mohammed, M.,
Kamange, J., Msangi, J., Mwaijande, B., Mtaka, I., Mhapa, M., Mlaganile, T., Mbaga, T., Yerbanga,
R.S., Samtouma, W., Sienou, A.A., Kabre, Z., Ouedraogo, W.J.M., Yarbanga, G.A.B., Zongo, I.,
Savadogo, H., Sanon, J., Compaore, J., Kere, I., Yoni, F.L., Sanre, T.M., Ouattara, S.B., Provstgaard-
Morys, S., Woods, D., Snow, R.W., Amek, N., Ngetsa, C.J., Ochola-Oyier, L.I., Musyoki, J., Munene,
M., Mumba, N., Adetifa, U.J., Muiruri, C.M., Mwawaka, J.S., Mwaganyuma, M.H., Ndichu, M.N.,
Weya, J.O., Njogu, K., Grant, J., Webster, J., Lakhkar, A., Ido, N.F.A., Traore, O., Tahita, M.C.,
Bonko, M. dit A., Rouamba, T., Ouedraogo, D.F., Soma, R., Millogo, A., Ouedraogo, E., Sorgho, F.,
Konate, F., Valea, I., 2024. Safety and efficacy of malaria vaccine candidate R21/Matrix-M in
African children: a multicentre, double-blind, randomised, phase 3 trial. The Lancet 403, 533–
544. https://doi.org/10.1016/S0140-6736(23)02511-4

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras,
T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21.
https://doi.org/10.1093/bioinformatics/bts635

Douglas, A.D., Williams, A.R., Illingworth, J.J., Kamuyu, G., Biswas, S., Goodman, A.L., Wyllie, D.H.,
Crosnier, C., Miura, K., Wright, G.J., Long, C.A., Osier, F.H., Marsh, K., Turner, A. V., Hill, A.V.S.,
Draper, S.J., 2011. The blood-stage malaria antigen PfRH5 is susceptible to vaccine-inducible
cross-strain neutralizing antibody. Nat Commun 2, 601. https://doi.org/10.1038/ncomms1615

Duraisingh, M.T., Maier, A.G., Triglia, T., Cowman, A.F., 2003. Erythrocyte-binding antigen 175
mediates invasion in Plasmodium falciparum utilizing sialic acid-dependent and -independent
pathways. Proc Natl Acad Sci U S A 100, 4796–4801. https://doi.org/10.1073/pnas.0730883100

Eisenhaber, B., Bork, P., Eisenhaber, F., 1999. Prediction of potential GPI-modification sites in
proprotein sequences. J Mol Biol 292, 741–58. https://doi.org/10.1006/jmbi.1999.3069

Escalante, A.A., Cepeda, A.S., Pacheco, M.A., 2022. Why Plasmodium vivax and Plasmodium
falciparum are so different? A tale of two clades and their species diversities. Malar J 21, 139.
https://doi.org/10.1186/s12936-022-04130-9

Feng, G., Boyle, M.J., Cross, N., Chan, J.-A., Reiling, L., Osier, F., Stanisic, D.I., Mueller, I., Anders, R.F.,
McCarthy, J.S., Richards, J.S., Beeson, J.G., 2018. Human Immunization With a Polymorphic
Malaria Vaccine Candidate Induced Antibodies to Conserved Epitopes That Promote Functional
Antibodies to Multiple Parasite Strains. J Infect Dis 218, 35–43.
https://doi.org/10.1093/infdis/jiy170

Ferreira, E. d'Avila, Alexandre, M.A., Salinas, J.L., de Siqueira, A.M., Benzecry, S.G., de Lacerda, M.V.G.,
Monteiro, W.M., 2015. Association between anthropometry-based nutritional status and
malaria: a systematic review of observational studies. Malaria J 14, 346.
https://doi.org/10.1186/s12936-015-0870-5

Fowkes, F.J.I., Richards, J.S., Simpson, J.A., Beeson, J.G., 2010. The Relationship between Anti-
merozoite Antibodies and Incidence of Plasmodium falciparum Malaria: A Systematic Review
and Meta-analysis. PLoS Med 7, e1000218. https://doi.org/10.1371/journal.pmed.1000218

Genton, B., Al-Yaman, F., Betuela, I., Anders, R.F., Saul, A., Baea, K., Mellombo, M., Taraika, J., Brown,
G. V., Pye, D., Irving, D.O., Felger, I., Beck, H.-P.P., Smith, T.A., Alpers, M.P., 2003. Safety and
immunogenicity of a three-component blood-stage malaria vaccine (MSP1, MSP2, RESA) against
Plasmodium falciparum in Papua New Guinean children. Vaccine 22, 30–41.
https://doi.org/https://doi.org/10.1016/S0264-410X(03)00536-X

Genton, B., Betuela, I., Felger, I., Al-Yaman, F., Anders, R.F., Saul, A., Rare, L., Baisor, M., Lorry, K.,
Brown, G. V, Pye, D., Irving, D.O., Smith, T.A., Beck, H.-P., Alpers, M.P., 2002. A Recombinant
Blood-Stage Malaria Vaccine Reduces Plasmodium falciparum Density and Exerts Selective
Pressure on Parasite Populations in a Phase 1-2b Trial in Papua New Guinea. J Infect Dis 185,
820–827.

Gilson, P.R., Nebl, T., Vukcevic, D., Moritz, R.L., Sargeant, T., Speed, T.P., Schofield, L., Crabb, B.S.,
2006. Identification and Stoichiometry of Glycosylphosphatidylinositol-anchored Membrane
Proteins of the Human Malaria Parasite *Plasmodium falciparum*. Molecular & Cellular
Proteomics 5, 1286–1299. https://doi.org/10.1074/mcp.M600035-MCP200

Griffiths, K., Dolezal, O., Cao, B., Nilsson, S.K., See, H.B., Pfleger, K.D.G., Roche, M., Gorry, P.R., Pow,
A., Viduka, K., Lim, K., Lu, B.G.C., Chang, D.H.C., Murray-Rust, T., Kvansakul, M., Perugini, M.A.,
Dogovski, C., Doerflinger, M., Zhang, Y., Parisi, K., Casey, J.L., Nuttall, S.D., Foley, M., 2016. i-
bodies, Human Single Domain Antibodies That Antagonize Chemokine Receptor CXCR4. J Biol
Chem 291, 12641–12657. https://doi.org/10.1074/jbc.M116.721050

Griffiths, K., Habiel, D.M., Jaffar, J., Binder, U., Darby, W.G., Hosking, C.G., Skerra, A., Westall, G.P.,
Hogaboam, C.M., Foley, M., 2018. Anti-fibrotic Effects of CXCR4-Targeting i-body AD-114 in
Preclinical Models of Pulmonary Fibrosis. Sci Rep 8, 3212. https://doi.org/10.1038/s41598-018-
20811-5

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates,
R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-

Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., Hassabis, D., 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596, 583–589. https://doi.org/10.1038/s41586-021-03819-2

Kals, E., Kals, M., Lees, R.A., Introini, V., Kemp, A., Silvester, E., Collins, C.R., Umrekar, T., Kotar, J., Cicuta, P., Rayner, J.C., 2024. Application of optical tweezer technology reveals that PfEBA and PfRH ligands, not PfMSP1, play a central role in Plasmodium falciparum merozoite-erythrocyte attachment. PLoS Pathog 20, e1012041. https://doi.org/10.1371/journal.ppat.1012041

Kocken, C.H., van der Wel, A.M., Dubbeld, M.A., Narum, D.L., van de Rijke, F.M., van Gemert, G.J., van der Linde, X., Bannister, L.H., Janse, C., Waters, A.P., Thomas, A.W., 1998. Precise timing of expression of a Plasmodium falciparum-derived transgene in Plasmodium berghei is a critical determinant of subsequent subcellular localization. J Biol Chem 273, 15119–15124. https://doi.org/10.1074/jbc.273.24.15119

Lamarque, M., Besteiro, S., Papoin, J., Roques, M., Vulliez-Le Normand, B., Morlon-Guyot, J., Dubremetz, J.-F., Fauquenoy, S., Tomavo, S., Faber, B.W., Kocken, C.H., Thomas, A.W., Boulanger, M.J., Bentley, G.A., Lebrun, M., 2011. The RON2-AMA1 Interaction is a Critical Step in Moving Junction-Dependent Invasion by Apicomplexan Parasites. PLoS Pathog 7, e1001276.

Li, X., Chen, H., Oo, T.H., Daly, T.M., Bergman, L.W., Liu, S.-C., Chishti, A.H., Oh, S.S., 2004. A co-ligand complex anchors Plasmodium falciparum merozoites to the erythrocyte invasion receptor band 3. J Biol Chem 279, 5765–5771. https://doi.org/10.1074/jbc.M308716200

Li, Y., Yang, L., Yang, L.-Q., 2024. Effects of intrinsically disordered regions in gp120 underlying HIV neutralization phenotypes. Biochem Biophys Res Commun 709, 149830. https://doi.org/10.1016/j.bbrc.2024.149830

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30, 923–930. https://doi.org/10.1093/bioinformatics/btt656

Liffner, B., Frölich, S., Heinemann, G.K., Liu, B., Ralph, S.A., Dixon, M.W.A., Gilberger, T.W., Wilson, D.W., 2020. PfCERLI1 is a conserved rhoptry associated protein essential for Plasmodium falciparum merozoite invasion of erythrocytes. Nat Commun 11. https://doi.org/10.1038/s41467-020-15127-w

Livak, K.J., Schmittgen, T.D., 2001. Analysis of Relative Gene Expression Data Using Real-Time

Quantitative PCR and the 2−ΔΔCT Method. Methods 25, 402–408.

https://doi.org/https://doi.org/10.1006/meth.2001.1262

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol 15, 550. https://doi.org/10.1186/s13059-014-0550-8

Lu, C., Zheng, X., Zhang, W., Zhao, H., MacRaild, C.A., Norton, R.S., Zhuang, Y., Wang, J., Zhang, X.,

2019. Interaction of merozoite surface protein 2 with lipid membranes. FEBS Lett 593, 288–295.

https://doi.org/10.1002/1873-3468.13320

MacRaild, C.A., Pedersen, M.Ø., Anders, R.F., Norton, R.S., 2012. Lipid interactions of the malaria

antigen merozoite surface protein 2. Biochim Biophys Acta 1818, 2572–8.

https://doi.org/10.1016/j.bbamem.2012.06.015

MacRaild, C.A., Zachrdla, M., Andrew, D., Krishnarjuna, B., Nováček, J., Žídek, L., Sklenář, V., Richards,

J.S., Beeson, J.G., Anders, R.F., Norton, R.S., 2015. Conformational dynamics and antigenicity in

the disordered malaria antigen merozoite surface protein 2. PLoS One 10, e0119899.

https://doi.org/10.1371/journal.pone.0119899

McCarthy, J.S., Marjason, J., Elliott, S., Fahey, P., Bang, G., Malkin, E., Tierney, E., Aked-Hurditch, H.,

Adda, C., Cross, N., Richards, J.S., Fowkes, F.J.I., Boyle, M.J., Long, C., Druilhe, P., Beeson, J.G.,

Anders, R.F., 2011. A phase 1 trial of MSP2-C1, a blood-stage malaria vaccine containing 2

isoforms of MSP2 formulated with montanide® ISA 720. PLoS One 6.

https://doi.org/10.1371/journal.pone.0024413

Ocampo, M., Urquiza, M., Guzmán, F., Rodriguez, L.E., Suarez, J., Curtidor, H., Rosas, J., Diaz, M.,

Patarroyo, M.E., 2003. Two MSA 2 peptides that bind to human red blood cells are relevant to

Plasmodium falciparum merozoite invasion. The Journal of Peptide Research 55, 216–223.

https://doi.org/10.1034/j.1399-3011.2000.00174.x

Orlandi, P.A., Klotz, F.W., Haynes, J.D., 1992. A malaria invasion receptor, the 175-kilodalton

erythrocyte binding antigen of Plasmodium falciparum recognizes the terminal Neu5Ac (alpha 2-

3) Gal-sequences of glycophorin A. J Cell Biol 116, 901–909.

Osier, F.H., Feng, G., Boyle, M.J., Langer, C., Zhou, J., Richards, J.S., McCallum, F.J., Reiling, L.,

Jaworowski, A., Anders, R.F., Marsh, K., Beeson, J.G., 2014. Opsonic phagocytosis of Plasmodium

falciparum merozoites: mechanism in human immunity and a correlate of protection against

malaria. BMC Med 12, 108. https://doi.org/10.1186/1741-7015-12-108

Osier, F.H.A., Murungi, L.M., Fegan, G., Tuju, J., Tetteh, K.K., Bull, P.C., Conway, D.J., Marsh, K., 2010.

Allele-specific antibodies to Plasmodium falciparum merozoite surface protein-2 and protection

against clinical malaria. Parasite Immunol 32, 193–201. https://doi.org/10.1111/j.1365-

3024.2009.01178.x

Peng, D., Tarleton, R., 2015. EuPaGDT: a web tool tailored to design CRISPR guide RNAs for eukaryotic

pathogens. Microb Genom 1. https://doi.org/10.1099/mgen.0.000033

Persson, K.E.M., Fowkes, F.J.I., McCallum, F.J., Gicheru, N., Reiling, L., Richards, J.S., Wilson, D.W.,

Lopaticki, S., Cowman, A.F., Marsh, K., Beeson, J.G., 2013. Erythrocyte-Binding Antigens of

Plasmodium falciparum Are Targets of Human Inhibitory Antibodies and Function To Evade

Naturally   Acquired   Immunity.   The   Journal   of   Immunology   191,   785–794.

https://doi.org/10.4049/jimmunol.1300444

Persson, K.E.M., McCallum, F.J., Reiling, L., Lister, N.A., Stubbs, J., Cowman, A.F., Marsh, K., Beeson,

J.G., 2008. Variation in use of erythrocyte invasion pathways by Plasmodium

falciparum mediates evasion of human inhibitory antibodies. J Clin Invest 118, 342–351.

https://doi.org/10.1172/JCI32138

Reiling, L., Boyle, M.J., White, M.T., Wilson, D.W., Feng, G., Weaver, R., Opi, D.H., Persson, K.E.M.,

Richards, J.S., Siba, P.M., Fowkes, F.J.I., Takashima, E., Tsuboi, T., Mueller, I., Beeson, J.G., 2019.

Targets of complement-fixing antibodies in protective immunity against malaria in children. Nat

Commun 10, 610. https://doi.org/10.1038/s41467-019-08528-z

Ressurreição, M., Thomas, J.A., Nofal, S.D., Flueck, C., Moon, R.W., Baker, D.A., van Ooij, C., 2020. Use

of a highly specific kinase inhibitor for rapid, simple and precise synchronization of Plasmodium

falciparum and Plasmodium knowlesi asexual blood-stage parasites. PLoS One 15, e0235798.

https://doi.org/10.1371/journal.pone.0235798

Richards, J.S., Arumugam, T.U., Reiling, L., Healer, J., Hodder, A.N., Fowkes, F.J.I., Cross, N., Langer, C.,

Takeo, S., Uboldi, A.D., Thompson, J.K., Gilson, P.R., Coppel, R.L., Siba, P.M., King, C.L., Torii, M.,

Chitnis, C.E., Narum, D.L., Mueller, I., Crabb, B.S., Cowman, A.F., Tsuboi, T., Beeson, J.G., 2013.

Identification and Prioritization of Merozoite Antigens as Targets of Protective Human Immunity

to Plasmodium falciparum Malaria for Vaccine and Biomarker Development. The Journal of

Immunology 191, 795 LP – 809.

RTSS Clinical Trials Partnership, 2015. Efficacy and safety of RTS,S/AS01 malaria vaccine with or

without a booster dose in infants and children in Africa: final results of a phase 3, individually

randomised,   controlled   trial.   Lancet   386,   31–45.
https://doi.org/http://dx.doi.org/10.1016/S0140-6736(15)60721-8

Sagara, I., Dicko, A., Ellis, R.D., Fay, M.P., Diawara, S.I., Assadou, M.H., Sissoko, M.S., Kone, M., Diallo,
A.I., Saye, R., Guindo, M.A., Kante, O., Niambele, M.B., Miura, K., Mullen, G.E.D., Pierce, M.,
Martin, L.B., Dolo, A., Diallo, D.A., Doumbo, O.K., Miller, L.H., Saul, A., 2009. A randomized
controlled phase 2 trial of the blood stage AMA1-C1/Alhydrogel malaria vaccine in children in
Mali. Vaccine 27, 3090–3098. https://doi.org/10.1016/j.vaccine.2009.03.014

Salanti, A., Staalsoe, T., Lavstsen, T., Jensen, A.T.R., Sowa, M.P.K., Arnot, D.E., Hviid, L., Theander, T.G.,
2003. Selective upregulation of a single distinctly structured var gene in chondroitin sulphate A-
adhering Plasmodium falciparum involved in pregnancy-associated malaria. Mol Microbiol 49,
179–191. https://doi.org/10.1046/j.1365-2958.2003.03570.x

Sanders, P.R., Kats, L.M., Drew, D.R., O'Donnell, R.A., O'Neill, M., Maier, A.G., Coppel, R.L., Crabb, B.S.,
2006. A Set of Glycosylphosphatidyl Inositol-Anchored Membrane Proteins of Plasmodium
falciparum Is Refractory to Genetic Deletion. Infect Immun 74, 4330–4338.
https://doi.org/10.1128/IAI.00054-06

Scally, S.W., Triglia, T., Evelyn, C., Seager, B.A., Pasternak, M., Lim, P.S., Healer, J., Geoghegan, N.D.,
Adair, A., Tham, W.-H., Dagley, L.F., Rogers, K.L., Cowman, A.F., 2022. PCRCR complex is essential
for invasion of human erythrocytes by Plasmodium falciparum. Nat Microbiol 7, 2039–2053.
https://doi.org/10.1038/s41564-022-01261-2

Sheehy, S.H., Duncan, C.J.A., Elias, S.C., Choudhary, P., Biswas, S., Halstead, F.D., Collins, K.A., Edwards,
N.J., Douglas, A.D., Anagnostou, N.A., Ewer, K.J., Havelock, T., Mahungu, T., Bliss, C.M., Miura,
K., Poulton, I.D., Lillie, P.J., Antrobus, R.D., Berrie, E., Moyle, S., Gantlett, K., Colloca, S., Cortese,
R., Long, C.A., Sinden, R.E., Gilbert, S.C., Lawrie, A.M., Doherty, T., Faust, S.N., Nicosia, A., Hill,
A.V.S., Draper, S.J., 2012. ChAd63-MVA-vectored blood-stage malaria vaccines targeting MSP1
and AMA1: assessment of efficacy against mosquito bite challenge in humans. Mol Ther 20,
2355–2368. https://doi.org/10.1038/mt.2012.223

Smythe, J.A., Coppel, R.L., Day, K.P., Martin, R.K., Oduola, A.M., Kemp, D.J., Anders, R.F., 1991.
Structural diversity in the Plasmodium falciparum merozoite surface antigen 2. Proceedings of
the National Academy of Sciences 88, 1751 LP – 1755. https://doi.org/10.1073/pnas.88.5.1751

Srinivasan, P., Beatty, W.L., Diouf, A., Herrera, R., Ambroggio, X., Moch, J.K., Tyler, J.S., Narum, D.L.,
Pierce, S.K., Boothroyd, J.C., Haynes, J.D., Miller, L.H., 2011. Binding of Plasmodium merozoite

proteins RON2 and AMA1 triggers commitment to invasion. Proceedings of the National
Academy of Sciences 108, 13275–13280. https://doi.org/10.1073/pnas.1110303108

Stanisic, D.I., Richards, J.S., McCallum, F.J., Michon, P., King, C.L., Schoepflin, S., Gilson, P.R., Murphy,
V.J., Anders, R.F., Mueller, I., Beeson, J.G., 2009. Immunoglobulin G Subclass-Specific Responses
against Plasmodium falciparum Merozoite Antigens Are Associated with Control of Parasitemia
and Protection from Symptomatic Illness. Infect Immun 77, 1165–1174.
https://doi.org/10.1128/IAI.01129-08

Stejskal, L., Kalemera, M.D., Lewis, C.B., Palor, M., Walker, L., Daviter, T., Lees, W.D., Moss, D.S.,
Kremyda-Vlachou, M., Kozlakidis, Z., Gallo, G., Bailey, D., Rosenberg, W., Illingworth, C.J.R.,
Shepherd, A.J., Grove, J., 2022. An entropic safety catch controls hepatitis C virus entry and
antibody resistance. Elife 11. https://doi.org/10.7554/eLife.71854

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version
11. Mol Biol Evol 38, 3022–3027. https://doi.org/10.1093/molbev/msab120

Taylor, H.M., McRobert, L., Grainger, M., Sicard, A., Dluzewski, A.R., Hopp, C.S., Holder, A.A., Baker,
D.A., 2010. The malaria parasite cyclic GMP-dependent protein kinase plays a central role in
blood-stage schizogony. Eukaryot Cell 9, 37–45. https://doi.org/10.1128/EC.00186-09

Thera, M.A., Doumbo, O.K., Coulibaly, D., Laurens, M.B., Ouattara, A., Kone, A.K., Guindo, A.B., Traore,
K., Traore, I., Kouriba, B., Diallo, D.A., Diarra, I., Daou, M., Dolo, A., Tolo, Y., Sissoko, M.S.,
Niangaly, A., Sissoko, M., Takala-Harrison, S., Lyke, K.E., Wu, Y., Blackwelder, W.C., Godeaux, O.,
Vekemans, J., Dubois, M.-C., Ballou, W.R., Cohen, J., Thompson, D., Dube, T., Soisson, L., Diggs,
C.L., House, B., Lanar, D.E., Dutta, S., Heppner, D.G., Plowe, C. V., 2011. A Field Trial to Assess a
Blood-Stage Malaria Vaccine. New England Journal of Medicine 365, 1004–1013.
https://doi.org/10.1056/NEJMoa1008115

Weiss, G.E., Gilson, P.R., Taechalertpaisarn, T., Tham, W.-H.H., de Jong, N.W.M.M., Harvey, K.L.,
Fowkes, F.J.I.I., Barlow, P.N., Rayner, J.C., Wright, G.J., Cowman, A.F., Crabb, B.S., 2015. Revealing
the Sequence and Resulting Cellular Morphology of Receptor-Ligand Interactions during
Plasmodium falciparum Invasion of Erythrocytes. PLoS Pathog 11, 284–295.
https://doi.org/10.1371/journal.ppat.1004670

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wilson, D.W., Goodman, C.D., Sleebs, B.E., Weiss, G.E., de Jong, N.W.M., Angrisano, F., Langer, C.,
Baum, J., Crabb, B.S., Gilson, P.R., McFadden, G.I., Beeson, J.G., 2015. Macrolides rapidly inhibit

red blood cell invasion by the human malaria parasite, Plasmodium falciparum. BMC Biol 13, 52.

https://doi.org/10.1186/s12915-015-0162-0

World Health Organization, 2024. World malaria report 2024: addressing inequity in the global malaria

response. World Health Organization, Geneva.

Yang, X., Adda, C.G., MacRaild, C.A., Low, A., Zhang, X., Zeng, W., Jackson, D.C., Anders, R.F., Norton,

R.S., 2010. Identification of key residues involved in fibril formation by the conserved N-terminal

region of Plasmodium falciparum merozoite surface protein 2 (MSP2). Biochimie 92, 1287–95.

https://doi.org/10.1016/j.biochi.2010.06.001

Zerebinski, J., Margerie, L., Han, N.S., Moll, M., Ritvos, M., Jahnmatz, P., Ahlborg, N., Ngasala, B.,

Rooth, I., Sjöberg, R., Sundling, C., Yman, V., Färnert, A., Plaza, D.F., 2024. Naturally acquired IgG

responses to Plasmodium falciparum do not target the conserved termini of the malaria vaccine

candidate Merozoite Surface Protein 2. Front Immunol 15, 1501700.

https://doi.org/10.3389/fimmu.2024.1501700

Zhang, X., Adda, C.G., Low, A., Zhang, J., Zhang, W., Sun, H., Tu, X., Anders, R.F., Norton, R.S., 2012.

Role of the helical structure of the N-terminal region of plasmodium falciparum merozoite

surface protein 2 in fibril formation and membrane interaction. Biochemistry 51, 1380–1387.

https://doi.org/10.1021/bi201880s

Zhang, X., Perugini, M.A., Yao, S., Adda, C.G., Murphy, V.J., Low, A., Anders, R.F., Norton, R.S., 2008.

Solution conformation, backbone dynamics and lipid interactions of the intrinsically

unstructured malaria surface protein MSP2. J Mol Biol 379, 105–21.

https://doi.org/10.1016/j.jmb.2008.03.039

1263

Table 1. Amino acids sequence properties for the N and C-terminal conserved, and N-terminal repeat, regions for available *P. falciparum* 3D7, *Laverania* (including 3 *Laverania* sequences of unknown speciation) *P. relictum* and *P. gallinaceum* MSP2 sequences.

Species	N terminal				Repeat regions						C terminal			
	Polar uncharged	Basic	Acidic	Hydrophobic	% G	% S	% T	% A	Number of repeats	Amino acid range	Polar uncharged	Basic	Acidic	Hydrophobic
P. falciparum 3D7	50%	17%	4%	29%	50%	25%	0%	25%	6	54-77	46%	16%	10%	28%
P. praefalciparum G01	54%	17%	4%	25%	29%	7%	43%	7%	2	62-75	48%	16%	12%	24%
P. reichenowi CDC	54%	17%	4%	25%	36%	24%	18%	13%	3+4^	51-105	48%	16%	12%	24%
P. reichenowi G01	54%	17%	4%	25%	33%	0%	33%	33%	5	67-81	48%	16%	12%	24%
P. reichenowi XM_012905494.2	54%	17%	4%	25%	29%	0%	29%	38%	6*	67-87	48%	16%	12%	24%
P. billcollinsi G01	54%	17%	4%	25%	31%	6%	25%	13%	4*	53-68	52%	18%	10%	20%
P. blacklocki G01	50%	21%	4%	25%	23%	31%	31%	8%	4*	52-64	40%	16%	16%	28%
P. adleri G01	50%	17%	4%	29%	19%	19%	25%	11%	5*	45-80	44%	10%	12%	34%
P. gaboni SY75	46%	17%	8%	29%	21%	33%	21%	17%	6	61-84	45%	12%	10%	33%
P. relictum	54%	13%	4%	29%	33%	0%	33%	33%	8	57-80	33%	12%	22%	33%
P. gallinaceum 8A	38%	25%	4%	33%	31%	6%	19%	25%	4*	55-70	33%	8%	25%	34%
Plasmodium sp. 1 MAP-2013 JX899678	50%	17%	4%	29%	26%	4%	30%	22%	4*	51-73	48%	16%	12%	24%
Plasmodium sp. 2 MAP-2013 JX893169	50%	17%	4%	29%	26%	4%	30%	22%	4*	55-77	48%	16%	12%	24%
Plasmodium sp. 2 MAP-2013 JX893170	46%	21%	4%	29%	25%	6%	31%	19%	4*	51-66	48%	16%	12%	24%

Conservation of amino acids between repeats

*insertion present
^two repeat sequences

Full	100% conservation of amino acid composition and positions
Partially degenerate	81-99% conservation of amino acid composition (between all pairs of repeats); full conservation of amino acid positions
Degenerate	50-80% conservation of amino acid composition in partially conserved positions (between all pairs of repeats)
Highly degenerate	50-80% conservation of amino acid composition to that in one selected repeat; no conservation of amino acid positions

Table 2: Genes significantly up or down-regulated with Pf3D7 MSP2 KO

Gene Name	log2FoldChange	p-value	Name
PF3D7_0202400	2.547546545	1.09E-36	translation-enhancing factor
PF3D7_1348000	1.604559487	0.000176	conserved Plasmodium protein, unknown function
PF3D7_1423500	1.411878628	4.31E-08	conserved Plasmodium protein, unknown function
PF3D7_1208200	1.384846624	3.59E-05	cysteine repeat modular protein 3
PF3D7_1461800	1.376899732	9.97E-05	conserved Plasmodium protein, unknown function
PF3D7_0909100	1.155215486	4.07E-05	conserved Plasmodium membrane protein, unknown function
PF3D7_1426500	1.021678659	1.52E-05	ABC transporter G family member 2
PF3D7_0322200	1.014455112	0.000313	conserved protein, unknown function
PF3D7_1372100	-2.436194076	9.31E-18	Plasmodium exported protein (PHISTb), unknown function
PF3D7_1149400	-2.748153616	1.72E-25	Plasmodium exported protein, unknown function
PF3D7_0206800	-4.324734054	9.44E-13	merozoite surface protein 2

Supplementary Figure 1 Alignment of full-length amino acid protein sequences for all *Pf*MSP2 like sequences on PlasmoDB. Alignment done using MUSCLE alignment with default settings.

Identity

1. P. falciparum 3D7
2. P. praefalciparum
3. P. reichenowi CDC
4. P. reichenowi G01
5. P. reichenowi XM_012905494.2
6. P. billcollinsi G01
7. P. blacklocki G01
8. P. adleri G01
9. P. gaboni SY75
10. P. relictum
11. P. gallinaceum 8A
12. P. sp. 1 MAP-2013 JX899678
13. P. sp. 2 MAP-2013 JX893169
14. P. sp. 2 MAP-2013 JX893170

Supplementary Figure 2 Alignment (Blosum 62) of Amino acid sequences for the N-terminal conserved region of *P. falciparum* 3D7, available *Laverania*, *P. relictum* and *P. gallinaceum* MSP2 like sequences on PlasmoDB and identified through a BLAST search on the NCBI server. P. sp. 1, 2 and 3 represent MSP2 sequences identified using a BLAST search that are indicated to be from an unspecified *Laverania* spp. parasite. Amino acids are coloured according to their properties (RasMol).

Supplementary Figure 3 Amino Acid sequences for the N-terminal repeat region of *P. falciparum* 3D7, available *Laverania*, *P. relictum* and *P. gallinaceum* MSP2 like sequences on PlasmoDB and identified through a BLAST search on the NCBI server. P. sp. 1, 2 and 3 represent MSP2 sequences identified using a BLAST search that are indicated to be from an unspecific *Laverania* spp. parasite. Sequences presented are not aligned, with the exceptions of the 3 *P. reichenowi* and 3 unspecified *Laverania* sequences in order to highlight the conserved sequence despite an insertion. Insertions that are considered to break up the small amino acid and/or repeat structure for the purposes of this comparison are underlined in Black. Respeats, both conserved and degenerate, are underlined in grey. Amino acids are coloured according to their properties (RasMol).

Supplementary Figure 4 Alignment of amino acid sequences for the region corresponding to the C-terminal conserved region of *P. falciparum* 3D7, available *Laverania*, *P. relictum* and *P. gallinaceum* MSP2 like sequences on PlasmoDB and identified through a BLAST search on the NCBI server. *P. sp.* 1, 2 and 3 represent MSP2 sequences identified using a BLAST search that are indicated to be from an unspecified *Laverania* spp. parasite. Initial alignment done using Blosum62 with default settings, with subsequent manual modification as required to align to the cysteine residues and other prominent areas of conservation between the *Laverania* and avian malarias. Amino acids are coloured according to their properties (RasMol).

Supplementary Figure 5 Coverage plot (IGV) showing absence of majority of *msp2* in schizont transcriptome of *Pf*3D7 DMSP2 parasites but *msp2* presence in *Pf*3D7 MSP2WT. Small section of *msp2* transcribe in *Pf*3D7 ΔMSP2 parasites corresponds with the end of the homology region and primarily covers the signal peptide.

Supplementary Table 1. Primers used to generate MSP2 KO lines.

5' flank MSP2 KO F	GGTCCGCGGCTCGACTAATCAATTTACAATTTC	Sac11
5' flank MSP2 KO R	GGTACTAGTGCCATACTTCTCCTTATACTC	Spe1
3' flank MSP2 KO F	GGTGAATTCCTCTTCATTTTAAAACATTGAC	Ecor1
3' flank MSP2 KO R	GGTCCATGGGTTTTTTCAATGCGTGC	Nco1
Dd2 MSP2 KO 5' flank F	GGTCCGCGGCTCGACTAATCAATTTACAATTTCC	Sac11
Dd2 MSP2 KO 5' flank R	GGTACTAGTCTAGTAGTATTAGAACCTTCATTTG	Spe1
Dd2 MSP2 KO 3' flank F	GGTGAATTCATTCTCTTCATTTTAAAACATTGAC	EcoR1
Dd2 MSP2 KO 3' flank R	GGTCCATGGGTACTTGAAGAAATATGGTACC	Nco1
PfMSP2 guide 1 F	TAAGTATATAATATTTGGTAATGGTGCAGATGCTGGTTTT AGAGCTAGAA	
PfMSP2 guide 1 R	TTCTAGCTCTAAAACCAGCATCTGCACCATTACCAAATATT ATATACTTA	
Dd2 MSP2 Guide 2 F	TAAGTATATAATATTGCACCAGAGAATAAAGGTACGTTTT AGAGCTAGAA	
Dd2 MSP2 Guide 2 R	TTCTAGCTCTAAAACGTACCTTTATTCTCTGGTGCAATAT TATATACTTA	
5' int 3D7 MSP2 KO F	CTTTTCATATAATAAATCAAATGCAC	
5' int 3D7 MSP2 KO R	TACAAAATGCTTAAGACAGATC	
3D7 WT 5' int check R	GGTAGTTGTGGTAGTAGC	
Dd2 5' check F	CAATTACGATATAAAACCTAGTATCTTTC	
Dd2 WT 5' int check R	CTATTTGTACTCCTTTGACTTCCAC	

Supplementary Table 2. Antibodies used in this study.

Serum/Antibody/ibody	Source
Anti-MSP2 mAb 2F2	Adda et al Infect Immun 2012
Anti-MSP2 Rb FC27	Robin Anders, LaTrobe University
Anti-EXP2 Rb	Bullen et al. J Biol Chem 2012
Anti-PfAldolase	Abcam
Anti-PfRh2b Rb 1055-3	Lopaticki et al Infect and Immun 2011
Anti-PfEBA175/Rh2b Rb 1056-3	Lopaticki et al Infect and Immun 2011
Anti-PfEBA175/Rh4 Rb 1049-3	Lopaticki et al Infect and Immun 2011
Anti-PfEBA175/Rh2b/Rh4 Rb 1067-3	Lopaticki et al Infect and Immun 2011
Anti-w2mef EBA175 Rb	Healer et al PLoS One 2013
Anti-MSP1-19 Rb (6858, 645, 647)	Paul Gilson, Burnet Institute
Anti-Rh5.1 Rt 494	Healer et al Front Cell Infect Microbiol 2022
Anti-PTRAMP nAb H8	Scally et al Nat Microbiol 2022
Anti-CSS nAb D2	Scally et al Nat Microbiol 2022
Anti-CSS nAb D2-FC	Scally et al Nat Microbiol 2022
Anti-AMA1 3D7 Rb (1072, 1149, 1161)	Drew et al. PLoS One 2012
Anti-AMA1 w2mef Rb (1151, 1164, 1163, 128/7)	Drew et al. PLoS One 2012
Anti-AMA1 mAb 1F9	Coley et al PEDS 2001
Anti-AMA1 mAb 4G2	Collins et al J. Biol. Chem. 2007
Anti-AMA1 ibody WD33	Angange et al Under Review
Anti-AMA1 ibody WD34	Angange et al Under Review
Anti-AMA1 WD34	Angange et al Under Review
Anti-AMA1 ibody WD34-FC	Angange et al Under Review

Rb= rabbit. mAb= mouse monoclonal. Rt= rat. nAb= camelid nanobody. I-body= single domain antibody derived from shark variable new antigen receptor (V_{NAR}).

Supplementary Table 3. Enzymes used to remove/inactivate RBC receptors and target residues of the enzyme.

Enzyme	Cleavage Target	Supplier
Neuraminidase (0.067 U/mL)	Sialic acid residues	Sigma-Aldrich
High Trypsin (1 mg/mL)	Lysine or Arginine residues, unless followed by a proline	Sigma-Aldrich
Low Trypsin (0.067 mg/mL)	Lysine or Arginine residues, unless followed by a proline	Sigma-Aldrich
Chymotrypsin (1 mg/mL)	Tyrosine, Tryptophan or Phenylalanine	Worthington Biochemical Corporation
Chymotrypsin (1 mg/mL)/ Trypsin (0.067 mg/mL)	Tyrosine, Tryptophan, Phenylalanine, Lysine or Arginine residues, unless followed by a proline	Worthington Biochemical Corporation & Sigma-Aldrich

Supplementary Table 4. Primers used for RT- qPCR of *P. falciparum* MSP2, MSP5, MSP4 and controls.

Name	Sequence
FrucBiAld 3D7 F qPCR [a, b]	TGTACCACCAGCCTTACCAG
FrucBiAld 3D7 R qPCR [a,b]	TTCCTTGCCATGTGTTCAAT
MSP2 3D7 F qPCR	TCCTACTGCACAACCTGAACAA
MSP2 3D7 R qPCR	ATGTCCATGTTGTCCTGTACCTT
MSP4 3D7 F qPCR	CAAAAGAATCCCAAATGGTTGATGATAA
MSP4 3D7 R qPCR	AACATGGCCACCTGAATTTGAT
MSP5 3D7 F qPCR	AAATTAATGAGAATGCAGAAATAGGTCAA
MSP5 3D7 R qPCR	CGCTATAATGTGGCACCTCAT
SUB1 3D7 F qPCR[c]	GGAATGAGGTAGATGCCGATGAA
SUB1 3D7 F qPCR[c]	TCCTTTACATCTTTCAGTTCCCTCAT

[a] FrucBisAld primers had been previously validated as an appropriate housekeeping gene by Salanti et. al. (2003). Primers listed here are the fructose-bisphosphate aldolase qPCR primers (primer pair 61) from Salanti et. al. (2003).

[b] fructose-bisphosphate aldolase is ubiquitously expressed throughout the parasite lifecycle and is used as a non-stage specific housekeeping gene.

[c] SUB1 is a schizont expressed protein and is used as a schizont-stage specific housekeeping gene.