An abundant merozoite surface protein of *Plasmodium falciparum* modulates
susceptibility to inhibitory antibodies

Isabelle G. Henshall[1], Jill Chmielewski[1], Dimuthu Angage[2], Ornella Romeo[1], Keng Heng -
Lai[1, 3], Kaitlin R. Turland[1, 3], Nicki Badii[2], Michael Foley[2, 4], Robin F. Anders[2], James Beeson[5,
6,7] and Danny W Wilson[1, 3, 5, *].

[1]Research Centre for Infectious Diseases, School of Biological Sciences, The University of
Adelaide, Australia 5005

[2]Department of Biochemistry and Chemistry, La Trobe Institute for Molecular Sciences, La
Trobe University, Victoria, Australia 3086

[3]Institute for Photonics and Advanced Sensing (IPAS), University of Adelaide, 5005, SA,
Australia

[4]AdAlta, 15/2 Park Drive, Bundoora, Victoria, Australia 3083

[5]Burnet Institute, 85 Commercial Road, Melbourne 3004, Victoria, Australia

[6]Department of Infectious Diseases, University of Melbourne, Victoria, Australia

[7] School of Translational Medicine and Department of Microbiology, Monash University, Victoria,
Australia

*Corresponding Author: danny.wilson@adelaide.edu.au

Abstract

Malaria merozoite surface proteins (MSPs), are thought to have important roles in red blood
cell (RBC) invasion and their exposure on the parasite surface makes them attractive vaccine
candidates. However, their role in invasion has not been directly demonstrated and their
biological functions are unknown. One of the most abundant proteins is *Pf*MSP2, which is
likely an ancestral protein that has been maintained in the *Plasmodium falciparum* lineage and
is a focus of vaccine development, whose function remains unknown. Using CRISPR-Cas9
gene-editing, we removed *Pf*MSP2 from two different *P. falciparum* lines with no impact on
parasite replication or phenotype *in vitro*, demonstrating that it is not essential for RBC

invasion. However, loss of *Pf*MSP2 led to increased inhibitory potency of antibodies targeting

other merozoite proteins involved in invasion, particularly *Pf*AMA1. In a solid-phase model,

increasing concentrations of *Pf*MSP2 protein reduced binding of different antibodies against

*Pf*AMA1 in a dose dependent manner. These data suggest that *Pf*MSP2 can modulate the

susceptibility of merozoites to protective inhibitory antibodies. The results of this study change

our understanding of the potential functions of *Pf*MSP2 and establishes a new concept in

malaria where a surface protein can reduce the protective efficacy of antibodies targeting a

different antigen. These findings have important implications for understanding malaria

immunity and informing vaccine development.

Introduction

Plasmodium falciparum malaria remains a global health challenge with over half a million

deaths annually (World Health Organization, 2024). Worryingly, increasing parasite resistance

to frontline antimalarials and mosquito insecticide resistance are decreasing the effectiveness

of key control measures (World Health Organization, 2024). Development of the RTS,S and

R21 vaccines targeting the *P. falciparum* circumsporozoite protein is a step forward for malaria

vaccine development, however further improvements in vaccine efficacy and longevity are

needed to protect those most at risk – children under 5 years – and enable malaria elimination

(Beeson et al., 2019; Datoo et al., 2024; RTSS Clinical Trials Partnership, 2015).

Understanding the function of vaccine targets remains important for prioritisation and

optimisation of candidates in the development pipeline. Merozoites, the parasite stage that

invades host red blood cells (RBCs), have long been targeted for vaccine development since

successful blocking of merozoite invasion would prevent the repeated cycles of blood-stage

parasite multiplication and associated disease. While much work has been done to understand

the processes and steps required for merozoite invasion, only a few merozoite proteins have

successfully had functions defined. To date, most vaccines based on merozoite antigens have

demonstrated limited efficacy (Beeson et al., 2016) at least in part due to the extensive

polymorphisms exhibited by many of these antigens.

The surface of the merozoite is covered in a fibril coat which is comprised of multiple proteins,

broadly known as merozoite surface proteins (MSPs). These proteins have been proposed to

mediate weak initial merozoite RBC contact (Bannister et al., 1986; Beeson et al., 2016;

Cowman et al., 2017; Weiss et al., 2015) and in *P. falciparum* are dominated by GPI-anchored

proteins, the most abundant being *Pf*MSP1 and 2 (Gilson et al., 2006). *Pf*MSP1 has been

suggested to have a role in merozoite attachment to the RBC (Li et al., 2004), but recent

evidence instead suggests that it has a role in merozoite rupture from schizonts and cellular

interaction studies using optical tweezers do not support a substantial role for this protein in

RBC binding (Das et al., 2015; Kals et al., 2024), although it may have another function in

invasion. These recent insights into *Pf*MSP1 function, the best studied MSP, highlight what

little is known of MSPs and their roles in parasite survival.

P. falciparum merozoite surface protein 2 (*Pf*MSP2), a protein proposed to be essential for

parasite growth (Sanders et al., 2006), has been of long-term interest as a vaccine candidate.

*Pf*MSP2 is a ~28 kDa intrinsically disordered protein that has conserved N- and C- termini,

along with a central variable region that defines two main allelic groups, 3D7-like and FC27-

like (Adda et al., 2012). While *Pf*MSP2 was theorised to have a mechanical role in the early

steps of invasion (Anders et al., 2010), there is minimal supporting evidence for this, in part

due to the lack of tools to study *Pf*MSP2 function. Studies with recombinant proteins suggest

that it is likely *Pf*MSP2 interacts with lipids and forms complexes with itself on the merozoite

surface (Adda et al., 2009; Lu et al., 2019; Zhang et al., 2012). *Pf*MSP2 peptides have also

been reported to bind the RBC and inhibit *P. falciparum* growth, which would support *Pf*MSP2

mediating merozoite-RBC interactions (Ocampo et al., 2003). Unlike the majority of the

merozoite surface coat proteins, *Pf*MSP2 is not present in other malaria parasites that infect

humans and has been postulated to have evolved specifically in the *Laverania* subgenus of

Plasmodium, which includes *P. falciparum* (Black et al., 2002).

*Pf*MSP2 has been trialled in a vaccine (Combination B) combined with fragments of *Pf*MSP1

and *Pf*RESA. A Phase 1/2b trial of this vaccine showed efficacy in reducing the parasite burden

in children, which was specific to the *Pf*MSP2 3D7-like allele used in the vaccine formulation

(Genton et al., 2003). Vaccine-induced antibodies from clinical trials or experiments in animals

have little or no inhibitory activity in standard growth inhibition assays (Boyle et al., 2015;

McCarthy et al., 2011), but did show functional activity through the recruitment of

complement, promoting opsonic phagocytosis and antibody-dependent cellular inhibition

(Boyle et al., 2015; Feng et al., 2018; McCarthy et al., 2011). Naturally acquired antibodies are

also known to target *Pf*MSP2 and have been linked to protection (Fowkes et al., 2010; Osier et

al., 2010; Stanisic et al., 2009; Zerebinski et al., 2024). Studies have highlighted that it is the

Fc-mediated functional activities of these naturally acquired *Pf*MSP2 antibodies, as opposed

to direct blocking of *Pf*MSP2 protein function, that correlates best with protection (Boyle et

al., 2015, 2014; Osier et al., 2014; Reiling et al., 2019). While *Pf*MSP2 is a key target of

naturally acquired malaria immunity, little is known about the function of *Pf*MSP2, hindering

its advancement as a potential vaccine candidate.

Here we take a reverse genetics approach to assess the function of *Pf*MSP2, quantifying the

impacts of *Pf*MSP2 deletion on parasite invasion and phenotype. Given our findings that

deletion of MSP2 did not impact invasion *in vitro*, we tested the alternative hypothesis that

merozoite surface proteins may function to modulate susceptibility of merozoites to inhibitory

antibodies.

Results

Avian malaria MSP2-like proteins are structurally similar to *Laverania* MSP2s and show

that MSP2 did not arise exclusively in this clade.

Genome annotations of *pfmsp2* show it lies between the merozoite surface protein *pfmsp5*

(PF3D7_0206900) and *pfmsp*4 (PF3D7_0207000), and the conserved purine biosynthesis

enzyme *adenylosuccinate lyase* (ADSL, PF3D7_0206700) (Figure 1A) (PlasmoDB;

(Aurrecoechea et al., 2009)), with this arrangement previously thought to be restricted to the

Laverania clade which includes parasites that infect great apes and *P. falciparum* (Black et al.,

2002; Ferreira et al., 2015). As described by Escalante et al. (2022), we found the genomes of

the two annotated avian *Plasmodium* spp. available on PlasmoDB, *P. relictum* and *P.*

gallinaceum, to also have a putative gene between ADSL and MSP5 with structural similarities

to *Laverania* MSP2s. The reduced length of the *P. relictum* (PRELSG_0415300) and *P.*

gallinaceum (PGAL8A_00017700) MSP2-like proteins compared to those of *Laverania*

parasites is accounted for by the absence of any extended central domains of the two avian

malaria parasites (Supplementary Figure 1).

Given the estimated that ~10 million years of evolution separate the avian malaria parasites *P.*

relictum and *P. gallinaceum* from the *Laverania* (Böhme et al., 2018), we explored whether

similarities in amino acid sequence and predicted conformation would support a shared lineage

and functional properties of these putative MSP2s. Although *Pf*MSP2 is generally considered

a disordered protein with limited conservation, the hydrophobic N- and C-terminal sequences
that correspond to secretion signals (SP) and GPI-attachment signals, as well as short regions
downstream and upstream of these signal sequences respectively, show significant
conservation across *P. falciparum* MSP2s. Here we compared the sequence similarity of these
conserved regions between the *Laverania* and avian MSPs and found consistently high levels
of conservation for the putative SP (>89% similarity) (Supplementary Figure 1), the 24 amino
acid N-terminal conserved region (78% similarity) that has a propensity for alpha-helical
structure and fibril formation (Yang et al., 2010) (Supplementary Figure 2), the 50 amino acid
residue C-terminal conserved region (62% similarity) (Supplementary Figure 3) and the C-
terminal GPI signal sequence (>79%), including the presence of short side chain amino acids
that mark predicted GPI cleavage sites (Eisenhaber et al., 1999)(Smythe et al., 1991)(Gilson et
al., 2006) (Supplementary Figure 1). *Laverania* MSP2s contain two C-terminal cysteine
residues that form an intramolecular disulphide bond (Zhang et al., 2008). This arrangement
of 2 C-terminal cysteines is found in *P. gallinaceum* MSP2, but not *P. relictum* which has only
one (Supplementary Figure 3). Overall, these N and C-terminal conserved regions show broad
conservation in properties across the *Laverania* and avian malarias which could contribute to
a conserved sub-cellular localisation.

Characteristic of *P. falciparum* MSP2s is the dimorphic central variable repeat region that are
grouped as 3D7 or FC27-like amino acid sequence structures. A defining feature of the 3D7-
like MSP2 family are 4 or 8-mer tandem repeats consisting of small amino acids (for 3D7: Gly,
Gly, Ser, Ala) (MacRaild et al., 2015; Smythe et al., 1991). The FC27-like MSP2 family also
has repeats of 12 or 32 amino acids in length, but these do not resemble those of the 3D7-like
family. We found that the non-*P. falciparum Laverania* and avian malaria MSP2 sequences
also contained >75% small amino acids (some of: Gly, Ser, Ala, Thr) repeats, but the exact
number (4 to 8 repeats), length (3 to 7 amino acids) and the level of conservation between
repeats differed between and within species (Supplementary Figure 4). Broadly speaking, these
Laverania and avian malaria MSP2 repeats had a 3D7-like repeat structure, with no example
of a FC27-like repeat structure identified even with an additional nine *Laverania* MSP2s from
the NCBI experimental database compared (data not shown). This comparison highlights the
surprising finding that a region of known dimorphic variability in *P. falciparum* MSP2 is likely
to retain functional constraints that favour small amino acid rich repeats immediately
downstream of the N-terminal conserved region.

Extensive studies on recombinant proteins of the *P. falciparum* 3D7 and FC27 variants of

*Pf*MSP2 have shown this merozoite surface protein to be intrinsically disordered. Given the

evidence for conservation of the primary structure of MSP2s from distantly related malaria

parasites, we used AlphaFold2 predictions (Jumper et al., 2021) to examine whether there were

likely to be also conformational similarities between *Laverania* MSP2s and those of *P. relictum*

and *P. gallinaceum*. The MSP2s in the database of AlphaFold2-predicted structures retain their

N- and C-terminal signal sequences, but as these are both absent from mature MSP2 on the

merozoite surface they were deleted from the MSP2 sequences used for the AlphaFold2

predictions reported here. The predicted structures of the two avian parasite MSP2s, and other

Laverania MSP2s, are very similar to that of *Pf*3D7 MSP2, with most of the polypetide chains

lacking predicted secondary structure and, from the colour coding, have a predicted local

distance test (pLDDT) consistent with that of an intrinsically disordered protein over the

majority of the proteins length (Figure 1B). In contrast to the rest of the polypeptide, some α-

helical structure was predicted in the conserved 20-residue N-terminal region of all the

Laverania and avian MSP2s. NMR studies with recombinant *Pf*3D7 and *Pf*FC27 MSP2

showed this conserved N-terminal sequence to have a propensity for forming an α-helix that

was stabilized in the presence of lipid mimetics (MacRaild et al., 2012). The low pLDDT

values are consistent with this region in the other *Laverania* and *P. gallinaceum* MSP2 having

some propensity for α-helix formation, whereas higher pLDDT values suggest the possible

presence of a more stable α-helix in this region of *P. relictum* MSP2.

Having established that MSP2 is present across different *Plasmodium* lineages sharing similar

predicted amino acid compositions and structural features, the phylogenetic relationship

between annotated *Laverania* and putative avian malaria parasite MSP2s was examined. A

phylogenetic tree was generated based on alignments of representative sequences from each

species. MSP2 sequences from species belonging to the *Laverania* subgenus all grouped

together with two smaller clusters, corresponding to the known *Laverania* clade A and clade

B species (Figure 1C). As expected, *P. relictum* and *P. gallinaceum* MSP2 also group together.

The phylogenetic tree generated based on MSP2 sequences mirrors the proposed evolution of

Plasmodium spp. (Figure 1C). As postulated by Escalante et al. (2022) , these data suggest that

MSP2 was likely present in the ancestral *Plasmodium* parasite prior to the split of the

mammalian infecting *Plasmodium* spp. from those that infect birds and lizards. Given the

shared amino acid sequence and predicted conformational similarities, including the high levels

of intrinsically disordered sequence outside of the N- and C-terminal regions, the *P. relictum*

and *P. gallinaceum* MSP2-like proteins appear likely to contain enough properties to
potentially be functionally equivalent to *Laverania* MSP2s. This conservation of MSP2-like
proteins across different species suggests it has an important function.

Figure 1. Presence and structure of MSP2 across different *Plasmodium* species. **(A)** Schematic of the gene arrangement surrounding *msp2* on *P. falciparum* chromosome 2. **(B)** AlphaFold2 structural predictions of example *Laverania* (*P. falciparum* (3D7), *P. reichenowi*, *P. billcollinsi*, *P. adleri*) and the two avian (*P. relictum*, *P. gallinaceum*) malaria species. The N-terminal signal peptide and C-terminal GPI anchor sequences were removed before the proteins structure was predicted. The most N-terminal amino acid is indicated. Colours represent the predicted local distance difference test (pLDDT) scores, with dark blue representing very high confidence (>90%), light blue high confidence (90 to >70), yellow low confidence (70 to 50) and orange very low confidence (<50). An enlarged modelled structure of the N-terminal helical region is provided for *P. falciparum* 3D7 MSP2 on the left of the panel (grey shading), and also for the other examples of *Laverania* and avian MSPs (various colours) superimposed on the structure of *P. falciparum* 3D7 MSP2 (green). **(C)** Maximum likelihood tree showing relationship of MSP2 protein sequences found in different *P. falciparum* isolates and other *Plasmodium* species. Grouping of sequences from *P. falciparum* isolates into two main allele types can be seen as well as the separation of the *Laverania* and avian malaria species into their expected groups. Tree robustness tested by bootstrap.

Loss of *Pf*MSP2 does not noticeably impact growth or invasion of different *P. falciparum* lines *in vitro*.

Given previous unsuccessful attempts to disrupt *pfmsp2* (Sanders et al., 2006), and its high abundance on the merozoite surface (Gilson et al., 2006), *Pf*MSP2 has been traditionally viewed as an essential *P. falciparum* protein with an essential function in merozoite invasion. We employed CRISPR-Cas9 gene editing to determine whether the reported inability to knock-out *pfmsp2 in vitro* was a result of this protein being essential for parasite survival or because of the lower efficiency of the previously used gene-editing system (Sanders et al., 2006). Unexpectedly, we confirmed successful disruption of *pfmsp2* by replacing the coding sequence between 132 bp and 819 bp of the gene with a hDHFR drug selection cassette in the 3D7 *P. falciparum* laboratory-adapted line (Figure 2A and B), resulting in *Pf*3D7 ΔMSP2 parasites. We confirmed that *Pf*MSP2 was no longer expressed in *Pf*3D7 ΔMSP2 parasites by western blot, which detected a MSP2 band around 50 kDa in *Pf*3D7 WT parasite material but not in *Pf*3D7 ΔMSP2 parasites (Figure 2C). Similarly, immunofluorescence microscopy using anti-*Pf*MSP2 rabbit polyclonal antibodies showed the expected surface localisation of MSP2 on

228 *Pf*3D7 WT merozoites but not on *Pf*3D7 ΔMSP2 parasites (Figure 2D). We next assessed

229 whether knock-out of *Pf*MSP2 impacted on parasite growth over one or two cycles of

230 development *in vitro* and found that deletion of *Pf*3D7 MSP2 did not cause any significant

231 change in growth (Figure 2E and F).

Figure 2. *Pf*MSP2 is not essential for growth *in vitro* of *Pf*3D7 blood stage parasites. **(A-B)** Schematic and agarose gel showing integration of knockout construct (band A+C) in the *msp2* gene locus and absence of the *msp2* sequence from the *Pf*3D7 ΔMSP2 line (band A+B). **(C)** Western Blot of late schizont protein extracts confirms no *Pf*MSP2 is expressed in the *Pf*3D7 ΔMSP2 line. *Pf*MSP2 detected by anti-*Pf*MSP2 2F2 3D7 mAb with *Pf*Aldolase (upper blot) or *Pf*GAP45 (lower blot) serving as loading and stage of expression controls respectively. Representative image shown. **(D)** Distribution of key merozoite surface proteins in the presence or absence of *Pf*MSP2 was visualised by immunofluorescence. *Pf*MSP2 (magenta), the nucleus stained by DAPI (cyan) and *Pf*AMA1 (yellow, top two rows) or *Pf*MSP1-19 (yellow, bottom two rows), and the coloured merge of the preceding panels. Scale bar = 0.7 µm. Representative images shown from a minimum of 10 schizonts imaged per condition. **(E-F)** Growth of *Pf*3D7 WT compared to *Pf*3D7 ΔMSP2 *P. falciparum* parasites, measured as fold increase in parasitaemia, over one (48 hrs) or two (96 hrs) cycles in either standard (still- (E)) or shaking (F) conditions, with no measurable difference between parasite growth rates seen between standard or shaking conditions. Parasitaemia was determined by flow cytometry at the start and end to calculate fold increase. Graph displays mean ± S.D. of three independent experiments performed with technical triplicates. Significance determined by unpaired t-test with p< 0.05 deemed significant.

As *Pf*MSP2 has been considered, to this point, as essential for *P. falciparum* growth *in vitro* (Sanders et al., 2006), we investigated whether *Pf*MSP2 could also be removed from *Pf*Dd2, an isolate of *P. falciparum* that differs from 3D7 in geographical origin, RBC receptor usage and allelic type of *pfmsp2*. Using CRISPR-Cas9 we successfully knocked-out *msp2* in *Pf*Dd2 (Figure 3 A and B) and confirmed the absence of the protein in *Pf*Dd2 ΔMSP2 parasites using western blot (Figure 3C). As we found for *Pf*3D7, deletion of *Pf*Dd2 *msp2* did not result in a significant impact on parasite growth over one or two cycles of parasite replication (Figure 3D).

Given the abundance of *Pf*MSP2 on the merozoite surface, we postulated that loss of this protein may impact on the timing of merozoite invasion of the RBC, which is not reflected in reduced growth in *in vitro* cultures. Previous studies using live-cell microscopy have reported that merozoites first contact the RBC and form weak initial interactions. The merozoite then

reorientates to bring apical organelles into position to release their contents onto the RBC,
which then leads to formation of the tight junction, merozoite invasion through the tight-
junction, formation of the parasitophorous vacuole and resealing of the RBC membrane post-
entry (Weiss et al., 2015). Given its surface localization, *Pf*MSP2 has been speculated to act in
the early phase of merozoite attachment. To assess whether knock-out of *pfmsp2* impacted on
the progression of invasion, we used live-cell microscopy of *Pf*Dd2 ΔMSP2 parasites and
compared the timing and key steps of invasion to *Pf*Dd2 WT parasites *in vitro*. This analysis
showed that, although there was a trend for *Pf*Dd2 ΔMSP2 knock-out parasites to have a higher
mean time to attach to the RBC, as well as for the length and strength of RBC deformation,
these trends did not reach significance. For those merozoites that did invade the RBC, on
average it took less time for *Pf*Dd2 ΔMSP2 knock-out parasites to invade then *Pf*Dd2 WT, but
this again did not reach significance (Figure 3 E-H). Together these data show *Pf*MSP2 is not
essential for blood-stage replication *in vitro* in two *P. falciparum* laboratory isolates from
different geographical regions and knock-out of *Pf*MSP2 does not seem to significantly impact
parasite growth or merozoite invasion *in vitro*. However, its conservation across diverse
Plasmodium spp. suggests it does play an important function or is advantageous for parasite
survival.

Figure 3. *Pf*Dd2 does not require MSP2 for asexual growth *in vitro*. **(A-B)** Successful

integration of KO construct (schematic in A) into the *msp2* gene locus of *Pf*Dd2 ΔMSP2 was

confirmed by PCR of genomic DNA (primers A+B amplify WT locus, primers D+E amplify

integrated KO construct). **(C)** Loss of *Pf*MSP2 expression was demonstrated by western blot

of schizont protein extract with *Pf*MSP2 detected by anti-*Pf*MSP2 FC27 and anti-*Pf*EXP2 as

loading control. Representative image shown. **(D)** Growth of *Pf*Dd2 WT *P. falciparum*

parasites and *Pf*Dd2 ΔMSP2 over one (48 hrs) or two (96 hrs) cycles. Parasitaemia was

determined by flow cytometry at the start and end to calculate fold increase. Graph displays

mean ± S.D. of three independent experiments performed with technical triplicates. **(E-H)**

Key parameters of merozoite invasion were measured for both *Pf*Dd2 WT and *Pf*Dd2

ΔMSP2 parasites that had successfully invaded a RBC using live cell imaging of merozoite

invasion. Time to merozoite attachment to RBCs **(E)**, length **(F)** and strength **(G)** of RBC

deformation, and time to complete merozoite invasion **(H)** were measured by live

microscopy. Deformation scores are as defined by Weiss et al (2015), with 1 = weak

deformation of the RBC membrane at the point of contact, 2 = strong deformation leading to

the RBC membrane extending up the sides of the merozoite and changes in RBC membrane

curvature beyond the point of contact and 3 = extreme deformation indicated by the

merozoite being deeply embedded in the RBC membrane and strong deformation of the RBC

well beyond the point of contact. A minimum of 30 invading merozoites for each line were

captured and assessed. Significance determined by unpaired t-test with $p < 0.05$ deemed

significant.

Impact of MSP2 gene deletion on gene expression and invasion pathway usage.

Because of the close proximity of the *pfmsp*2, 4 and 5 genes on chromosome 2, similarities in

their structure (e.g. GPI-anchor intrinsically disordered protein) and shared merozoite surface

localization, we used qPCR to assess whether knock-out of *Pf*3D7 MSP2 impacted the

expression levels of *Pf*3D7 MSP4 or MSP5. We found no change in MSP4 and MSP5

expression levels between *Pf*3D7 ΔMSP2 parasites and *Pf*3D7 WT parasites (Figure 4A). As

expected, *Pf*MSP2 was not detected in the *Pf*3D7 ΔMSP2 parasites.

Secreted *P. falciparum* antigens belonging to the erythrocyte binding antigen (EBA) and

reticulocyte binding homologue (Rh) families are known to facilitate host-cell invasion through

specific receptors on the RBC surface. Reliance on specific EBA or Rh proteins, or their

cognate host-cell receptor, define different invasion pathways that can be crudely described by

the sensitivity of merozoite invasion to cleavage of RBC surface proteins with trypsin,

chymotrypsin and neuraminidase (Baum et al., 2005; Duraisingh et al., 2003; Orlandi et al.,

1992).When we assessed whether *Pf*MSP2 knock-out impacted merozoite invasion pathway

usage, we found that *Pf*3D7 ΔMSP2 parasites grew equally as well as *Pf*3D7 WT parasites in

all enzyme-treated RBCs (Figure 4B).

To investigate whether *Pf*MSP2 knock-out caused any changes in gene-expression of proteins

that may have a role in merozoite invasion and possibly compensate for loss of *Pf*MSP2, we

undertook RNA sequencing and differential gene expression analysis of *Pf*3D7 ΔMSP2 and

*Pf*3D7 WT parasites. As expected, the *Pf*MSP2 transcript, which was targeted for removal by

our CRISPR-Cas9 strategy, was not detected in the *Pf*3D7 ΔMSP2 parasites between 132 bp

and 819 bp (Supplementary Figure 5). After removal of genes belonging to variant surface

antigen families from the data, eight transcripts were found to have a log2fold expression

increase above 1; none of these proteins are annotated as a merozoite surface or secreted protein

linked to merozoite invasion (Figure 4C, Table 1). Five of these eight upregulated proteins are

of unknown function and expressed at schizont stages, and only one (*Pf*3D7_0909100) has a

transmembrane domain that could possibly anchor it to a membrane surface as the GPI anchor

does for *Pf*MSP2 (Table 1). Three proteins were found to have a log2fold expression of <-1,

indicating a significant reduction in expression with *Pf*MSP2 knock-out, with *Pf*MSP2 being

one of these and the other two predicted to be exported proteins (Figure 4C, Table 1). While

these data do not exclude any of the *Plasmodium* proteins with unknown function that are

upregulated in *Pf*3D7 ΔMSP2 parasites as having a role that compensates for the loss of

*Pf*MSP2, only one is predicted to have a transmembrane domain that would bind it to a

membrane and no protein identified as up or down-regulated with *Pf*MSP2 knock-out has

previously been linked to merozoite invasion. Therefore, *Pf*MSP2 knock-out does appear to

have led to changes in gene expression, but does not appear to have a significant impact on the

regulation of known merozoite surface or secreted protein gene-expression that would suggest

loss of *Pf*MSP2 is compensated for by another protein.

Figure 4. The impact of the loss of *Pf*MSP2 on expression of known merozoite invasion genes
and invasion pathway utilization. **(A)** Impact of *Pf*MSP2 KO on schizont transcript abundance
was assessed by qPCR for genes located in proximity to *Pfmsp2* on chromosome 2. Changes
in expression between *Pf*3D7 WT and *Pf*3D7 ΔMSP2 parasites was determined by qPCR

relative to *pfaldolase* expression with *pfsub1* serving as a schizont stage control. Graph displays

mean ± S.D. of three independent RNA harvests. **(B)** Selective enzymatic cleavage of key RBC

receptors showed no difference in invasion preference between *Pf*3D7MSP2 WT and *Pf*3D7

ΔMSP2 parasites. Parasitaemia was determined by flow cytometry and compared to growth in

non-treated control RBCs. Graph displays mean ± S.D. of three independent experiments. **(C)**

Log2(fold change) for differentially expressed genes, including multigene families, between

the transcriptome of *Pf*3D7 WT and *Pf*3D7 ΔMSP2 schizonts. Plot represents the results for

one of four independent schizont RNA harvests for *Pf*3D7 WT and *Pf*3D7 ΔMSP2 parasites

and red lines differentiate genes with a log2 (fold change) > 0.5 and < -0.5 with adjusted p-

value < 0.01. Genes shaded blue represent those genes that were found to have an average log2

(fold change) > 0.5 (dark blue) or < -0.5 (light blue) across the four replicate samples compared.

Significance determined as below p< 0.05 after correction for multiple testing.

Impact of *Pf*MSP2 disruption on antibodies and inhibitors targeting secreted surface

exposed antigens

Merozoites are key targets of antibodies during malaria infection with antibodies either able to

directly block protein function and thus merozoite invasion, or recruit effectors such as

complement leading to the destruction of the merozoite (Beeson et al., 2016). Given the

prominence of *Pf*MSP2 on the merozoite surface we asked the question; how does the loss of

*Pf*MSP2 affect antibody efficacy against merozoites?

*Pf*EBAs and *Pf*Rhs moderate early steps in invasion with a degree of redundancy between these

proteins. Several EBAs and Rhs can be targeted with invasion inhibitory antibodies and are

targets of acquired human antibodies (Persson et al., 2013, 2008; Richards et al., 2013). We

found no significant difference in the ability of purified rabbit immunoglobulin raised against

single (*Pf*Rh2b) or combinations of *Pf*EBAs and *Pf*Rhs (*Pf*EBA175/*Pf*Rh2b,

*Pf*EBA175/*Pf*Rh4, *Pf*EBA175/*Pf*Rh2b/*Pf*Rh4) to inhibit growth of *Pf*3D7 WT or *Pf*3D7

ΔMSP2 parasites (Figure 5A). Similarly, knock-out of MSP2 in *Pf*Dd2 did not significantly

change the parasites sensitivity to rabbit antibodies raised against isogeneic W2mef EBA175

(Figure 5B). Broadly speaking, these results mirror what was seen with selective cleavage of

the RBC ligands of *Pf*EBAs and *Pf*Rhs using enzyme treatment of RBCs (Figure 4B), with no

evidence that loss of *Pf*MSP2 changed either the importance of these secreted merozoite

antigens during invasion or their sensitivity to antibodies targeting different antigens.

The interaction of *Pf*Rh5 with RBC surface receptor basigin is a critical, non-redundant interaction for *P. falciparum* merozoite invasion and a known target of direct invasion-inhibitory antibodies (Alanine et al., 2019; Crosnier et al., 2011; Douglas et al., 2011). *Pf*Rh5 was the initial member of the PCRCR complex to be described, with subsequent studies demonstrating that antibodies against other members of this complex could also block merozoite invasion of the RBC (Scally et al., 2022). Given the importance of these secreted antigens to current vaccine development efforts, we next investigated whether loss of *Pf*MSP2 impacted on the parasite's sensitivity to antibodies targeting these antigens. There was no difference in sensitivity between *Pf*3D7 WT and *Pf*3D7 ΔMSP2 parasites when treated with invasion-inhibitory rabbit antibodies targeting *Pf*Rh5 (Figure 5C) or nanobodies targeting PCRCR complex component *Pf*PTRAMP (H8) (Figure 5D). Nanobodies targeting the PCRCR complex component CSS (D2) showed a slight, but non-significant increase in invasion-inhibitory activity in the absence of *Pf*MSP2 (1.4-fold increase; IC_{50} value for *Pf*3D7 WT is 57 μg/mL; IC_{50} value for *Pf*3D7 ΔMSP2 is 42 μg/mL; p=0.3; Figure 5E). However, the *Pf*3D7 ΔMSP2 specific increase in invasion-inhibitory activity was amplified with the addition of the human Fc domain to the anti-CSS nanobody (D2-Fc) to be 3-fold greater for *Pf*3D7 ΔMSP2 parasites (IC_{50} *Pf*3D7 WT 138 μg/mL; IC_{50} *Pf*3D7 ΔMSP2 46 μg/mL; p=0.0001; Figure 5F).

The glycosaminoglycan heparin is a specific inhibitor of invasion and has been observed to bind to the 42 kDa C-terminal region of *Pf*MSP1 the most abundant GPI-anchored protein on the merozoite surface, marking this as a possible mechanism for its invasion inhibitory activity (Boyle et al., 2010a). However, the highly charged state of heparin means that it could block invasion through binding additional proteins. When tested against *Pf*3D7 WT and *Pf*3D7 ΔMSP2 parasites, we observed a small increase in inhibitory potency of heparin for parasites lacking *Pf*MSP2 (1.4-fold; IC_{50} *Pf*3D7 WT 0.65 IU/mL; IC_{50} *Pf*3D7 ΔMSP2 0.47 IU/mL; p=0.0003; Figure 5G). We next tested rabbit antibodies raised against the C-terminal 19 kDa fragment of *Pf*MSP1. We found a small but non-significant trend towards increased inhibition of growth in parasites that lacked *Pf*MSP2 across repeat experiments (Figure 5H). These data suggest that loss of *Pf*MSP2 can impact on the invasion-inhibitory potency of some antibodies that target secreted or surface antigens.

Figure 5. Impact of *Pf*MSP2 removal on efficacy of antibodies targeting other merozoite

surface-exposed antigens. Changes in antibody efficacy in the absence of *Pf*MSP2 was

assessed by measuring changes in antibody invasion inhibition and subsequent growth

compared to growth in the absence of antibody for both *P. falciparum* WT and ΔMSP2

parasites over 2 cycles. **(A)** Rabbit (Rb) IgG raised against merozoite antigens of the *Pf*3D7

EBA/Rh family. **(B)** Rabbit sera raised against *Pf*Dd2 EBA175. **(C)** Rabbit IgG raised against

*Pf*3D7 Rh5. **(D)** Nanobody (nAb) to *Pf*3D7 PTRAMP. **(E-F)** Nanobody and Fc-tagged

nanobody to *Pf*3D7 CSS. **(G)** The invasion inhibitory glycosaminoglycan heparin. **(H)** Rabbit

sera raised against *Pf*3D7 MSP1-19 (different vaccinated rabbit sera identified by numbers).

Graph displays mean ± S.D. of three different experiments. Significance was determined by

unpaired t-test when only a single concentration point was tested and for IC_{50} comparisons an

extra Sum-of-Squares F Test (best-fit $LogIC_{50}$) was performed with $p < 0.05$ deemed

significant.

Loss of *Pf*MSP2 consistently potentiates AMA1 invasion inhibitory antibodies

Subsequent to *Pf*Rh5 -basigin binding in the steps of merozoite invasion is the formation of the

tight-junction between *Pf*AMA1, secreted from the micronemes onto the merozoite surface,

and the rhoptry neck protein *Pf*RON2, which is inserted into the RBC membrane (Lamarque

et al., 2011; Srinivasan et al., 2011). In lines where *Pf*MSP2 was disrupted, we found a

consistent potentiation of antibodies that had anti-*Pf*AMA1 invasion-inhibitory activity.

Invasion-inhibitory polyclonal rabbit serum antibodies and purified immunoglobulin raised

against *Pf*AMA1 of *Pf*3D7 and *Pf*W2mef (isogenic to *Pf*Dd2), which express different alleles

of both *Pf*MSP2 and *Pf*AMA1, were found to be more inhibitory with *Pf*MSP2 knock-out

(Figure 6A and B). The polyclonal antibody with the greatest potentiation of inhibitory activity

was an IgG preparation purified from a rabbit antiserum raised against *Pf*W2mef AMA1. This

had a 3-fold increased potency against *Pf*Dd2 ΔMSP2 compared to *Pf*Dd2 WT (IC_{50} *Pf*Dd2

WT 0.015 mg/mL; IC_{50} *Pf*Dd2 ΔMSP2 0.005 mg/mL; $p < 0.0001$) over two cycles of growth

(Figure 6B, C).

The mouse 1F9 (binds to the RON2 binding pocket and polymorphic loop 1d (Coley et al.,

2007)) and rat 4G2 (binds to a conserved epitope in the ectodomain (Kocken et al., 1998))

monoclonal antibodies have been shown to target different regions of *Pf*AMA1 and exhibit

selective inhibition against *Pf*3D7 AMA1. Here we found potentiation of invasion inhibitory

activity with *Pf*3D7 MSP2 knock-out for both 1F9 (2.1-fold; IC_{50} *Pf*3D7 WT 0.27 mg/mL; IC_{50}

*Pf*3D7 ΔMSP2 0.13 mg/mL; $p < 0.0001$; Figure 6D) and 4G2 (3.2-fold; IC_{50} *Pf*3D7 WT 1.15

mg/mL; IC_{50} *Pf*3D7 ΔMSP2 0.36 mg/mL; $p < 0.0001$; Figure 6E).

We also tested the invasion inhibitory effect of i-bodies, which are smaller single-domain

antibody-like molecules inspired from the shark variable new antigen receptor (V_{NAR})

(Griffiths et al., 2016a, 2018). When we tested the i-body WD34 (Angage et al., 2024) which

binds a highly conserved epitope that includes the *Pf*RON2-binding pocket on *Pf*AMA1, we

observed a small potentiation of *Pf*AMA1 specific activity with knock-out of *Pf*MSP2 in *Pf*3D7

(1.3-fold; IC_{50} *Pf*3D7 WT 0.012 mg/mL; IC_{50} *Pf*3D7 ΔMSP2 0.009 mg/mL; p=0.08 Figure 6F).

Given WD34 was also found to be inhibitory to growth of W2mef parasites (Angage et al.,

2024) we also tested its activity against our *Pf*Dd2 ΔMSP2 parasites and found potentiation of

invasion inhibitory activity with loss of MSP2 in *Pf*Dd2 parasites (2-fold; IC_{50} *Pf*Dd2 WT

0.016 mg/mL; IC_{50} *Pf*Dd2 ΔMSP2 0.008 mg/mL; p=0.004 Figure 6G). As was observed for

the anti-CSS nanobody, addition of a human Fc domain to WD34 amplified the inhibitory

phenotype, with WD34 Fc having a 2.3-fold greater inhibitory activity against *Pf*3D7 ΔMSP2

than against *Pf*3D7 WT (IC_{50} *Pf*3D7 WT 0.1 mg/mL; IC_{50} *Pf*3D7 ΔMSP2 0.04 mg/mL;

p=0.0004; Figure 6H). A second i-body, WD33 (Angage et al., 2024), which binds AMA1

between domain II and domain III, had very limited invasion inhibitory activity against *Pf*3D7

parasites and did not show improved potency with knock-out of *Pf*3D7 MSP2 (0.9-fold; IC_{50}

*Pf*3D7 WT 1.02 mg/mL; IC_{50} *Pf*3D7 ΔMSP2 1.1 mg/mL; p=0.8; Figure 6I). Although the

limited inhibitory activity for WD33 tagged with the Fc receptor prevented us determining an

IC_{50} at the concentrations feasible to test, there was no increased inhibition of *Pf*3D7 ΔMSP2

parasites compared to the low level seen with *Pf*3D7 WT parasites (data not shown). These

data suggest that the potentiation of *Pf*AMA1 targeted antibody inhibition with MSP2 knock-

out is *Pf*AMA1 epitope dependent.

To account for the differences in i-body sizes, we compared the WD34 and WD34 Fc i-bodies

on a molar basis and found that they had similar activities against *Pf*3D7 MSP2 WT parasites

(IC_{50} 0.94 μM and 1 μM respectively) indicating that the Fc-tag itself did not modify epitope

binding properties. Rather, it is the absence of *Pf*MSP2 that results in increased inhibition of

AMA1 function by WD34, and this is further potentiated by antibody size.

Figure 6. Absence of *Pf*MSP2 from the merozoite surface impacts invasion inhibition by *Pf*AMA1 antibodies. *Pf*3D7 (**A, D-F, H, I**) and *Pf*Dd2 (**B, C, G**) express different *Pf*MSP2 alleles and different *Pf*AMA1 alleles, yet both showed altered anti-AMA1 antibody growth inhibition in the absence of *Pf*MSP2. The effect was seen with serum (**A-B**; different vaccinated rabbit (Rb) sera identified by numbers), purified rabbit and mouse monoclonal

(mAb) antibodies **(C-E)** and i-bodies (ibA) **(F-J)**. Final parasitaemia was determined by flow

cytometry and compared to control. Graph displays mean ± S.D. of three or four independent

experiments. Significance was determined by unpaired t-test when only a single concentration

point was tested and for IC_{50} comparisons an extra Sum-of-Squares F Test (best-fit $LogIC_{50}$)

was performed with $p < 0.05$ deemed significant.

In light of these results, we hypothesised that removal of PfMSP2 may improve antibody access

to other antigens that are targets of inhibitory antibodies. To assess whether this was the case,

we first tested whether increased PfAMA1 polyclonal antibody binding to the surface of

merozoites could be observed for Pf3D7 ΔMSP2 compared to Pf3D7 MSP2 WT parasites using

a fluorescently tagged WD34 i-body (WD34-mCherry) for quantitative immunofluorescence

assays of late stage schizonts labelled with a single incubation and wash step. We observed

WD34-mCherry to have a significantly higher mean fluorescence intensity for Pf3D7 ΔMSP2

compared to Pf3D7 WT (Figure 7A), supporting that the PfAMA1 i-bodies may have better

access in the absence of PfMSP2. As a control, we tested labelling using a fluorescently tagged

WD33 i-body which binds to AMA1 (Angage et al., 2024), but we found did not have improved

inhibitory activity in the absence of MSP2 (Figure 6I). Mirroring the results of the growth

assay, there was no difference observed for WD33-eGFP binding fluorescence for Pf3D7

ΔMSP2 compared to Pf3D7 WT parasites (Figure 7B).

We next explored whether the addition of recombinant PfMSP2 could impact on the inhibitory

antibody binding on-rate against substrate-bound PfAMA1 recombinant protein using Surface

Plasmon Resonance (SPR). For the anti-PfAMA1 targeting mAb 4G2, there was a non-

significant trend toward a lower on-rate in the presence of recombinant PfMSP2 (Figure 7C).

For the i-body WD34-Fc, there was minimal difference in detectable binding to PfAMA1 in

the presence or absence of PfMSP2 (Figure 7C). Using a similar principle of testing antibody

binding to substrate-bound PfAMA1 protein in the presence, or absence of increasing

concentrations of PfMSP2, we assessed changes in levels of anti-PfAMA1 antibody binding

using an enzyme linked immunosorbent assay (ELISA). With increasing concentrations of

PfMSP2 added to wells coated with AMA1, we found decreased binding of anti-PfAMA1 mAb

4G2 and i-body WD34-Fc (mouse Fc) by ELISA (Figure 7D). Concentrations where the

PfMSP2 protein was barely detectable above background using an MSP2 mAb were found to

be enough to impact on the binding of PfAMA1 antibodies. In contrast, increasing

concentrations of the intrinsically disordered MSP4 from *P. falciparum* 3D7 (40 kDa) and the

highly structured immunoglobulin domain of neural cell adhesion molecule (NCAM, CD56,

16 kDa) recombinant proteins did not impact on binding of anti-*Pf*AMA1 antibodies to

recombinant AMA1 (Supplementary Figure 6). By looking at the impacts of removing *Pf*MSP2

from the merozoite surface, or adding this protein in to recombinant assays, these data provide

additional support that inhibitory antibodies targeting *Pf*AMA1 bind at a higher rate in the

absence of *Pf*MSP2.

Figure 7. Quantitative fluorescence microscopy to assess whether differential binding may

explain the increased potency of anti-*Pf*AMA1 invasion-inhibitory antibodies in the absence

of *Pf*MSP2. Fluorescence intensity of fluorescently tagged anti-*Pf*AMA1 i-body (ibA) WD34-

mCherry **(A)** and WD33-eGFP **(B)** for both *Pf*3D7 WT and *Pf*3D7 ΔMSP2, with a
representative image for i-body WD34-mCherry. Nucleus in blue, i-body signal in red. Scale
bar = 4 μm. Two independent experiments were performed with significance determined by
unpaired t-test with p< 0.05 deemed significant. The lower overall mCherry signal required a
higher antibody concentration (240 ng/mL) to have a comparable intensity measure to the
eGFP tagged antibody (120 ng/mL) for *Pf*3D7 WT merozoites. **(C)** Read-out of the Surface
plasmon resonance (SPR) antibody on-rate (association constant) for anti-*Pf*AMA1 mAb 4G2
and i-body WD34-Fc (mouse Fc) binding to *Pf*AMA1 in the presence or absence of *Pf*MSP2
protein. Data represents the mean of 3 experiments with significance determined by unpaired
t-test with p< 0.05 deemed significant. **(D)** ELISA based assessment of the anti-*Pf*AMA1 mAb
4G2 and i-body WD34-Fc antibody binding levels to recombinant *Pf*3D7 AMA1 in the
presence or absence of recombinant *Pf*3D7 MSP2. PBS control demonstrates background
fluorescence. Dashed orange line provides a guide for peak absorbance levels. Anti-*Pf*MSP2
mAb shows increasing concentrations of *Pf*MSP2 protein results in decreased binding of mAb
4G2 and i-body WD34-Fc. Data represents the mean of 3 experiments and error bars are ±S.D.

Discussion

Despite decades of research and interest, the function of most merozoite surface proteins
remains unclear and targeting merozoite antigens in vaccine development has not achieved
high efficacy in clinical trials. Additionally, it has generally proven difficult to generate highly
inhibitory antibodies through vaccination with merozoite antigens. *Pf*MSP2 is one of the most
abundant proteins on the merozoite surface, has shown promise as a vaccine candidate and is
an established target of protective cytophilic antibodies (Boyle et al., 2015; Genton et al., 2002;
McCarthy et al., 2011; Osier et al., 2014, 2010). However, its function remains unclear. Our
findings here suggest that *Pf*MSP2 does not have an essential role in invasion *in vitro*, but can
modulate the sensitivity of merozoites to inhibitory antibodies targeting other antigens.

Despite its abundance on the merozoite surface and previous work suggesting a role in RBC
invasion, we found merozoites invade normally in the absence of this protein after gene
deletion. This does not exclude a role for *Pf*MSP2 *in vivo* where there are additional pressures,
such as immune-effector mechanisms and flow dynamics, on merozoite invasion. However,
given we have knocked-out *Pf*MSP2 from two different *P. falciparum* isolates, our findings
do not currently support a major role for *Pf*MSP2 in the mechanics of merozoite invasion. Thus,

it appears that the function of the two most abundant proteins on the merozoite surface, *Pf*MSP1 (Das et al., 2015; Kals et al., 2024) and *Pf*MSP2, are not directly linked to merozoite binding to the RBC and subsequent invasion. Deletion of MSP2 did lead to significant changes in the expression of several genes, but there were no proteins with a clear link to merozoite invasion and the biological relevance of these changes is not currently understood and requires future investigation.

Both rabbit polyclonal and mouse monoclonal anti-*Pf*AMA1 antibodies tested were consistently more inhibitory in the absence of *Pf*MSP2. Of the two *Pf*AMA1 targeting i-bodies tested (Angage et al., 2024), the most potent WD34 was also more potent in the absence of *Pf*MSP2, with this activity increasing significantly with the ~5.6-fold increase in size that occurred upon addition of a human Fc region. Similarly, activity of a nanobody targeting the PCRCR complex protein CSS showed increased potency in the absence of *Pf*MSP2, and again this was increased further with antibody enlargement through addition of an Fc region. These data suggest that *Pf*MSP2 may act to dampen the inhibitory activity of antibodies targeting other antigens on the surface of the merozoite. Such immune dampening (called conformational masking) by intrinsically disordered proteins, or protein regions, of host-cell invasion related proteins has recently been demonstrated by modelling and gene-editing studies in the viral pathogens HIV and HCV, respectively (Li et al., 2024; Stejskal et al., 2022). These studies, along with our demonstration of increase antibody inhibitory efficacy with loss of the intrinsically disordered *Pf*MSP2, support that this may be a widespread approach used by pathogens to protect important protein functional sites from inhibitory antibodies.

A role in immune evasion has previously been suggested for *Pf*MSP2 and speculatively an alternate explanation for the sensitisation observed is that this protein may shield key *Pf*AMA1 epitopes from antibody inhibition. For the *P. falciparum* 3D7 line, we found that *Pf*MSP2 knock-out led to higher *Pf*AMA1 binding as measured by increased fluorescence intensity for the mCherry tagged i-body WD34 in the absence of *Pf*MSP2. Using a solid-phase model and titrating recombinant *Pf*MSP2, we found that increasing concentrations of *Pf*MSP2 reduced binding of an anti-*Pf*AMA1 mAb and i-body, even at concentrations where *Pf*MSP2 was not detectable itself by antibodies. These data support that the presence of *Pf*MSP2 may reduce antibody access (masking) to important invasion proteins exposed on the merozoites surface. However, we acknowledge that the antibody-binding assays used (immunofluorescence assays, SPR, ELISA) might not be sensitive enough to observe all potential changes in binding, temporal changes when binding epitopes are exposed or differences in avidity that could lead

to increased antibody potency during the course of parasite invasion and growth, and these
factors may also contribute. Future studies of the merozoite surface and interactions between
proteins may help decipher in full how *Pf*MSP2 is protecting essential merozoite antigens from
invasion inhibitory antibody activity and reveal strategies for enhanced vaccine design.

Our observation that loss of *Pf*MSP2 potentiates the invasion-inhibitory activity of antibodies
targeting other antigens opens an additional avenue for vaccine development, such as targeting
*Pf*MSP2 to interfere with its potential role in immune evasion and thus potentiating antibodies
against other merozoite antigens. Additionally, our results suggest that vaccines based on
*Pf*AMA1 may need to be designed to specifically avoid the effect of *Pf*MSP2 modulating
inhibitory antibodies, such as focussing on specific epitopes or structural features. To date,
vaccines based on AMA1 have failed to achieve substantial efficacy (Sagara et al., 2009;
Sheehy et al., 2012; Thera et al., 2011). We can speculate that one possible protective
mechanism of action of the Combination B vaccine (Genton et al., 2002; McCarthy et al.,
2011), which combined *Pf*MSP2, *Pf*MSP1 and *Pf*RESA, could be that the inhibitory activity
of vaccine-induced antibodies raised against *Pf*MSP2 were then able to potentiate the activity
of antibodies targeting other antigens, potentially *Pf*MSP1 which was also a target of the
vaccine. Here we show consistent potency improvement with *Pf*MSP2 knock-out for growth
inhibitory rabbit, mouse monoclonal and i-body antibodies targeting *Pf*AMA1, as well as
demonstrate improved activity for and Fc-tagged nanobody targeting *Pf*CSS, indicating that
these are not outlier results from a single antibody or antibody type. However, increased
antibody potency was not shared across all antibodies tested, possibly because the specific
function or localisation of a target protein, the region that an antibody binds to or the functional
activity (or lack thereof) of an antibody may all play a role in determining whether loss of
*Pf*MSP2 can potentiate growth inhibitory activity. Further investigation using the parasite lines
developed in this study and a wider panel of antibodies that target different stages of the
merozoite invasion process could shed more light on this potentially novel mechanism of
vaccine derived antibody efficacy.

The results of this study and work done by Escalante et al. (2022) demonstrate that MSP2-like
sequences are found in the avian malarias *P. galinaceum* and *P. relictum*, which differs from
the fields previous understanding that *Pf*MSP2 arose in the *Laverania* lineage of malaria
parasites. This observation indicates that MSP2 was likely present in the ancestral malaria
parasite before being lost in primate and rodent *Plasmodium* spp. Further insights into the
evolution and loss of *Pf*MSP2 are likely to be found with additional whole genome sequences

of bird and lizard malaria species. The presence of *msp2*-like sequences in some avian

Plasmodium species and *Laverania* suggest a retention of *msp2* sequences in the *Laverania*

which indicates an important role for MSP2 for these parasites. Indeed, our amino acid and

structural prediction level comparison demonstrates that much of the protein properties

conserved in the *Laverania* MSP2s are also recognisable, and in many cases conserved, in the

avian malaria MSP2s, despite 10 million years of evolutionary divergence. This suggests that

having an MSP2-like protein is beneficial for these parasites with very different hosts.

However, CRISPR-Cas9 gene editing used in this work has shown that, in contrast to previous

work, *Pf*MSP2 is not essential for *P. falciparum* blood stage parasite growth *in vitro*. Our

findings raise the possibility that the retention of *Pf*MSP2 in the *Laverania*, and by association

potentially the avian malaria parasites, may be linked to its apparent capacity to modulate the

impact of antibodies against other merozoite surface-exposed antigens, rather than through a

direct mechanistic role in host-cell invasion. If MSP2 was to fulfill such a role in the *Laverania*

and avian malaria parasites, it raises the question of whether a similar protective system exists

in other *Plasmodium* spp. and what the key protein/s are that fulfil this role in these parasites.

The *Pf*3D7 MSP2 and *Pf*Dd2 MSP2 lines developed in this study will provide useful reagents

for investigating the potential of targeting a single, or both, *Pf*MSP2 allelic types with vaccines

that elicit different immune responses. These parasites could also be used to explore whether

vaccine-induced antibodies against other merozoite antigens could also be potentiated if any

immune evasion mechanism provided by *Pf*MSP2 is removed. This may be achievable with a

combination vaccine or through targeting an epitope of another merozoite vaccine candidate

that is not protected by *Pf*MSP2. Such information, investigated using the *Pf*MSP2 knock-out

parasites developed in this study, could be useful in prioritising vaccine candidates, specific

epitopes and combinations that target the merozoite.

Conclusion

Advancements in gene-editing techniques in *P. falciparum* have allowed us to show for the

first time, and contrary to previous work, that *Pf*MSP2 is not essential for *P. falciparum* growth

in vitro. Instead, we present a new concept that MSP2 can modulate the activity of invasion-

inhibitory antibodies and may have evolved for this purpose. It is possible that other merozoite

surface proteins may have similar roles and should be investigated in future studies. Our

observation that loss of *Pf*MSP2 potentiates the inhibitory activity of antibodies targeting other

merozoite surface exposed proteins reveals a new avenue to explore in vaccine development

by blocking or bypassing this function to improve the activity of vaccines targeting these

antigens.

Materials and Methods

Bioinformatic analysis of *Pf*MSP2, *Pf*MSP4 and *Pf*MSP5 like proteins in *Plasmodium* spp.

The PlasmoDB database (Aurrecoechea et al., 2009) was used to identify in the available

Plasmodium genomes the region between Adenylosuccinate lyase (PF3D7_0206700) and a

conserved protein of unknown function (PF3D7_0207100) where *Pf*MSP2, *Pf*MSP4 and

*Pf*MSP5 are localised in *P. falciparum*. All annotated MSP2, MSP4 and MSP5 protein

sequences were downloaded. Translated sequences of unannotated genes and putative

annotations in this region were also downloaded and aligned using the MUSCLE alignment

tool MEGA11: Molecular Evolutionary Genetics Analysis version 11 (Tamura et al., 2021) or

the Geneious Prime 2019.2.1 MUSCLE alignment tool (Biomatters Ltd.) and determined to be

MSP2, MSP4 or MSP5 based on sequence similarity in the conserved regions of these genes.

A BLAST (NCBI) search with the N-terminal and C-terminal conserved regions of MSP2 was

also performed to capture any sequences missed by the above approach. A maximum likelihood

phylogenetic tree of identified MSP2 sequences was generated using the default maximum

likelihood settings of MEGA11 with 500 bootstraps and the *P. gallinaceum* MSP2 as an

outgroup.

Modelling of MSP structure across *Plasmodium* spp. using Alpha-fold

In order to obtain models of the 3-dimensional protein structures of MSP2 for *P. falciparum*,

P. reichenowi, *P. billcollinsi*, *P. adleri*, *P. relictum* and *P. gallinaceum*, we first removed the

predicted N-terminal signal sequences and C-terminal GPI anchor sequence. The resulting

sequences were submitted to AlphaFold2 (Jumper et al., 2021). The resulting models were

subsequently analysed and visualised using PyMOL 2.4.0 (Schrödinger, USA).

Culture and synchronisation of *P. falciparum*

Plasmodium falciparum 3D7 and Dd2 was cultured in RPMI-HEPES media (Gibco) containing

0.5% Albumax ll (Gibco), 25 µM Gentamicin (Gibco), 367 µM hypoxanthine (Sigma Aldrich),

2 mM L-Glutamax (Gibco), 0.17% sodium bicarbonate (Sigma Aldrich) and O positive human

red blood cells (Lifeblood, Australia) at 3% parasitemia, 3% haematocrit. Cultures were grown

at 37°C under 1% oxygen, 5% CO_2 (BOC gas), in a humidified chamber. For gene-edited lines,

WR99210 (Jakobus Pharmaceuticals) was added to culture media. Sorbitol lysis (5%) (Sigma

Aldrich) to select for ring-stage parasites, Percoll Plus (Sigma Aldrich) selection of late-stage

parasites and heparin treatment (Pfizer) (Boyle et al., 2010b) were used to maintain

synchronicity of parasite cultures.

Generation of *Pf*MSP2 knockout line using CRISPR Cas9

Briefly, a flank region incorporating the 5' region upstream of *Pf*MSP2, the first 44 amino

acids of the protein, and a flank of the region immediately 3' of *Pf*MSP2 was inserted into the

pCC1 plasmid by restriction enzyme cloning. A 20 bp guide sequence was designed using

EuPaGDT (Peng and Tarleton, 2015), annealed and inserted in the pUF_Cas9 plasmid by

infusion cloning (Takara Bio). pUF_Cas9 guide plasmid (20 µg) and pCC1 *Pf*MSP2 repair

plasmid (60 µg) was transfected directly into *P. falciparum* schizonts. Early schizonts were

purified using a 70% percoll gradient and treated for 4 hrs with compound 1 (Taylor et al.,

2010) at 2 µM. After incubation treated schizonts were washed twice and left shaking for 20

minutes before pelleting and resuspension in Complete Cytomix (plasmids plus 0.895% KCl,

0.0017% $CaCl_2$, 0.076% EGTA, 0.102% $MgCl_2$, 0.0871% K_2HPO_4, 0.068% KH_2PO_4, 0.708%

HEPES). Schizonts were electroporated in 0.2 cm cuvettes (Biorad) at 800 V and 25 µF.

Electroporated schizonts were mixed with warmed media and fresh RBCs and shaken to

promote invasion for 20 minutes after which they were placed in a culture dish. Transfected

parasites were then treated with WR99210 to select for integration of the hDHFR drug selection

cassette and MSP2 knock-out.

To confirm integration gDNA was collected by saponin lysis of schizont stage parasite culture

and gDNA extracted using the PureLink Genomic DNA Mini Kit (Invitrogen) as per

manufacturer's instructions. Integration of the *Pf*MSP2 knockout construct into a portion of

parasites was confirmed by PCR (Supplementary Table 1). Once confirmed cultures were

cycled on and off WR99210 (Jakobus Pharmaceuticals) and 1 µM 5FC (Sigma Aldrich) to

select for integrated parasites no longer expressing the guide plasmid. Limited dilution cloning
was then performed to obtain parasite clones which had integrated the knockout construct.

Western blotting to detect *Pf*MSP2

Synchronised schizonts were harvested and the RBCs lysed by saponin. Briefly ~10 mL
schizont (38-44 hrs) culture was lysed on ice for 10 minutes with 0.15% w/v saponin before
pelleting by centrifugation and washing once in 0.075% w/v saponin followed by three washes
in PBS containing protease inhibitors (CØmplete, Roche). Saponin treated schizont pellets
were treated with DNAseI (Qiagen) for 10 minutes at room temperature before resuspension
in reducing sample buffer (0.125 M Tris-HCL, 4% SDS, 20% glycerol, 10% beta-
mercaptoethanol and 0.002% bromophenol blue). Proteins were separated by size using SDS-
PAGE 4-12% Bis-Tris Gels (Bolt, Invitrogen) at 110V for 80 minutes before transfer to a
nitrocellulose membrane (iBlot, Invitrogen) at 20V for 7 minutes. Blots were blocked from 1
hr to overnight in 3% skim milk PBS (Sigma Aldrich) before incubation with primary
antibodies (1/75000 mouse 2F2 anti-*Pf*MSP2 3D7 (Supplementary Table 2; (Adda et al.,
2012)) and 1/75000 rabbit anti-*Plasmodium* aldolase (Abcam) or 1/75000 rabbit anti-FC27 and
1/10000 mouse anti- EXP2 (a gift of Paul Gilson, Burnet Institute, Melbourne) for 1- 2 hrs.
IRDye 800CW goat anti-mouse (1/4000, LI-COR Biosciences) or IRDye 680RD goat anti-
rabbit (1/4000, LI-COR Biosciences) secondary antibodies were used for detection on the
Odyssey Infrared imaging system (LI-COR Biosciences). Quantification was performed using
Image Studio Lite 5.2.5 (LI-COR Biosciences).

Quantification of parasite expansion rate and invasion inhibition

Potential growth defects resulting from gene editing was assessed by flow cytometry. The
initial parasitemia of cultures was determined by flow cytometry and then measured again after
the 50 μL cultures in 96 well plates were maintained under standard (still) or shaking (50 rpm)
conditions for 48 hrs or 96 hrs of growth. To determine parasitemia by flow cytometry
Ethidium Bromide (10 μg/mL BioRad) was added to 50 μL of 1% haematocrit culture for 30
mins to allow for staining of parasite DNA. The stain was washed off and wells resuspended
in 200μL of PBS. Data was collected on a BD Accuri C6 Plus Flow Cytometer (BD
Biosciences) and analysed on FlowJo (FlowJo LLC). Briefly, forward scatter and side scatter
was used to determine the RBC population. From this population infected RBCs were gated

on as highly fluorescent in the PE channel. Final parasitemia was compared back to initial

parasitemia to determine the fold increase in parasitemia for each line.

To examine the invasion inhibitory effect of antibodies (Supplementary Table 2), 5 µL of

potential inhibitor was mixed with 45 µL of 0.2% schizont parasitemia at 1% haematocrit for

96 hrs before end parasitemia was determined by flow cytometry. Data is displayed as %

Growth of media only controls for each line.

To examine differences in RBC receptor preference RBCs were treated with different enzymes

that cleave different residues of RBC receptors which is used as a marker for which RBC

receptor/s are preferred by *P. falciparum* (Duraisingh et al., 2003). Infected RBCs at 1% ring

stage parasitemia were washed three times in RPMI-HEPES to remove excess protein. Packed

RBCs (20 µL) at 1% parasitemia were then resuspended with 5X RBC volume of pre-warmed

enzymes at desired concentration (Supplementary Table 3). RBC-enzyme mix was incubated

at 37°C for 45 mins with rotation to keep RBCs resuspended. Following incubation, RBCs

were washed three times and added to fresh complete media at 1% haematocrit in a U- bottom

96 well plate in technical duplicate. Plates were incubated for 72 hrs, sufficient time for *P.

falciparum* to complete one full cycle of growth and reach trophozoite stage again. Final

parasitemia was determined by flow cytometry, as detailed above, and compared to non-treated

controls. IC_{50} calculations data was log transformed then a nonliner regression log-(inhibitor)-

versus-response curve was calculated with Extra Sum-of-Squares F Test (best-fit $LogIC_{50}$)

used to compared IC_{50}s between conditions.

Immunofluorescence assay of Wildtype and *Pf*MSP2 KO parasites

Schizonts ~38 hrs old were treated with E64 (Sigma-Aldrich) for 5 hrs to allow development

of very mature schizonts desired for imaging. After E64 treatment, parasite cultures were fixed

in 4% v/v paraformaledehyde (PFA, Sigma-Aldrich), 0.0075% v/v glutaraldehyde (pH 7.5,

Electron Microscopy Sciences) solution for 30 min at room temperature shaking gently. Fixed

parasites were washed in 1X PBS and then resuspended in PBS at 1% haematocrit. Coverslips

(#1.5H high-precision coverslips, Carl Zeiss, Oberkochen, Germany) were coated in 0.01%

Poly-L-lysine (Sigma Aldrich) for 1 hr at room temperature, washed in miliQ water before the

fixed parasite culture was allowed to adhere for 1 hr. Cells were permeabilised by 0.1% Triton

X-100 PBS for 10 mins before blocking for 1 hr to overnight in 3% BSA 0.05% Tween 20

PBS. Primary antibodies were diluted in 1% BSA 0.05% Tween 20 PBS and incubated for 2

hrs. Coverslips were washed in 0.05% Tween 20 PBS three times before incubation with the
secondary antibody (1/500 goat anti-chicken/mouse/rabbit Alexa Fluor coupled secondary
antibodies -488 nm, 594 nm, 647 nm; Life Technologies) for 1 hr. After the secondary antibody
was washed off coverslips underwent a secondary fix in 4% v/v PFA, 0.0075% v/v
glutaraldehyde for 5 mins. The fixative was washed off and the coverslips dehydrated with
sequential 3 min ethanol treatment (70%, 90% and 100%). Coverslips were allowed to dry and
mounted with ProLong Gold antifade solution (refractive index 1.4) which contains 4', 6-
diamidino-2phenylindole, dihydrochloride (DAPI) (ThermoFisher Scientific) and allowed to
set overnight. Imaging was performed on the Zeiss LSM 800 using the Airyscan super-
resolution mode (Carl Zeiss, Obekochen, Germany).

Live cell microscopic analysis of merozoite RBC invasion

Highly synchronised PfDd2 and PfDd2 ΔMSP2 parasites at schizont stages were adjusted to
0.25% Haematocrit in complete culture medium. A volume of 200 μL was added to a well of
an iBidi 15 μ-Slide 8 well, glass bottom slide (iBidi 80827) and the slide was promptly returned
to sit on a prewarmed water block within a gassed box and left to incubate at 37°C to allow
iRBCs to settle. The slide was then transported directly to a prewarmed Nikon TiE microscope
with an environmental chamber heated to 37°C and gas mixture of 1% O_2, 5% CO_2 and 94%
nitrogen. Filming was conducted using either a 60X water or 100X oil objective. Differential
interference contrast (DIC) imaging was captured at 3 – 3.5 Volts with a camera exposure of
60 – 180 milliseconds.

Captured footage was processed using the Nikon analysis software (NIS-Elements AR
Analysis version 5.21.01) and analysis of the schizont rupture and merozoite invasion events
was performed manually. The merozoites which invaded successfully were tracked and
attachment time, reorientation time (if obvious), deformation start and end times, the
deformation score (Weiss et al., 2015; Wilson et al., 2015), invasion initiation and completion
times and echinocytosis start times were recorded.

RNA extraction

Highly synchronised cultures were pelleted, resuspended in TriZol (Invitrogen) and incubated
for 5 minutes at 37°C. Chloroform (Sigma Aldrich) was added, mixed and then spun at 12,000g

for 30 minutes at 4°C. Supernatant was collected and mixed with an equal volume of 70%

Ethanol and processed using the RNeasy Kit (Qiagen) to extract purified RNA. RNA was

DNAse l (Qiagen) treated for 30 mins at room temperature and cleaned up on RNeasy columns

according to the manufacturers protocol. Absence of gDNA was checked by qPCR of gDNA

with primers (*Pf*SUB1) and DNAse I treatment repeated if necessary.

qPCR of gene expression in schizonts

DNAse treated RNA was added to 40 mM dNTPs (Qiagen), 0.4 mg/mL random hexamer

(Qiagen) and incubated at 65°C for 5 mins. After incubation 5x Superscript Buffer (Invitrogen),

100 mM DTT (Invitrogen), 40 U/μL RNAseOUT (Invitrogen) and 200 μ/mL Superscript lll

Reverse Transcriptase (Invitrogen) was added and cycled 25°C 5 mins, 50°C for 60 mins and

70°C for 15 mins. Primers (Supplementary Table 4) were designed to detect *Pf*MSP2, *Pf*MSP4,

*Pf*MSP5, *Pf*SUB1 and Fructose Bisphosphate Aldolase as a reference gene (Salanti et al.,

2003). qPCR was performed on cDNA from WT and knockout parasites with PowerUp SYBR

Green Master Mix (Applied Biosystems) and cycling parameters; 95°C for 10 mins followed

by 40 cycles 95°C for 15 mins, 60°C for 14 mins before a dissociation step at 95°C for 2 mins

followed by 60°C for 2 mins with a 2% ramp to 95°C for 2 mins on the QuantStudio 7 Flex

System (ThermoFisher). qPCR was performed in triplicate for each of the three independent

experiments. Change in gene expression in knockout lines compared to WT was quantified by

the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

Differential gene expression in *Pf*MSP2 KO compared to WT schizonts

RNA sequencing of the schizont stage transcriptome of *Pf*3D7 MSP2 KO and *Pf*3D7 WT was

performed using the Illumina total RNA with Ribo Zero plus for library preparation and

sequenced on the NovaSeq 6000 platform (Victorian Clinical Genetics Services) with 2x150

bp paired end reads (Supplementary Table 5). Reads were aligned to the *Pf*3D7 reference

genome (PlasmoDB v61) using STAR aligner (Dobin et al., 2013). Reads were called using

Rsubread FeatureCounts (Liao et al., 2014) with reads counted by CDS and summarised by

gene. Differential gene analysis was performed with DESeq2 (Love et al., 2014) and visualized

in R studio using ggplots2 (Wickham, 2016). MSP2 RNASeq data is available at ArrayExpress

through accession number E-MTAB-15427.

Quantitative Immunofluorescence assay

Synchronous Pf3D7 MSP2 KO and Pf3D7 WT parasite cultures were treated with ML10 (Ressurreição et al., 2020) to allow mature schizonts to develop. Thin blood smears were then made and fixed in 100%, -20°C methanol for 5 mins. Blocking of smears was performed in 1% BSA PBS, then fluorophore conjugated WD34 (240 ng/mL) and WD33 (120 ng/mL) i-bodies (Angage et al., 2024) in 1% BSA PBS were added and incubated at room temperature for 1 hr. Smears were then washed, ethanol dehydrated and a coverslip mounted (ProLong Diamond Antifade mountant with DAPI (refractive index 1.47, ThermoFisher Scientific)) and allowed to cure for 48 hrs at room temperature. Fluorescence images were acquired using an Olympus FV3000 confocal microscope equipped with a 100x oil objective (NA 1.4). Quantification of parasite associated fluorescence was performed using a modified image analysis pipeline (Liffner et al., 2020) implemented in Imaris^{TM} (9.9 version) where non-specific background fluorescence was removed using rolling ball background subtraction and a thresholding algorithm applied to the image to generate binary masks, separating objects exceeding the threshold (putative parasites) from the background. Binary masks were further refined to exclude objects smaller or larger than a defined size range (< 3 μm and > 15 μm), corresponding to the expected size of AMA i-body stained schizonts. Following segmentation the surface area containing antibody signal were calculated and these values compared between Pf3D7 MSP2 KO and Pf3D7 WT parasites.

Surface Plasmon Resonance (SPR) Assay

Surface plasmon resonance (SPR) was employed to determine the association rate constant (K_a) of AMA1-antibody interactions in the presence of MSP2 using the Biacore T200 system (Cytiva). Flow cells 1, 2, and 3 were activated for 14 minutes using a 1:1 mixture of 0.1 M N-hydroxysuccinimide (NHS) and 0.4 M N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDC) at a flow rate of 5 μL/min. Flow cell one was immobilised with bovine serum albumin (BSA) to serve as a reference surface. AMA1 was immobilised at a concentration of 80 μg/mL in 10 mM sodium acetate (pH 4.5) to achieve a surface density of 1000 response units (RU) on flow cells Two and three. MSP2 was co-immobilised in flow cell three to a surface density of 2000 RU. All surfaces were subsequently blocked with a 7-minute injection of 1 M ethanolamine (pH 8.0). Purified antibodies were prepared in PBS containing 0.005% Tween-20 (pH 7.4) and injected over the flow cells at a flow rate of 60

µL/min at 25°C. Single-cycle kinetics, using a bivalent analyte model, was applied to analyse
the interactions.

Enzyme-Linked Immunosorbent Assay (ELISA)

ELISAs to evaluate the effect of MSP2 on the binding of AMA1-specific antibodies were

carried out in a volume of 100 µL per well, either at room temperature for 1 hr or overnight

at 4°C. All washing steps consisted of three washes with PBS containing 0.1% Tween-20

(PBS-T). Nunc Maxisorp plates (Thermo Fisher Scientific, USA) were coated with AMA1

at a concentration of 0.8 µg/mL, followed by washing and subsequent coating with

increasing concentrations of recombinant HIS-tagged *Pf*MSP2, HIS-tagged *Pf*MSP4 or HIS-

tagged NCAM. Wells were then blocked with 5% (w/v) skim milk in PBS and then

incubated for 1 hr at room temperature with one of mouse Fc-conjugated WD34 anti-AMA1

i-body (Angage et al., 2024), mouse monoclonal anti-AMA1 antibody (mAb) 4G2 (Kocken

et al., 1998), anti-MSP2 mAb 9G8 (Adda et al., 2012), anti-HIS antibody (Sigma Aldrich) to

detect *Pf*MSP4 or anti-NCAM antibody (21H5, (Griffiths et al., 2016)) at a concentration of

2.5 µg/mL. The primary antibody was washed off and the HRP-conjugated secondary

antibody (1:5000, Sigma Aldrich) was then applied for 1 hr at room temperature before

washing. For signal development to detect bound primary antibodies, wells were incubated

with 1-Step^{TM} Ultra TMB-ELISA Substrate Solution (Thermo Fisher Scientific, USA)

according to the manufacturers instructions. The reaction was stopped by adding 1 M

sulfuric acid, and the absorbance at 450 nm was measured using a microplate reader.

Statistical Analysis

Three independent experiments were performed for all experiments in technical duplicate,

unless otherwise stated. Data was graphed and statistical tests performed in Prism GraphPad

v9 or R.

Acknowledgements

The authors would like to thank Prof Alan Cowman, Dr Stephen Scally and Jennifer Thompson

(WEHI, Melbourne) for provision of PCRCR complex antibodies. Assoc Prof Paul Gilson

(Burnet Institute, Melb) for EXP2 and MSP1 antibodies. Dr Chris MacRaild (Monash

University) for helpful discussion. Dr Adam Thomas for the *Pf*MSP4 protein (Burnet Institute).

Dr Sonja Frölich (the University of Adelaide) for assistance on quantitative

immunofluorescence assays and Adelaide Microscopy for training and use of microscopy

instruments. We thank Lifeblood, Australia, for provision of RBCs. DWW was supported by

a Hospital Research Foundation (C-MCF-52-2019) and Australian Research Council

(FT240100420) Future Fellowship. DWW and JGB by a Hospital Research Foundation

Collaborative Grant (S-03-EOI-2021). JGB was supported by the National Health and Medical

Research Council of Australia (Research Fellowship and Investigator Grant to JGB, Australian

Centre for Research Excellence in Malaria Research). IGH and JC were supported by an

Australian Research Council RTP scholarship. The Burnet Institute is supported by the

NHMRC for Independent Research Institutes Infrastructure Support Scheme and the Victorian

State Government Operational Infrastructure Support.

Author Contributions

919 Conceived and designed the experiments: DWW, IGH, JC, JGB, DA, MF, RA. Performed the

920 experiments and phylogenetic analysis: IGH, JC, KHL, OR, DA, KRT, NB. Analysed the data:

921 IGH, DWW, JC, KHL, OR, DA, MF. Provided key discussions on PfMSP2 and/or key

922 reagents: RA, MF, DA. Paper writing: DWW, JC, IGH, JGB with critical input from RA, MF,

923 DA, OR, KRT and based on the PhD thesis of IGH.

Competing Interests

926 M.F. is the founding chief scientist and a shareholder in AdAlta Ltd., and R.F.A. is also a

927 shareholder in AdAlta. The other authors have declared no competing interests.

References

Adda, C.G., MacRaild, C.A., Reiling, L., Wycherley, K., Boyle, M.J., Kienzle, V., Masendycz,

P., Foley, M., Beeson, J.G., Norton, R.S., Anders, R.F., 2012. Antigenic characterization

of an intrinsically unstructured protein, Plasmodium falciparum merozoite surface protein

2. Infect Immun 80, 4177–4185. https://doi.org/10.1128/IAI.00665-12

Adda, C.G., Murphy, V.J., Sunde, M., Waddington, L.J., Schloegel, J., Talbo, G.H., Vingas,

K., Kienzle, V., Masciantonio, R., Howlett, G.J., Hodder, A.N., Foley, M., Anders, R.F.,

2009. Plasmodium falciparum merozoite surface protein 2 is unstructured and forms

amyloid-like fibrils. Mol Biochem Parasitol 166, 159–171.

https://doi.org/10.1016/j.molbiopara.2009.03.012

Alanine, D.G.W., Quinkert, D., Kumarasingha, R., Mehmood, S., Donnellan, F.R., Minkah,

N.K., Dadonaite, B., Diouf, A., Galaway, F., Silk, S.E., Jamwal, A., Marshall, J.M.,

Miura, K., Foquet, L., Elias, S.C., Labbé, G.M., Douglas, A.D., Jin, J., Payne, R.O.,

Illingworth, J.J., Pattinson, D.J., Pulido, D., Williams, B.G., de Jongh, W.A., Wright, G.J.,

Kappe, S.H.I., Robinson, C. V, Long, C.A., Crabb, B.S., Gilson, P.R., Higgins, M.K.,

Draper, S.J., 2019. Human Antibodies that Slow Erythrocyte Invasion Potentiate Malaria-

Neutralizing Antibodies. Cell 178, 216-228.e21.

https://doi.org/10.1016/j.cell.2019.05.025

Anders, R.F., Adda, C.G., Foley, M., Norton, R.S., 2010. Recombinant protein vaccines against

the asexual blood-stages of Plasmodium falciparum. Hum Vaccin 6, 39–53.

https://doi.org/10.4161/hv.6.1.10712

Angage, D., Chmielewski, J., Maddumage, J.C., Hesping, E., Caiazzo, S., Lai, K.H., Yeoh,

L.M., Menassa, J., Opi, D.H., Cairns, C., Puthalakath, H., Beeson, J.G., Kvansakul, M.,

Boddey, J.A., Wilson, D.W., Anders, R.F., Foley, M., 2024. A broadly cross-reactive i-

body to AMA1 potently inhibits blood and liver stages of Plasmodium parasites. Nat

Commun 15, 7206. https://doi.org/10.1038/s41467-024-50770-7

Aurrecoechea, C., Brestelli, J., Brunk, B.P., Dommer, J., Fischer, S., Gajria, B., Gao, X.,

Gingle, A., Grant, G., Harb, O.S., Heiges, M., Innamorato, F., Iodice, J., Kissinger, J.C.,

Kraemer, E., Li, W., Miller, J.A., Nayak, V., Pennington, C., Pinney, D.F., Roos, D.S.,

Ross, C., Stoeckert, J.C.J., Treatman, C., Wang, H., 2009. PlasmoDB: a functional

genomic database for malaria parasites. Nucleic Acids Res 37, D539–D543.

https://doi.org/10.1093/nar/gkn814

Bannister, L.H., Mitchell, G.H., Butcher, G.A., Gennis, E.D., Cohen, S., 1986. Structure and

development of the surface coat of erythrocytic merozoites of Plasmodium knowlesi. Cell

Tissue Res 245, 281–290.

Baum, J., Maier, A.G., Good, R.T., Simpson, K.M., Cowman, A.F., 2005. Invasion by P.
falciparum Merozoites Suggests a Hierarchy of Molecular Interactions. PLoS Pathog 1,
e37.

Beeson, J.G., Drew, D.R., Boyle, M.J., Feng, G., Fowkes, F.J.I.I., Richards, J.S., 2016.
Merozoite surface proteins in red blood cell invasion, immunity and vaccines against
malaria. FEMS Microbiol Rev 40, 343–372. https://doi.org/10.1093/femsre/fuw001

Beeson, J.G., Kurtovic, L., Dobaño, C., Opi, D.H., Chan, J.-A., Feng, G., Good, M.F., Reiling,
L., Boyle, M.J., 2019. Challenges and strategies for developing efficacious and long-
lasting   malaria   vaccines.   Sci   Transl   Med   11.
https://doi.org/10.1126/scitranslmed.aau1458

Black, C.G., Barnwell, J.W., Huber, C.S., Galinski, M.R., Coppel, R.L., 2002. The
Plasmodium vivax homologues of merozoite surface proteins 4 and 5 from Plasmodium
falciparum are expressed at different locations in the merozoite. Mol Biochem Parasitol
120, 215–224. https://doi.org/https://doi.org/10.1016/S0166-6851(01)00458-3

Böhme, U., Otto, T.D., Cotton, J.A., Steinbiss, S., Sanders, M., Oyola, S.O., Nicot, A., Gandon,
S., Patra, K.P., Herd, C., Bushell, E., Modrzynska, K.K., Billker, O., Vinetz, J.M., Rivero,
A., Newbold, C.I., Berriman, M., 2018. Complete avian malaria parasite genomes reveal
features associated with lineage-specific evolution in birds and mammals. Genome Res
28, 547–560. https://doi.org/10.1101/gr.218123.116

Boyle, M.J., Langer, C., Chan, J.-A., Hodder, A.N., Coppel, R.L., Anders, R.F., Beeson, J.G.,
2014. Sequential Processing of Merozoite Surface Proteins during and after Erythrocyte
Invasion   by   Plasmodium   falciparum.   Infect   Immun   82,   924–936.
https://doi.org/10.1128/IAI.00866-13

Boyle, M.J., Reiling, L., Feng, G., Langer, C., Osier, F.H., Aspeling-Jones, H., Cheng, Y.S.,
Stubbs, J., Tetteh, K.K.A., Conway, D.J., McCarthy, J.S., Muller, I., Marsh, K., Anders,
R.F., Beeson, J.G., 2015. Human antibodies fix complement to inhibit plasmodium
falciparum invasion of erythrocytes and are associated with protection against malaria.
Immunity 42, 580–590. https://doi.org/10.1016/j.immuni.2015.02.012

Boyle, M.J., Richards, J.S., Gilson, P.R., Chai, W., Beeson, J.G., 2010a. Interactions with
heparin-like molecules during erythrocyte invasion by Plasmodium falciparum
merozoites. Blood 115, 4559–4568. https://doi.org/10.1182/blood-2009-09-243725

Boyle, M.J., Wilson, D.W., Richards, J.S., Riglar, D.T., Tetteh, K.K.A., Conway, D.J., Ralph, S.A., Baum, J., Beeson, J.G., 2010b. Isolation of viable Plasmodium falciparum merozoites to define erythrocyte invasion events and advance vaccine and drug development. Proc Natl Acad Sci U S A 107, 14378–14383. https://doi.org/10.1073/pnas.1009198107

Coley, A.M., Gupta, A., Murphy, V.J., Bai, T., Kim, H., Anders, R.F., Foley, M., Batchelor, A.H., 2007. Structure of the Malaria Antigen AMA1 in Complex with a Growth-Inhibitory Antibody. PLoS Pathog 3, e138.

Cowman, A.F., Tonkin, C.J., Tham, W.-H., Duraisingh, M.T., 2017. The Molecular Basis of Erythrocyte Invasion by Malaria Parasites. Cell Host Microbe 22, 232–245. https://doi.org/10.1016/j.chom.2017.07.003

Crosnier, C., Bustamante, L.Y., Bartholdson, S.J., Bei, A.K., Theron, M., Uchikawa, M., Mboup, S., Ndir, O., Kwiatkowski, D.P., Duraisingh, M.T., Rayner, J.C., Wright, G.J., 2011. Basigin is a receptor essential for erythrocyte invasion by Plasmodium falciparum. Nature 480, 534–537. https://doi.org/10.1038/nature10606

Das, S., Hertrich, N., Perrin, A.J., Withers-Martinez, C., Collins, C.R., Jones, M.L., Watermeyer, J.M., Fobes, E.T., Martin, S.R., Saibil, H.R., Wright, G.J., Treeck, M., Epp, C., Blackman, M.J., 2015. Processing of Plasmodium falciparum Merozoite Surface Protein MSP1 Activates a Spectrin-Binding Function Enabling Parasite Egress from RBCs. Cell Host Microbe 18, 433–444. https://doi.org/10.1016/j.chom.2015.09.007

Datoo, M.S., Dicko, A., Tinto, H., Ouédraogo, J.B., Hamaluba, M., Olotu, A., Beaumont, E., Ramos Lopez, F., Natama, H.M., Weston, S., Chemba, M., Compaore, Y.D., Issiaka, D., Salou, D., Some, A.M., Omenda, S., Lawrie, A., Bejon, P., Rao, H., Chandramohan, D., Roberts, R., Bharati, S., Stockdale, L., Gairola, S., Greenwood, B.M., Ewer, K.J., Bradley, J., Kulkarni, P.S., Shaligram, U., Hill, A.V.S., Mahamar, A., Sanogo, K., Sidibe, Y., Diarra, K., Samassekou, M., Attaher, O., Tapily, A., Diallo, M., Dicko, O.M., Kaya, M., Maguiraga, S.O., Sankare, Y., Yalcouye, H., Diarra, S., Niambele, S.M., Thera, I., Sagara, I., Sylla, M., Dolo, A., Misidai, N., Simando, S., Msami, H., Juma, O., Gutapaka, N., Paul, R., Mswata, S., Sasamalo, I., Johaness, K., Sultan, M., Alexander, A., Kimaro, I., Lwanga, K., Mtungwe, M., Khamis, K., Rugarabam, L., Kalinga, W., Mohammed, M., Kamange, J., Msangi, J., Mwaijande, B., Mtaka, I., Mhapa, M., Mlaganile, T., Mbaga, T., Yerbanga, R.S., Samtouma, W., Sienou, A.A., Kabre, Z., Ouedraogo, W.J.M., Yarbanga, G.A.B.,

Zongo, I., Savadogo, H., Sanon, J., Compaore, J., Kere, I., Yoni, F.L., Sanre, T.M., Ouattara, S.B., Provstgaard-Morys, S., Woods, D., Snow, R.W., Amek, N., Ngetsa, C.J., Ochola-Oyier, L.I., Musyoki, J., Munene, M., Mumba, N., Adetifa, U.J., Muiruri, C.M., Mwawaka, J.S., Mwaganyuma, M.H., Ndichu, M.N., Weya, J.O., Njogu, K., Grant, J., Webster, J., Lakhkar, A., Ido, N.F.A., Traore, O., Tahita, M.C., Bonko, M. dit A., Rouamba, T., Ouedraogo, D.F., Soma, R., Millogo, A., Ouedraogo, E., Sorgho, F., Konate, F., Valea, I., 2024. Safety and efficacy of malaria vaccine candidate R21/Matrix-M in African children: a multicentre, double-blind, randomised, phase 3 trial. The Lancet 403, 533–544. https://doi.org/10.1016/S0140-6736(23)02511-4

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21. https://doi.org/10.1093/bioinformatics/bts635

Douglas, A.D., Williams, A.R., Illingworth, J.J., Kamuyu, G., Biswas, S., Goodman, A.L., Wyllie, D.H., Crosnier, C., Miura, K., Wright, G.J., Long, C.A., Osier, F.H., Marsh, K., Turner, A. V., Hill, A.V.S., Draper, S.J., 2011. The blood-stage malaria antigen PfRH5 is susceptible to vaccine-inducible cross-strain neutralizing antibody. Nat Commun 2, 601. https://doi.org/10.1038/ncomms1615

Duraisingh, M.T., Maier, A.G., Triglia, T., Cowman, A.F., 2003. Erythrocyte-binding antigen 175 mediates invasion in Plasmodium falciparum utilizing sialic acid-dependent and -independent pathways. Proc Natl Acad Sci U S A 100, 4796–4801. https://doi.org/10.1073/pnas.0730883100

Eisenhaber, B., Bork, P., Eisenhaber, F., 1999. Prediction of potential GPI-modification sites in proprotein sequences. J Mol Biol 292, 741–58. https://doi.org/10.1006/jmbi.1999.3069

Escalante, A.A., Cepeda, A.S., Pacheco, M.A., 2022. Why Plasmodium vivax and Plasmodium falciparum are so different? A tale of two clades and their species diversities. Malar J 21, 139. https://doi.org/10.1186/s12936-022-04130-9

Feng, G., Boyle, M.J., Cross, N., Chan, J.-A., Reiling, L., Osier, F., Stanisic, D.I., Mueller, I., Anders, R.F., McCarthy, J.S., Richards, J.S., Beeson, J.G., 2018. Human Immunization With a Polymorphic Malaria Vaccine Candidate Induced Antibodies to Conserved Epitopes That Promote Functional Antibodies to Multiple Parasite Strains. J Infect Dis 218, 35–43. https://doi.org/10.1093/infdis/jiy170

Ferreira, E. d'Avila, Alexandre, M.A., Salinas, J.L., de Siqueira, A.M., Benzecry, S.G., de
Lacerda, M.V.G., Monteiro, W.M., 2015. Association between anthropometry-based
nutritional status and malaria: a systematic review of observational studies. Malaria J 14,
346. https://doi.org/10.1186/s12936-015-0870-5

Fowkes, F.J.I., Richards, J.S., Simpson, J.A., Beeson, J.G., 2010. The Relationship between
Anti-merozoite Antibodies and Incidence of Plasmodium falciparum Malaria: A
Systematic   Review   and   Meta-analysis.   PLoS   Med   7,   e1000218.
https://doi.org/10.1371/journal.pmed.1000218

Genton, B., Al-Yaman, F., Betuela, I., Anders, R.F., Saul, A., Baea, K., Mellombo, M.,
Taraika, J., Brown, G. V., Pye, D., Irving, D.O., Felger, I., Beck, H.-P.P., Smith, T.A.,
Alpers, M.P., 2003. Safety and immunogenicity of a three-component blood-stage malaria
vaccine (MSP1, MSP2, RESA) against Plasmodium falciparum in Papua New Guinean
children.   Vaccine   22,   30–41.   https://doi.org/https://doi.org/10.1016/S0264-
410X(03)00536-X

Genton, B., Betuela, I., Felger, I., Al-Yaman, F., Anders, R.F., Saul, A., Rare, L., Baisor, M.,
Lorry, K., Brown, G. V, Pye, D., Irving, D.O., Smith, T.A., Beck, H.-P., Alpers, M.P.,
2002. A Recombinant Blood-Stage Malaria Vaccine Reduces Plasmodium falciparum
Density and Exerts Selective Pressure on Parasite Populations in a Phase 1-2b Trial in
Papua New Guinea. J Infect Dis 185, 820–827.

Gilson, P.R., Nebl, T., Vukcevic, D., Moritz, R.L., Sargeant, T., Speed, T.P., Schofield, L.,
Crabb, B.S., 2006. Identification and Stoichiometry of Glycosylphosphatidylinositol-
anchored Membrane Proteins of the Human Malaria Parasite *Plasmodium falciparum*.
Molecular & Cellular Proteomics 5, 1286–1299. https://doi.org/10.1074/mcp.M600035-
MCP200

Griffiths, K., Dolezal, O., Cao, B., Nilsson, S.K., See, H.B., Pfleger, K.D.G., Roche, M., Gorry,
P.R., Pow, A., Viduka, K., Lim, K., Lu, B.G.C., Chang, D.H.C., Murray-Rust, T.,
Kvansakul, M., Perugini, M.A., Dogovski, C., Doerflinger, M., Zhang, Y., Parisi, K.,
Casey, J.L., Nuttall, S.D., Foley, M., 2016a. i-bodies, Human Single Domain Antibodies
That Antagonize Chemokine Receptor CXCR4. J Biol Chem 291, 12641–12657.
https://doi.org/10.1074/jbc.M116.721050

Griffiths, K., Habiel, D.M., Jaffar, J., Binder, U., Darby, W.G., Hosking, C.G., Skerra, A.,
Westall, G.P., Hogaboam, C.M., Foley, M., 2018. Anti-fibrotic Effects of CXCR4-

Targeting i-body AD-114 in Preclinical Models of Pulmonary Fibrosis. Sci Rep 8, 3212. https://doi.org/10.1038/s41598-018-20811-5

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., Hassabis, D., 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596, 583–589. https://doi.org/10.1038/s41586-021-03819-2

Kals, E., Kals, M., Lees, R.A., Introini, V., Kemp, A., Silvester, E., Collins, C.R., Umrekar, T., Kotar, J., Cicuta, P., Rayner, J.C., 2024. Application of optical tweezer technology reveals that PfEBA and PfRH ligands, not PfMSP1, play a central role in Plasmodium falciparum merozoite-erythrocyte attachment. PLoS Pathog 20, e1012041. https://doi.org/10.1371/journal.ppat.1012041

Kocken, C.H., van der Wel, A.M., Dubbeld, M.A., Narum, D.L., van de Rijke, F.M., van Gemert, G.J., van der Linde, X., Bannister, L.H., Janse, C., Waters, A.P., Thomas, A.W., 1998. Precise timing of expression of a Plasmodium falciparum-derived transgene in Plasmodium berghei is a critical determinant of subsequent subcellular localization. J Biol Chem 273, 15119–15124. https://doi.org/10.1074/jbc.273.24.15119

Lamarque, M., Besteiro, S., Papoin, J., Roques, M., Vulliez-Le Normand, B., Morlon-Guyot, J., Dubremetz, J.-F., Fauquenoy, S., Tomavo, S., Faber, B.W., Kocken, C.H., Thomas, A.W., Boulanger, M.J., Bentley, G.A., Lebrun, M., 2011. The RON2-AMA1 Interaction is a Critical Step in Moving Junction-Dependent Invasion by Apicomplexan Parasites. PLoS Pathog 7, e1001276.

Li, X., Chen, H., Oo, T.H., Daly, T.M., Bergman, L.W., Liu, S.-C., Chishti, A.H., Oh, S.S., 2004. A co-ligand complex anchors Plasmodium falciparum merozoites to the erythrocyte invasion receptor band 3. J Biol Chem 279, 5765–5771. https://doi.org/10.1074/jbc.M308716200

Li, Y., Yang, L., Yang, L.-Q., 2024. Effects of intrinsically disordered regions in gp120 underlying HIV neutralization phenotypes. Biochem Biophys Res Commun 709, 149830. https://doi.org/10.1016/j.bbrc.2024.149830

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30, 923–930.
https://doi.org/10.1093/bioinformatics/btt656

Liffner, B., Frölich, S., Heinemann, G.K., Liu, B., Ralph, S.A., Dixon, M.W.A., Gilberger,
T.W., Wilson, D.W., 2020. PfCERLI1 is a conserved rhoptry associated protein essential
for Plasmodium falciparum merozoite invasion of erythrocytes. Nat Commun 11.
https://doi.org/10.1038/s41467-020-15127-w

Livak, K.J., Schmittgen, T.D., 2001. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. Methods 25, 402–408.
https://doi.org/https://doi.org/10.1006/meth.2001.1262

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol 15, 550. https://doi.org/10.1186/s13059-
014-0550-8

Lu, C., Zheng, X., Zhang, W., Zhao, H., MacRaild, C.A., Norton, R.S., Zhuang, Y., Wang, J.,
Zhang, X., 2019. Interaction of merozoite surface protein 2 with lipid membranes. FEBS
Lett 593, 288–295. https://doi.org/10.1002/1873-3468.13320

MacRaild, C.A., Pedersen, M.Ø., Anders, R.F., Norton, R.S., 2012. Lipid interactions of the
malaria antigen merozoite surface protein 2. Biochim Biophys Acta 1818, 2572–8.
https://doi.org/10.1016/j.bbamem.2012.06.015

MacRaild, C.A., Zachrdla, M., Andrew, D., Krishnarjuna, B., Nováček, J., Žídek, L., Sklenář,
V., Richards, J.S., Beeson, J.G., Anders, R.F., Norton, R.S., 2015. Conformational
dynamics and antigenicity in the disordered malaria antigen merozoite surface protein 2.
PLoS One 10, e0119899. https://doi.org/10.1371/journal.pone.0119899

McCarthy, J.S., Marjason, J., Elliott, S., Fahey, P., Bang, G., Malkin, E., Tierney, E., Aked-
Hurditch, H., Adda, C., Cross, N., Richards, J.S., Fowkes, F.J.I., Boyle, M.J., Long, C.,
Druilhe, P., Beeson, J.G., Anders, R.F., 2011. A phase 1 trial of MSP2-C1, a blood-stage
malaria vaccine containing 2 isoforms of MSP2 formulated with montanide® ISA 720.
PLoS One 6. https://doi.org/10.1371/journal.pone.0024413

Ocampo, M., Urquiza, M., Guzmán, F., Rodriguez, L.E., Suarez, J., Curtidor, H., Rosas, J.,
Diaz, M., Patarroyo, M.E., 2003. Two MSA 2 peptides that bind to human red blood cells

1151  are relevant to Plasmodium falciparum merozoite invasion. The Journal of Peptide
1152  Research 55, 216–223. https://doi.org/10.1034/j.1399-3011.2000.00174.x

1153 Orlandi, P.A., Klotz, F.W., Haynes, J.D., 1992. A malaria invasion receptor, the 175-kilodalton
1154  erythrocyte binding antigen of Plasmodium falciparum recognizes the terminal Neu5Ac
1155  (alpha 2-3) Gal-sequences of glycophorin A. J Cell Biol 116, 901–909.

1156 Osier, F.H., Feng, G., Boyle, M.J., Langer, C., Zhou, J., Richards, J.S., McCallum, F.J.,
1157  Reiling, L., Jaworowski, A., Anders, R.F., Marsh, K., Beeson, J.G., 2014. Opsonic
1158  phagocytosis of Plasmodium falciparum merozoites: mechanism in human immunity and
1159  a correlate of protection against malaria. BMC Med 12, 108.
1160  https://doi.org/10.1186/1741-7015-12-108

1161 Osier, F.H.A., Murungi, L.M., Fegan, G., Tuju, J., Tetteh, K.K., Bull, P.C., Conway, D.J.,
1162  Marsh, K., 2010. Allele-specific antibodies to Plasmodium falciparum merozoite surface
1163  protein-2 and protection against clinical malaria. Parasite Immunol 32, 193–201.
1164  https://doi.org/10.1111/j.1365-3024.2009.01178.x

1165 Peng, D., Tarleton, R., 2015. EuPaGDT: a web tool tailored to design CRISPR guide RNAs
1166  for eukaryotic pathogens. Microb Genom 1. https://doi.org/10.1099/mgen.0.000033

1167 Persson, K.E.M., Fowkes, F.J.I., McCallum, F.J., Gicheru, N., Reiling, L., Richards, J.S.,
1168  Wilson, D.W., Lopaticki, S., Cowman, A.F., Marsh, K., Beeson, J.G., 2013. Erythrocyte-
1169  Binding Antigens of Plasmodium falciparum Are Targets of Human Inhibitory Antibodies
1170  and Function To Evade Naturally Acquired Immunity. The Journal of Immunology 191,
1171  785–794. https://doi.org/10.4049/jimmunol.1300444

1172 Persson, K.E.M., McCallum, F.J., Reiling, L., Lister, N.A., Stubbs, J., Cowman, A.F., Marsh,
1173  K., Beeson, J.G., 2008. Variation in use of erythrocyte invasion pathways by Plasmodium
1174  falciparum mediates evasion of human inhibitory antibodies. J Clin Invest 118, 342–351.
1175  https://doi.org/10.1172/JCI32138

1176 Reiling, L., Boyle, M.J., White, M.T., Wilson, D.W., Feng, G., Weaver, R., Opi, D.H., Persson,
1177  K.E.M., Richards, J.S., Siba, P.M., Fowkes, F.J.I., Takashima, E., Tsuboi, T., Mueller, I.,
1178  Beeson, J.G., 2019. Targets of complement-fixing antibodies in protective immunity
1179  against malaria in children. Nat Commun 10, 610. https://doi.org/10.1038/s41467-019-
1180  08528-z

Ressurreição, M., Thomas, J.A., Nofal, S.D., Flueck, C., Moon, R.W., Baker, D.A., van Ooij, C., 2020. Use of a highly specific kinase inhibitor for rapid, simple and precise synchronization of Plasmodium falciparum and Plasmodium knowlesi asexual blood-stage parasites. PLoS One 15, e0235798. https://doi.org/10.1371/journal.pone.0235798

Richards, J.S., Arumugam, T.U., Reiling, L., Healer, J., Hodder, A.N., Fowkes, F.J.I., Cross, N., Langer, C., Takeo, S., Uboldi, A.D., Thompson, J.K., Gilson, P.R., Coppel, R.L., Siba, P.M., King, C.L., Torii, M., Chitnis, C.E., Narum, D.L., Mueller, I., Crabb, B.S., Cowman, A.F., Tsuboi, T., Beeson, J.G., 2013. Identification and Prioritization of Merozoite Antigens as Targets of Protective Human Immunity to Plasmodium falciparum Malaria for Vaccine and Biomarker Development. The Journal of Immunology 191, 795 LP – 809.

RTSS Clinical Trials Partnership, 2015. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose in infants and children in Africa: final results of a phase 3, individually randomised, controlled trial. Lancet 386, 31–45. https://doi.org/http://dx.doi.org/10.1016/S0140-6736(15)60721-8

Sagara, I., Dicko, A., Ellis, R.D., Fay, M.P., Diawara, S.I., Assadou, M.H., Sissoko, M.S., Kone, M., Diallo, A.I., Saye, R., Guindo, M.A., Kante, O., Niambele, M.B., Miura, K., Mullen, G.E.D., Pierce, M., Martin, L.B., Dolo, A., Diallo, D.A., Doumbo, O.K., Miller, L.H., Saul, A., 2009. A randomized controlled phase 2 trial of the blood stage AMA1-C1/Alhydrogel malaria vaccine in children in Mali. Vaccine 27, 3090–3098. https://doi.org/10.1016/j.vaccine.2009.03.014

Salanti, A., Staalsoe, T., Lavstsen, T., Jensen, A.T.R., Sowa, M.P.K., Arnot, D.E., Hviid, L., Theander, T.G., 2003. Selective upregulation of a single distinctly structured var gene in chondroitin sulphate A-adhering Plasmodium falciparum involved in pregnancy-associated malaria. Mol Microbiol 49, 179–191. https://doi.org/10.1046/j.1365-2958.2003.03570.x

Sanders, P.R., Kats, L.M., Drew, D.R., O'Donnell, R.A., O'Neill, M., Maier, A.G., Coppel, R.L., Crabb, B.S., 2006. A Set of Glycosylphosphatidyl Inositol-Anchored Membrane Proteins of Plasmodium falciparum Is Refractory to Genetic Deletion. Infect Immun 74, 4330–4338. https://doi.org/10.1128/IAI.00054-06

Scally, S.W., Triglia, T., Evelyn, C., Seager, B.A., Pasternak, M., Lim, P.S., Healer, J., Geoghegan, N.D., Adair, A., Tham, W.-H., Dagley, L.F., Rogers, K.L., Cowman, A.F.,

2022. PCRCR complex is essential for invasion of human erythrocytes by Plasmodium

falciparum. Nat Microbiol 7, 2039–2053. https://doi.org/10.1038/s41564-022-01261-2

Sheehy, S.H., Duncan, C.J.A., Elias, S.C., Choudhary, P., Biswas, S., Halstead, F.D., Collins,

K.A., Edwards, N.J., Douglas, A.D., Anagnostou, N.A., Ewer, K.J., Havelock, T.,

Mahungu, T., Bliss, C.M., Miura, K., Poulton, I.D., Lillie, P.J., Antrobus, R.D., Berrie,

E., Moyle, S., Gantlett, K., Colloca, S., Cortese, R., Long, C.A., Sinden, R.E., Gilbert,

S.C., Lawrie, A.M., Doherty, T., Faust, S.N., Nicosia, A., Hill, A.V.S., Draper, S.J., 2012.

ChAd63-MVA-vectored blood-stage malaria vaccines targeting MSP1 and AMA1:

assessment of efficacy against mosquito bite challenge in humans. Mol Ther 20, 2355–

2368. https://doi.org/10.1038/mt.2012.223

Smythe, J.A., Coppel, R.L., Day, K.P., Martin, R.K., Oduola, A.M., Kemp, D.J., Anders, R.F.,

1991. Structural diversity in the Plasmodium falciparum merozoite surface antigen 2.

Proceedings of the National Academy of Sciences 88, 1751 LP – 1755.

https://doi.org/10.1073/pnas.88.5.1751

Srinivasan, P., Beatty, W.L., Diouf, A., Herrera, R., Ambroggio, X., Moch, J.K., Tyler, J.S.,

Narum, D.L., Pierce, S.K., Boothroyd, J.C., Haynes, J.D., Miller, L.H., 2011. Binding of

Plasmodium merozoite proteins RON2 and AMA1 triggers commitment to invasion.

Proceedings of the National Academy of Sciences 108, 13275–13280.

https://doi.org/10.1073/pnas.1110303108

Stanisic, D.I., Richards, J.S., McCallum, F.J., Michon, P., King, C.L., Schoepflin, S., Gilson,

P.R., Murphy, V.J., Anders, R.F., Mueller, I., Beeson, J.G., 2009. Immunoglobulin G

Subclass-Specific Responses against Plasmodium falciparum Merozoite Antigens Are

Associated with Control of Parasitemia and Protection from Symptomatic Illness. Infect

Immun 77, 1165–1174. https://doi.org/10.1128/IAI.01129-08

Stejskal, L., Kalemera, M.D., Lewis, C.B., Palor, M., Walker, L., Daviter, T., Lees, W.D.,

Moss, D.S., Kremyda-Vlachou, M., Kozlakidis, Z., Gallo, G., Bailey, D., Rosenberg, W.,

Illingworth, C.J.R., Shepherd, A.J., Grove, J., 2022. An entropic safety catch controls

hepatitis C virus entry and antibody resistance. Elife 11.

https://doi.org/10.7554/eLife.71854

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics

Analysis Version 11. Mol Biol Evol 38, 3022–3027.

https://doi.org/10.1093/molbev/msab120

Taylor, H.M., McRobert, L., Grainger, M., Sicard, A., Dluzewski, A.R., Hopp, C.S., Holder,
A.A., Baker, D.A., 2010. The malaria parasite cyclic GMP-dependent protein kinase plays
a central role in blood-stage schizogony. Eukaryot Cell 9, 37–45.
https://doi.org/10.1128/EC.00186-09

Thera, M.A., Doumbo, O.K., Coulibaly, D., Laurens, M.B., Ouattara, A., Kone, A.K., Guindo,
A.B., Traore, K., Traore, I., Kouriba, B., Diallo, D.A., Diarra, I., Daou, M., Dolo, A.,
Tolo, Y., Sissoko, M.S., Niangaly, A., Sissoko, M., Takala-Harrison, S., Lyke, K.E., Wu,
Y., Blackwelder, W.C., Godeaux, O., Vekemans, J., Dubois, M.-C., Ballou, W.R., Cohen,
J., Thompson, D., Dube, T., Soisson, L., Diggs, C.L., House, B., Lanar, D.E., Dutta, S.,
Heppner, D.G., Plowe, C. V., 2011. A Field Trial to Assess a Blood-Stage Malaria
Vaccine. New England Journal of Medicine 365, 1004–1013.
https://doi.org/10.1056/NEJMoa1008115

Weiss, G.E., Gilson, P.R., Taechalertpaisarn, T., Tham, W.-H.H., de Jong, N.W.M.M., Harvey,
K.L., Fowkes, F.J.I.I., Barlow, P.N., Rayner, J.C., Wright, G.J., Cowman, A.F., Crabb,
B.S., 2015. Revealing the Sequence and Resulting Cellular Morphology of Receptor-
Ligand Interactions during Plasmodium falciparum Invasion of Erythrocytes. PLoS
Pathog 11, 284–295. https://doi.org/10.1371/journal.ppat.1004670

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wilson, D.W., Goodman, C.D., Sleebs, B.E., Weiss, G.E., de Jong, N.W.M., Angrisano, F.,
Langer, C., Baum, J., Crabb, B.S., Gilson, P.R., McFadden, G.I., Beeson, J.G., 2015.
Macrolides rapidly inhibit red blood cell invasion by the human malaria parasite,
Plasmodium falciparum. BMC Biol 13, 52. https://doi.org/10.1186/s12915-015-0162-0

World Health Organization, 2024. World malaria report 2024: addressing inequity in the global
malaria response. World Health Organization, Geneva.

Yang, X., Adda, C.G., MacRaild, C.A., Low, A., Zhang, X., Zeng, W., Jackson, D.C., Anders,
R.F., Norton, R.S., 2010. Identification of key residues involved in fibril formation by the
conserved N-terminal region of Plasmodium falciparum merozoite surface protein 2
(MSP2). Biochimie 92, 1287–95. https://doi.org/10.1016/j.biochi.2010.06.001

Zerebinski, J., Margerie, L., Han, N.S., Moll, M., Ritvos, M., Jahnmatz, P., Ahlborg, N.,
Ngasala, B., Rooth, I., Sjöberg, R., Sundling, C., Yman, V., Färnert, A., Plaza, D.F., 2024.
Naturally acquired IgG responses to Plasmodium falciparum do not target the conserved

termini of the malaria vaccine candidate Merozoite Surface Protein 2. Front Immunol 15,
1501700. https://doi.org/10.3389/fimmu.2024.1501700

Zhang, X., Adda, C.G., Low, A., Zhang, J., Zhang, W., Sun, H., Tu, X., Anders, R.F., Norton,
R.S., 2012. Role of the helical structure of the N-terminal region of plasmodium
falciparum merozoite surface protein 2 in fibril formation and membrane interaction.
Biochemistry 51, 1380–1387. https://doi.org/10.1021/bi201880s

Zhang, X., Perugini, M.A., Yao, S., Adda, C.G., Murphy, V.J., Low, A., Anders, R.F., Norton,
R.S., 2008. Solution conformation, backbone dynamics and lipid interactions of the
intrinsically unstructured malaria surface protein MSP2. J Mol Biol 379, 105–21.
https://doi.org/10.1016/j.jmb.2008.03.039

1286