Spontaneous spiking statistics form unique area-specific fingerprints and reflect the hierarchy of cerebral cortex

Aitor Morales-Gregorio[1,2*], Robin Gutzen[1,3], Sofia Paneri[4,5], Panagiotis Sapountzis[4,5], Alexander Kleinjohann[1,6], Sonja Grün[1,7,8], Alexa Riehle[9], Xing Chen[10,11], Thomas Brochier[9], Georgia G. Gregoriou[4,5], Bjørg E. Kilavik[9], Sacha J. van Albada[1,12]

[1]Institute for Advanced Simulation (IAS-6), Jülich Research Centre, Jülich, Germany. [2]Faculty of Mathematics and Physics, Charles University, Prague, Czechia. [3]Center for Data Science, New York University, New York, USA. [4]Institute of Applied and Computational Mathematics, Foundation for Research and Technology Hellas, Heraklion, Greece. [5]Department of Basic Sciences, Medical School, University of Crete, Heraklion, Greece. [6]RWTH Aachen University, Aachen, Germany. [7]JARA-Institut Brain Structure-Function Relationships (INM-10), Jülich Research Centre, Jülich, Germany. [8]Theoretical Systems Neurobiology, RWTH Aachen University, Aachen, Germany. [9]Institut de Neurosciences de la Timone (INT), Aix-Marseille Université, CNRS, Marseille, France [10]Department of Vision & Cognition, Netherlands Institute for Neuroscience, Meibergdreef 47, 1105 BA, Amsterdam, Netherlands [11]Department of Ophthalmology, University of Pittsburgh, 1622 Locust St, PA 15219, Pittsburgh, US [12]Institute of Zoology, University of Cologne, Cologne, Germany.

[*]Corresponding author: aitor.morales-gregorio@matfyz.cuni.cz

May 13, 2025

The cerebral cortex, from sensory to higher cognitive areas, is hierarchically organised[1–4]. Several dynamical[5–7] and anatomical[1–4] measures, such as timescales and neurotransmitter receptor expression, have independently been linked to the cortical hierarchy. However, a systematic quantitative link between anatomical markers of the cortical hierarchy and single-neuron dynamics has not yet been established. Here, we hypothesise that the single-neuron spontaneous spiking statistics uniquely characterise each cortical area, and that they quantitatively correlate with the cortical hierarchy. We consider the spontaneous activity of neurons in seven cortical areas (V1, V4, DP, 7A, M1, PMd, PFC) in macaques[8–11] in the eyes-open and eyes-closed conditions. First, we show that the firing rate, inter-spike interval variation, and cross-correlation form a unique fingerprint of the cortical areas, but only when considering them in combination. Second, we find that the differences between the spiking statistics strongly correlate with multiple anatomical markers[1,2,4,12–16] of the cortical hierarchy, especially in the eyes-closed condition. We also observe a correlation between autocorrelation timescales and the anatomical hierarchy, consistent with previous findings[5,17,18]. In conclusion, we demonstrate that spontaneous single-neuron spiking activity reflects the hierarchical organisation of the cerebral cortex: distinct spiking statistics for hierarchically distant areas; similar statistics for nearby areas. Our results thus add a new dynamical dimension to the concept of the cortical hierarchy.

Keywords: electrophysiology, spontaneous activity, macaque, cerebral cortex, hierarchy, anatomy

Introduction

Information in the cerebral cortex is processed in multiple stages, following a hierarchy[1]. The cortical hierarchy is thought to support information processing, with lower-order regions handling localized sensory inputs and higher-order regions integrating signals into abstract representations. Several anatomical features, such as structural connectivity[1–3,19], neuron[12,13] and receptor[4,14,20] densities, spine counts[15,21], and myelination[16], reflect the cortical hierarchy. All these anatomical markers of the hierarchy are known to strongly correlate with each other[4,14], suggesting a complex relationship between the structural anatomy and information processing along the hierarchy.

Neural activity also varies along the cortical hierarchy. For example, inter-spike interval (ISI) variation[22] in spontaneous activity differs significantly across areas at different hierarchical levels and is surprisingly consistent across mammals for the same area[6]. Functional connectivity[23,24] and visual response measures[7] have been reported to align with the hierarchical organisation. Autocorrelation timescales also follow a hierarchical gradient in macaques[5,25–27], humans[17,26,28,29], rats[27], and less clearly in mice[7,18,26,27]. While prior studies have independently examined anatomical markers and neural dynamics, the relationship between these features remains largely unexplored. Specifically, no study reports systematic quantitative correlation between spontaneous spiking activity at the single-neuron level and cortical hierarchy across many areas in primates.

Here, we test and confirm the hypothesis that cortical spontaneous activity forms a unique fingerprint of each area and follows an anatomical hierarchical gradient. We use extracellular electrophysiological data from seven cortical areas (V1, V4, DP, 7A, M1, PMd, PFC) in several macaques[8–11] ($N = 6$, *Macaca mulatta*), and measure single-neuron statistics; i.e. firing rate, ISI variation, and cross-correlation. We study spontaneous activity because it should be less biased toward external stimuli and more closely related to the underlying anatomical structure than evoked activity. We test whether the spontaneous activity varies across cortical areas, and find significant differences in the spontaneous spiking statistics across all areas, especially when the spiking statistics are considered multivariately. These differences strongly correlate with several anatomical markers of the cortical hierarchy: neuron densities[12], receptor densities[4], spine counts[15], and hierarchical levels estimated from laminar cortico-cortical connectivity[2]. We further confirm[5,25–27] that autocorrelation timescales also exhibit a consistent correlation with the hierarchy. All in all, the results suggest a key role for cortical anatomy in shaping spontaneous spiking activity and add a new dynamical dimension to the concept of cortical hierarchies.

Results

Multivariate spiking statistics of spontaneous activity

Does spontaneous neural activity systematically vary across cortical areas? To answer this question, we studied spontaneous intracortical electrophysiology recordings from various experiments in macaques[8–11]. The data were recorded from seven areas along the visuomotor hierarchy (V1, V4, DP, 7A, PFC, PMd, and M1). Approximate recording locations are shown in Figure 1a. The data were spike sorted based on their waveforms to retain only single-unit activity (SUA), after which any SUA suspected of originating from hardware noise or cross-talk was excluded[30] (Figure 1b). The different experiments varied in recording length (Figure S4a,b), hence we split the data into 5-s segments to allow for unbiased comparisons. To reduce nonstationarities, segments were only included if the monkey remained in the same behavioural state (eyes-open or eyes-closed) throughout the whole segment. See Electrophysiological data collection of spontaneous activity for details on the data collection and preprocessing.

To characterize the spiking activity, we calculated single-neuron statistics for each data segment and assembled them into a multivariate point cloud (Figure 1c). The statistics were firing rate (FR), revised local variation (LvR) of inter-spike intervals[22], and the mean and standard deviation of each neuron's cross-correlation with all other neurons in the same area and subject (CC_{avg} and CC_{std}). The LvR measures the irregularity of spike trains and it has been shown to vary across brain areas in several species[6]. The cross-correlation statistics are related to the dimensionality of the population activity[31–35]. Since the FR, CC_{avg} and CC_{std} had highly skewed distributions, we took their base-10 logarithm, rendering the distributions more symmetric. See Statistics of spiking neuronal data for details.

Note that we did not include the single-neuron timescales here, because they are difficult to estimate from short recordings[36–38], as we illustrate in Figure S4. Instead, we studied timescales on their own (see Autocorrelation timescales also reflect hierarchy).

We visualized the four-dimensional cloud of single-neuron statistics (Figure 1c) as a pair plot, where each subplot displays two statistics, and the diagonal subplots show the kernel density estimates (KDE) of individual metric distributions. The spiking statistics for each animal and area are shown individually in Figure S1-S3. To estimate whether our chosen statistics were relevant and linearly independent, we estimated the explained

Figure 1: Sample of available experimental data and their spontaneous spiking statistics. Same color scheme across panels. **a)** Schematic representation of the data recording locations. **b)** Sample recordings of simultaneous spike trains from each dataset for a 5-s segment; superscripts indicate the subject identity. **c)** Pair plot of single-neuron spiking statistics. Panels along the diagonal show the kernel density estimates (KDE) of individual distributions, whereas off-diagonal panels show scatter plots where each point is calculated from a 5-s spike train from one neuron. **d)** (Top) Variance explained by each of the four principal components (PC) of the multivariate spiking statistics. (Bottom) Loadings of each spiking statistic (average across datasets ± standard deviation). PCA performed separately for each dataset.

variance and loadings using PCA. The principal components each explained a substantial proportion of variance (at least 5–20%, (Figure 1d), demonstrating that all the selected spiking statistics capture independent aspects of neural dynamics.

The experiments were highly heterogeneous, with different subjects, hardware setups, neuron counts, and recording durations. Calculating the multivariate distribution of single-neuron spontaneous spiking statistics allows comparison of heterogeneous datasets by summarizing them in a common space.

Multivariate spiking statistics form a unique fingerprint of each cortical area

Visual inspection of the spiking statistics (Figure 1c) revealed noticeable differences in their distributions across brain areas. We hypothesized that the spiking statistics differ between areas, due to the differences in the underlying anatomical structure such as connectivity, neuron densities, receptors etc. Moreover, the inclusion of multiple spiking statistics provides a more comprehensive characterization of each area, since each statistic captures some unique information about the area. Hence, we expected the combined spiking statistics to show more differences between areas than each single statistic on their own. For example, while FR may not always uniquely characterise each area, the combination of all four statistics could. We tested this hypothesis, by testing whether the means of the distributions of spiking statistics were significantly different across areas and subjects. First, we performed a pairwise univariate analysis of variance (ANOVA) for each of our four statistics (Figure 2a–d). Second, we performed pairwise multivariate ANOVA (MANOVA) tests (Figure 2e), where all four spiking statistics were considered in combination. See Statistical testing with ANOVA and MANOVA for details about the tests.

The univariate ANOVA tests (Figure 2a–d) revealed that no single spiking statistic consistently differentiates activity across all areas. However, when all the spiking statistics were aggregated and tested as an ensemble with the MANOVA test (Figure 2e) nearly all dataset pairs became statistically different. The only exception was the

Figure 2: Uni- and multivariate pairwise comparisons of spontaneous spiking statistics. We test the null hypothesis that two or more datasets have the same mean. In all panels, lower triangular entries show the base-10 logarithm of the p-values and the upper triangular entries show the F-statistic. Significance levels ($\alpha = 0.05/k$) are corrected for multiple testing following the Bonferroni correction ($k = 630$, including the tests from Figure S8a). In all panels, the significance level is color-coded, with red p-values (orange F-statistics) denoting non-significant results. **a-d)** Univariate pairwise analysis of variance (ANOVA) tests. **e)** Multivariate pairwise analysis of variance (MANOVA) tests.

MANOVA test between $V4^F \leftrightarrow V4^L$, for data originating from the same area in two different subjects recorded with different hardware and in different laboratories.

The statistical tests support our hypothesis that the spiking statistics are area-specific, but only if the spiking statistics are considered multivariately. Thus, the multivariate spiking statistics form a dynamical fingerprint of each cortical area.

Dynamical dissimilarity reveals hierarchical organisation of cortical areas

The statistical tests so far revealed that the multivariate spiking statistics differ between areas, but they provided no information as to how similar or different they are. To quantify those differences, we introduce the dynamical dissimilarity, which we define as the Wasserstein distance (WS) between the multivariate spiking statistics of two areas (Figure 3a). We chose to use the Wasserstein distance because it is symmetric, non-parametric, and numerically robust on point clouds. The dynamical dissimilarity takes a positive value between zero (exactly the same) and infinity, and can be applied to compare any two point distributions both univariately or multivariately. Low values of the dynamical dissimilarity indicate high similarity. We calculated all within- and across-area dynamical dissimilarities (Figure 3a), and found that the within-area dissimilarities were consistently smaller than across-area dissimilarities (Figure 3a inset). See Dynamical dissimilarity for comparison of multivariate and univariate distributions for a detailed description.

Moreover, the dynamical dissimilarities were highly structured (Figure 3a). They were smallest for neighbouring ($7A \leftrightarrow DP$ or $PMd \leftrightarrow M1$) or equivalent areas in different subjects ($M1/PMd^N \leftrightarrow M1/PMd^E$ or $V4^F \leftrightarrow V4^L$), and were largest between functionally distant areas (e.g. $V1 \leftrightarrow M1/PMd$). This trend suggests a hierarchical organisation of dynamical differences.

To test the hierarchical organisation of dynamical dissimilarities we compared them to inter-area differences in several known anatomical markers of hierarchy in the primate cerebral cortex: neuron densities[12], receptor densities[4], spine counts[15], and hierarchical levels estimated from laminar cortico-cortical connectivity[2]. See

Figure 3: Dynamical dissimilarity between spiking activity statistics strongly correlates with anatomical markers of cortical hierarchy. **a)** Dynamical dissimilarity between all datasets. The inset diagram schematically illustrates the Wasserstein distance (WS). The inset histogram shows the distribution of dynamical dissimilarity within and across datasets. **b)** Anatomical dissimilarity between all areas, defined from the combination of neuron density, receptor density, spine counts, and connectivity patterns. **c)** Dynamical dissimilarity strongly correlates with anatomical dissimilarity; Pearson correlation coefficient and corresponding t-test p-value are shown. Note that myelination was excluded from the anatomical dissimilarity; including myelination slightly lowers the correlation ($r = 0.74$, $p < 10^{-8}$). **d)** Pearson r between anatomical dissimilarity and modified versions of the dynamical dissimilarity: excluding one metric at a time (leave-one-out) or only including a single metric. Stars indicate result of t-test: $p > 0.05/k$ (n.s.), $0.05/k > p > 0.01/k$ (*), $0.01/k > p > 0.001/k$ (**), $p < 0.001/k$ (***). Bonferroni corrected for multiple testing ($k = 61$). **e)** Estimates of the hierarchy from the dynamical and anatomical dissimilarities. **f)** Boxplot of the spiking statistics for each area compared to their position in the anatomical hierarchy. Weighted least squares (WLS) fit, its coefficient of determination R^2 and t-test p-value are shown.

 for a detailed description of the anatomical data.

We related the anatomical markers to the dynamical dissimilarity by calculating the Euclidean distance

between the 4D data points representing the anatomical markers for each pair of areas (Figure 3b). Each marker

was normalized between 0 and 1 prior to measuring the anatomical dissimilarity. We found a strong correlation

(Pearson's $r = 0.78$) between the dynamical and anatomical dissimilarities (Figure 3c).

The receptor densities were available for 14 different receptor types, but we included only their average in our

anatomical dissimilarity, so as not to eclipse all other anatomical markers. We also tested the individual receptor

densities and most of them strongly correlated with the full dynamical dissimilarity (Figure S5).

To analyze the individual contributions of each spiking statistic, we compared the anatomical dissimilarity to

the dynamical dissimilarity when one metric was excluded ('Exclude one metric' in Figure 3d) and also to the

dynamical dissimilarity when a single spiking statistic was included ('Only one metric' in Figure 3d). Furthermore,

we extended the comparisons to each individual anatomical marker (Figure 3d); corresponding scatter plots

are shown in Figure S6. This analysis reveals which spiking statistic relates most to which anatomical feature. Notably, log(FR) did not correlate with any anatomical markers when considered univariately. Furthermore, we also tested the correlation with myelination levels[16], but found no significant relationship with any spiking statistics. Thus, we did not include myelination in the anatomical dissimilarity.

We embed the dynamical and anatomical dissimilarities into a 1-dimensional line, representing the hierarchy (see Estimation of hierarchies from dynamical and anatomical dissimilarities, Figure 3e). A similar (but not equal) area ordering emerged from both dynamics and anatomy (Figure 3e).

To study how dynamics change along the hierarchy we compared the mean of each spiking statistic against the anatomical hierarchy using Weighted Least Squares (WLS), weighted by the inverse of the variance of each statistic (Figure 3f). The WLS indicated that LvR decreases along the hierarchy: the spike trains in V1 were very irregular (high LvR), and the spike trains in motor areas tend to be more regular (lower LvR). Furthermore, the WLS suggested that $\log(CC_{avg})$ and $\log(CC_{std})$ decrease along the hierarchy. No significant trend was found for log(FR).

In conclusion, the spontaneous spiking statistics robustly correlate with anatomical markers of the cortical hierarchy, strongly supporting our hypothesis that cortical spontaneous activity follows an anatomical hierarchical gradient.

Correspondence between dynamics and anatomy is strongest with the eyes closed, and nearly disappears with the eyes open

Figure 4: Dynamical dissimilarity changes between eyes-open (EO) and eyes-closed (EC) epochs. **a** Difference between the EO and EC dynamical dissimilarity. Negative values (blue) indicate higher dissimilarity in EC, conversely positive values (red) indicate higher dissimilarity in EO. **b** Correlation between dynamical and anatomical dissimilarities in EC and EO. The correlation is much weaker for EO. **c** Histograms of the dynamical dissimilarity distributions for EO and EC. Dashed lines indicate the mean value. The EO and EC distributions are not statistically different (two sided t-test).

The results until this point in the paper combined all behavioural states. However, the spiking activity is known to vary between different behaviours[35,39]. We expect stronger external inputs during the eyes-open (EO) than eyes-closed (EC) periods, which would globally spread across the entire brain and modify the spontaneous activity. During EC, the activity would more closely reflect the local anatomical structure of each area. Analyzing the differences between EO and EC thus provides potential information about the relative contributions of local and global factors to the observed hierarchical organisation. We thus split the activity by behaviour and study the changes in the dynamical dissimilarity (Figure 4).

First, we measured the dynamical dissimilarities for samples during eyes-open (EO) or eyes-closed (EC) periods; movement (M) periods could not be analyzed as there were too few of them. We found that visual areas, especially V1, have higher dynamical dissimilarity in EC than EO; whereas motor areas have higher dissimilarities in EO than EC (Figure 4a). Second, we compared the correlation between dynamical and anatomical dissimilarities in EC and EO (Figure 4b). Notably, this correlation is much stronger during EC ($r = 0.81$, $p < 10^{-10}$), than EO ($r = 0.42$, $p = 0.004$). This is in line with our hypothesis that externally driven input during EO would suppress the dynamics originating from local anatomy. In spite of these differences, the distribution of dynamical dissimilarities is statistically indistinguishable between EC and EO (t-test in Figure 4c). Therefore, the dynamics during both EC and EO are different across areas (confirmed by MANOVA tests in Figure S8a), yet only the EC dynamics strongly correlate with the anatomical differences.

To fully understand the changes to dynamics caused by behaviour we also compared the univariate distributions of the spiking statistics between EC and EO (Figure S8b,c). We found the biggest changes in the average cross-

[143] correlation $\log(\mathrm{CC}_{\mathrm{avg}})$ especially for $\mathrm{V1^L}$, $\mathrm{V4^L}$ and $\mathrm{M1/PMd^{N\&E}}$ (Figure S8c,d), in agreement with previous
[144] reports[35,39].

Autocorrelation timescales also reflect hierarchy

Figure 5: Autocorrelation timescales vary across the cerebral cortex and correlate with the anatomical hierarchy.
a) Autocorrelation function for single neurons with an exponential fit for each area. The mean autocorrelation across neurons (dark line) and 5th to 95th percentile (shading) is shown. Insets show the distribution of the estimated timescales and the number of single neurons with a valid fit ($R^2 > 0.8$). **b)** Boxplot of the estimated timescales for each area compared to their position in the anatomical hierarchy. WLS fit, its R^2 and t-test p-value are shown. **c)** Timescale dissimilarity: Wasserstein distance between the timescale distributions. **d)** Pearson correlation coefficient between the timescale dissimilarity and other metrics: dynamical dissimilarity, anatomical dissimilarity, and inter-area differences in all the individual anatomical markers. Stars indicate result of t-test: $p > 0.05/k$ (n.s.), $0.05/k > p > 0.01/k$ (*), $0.01/k > p > 0.001/k$ (**), $p < 0.001/k$ (***). Bonferroni corrected for multiple testing ($k = 61$).

[145]   Autocorrelation timescales are a common metric for cortical hierarchy[5,17]. Previous studies have shown[5,17]
[146] that sensory areas have shorter timescales, whereas areas higher in the hierarchy have longer timescales. Our
[147] dynamical dissimilarity does not include the autocorrelation timescales, because these are notoriously difficult
[148] to estimate from limited data[36,37] (see Figure S4). To overcome this problem, we estimated the timescales
[149] using longer data segments ($t_{\mathrm{segment}} = 300$ s). We fitted an exponential function to the autocorrelation function
[150] (Figure 5a, see sample fits in Figure S9), analogous to previous studies[5,40]. See Autocorrelation timescales for a
[151] detailed description of the timescale estimation.
[152]   As expected, we observed shorter timescales in sensory areas and longer timescales in motor areas (Figure 5a,b).
[153] Indeed, the timescales have a positive correlation with the anatomical hierarchy (Figure 5b). To compare the
[154] timescales to the other spiking statistics we measured the timescale dissimilarity, by taking the Wasserstein distance
[155] between the timescale distributions of every pair of areas (Figure 5c). We found a significant correlation between
[156] the timescale dissimilarity and the other hierarchy markers (Figure 5d). As with the other spiking statistics
[157] (Figure 3d), timescale dissimilarities did not correlate significantly with myelination differences (Figure 5d). The
[158] correlation between the anatomical and timescale dissimilarities is very similar (Pearson's $r = 0.76$, Figure 5d)
[159] to the correlation between the anatomical and dynamical dissimilarities (Pearson's $r = 0.78$, Figure 3c). These
[160] results suggest that the timescales and the other spiking statistics are comparably robust indicators of the cortical

hierarchy; but the spiking statistics can be measured faster and more robustly using far less data than the timescales (Figure S4).

Discussion

This work has two main findings. First, we show that single-neuron spontaneous spiking statistics form a unique fingerprint of cortical areas, but only when considering multiple spiking statistics in combination. Second, our results reveal a strong correlation between the spontaneous spiking statistics and anatomical markers of cortical hierarchy, suggesting that the anatomical differences along the cortical hierarchy are imprinted within the spontaneous spiking activity, especially when the eyes are closed.

Our findings uncover rich and varied spiking statistics across different cortical areas, in agreement with previous reports[5,6,22]. Yet, the firing rate (FR) distributions were very similar across all cortical areas (Figure 2a). When considered alone, the FR did not correlate with any of the anatomical markers of the cortical hierarchy (Figure 3d). This suggests that the FR distribution from spontaneous activity is not predominantly shaped by the local structure of the cerebral cortex, even though the FR is known to vary in an area-specific way during task-related activity.

Furthermore, we found no significant correlation between the myelination of cortical areas and their dynamics (Figure 3d and Figure 5d). This is surprising, since the myelination is known to correlate with the other anatomical markers of the cortical hierarchy in macaques[4] and myelination has also been shown to correlate with autocorrelation timescales in human[17,29]. This discrepancy raises questions about the role of myelination in shaping cortical dynamics. A potential explanation could be that the commonly used myelination proxy (T1w/T2w MRI signal) can include traversing axons that do not connect to nor originate from the measured area, thus obscuring the hierarchy. Additional spiking data and new direct measurements of the myelination (e.g. histology or polarized light imaging[41]) could help clarify this inconsistency.

Topographic connectivity between areas is ubiquitous in the brain[42], such as retinotopic connectivity in visual areas[43] or somatotopic connectivity in motor and somatosensory areas[44]. Cortical sites sharing more direct connections are known to have more similar spiking dynamics[43]. However, it was not possible to take these topographic connections into account for this study.

The hierarchical order derived from dynamical and anatomical dissimilarities have a few notable differences (Figure 3e). The dynamical hierarchy positions of $M1^T$ and PMd^T are much lower than expected from the anatomical hierarchy. The reason for this could be that these two datasets include spiking activity from all cortical layers, whereas the other recordings contain data from a single layer. Since we did not have spiking data from all layers for most areas, we did not perform a layer-wise analysis.

The observed differences in spiking statistics between areas depend on both global influences from other brain areas and characteristics of each individual area. However, it is not possible to fully disentangle the global and local contributions *in vivo*. To prevent global stimulus-specific activity from dominating the dynamics, we study spontaneous activity, which should be less biased toward external stimuli. We also studied eyes-open and eyes-closed activity separately (Figure 4), revealing a stronger correlation between the dynamical and anatomical dissimilarities with the eyes closed. We believe the weaker or absent external stimuli to be a partial reason for this observation, suggesting that the changes caused by external visual stimuli modify the dynamics with the eyes open, suppressing the differences caused by the local structural differences between areas. If this is the case, then our results suggest that the global component to the dynamics is not hierarchical across areas, only the local component. To eliminate the global component completely and study local dynamics, future work could employ the approach and metrics introduced in the current study on *in vitro* brain slice activity.

The hierarchical organisation of timescales has recently come into question with contradicting results in mice[7,18]. However, timescales in spiking activity have been reported to vary along the cortical hierarchy in the macaque brain[5,25]. As expected, our analysis found a significant correlation between the anatomical markers of the hierarchy and timescales in macaques, confirming previous findings. Interestingly, Cusinato el al[29] report large changes to the hierarchy of timescales between wakefulness and different sleep stages from human intracortical EEG recordings. Particularly, they report that timescales estimated from gamma-band signals (30-80 Hz) exhibit a hierarchy of timescales in sleep, but not during wakefulness, similar to our observations for macaque dynamical dissimilarities between EO and EC (Figure 4). It was not possible to determine whether the macaques were asleep in our data; it is possible that they were drowsy or even entered the first stages of sleep in some cases. We did not test whether the timescales change with behaviour in our data, since the periods of EO and EC are relatively short and do not allow for reliable timescale estimates (Figure S4).

All our results focus on the pairwise differences between areas, e.g., the dynamical dissimilarity between V1 and M1 compared to the receptor density ratio between V1 and M1. However, the values for each area could carry similar information, e.g., the mean LvR in V1 compared to the mean receptor density in V1. Comparing the anatomical markers against the mean spiking statistics in each area (Figure S7) reveals some correlations, but

most correlations are not statistically significant. Notably, no single statistic significantly correlates with all the anatomical markers. Thus, inter-area differences in spiking statistics more robustly reflect hierarchical patterns.

Comparing brain dynamics usually involves some metric in high dimensions, such as representational similarity via cross-correlation[45], canonical correlation analysis[46], hyperalignment[47] or some shape metric[48] (e.g. Procrustes distance). All these approaches are usually applied to spatio-temporal activity data. In contrast, our work explores an alternative way of comparing brain activity data, collapsing the time component and focusing on the space of a few relevant spiking statistics (firing rate, inter-spike interval variability, cross-correlation). One strength of our approach is its ability to compare heterogeneous datasets with varying electrode counts, recording duration, and experimental setup. Despite the heterogeneous data, the spiking statistics reliably distinguish cortical areas from each other (Figure 2) and reveal their link to the anatomical hierarchy (Figure 3).

Moreover, our approach is well-suited for data-to-model comparisons, which is usually challenging due to the vast subsampling in brain recordings. The multivariate spiking statistics distribution is unique for each brain area (Figure 2), which could be used as a target distribution for parameter inference in spiking neuron models.

While this study focused on areas within the visuomotor processing stream, our methods could readily be applied to other brain areas, such as those in the auditory and somatosensory processing streams. This could potentially uncover analogies and key differences between sensory modalities.

All in all, our findings reveal a fundamental link between cortical structure and spontaneous dynamics along the cortical hierarchy.

Materials and methods

Electrophysiological data collection of spontaneous activity

All data were collected from the cerebral cortex of macaque monkeys (*Macaca Mulatta*, N = 6) in the resting state, reflecting spontaneous cortical activity. The macaques were sitting in a dim-lit room, with no particular task, while the continuous activity from the cerebral cortex was recorded. The relevant behaviour in each experiment was also tracked using video recordings and/or eye tracking systems.

Table 1: Summary of subjects and recordings included in this study. Identity of the subject, cortical area, number of sessions, number of simultaneously recorded electrode sites in each session, and number of detected single-unit activities (SUA) are indicated.

Subject	Area	# Sessions	# Contacts	# SUA
L	V1	1	896	239
	V4	1	128	81
F	V4	59	4	375
	PFC	59	4	209
Y	DP	1	36	39
	7A	1	36	47
T	M1	7	24	89
	PMd	11	24	57
N	M1/PMd	2	96	310
E	M1/PMd	2	96	277

Data from macaque L

The data from macaque L were recorded from visual areas V1 and V4 (N=1 subject, N=1 session of 20 min duration). Chronic recordings were made using 16 8×8 electrode Utah arrays (Blackrock microsystems), two of them in visual area V4 and the rest in the primary visual cortex (V1), with a total of 1024 electrodes. The electrodes were 1.5 mm long; thus, the recordings were likely made from the deeper layers L5 and L6. A full description of the experimental setup and the data collection and preprocessing has already been published[8].

Pupil position and diameter data were collected using an infrared camera to determine the direction of gaze and eye closure of the macaques. In addition to the spontaneous activity recordings, a visual response task was also performed. The visual response data were used to calculate the signal-to-noise ratio (SNR) of each electrode and all electrodes with an SNR lower than 2 were excluded from further analysis.

The raw data were spike-sorted using a semi-automatic workflow, as described in a previous publication[35].

All experimental and surgical procedures for macaque L complied with the NIH Guide for Care and Use of Laboratory Animals, and were approved by the institutional animal care and use committee of the Royal Netherlands Academy of Arts and Sciences (approval number AVD-8010020171046).

Data from macaque F

The data from macaque F were recorded from visual area V4 and prefrontal cortex (PFC) including Brodmann areas 8, 45A, and 46 below the principal sulcus (N=1 subject, N=59 sessions of 5 min duration). Acute recordings were made using the Omniplex system (Plexon Inc., Dallas, TX). On a given day, up to four glass-coated tungsten microelectrodes (impedance 1-1.5 MOhm, Alpha-Omega Engineering) were positioned through a grid system over each area (PFC and V4) and were advanced through the dura by a four-channel microdrive system (NAN Instruments, Nazareth, Israel). Signals were filtered between 300 Hz (4-pole high pass Bessel) and 8 kHz (2-pole high pass Bessel), amplified, and digitized at 40 kHz to obtain spike data. Recordings were carried out within the first 1-1.5mm after encountering the first activity; thus, the majority of data were obtained from the middle and superficial layers (L2/3/4). The spontaneous activity was recorded while the macaque was not performing any particular task. A full description of the experimental setup and the data collection and preprocessing—for the accompanying experiment in the same subject—has already been published[11].

Eye position was monitored by an infrared-based tracking system (ETL-200, ISCAN Inc., Woburn, MA) and was sampled at 240 Hz. The x and y position were recorded and used for behavioural segmentation into eyes-open and eyes-closed epochs. Offline spike sorting was carried out manually using template matching in Offline Sorter (Plexon Inc). Four to ten clean single units were identified per area and session.

All experimental and surgical procedures for macaque F were approved by the Institutional Experimental Protocol Evaluation Committee (Approval Number 108596/30-5-2017).

Data from macaque Y

The data from macaque Y were obtained from two chronic Utah arrays (Blackrock microsystems), one in dorsal prelunate cortex (area DP) and another one in parietal area 7A, for a total of 72 electrodes (N=1 subject, N=2 sessions of 10 min duration). The recording apparatus is described elsewhere[10]. The electrodes were 1 mm long, thus recording from the central layers, likely layer 4. The recording system recorded the electric potential at each electrode with a sampling rate of 30 kHz. Spontaneous activity data from the same experiment have previously been analysed and published[35].

The raw data were spike-sorted for this study using a semi-automatic workflow using spikeinterface[49]. After the automatic sorting the quality of the waveform clusters was manually evaluated and only clean single-unit activity (SUA) waveforms were included in this study.

All experimental and surgical procedures for macaque Y were approved by the local (INT, Marseille) ethical committee (C2EA 71; authorization Apafis#13894-2018030217116218v4) and conformed to the European and French government regulations.

Data from macaque T

The data from macaque T were recorded from premotor cortex (PMd) and primary motor cortex (M1) (N=1 subject, N=11 sessions of 15 min duration). Acute recordings were made with laminar probes (Plexon and Alpha Omega, 24 contacts, 100 and 200 μm pitch). The laminar probes enabled recording from across the cortical layers. The 200 μm pitch probes could record all layers simultaneously, while the 100 μm pitch probes did not span the entirety of the cortical gray matter. The motor cortex gray matter is known to be approximately 3.5 mm thick[50], with the superficial and deep layers roughly equal in thickness. A full description of the experimental setup and the data collection and preprocessing—for the accompanying experiment in the same subject—has already been published[51].

The raw data was spike sorted offline. Spike sorting identified 5-13 clean single units per probe and session.

In addition to the spiking data, surface Electromyography (EMG) of the contralateral deltoid muscle, the heart rate with an ear clip, and a video of the macaque behaviour were recorded. In all the behavioural videos the screen LEDs were used to send a 1 s long blink every minute that can be used to realign the video with the neural recordings. We performed a video-based segmentation into behavioural epochs: eyes-open, eyes-closed, and movement periods.

Care and treatment of macaque T during all stages of the experiments conformed to the European Commission Regulations (Directive 2010/63/EU on the protection of animals used for scientific purposes) applied to French laws (decision of the 1st of February 2013). The experimental protocol was evaluated by the local Ethics Committee (CEEA 071) and carried out in a licensed institution (B1301404) under the authorization 03383.02 delivered by the French Ministry of High Education and Research.

Data from macaques N & E

The data from macaques N & E were recorded from the interface between premotor (PMd) and primary motor (M1) cortex (N=2 subjects, N=2 sessions of 15-20 min duration per subject). The recordings were chronic using one 10x10 electrode Utah array (Blackrock microsystems).

A full description of the experimental setup and the data collection and preprocessing—for the accompanying experiment in the same subject—has already been published[9]. An extensive analysis of the spontaneous data has also been published[39].

In addition to the registration of brain activity, the monkey's behaviour was video recorded and synchronized with the electrophysiology recording.

All experimental and surgical procedures for macaque N & E were approved by the local (INT, Marseille) ethical committee (C2EA 71; authorization A1/10/12) and conformed to the European and French government regulations.

Spiking data preprocessing

To ensure reliable analysis, several preprocessing steps common to all datasets were taken.

The data were sliced into segments of 5-second length. The segments were fully contained within one behavioural epoch: eyes-open, eyes-closed, or movement. The behaviour was estimated from video-tracking, eye-tracking and/or muscle activity, depending on the available setup in each experiment. Spontaneous spiking

activity is known to vary strongly between different behaviours[35,39]. Including samples uniquely in one epoch ensured that major activity transients from behavioural transitions were excluded.

Furthermore, electrophysiology recordings are prone to artifacts from external noise and cross-talk between recording channels[30]. These artifacts can be detected as higher-than-chance synchronous activity at sampling resolution (30 kHz)[8,30]. Using the methods proposed by Oberste-Frielinghaus et al[30], we estimated the excess synchrony and removed all units suspected to originate from common noise or cross-talk, which would especially affect the correlation statistics in our analysis.

Statistics of spiking neuronal data

The experimental spike trains were sliced into 5-second segments. We computed a number of statistics for each one of these data samples and constructed a multivariate cloud of neuronal spiking statistics. These multivariate spiking statistics are used to characterise a given area and assess the similarities across areas.

We used the following statistics of spike trains:

- Mean firing rate (FR), given as the total number of spikes divided by the duration of the recording sample. Since the FR has a right-skewed distribution, we take its logarithm throughout this paper. The distribution of log(FR) better resembles a normal distribution.

- Revised local variation of the inter-spike intervals (LvR)[6],

$$\text{LvR} = \frac{3}{N-1} \sum_{i=1}^{N-1} \left(1 - \frac{4 I_i I_{i+1}}{(I_i + I_{i+1})^2} \right) \left(1 + \frac{4R}{I_i + I_{i+1}} \right) \tag{1}$$

  where N is the total number of spikes, I_i is the ith inter-spike interval, and $R = 5$ ms is the refractoriness constant. A value of $\text{LvR} = 0$ corresponds to perfectly rhythmic spiking, whereas $\text{LvR} = 1$ corresponds to the irregularity of a Poisson point process.

- The average of the Pearson cross-correlation of the spike counts (bin width = 10 ms) with all other simultaneously recorded neurons (CC_{avg}). This metric quantifies whether the measured neuron coordinates its activity with many of the other neurons (i.e. if it is a chorister) or if it is spiking independently (i.e. is a soloist)[31].

- The standard deviation of the Pearson cross-correlation of the spike counts (bin width = 10 ms) with all other simultaneously recorded neurons (CC_{std}).

Further statistics were considered but not used in our analysis, because they were strongly correlated with other statistics. If strongly correlated statistics were introduced, some spiking properties would be over-represented. For example, the coefficient of variation (Cv, Cv2)[52] and local variation (Lv)[22] of the inter-spike intervals are excluded due to their similarity to the LvR. Introducing all of Cv, Cv2, Lv and LvR would significantly increase the representation of the inter-spike interval variation, without a large increase in the explained variance. Another statistic that was considered was the spike-triggered population response (stPR)[31], but it was not included due to its similarity with CC_{avg}.

To exclude outliers, potential artifacts, or numerical issues the statistics were bounded to the following empirically determined ranges: $\text{FR} > 1$ spikes/s, $\text{LvR} \in [0,3]$, $\text{CC}_{\text{avg}} > 10^{-4}$, and $\text{CC}_{\text{std}} > 0.01$. We ensured that the selection of these ranges did not distort the underlying distributions or truncate any long tail, instead excluding only outliers. See Figure S1-S3 for the full distribution of included spiking statistics for all datasets.

All metrics were computed using their implementation in the elephant toolbox[53], within the NetworkUnit reproducible testing framework[54].

Statistical testing with ANOVA and MANOVA

For all ANOVA and MANOVA tests, we used a fixed number of points from the multivariate spiking statistics ($N = 500$, randomly sampled), since the tests are robust to the normality assumption if the samples are large (generally $N > 30$) and robust to the assumption of equal (co-)variance (homoscedasticity) if the sample sizes are equal[55]. The significance level was Bonferroni corrected for multiple testing ($k = 90$ experiment pairs $\cdot$ 5 test types $= 630$), such that $\alpha = 0.05/k = 1.11 \cdot 10^{-4}$.

Dynamical dissimilarity for comparison of multivariate and univariate distributions

To compare the distributions of spiking statistics to each other we used the Wasserstein distance (WS). The
WS is defined as an optimal transport distance between two observed distributions. Each multivariate spiking
statistics cloud is an $N \times M$ matrix of N samples (spike train segments) and M spiking statistics. The compared
distributions must have the same number of spiking statistics M, but can have different sample sizes N.

We normalized (z-scored) the statistics, to ensure that all metrics M were equally weighted for the distance
measurement. We assigned equal weights w to each neuron within each distribution, such that $\sum^N w = 1$. Finally,
we searched for the optimal transport between the two distributions. Note that it is not necessarily a one-to-one
map since the weight from one node can be redistributed to various nodes. The Wasserstein distance (WS) is the
work normalized by the transported weight, which we defined to $\sum^N w = 1$, thus $\sum_i^{N_i} \sum_j^{N_j} w_{i,j} = 1$, and the
WS is therefore

$$\mathrm{WS} = \min \sum_i^{N_i} \sum_j^{N_j} w_{i,j} \cdot d_{i,j} \tag{2}$$

367 where N_i, N_j are the number of samples in the two compared spiking statistics clouds, $w_{i,j}$ is the weight
368 transported between points i and j, and $d_{i,j}$ is the Euclidean distance between them. To find the optimal
369 transport configuration we used a simplex algorithm implemented in OpenCV for this purpose[56].

370 We studied both the multivariate and univariate distributions using the WS distance.

Anatomical data for hierarchical characterization

The anatomical data for hierarchical characterization of the cortical areas was collated from various published
sources in the literature.

Neuron densities were estimated from Nissl and NeuN stained histological slices of macaque[12,57]. Since
the NeuN method tends to underestimate neuron densities[58], we linearly scaled NeuN estimates to match the
Nissl-stain estimates of neuron density[13,59].

Receptor densities were measured via *in vitro* receptor autoradiography for 14 different receptor types using
receptor-specific protocols[4]. The 14 receptor types included the following; Glutamate: AMPA, kainate and
NMDA; GABA: $\mathrm{GABA_A}$, $\mathrm{GABA_A}$/BZ and $\mathrm{GABA_B}$; acetylcholine: $\mathrm{M_1}$, $\mathrm{M_2}$ and $\mathrm{M_3}$; serotonin: $5-\mathrm{HT_{1A}}$ and
$5-\mathrm{HT_{2A}}$; noradrenaline: α_1 and α_2; and dopamine: $\mathrm{D_1}$. We used the average receptor density as a proxy for the
cortical hierarchy (Figure 3), and also report the correlations for each individual receptor (Figure S5).

Spine counts for several cortical areas in the macaque were previously described by Elston[15], later mapped to
a common parcellation and openly shared by Froudist-Walsh et al.[14].

The ratio of T1-weighted to T2-weighted (T1w/T2w) MRI signal is a proposed marker of myelination in the
cortical gray matter. T1w/T2w was measured by Donahue et al.[16], later mapped to a common parcellation and
openly shared by Froudist-Walsh et al.[4].

The hierarchical level of cortical areas is classically defined from the structural connectivity between cortical
areas[1]. In particular, the laminar connectivity patterns have been shown to follow a hierarchical organisation[2].
Here, we use the hierarchical levels estimated and shared by Froudist-Walsh et al.[14].

Many anatomical markers are known to have heavy-tailed distributions, such as the lognormal distribution[13,60].
Indeed, the neuron density, mean receptor density, spine count, and myelination have right-skewed distributions;
thus, we took their base-10 logarithm. The logarithm of a lognormal distribution is a normal distribution, which
is better suited for statistical testing and correlation measurements.

For this work, we calculated the anatomical dissimilarity by taking the Euclidean distance between the 4D
points reflecting the neuron density, mean receptor density, spine count, and hierarchical levels from connectivity
patterns. The individual anatomical markers were first normalized between 0 and 1. We excluded myelination
from the anatomical dissimilarity.

Estimation of hierarchies from dynamical and anatomical dissimilarities

397 To estimate the hierarchical position of each area from the dynamical or anatomical dissimilarities, we embed
398 them into a 1-dimensional line using Isomap ($k = 3$ neighbours). The resulting embedding is normalized between
399 0 and 1. For the embedding we use the full symmetric dissimilarity matrices, not just the lower triangular
400 matrices shown in Figure 3a,b.

Autocorrelation timescales

401 To estimate the autocorrelation timescale, we first estimated the firing rate of all neurons using a 10-ms bin
402 size. Then, we z-scored the firing rate, calculated the autocorrelation by convolving the signal with itself,

and normalized the autocorrelation by its maximum value. Finally, we fitted an exponential function to the autocorrelation using least-squares (scipy.optimize.curve_fit). The exponential function is of the form:

$$f(x) = ae^{-x/\tau} + b \tag{3}$$

where a and b are arbitrary scaling constants and τ is the autocorrelation timescale. Following Fontanier et al[40], if there was an autocorrelation peak within the first 100-ms of time lag, we fit only starting from that peak and not from zero lag. One fit was calculated for each neuron. We consider the neuron to lack a valid (measurable) τ when the exponential function fit does not converge or it has a coefficient of determination $R^2 \leq 0.8$. For the main results (Figure 5) we split the spike trains of each neuron into 300 s data segments. Shorter data segments yielded nearly no valid measurements of τ (Figure S4). Example autocorrelation functions and fits with valid τ for a few randomly selected neurons are shown in Figure S9.

Acknowledgements

We thank Jon Martinez Corral for proofreading the manuscript; Sarah Palmis for support with data collection from macaque T; Anno Kurth for helpful discussions about statistics and high-dimensional spaces; Nilanjana Nandi for helpful discussions about autocorrelation timescales; Ján Antolík and Karolína Korvasová for feedback on the manuscript about the interpretability of the results.

This work received funding from the Programme Johannes Amos Comenius (OP JAK) under the project 'MSCA Fellowships CZ - UK3' (reg. n. CZ.02.01.01/00/22_010/0008220); the DFG Priority Program (SPP 2041 "Computational Connectomics") [S.J.van Albada: AL 2041/1-1]; the EU's Horizon 2020 Framework Grant Agreement No. 785907 (HBP-SGA2) and No. 945539 (HBP-SGA3); the DFG (RTG 2416 "MultiSenses-MultiScales"); the CNRS Multidisciplinary Exploratory Projects initiative; and a grant co-financed by Greece and the European Union (European Regional Development Fund) through the Operational Programme "Competitiveness Entrepreneurship Innovation 2014–2020" (to G.G. Gregoriou project MIS 5070462) in the context of the FLAG-ERA project PrimCorNet.

Author contributions

Conceptualization AMG. Data collection XC (macaque L); SP, PS, GG (macaque F); BEK (macaque T). Spike sorting AMG (macaque L); AKJ (macaque Y); SP, PS, GG (macaque F); BEK (macaque T). Data curation and preprocessing AMG. Methodology AMG, RG, BEK, SvA. Software AMG, RG. Formal Analysis AMG. Visualization AMG. Writing - original draft AMG. Writing - review & editing all authors. Supervision SvA, BEK. Resources GG, TB, BEK, SvA. Funding acquisition AMG, GG, TB, BEK, SvA.

Competing interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

XC is a co-founder and shareholder of a neurotechnology start-up, Phosphoenix BV (https://phosphoenix.nl).

Data and code availability

Spike sorted activity recordings and the code required to reproduce this analysis will be made available upon publication in a peer-reviewed journal.

The anatomical data were openly available from the literature, see Anatomical data for hierarchical characterization for the references.

References

[1] D. J. Felleman and D. C. Van Essen. "Distributed Hierarchical Processing in the Primate Cerebral Cortex" (1991).

[2] N. T. Markov et al. "Anatomy of hierarchy: Feedforward and feedback pathways in macaque visual cortex". *Journal of Comparative Neurology* 522.1 (2014), pp. 225–259.

[3] J. A. Harris et al. "Hierarchical organization of cortical and thalamic connectivity". *Nature* 575.7781 (2019), pp. 195–202.

[4]   S. Froudist-Walsh et al. "Gradients of neurotransmitter receptor expression in the macaque cortex". *Nature Neuroscience* 26.7 (2023), pp. 1281–1294.

[5]   J. D. Murray et al. "A hierarchy of intrinsic timescales across primate cortex". *Nature Neuroscience* 17.12 (2014), pp. 1661–1663.

[6]   Y. Mochizuki et al. "Similarity in Neuronal Firing Regimes across Mammalian Species". *Journal of Neuroscience* 36.21 (2016), pp. 5736–5747.

[7]   J. H. Siegle et al. "Survey of spiking in the mouse visual system reveals functional hierarchy". *Nature* 592.7852 (2021), pp. 86–92.

[8]   X. Chen et al. "1024-channel electrophysiological recordings in macaque V1 and V4 during resting state". *Scientific Data* 9.1 (2022), p. 77.

[9]   T. Brochier et al. "Massively parallel recordings in macaque motor cortex during an instructed delayed reach-to-grasp task". *Scientific Data* 5.1 (2018), p. 180055.

[10]   M. J. de Haan et al. "Real-time visuomotor behavior and electrophysiology recording setup for use with humans and monkeys". *Journal of Neurophysiology* 120.2 (2018), pp. 539–552.

[11]   P. Sapountzis et al. "Dynamic and stable population coding of attentional instructions coexist in the prefrontal cortex". *Proceedings of the National Academy of Sciences* 119.40 (2022), e2202564119.

[12]   S. F. Beul and C. C. Hilgetag. "Neuron density fundamentally relates to architecture and connectivity of the primate cerebral cortex". *NeuroImage* 189 (2019), pp. 777–792.

[13]   A. Morales-Gregorio, A. Van Meegen, and S. J. Van Albada. "Ubiquitous lognormal distribution of neuron densities in mammalian cerebral cortex". *Cerebral Cortex* 33.16 (2023), pp. 9439–9449.

[14]   S. Froudist-Walsh et al. "A dopamine gradient controls access to distributed working memory in the large-scale monkey cortex". *Neuron* 109.21 (2021), 3500–3520.e13.

[15]   G. Elston. "Specialization of the Neocortical Pyramidal Cell during Primate Evolution". In: *Evolution of Nervous Systems.* Elsevier, 2007, pp. 191–242.

[16]   C. J. Donahue et al. "Using Diffusion Tractography to Predict Cortical Connection Strength and Distance: A Quantitative Comparison with Tracers in the Monkey". *The Journal of Neuroscience* 36.25 (2016), pp. 6758–6770.

[17]   R. Gao et al. "Neuronal timescales are functionally dynamic and shaped by cortical microarchitecture". *eLife* 9 (2020), e61277.

[18]   L. Rudelt et al. "Signatures of hierarchical temporal processing in the mouse visual system". *PLOS Computational Biology* 20.8 (2024), e1012355.

[19]   N. T. Markov et al. "A weighted and directed interareal connectivity matrix for macaque cerebral cortex". *Cerebral Cortex* 24.1 (2012), pp. 17–36.

[20]   A. Goulas et al. "The natural axis of transmitter receptor distribution in the human cerebral cortex". *Proceedings of the National Academy of Sciences* 118.3 (2021), e2020574118.

[21]   G. N. Elston. "Cortex, Cognition and the Cell: New Insights into the Pyramidal Neuron and Prefrontal Function". *Cerebral Cortex* 13.11 (2003), pp. 1124–1138.

[22]   S. Shinomoto, K. Shima, and J. Tanji. "Differences in Spiking Patterns Among Cortical Neurons". *Neural Computation* 15.12 (2003), pp. 2823–2842.

[23]   A. S. Bock et al. "Resting-State Retinotopic Organization in the Absence of Retinal Input and Visual Experience". *Journal of Neuroscience* 35.36 (2015), pp. 12366–12382.

[24]   N. Gravel et al. "Cortical connective field estimates from resting state fMRI activity". *Frontiers in Neuroscience* 8 (2014).

[25]   C. A. Runyan et al. "Distinct timescales of population coding across cortex". *Nature* 548.7665 (2017), pp. 92–96.

[26]   Z. R. Zeisler et al. "Consistent Hierarchies of Single-Neuron Timescales in Mice, Macaques, and Humans". *The Journal of Neuroscience* 45.19 (2025), e2155242025.

[27]   M. Song et al. "Hierarchical gradients of multiple timescales in the mammalian forebrain". *Proceedings of the National Academy of Sciences* 121.51 (2024), e2415695121.

[28]   R. Cusinato et al. "Intrinsic neural timescales in the temporal lobe support an auditory processing hierarchy". *The Journal of Neuroscience* (2023), JN–RM–1941–22.

[29] R. Cusinato et al. "Sleep modulates neural timescales and spatiotemporal integration in the human cortex". *The Journal of Neuroscience* 45.15 (2025), e1845242025.

[30] J. Oberste-Frielinghaus et al. *Detection and Removal of Hyper-synchronous Artifacts in Massively Parallel Spike Recordings*. 2024.

[31] M. Okun et al. "Diverse coupling of neurons to populations in sensory cortex". *Nature* 521.7553 (2015), pp. 511–515.

[32] L. Mazzucato, A. Fontanini, and G. La Camera. "Stimuli Reduce the Dimensionality of Cortical Activity". *Frontiers in Systems Neuroscience* 10 (2016).

[33] D. Dahmen et al. "Strong and localized recurrence controls dimensionality of neural activity across brain areas". *bioRxiv* (v2 2020).

[34] Y. Hu and H. Sompolinsky. "The spectrum of covariance matrices of randomly connected recurrent neuronal networks with linear dynamics". *PLOS Computational Biology* 18.7 (2022), e1010327.

[35] A. Morales-Gregorio et al. "Neural manifolds in V1 change with top-down signals from V4 targeting the foveal region". *Cell Reports* 43.7 (2024), p. 114371.

[36] F. H. C. Marriott and J. A. Pope. "Bias in the estimation of autocorrelations". *Biometrika* 41.3-4 (1954), pp. 390–402.

[37] B. E. Huitema and J. W. McKean. "Autocorrelation estimation and inference with small samples." *Psychological Bulletin* 110.2 (1991), pp. 291–304.

[38] R. Zeraati, T. A. Engel, and A. Levina. "A flexible Bayesian framework for unbiased estimation of timescales". *Nature Computational Science* 2.3 (2022), pp. 193–204.

[39] P. A. Dabrowska et al. "On the Complexity of Resting State Spiking Activity in Monkey Motor Cortex". *Cerebral Cortex Communications* 2.3 (2021), tgab033.

[40] V. Fontanier et al. "Inhibitory control of frontal metastability sets the temporal signature of cognition". *eLife* 11 (2022), e63795.

[41] M. Axer et al. "A novel approach to the human connectome: Ultra-high resolution mapping of fiber tracts in the brain". *NeuroImage* 54.2 (2011), pp. 1091–1101.

[42] S. B. Eickhoff, R. T. Constable, and B. T. Yeo. "Topographic organization of the cerebral cortex and brain cartography". *NeuroImage* 170 (2018), pp. 332–347.

[43] E. Genç et al. "Functional Connectivity Patterns of Visual Cortex Reflect its Anatomical Organization". *Cerebral Cortex* 26.9 (2015), pp. 3719–3731.

[44] E. M. Gordon et al. "A somato-cognitive action network alternates with effector regions in motor cortex". *Nature* 617.7960 (2023), pp. 351–359.

[45] N. Kriegeskorte. "Representational similarity analysis – connecting the branches of systems neuroscience". *Frontiers in Systems Neuroscience* (2008).

[46] J. A. Gallego et al. "Long-term stability of cortical population dynamics underlying consistent behavior". *Nature Neuroscience* 23.2 (2020), pp. 260–270.

[47] J. V. Haxby et al. "Hyperalignment: Modeling shared information encoded in idiosyncratic cortical topographies". *eLife* 9 (2020), e56601.

[48] J. Barbosa et al. "Quantifying Differences in Neural Population Activity With Shape Metrics". *bioRxiv* (2025).

[49] A. P. Buccino et al. "SpikeInterface, a unified framework for spike sorting". *eLife* 9 (2020), e61834.

[50] B.-B. Koo et al. "Age-related effects on cortical thickness patterns of the Rhesus monkey brain". *Neurobiology of Aging* 33.1 (2012), 200.e23–200.e31.

[51] S. Nougaret et al. "Low and high beta rhythms have different motor cortical sources and distinct roles in movement control and spatiotemporal attention". *PLOS Biology* 22.6 (2024), e3002670.

[52] G. R. Holt et al. "Comparison of discharge variability in vitro and in vivo in cat visual cortex neurons". *Journal of Neurophysiology* 75.5 (1996), pp. 1806–1814.

[53] M. Denker et al. *Elephant 0.11.1*. 2022.

[54] R. Gutzen et al. "Reproducible Neural Network Simulations: Statistical Methods for Model Validation on the Level of Network Activity Data". *Frontiers in Neuroinformatics* 12 (2018), p. 90.

[55] H. R. Lindman. *Analysis of Variance in Experimental Design*. New York: Springer, 1992.

[56] G. Bradski. "The OpenCV Library". *Dr. Dobb's Journal of Software Tools* (2000).

[57] C. C. Hilgetag et al. "The primate connectome in context: Principles of connections of the cortical visual system". *NeuroImage* 134 (2016), pp. 685–702.

[58] S. J. van Albada et al. *Bringing Anatomical Information into Neuronal Network Models.* Ed. by M. Giugliano, M. Negrello, and D. Linaro. Vol. 1359. Springer, 2021, pp. 201–234.

[59] M. Schmidt et al. "Multi-scale account of the network structure of macaque visual cortex". *Brain Structure and Function* (2018).

[60] G. Buzsáki and K. Mizuseki. "The log-dynamic brain: how skewed distributions affect network operations". *Nature Reviews Neuroscience* 15.4 (2014), pp. 264–278.

Supplementary figures

Figure S1: Multivariate spiking statistics for macaque L, areas V1 (top left) and V4 (top right), and macaque F, areas V4 (bottom left) and PFC (bottom right). Sample spike trains shown.

Figure S2: Multivariate spiking statistics for macaque Y, areas DP (top left) and 7A (top right), and macaque T, areas PMd (bottom left) and M1 (bottom right). Sample spike trains shown.

Figure S3: Multivariate spiking statistics for macaque N (left) and macaque E (right), area M1/PMd in both cases. Sample spike trains shown.

Figure S4: Estimation of autocorrelation timescales heavily depends on data length. **a)** Total number of samples when data is sliced at different lengths, each segment is only allowed to contain one behavioral epoch. **b)** Total amount of available data (when $t_{segment} = 5$ s), given in hours. **c)** Fraction of samples with a valid metric as a function of segment lengths $t_{segment}$. FR, CC_{avg} and CC_{std} are always well defined, even when no spikes are present. LvR cannot be calculated in segments with fewer than 3 spikes, which happens in less than 20% of samples. The autocorrelation timescale τ is only valid in a small fraction of the data samples (we consider τ invalid when the exponential function fit does not converge or it has $R^2 \leq 0.8$). Increasing $t_{segment}$ drastically reduces the number of samples and for many, it is still not possible to estimate τ. **d)** Selected single-neuron examples where timescales fail to converge in short segments, but produce robust values for very long data segments.

Figure S5: Correlation between the multivariate dynamical dissimilarity and the log density ratio for all 14 receptors individually. Lower barplot summarizes the Pearson r coefficients. All receptors strongly correlate with the multivariate dynamical dissimilarity except GABAa and M2.

Figure S6: Detailed scatter plots corresponding to the correlations reported in Figure 3d. All the variants of dynamical dissimilarity (excluding one, or including only one metric) and all the individual anatomical differences shown. For each pair of variables, Pearson's r and the associated p-value are shown, which correspond to the values from Figure 3d.

Figure S7: Correlation between the individual anatomical markers (x-axis) and the spiking statistics (y-axis). The error bars indicate the variance of the spiking statistics. A linear fit is performed via weighted least squares, where the weights are the inverse of the variance. R^2 of the fit and the results of the corresponding t-test are shown. The correlation is considered non-significant (n.s.) if $p > 0.05$, which is the case in the panels with gray shading and an 'n.s.' watermark. No single spiking statistic correlates with all anatomical markers. Note that here we compare each area individually, whereas in Figure 3a–d and Figure 5c–d the areas are compared in a pairwise manner.

Figure S8: Supplementary analysis to the differences between eyes-open (EO) and eyes-closed (EC). **a** MANOVA test between all areas when including exclusively EC or EO data. Lower triangular entries show the base-10 logarithm of the p-values and the upper triangular entries show the F-statistic. Significance levels ($\alpha = 0.05/k$) are corrected for multiple testing following the Bonferroni correction ($k = 630$ including all the tests from [Figure 2](#)). The significance level is color-coded, with red p-values (orange F-statistics) denoting non-significant results. The test concludes that nearly all area dynamics are different from each other both in EC and EO. **b** Comparison of mean and std (errorbar) between EO and EC for all spiking statistics. For cases where the distributions do not change between EO and EC the mean should lie along the diagonal; conversely significant changes diverge from the diagonal. **c** Measurement of EO/EC differences for each spiking statistic and area. We report $\theta = (\mu_{EO} - \mu_{EC})/\sqrt{(\sigma_{EO} + \sigma_{EC})/2}$, which measures the changes to the mean in terms of the standard deviation. Most cases have small or even non-noticeable changes. **d** Sample distributions of $\log(CC_{avg})$ for the four areas with the largest differences between EO and EC.

Figure S9: Example autocorrelation functions of single neuron spike trains and the corresponding exponential fits (R^2 and τ shown). The examples were randomly selected out of the neurons that had a good fit ($R^2 > 0.8$).