1

Genomic differentiation of *Mannheimia haemolytica* genotypes

Darien Deschner, Janet E. Hill*

Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon, Canada

*Corresponding author

E-mail: Janet.Hill@usask.ca

Abstract

Mannheimia haemolytica is an opportunistic bacterial pathogen associated with the economically costly bovine respiratory disease. Two genotypes have been described, of which genotype 2 is more strongly associated with disease. Several previous studies have investigated the genomic differences between the genotypes and/or the major serotypes (1, 2 and 6) of *M. haemolytica*, however we still lack a clear basis for the greater disease association of genotype 2 (serotypes 1 and 6) and demonstrations of phenotypic differences are scarce. This work builds upon previous investigations to identify genes that differentiate the two genotypes with a particular focus on genes that may play a role in virulence and fitness in the respiratory tract microbiome. We identified 422 genotype differentiating genes in a collection of 208 *M. haemolytica* genomes (62 genotype 1, 146 genotype 2). Genotype differentiating genes including genotype-associated variants of a TonB-dependent siderophore receptor homolog, transferrin binding protein B, leukotoxin A, and IgA1 proteases. We also identified a genotype 1 associated lytic transglycosylase, and a genotype 2 specific highly immunogenic outer membrane lipoprotein. These results expand our knowledge of what differentiates the genotypes of *M. haemolytica* and provides information that can be utilized to inform diagnostic strategies. The results also highlight the relative lack of information regarding *M. haemolytica* from sheep and goats and geographic regions other than Canada and the United States.

Introduction

Mannheimia haemolytica is a gram-negative opportunist bacterial pathogen associated with bovine respiratory disease (BRD) [1,2]. The beef industry is estimated to lose more than $3

billion per year globally due to BRD [3,4]. Two genotypes of *M. haemolytica* have been

described based on whole genome sequencing and single-nucleotide polymorphism (SNP)

analysis and multiple serotypes have also been defined [5,6]. The three most common serotypes

(1, 2 &, 6) have been found to correspond to genotypes with serotypes 1 & 6 corresponding to

genotype 2 (G2) and serotype 2 corresponding to genotype 1 (G1) [1,5,7,8]. Genotype 2 has been

more frequently associated with clinical cases of BRD, however the genetic and phenotypic basis

for this apparent association is unclear [1,9–14].

Differences in disease association between strains of the same species of bacteria is

relatively common. For example, commensal and pathogenic strains of *E. coli* differ in their

possession of virulence factors such as adhesins, toxins such as the Shiga toxin, and other

effectors such as the cytotoxic necrotizing factors [15]. Other species known to exhibit

differences in disease outcome based on strain diversity include *Staphylococcus aureus*,

Mycobacterium tuberculosis, *Campylobacter jejuni*, *Listeria monocytogenes*, and *Salmonella*

enterica [16]. In addition to virulence factors, adaptations that confer increased fitness, such as

greater ability to acquire iron and other nutrients, can contribute to disease association by

allowing a pathogenic strain to outcompete non-pathogenic strain(s) present in the same

environment [17].

Previous research into the virulence factors of *M. haemolytica* has focused on leukotoxin,

lipopolysaccharide, adhesins and various proteases [11,18–20]. The response of *M. haemolytica*

to iron limitation has also been studied, however most previous studies have only examined the

virulence factors and metabolism of serotype 1 isolates [21]. Serotype associated differences in

leukotoxin have been demonstrated, including differing degrees of cytoxic activity [1,22,23].

Serotype 1 has also been demonstrated to invade differentiated bovine bronchial epithelial cells

in vitro while serotype 2 was unable to, suggesting that serotype 1 (included in G2) likely

possesses virulence factors facilitating this invasion that serotype 2 (included in G1) lacks [24].

Expanding on these previous observations and examining them through a genotype rather than

serotype lens will help elucidate the clinical relevance of genotype in *M. haemolytica*. Also, as

diagnostic laboratories continue to move toward molecular methods and away from techniques

like serology for many pathogens, understanding the association of genotype with pathogenicity

is increasingly important.

Several previous studies have investigated the phenotypic and genetic differences

between the two genotypes of *M. haemolytica*. Phenotypic differences described to date are

distinct matrix-assisted laser desorption/ionization - time of flight mass spectrometry (MALDI-

TOF) profiles and colony morphology [25,26]. A pangenome analysis of 69 *M. haemolytica*

genomes identified 112 G1 specific, and 179 G2 specific genes including seven G2 specific

predicted outer membrane protein encoding genes that were hypothesized to play a role in the

greater disease association of genotype 2 [27]. Another study examined a collection of 11 *M.*

haemolytica genomes for potential differences in virulence factors between the serotypes 1, 2

and 6 [28]. The authors reported more prophages in serotypes 1 and 6 than in serotype 2, which

could influence virulence due to the presence of virulence factors in these prophages. Self-

targeting CRISPR spacer sequences were detected in serotypes 1 and 6, which may enhance

immune evasion or adhesion during infection by modulating sialic acid expression on the cells

surface. ARG-carrying ICEs were more prevalent in serotypes 1 and 6 than serotype 2, which

could enhance survival of antibiotic exposure [28]. In our own previous investigation of

macrolide resistance in *M. haemolytica,* we found that genotype was not associated with

differences in phenotypic resistance after accounting for the prevalence of ICEs, however we did

identify a resistance-associated single nucleotide polymorphism in the L22 ribosomal protein that was exclusive to G2 [29].

In addition to the demonstration of genotype differentiation by MALDI-TOF [25], a loop-mediated amplification (LAMP) assay has also been developed to differentiate the *M. haemolytica* genotypes [30]. The rationale for application of such genotype specific diagnostics beyond research studies and the potential value of their adoption in routine diagnostic investigations depends on a better understanding of the roles of *M. haemolytica* genotypes in bovine respiratory disease. In the current study, we take advantage of the greater number of *M. haemolytica* whole genome sequences now available to identify genetic differences between the two genotypes. In our examination of >200 *M. haemolytica* genomes, particular attention was given to genotype differentiating genes that are predicted to encode virulence factors and those implicated in processes that affect fitness in the respiratory tract such as iron acquisition and carbohydrate utilization.

Materials & Methods

Genome sequences

M. haemolytica whole genome sequences defined as complete by NCBI were downloaded from the NCBI Genome database on March 25, 2024, using the NCBI Datasets command line tools [31]. To avoid duplication all .gca files that had a matching .gcf file were removed. The remaining genome sequences were genotyped using an *in silico* PCR assay in Unipro UGENe v45.1 and primers from a genotype specific LAMP assay [30,32]. Primers from the LAMP assay included a pair targeting G1 specific adhesin pseudogene B1, G2 adhesin G

(CP017538.1 locus tag: BG586_06285; Forward: 5'-GCA CAT TAA AAT ATA GCA GCT TTG-

3'; Reverse: 5'-CTA AGC CAG AGT GAT CCG-3'), and leukotoxin D (*lktD*) (CP005972.1

locus tag: F382_07400; Forward: 5'-ACT GAA AA TTC ACA CTA TAG GTG-3'; Reverse: 5'-

GCT AAT ATG TTT AAT TCG ACC AGT T-3'). The *lktD* primers were used to confirm identity

of genomes as *M. haemolytica*. In initial testing G1 specific primers failed to produce any

predicted products in our G1 reference strain and so were discarded. New G1 specific primers

targeting the G1 specific adhesin pseudogene B1 were developed using Primer 3 v 4.1.0

(CP017495.1 locus tag BG548_01640; Forward: 5'-ACT AAT CTG AAG AGC GGC GT-3';

Reverse: 5'-TTG GTG GTT GCT GCT TGT AC-3') [33–35]. Successfully genotyped NCBI

genomes were combined with a set of 99 genome sequences previously generated by our lab

(BioProject: PRJNA1088094)(30) and reference genomes (G1 = CP017518; G2 = CP017519)

for both genotypes to generate the final collection of 209 genomes (62 G1, 147 G2) (S1 Table).

Pangenome construction, annotation, and classification

The pangenome of 209 genomes was constructed using PanTools v 4.3.2 with FASTA

files, gff3 annotation files, and genotype designation as inputs [36–38]. Genomes were assigned

a new unique identifier based on their input order during pangenome construction (S1 Table).

Pantools groups protein sequences from the input genomes into homology groups based

on their similarity. Homology groups also contain the nucleotide sequences encoding the

constituent protein sequence(s). Homology groups were generated using the group command

with the relaxation parameter set to 2. The relaxation parameter is a combination of four sub-

parameters (intersection rate, similarity threshold, mcl inflation, and contrast) where a lower

number represents a stricter set of thresholds. Given the large number of genomes and

anticipated sequence variation, we set the relaxation parameter to 2 to reduce the chances of

minor differences preventing similar proteins from being grouped together.

Genome metrics such as genome size, GC content, number of genes, etc. were

determined using the metrics command in PanTools [36]. One genome (Genome 130/38241 S-

19) was approximately twice as large as all others and appeared to represent a chimeric

assembly. This genome was excluded from all subsequent analyses reducing the genome

collection to a total of 208 (62 G1, 146 G2).

Homology groups were classified as core (present in $\geq$ 90%, 187/208 genomes), unique

(present in only one single genome), or accessory (present in 2-186 genomes). Homology groups

were additionally classified as either shared (present in $\geq$ 90% genomes of one genotype),

exclusive (solely present in genomes of one genotype), or specific (present in $\geq$ 90% genomes of

one genotype, and no genomes of the other genotype) [36]. To allow for instances in which a

homology group was strongly associated with one genotype yet still present in small numbers of

the other genotype, all homology groups not initially classified as core or unique were further

manually examined. Homology groups that were present in $\geq$ 95% (60/62 G1, 138/146 G2) of

one genotype while also being present in $\leq$ 5% (3/62 G1, 7/146 G2) of the other genotype were

classified as being genotype associated.

Genomes were functionally annotated using InterProScan v 5.64-96.0 with SignalP v 4.1

database integrated [39–41]. Biosynthetic gene cluster annotations were generated using

antiSMASH 7.1.0 with default parameters [42]. InterPro and antiSMASH annotations were

added to the pangenome, with InterProScan annotations including InterPro, Pfam, GO, and

SignalP annotations [36].

Multiple sequence alignments of constituent nucleotide sequences were generated for all

homology groups in the pangenome with MAFFT and consensus sequences from the alignments

were generated to represent the genotype specific and associated homology groups [36,43].

Consensus sequences were used as queries for BLASTx searches using the bacterial/archaeal

translation code against the NCBI refseq_select database [44–48]. All other parameters were kept

as default. Searches were performed March 20-25, 2025. The top hit for each search was

manually recorded to provide further evidence of the predicted function, facilitate human

interpretation, and identify functions that may not have been annotated by any of the other

approaches. The entire pangenome database is available in S1 Dataset [43].

The nucleotide sequences of several previously characterized genes upregulated in

response to iron limitation, and previously identified proteases were used as BLAST queries

against the pangenome using the blast command in PanTools to determine if these previously

characterized genes were present in our dataset [18,21,36]. Both alignment threshold and

minimum identity were set to 90. Query nucleotide sequences are available S2 Table.

Tet(H) gene abundance

Tet(H) genes were used as a proxy for the presence of integrative and conjugative

elements (ICE) due to the strong correlation between presence of tet(H) and presence of ICE

[12,49]. The reference tet(H) nucleotide sequence (Accession Number: Y15510.1 (nucleotides

2099-3301)) was downloaded from the comprehensive antibiotic resistance database (CARD) on

July 26, 2024, and used as nucleotide BLAST query against the pangenome database using the

inbuilt PanTools blast command. Minimum identity and alignment threshold were both set to

90%.

Leukotoxin operons

Reference nucleotide sequences for each of the component genes of the *M. haemolytica* leukotoxin operon (*lktCABD*) were downloaded from GenBank (AF314508.2) and used as queries against the pangenome BLAST database with percent identity and alignment threshold both set to 90%. Consensus nucleotide sequences of G1 and G2 versions of all leukotoxin operon genes were extracted and translated in Geneious 2024.0.5. Predicted protein sequences for the G1 and G2 versions of all leukotoxin operon genes were aligned against one another in Geneious 2024.0.5.

Annotation of carbohydrate active enzymes

Carbohydrate active enzymes (CAZymes) were annotated using a local installation of dbCAN v 4.1.1 software with integrated SignalP v 4.1 database and DIAMOND, HMMER, and HMMER dbcan_sub prediction algorithms [50]. Annotation was performed on each genome sequence individually using a custom BASH script [43]. dbCAN annotations that had supporting annotations from at least three tools, and contained a predicted signal peptide were selected for further analysis.

Protein structure prediction and visualization

Protein sequences were submitted to the AlphaFold Server for structural prediction [51]. Models with the highest-ranking score were downloaded and visualized in PyMol version 2.5.2 (Schrödinger, LLC).

Statistics

A Fisher's exact test and odds ratio comparison was performed to determine if the prevalence of tet(H) was different between genotypes. Both tests were performed in Microsoft Excel for Microsoft 365 using Real Statistics Resource Pack. A p-value ≤ 0.05 was considered significant.

A one-way ANOVA (Kruskal-Wallis), and post-hoc pairwise Mann-Whitney t-tests were performed in GraphPad PRISM v 10.4.1 to determine if there was a difference in genome size between G1 and G2 with or without ICE. A p-value ≤ 0.05 was considered significant.

Results & Discussion

Description of genome collection

M. haemolytica genomes designated as complete were retrieved from NCBI and after removal of duplicates, 121 genomes were available for further analysis. *In silico* PCR with modified primers from a previously published genotyping assay resulted in the expected amplicons for 108/121 genomes. No *in silico* genotyping PCR products were obtained for 13 genomes and so they were removed from further analyses leaving 108 genomes.

Three of the 13 isolates that could not be genotyped also failed to produce any predicted product for the *lktD in silico* PCR. Two of these genomes have since been removed by the RefSeq curation team for either containing too many frameshift mutations (GCF_900474405) or failing to pass a completeness check (GCF_023516295). All three genome sequences that failed to produce an *lktD* product were from sheep isolates (NCTC 10643, NCTC 10609) or from

culture collections from the 1960s (ATCC 29701). Of the other ten genomes that were not

genotyped by *in silico* PCR, four were from a German study examining macrolide resistance in

M. haemolytica [49], and three were from sheep isolates collected in 1951 or 1968 and

sequenced as part of the United States Department of Agriculture (USDA) *M. haemolytica*

sequencing project (BioProject: PRJNA172854). The last three isolates were all generated from

non-nasopharyngeal cattle samples as part of either the USDA sequencing project

(GCF_000819525, GCF_023516255) or a German study (GCF_020971705) and had a CheckM

completeness <92.5%.

Given the variety of the sources of the isolates we were unable to genotype it is difficult

to determine a general cause, however the assay developed may struggle with genomes derived

from ovine isolates, non-North American isolates, and non-nasopharyngeal isolates. Whether the

failure of the *in silico* assay to genotype isolates from these origins represents true genotypic

differences is unclear and requires further investigation. Difficulties in genotyping these isolates

likely reflects the lack of diversity in available *M. haemolytica* genomes. For example, of the 158

annotated complete *M. haemolytica* genome sequences available in the NCBI genome database

as of April 4, 2025, only 22 were from sources other than collections from the USDA or the

University of Saskatchewan.

The successfully genotyped genomes were combined with 99 genomes from our previous

work (BioProject Accession PRJNA1088094) and reference genomes for G1 and G2 as

described in the Methods. One additional genome was removed following discovery that it was

likely a chimera, leaving a final dataset 208 *M. haemolytica* whole genomes with 62 G1 and 146

G2 representatives (S1 Table). Most genomes were from isolates originating from feedlot cattle

in either Canada or the United States between 2003 and 2021. The lack of geographic diversity

and absence of genomes from ovine isolates means that the results presented in this work may

not represent the characteristics of *M. haemolytica* genomes outside of the Canadian and

American feedlot industries.

Genome properties & Tet(H) abundance

The two genotypes were compared to one another on a broad set of genome metrics, such

as genome size, GC content, etc., to determine if there were any high-level differences in the

genomic structure and content of the two genotypes. G2 genomes were larger, had more

predicted protein-coding sequences, and had slightly lower GC content than G1 genomes. No

differences were identified in the number of tRNA or rRNA genes or the proportion of the

genome covered by genes (Table 1).

Table 1. Comparison of genome features of G1 and G2. Values are the average ± standard
deviation. P-value ≤ 0.05 was considered significant.

Metric	Genotype 1 (n = 62)	Genotype 2 (n = 146)	t-test *P*-value
Genome Size (Mb)	2.592 ± 0.045	2.719 ± 0.048	<2e-16
GC Content (%)	41.14 ± 0.026	41.04 ± 0.031	<2e-16
Gene Count	2,561.7 ± 100.9	2,798.9 ± 116.4	<2e-16
Genome Covered by Genes (%)	88.17 ± 1.5	88.73 ± 1.2	0.0097
tRNA Count	64.31 ± 1.70	64.17 ± 2.04	0.61
rRNA Count	18.98 ± 4.3	19.68 ± 2.4	0.24

The larger size of G2 genomes may be attributable to a greater prevalence of integrative

and conjugative elements (ICEs), large (~20 kb to >500 kb) mobile genetic elements that are

known to be associated with antimicrobial resistance and are more frequently identified in G2

genomes [5,29]. To investigate this possibility, we used the proxy measure of the presence of

tetracycline resistance gene tet(H) due to its strong association with ICEs [12,49]. Tet(H) was

identified significantly more frequently in G2 (67/146, 45.9%) than G1 (1/62, 1.6%) (Fisher's *p*-

value = 4.39 x 10^{-12} Odds Ratio: 0.02 ± 1.02), however, G2 genomes were larger than G1

regardless of the presence of ICE (Figure 1). The larger size of G2 genomes regardless of

presence of ICE suggests that G2 genomes carry a wider repertoire of genes that may allow them

to be more adaptable to a wider variety of environments than the smaller G1 genomes.

Considering average bacterial gene sizes, the difference in the median genome size of G2

without ICE and G1 (~100 kb) could represent dozens of G2-specific genes. Additionally, the

greater prevalence of ICEs in G2 suggests that this more disease associated genotype is also

more likely to possess antimicrobial resistance genes.

Figure 1. Genome sizes of Genotype 1 (G1), Genotype 2 (G2), G2 possessing integrative and conjugative elements (ICEs), and G2 without ICEs (no ICE). Red bars indicate mean. Significant differences (P < 0.0001) are indicated (one way ANOVA (Kruskal-Wallis), post-hoc pairwise Mann-Whitney t-tests).

Pangenome analysis

The pangenome calculation resulted in the identification of "homology groups": clusters of sequences from input genomes that exceed a given similarity threshold. We then examined the distribution of homology groups across all *M. haemolytica* genomes to identify those that are core, unique, accessory, shared, genotype specific or genotype associated as defined in the methods. A total of 4,038 homology groups were identified of which 2,007 (49.7%) were core,

1,473 (36.48%) were accessory, and 558 (13.82%) were unique. Subsequent analyses focused on

accessory homology groups as these groups were most likely to differ in prevalence between the

two genotypes. We identified 280 G1 exclusive, 2,169 G1 shared, and 135 G1 specific homology

groups and 790 G2 exclusive, 2,342 G2 shared, and 233 G2 specific homology groups. To focus

our attention on the homology groups most likely to define and differentiate genotypes, the

shared or exclusive categories were further examined to identify those homology groups that

were present in ≥ 95% of one genotype and ≤ 5% of the other genotype, resulting in 21 G1

associated, and 33 G2 associated homology groups (S3-S4 Tables). Identification of genotype

specific and associated genes within this relatively large genome collection further reinforces

previous findings that the two genotypes carry distinct sets of genes [27]. A substantial

proportion of the proteins encoded by the specific and associated homology groups were mobile

genetic element (MGE) associated (transposases, toxin-antitoxin systems, prophage associated

proteins, etc.) or hypothetical or unknown function proteins (Figure 2).

The larger number of G2 specific or associated homology groups compared to G1

specific or associated groups is consistent with the larger size of G2 genomes (Figure 1) and the

larger number of specific and associated homology groups encoding MGE-associated proteins in

G2 suggests that these elements are more common in G2 than G1, consistent with previous

observations [9,12,27,28,49]. The number of hypothetical/unknown proteins is also unsurprising

as ~22□% of all domain families in Pfam v32.0 are defined as "domains of unknown function"

and a recent examination of >30,000 bacterial genomes and metagenomes showed that 40-60%

of genes are of unknown function [52,53]. Homology groups predicted to encode

unknown/hypothetical proteins, and MGE-associated proteins were excluded from further

analysis in this study but certainly represent an opportunity for future investigation.

Figure 2. Genotype specific/associated homology groups. Mobile Genetic Element (MGE) associated proteins included transposases, plasmid toxin-antitoxin systems, prophage associated proteins, etc. Hypothetical proteins included those annotated as hypothetical proteins as well as proteins annotated with a domain of unknown function (DUF) which did not appear to be MGE-associated and did not have any putative functions. Proteins that did not fall into either of the two preceding categories were placed into the Other category. G1 = Genotype 1 G2 = Genotype 2.

Based on inspection of the complete inventory of pangenome homology groups (S3-4 Tables), we identified a subset of specific/associated homology groups that may be associated with reported differences between disease association of G1 and G2 (Table 2).

Table 2. Genotype association and predicted functions of selected homology groups. Annotation notes summarize the functional annotations from InterProScan for each homology group. BLASTX search used consensus nucleotide sequence of the homology group as query against the refseq_select database. Reported top BLASTx hits were from *M. haemolytica* unless otherwise stated.

Homolog group	Genotype association	InterProScan Annotation	BLASTx hit (top)	Predicted Function
2042480	G2 specific	TonB-dependent receptor.	TonB-dependent siderophore receptor	Iron Acquisition
2042481	G1 associated	TonB dependent receptor protein.	TonB-dependent siderophore receptor	Iron Acqusition
2041941	G2 specific	TonB Dependent Receptor	TonB-dependent receptor plug-domain containing protein	Iron Acquisition
2944545	G2 specific	Transferrin Binding	Transferrin-binding protein-like solute	Iron Acquisition

		Protein (TpbB)	binding protein	
2040773	G1 specific	Transferrin Binding Protein	Transferrin-binding protein-like solute binding protein. *Mannheimia pernigra*	Iron acquisition
2042189	G2 specific	Peptidase S6/Autotransporter/IgA1 protease	S6 Family Peptidase	Host immune evasion
2040817	G1 associated	Peptidase/Pectate Lyase/IgA1 Protease	S6 family peptidase	Host immune evasion
2044609	G2 specific	N/A	Outer membrane lipoprotein PlpE.	Highly immunogenic outer membrane protein
2042729	G1 associated	MltB-like Transglycosylase	Lytic murein transglycosylase. *Mannheimia glucosida*	Cell wall remodeling/recycling

Homology groups 2042480 (G2 specific) and 2042481 (G1 associated) were both

annotated as encoding a TonB-dependent siderophore receptor. The consensus amino acid

sequences of these homology groups were 84.4% identical, and both contained consensus TonB-

dependent transporter (TBDT) motifs. In addition, the predicted 3D structures demonstrated the

canonical 22 β-strand barrel structure with an N-terminal plug domain characteristic of TBDTs

(Figure 3A). The amino acid sequence differences between the G1 and G2 sequences were

concentrated in the extracellular loops responsible for substrate binding, suggesting that these

two proteins may differ in their ability to bind their substrate [54,55] or they may bind different

substrates (Figure 3B). TonB dependent receptors (TBDTs) are transporters often involved in

iron acquisition, specifically in the uptake of small, secreted iron scavenging molecules known

as siderophores [54,55]. Homology groups 2042480 and 2042481 also produced BLAST hits to a

previously characterized siderophore (ferric enterobactin) receptor (*frpB*) homolog that was

upregulated in *M. haemolytica* in response to iron limitation [21]. Identification of a putative

siderophore receptor in *M. haemolytica* is unexpected as *M. haemolytica* is not known to produce

or utilize siderophores and has been demonstrated to be unable to utilize ferric enterobactin so it

is likely that the identified TBDTs have some other function besides siderophore uptake [11,21].

Figure 3. A) Predicted 3D structures of Genotype 1 (G1, Purple) & Genotype 2 (G2, Orange) ferric enterobactin receptor (frpB) homolog proteins. 3D structure conforms to canonical TonB-dependent receptor structures. B) 2D representation of tertiary structure of G1 and G2 frpB homologs. Each grey rectangle represents one of the 22 β-strand sheets, and the size of the extracellular/periplasmic loops are shown. Differences in the amino acid sequence for select extracellular loops are shown in further detail. Difference in extracellular loops may contribute to differences in substrate binding efficiency and/or result in interactions with completely different substrates.

G2 specific homology group 2041941 was also predicted to encode a TBDT, and no

similar G1 specific/associated protein was identified and the consensus sequence of 2041941 did

not correspond to previously identified genes upregulated in response to iron limitation [21]. The

identification of another G2 specific TBDT suggests that G2 may have a greater ability to uptake

iron and/or other substrates. TBDT have been implicated in transport of a variety of substrates in

Gram-negative bacteria including siderophores as well as vitamin B_{12} and carbohydrates [55].

Further examination is required to determine the function of this TBDT and whether it provides a

competitive advantage to G2.

Other iron acquisition proteins include transferrin binding proteins A and B (*tpbAB*), both

of which are found in *M. haemolytica* [11]. Transferrin is an iron-carrying glycoprotein found in

mammalian circulatory systems that transports iron throughout the body and serves as a sink for

any excess free iron released into the circulatory system [56]. TpbB is a surface lipoprotein that

extends outwards from the outer membrane to the extracellular space where it binds to transferrin

and brings the transferrin closer to TbpA, a TBDT in the outer membrane that transports the

transferrin into the periplasm [56]. To investigate any genotype associations of transferrin

binding proteins, previously characterized *tbpA* and *tbpB* nucleotide sequences from *M.*

haemolytica were used as BLAST queries against the pangenome database. Core homology

group 2044546 contained hits for *tbpA* while *tbpB* hits were solely identified in G2 specific

homology group 2044545 with no hits in any G1 genomes. These results suggest that *tbpA* is

highly conserved within *M. haemolytica* and likely plays a key role in iron metabolism. In

contrast while no BLAST hits were produced to the query *tbpB* nucleotide sequence in G1

genomes, G1 specific homology group 2040773 was predicted to encode for a transferrin binding

protein. The amino acid sequences of G1 and G2 variants of the putative tbpB (homology groups

2040773 and 2044545) were compared and found to be 56.5% identical and 81.2% similar.

Despite the relatively low similarity of these two proteins, the predicted 3D structures were very

similar to one another as well as the crystal structure for the TbpB from *Actinobacillus*

pneumoniae (S1 Figure). A difference in the TbpB structure between the genotypes was observed

in the N lobe with G2 appearing to have a longer alpha-helix structure in the N-lobe, which is the

region of the protein involved in transferrin binding [56]. Genotype variation of a transferrin

binding protein further suggests they may differ in their ability to acquire iron, which may

partially explain observed differences in disease association between the two genotypes. The

genotype variants of TbpB identified here may also reflect host preferences of the genotypes as

TbpB is selective for host transferrin and differences between TbpB proteins has been partially

attributed to this host selectivity [56].

IgA proteases are metalloproteases that cleave immunoglobulin A (IgA), a critical

component of the mucosal immune system. Two IgA proteases (IgA1 and IgA2) have been

previously identified and characterized in a G2 (serotype 1) *M. haemolytica* isolate [18]. We

identified a G2 specific IgA protease (homology group 2042189) and a G1 associated IgA

protease (2040817) (Table 2). To determine the relationship of these sequences to the previously

reported IgA proteases, a BLAST search was performed using previously reported nucleotide

sequences for IgA1 and IgA2 as queries against the pangenome database (IgA1:NZ_DS264709

Locus Tag: MHA_RS00165IgA2: NZ_DS264612.1 Locus Tag: MHA_RS09765). At least one

BLAST hit was returned for IgA2 in all but five of the included genomes with hits grouped into

five homology groups (2041035, 2041036, 2041399, 2041400, and 2043723). Homology groups

2041305 and 2041306 were accessory, while 2041399 and 2041400 were unique suggesting

some minor variations are likely present within the pangenome for this IgA2 protease. Homology

group 2043723, however, was a core group suggesting that this IgA2 protease is a core gene of

M. haemolytica and is not genotype differentiating. In contrast, a BLAST hit was returned for

IgA1 in 146/209 of the genomes with hits grouped into one of three homology groups (2040817,

2041298, and 2042189). Homology group 2041298 was found in only one genome so was not

examined any further. Group 2040817 is a G1 associated group while 2042189 is a G2 specific

group and thus these groups appear to represent genotype variants of an IgA protease.

Comparison of the consensus sequences of the two IgA1-like groups revealed that the

nucleotide sequence for 2040817 is 2,281 bp while the sequence for 2042189 is 4,512 bp (94%

identical across 2,281 shared positions). The G1 associated consensus sequence is much smaller

than the previously reported 1,503 amino acids (4,509 bp) for IgA1 and 1,398 amino acids

(4,194 bp) for IgA2 while the G2 specific consensus sequence is slightly larger than both

previously reported IgA protease sequences. The smaller size of the G1 sequence suggests that

this sequence may represent a truncated protease or difficulties in the gene finding process due to

the large size of these genes [18]. The slightly larger size of the G2 specific sequence may

similarly be due to gene finding issues or in generating the consensus sequence. While the

identified G1 specific sequence may not represent a complete protease sequence, the observation

of a G2 specific sequence and possibility of genotype variation remains of interest and warrants

further investigation. Should both genotypes have differing IgA protease sequences the potential

difference(s) in their ability to evade the host immune system may contribute to differences in

disease association.

Pasteurella lipoproteins (Plp) are a group of several immunogenic outer membrane

lipoproteins identified in *M. haemolytica* [57]. G2 specific homology group 2044609 was

predicted to encode PlpE, which has been investigated as a potential vaccine target for *M.*

haemolytica because of its immunogenicity [58–60]. Aside from the immunogenic role of PlpE

relatively little is known regarding its function although it is important in complement-mediated

killing of *M. haemolytica* cells in bovine hosts. Previous investigations of PlpE sequence

diversity across serotypes showed that the serotype 1 and 6 (G2) sequences were highly

conserved whereas serotype 2 (G1) PlpE sequences were extremely heterogenous, however,

antibodies raised against serotype 1 PlpE cross reacted with serotype 2 bacteria [61]. No G1

associated or specific *plpE* gene was identified in this study which may be due to the previously

observed heterogeneity of this sequence in G1 genomes or potentially due to the relative lack of

available G1 genomes compared to G2. The identification of G2 specific PlpE suggests that G2

isolates may induce stronger immune reactions than G1 isolates, however further research is

needed to investigate the prevalence of PlpE in G1and the immunogenicity of G1 PlpE compared

to G2-derived PlpE.

Homology group 2040817 (G1 specific) was predicted to encode a lytic transglycosylase

and is explored in more depth below in the discussion of carbohydrate active enzymes

(CAZymes).

Leukotoxin operon

Leukotoxin (Lkt) is a critical virulence factor in *M. haemolytica* and is produced by both

genotypes (all serotypes) [62]. Lkt is encoded in an operon consisting of four genes *(lktCABD)*.

lktA encodes the protoxin, *lktC* encodes the acylating enzyme essential for post-translational

modification and activation of the protoxin, and the products of *lktB* and *lktD* are involved in

secretion of the toxin [23,63]. *lktCABD* genes were identified and the predicted protein

sequences for the Lkt operon genes from G1 and G2 were aligned (Figure 4A). Within the Lkt

operon the predicted proteins of the *lktC, lktB* and *lktD* genes were 95.2%, 99.6% and 100%

identical, respectively, suggesting these genes are under conservation pressure due to their roles

in regulating the secretion and function of Lkt [63,64]. In contrast, LktA protein sequences were

only 89% identical. Differences in the predicted LktA proteins of G1 and G2 were concentrated

in the N-terminal and C-terminal regions of the protein (Figure 4B). In leukotoxins the N-

terminus is involved in pore-forming activities while the C-terminus is involved in export and

host immune system recognition [63]. As such any differences in the N-terminal and/or C-

terminal regions may potentially influence the activities of the Lkt and thus may result in

virulence differences between the two genotypes.

430
431 **Figure 4.** A) Percent identity between the predicted protein sequence of each of the four genes in
432 the leukotoxin operon in G1 and G2. B) Sequence alignment of G1 and G2 LktA amino acid
433 sequence with a sliding window size of 10. A larger peak indicates a greater number of matching
434 amino acids within the window for that position.
435

In a previous study, Davies *et al*. examined the sequence diversity of leukotoxin in 31 *M.*

haemolytica, 6 *Mannheimia glucosida,* and 4 *Pasteurella* [*Bibersteinia*] *trehalosi* isolates

identified eight *M. haemolytica* allelic variants of *lktA* [23]. Five of the alleles (A6-A10) were

identified solely in ovine isolates, allele A3 was only identified in bovine isolates, while A1 and

A2 alleles were identified in both ovine and bovine isolates. The A1 allele was subdivided into 5

groups (A1.1-A1.5) with A1.1 solely occurring in bovine isolates from serotype 1 (n = 3) and 6

(n = 1) isolates (corresponding to G2). In bovine isolates from serotype 2 (corresponding to G1),

the authors reported detection of alleles A2, A2.1, A2.2, and A3. LktA sequences of study

genomes were compared with representative amino acid sequences for each allele and subgroup

downloaded from NCBI Genbank. Within G2, 141/146 LktA sequences were 100% identical to

the A1.1 allele and another 4/146 were >99.5% identical to A1.1, consistent with the

observations of Davies *et al*. (S5-7 Tables). The only exception was the LktA from Genome 2,

that shared 100% amino acid identity with A1.3 and A1.5 (S7 Table). Interestingly, all but one of

the G1 LktA sequences were 99.5-100% identical to the A8.1 allele (Genome 157 was identical

to A2) (S5 and S8-9 Tables). The A8.1 allele was found in serotype 2 (G1) isolates by Davies *et*

al., however, all the isolates in that study were ovine (n = 3), and A8.1 was also detected in ovine

serotype 7 isolates (n = 3). The identification of A8.1 across many bovine origin genomes and

nearly complete absence of A2 alleles suggests that G1 from Canada and the U.S.A has diverged

from the G1 found in the United Kingdom where most of the isolates used by Davies *et al*. were

collected and further demonstrates that A8.1 is not restricted to ovine isolates [23]. Whether there

is a relationship between specific LktA alleles and respiratory disease severity in cattle warrants

further investigation.

CAZymes

Carbohydrate active enzymes (CAZymes) are enzymes whose function has been

biochemically verified to be involved in the metabolism of carbohydrates. We chose to

investigate any differences in CAZymes between the two genotypes since these may indicate

differences in ability to metabolize certain carbohydrates, and thus in the nutritional niches of the

genotypes [65]. We were particularly interested in enzymes predicted to be secreted into either

the periplasmic or extracellular space (i.e. contain a signal peptide) and are thus most likely to be

involved in interactions with the external environment. A total of six CAZymes containing signal

peptides were identified in the pangenome (S10 Table) (50). At least one CAZyme was predicted

in all genomes except for Genome 86/GCF_022404935, which was removed from this analysis.

CAZymes identified were members of families GH23 (lytic transglycosylase), GH33 (sialidase),

GH13_19 (amylose glucosidase), GT9 (glycosyltransferase), GH102 (lytic transglycosylase) and

GH103 (lytic transglycosylase). Only GH103 (homology group 2042729) was genotype

associated, present in 60/61 (98.4%) G1 genomes and 2/146 (1.37%) G2 genomes. The GH103

family consists of a group of lytic transglycosylases that cleave the β-1,4 linkage between *N*-

acetylmuramoyl and *N*-acetylglucosaminyl residues in peptidoglycan and belong to family 3 of

the transglycosylases with structures similar to lysozyme [66,67]. Numerous biological functions

have been attributed lytic transglycosylases, however the only clear role is that they are involved

in cell wall remodelling [66–69]. Whether this genotype difference in lytic transglycosylases is

associated with phenotypic differences in colony morphology of the two genotypes [26] or other

differences in cell wall maintenance will require further investigation.

Conclusions

We identified a total of 422 *Mannheimia haemolytica* genotype differentiating genes (135

G1 specific, 21 G1 associated, 233 G2 specific, and 33 G2 associated), including 125 with

unknown functions and 132 that are associated with mobile genetic elements. Genotype

associated genes and variants that may contribute to differences in disease association include

TonB-dependent transporters, IgA proteases, a highly immunogenic outer membrane protein

(PlpE), a lytic transglycosylase, and genotype variants of the leukotoxin gene. The list of

genotype differentiating genes provides a shortlist of targets that warrant further investigation to

provide a mechanistic understanding of phenotypic differences between the *M. haemolytica*

genotypes and inform rational strategies for *M. haemolytica* and bovine respiratory disease

diagnostics.

Acknowledgements

491 The authors are grateful to Dr. David Palmer (Department of Chemistry, University of

492 Saskatchewan) for assistance with the protein structure alignments, and to Dr. Maarten

493 Voordouw (Department of Veterinary Microbiology, University of Saskatchewan) for assistance

494 with statistical analysis and approaches.

Supporting Information

S1 Table. Summary of Genomes Included in Pangenome Analysis

S2 Table. Accession numbers of sequences used as BLAST queries against pangenome database.

S3 Table. Overview of pangenome classification of all homology groups. Genes matching is the

number of genes that were found to correspond to the given homology group, as this referred to

genes the number may be greater than or less than the number of genomes examined. For

pangenome classification: core = group was present in ≥90% of all genomes (187/209); unique =

Group was present in exactly one genome; Accessory = Group was present in somewhere

between 2-186 genomes. For phenotype (Genotype) classification: Shared = Group was present

in ≥90% of the genomes of the given genotype; exclusive = Group was present solely in

genomes of the given genotype; Specific = Group was exclusive to one genotype and present in

≥ 90% of that genotype; Associated = Group was present in ≥90% of the genomes of one

genotype and ≤5% of the other genotype.

S4 Table. Genotype Specific and Associated Homology Groups. Annotation notes column

provides summary of the interpro, pfam and GO functional annotations. Genes matching is the

number of genes that were found to correspond to the given homology group, as this referred to

genes the number may be greater than or less than the number of genomes examined. For

pangenome classification: core = group was present in ≥90% of all genomes (187/209); unique =

Group was present in exactly one genome; Accessory = Group was present in somewhere

between 2-186 genomes. For phenotype (Genotype) classification: Shared = Group was present

in ≥90% of the genomes of the given genotype; exclusive = Group was present solely in

genomes of the given genotype; Specific = Group was exclusive to one genotype and present in

≥ 90% of that genotype; Associated = Group was present in ≥90% of the genomes of one

genotype and ≤5% of the other genotype.

S5 Table. Pairwise comparisons of amino acid sequences of genotype 1 (G1) and genotype 2

(G2) Leukotoxin A consensus sequences against representatives of LktA alleles from Davies *et*

al. 2001

S6 Table. Pairwise comparison of all genotype 1 LktA nucleotide sequences.

S7 Table. Pairwise comparison of genotype 1 LktA amino acid sequences that had a divergent

LktA nucleotide sequence to representative sequences of LktA alleles identified in Davies *et al*.

2001.

S8 Table. Pairwise comparison of all genotype 2 LktA nucleotide sequences.

S9 Table. Pairwise comparison of genotype 1 LktA amino acid sequences that had a divergent

LktA nucleotide sequence to representative sequences of LktA alleles identified in Davies *et al*.

2001.

S10 Table. Summary of CAZyme Analysis Results. Only CAZymes that were predicted to

contain a signal peptide and were annotated by 3 of the utilized annotation tools (dbCAN_sub,

DIAMOND, & HMMER) were included. No CAZymes were predicted in genome

86/GCF_022404935 and so was excluded from this analysis.

S1 Figure. Predicted structures of G1 and G2 variants of a predicted transferrin binding protein

B (TbpB). (A) The crystal structure of TbpB from *Actinobacillus pleuropneumoniae* (PDB

accession 3HOL) was used as a reference. Major domains include the N-lobe and C-lobe, each

containing a beta-barrel and a "handle". The reference structure is "rainbow" coloured from N

terminus (blue) to C terminus (red). (B) Alignment of G1 and G2 variants. (C) G2 aligned to

reference. (D) G1 aligned to reference. The G1 variant has an elongated alpha-helix in the N-lobe

handle domain relative to G2 and the reference. The N-lobe has been demonstrated to interact

with transferrin. Structures and alignments were visualized in PyMol.

References

1. Rice JA, Carrasco-Medina L, Hodgins DC, Shewen PE. *Mannheimia haemolytica* and

bovine respiratory disease. Animal Health Research Reviews. 2007;8: 117–128.

doi:10.1017/S1466252307001375

2. Highlander SK. Molecular genetic analysis of virulence in *Mannheimia (Pasteurella)*

haemolytica. FBL. 2001;6: 1128–1150. doi:10.2741/highland

3. DeDonder KD, Apley MD. A literature review of antimicrobial resistance in pathogens

associated with bovine respiratory disease. Animal Health Research Reviews. 2015;16:

125–134. doi:10.1017/S146625231500016X

4. Blakebrough-Hall C, McMeniman JP, González LA. An evaluation of the economic effects

of bovine respiratory disease on animal performance, carcass traits, and economic outcomes

in feedlot cattle defined using four BRD diagnosis methods. Journal of Animal Science.

2020;98: skaa005. doi:10.1093/jas/skaa005

5. Clawson ML, Murray RW, Sweeney MT, Apley MD, DeDonder KD, Capik SF, et al.

Genomic signatures of *Mannheimia haemolytica* that associate with the lungs of cattle with

respiratory disease, an integrative conjugative element, and antibiotic resistance genes.

BMC Genomics. 2016;17: 982. doi:10.1186/s12864-016-3316-8

6. Biberstein EL, Gills M, Knight H. Serological types of Pasteurella hemolytica. Cornell Vet.

1960;50: 283–300.

7.   Al-Ghamdi GM, Ames TR, Baker JC, Walker R, Chase CCL, Frank GH, et al. Serotyping

of *Mannheimia (Pasteurella) haemolytica* Isolates from the Upper Midwest United States. J

VET Diagn Invest. 2000;12: 576–578. doi:10.1177/104063870001200617

8.   Mason C, Errington J, Foster G, Thacker J, Grace O, Baxter-Smith K. *Mannheimia*

haemolytica serovars associated with respiratory disease in cattle in Great Britain. BMC

Veterinary Research. 2022;18: 5. doi:10.1186/s12917-021-03121-3

9.   Klima CL, Alexander TW, Hendrick S, McAllister TA. Characterization of *Mannheimia*

haemolytica isolated from feedlot cattle that were healthy or treated for bovine respiratory

disease. Can J Vet Res. 2014;78: 38–45.

10.   Arnal JL, Fernández A, Vela AI, Sanz C, Fernández-Garyzábal JF, Cid D. Capsular type

diversity of *Mannheimia haemolytica* determined by multiplex real-time PCR and indirect

hemagglutination in clinical isolates from cattle, sheep, and goats in Spain. Veterinary

Microbiology. 2021;258: 109121. doi:10.1016/j.vetmic.2021.109121

11.   Confer AW, Martino P. *Mannheimia haemolytica*. CABI Compendium. 2017;CABI

Compendium: 70912. doi:10.1079/cabicompendium.70912

12.   Klima CL, Zaheer R, Cook SR, Booker CW, Hendrick S, Alexander TW, et al. Pathogens of

Bovine Respiratory Disease in North American Feedlots Conferring Multidrug Resistance

via Integrative Conjugative Elements. Journal of Clinical Microbiology. 2014;52: 438–448.

doi:10.1128/jcm.02485-13

13.   Holman DB, McAllister TA, Topp E, Wright A-DG, Alexander TW. The nasopharyngeal

microbiota of feedlot cattle that develop bovine respiratory disease. Vet Microbiol.

2015;180: 90–95. doi:10.1016/j.vetmic.2015.07.031

14.   Noyes NR, Benedict KM, Gow SP, Booker CW, Hannon SJ, McAllister TA, et al.

Mannheimia haemolytica in feedlot cattle: prevalence of recovery and associations with

antimicrobial use, resistance, and health outcomes. J Vet Intern Med. 2015;29: 705–713.

doi:10.1111/jvim.12547

15.   Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. Nat Rev Microbiol. 2004;2:

123–140. doi:10.1038/nrmicro818

16.   Nysten J, Sofras D, Dijck PV. One species, many faces: The underappreciated importance

of strain diversity. PLOS Pathogens. 2024;20: e1011931. doi:10.1371/journal.ppat.1011931

17.   Hacker J, Carniel E. Ecological fitness, genomic islands and bacterial pathogenicity. A

Darwinian view of the evolution of microbes. EMBO Rep. 2001;2: 376–381.

doi:10.1093/embo-reports/kve097

18.   Ayalew S, Murdock BK, Snider TA, Confer AW. *Mannheimia haemolytica* IgA-specific

proteases. Vet Microbiol. 2019;239: 108487. doi:10.1016/j.vetmic.2019.108487

19.   Singh K, Ritchey JW, Confer AW. *Mannheimia haemolytica*: Bacterial–Host Interactions in

Bovine Pneumonia. Vet Pathol. 2011;48: 338–348. doi:10.1177/0300985810377182

20. Kisiela DI, Czuprynski CJ. Identification of *Mannheimia haemolytica* Adhesins Involved in

Binding to Bovine Bronchial Epithelial Cells. Infection and Immunity. 2009;77: 446–455.

doi:10.1128/iai.00312-08

21. Roehrig SC, Tran HQ, Spehr V, Gunkel N, Selzer PM, Ullrich HJ. The response of

Mannheimia haemolytica to iron limitation: Implications for the acquisition of iron in the

bovine lung. Veterinary Microbiology. 2007;121: 316–329.

doi:10.1016/j.vetmic.2006.12.013

22. Davies RL, Baillie S. Cytotoxic activity of *Mannheimia haemolytica* strains in relation to

diversity of the leukotoxin structural gene lktA. Vet Microbiol. 2003;92: 263–279.

doi:10.1016/s0378-1135(02)00408-x

23. Davies RL, Whittam TS, Selander RK. Sequence Diversity and Molecular Evolution of the

Leukotoxin (lktA) Gene in Bovine and Ovine Strains of *Mannheimia (Pasteurella)*

haemolytica. J Bacteriol. 2001;183: 1394–1404. doi:10.1128/jb.183.4.1394-1404.2001

24. Cozens D, Sutherland E, Lauder M, Taylor G, Berry CC, Davies RL. Pathogenic

Mannheimia haemolytica Invades Differentiated Bovine Airway Epithelial Cells. Infect

Immun. 2019;87: 10.1128/iai.00078-19. doi:10.1128/iai.00078-19

25. Loy JD, Clawson ML. Rapid typing of *Mannheimia haemolytica* major genotypes 1 and 2

using MALDI-TOF mass spectrometry. J Microbiol Methods. 2017;136: 30–33.

doi:10.1016/j.mimet.2017.03.002

26. Wynn EL, Schuller G, Loy JD, Workman AM, McDaneld TG, Clawson ML. Differentiation

of *Mannheimia haemolytica* genotype 1 and 2 strains by visible phenotypic characteristics

on solid media. Journal of Microbiological Methods. 2020;171: 105877.

doi:10.1016/j.mimet.2020.105877

27. Clawson ML, Schuller G, Dickey AM, Bono JL, Murray RW, Sweeney MT, et al.

Differences between predicted outer membrane proteins of genotype 1 and 2 *Mannheimia*

haemolytica. BMC Microbiology. 2020;20: 250. doi:10.1186/s12866-020-01932-2

28. Klima CL, Cook SR, Zaheer R, Laing C, Gannon VP, Xu Y, et al. Comparative Genomic

Analysis of *Mannheimia haemolytica* from Bovine Sources. PLOS ONE. 2016;11:

e0149520. doi:10.1371/journal.pone.0149520

29. Deschner D, Voordouw MJ, Fernando C, Campbell J, Waldner CL, Hill JE. Identification of

genetic markers of resistance to macrolide class antibiotics in *Mannheimia haemolytica*

isolates from a Saskatchewan feedlot. Appl Environ Microbiol. 2024;90: e00502-24.

doi:10.1128/aem.00502-24

30. Dassanayake RP, Clawson ML, Tatum FM, Briggs RE, Kaplan BS, Casas E. Differential

identification of *Mannheimia haemolytica* genotypes 1 and 2 using colorimetric loop-

mediated isothermal amplification. BMC Research Notes. 2023;16: 4. doi:10.1186/s13104-

023-06272-8

31. O'Leary NA, Cox E, Holmes JB, Anderson WR, Falk R, Hem V, et al. Exploring and

retrieving sequence and metadata for species across the tree of life with NCBI Datasets. Sci

Data. 2024;11: 732. doi:10.1038/s41597-024-03571-y

32.   Okonechnikov K, Golosova O, Fursov M, UGENE team. Unipro UGENE: a unified

bioinformatics toolkit. Bioinformatics. 2012;28: 1166–1167.

doi:10.1093/bioinformatics/bts091

33.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3--

new capabilities and interfaces. Nucleic Acids Res. 2012;40: e115. doi:10.1093/nar/gks596

34.   Kõressaar T, Lepamets M, Kaplinski L, Raime K, Andreson R, Remm M. Primer3_masker:

integrating masking of template sequence with primer design software. Bioinformatics.

2018;34: 1937–1938. doi:10.1093/bioinformatics/bty036

35.   Koressaar T, Remm M. Enhancements and modifications of primer design program

Primer3. Bioinformatics. 2007;23: 1289–1291. doi:10.1093/bioinformatics/btm091

36.   Jonkheer EM, van Workum D-JM, Sheikhizadeh Anari S, Brankovics B, de Haan JR, Berke

L, et al. PanTools v3: functional annotation, classification and phylogenomics.

Bioinformatics. 2022;38: 4403–4405. doi:10.1093/bioinformatics/btac506

37.   Sheikhizadeh S, Schranz ME, Akdel M, de Ridder D, Smit S. PanTools: representation,

storage and exploration of pan-genomic data. Bioinformatics. 2016;32: i487–i493.

doi:10.1093/bioinformatics/btw455

38.   Anari SS, Ridder D de, Schranz ME, Smit S. Pangenomic read mapping. bioRxiv; 2019. p.

813634. doi:10.1101/813634

39.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-

scale protein function classification. Bioinformatics. 2014;30: 1236–1240.

doi:10.1093/bioinformatics/btu031

40.   Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The

InterPro protein families and domains database: 20 years on. Nucleic Acids Res. 2021;49:

D344–D354. doi:10.1093/nar/gkaa977

41.   Nielsen H. Predicting Secretory Proteins with SignalP. Methods Mol Biol. 2017;1611: 59–

73. doi:10.1007/978-1-4939-7015-5_6

42.   Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, et al. antiSMASH 7.0:

new and improved predictions for detection, regulation, chemical structures and

visualisation. Nucleic Acids Res. 2023;51: W46–W50. doi:10.1093/nar/gkad344

43.   Deschner D, Hill JE. Genomic Differentiation of *Mannheimia haemolytica* genotypes.

FigShare; 2025. doi:10.6084/m9.figshare.28872536

44.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics. 2009;10: 421. doi:10.1186/1471-2105-

10-421

45.   Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST

and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res.

1997;25: 3389–402.

46. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J

Mol Biol. 1990;215: 403–10.

47. Zhang Z, Schwartz S, Wagner L, Miller W. A greedy algorithm for aligning DNA

sequences. J Comput Biol. 2000;7: 203–214. doi:10.1089/10665270050081478

48. Morgulis A, Coulouris G, Raytselis Y, Madden TL, Agarwala R, Schäffer AA. Database

indexing for production MegaBLAST searches. Bioinformatics. 2008;24: 1757–1764.

doi:10.1093/bioinformatics/btn322

49. Kostova V, Hanke D, Kaspar H, Fiedler S, Schwarz S, Krüger-Haker H. Macrolide

resistance in *Mannheimia haemolytica* isolates associated with bovine respiratory disease

from the German national resistance monitoring program GERM-Vet 2009 to 2020. Front

Microbiol. 2024;15. doi:10.3389/fmicb.2024.1356208

50. Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y. dbCAN3: automated carbohydrate-active

enzyme and substrate annotation. Nucleic Acids Research. 2023;51: W115–W121.

doi:10.1093/nar/gkad328

51. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure

prediction of biomolecular interactions with AlphaFold 3. Nature. 2024;630: 493–500.

doi:10.1038/s41586-024-07487-w

52. Vanni C, Schechter MS, Acinas SG, Barberán A, Buttigieg PL, Casamayor EO, et al.

Unifying the known and unknown microbial coding sequence space. Brown CT, Storz G,

Brown CT, Smith B, editors. eLife. 2022;11: e67667. doi:10.7554/eLife.67667

53. Lobb B, Tremblay BJ-M, Moreno-Hagelsieb G, Doxey AC. PathFams: statistical detection

of pathogen-associated protein domains. BMC Genomics. 2021;22: 663.

doi:10.1186/s12864-021-07982-8

54. Ferguson AD, Deisenhofer J. TonB-dependent receptors-structural perspectives. Biochim

Biophys Acta. 2002;1565: 318–332. doi:10.1016/s0005-2736(02)00578-3

55. Silale A, Van Den Berg B. TonB-Dependent Transport Across the Bacterial Outer

Membrane. Annu Rev Microbiol. 2023;77: 67–88. doi:10.1146/annurev-micro-032421-

111116

56. Pogoutse AK, Moraes TF. Iron acquisition through the bacterial transferrin receptor. Crit

Rev Biochem Mol Biol. 2017;52: 314–326. doi:10.1080/10409238.2017.1293606

57. Ayalew S, Shrestha B, Montelongo M, Wilson AE, Confer AW. Identification and

immunogenicity of *Mannheimia haemolytica* S1 outer membrane lipoprotein PlpF. Vaccine.

2011;29: 8712–8718. doi:10.1016/j.vaccine.2011.08.074

58. Ayalew S, Confer AW, Blackwood ER. Characterization of immunodominant and

potentially protective epitopes of *Mannheimia haemolytica* serotype 1 outer membrane

lipoprotein PlpE. Infect Immun. 2004;72: 7265–7274. doi:10.1128/IAI.72.12.7265-

7274.2004

59. Confer AW, Ayalew S, Panciera RJ, Montelongo M, Wray JH. Recombinant *Mannheimia*

haemolytica serotype 1 outer membrane protein PlpE enhances commercial *M. haemolytica*

vaccine-induced resistance against serotype 6 challenge. Vaccine. 2006;24: 2248–2255.

doi:10.1016/j.vaccine.2005.11.036

60.   Pandher K, Confer AW, Murphy GL. Genetic and immunologic analyses of PlpE, a

lipoprotein important in complement-mediated killing of Pasteurella haemolytica serotype

1. Infect Immun. 1998;66: 5613–5619. doi:10.1128/IAI.66.12.5613-5619.1998

61.   Ayalew S, Blackwood ER, Confer AW. Sequence diversity of the immunogenic outer

membrane lipoprotein PlpE from *Mannheimia haemolytica* serotypes 1, 2, and 6. Vet

Microbiol. 2006;114: 260–268. doi:10.1016/j.vetmic.2005.11.067

62.   Oppermann T, Busse N, Czermak P. *Mannheimia haemolytica* growth and leukotoxin

production for vaccine manufacturing — A bioprocess review. Electronic Journal of

Biotechnology. 2017;28: 95–100. doi:10.1016/j.ejbt.2017.06.001

63.   Narayanan SK, Nagaraja TG, Chengappa MM, Stewart GC. Leukotoxins of gram-negative

bacteria. Vet Microbiol. 2002;84: 337–356. doi:10.1016/s0378-1135(01)00467-9

64.   Larsen J, Pedersen AG, Davies RL, Kuhnert P, Frey J, Christensen H, et al. Evolution of the

leukotoxin promoter in genus *Mannheimia*. BMC Evolutionary Biology. 2009;9: 121.

doi:10.1186/1471-2148-9-121

65.   Drula E, Garron M-L, Dogan S, Lombard V, Henrissat B, Terrapon N. The carbohydrate-

active enzyme database: functions and literature. Nucleic Acids Res. 2022;50: D571–D577.

doi:10.1093/nar/gkab1045

66.   Dik DA, Marous DR, Fisher JF, Mobashery S. Lytic transglycosylases: concinnity in

concision of the bacterial cell wall. Crit Rev Biochem Mol Biol. 2017;52: 503–542.

doi:10.1080/10409238.2017.1337705

67.   Martinez-Bond EA, Soriano BM, Williams AH. The mechanistic landscape of Lytic

transglycosylase as targets for antibacterial therapy. Curr Opin Struct Biol. 2022;77:

102480. doi:10.1016/j.sbi.2022.102480

68.   Suvorov M, Lee M, Hesek D, Boggess B, Mobashery S. Lytic Transglycosylase MltB of

Escherichia coli and Its Role in Recycling of Peptidoglycan Strands of Bacterial Cell Wall.

J Am Chem Soc. 2008;130: 11878–11879. doi:10.1021/ja805482b

69.   Scheurwater E, Reid CW, Clarke AJ. Lytic transglycosylases: bacterial space-making

autolysins. Int J Biochem Cell Biol. 2008;40: 586–591. doi:10.1016/j.biocel.2007.03.018