1

Identification of genes that differentiate *Mannheimia haemolytica* genotypes 1 and 2 using a

pangenome approach

Darien Deschner, Janet E. Hill*

Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon, Canada

*Corresponding author

E-mail: Janet.Hill@usask.ca

Abstract

Mannheimia haemolytica is an opportunistic bacterial pathogen associated with the economically costly bovine respiratory disease. Two genotypes have been described, of which genotype 2 is more strongly associated with disease. Several previous studies have investigated the genomic differences between the genotypes and/or the major serotypes (1, 2 and 6) of *M. haemolytica*, however we still lack a clear basis for the greater disease association of genotype 2 (serotypes 1 and 6) and demonstrations of phenotypic differences are scarce. This work builds upon previous investigations to identify genes that differentiate the two genotypes with a particular focus on genes that may play a role in virulence and fitness in the respiratory tract microbiome. We identified 422 genotype differentiating genes in a collection of 206 unique *M. haemolytica* genomes (61 genotype 1, 145 genotype 2). Genotype differentiating genes included genotype-associated variants of a TonB-dependent siderophore receptor homolog, transferrin binding protein B, leukotoxin A, and IgA1 proteases. We also identified a genotype 1 associated lytic transglycosylase, and a genotype 2 specific highly immunogenic outer membrane lipoprotein. Genotype 2 genomes were significantly larger in size and contained more predicted protein coding genes than genotype 1 genomes. These results expand our knowledge of what differentiates the genotypes 1 and 2 of *M. haemolytica* and provides information that can be used as the basis for laboratory investigations of corresponding phenotypic differences.

Introduction

Mannheimia haemolytica is a gram-negative opportunist bacterial pathogen associated with bovine respiratory disease (BRD) [1,2]. The beef industry is estimated to lose more than $3

billion per year globally due to BRD [3,4]. Two genotypes of *M. haemolytica* have been

described based on whole genome sequencing and single-nucleotide polymorphism (SNP)

analysis and multiple serotypes have also been defined [5,6]. The three most common serotypes

(1, 2 &, 6) have been found to correspond to genotypes with serotypes 1 & 6 corresponding to

genotype 2 (G2) and serotype 2 corresponding to genotype 1 (G1) [1,5,7,8]. Genotype 2 has been

more frequently associated with clinical cases of BRD, however the genetic and phenotypic basis

for this apparent association is unclear [1,9–14]. In addition to virulence factors such as

leukotoxin, lipopolysaccharide, adhesins and proteases [11,15–17], adaptations that confer

increased fitness, such as greater ability to acquire iron and other nutrients, could contribute to

disease association by allowing a pathogenic strain to outcompete non-pathogenic strain(s)

present in the same environment [18].

Reports of specific phenotypic differences between *M. haemolytica* G1 and G2 are

scarce. G2 strains corresponding to serotype 1 have been demonstrated to invade differentiated

bovine bronchial epithelial cells *in vitro* while G1 strains (serotype 2) could not [19]. The only

other phenotypic differences described to date are distinct matrix-assisted laser

desorption/ionization - time of flight mass spectrometry (MALDI-TOF) profiles and colony

morphology [20,21].

Genotypic differences between G1 and G2 have been investigated previously using whole

genome sequences. A pangenome analysis of 69 *M. haemolytica* genomes identified 112 G1

specific, and 179 G2 specific genes. This study was focused on outer membrane protein

encoding genes and identified seven G2 specific predicted outer membrane protein encoding

genes that were hypothesized to play a role in the greater disease association of G2 [22]. Another

study examined 11 *M. haemolytica* genomes for potential differences in virulence factors

between the serotypes 1, 2 and 6 [23]. The authors reported more prophages in serotypes 1 and 6

than in serotype 2, which could influence virulence due to the presence of virulence factors in

these prophages. Self-targeting CRISPR spacer sequences were detected in serotypes 1 and 6,

which may enhance immune evasion or adhesion during infection by modulating sialic acid

expression on the cells surface. ARG-carrying ICEs were more prevalent in serotypes 1 and 6

than serotype 2, which could enhance survival of antibiotic exposure [23].

The objective of our current study was to take advantage of the greater number of *M.*

haemolytica whole genome sequences now available to identify genetic differences between the

two genotypes. In our examination of >200 *M. haemolytica* genomes, we gave particular

attention to genes encoding proteins that may contribute to the reported differences in disease

association such as those involved in virulence and/or fitness in the bovine respiratory tract. We

designed an approach that allowed the detection of both genes with genotype associated

presence/absence patterns and sequence variants of genes conserved in both genotypes. This

contrasts with previous studies that have largely focused on the presence/absence of genes within

the genotypes. Our results showed that G2 genomes are significantly larger in size and contained

more predicted protein coding genes than genotype 1 genomes. We identified a total of 422

genotype differentiating genes, including several whose predicted functions may partially explain

the differences in observed disease association and that have not to our knowledge been reported

previously. This list of genes will inform future research to demonstrate phenotypic differences

between G1 and G2.

Materials & Methods

Genome sequences

M. haemolytica whole genome sequences defined as complete by NCBI were downloaded from the NCBI Genome database on March 25, 2024, using the NCBI Datasets command line tools [24]. NCBI defines "complete" as all chromosomes are gapless and contain runs of nine or less ambiguous bases, no unplaced or unlocalized scaffolds, and all expected chromosomes present. To avoid duplication all .gca files that had a matching .gcf file were removed. To identify G1 and G2 genomes, the sequences were genotyped using an *in silico* PCR assay in Unipro UGENe v45.1 and primers from a genotype specific LAMP assay with default parameters (Mismatches = 3, Min Perfect Match =15, Max Product Size = 5000, Use ambiguous bases = True, Extract annotations = Inner, and Temperature settings = Primer 3) [25,26]. Primers from the LAMP assay included a pair targeting G1 specific adhesin pseudogene B1, G2 adhesin G (CP017538.1 locus tag: BG586_06285; Forward: 5'-GCA CAT TAA AAT ATA GCA GCT TTG-3'; Reverse: 5'-CTA AGC CAG AGT GAT CCG-3'), and leukotoxin D (*lktD*) (CP005972.1 locus tag: F382_07400; Forward: 5'-ACT GAA AA TTC ACA CTA TAG GTG-3'; Reverse: 5'-GCT AAT ATG TTT AAT TCG ACC AGT T-3'). The *lktD* primers were used to confirm identity of genomes as *M. haemolytica*. In initial testing G1 specific primers failed to produce any predicted products in our G1 reference strain and so were discarded. New G1 specific primers targeting the G1 specific adhesin pseudogene B1 were developed using Primer 3 v 4.1.0 (CP017495.1 locus tag BG548_01640; Forward: 5'-ACT AAT CTG AAG AGC GGC GT-3'; Reverse: 5'-TTG GTG GTT GCT GCT TGT AC-3') [27–29]. G1 and G2 NCBI genomes were combined with a set of G1 and G2 genome sequences previously generated by our lab

97   (BioProject: PRJNA1088094)(30) and reference genomes (G1 = CP017518; G2 = CP017519)

98   for both genotypes.

99   A phylogenetic tree was constructed using MASH with the ani command in PanTools to

100   verify genotype designations [30].

Pangenome construction, annotation, and classification

A pangenome was constructed using PanTools v 4.3.2 with FASTA files, gff3 annotation

files, and genotype designation as inputs [30–32]. Genomes were assigned a new unique

identifier based on their input order during pangenome construction (S1 Table). Genome metrics

such as genome size, GC content, number of genes, etc. were determined using the metrics

command.

Pantools groups protein sequences from the input genomes into homology groups based

on their similarity. Homology groups also contain the nucleotide sequences encoding the

constituent protein sequence(s). Homology groups were generated using the group command

with the relaxation parameter set to 2. The relaxation parameter is a combination of four sub-

parameters (intersection rate, similarity threshold, mcl inflation, and contrast) where a lower

number represents a stricter set of thresholds. We elected to utilize a relaxation parameter of 2 to

allow for detection of minor variations that might be expected when examining sub-populations

of the same species and avoid grouping of more divergent genes as might occur with a looser

parameter.

Homology groups were classified by PanTools as core (present in $\geq$ 90% of all genomes),

unique (present in only one single genome), or accessory (present in >1 genome and <90% of all

genomes). Based on their prevalence within each genotype, homology groups were classified as

either shared (present in ≥ 90% genomes of one genotype, regardless of their prevalence in the

other genotype), exclusive (solely present in genomes of one genotype), or specific (present in ≥

90% genomes of one genotype, and no genomes of the other genotype) [30]. To capture

homology groups strongly associated with one genotype that did not meet the strict PanTools

definition of specific we used the classification "genotype associated" for homology groups that

were present in ≥ 95% of one genotype while also being present in > 0% and ≤ 5% of the other

genotype.

Genomes were functionally annotated using InterProScan v 5.64-96.0 with SignalP v 4.1

database integrated [33–35]. Biosynthetic gene cluster annotations were generated using

antiSMASH 7.1.0 with default parameters [36]. InterPro and antiSMASH annotations were

added to the pangenome, with InterProScan annotations including InterPro, Pfam, GO, and

SignalP annotations [30].

Multiple sequence alignments of constituent nucleotide sequences were generated for all

homology groups in the pangenome with MAFFT and consensus sequences from the alignments

were generated to represent the genotype specific and associated homology groups [30,37].

Consensus sequences were used as queries for BLASTx searches using the bacterial/archaeal

translation code against the NCBI refseq_select database [38–42]. All other parameters were kept

as default. Searches were performed March 20-25, 2025. The top hit for each search was

manually recorded to provide further evidence of the predicted function, facilitate human

interpretation, and identify functions that may not have been annotated by any of the other

approaches. The entire pangenome database is available in S1 Dataset [37].

The nucleotide sequences of several previously characterized genes upregulated in
response to iron limitation, and previously identified proteases were used as BLAST queries
against the pangenome using the blast command in PanTools to determine if these previously
characterized genes were present in our dataset [15,30,43]. Both alignment threshold and
minimum identity were set to 90. Query nucleotide sequences are available S2 Table.

Tet(H) gene abundance

Tet(H) genes were used as a proxy for the presence of integrative and conjugative
elements (ICE) due to the strong correlation between presence of tet(H) and presence of ICE
[12,44]. The reference tet(H) nucleotide sequence (Accession Number: Y15510.1 (nucleotides
2099-3301)) was downloaded from the comprehensive antibiotic resistance database (CARD) on
July 26, 2024, and used as nucleotide BLAST query against the pangenome database using the
inbuilt PanTools blast command. Minimum identity and alignment threshold were both set to
90%.

Leukotoxin operons

Reference nucleotide sequences for each of the component genes of the *M. haemolytica*
leukotoxin operon (*lktCABD*) were downloaded from GenBank (AF314508.2) and used as
queries against the pangenome BLAST database with percent identity and alignment threshold
both set to 90%. Consensus nucleotide sequences of G1 and G2 versions of all leukotoxin operon
genes were extracted and translated in Geneious 2024.0.5. Predicted protein sequences for the
G1 and G2 versions of all leukotoxin operon genes were aligned against one another in Geneious
2024.0.5.

Annotation of carbohydrate active enzymes

Carbohydrate active enzymes (CAZymes) were annotated using a local installation of dbCAN v 4.1.1 software with integrated SignalP v 4.1 database and DIAMOND, HMMER, and HMMER dbcan_sub prediction algorithms [45]. Annotation was performed on each genome sequence individually using a custom BASH script [37]. dbCAN annotations that had supporting annotations from at least three tools and contained a predicted signal peptide were selected for further analysis.

Protein structure prediction and visualization

Protein sequences were submitted to the AlphaFold Server for structural prediction [46]. Models with the highest ranking score were downloaded and visualized in PyMol version 2.5.2 (Schrödinger, LLC).

Statistics

Welch's t-test for unpaired data was used to compare G1 and G2 genome characteristics with $P \leq 0.05$ considered significant and 95% confidence intervals for the difference in means were reported. A one-way ANOVA (Kruskal-Wallis) with Dunn's test for multiple comparisons was used to determine if there was a difference in genome size between G1 and G2 with or without ICE. $P \leq 0.05$ was considered significant. These analyses were performed in GraphPad PRISM v 10.4.1.

A Fisher's exact test and odds ratio comparison was performed to determine if the prevalence of tet(H) was different between genotypes. Both tests were performed in Microsoft

182 Excel for Microsoft 365 using Real Statistics Resource Pack. $P \leq 0.05$ was considered

183 significant.

Results & Discussion

Description of genome collection

M. haemolytica genomes designated as complete were retrieved from NCBI and after

removal of duplicates, 121 genomes were available for further analysis. *In silico* PCR with

modified primers from a previously published genotyping assay resulted in the expected

amplicons for 108/121 genomes. No *in silico* genotyping PCR products were obtained for 13

genomes and so they were removed from further analyses leaving 108 genomes.

Three of the 13 isolates that could not be genotyped also failed to produce any predicted

product for the *lktD in silico* PCR. Two of these genomes have since been removed by the

RefSeq curation team for either containing too many frameshift mutations (GCF_900474405) or

failing to pass a completeness check (GCF_023516295). All three genome sequences that failed

to produce an *lktD* product were from sheep isolates (NCTC 10643, NCTC 10609) or from

culture collections from the 1960s (ATCC 29701). Of the other ten genomes that were not

genotyped by *in silico* PCR, four were from a German study examining macrolide resistance in

M. haemolytica [44], and three were from sheep isolates collected in 1951 or 1968 and

sequenced as part of the United States Department of Agriculture (USDA) *M. haemolytica*

sequencing project (BioProject: PRJNA172854). The last three isolates were all generated from

non-nasopharyngeal cattle samples as part of either the USDA sequencing project

(GCF_000819525, GCF_023516255) or a German study (GCF_020971705) and had a CheckM

completeness <92.5%.

The *in silico* PCR assay we used was selected for identification of G1 and G2 genomes as

defined by Clawson et al. [5]. The original LAMP assay from which we selected primers for this

process was developed using *M. haemolytica* isolates from North American cattle [25]. Thus, the

most likely explanation for genomes to be un-typable by this approach (besides sequence quality

issues) is that they represent genotypes other than G1 or G2.

The NCBI genomes were combined with 99 genomes from our previous work

(BioProject Accession PRJNA1088094) and reference genomes for G1 and G2 as described in

the Methods. One additional genome was removed following discovery that it was likely a

chimera, leaving a final dataset 208 *M. haemolytica* whole genomes with 62 G1 and 146 G2

representatives (S1 Table). The reference genomes for G1 and G2 were also represented in the

collection of genomes downloaded from NCBI resulting in the collection representing 206

unique *M. haemolytica* genomes. Most genomes were from isolates originating from feedlot

cattle in either Canada or the United States between 2003 and 2021. Consistent with the *in silico*

PCR identifications, a pangenome derived phylogenetic tree showed two distinct clades

corresponding to isolates classified as G1 or G2 (Figure S1).

Genome properties & Tet(H) abundance

The two genotypes were compared to one another on a broad set of genome metrics, such

as genome size, GC content, etc., to determine if there were any high-level differences in the

genomic structure and content of the two genotypes. G2 genomes were larger, had more

predicted protein-coding sequences, and had slightly lower GC content than G1 genomes. No

differences were identified in the number of tRNA or rRNA genes or the proportion of the

genome covered by genes (Table 1).

Table 1. Comparison of genome features of G1 and G2. Values are mean ± standard deviation.
P-values are from a Welch's t-test with *P*<0.05 considered significant.

Metric	Genotype 1 (n = 62)	Genotype 2 (n = 146)	*P*-value	95% CI difference between means
Genome size (bp)	2593014 ± 44574	2718798 ± 47361	<0.0001	[112154, 139415]
GC content (%)	41.14 ± 0.03	41.04 ± 0.03	<0.0001	[0.10, 0.11]
Gene count	2567 ± 108	2798 ± 116	<0.0001	[198, 264]
Genome covered by Genes (%)	88.17 ± 1.48	88.73 ± 1.23	0.0093	[0.14, 0.99]
tRNA count	64.3 ± 1.7	64.2 ± 2.0	0.6088	[-0.68, 0.40]
rRNA count	19.0 ± 4.3	19.7 ± 2.4	0.2500	[-0.49, 1.84]

We hypothesized that the larger size of G2 genomes was attributable to a greater

prevalence of integrative and conjugative elements (ICEs), large (~20 kb to >500 kb) mobile

genetic elements that are known to be associated with antimicrobial resistance and are more

frequently identified in G2 genomes [5,47]. To test this hypothesis, we used the proxy measure

of the presence of tetracycline resistance gene tet(H) to determine the prevalence of ICEs due to

the strong association of tet(H) with ICEs [12,44].

Tet(H) was identified significantly more frequently in G2 (67/146, 45.9%) than G1 (1/62,

1.6%) (Fisher's *p*-value = 4.39 x 10^{-12} Odds Ratio: 0.02 ± 1.02), however, G2 genomes were

larger than G1 regardless of the presence of ICE (Figure 1). The larger size and higher gene

count of G2 genomes regardless of presence of ICE (Table 1) could indicate that G2 genomes

carry a larger repertoire of genes that may allow them to be more adaptable to a wider variety of

environments than the smaller G1 genomes. Considering average bacterial gene sizes, the

difference in the median genome size of G2 without ICE and G1 (~100 kb) could represent

dozens of G2-specific genes. Additionally, the greater prevalence of ICEs in G2 suggests that this

more disease associated genotype is also more likely to possess antimicrobial resistance genes.

244
245
246   **Figure 1.** Genome sizes of Genotype 1 (G1, n = 62), Genotype 2 (G2, n = 146), G2 possessing
247   integrative and conjugative elements (ICEs, n = 66), and G2 without ICEs (no ICE, n = 80). One
248   way ANOVA (Kruskal-Wallis) followed by Dunn's multiple comparisons test (corrected *P* values
249   are shown). Red bars indicate mean.
250

Pangenome analysis

The pangenome calculation resulted in the identification of "homology groups": clusters

of sequences from input genomes that exceed a given similarity threshold. We then examined the

distribution of homology groups across all *M. haemolytica* genomes to identify those that are

core, unique, accessory, shared, genotype specific or genotype associated as defined in the

methods. A total of 4,038 homology groups were generated of which 2,007 (49.7%) were core,

1,473 (36.5%) were accessory, and 558 (13.8%) were unique (Pangenome_Classification in S3

Table).

For the current study, our interest was homology groups that differentiate the genotypes:

those that are found in the large majority of representatives of one genotype and in none, or a

small minority, of the other genotype. PanTools identified 135 G1 specific and 233 G2 specific

homology groups. The PanTools specific category definition is strict, however, requiring a

homology group to be present in >90% of one genotype and zero of the other. To identify

homology groups that do not meet this strict criterion but are nevertheless strongly associated

with one genotype, we examined accessory groups not initially classified as specific for those

that were present in ≥ 95% of the one genotype and > 0% and ≤ 5% of the other genotype. These

were classified as G1 associated (n = 21) or G2 associated (n = 34) (S3 and S4 Tables).

Identification of genotype specific and associated genes within this relatively large genome

collection reinforces previous findings that the two genotypes carry distinct sets of genes [22].

The pangenome results also provide evidence of the internal coherence of the genotypes, since

when considered separately, G1 and G2 genomes included 2,169 and 2,342 "genotype-core"

homology groups (present in >90% of genomes of the genotype), respectively.

To provide an overview of the potential phenotypic differences between the two

genotypes, all specific and associated homology groups were assigned to one of nine general

functional categories based on functional annotations and the results of manual BLASTx

searches against the NCBI refseq_select database (Figure 2, S4 Table, S5 Table). For both

genotypes, the largest number of specific and associated homology groups were classified as

MGE proteins (transposases, toxin-antitoxin plasmid maintenance systems, prophage associated

proteins, etc.) or hypothetical. The larger number of G2 specific and associated MGE proteins

and antimicrobial resistance genes is consistent with greater ICE prevalence in G2

[9,12,22,23,44]. The large numbers of genotype specific and associated hypothetical proteins in

both genotypes is also unsurprising as ~22□% of all domain families in Pfam v32.0 are defined

as "domains of unknown function" and a recent examination of >30,000 bacterial genomes and

metagenomes showed that 40-60% of genes are of unknown function [48,49]. Homology groups

predicted to encode unknown/hypothetical proteins, and MGE proteins were excluded from

further analysis in this study but certainly represent an opportunity for future investigation.

287
288
Figure 2. Predicted functional categories of genotype specific/associated homology groups (n =
156 for G1; 266 for G2). G1 = Genotype 1, G2 = Genotype 2. Specific groups are those found in
≥90% of the genomes of the given genotype and no genomes of the other genotype. Associated
groups are present in ≥95% of the genomes of one genotype > 0% and ≤5% of the genomes of
the other genotype. MGE proteins include transposases, toxin-antitoxin plasmid maintenance
systems, prophage associated proteins, etc. Hypothetical proteins include those annotated as
hypothetical or domain of unknown function.

Based on inspection of the complete inventory of pangenome homology groups (Table

S3, S4), we identified a subset of specific/associated homology groups with functional

annotations that may be associated with reported differences between disease association of G1

and G2 (Table 2). These genome features were related to iron acquisition, host immune evasion,

immunogenicity and carbohydrate modification.

Table 1. Genotype association and predicted functions of selected homology groups. Annotation
notes summarize the functional annotations from InterProScan for each homology group.
BLASTx search used consensus nucleotide sequence of the homology group as query against the
refseq_select database. Reported top BLASTx hits were from *M. haemolytica* unless otherwise
stated.

Homology group	Genotype association	InterProScan Annotation	BLASTx hit (top)	Predicted Function
2042480	G2 specific	TonB-dependent receptor	TonB-dependent siderophore receptor	Iron Acquisition
2042481	G1 associated	TonB-dependent receptor	TonB-dependent siderophore receptor	Iron Acquisition
2041941	G2 specific	TonB-dependent receptor	TonB-dependent receptor plug-domain containing protein	Iron Acquisition
2944545	G2 specific	Transferrin binding protein (TpbB)	Transferrin-binding protein-like solute binding protein	Iron Acquisition
2040773	G1 specific	Transferrin binding protein	Transferrin-binding protein-like solute binding protein. *Mannheimia pernigra*	Iron acquisition
2042189	G2 specific	Peptidase S6/Autotransporter/IgA1 protease	S6 Family Peptidase	Host immune evasion
2040817	G1 associated	Peptidase/Pectate Lyase/IgA1 Protease	S6 family peptidase	Host immune evasion
2044609	G2 specific	N/A	Outer membrane lipoprotein PlpE.	Highly immunogenic outer membrane protein
2042729	G1 associated	MltB-like Transglycosylase	Lytic murein transglycosylase. *Mannheimia glucosida*	Cell wall remodeling/recycling

Iron acquisition

Iron is essential for bacterial growth and bacterial pathogens must compete with each

other and with the host for the scant iron typically available in their environment [50,51]. Animal

pathogens like *M. haemolytica* are faced with concentrations of available iron well below their

needs, which has necessitated the evolution of mechanisms for acquiring iron including

siderophores, transferrin binding proteins, and haem binding proteins [52]. Given the essential

nature of iron, and the generally iron limited conditions of the respiratory tract, the ability to

acquire iron more readily than other bacteria provides a considerable advantage. The

identification of several genotype associated, and specific homology groups potentially involved

in iron acquisition was of interest since these differences may be associated with fitness in the

respiratory tract and thus with disease association (Table 2).

Homology groups 2042480 (G2 specific) and 2042481 (G1 associated) were both

annotated as encoding a TonB-dependent siderophore receptor. TonB dependent receptors

(TBDTs) are transporters often involved in iron acquisition, specifically in the uptake of small,

secreted iron scavenging molecules known as siderophores [53,54]. Homology groups 2042480

and 2042481 also produced BLAST hits to a previously characterized siderophore (ferric

enterobactin) receptor (*frpB*) homolog that was upregulated in *M. haemolytica* in response to iron

limitation [43]. The consensus amino acid sequences of these homology groups were 84.4%

identical, and both contained consensus TonB-dependent transporter (TBDT) motifs. Predicted

3D structures demonstrated the canonical 22 β-strand barrel structure with an N-terminal plug

domain characteristic of TBDTs (AlphaFold ranking_score = 0.95 for both) (Figure 3A). The

amino acid sequence differences between the G1 and G2 sequences were concentrated in the

extracellular loops responsible for substrate binding, suggesting that these two proteins may

differ in their ability to bind their substrate [53,54] or they may bind different substrates (Figure

3B). Identification of a putative siderophore receptor in *M. haemolytica* is unexpected as *M.*

haemolytica is not known to produce or utilize siderophores and has been demonstrated to be

unable to utilize ferric enterobactin so it is likely that the identified TBDTs have some other

function besides siderophore uptake [11,43].

336
337
Figure 3. A) Predicted 3D structures of Genotype 1 (G1, Purple) & Genotype 2 (G2, Orange)
ferric enterobactin receptor (frpB) homolog proteins. 3D structure conforms to canonical TonB-
dependent receptor structures. B) 2D representation of tertiary structure of G1 and G2 frpB
homologs. Each grey rectangle represents one of the 22 β-strand sheets, and the size of the
extracellular/periplasmic loops are shown. Differences in the amino acid sequence for select
extracellular loops are shown in further detail. Difference in extracellular loops may contribute to
differences in substrate binding efficiency and/or result in interactions with completely different
substrates. G1 AlphaFold ranking_score = 0.95; G2 AlphaFold ranking_score = 0.95.
346

G2 specific homology group 2041941 was also predicted to encode a TBDT. No similar

G1 specific/associated protein was identified and the consensus sequence of 2041941 did not

correspond to previously identified genes upregulated in response to iron limitation [43]. The

identification of another G2 specific TBDT suggests that G2 may have a greater ability to uptake

iron and/or other substrates. TBDT have been implicated in transport of a variety of substrates in

gram-negative bacteria including siderophores as well as vitamin B_{12} and carbohydrates [54].

Further examination is required to determine the function of this TBDT and whether it provides a

competitive advantage to G2 over G1 strains.

Other iron acquisition proteins include transferrin binding proteins A and B (*tpbAB*), both

of which are found in *M. haemolytica* [11]. Transferrin is an iron-carrying glycoprotein found in

mammalian circulatory systems that transports iron throughout the body and serves as a sink for

any excess free iron released into the circulatory system [55]. TpbB is a surface lipoprotein that

extends outwards from the outer membrane to the extracellular space where it binds to transferrin

and brings the transferrin closer to TbpA, a TBDT in the outer membrane that transports the

transferrin into the periplasm [55]. To investigate any genotype associations of transferrin

binding proteins, previously characterized *tbpA* and *tbpB* nucleotide sequences from *M.*

haemolytica were used as BLAST queries against the pangenome database. Core homology

group 2044546 contained hits for *tbpA* while *tbpB* hits were solely identified in G2 specific

homology group 2044545 with no hits in any G1 genomes. These results suggest that *tbpA* is

highly conserved within *M. haemolytica* and likely plays a key role in iron metabolism. In

contrast, while no BLAST hits were produced to the query *tbpB* nucleotide sequence in G1

genomes, G1 specific homology group 2040773 was predicted to encode for a transferrin binding

protein. The amino acid sequences of G1 and G2 variants of the putative TbpB proteins

(homology groups 2040773 and 2044545) were compared and found to be 56.5% identical and

81.2% similar. Despite the relatively low similarity of these two proteins, the predicted 3D

structures were very similar to one another as well as the crystal structure for the TbpB from

Actinobacillus pleuropneumoniae (G1 ranking_score = 0.88, G2 ranking_score = 0.93, Figure 4).

A difference in the TbpB structure between the genotypes was observed in the N lobe with G2

appearing to have a longer alpha-helix structure in the N-lobe handle region than G1 or *A.*

pleuropneumoniae. This region of the protein contains residues important for binding of iron-

loaded transferrin [56]. Genotype variation of a transferrin binding protein further suggests they

may differ in their ability to acquire iron, which may partially explain observed differences in

disease association between the two genotypes. The genotype variants of TbpB identified here

may also reflect host preferences of the genotypes as TbpB is selective for host transferrin and

differences between TbpB proteins has been partially attributed to this host selectivity [55].

382
383
Figure 4. Predicted structures of G1 and G2 variants of a predicted transferrin binding protein B
(TbpB). (A) The crystal structure of TbpB from *Actinobacillus pleuropneumoniae* (PDB
accession 3HOL [56]) was used as a reference. Major domains include the N-lobe and C-lobe,
each containing a beta-barrel and a "handle". The reference structure is rainbow coloured from N
terminus (blue) to C terminus (red). (B) Alignment of G1 and G2 variants. (C) G2 aligned to
reference. (D) G1 aligned to reference. The G1 variant has an elongated alpha-helix in the N-lobe

handle domain relative to G2 and the reference. (G1 ranking_score = 0.88, G2 ranking_score =
0.93).

IgA protease

IgA proteases are metalloproteases that cleave immunoglobulin A (IgA), a critical

component of the mucosal immune system. Two IgA proteases (IgA1 and IgA2) have been

previously identified and characterized in a G2 (serotype 1) *M. haemolytica* isolate [15]. We

identified a G2 specific IgA protease (homology group 2042189) and a G1 associated IgA

protease (2040817) (Table 2). To determine the relationship of these sequences to the previously

reported IgA proteases, a BLAST search was performed using previously reported nucleotide

sequences for IgA1 and IgA2 as queries against the pangenome database (IgA1:NZ_DS264709

Locus Tag: MHA_RS00165IgA2: NZ_DS264612.1 Locus Tag: MHA_RS09765). At least one

BLAST hit was returned for IgA2 in all but five of the included genomes with hits grouped into

five homology groups (2041035, 2041036, 2041399, 2041400, and 2043723). Homology groups

2041305 and 2041306 were accessory, while 2041399 and 2041400 were unique suggesting

some minor variations are likely present within the pangenome for this IgA2 protease. Homology

group 2043723, however, was a core group suggesting that this IgA2 protease is a core gene of

M. haemolytica and is not genotype differentiating. In contrast, a BLAST hit was returned for

IgA1 in 146/209 of the genomes with hits grouped into one of three homology groups (2040817,

2041298, and 2042189). Homology group 2041298 was found in only one genome so was not

examined any further. Group 2040817 is a G1 associated group while 2042189 is a G2 specific

group and thus these groups appear to represent genotype variants of an IgA protease.

Comparison of the consensus sequences of the two IgA1-like homology groups revealed

that the nucleotide sequence of 2042189 (G2 specific) is 4,512 bp consistent with previous

descriptions of IgA1 in G2 *M. haemolytica* [15], while the sequence of 2040817 (G1 associated)

was only 2,281 bp. An alignment of the two variants showed that they were 94% identical over

2,281 shared positions and that the G1 associated sequence is a truncated version of the other.

The *M. haemolytica* IgA1 protease contains three functional domains: a large, N-terminal

peptidase domain, pertactin-like passenger domain and a C-terminal autotransporter domain [15].

The product of the truncated G1 associated gene would include the peptidase domain, but lack

part of the pertactin-like passenger domain and the autotransporter, and thus may not encode a

complete or functional IgA1 protease.

The observation of a G2 specific IgA1 sequence and possibility of genotype variation

remains of interest and warrants further investigation since differences in IgA protease activity

could relate to difference(s) in their ability to evade the host immune system and thus may

contribute to differences in disease association.

Highly immunogenic outer member lipoprotein

Pasteurella lipoproteins (Plp) are a group of several immunogenic outer membrane

lipoproteins identified in *M. haemolytica* [57] and PlpE has been investigated as a potential

vaccine target for *M. haemolytica* because of its immunogenicity [58–60]. Aside from the

immunogenic role of PlpE relatively little is known regarding its function although it is

important as a vaccine target and thus clinically and diagnostically significant.

G2 specific homology group 2044609 was predicted to encode PlpE, however no

corresponding G1 homology group was identified. Previous investigations of PlpE sequence

diversity across serotypes showed that the serotype 1 and 6 (G2) sequences were highly

conserved whereas serotype 2 (G1) PlpE sequences were extremely heterogenous, however,

antibodies raised against serotype 1 PlpE cross reacted with serotype 2 bacteria [61]. To identify

potential G1 variants of the G2 *plpE*, a reference *plpE* gene (NZ_CP023044.1, c780551-779481,

locus tag: CKG22_RS03975) was used in a BLASTx search with relaxed parameters. When

percent identity threshold was reduced to 70% and alignment threshold to 50% a hit to G1

specific homology group 2042820 was observed.

The identification of a sequence encoding a G2 specific PlpE and possible G1 variant

suggests that the genotypes may differ in immunogenicity, however further research is needed to

investigate the prevalence and immunogenicity of Plp variants across *M. haemolytica* genotypes.

Leukotoxin operon

Leukotoxin (Lkt) is a critical virulence factor in *M. haemolytica* and is produced by both

genotypes (all serotypes) [62]. Lkt is encoded in an operon consisting of four genes *(lktCABD)*.

lktA encodes the protoxin, *lktC* encodes the acylating enzyme essential for post-translational

modification and activation of the protoxin, and the products of *lktB* and *lktD* are involved in

secretion of the toxin [63,64]. *lktCABD* genes were identified and the predicted protein

sequences for the Lkt operon genes from G1 and G2 were aligned (Figure 5). The predicted

proteins LktC, LktB and LktD were 95.2%, 99.6% and 100% identical, respectively, suggesting

these proteins are under conservation pressure, likely due to their roles in regulating the secretion

and function of Lkt [64,65]. In contrast, the predited LktA protein sequences were only 89%

identical. Differences in the predicted LktA proteins of G1 and G2 were concentrated in the N-

terminal and C-terminal regions of the protein (Figure 5). In leukotoxins the N-terminus is

involved in pore-forming activities while the C-terminus is involved in export and host immune

system recognition [64]. As such any differences in the N-terminal and/or C-terminal regions

may potentially influence the activities of the Lkt and thus may result in virulence differences

between the two genotypes.

460
461
462   **Figure 5.** Sequence alignment of G1 and G2 LktA amino acid sequences. Percent sequence
463   identities were calculated with a sliding window size of 10. A larger peak indicates a greater
464   number of matching amino acids within the window for that position. Locations of pore forming
465   domain and export signal domain are indicated above the identity plot. Schematic below the
466   identity plot shows identical (black) and non-identical (light grey) residues in the alignment.
467

Sequence diversity in LktA within *M. haemolytica* has recently been reported to be

greater than within other species in the genus [66]. Davies *et al*. examined the sequence diversity

of leukotoxin in 31 *M. haemolytica,* 6 *Mannheimia glucosida,* and 4 *Pasteurella* [*Bibersteinia*]

trehalosi isolates identified eight *M. haemolytica* allelic variants of *lktA* [63]. Five of the alleles

(A6-A10) were identified solely in ovine isolates, allele A3 was only identified in bovine

isolates, while A1 and A2 alleles were identified in both ovine and bovine isolates. The A1 allele

was subdivided into 5 groups (A1.1-A1.5) with A1.1 solely occurring in bovine isolates from

serotype 1 (n = 3) and 6 (n = 1) isolates (corresponding to G2). In bovine isolates from serotype

2 (corresponding to G1), the authors reported detection of alleles A2, A2.1, A2.2, and A3. LktA

sequences from genomes in our current study were compared with representative amino acid

sequences for each allele and subgroup downloaded from NCBI Genbank. Within G2, 141/146

LktA sequences were 100% identical to the A1.1 allele and another 4/146 were >99.5% identical

to A1.1, consistent with the observations of Davies *et al*. (S6-S8 Tables). The only exception was

the LktA from Genome 2, that shared 100% amino acid identity with A1.3 and A1.5 (S9 Table).

Interestingly, all but one of the G1 LktA sequences were 99.5-100% identical to the A8.1 allele

(Genome 157 was identical to A2) (S6 and S9-10 Tables). The A8.1 allele was found in serotype

2 (G1) isolates by Davies *et al*., however, all the isolates in that study were ovine (n = 3), and

A8.1 was also detected in ovine serotype 7 isolates (n = 3). The identification of A8.1 across

many bovine origin genomes and nearly complete absence of A2 alleles suggests that G1 from

Canada and the U.S.A has diverged from the G1 found in the United Kingdom where most of the

isolates used by Davies *et al*. were collected and further demonstrates that A8.1 is not restricted

to ovine isolates [63]. Whether there is a relationship between specific LktA alleles and

respiratory disease severity in cattle warrants further investigation.

CAZymes

Carbohydrate active enzymes (CAZymes) are enzymes whose function has been

biochemically verified to be involved in the metabolism of carbohydrates. We chose to

investigate any differences in CAZymes between the two genotypes since these may indicate

differences in ability to metabolize certain carbohydrates, and thus in the nutritional niches of the

genotypes [67]. We were particularly interested in enzymes predicted to be secreted into either

the periplasmic or extracellular space (i.e. contain a signal peptide) and are thus most likely to be

involved in interactions with the external environment. A total of six CAZymes containing signal

peptides were identified in the pangenome (S11 Table) (50). At least one CAZyme was predicted

in all genomes except for Genome 86/GCF_022404935, which was removed from this analysis.

CAZymes identified were members of families GH23 (lytic transglycosylase), GH33 (sialidase),

GH13_19 (amylose glucosidase), GT9 (glycosyltransferase), GH102 (lytic transglycosylase) and

GH103 (lytic transglycosylase). Only GH103 (homology group 2042729) was genotype

associated, present in 60/61 (98.4%) G1 genomes and 2/146 (1.37%) G2 genomes (Table 2). The

GH103 family consists of a group of lytic transglycosylases that cleave the β-1,4 linkage

between *N*-acetylmuramoyl and *N*-acetylglucosaminyl residues in peptidoglycan and belong to

family 3 of the transglycosylases with structures similar to lysozyme [68,69]. Numerous

biological functions have been attributed lytic transglycosylases, however the only clear role is

that they are involved in cell wall remodelling [68–71]. Whether this genotype difference in lytic

transglycosylases is associated with phenotypic differences in colony morphology of the two

genotypes [21] or other differences in cell wall maintenance will require further investigation.

Conclusions

We identified a total of 422 *Mannheimia haemolytica* genotype differentiating genes (135

G1 specific, 21 G1 associated, 233 G2 specific, and 33 G2 associated), including 125 with

unknown functions and 132 that are associated with mobile genetic elements. Genotype

associated genes and variants that may contribute to differences in disease association include

TonB-dependent transporters, IgA proteases, a highly immunogenic outer membrane protein

(PlpE), a lytic transglycosylase, and genotype variants of the leukotoxin gene. The list of

genotype differentiating genes provides a shortlist of targets that warrant further investigation to

provide a mechanistic understanding of phenotypic differences between the *M. haemolytica*

genotypes and inform rational strategies for *M. haemolytica* and bovine respiratory disease

diagnostics.

Acknowledgements

The authors are grateful to Dr. David Palmer (Department of Chemistry, University of Saskatchewan) for assistance with the protein structure alignments, and to Dr. Maarten Voordouw (Department of Veterinary Microbiology, University of Saskatchewan) for assistance with statistical analysis and approaches.

Supporting Information

S1 Dataset. Pangenome database. https://doi.org/10.6084/m9.figshare.28872536.v1

S1 Table. Summary of Genomes Included in Pangenome Analysis

S2 Table. Accession numbers of sequences used as BLAST queries against pangenome database.

S3 Table. Overview of pangenome classification of all homology groups. Genes matching is the number of genes that were found to correspond to the given homology group, as this referred to genes the number may be greater than or less than the number of genomes examined. For pangenome classification: core = group was present in ≥90% of all genomes (187/208); unique = Group was present in exactly one genome; Accessory = Group was present in somewhere between 2-186 genomes. For Genotype classification: Shared = Group was present in ≥90% of the genomes of the given genotype with no consideration of the presence/absence pattern of the other genotype; exclusive = Group was present solely in genomes of the given genotype; Specific = Group was exclusive to one genotype and present in ≥ 90% of that genotype;

Associated = Group was present in ≥95% of the genomes of one genotype and >0% and ≤5% of

the other genotype.

S4 Table. Genotype Specific and Associated Homology Groups. Annotation notes column

provides summary of the interpro, pfam and GO functional annotations. Genes matching is the

number of genes that were found to correspond to the given homology group, as this referred to

genes the number may be greater than or less than the number of genomes examined. For

pangenome classification: core = group was present in ≥90% of all genomes (187/208); unique =

Group was present in exactly one genome; Accessory = Group was present in somewhere

between 2-186 genomes. For phenotype (Genotype) classification: Shared = Group was present

in ≥90% of the genomes of the given genotype with no consideration of the presence/absence

pattern of the other genotype; exclusive = Group was present solely in genomes of the given

genotype; Specific = Group was exclusive to one genotype and present in ≥ 90% of that

genotype; Associated = Group was present in ≥95% of the genomes of one genotype and >0%

and ≤5% of the other genotype. For manual function categories: Metal Related = Predicted

protein function involved in metal metabolism and/or heavy metal resistance, Transporters &

Adhesins = Predicted protein is a transporter or adhesin, AMR = Predicted protein is an

antimicrobial resistance gene. Regulatory = Predicted protein has a regulatory function,

Restriction Endonuclease = Predicted protein is a restriction endonuclease or related to

restriction modification, Metabolism = Predicted protein has a function in metabolism such as

enzymes involved in carbohydrate degradation, MGE = Predicted proteins are characteristic of

mobile genetic elements such as transposases, toxin-antitoxin plasmid maintenance systems,

prophage associated proteins etc., Hypothetical = Predicted proteins are hypothetical/putative or

a protein with a domain of unknown function (DUF) that lacks any further functional

information, Other = Predicted protein could not be categorized into one of the other categories

S5 Table. Overview of the number of specific and associated homology groups assigned to each

functional category split by genotype classification. Specific = Group was exclusive to the given

genotype and present in ≥ 90% of genomes of that genotype, Associated = Group was present in

≥95% of the genomes of one genotype and □0% and ≤5% of the other genotype. For manual

function categories: Metal Related = Predicted protein function involved in metal metabolism

and/or heavy metal resistance, Transporters & Adhesins = Predicted protein is a transporter or

adhesin, AMR = Predicted protein is an antimicrobial resistance gene. Regulatory = Predicted

protein has a regulatory function, Restriction Endonuclease = Predicted protein is a restriction

endonuclease or related to restriction modification, Metabolism = Predicted protein has a

function in metabolism such as enzymes involved in carbohydrate degradation, MGE = Predicted

proteins are characteristic of mobile genetic elements such as transposases, toxin-antitoxin

plasmid maintenance systems, prophage associated proteins etc., Hypothetical = Predicted

proteins are hypothetical/putative or a protein with a domain of unknown function (DUF) that

lacks any further functional information, Other = Predicted protein could not be categorized into

one of the other categories

S6 Table. Pairwise comparisons of amino acid sequences of genotype 1 (G1) and genotype 2

(G2) Leukotoxin A consensus sequences against representatives of LktA alleles from Davies *et*

al. 2001

S7 Table. Pairwise comparison of all genotype 2 LktA nucleotide sequences.

S8 Table. Pairwise comparison of genotype 2 LktA amino acid sequences that had a divergent

LktA nucleotide sequence to representative sequences of LktA alleles identified in Davies *et al*.

2001.

S9 Table. Pairwise comparison of all genotype 1 LktA nucleotide sequences.

S10 Table. Pairwise comparison of genotype 1 LktA amino acid sequences that had a divergent

LktA nucleotide sequence to representative sequences of LktA alleles identified in Davies *et al*.

2001.

S11 Table. Summary of CAZyme Analysis Results. Only CAZymes that were predicted to

contain a signal peptide and were annotated by 3 of the utilized annotation tools (dbCAN_sub,

DIAMOND, & HMMER) were included. No CAZymes were predicted in genome

86/GCF_022404935 and so was excluded from this analysis.

S1 Figure. Phylogenetic tree of study genomes created using MASH with the ani command in

PanTools. G1 and G2 clusters are indicated.

References

1. Rice JA, Carrasco-Medina L, Hodgins DC, Shewen PE. *Mannheimia haemolytica* and bovine respiratory disease. Animal Health Research Reviews. 2007;8: 117–128. doi:10.1017/S1466252307001375

2. Highlander SK. Molecular genetic analysis of virulence in *Mannheimia (Pasteurella) haemolytica*. FBL. 2001;6: 1128–1150. doi:10.2741/highland

3. DeDonder KD, Apley MD. A literature review of antimicrobial resistance in pathogens associated with bovine respiratory disease. Animal Health Research Reviews. 2015;16: 125–134. doi:10.1017/S146625231500016X

4. Blakebrough-Hall C, McMeniman JP, González LA. An evaluation of the economic effects of bovine respiratory disease on animal performance, carcass traits, and economic outcomes in feedlot cattle defined using four BRD diagnosis methods. Journal of Animal Science. 2020;98: skaa005. doi:10.1093/jas/skaa005

5. Clawson ML, Murray RW, Sweeney MT, Apley MD, DeDonder KD, Capik SF, et al. Genomic signatures of *Mannheimia haemolytica* that associate with the lungs of cattle with respiratory disease, an integrative conjugative element, and antibiotic resistance genes. BMC Genomics. 2016;17: 982. doi:10.1186/s12864-016-3316-8

6. Biberstein EL, Gills M, Knight H. Serological types of *Pasteurella hemolytica*. Cornell Vet. 1960;50: 283–300.

7.   Al-Ghamdi GM, Ames TR, Baker JC, Walker R, Chase CCL, Frank GH, et al. Serotyping

of *Mannheimia (Pasteurella) haemolytica* Isolates from the Upper Midwest United States. J

VET Diagn Invest. 2000;12: 576–578. doi:10.1177/104063870001200617

8.   Mason C, Errington J, Foster G, Thacker J, Grace O, Baxter-Smith K. *Mannheimia*

haemolytica serovars associated with respiratory disease in cattle in Great Britain. BMC

Veterinary Research. 2022;18: 5. doi:10.1186/s12917-021-03121-3

9.   Klima CL, Alexander TW, Hendrick S, McAllister TA. Characterization of *Mannheimia*

haemolytica isolated from feedlot cattle that were healthy or treated for bovine respiratory

disease. Can J Vet Res. 2014;78: 38–45.

10.   Arnal JL, Fernández A, Vela AI, Sanz C, Fernández-Garyzábal JF, Cid D. Capsular type

diversity of *Mannheimia haemolytica* determined by multiplex real-time PCR and indirect

hemagglutination in clinical isolates from cattle, sheep, and goats in Spain. Veterinary

Microbiology. 2021;258: 109121. doi:10.1016/j.vetmic.2021.109121

11.   Confer AW, Martino P. *Mannheimia haemolytica*. CABI Compendium. 2017;CABI

Compendium: 70912. doi:10.1079/cabicompendium.70912

12.   Klima CL, Zaheer R, Cook SR, Booker CW, Hendrick S, Alexander TW, et al. Pathogens of

Bovine Respiratory Disease in North American Feedlots Conferring Multidrug Resistance

via Integrative Conjugative Elements. Journal of Clinical Microbiology. 2014;52: 438–448.

doi:10.1128/jcm.02485-13

13.   Holman DB, McAllister TA, Topp E, Wright A-DG, Alexander TW. The nasopharyngeal

649   microbiota of feedlot cattle that develop bovine respiratory disease. Vet Microbiol.

650   2015;180: 90–95. doi:10.1016/j.vetmic.2015.07.031

651   14.   Noyes NR, Benedict KM, Gow SP, Booker CW, Hannon SJ, McAllister TA, et al.

652   *Mannheimia haemolytica* in feedlot cattle: prevalence of recovery and associations with

653   antimicrobial use, resistance, and health outcomes. J Vet Intern Med. 2015;29: 705–713.

654   doi:10.1111/jvim.12547

655   15.   Ayalew S, Murdock BK, Snider TA, Confer AW. *Mannheimia haemolytica* IgA-specific

656   proteases. Vet Microbiol. 2019;239: 108487. doi:10.1016/j.vetmic.2019.108487

657   16.   Singh K, Ritchey JW, Confer AW. *Mannheimia haemolytica*: Bacterial–Host Interactions in

658   Bovine Pneumonia. Vet Pathol. 2011;48: 338–348. doi:10.1177/0300985810377182

659   17.   Kisiela DI, Czuprynski CJ. Identification of *Mannheimia haemolytica* Adhesins Involved in

660   Binding to Bovine Bronchial Epithelial Cells. Infection and Immunity. 2009;77: 446–455.

661   doi:10.1128/iai.00312-08

662   18.   Hacker J, Carniel E. Ecological fitness, genomic islands and bacterial pathogenicity. A

663   Darwinian view of the evolution of microbes. EMBO Rep. 2001;2: 376–381.

664   doi:10.1093/embo-reports/kve097

665   19.   Cozens D, Sutherland E, Lauder M, Taylor G, Berry CC, Davies RL. Pathogenic

666   *Mannheimia haemolytica* Invades Differentiated Bovine Airway Epithelial Cells. Infect

667   Immun. 2019;87: 10.1128/iai.00078-19. doi:10.1128/iai.00078-19

20. Loy JD, Clawson ML. Rapid typing of *Mannheimia haemolytica* major genotypes 1 and 2

using MALDI-TOF mass spectrometry. J Microbiol Methods. 2017;136: 30–33.

doi:10.1016/j.mimet.2017.03.002

21. Wynn EL, Schuller G, Loy JD, Workman AM, McDaneld TG, Clawson ML. Differentiation

of *Mannheimia haemolytica* genotype 1 and 2 strains by visible phenotypic characteristics

on solid media. Journal of Microbiological Methods. 2020;171: 105877.

doi:10.1016/j.mimet.2020.105877

22. Clawson ML, Schuller G, Dickey AM, Bono JL, Murray RW, Sweeney MT, et al.

Differences between predicted outer membrane proteins of genotype 1 and 2 *Mannheimia*

haemolytica. BMC Microbiology. 2020;20: 250. doi:10.1186/s12866-020-01932-2

23. Klima CL, Cook SR, Zaheer R, Laing C, Gannon VP, Xu Y, et al. Comparative Genomic

Analysis of *Mannheimia haemolytica* from Bovine Sources. PLOS ONE. 2016;11:

e0149520. doi:10.1371/journal.pone.0149520

24. O'Leary NA, Cox E, Holmes JB, Anderson WR, Falk R, Hem V, et al. Exploring and

retrieving sequence and metadata for species across the tree of life with NCBI Datasets. Sci

Data. 2024;11: 732. doi:10.1038/s41597-024-03571-y

25. Dassanayake RP, Clawson ML, Tatum FM, Briggs RE, Kaplan BS, Casas E. Differential

identification of *Mannheimia haemolytica* genotypes 1 and 2 using colorimetric loop-

mediated isothermal amplification. BMC Research Notes. 2023;16: 4. doi:10.1186/s13104-

023-06272-8

26.   Okonechnikov K, Golosova O, Fursov M, UGENE team. Unipro UGENE: a unified

bioinformatics toolkit. Bioinformatics. 2012;28: 1166–1167.

doi:10.1093/bioinformatics/bts091

27.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3--

new capabilities and interfaces. Nucleic Acids Res. 2012;40: e115. doi:10.1093/nar/gks596

28.   Kõressaar T, Lepamets M, Kaplinski L, Raime K, Andreson R, Remm M. Primer3_masker:

integrating masking of template sequence with primer design software. Bioinformatics.

2018;34: 1937–1938. doi:10.1093/bioinformatics/bty036

29.   Koressaar T, Remm M. Enhancements and modifications of primer design program

Primer3. Bioinformatics. 2007;23: 1289–1291. doi:10.1093/bioinformatics/btm091

30.   Jonkheer EM, van Workum D-JM, Sheikhizadeh Anari S, Brankovics B, de Haan JR, Berke

L, et al. PanTools v3: functional annotation, classification and phylogenomics.

Bioinformatics. 2022;38: 4403–4405. doi:10.1093/bioinformatics/btac506

31.   Sheikhizadeh S, Schranz ME, Akdel M, de Ridder D, Smit S. PanTools: representation,

storage and exploration of pan-genomic data. Bioinformatics. 2016;32: i487–i493.

doi:10.1093/bioinformatics/btw455

32.   Anari SS, Ridder D de, Schranz ME, Smit S. Pangenomic read mapping. bioRxiv; 2019. p.

813634. doi:10.1101/813634

33. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-

scale protein function classification. Bioinformatics. 2014;30: 1236–1240.

doi:10.1093/bioinformatics/btu031

34. Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The

InterPro protein families and domains database: 20 years on. Nucleic Acids Res. 2021;49:

D344–D354. doi:10.1093/nar/gkaa977

35. Nielsen H. Predicting Secretory Proteins with SignalP. Methods Mol Biol. 2017;1611: 59–

73. doi:10.1007/978-1-4939-7015-5_6

36. Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, et al. antiSMASH 7.0:

new and improved predictions for detection, regulation, chemical structures and

visualisation. Nucleic Acids Res. 2023;51: W46–W50. doi:10.1093/nar/gkad344

37. Deschner D, Hill JE. Genomic Differentiation of *Mannheimia haemolytica* genotypes.

FigShare; 2025. doi:10.6084/m9.figshare.28872536

38. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics. 2009;10: 421. doi:10.1186/1471-2105-

10-421

39. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST

and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res.

1997;25: 3389–402.

40. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990;215: 403–10.

41. Zhang Z, Schwartz S, Wagner L, Miller W. A greedy algorithm for aligning DNA
sequences. J Comput Biol. 2000;7: 203–214. doi:10.1089/10665270050081478

42. Morgulis A, Coulouris G, Raytselis Y, Madden TL, Agarwala R, Schäffer AA. Database
indexing for production MegaBLAST searches. Bioinformatics. 2008;24: 1757–1764.
doi:10.1093/bioinformatics/btn322

43. Roehrig SC, Tran HQ, Spehr V, Gunkel N, Selzer PM, Ullrich HJ. The response of
Mannheimia haemolytica to iron limitation: Implications for the acquisition of iron in the
bovine lung. Veterinary Microbiology. 2007;121: 316–329.
doi:10.1016/j.vetmic.2006.12.013

44. Kostova V, Hanke D, Kaspar H, Fiedler S, Schwarz S, Krüger-Haker H. Macrolide
resistance in *Mannheimia haemolytica* isolates associated with bovine respiratory disease
from the German national resistance monitoring program GERM-Vet 2009 to 2020. Front
Microbiol. 2024;15. doi:10.3389/fmicb.2024.1356208

45. Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y. dbCAN3: automated carbohydrate-active
enzyme and substrate annotation. Nucleic Acids Res. 2023;51: W115–W121.
doi:10.1093/nar/gkad328

46. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure
prediction of biomolecular interactions with AlphaFold 3. Nature. 2024;630: 493–500.
doi:10.1038/s41586-024-07487-w

47.   Deschner D, Voordouw MJ, Fernando C, Campbell J, Waldner CL, Hill JE. Identification of

747   genetic markers of resistance to macrolide class antibiotics in *Mannheimia haemolytica*

748   isolates from a Saskatchewan feedlot. Appl Environ Microbiol. 2024;90: e00502-24.

749   doi:10.1128/aem.00502-24

750   48.   Vanni C, Schechter MS, Acinas SG, Barberán A, Buttigieg PL, Casamayor EO, et al.

751   Unifying the known and unknown microbial coding sequence space. Brown CT, Storz G,

752   Brown CT, Smith B, editors. eLife. 2022;11: e67667. doi:10.7554/eLife.67667

753   49.   Lobb B, Tremblay BJ-M, Moreno-Hagelsieb G, Doxey AC. PathFams: statistical detection

754   of pathogen-associated protein domains. BMC Genomics. 2021;22: 663.

755   doi:10.1186/s12864-021-07982-8

756   50.   Messenger AJM, Barclay R. Bacteria, iron and pathogenicity. Biochemical Education.

757   1983;11: 54–63. doi:10.1016/0307-4412(83)90043-2

758   51.   Frawley ER, Fang FC. The ins and outs of bacterial iron metabolism. Molecular

759   Microbiology. 2014;93: 609–616. doi:10.1111/mmi.12709

760   52.   Sheldon JR, Laakso HA, Heinrichs DE. Iron Acquisition Strategies of Bacterial Pathogens.

761   Microbiology Spectrum. 2016;4: 10.1128/microbiolspec.vmbf-0010–2015.

762   doi:10.1128/microbiolspec.vmbf-0010-2015

763   53.   Ferguson AD, Deisenhofer J. TonB-dependent receptors-structural perspectives. Biochim

764   Biophys Acta. 2002;1565: 318–332. doi:10.1016/s0005-2736(02)00578-3

54.   Silale A, Van Den Berg B. TonB-Dependent Transport Across the Bacterial Outer

Membrane. Annu Rev Microbiol. 2023;77: 67–88. doi:10.1146/annurev-micro-032421-

111116

55.   Pogoutse AK, Moraes TF. Iron acquisition through the bacterial transferrin receptor. Crit

Rev Biochem Mol Biol. 2017;52: 314–326. doi:10.1080/10409238.2017.1293606

56.   Moraes TF, Yu R, Strynadka NCJ, Schryvers AB. Insights into the Bacterial Transferrin

Receptor: The Structure of Transferrin-Binding Protein B from *Actinobacillus*

pleuropneumoniae. Molecular Cell. 2009;35: 523–533. doi:10.1016/j.molcel.2009.06.029

57.   Ayalew S, Shrestha B, Montelongo M, Wilson AE, Confer AW. Identification and

immunogenicity of *Mannheimia haemolytica* S1 outer membrane lipoprotein PlpF. Vaccine.

2011;29: 8712–8718. doi:10.1016/j.vaccine.2011.08.074

58.   Ayalew S, Confer AW, Blackwood ER. Characterization of immunodominant and

potentially protective epitopes of *Mannheimia haemolytica* serotype 1 outer membrane

lipoprotein PlpE. Infect Immun. 2004;72: 7265–7274. doi:10.1128/IAI.72.12.7265-

7274.2004

59.   Confer AW, Ayalew S, Panciera RJ, Montelongo M, Wray JH. Recombinant *Mannheimia*

haemolytica serotype 1 outer membrane protein PlpE enhances commercial *M. haemolytica*

vaccine-induced resistance against serotype 6 challenge. Vaccine. 2006;24: 2248–2255.

doi:10.1016/j.vaccine.2005.11.036

60. Pandher K, Confer AW, Murphy GL. Genetic and immunologic analyses of PlpE, a

lipoprotein important in complement-mediated killing of *Pasteurella haemolytica* serotype

1. Infect Immun. 1998;66: 5613–5619. doi:10.1128/IAI.66.12.5613-5619.1998

61. Ayalew S, Blackwood ER, Confer AW. Sequence diversity of the immunogenic outer

membrane lipoprotein PlpE from *Mannheimia haemolytica* serotypes 1, 2, and 6. Vet

Microbiol. 2006;114: 260–268. doi:10.1016/j.vetmic.2005.11.067

62. Oppermann T, Busse N, Czermak P. *Mannheimia haemolytica* growth and leukotoxin

production for vaccine manufacturing — A bioprocess review. Electronic Journal of

Biotechnology. 2017;28: 95–100. doi:10.1016/j.ejbt.2017.06.001

63. Davies RL, Whittam TS, Selander RK. Sequence Diversity and Molecular Evolution of the

Leukotoxin (lktA) Gene in Bovine and Ovine Strains of *Mannheimia (Pasteurella)*

haemolytica. J Bacteriol. 2001;183: 1394–1404. doi:10.1128/jb.183.4.1394-1404.2001

64. Narayanan SK, Nagaraja TG, Chengappa MM, Stewart GC. Leukotoxins of gram-negative

bacteria. Vet Microbiol. 2002;84: 337–356. doi:10.1016/s0378-1135(01)00467-9

65. Larsen J, Pedersen AG, Davies RL, Kuhnert P, Frey J, Christensen H, et al. Evolution of the

leukotoxin promoter in genus *Mannheimia*. BMC Evolutionary Biology. 2009;9: 121.

doi:10.1186/1471-2148-9-121

66. Christensen H, Kuhnert P, Foster G, Bisgaard M. *Mannheimia indoligenes* sp. nov.,

proposed for clade V organisms of *Mannheimia*. International Journal of Systematic and

Evolutionary Microbiology. 2024;74: 006370. doi:10.1099/ijsem.0.006370

67. Drula E, Garron M-L, Dogan S, Lombard V, Henrissat B, Terrapon N. The carbohydrate-

active enzyme database: functions and literature. Nucleic Acids Res. 2022;50: D571–D577.

doi:10.1093/nar/gkab1045

68. Dik DA, Marous DR, Fisher JF, Mobashery S. Lytic transglycosylases: concinnity in

concision of the bacterial cell wall. Crit Rev Biochem Mol Biol. 2017;52: 503–542.

doi:10.1080/10409238.2017.1337705

69. Martinez-Bond EA, Soriano BM, Williams AH. The mechanistic landscape of Lytic

transglycosylase as targets for antibacterial therapy. Curr Opin Struct Biol. 2022;77:

102480. doi:10.1016/j.sbi.2022.102480

70. Suvorov M, Lee M, Hesek D, Boggess B, Mobashery S. Lytic Transglycosylase MltB of

Escherichia coli and Its Role in Recycling of Peptidoglycan Strands of Bacterial Cell Wall.

J Am Chem Soc. 2008;130: 11878–11879. doi:10.1021/ja805482b

71. Scheurwater E, Reid CW, Clarke AJ. Lytic transglycosylases: bacterial space-making

autolysins. Int J Biochem Cell Biol. 2008;40: 586–591. doi:10.1016/j.biocel.2007.03.018