Estimating the additive genetic variance for relative fitness from changes in allele frequency

Manas Geeta Arun[*1], Aidan Angus-Henry[1,2], Darren J. Obbard[1], and Jarrod D. Hadfield[1]

[1]Institute of Ecology and Evolution, The University of Edinburgh, Ashworth Laboratories Charlotte Auerbach Road, Edinburgh, EH9 3FL, United Kingdom.
[2]Charité - Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.

Running head

Estimating heritable variance in fitness

Keywords

Fisher's Fundamental Theorem of Natural Selection, rate of adaptation, quantitative genetics, evolve and resequence.

*Corresponding author: manas.geetaarun@ed.ac.uk

Abstract

The rate of adaptation is equal to the additive genetic variance for relative fitness (V_A) in the population. Estimating V_A typically involves obtaining suitable measures of fitness on a large number of individuals with known pairwise relatedness. Such data are hard to collect and the results are often sensitive to the definition of fitness used. Here, we present a new method for estimating V_A that does not involve making measurements of fitness on individuals, but instead tracks changes in the genetic composition of the population. First, we show that V_A can readily be expressed as a function of the genome-wide diversity/linkage disequilibrium matrix and genome-wide expected change in allele frequency due to selection. We then show how independent experimental replicates can be used to infer the expected change in allele frequency due to selection and then estimate V_A via a linear mixed model. Finally, using individual-based simulations, we demonstrate that our approach yields precise and accurate estimates over a range of biologically plausible scenarios.

Article summary

Conventional approaches for estimating the heritable component of fitness variation (V_A) have steep methodological, statistical, and even definitional challenges. Here, the authors present a new method that overcomes many of these issue by modelling V_A using selection-induced changes to a population's genetic composition. The authors develop novel mathematical theory and an inference approach that uses independent experimental populations derived from the same ancestral population. Individual based simulations show that this method provides unbiased and precise estimates of V_A. This opens the door for future studies investigating the genomic distribution of V_A, a key factor driving Darwinian evolution.

Introduction

Despite its simplicity, the fundamental theorem of natural selection (FTNS) (Fisher, 1930, 1958) is arguably one of the most central results in evolutionary biology, providing a concise mathematical statement of how quickly a population is expected to adapt. It describes the per generation gain in the mean fitness of a population as a result of natural selection, assuming the 'environment' (including variables intrinsic to the population, such as density or allele frequencies) are held constant (Ewens, 1989; Frank and Slatkin, 1992). The crucial insight the FTNS provides is that the

rate of this increase in mean fitness is exactly equal to the additive genetic variance for relative fitness (V_A) in the population (Burt, 1995; Grafen, 2015). Consequently, estimating V_A is one of the '*holy grails*' of evolutionary genetics (Walsh, 2022), despite the FTNS being criticised for its inability to predict the actual gain in mean fitness (Price, 1972; Ewens, 1989, 2024) (also see Edwards (1994) for an historical review of the debate around FTNS).

A number of attempts have been made to measure V_A in wild populations. Typically, these have involved long term studies on natural populations in which the lifetime reproductive success of a large number of individuals has been measured. Combining these fitness data with information on the relatedness among individuals in a (generalised) linear mixed model approach yields estimates of V_A (Kruuk, 2004). This is far from straightforward in natural populations, as it can be notoriously difficult to tease apart additive genetic effects from common environmental effects, such as parental effects (Kruuk and Hadfield, 2007; Shaw and Shaw, 2014). In addition, wild study systems are rarely closed, meaning emigration can be misinterpreted as mortality and offspring sired outside the study area can be overlooked. Furthermore, many studies on wild populations lack genetic pedigrees, and as a consequence may miss substantial fitness variation acquired through undetected polygamy (Vedder *et al.*, 2011; Charmantier and Sheldon, 2006). In addition to these biases, estimates from wild populations also tend to come with considerable uncertainty. Burt (1995) reviewed studies estimating V_A in three species of plants and three species of animals, and found that most estimates of V_A were not significantly different from zero. They argued that the upper bound for estimates of V_A could be as high as 0.3, but most likely less than 0.1. Consistent with this, and using a larger data set, Hendry *et al.* (2018) reported that estimates of V_A varied between zero and 0.85, with the vast majority of estimates (73%) being less than 0.2. Overall, the mean $\widehat{V_A}$ across studies was 0.08. In a recent meta-analysis, Bonnet *et al.* (2022) applied Bayesian quantitative genetic methods to data obtained from 19 long term studies on wild vertebrate populations. They reported a meta-analytic mean V_A of 0.185 across studies, considerably higher than those obtained by Burt (1995) and Hendry *et al.* (2018). Surprisingly, estimates of V_A in populations of Spotted Hyenas (*Crocuta crocuta*), as well as two of the three populations of Blue Tits (*Cyanistes caeruleus*) were higher than 0.4. This is a remarkable finding, since it suggests that growth rates in these populations should increase nearly 1.5 fold every generation due to selection, provided the environment remains constant. All three meta-analyses investigating V_A (Burt, 1995; Hendry *et al.*, 2018; Bonnet *et al.*, 2022) have detected substantial variability between study systems in their estimates of V_A. Although most of this variability will be sampling error, the best estimate suggests the variability in actual

V_A across populations is also large with a standard deviation of 0.11, although there is substantial uncertainty about its exact value (95% credible intervals: 0.01–0.26) (Bonnet *et al.*, 2022).

Measuring V_A in the laboratory is considerably more straightforward, and involves either quantitative genetic breeding designs such as diallel crosses and full-sib half-sib experiments (Falconer and Mackay, 1996; Lynch and Walsh, 1998), or experimental techniques such as hemiclonal analysis (Abbott and Morrow, 2011). While the controlled environment of the laboratory can, to a large extent, help overcome some of the challenges faced by field studies, laboratory environments often lack important features that are likely to generate fitness variation, such as parasites, predators, and competitors. Therefore, it is not entirely clear if laboratory estimates of V_A are particularly relevant. Furthermore, many laboratory studies standardise their fitness measurements in a way that makes it hard to infer estimates of V_A (eg. Ruzicka *et al.* (2019)) or work with absolute fitness but then fail to report V_A (eg. Singh *et al.* (2023)). However, these studies do suggest that genetic variance for fitness is likely highly dependent on the specific environment in which fitness is assayed (eg. Punzalan *et al.* (2014)). In one of the few laboratory studies that estimated the genetic variance for *relative* fitness, Martinossi-Allibert *et al.* (2018) reported estimates from isofemale lines of the bean beetle, *Acanthoscelides obtectus*, that were highly sensitive to evolutionary history, assay environment, and sex, being higher in males (0.13-0.42) than in females (0.013-0.056). A compromise between the precision of the laboratory environment and the biotic and abiotic complexity encountered by wild populations can be achieved by working with experimental populations established in the field, an approach especially tractable in annual plants. Working with field populations of the annual legume *Chamaecrista fasciculata*, Kulbaba *et al.* (2019) found estimates of V_A that varied considerably among populations and years. Interestingly, many of their estimates were appreciably larger, with the largest estimate (calculated from Table 1 in the correction) being 3.05.

A common difficulty for current field and laboratory approaches is that, while Darwinian fitness is a deceptively intuitive concept, there is little consensus on its precise definition. In fact, it has been argued that the appropriate definition of fitness can vary depending on the context (Hendry *et al.*, 2018). In the absence of a universal definition, empiricists can only use a measure of fitness deemed most appropriate for their study system, and it is reasonable to assume that estimates of V_A are likely to be sensitive to the definition of fitness used. A useful illustration of this point is provided by two studies that estimated V_A in a wild population of Red Deer (*Cervus elaphus*), using largely overlapping datasets, but markedly different definitions of fitness (Kruuk *et al.*, 2000; Foerster *et al.*, 2007). Kruuk *et al.* (2000)

defined fitness as the total number of progeny produced by an individual in its lifetime and estimated V_A to be 0.1, whereas Foerster *et al.* (2007) employed an alternative definition of fitness that measured an individual's contribution to population growth (Coulson *et al.*, 2006) and obtained the appreciably higher estimate of 0.64.

Some of the definitional difficulties of measuring V_A can be overcome by measuring V_A as the rate of adaptation, rather than comparing measures of fitness among relatives. One of the earliest attempts to explore this idea was made by Fowler *et al.* (1997) (also see Gardner *et al.* (2005)) in laboratory populations of *Drosophila melanogaster*. Using balancer chromosomes that allow recombination to be suppressed on the third chromosome, Fowler *et al.* (1997) could track the frequency trajectories of newly introduced wild-type third chromosomes over the course of 43 weeks. By modelling fitness using genotype frequency data for a number of different wild-type third chromosomes (Barton and Partridge, 2000), they demonstrated the presence of substantial V_A on the third chromosome. More recently, in a landmark study, Buffalo and Coop (2019) developed a method to estimate the amount of genome-wide allele frequency change that can be attributed to selection. The linchpin of their theory is the idea that linked selection induces across-generation covariances in allele frequency change at neutral loci, since associations between these neutral loci and their respective non-neutral backgrounds persist across generations. This new theoretical framework has the potential to pave the way for a powerful empirical tool to detect genomic signatures of linked selection (Buffalo and Coop, 2020; Simon and Coop, 2024). Of particular relevance here, Buffalo and Coop (2019) also show that their method can be used to obtain estimates of V_A, albeit under some potentially restrictive assumptions.

In this study, we present an alternative theoretical framework that relates V_A to genome-wide changes in allele frequency. Using mathematical identities only, we show how V_A can be obtained from an initial linkage disequilibrium (LD) matrix and expected changes in allele frequency due to selection, without making any assumptions about patterns of gene action or the relationships between genotype fitnesses and genotype frequencies. Our approach, like that of Buffalo and Coop (2019), relies on temporal genomic data and does not necessitate measuring fitness in individuals. However, in contrast to the 'bottom-up' population genetic approach of Buffalo and Coop (2019), we use a 'top-down' quantitative genetic approach which simplifies and generalises some aspects of the problem.

Our aim here is three-fold. First, we derive our central theoretical result from first principles. Second, we develop the statistical machinery required to apply our result to real biological data, and validate it with individual based simulations that permit dominance effects, but assume random mating and an absence of epistasis.

Third, we make a detailed comparison of our method with that of Buffalo and Coop (2019).

Materials and methods

Table 3 summarises the notation used in the following text and the subsection 'Assumptions retained in our approach' (see below) summarises the assumptions involved in our approach.

Outline of the theory

We consider a population consisting of N diploid individuals. We assume that there are n_L segregating biallelic loci in the population. Let $c_{k,i}$ and α_i represent the proportion of copies of an arbitrarily chosen reference allele at locus i in individual k and Fisher's average effect (Fisher, 1941) for relative fitness at locus i, respectively. The widely accepted mathematical definition of the α's are the regression coefficients obtained from a multiple regression of the c's on relative fitness, w (Fisher, 1941; Lee and Chow, 2013). The vector $\boldsymbol{\alpha}$ can be expressed as

$$\boldsymbol{\alpha} \ = \mathbf{L}^{-1} COV(\mathbf{c}, w) \tag{1}$$

where $\mathbf{c}$ is the vector of predictors representing the c's at all loci for an individual and $\mathbf{L}$ is a symmetric $n_L \times n_L$ matrix whose i^{th} (diagonal) element is the variance in the c's at locus i computed over individuals, while the ij^{th} (off-diagonal) element describes the covariance between the c's at locus i and locus j computed over individuals.

The breeding value for the relative fitness of individual k is then

$$u_k = \mathbf{c}_k^\top \boldsymbol{\alpha}$$

and the additive genetic variance for relative fitness is the variance of this quantity across individuals:

$$\begin{aligned} V_A \ &= VAR(\mathbf{c}^\top \boldsymbol{\alpha}) \\ &= \boldsymbol{\alpha}^\top VAR(\mathbf{c}, \mathbf{c}^\top) \boldsymbol{\alpha} \\ &= \boldsymbol{\alpha}^\top \mathbf{L} \boldsymbol{\alpha} \end{aligned} \tag{2}$$

This follows from the fact that at any given point in time, α's are constant across individuals. In the absence of mutation and meiotic drive, the allele frequency in parents is transmitted to offspring without bias, such that the vector of expected

change in allele frequencies due to selection can be expressed as Robertson's covariance (Robertson, 1966; Price, 1970; Queller, 2017):

$$
\begin{aligned}
E[\Delta\mathbf{p}] &= E[\Delta\bar{\mathbf{c}}] \\
&= COV(\mathbf{c}, w)
\end{aligned}
\tag{3}
$$

where the expectation is taken over the evolutionary process.

Substituting Equation 1 into Equation 3 gives $E[\Delta\mathbf{p}] = \mathbf{L}\boldsymbol{\alpha}$, which is the multivariate analogue of Equation 10 in Kirkpatrick *et al.* (2002). Combining this with Equation 2 yields,

$$
\begin{aligned}
V_A &= \left(\mathbf{L}^{-1}E[\Delta\mathbf{p}]\right)^{\top}\mathbf{L}\left(\mathbf{L}^{-1}E[\Delta\mathbf{p}]\right) \\
&= E[\Delta\mathbf{p}]^{\top}\mathbf{L}^{-1}E[\Delta\mathbf{p}]
\end{aligned}
\tag{4}
$$

Equation 4 is a general result and involves no assumptions about the patterns of dominance or epistasis for fitness, or about patterns of mating. An intuitive explanation of why V_A can be calculated this way is to note Fisher's Fundamental Theorem states that V_A is equal to the (partial) increase in mean fitness caused by evolutionary change through natural selection. Equation 2 can be expressed as a sum of $\alpha_i E[\Delta p_i]$ over all loci, i:

$$
V_A = \boldsymbol{\alpha}^{\top}E[\Delta\mathbf{p}]
\tag{5}
$$

In this, α represents the proportional change in the population mean fitness that is caused by a unit change in allele frequency at a locus (Fisher, 1941; Kojima, 1959; Lee and Chow, 2013). Therefore, multiplying α by the actual change caused by natural selection, $E[\Delta p]$, we get the proportional change in mean fitness caused by evolutionary change by natural selection at that locus (see Eq 51.4 Price, 1972, also). If we then add these changes at every locus in the genome, we obtain the total proportional change in mean fitness due to evolutionary change by natural selection, and hence V_A.

Extending our approach to practical situations

Our theoretical approach assumes $\boldsymbol{\alpha}$ or $E[\Delta\mathbf{p}]$ are known. In reality, neither can be directly observed and must be inferred from data on observed allele frequency change, $\Delta\mathbf{p}$. Since $\Delta\mathbf{p}$ will vary around $E[\Delta\mathbf{p}]$ due to genetic drift, $E[\Delta\mathbf{p}]$ must be inferred using replicate observations of $\Delta\mathbf{p}$. Since time cannot be replayed, we infer $E[\Delta\mathbf{p}]$ through experimental replicates (see Buffalo and Coop (2020) also). Our theoretical model also assumes $\mathbf{L}$ is known, but since it is hard to generate experimental replicates without at least one round of reproduction, we condition on $\mathbf{L}_0$

₂₁₃ (**L** in a generation prior to the first generation over which allele frequency change is
₂₁₄ measured). In what follows we will refer to the population at time zero (generation
₂₁₅ 0) as the 'base population'. Our aim is then to approximate the additive genetic
₂₁₆ variance for fitness in the base population $V_A(0)$ as $V_{\bar{A}}(0) = \bar{\boldsymbol{\alpha}}^\top \mathbf{L}_0 \bar{\boldsymbol{\alpha}}$, where $\bar{\boldsymbol{\alpha}}$ is
₂₁₇ the mean vector of average effects averaged over time and replicates. Note that if
₂₁₈ average effects are constant then $V_{\bar{A}}(0) = V_A(0)$, but if the average effects vary then
₂₁₉ $V_A(0)$ will in general exceed $V_{\bar{A}}(0)$ which can be interpreted as the additive genetic
₂₂₀ *covariance* in fitness between replicates/time points for a population with genotypic
₂₂₁ structure equal to that in the base population (see Supplementary information S2).
₂₂₂ It is important to note that conventional methods for estimating V_A – such as those
₂₂₃ relying on pedigreed individuals – also estimate additive genetic *covariances* among
₂₂₄ environments in which relatives are assayed (Vehviläinen *et al.*, 2008) and these co-
₂₂₅ variances equal V_A only when the average effects remain constant.

₂₂₇   We also allow allele frequency changes to be measured over multiple generations,
₂₂₈ rather than a single generation. Thus, $\Delta \mathbf{p}_m = \Delta \mathbf{p}_{t_m,m} + \Delta \mathbf{p}_{t_m+1,m} \ldots \Delta \mathbf{p}_{\tau_m-1,m}$ is
₂₂₉ the observed change in allele frequency from generation t_m to τ_m in replicate m, with
₂₃₀ $\Delta \mathbf{p}_{t,m}$ being the change from time t to $t+1$. Note that t_m (i.e. the generation in
₂₃₁ which we start recording allele frequency changes in replicate m) can be any positive
₂₃₂ integer. If replicate populations are derived from the base population with exactly
₂₃₃ one round of reproduction, t_m would be 1 for all m. τ_m can be any integer greater
₂₃₄ than t_m. We first note that the total change in allele frequency in replicate m between
₂₃₅ times t_m and τ_m is

$$\Delta \mathbf{p}_m = \sum_{t=t_m}^{\tau_m-1} \left(\mathbf{L}_{t,m} \boldsymbol{\alpha}_{t,m} + \underset{D}{\Delta} \mathbf{p}_{t,m} \right) \tag{6}$$

₂₃₆   where $\mathbf{L}_{t,m} \boldsymbol{\alpha}_{t,m} = E[\Delta \mathbf{p}_{t,m}]$ is the expected change in allele frequency due to
₂₃₇ selection in replicate m between generation t and $t+1$ and $\underset{D}{\Delta} \mathbf{p}_{t,m}$ is the change due
₂₃₈ to drift. It is important to note that $\mathbf{L}_{t,m} \boldsymbol{\alpha}_{t,m}$ captures responses to both direct
₂₃₉ and indirect selection. **L** will vary over time and we decompose **L** at a particular
₂₄₀ time/replicate into a part that can be predicted by $\mathbf{L}_0$ and the action of drift and
₂₄₁ recombination, and a part that cannot be predicted. To do this we decompose **L**
₂₄₂ into $\mathbf{L}'$ and $\mathbf{L}''$, following the notation of Buffalo and Coop (2019). $\mathbf{L}'$ represents
₂₄₃ the (co)variances in the c's computed over the haploid genomes of all $2N$ gametic
₂₄₄ contributions that constitute the population. Thus, the diagonal elements of $\mathbf{L}'$ are
₂₄₅ proportional to gametic (gene) diversity and the off-diagonals are proportional to
₂₄₆ gametic-phase disequilibrium. On the other hand, $\mathbf{L}''$ represents the (co)variances

₂₄₇ that arise due to alleles within the different gametic contributions of a genotype, and
₂₄₈ thus the diagonal elements are proportional to the additive coefficients of Hardy-
₂₄₉ Weinberg disequilibrium (Bulmer (1980) and Chapter 12 in Weir and Cockerham
₂₅₀ (1989)), and the off-diagonal elements are proportional to the nongametic-phase
₂₅₁ disequilibrium. Given this decomposition, the dynamics of $\mathbf{L}$ under drift and recom-
₂₅₂ bination are (See Supplementary information S1 for details and Hill and Robertson
₂₅₃ (1968) and Santiago and Caballero (1998)):

$$\mathbf{L}_{t,m} = \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0 + \Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m} \tag{7}$$

₂₅₄ where $\circ$ is the Hadamard (i.e. element-wise) product, $\tilde{\mathbf{L}}_0$ is the weighted sum of $\mathbf{L}'_0$
₂₅₅ and $\mathbf{L}''_0$, with the ij^{th} element of $\mathbf{L}''_0$ weighted by $r_{ij}/(1-r_{ij})$ and $\mathbf{N}_{t,m}$ is a matrix with
₂₅₆ the ij^{th} element being $(1-r_{ij})^t \prod_{k=0}^{t-1}(1-\frac{1}{2N_{e_{k,m}}})$. r_{ij} is the recombination rate between
₂₅₇ the two loci and $N_{e_{k,m}}$ is the effective population size in generation k in replicate m.
₂₅₈ $\Delta\mathbf{L}'_{t,m}$ is a stochastic term that represents the accumulated change in $\mathbf{L}'$ between
₂₅₉ generations 0 and t in replicate m that cannot be predicted. $\mathbf{L}''_{t,m}$ is the matrix
₂₆₀ of nongametic-phase disequilbria that arises in generation t in replicate m. In the
₂₆₁ absence of selection, $\Delta\mathbf{L}'_{t,m}$ and $\mathbf{L}''_{t,m}$ have zero expectation. In the following, it will
₂₆₂ be useful to designate $\boldsymbol{\mathcal{L}}_{t,m} = \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0$ as the predicted $\mathbf{L}_{t,m}$, for $t > 0$, conditional
₂₆₃ on $\mathbf{L}'_0$ and $\mathbf{L}''_0$. Also, we represent the deviation of $\mathbf{L}_{t,m}$ from this prediction as
₂₆₄ $\Delta\mathbf{L}_{t,m} = \Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m}$. It will also be useful to denote $\boldsymbol{\alpha}_{t,m} = \bar{\boldsymbol{\alpha}} + \Delta\boldsymbol{\alpha}_{t,m}$ where
₂₆₅ $\Delta\boldsymbol{\alpha}_{t,m}$ is the deviation of the average effects (in generation t in replicate m) from
₂₆₆ the mean of the average effects across time and replicates. We can then decompose
₂₆₇ the change due to selection into two terms:

$$
\begin{aligned}
\Delta\mathbf{p}_m =\ & \sum_{t=t_m}^{\tau_m-1}\left(\boldsymbol{\mathcal{L}}_{t,m}\bar{\boldsymbol{\alpha}} + \Delta\mathbf{L}_{t,m}\bar{\boldsymbol{\alpha}} + \boldsymbol{\mathcal{L}}_{t,m}\Delta\boldsymbol{\alpha}_{t,m} + \Delta\mathbf{L}_{t,m}\Delta\boldsymbol{\alpha}_{t,m} + \underset{D}{\Delta}\mathbf{p}_{t,m}\right) \\
=\ & \sum_{t=t_m}^{\tau_m-1}\left(\boldsymbol{\mathcal{L}}_{t,m}\bar{\boldsymbol{\alpha}} + \underset{U}{\Delta}\mathbf{p}_{t,m} + \underset{D}{\Delta}\mathbf{p}_{t,m}\right) \\
=\ & \boldsymbol{\mathcal{L}}_m\bar{\boldsymbol{\alpha}} + \underset{U}{\Delta}\mathbf{p}_m + \underset{D}{\Delta}\mathbf{p}_m
\end{aligned}
\tag{8}
$$

₂₆₈ Here quantities subscripted with just m are the sums of the relevant quantity over
₂₆₉ t (for example, $\boldsymbol{\mathcal{L}}_m = \sum_{t=t_m}^{\tau_m-1}\boldsymbol{\mathcal{L}}_{t,m}$). The predictable change due to selection is
₂₇₀ $\boldsymbol{\mathcal{L}}_m\bar{\boldsymbol{\alpha}}$ and is similar to that derived for a single locus in Santiago and Caballero
₂₇₁ (1998). The unpredictable change due to selection caused by stochastic changes
₂₇₂ in $\mathbf{L}$ and/or $\boldsymbol{\alpha}$ is $\underset{U}{\Delta}\mathbf{p}_m = \sum_{t=t_m}^{\tau_m-1}(\Delta\mathbf{L}_{t,m}\bar{\boldsymbol{\alpha}} + \boldsymbol{\mathcal{L}}_{t,m}\Delta\boldsymbol{\alpha}_{t,m} + \Delta\mathbf{L}_{t,m}\Delta\boldsymbol{\alpha}_{t,m})$. The total
₂₇₃ change due to drift is $\underset{D}{\Delta}\mathbf{p}_m$. The $\underset{D}{\Delta}\mathbf{p}_m$ have zero expectation and are independent
₂₇₄ across replicates (Buffalo and Coop, 2019). In what follows we also assume the $\underset{U}{\Delta}\mathbf{p}_m$
₂₇₅ have zero expectation, although we believe there are two primary mechanisms by

which this assumption can fail. First, the model for the predicted changes in $\mathbf{L}$ (i.e.
$\boldsymbol{\mathcal{L}}_m$) may be inaccurate such that the $\Delta \mathbf{L}_{t,m}$ have non-zero expectation. Since $\boldsymbol{\mathcal{L}}_m$ is
derived under the assumptions of drift and recombination only, directional changes in
$\mathbf{L}$ induced by selection is an obvious mechanism. However, if the genetic architecture
of fitness is sufficiently polygenic, such that selection coefficients associated with
individual loci are small, selection-induced changes in $\mathbf{L}$ may be safely ignored. In
other words, selection-induced changes in $\mathbf{L}$ may cause the total allele frequency
change due to selection to be negligibly different from $\boldsymbol{\mathcal{L}}_m \bar{\boldsymbol{\alpha}}$, the expected allele
frequency change due to selection. Second, the product $\Delta \mathbf{L}_{t,m} \Delta \boldsymbol{\alpha}_{t,m}$ may have non-
zero expectation if changes in the average effects are correlated with changes in
patterns of diversity/LD. Again, selection-induced changes in $\Delta \mathbf{L}_{t,m}$ is a possible
mechanism. Additionally, unless all gene action is additive, changes in $\mathbf{L}$ also induce
changes in $\boldsymbol{\alpha}$ since then average effects depend on allele frequencies (Supplementary
information S8).

However, calculating change in allele frequency over a single generation (i.e. $\tau_m -$
$t_m = 1$) would minimise any bias since any change in $\mathbf{L}$ due to selection should be
minimised and the approximations that follow will be more accurate. Although the
approximations will hold better when $\tau_m - t_m = 1$, increasing $\tau_m - t_m$ (i.e. calculating
change in allele frequency over multiple generations) will increase power since the
changes in allele frequency due to selection will be larger.

Assuming both $\underset{U}{\Delta}\mathbf{p}_m$ and $\underset{D}{\Delta}\mathbf{p}_m$ have zero expectation we can derive the mean
and covariance structure (computed over the evolutionary process) of allele frequency
change in a replicate as:

$$E\left[\Delta \mathbf{p}_m\right] = \boldsymbol{\mathcal{L}}_m \bar{\boldsymbol{\alpha}} \tag{9}$$

$$VAR\left(\Delta \mathbf{p}_m\right) = \boldsymbol{\mathcal{D}}_m + \boldsymbol{\mathcal{U}}_m \tag{10}$$

where $\boldsymbol{\mathcal{D}}_m$ and $\boldsymbol{\mathcal{U}}_m$ are the covariances due to the cumulative action of drift
and the unpredictable response to selection, respectively. The drift (co)variances
have expectation equal to $\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)}$ where $\mathbf{M}^{(m)} = \sum_{t=t_m}^{\tau_m - 1} \mathbf{M}_{t,m} \circ \mathbf{N}_{t,m}$ and the
ij^{th} element of $\mathbf{M}_{t,m}$ is $(1 - r_{ij})/N_{E_{t,m}}$ (Supplementary information S2). Note that
since the predictable response to selection includes both direct and indirect selection,
the relevant effective population size for the drift covariance in allele frequency does
not include the effects of linked selection and for this reason we denote it as N_E
rather than N_e. Using this notation, $N_E = 4N/(2 + V_o)$ where N is the census
population size and V_o is the variance in offspring number in the absence of additive
genetic fitness variation (Wright, 1938). Although the dynamics of $\mathbf{L}$ (Equation 7,

also see Supplementary information S1) were derived under drift and recombination
in the absence of selection, and hence $N_e = N_E$, the dynamical equations for $\mathbf{L}$ will
better approximate reality when an effective population size that incorporates all
excess variation in fitness is used. Consequently, we use N_e and N_E to distinguish
the effective population sizes that are relevant for stochastic changes in $\mathbf{L}$ and allele
frequency, respectively.

There is no easy form for the covariance due to the cumulative unpredictable re-
sponse to selection and we simply denote it as $\boldsymbol{\mathcal{U}}_m$ (Supplementary information S2).
While the drift terms will be independent across replicates, we also need to assume
that $\underset{U}{\Delta}\mathbf{p}_m$ are independent across replicates for the covariance between replicates to
be zero. An obvious source of non-independence would be if replicates are not ini-
tiated from individuals independently generated from Generation 0 individuals (i.e.
$t_m \neq 1$ for at least one m). If this were the case, the $\Delta\mathbf{L}$ will be dependent across
replicates due to the shared changes in $\mathbf{L}$ from generation 0 to the generation from
which the replicates are derived. This could be minimised by initialising replicates
as early as possible (ideally at $t_m = 1$).

Inference outline

In the previous section, the mean average effects, $\bar{\boldsymbol{\alpha}}$, are treated as fixed and expecta-
tions and variances are taken over the evolutionary process, including variation in the
average effects over time and replicates. When making inferences, we treat the mean
average effects as random variables rather than fixed quantities (See Supplementary
information S3 and Gianola *et al.* (2009) for a discussion on what this implies) and
derive expectations and (co)variances taken over both the evolutionary process and
the distribution of $\bar{\boldsymbol{\alpha}}$. Using the laws of total expectation and (co)variance we obtain:

$$E\left[\Delta\mathbf{p}_m\right] = \boldsymbol{\mathcal{L}}_m\boldsymbol{\mu}_{\bar{\alpha}} \tag{11}$$

334   and

$$VAR\left(\Delta\mathbf{p}_m\right) = \boldsymbol{\mathcal{L}}_m\mathbf{V}_{\bar{\alpha}}\boldsymbol{\mathcal{L}}_m + \boldsymbol{\mathcal{D}}_m + \boldsymbol{\mathcal{U}}_m \tag{12}$$

335 where $\boldsymbol{\mu}_{\bar{\alpha}}$ and $\mathbf{V}_{\bar{\alpha}}$ and are the mean and covariance structure of the mean average
336 effects respectively. Critically, when deconditioning on $\bar{\boldsymbol{\alpha}}$ the cross-replicate covari-
337 ances become non-zero:

$$COV(\Delta\mathbf{p}_m, \Delta\mathbf{p}_n^{\top}) = \boldsymbol{\mathcal{L}}_m\mathbf{V}_{\bar{\alpha}}\boldsymbol{\mathcal{L}}_n \tag{13}$$

³³⁸ where m and n are a pair of replicates.

³⁴⁰ In Supplementary information S3 we determine permissible models for $\boldsymbol{\mu}_{\bar{\alpha}}^{\top}$ and
³⁴¹ $\mathbf{V}_{\bar{\alpha}}$. Of those, we identify the sensible biological model for the mean average effect:

$$\boldsymbol{\mu}_{\bar{\alpha}} = \beta_{\bar{\alpha}}^{(1)}(\mathbf{p}_0 - \mathbf{q}_0) \tag{14}$$

³⁴² where $\mathbf{p}_0$ is the frequency of the reference alleles at each locus in the base population
³⁴³ and $\mathbf{q}_0 = \mathbf{1} - \mathbf{p_0}$. For the variance of the average effects, we identify the model

$$\mathbf{V}_{\bar{\alpha}} = \sigma_{\bar{\alpha}}^2 \mathbf{L}_0^{p_{\bar{\alpha}}} \tag{15}$$

³⁴⁴ although in our analysis of simulated data we set the off-diagonal elements of $\mathbf{L}_0$ to
³⁴⁵ zero in the above equation, such that the variance in average effects is simply a power
³⁴⁶ function of the genetic diversities (Zeng *et al.*, 2018). When $p_{\bar{\alpha}} = 0$ average effects
³⁴⁷ are independent of genetic diversity, and when $p_{\bar{\alpha}} = -1$ average effects scale inversely
³⁴⁸ with genetic diversity. $p_{\bar{\alpha}} = -1$ is a common assumption in many related approaches
³⁴⁹ (e.g. Yang *et al.*, 2011) and under this assumption $\sigma_{\bar{\alpha}}^2$ is the average contribution of
³⁵⁰ a locus to the additive genic variance.

³⁵¹ By applying sum of squares theory (Searle, 2006, page 355) to Equation 2 we can
³⁵² obtain the (posterior) expectation of $V_{\bar{A}}(0)$ after averaging over the distribution of
³⁵³ average effects (Supplementary information S3 and Gianola *et al.* (2009)):

$$\begin{aligned} E[V_{\bar{A}}(0)] &= E[\bar{\boldsymbol{\alpha}}^{\top}\mathbf{L}_0\bar{\boldsymbol{\alpha}}] \\ &= Tr(\mathbf{L}_0\mathbf{V}_{\bar{\alpha}}) + \boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0\boldsymbol{\mu}_{\bar{\alpha}} \end{aligned} \tag{16}$$

³⁵⁴ where we aim to estimate $\boldsymbol{\mu}_{\bar{\alpha}}$ and $\mathbf{V}_{\bar{\alpha}}$ through Equations 11–13 using multiple evo-
³⁵⁵ lutionary replicates starting from a common base population.

³⁵⁶ Rather than working with the allele frequency changes directly, we project them
³⁵⁷ on to a new (reduced) basis and denote this new vector of changes as $\Delta\overrightarrow{\mathbf{p}} = \mathbf{P}\Delta\mathbf{p}$
³⁵⁸ where $\mathbf{P}$ is some projection matrix. We chose a projection that collapses allele fre-
³⁵⁹ quency changes into the non-null subspace of $\mathbf{L}_0$, since $V_{\bar{A}}(0)$ only depends on this
³⁶⁰ subspace (Supplementary information S4 and Supporting Information in de Los Cam-
³⁶¹ pos *et al.* (2015)). To do this, let $\mathbf{U_L}$ be the eigenvectors of $\mathbf{L}_0$ with non-zero eigen-
³⁶² values and then the drift covariance in the reduced subspace is $\mathbf{U_L}^{\top}(\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)})\mathbf{U_L}$.
³⁶³ If we have $\mathbf{U}_2$ and $\mathbf{D}_2$ as the eigenvectors and a diagonal matrix of square-rooted
³⁶⁴ eigenvalues of this matrix, then the projection matrix $\mathbf{P} = \mathbf{D}_2^{-1}\mathbf{U}_2^{\top}\mathbf{U_L}^{\top}$ results in
³⁶⁵ projected allele frequency changes that are identically and independently distributed
³⁶⁶ under drift in the reduced subspace only (see Supplementary information S4).

³⁶⁷ The mean and covariance of the projected allele frequency changes due to pre-
³⁶⁸ dictable selection are:

$$E[\Delta \overrightarrow{\mathbf{p}}_m] = \mathbf{P}_m \boldsymbol{\mathcal{L}}_m \boldsymbol{\mu}_{\bar{\alpha}} \tag{17}$$

and

$$COV(\Delta \overrightarrow{\mathbf{p}}_m, \Delta \overrightarrow{\mathbf{p}}_n) = \mathbf{P}_m \boldsymbol{\mathcal{L}}_m \mathbf{V}_{\bar{\alpha}} \boldsymbol{\mathcal{L}}_n \mathbf{P}_n^\top \tag{18}$$

With $p_{\bar{\alpha}}$ known, the model is a linear mixed model with covariance structure due to the predictable response to selection being proportional (by a factor $\sigma_{\bar{\alpha}}^2$ that is to be estimated) to

$$\overrightarrow{\mathbf{V}}_{m,n} \propto \mathbf{P} \boldsymbol{\mathcal{L}}_m \mathbf{L}_0^{p_{\bar{\alpha}}} \boldsymbol{\mathcal{L}}_n \mathbf{P}^\top \tag{19}$$

between replicates m and n. When $\boldsymbol{\mathcal{L}}^{(m)} = \boldsymbol{\mathcal{L}}^{(n)}$ for all n and m, then this can more easily be fitted by incorporating locus as a random effect with the above covariance structure. The vector of expected values (shown for replicate m) is also proportional to

$$\overrightarrow{\boldsymbol{\mu}}_m \propto \mathbf{P} \boldsymbol{\mathcal{L}}^{(m)} (\mathbf{p}_0 - \mathbf{q}_0) \tag{20}$$

with the associated coefficient $\beta_{\bar{\alpha}}^{(1)}$ to be estimated.

We estimate the parameters of the model by treating it as a separable linear mixed model problem (Richards, 1961). Conditional on a value of $p_{\bar{\alpha}}$ the model is linear mixed model and the conditional (restricted) maximum likelihood can be obtained from asreml (Butler *et al.*, 2023). In order to maximise the (unconditional) likelihood we use the R function optim to find the value of $p_{\bar{\alpha}}$ that results in the highest conditional likelihood. Note that the residual variance should equal one if there is no unpredictable response to selection.

Although linear (mixed) models generate unbiased estimates of $\boldsymbol{\mu}_{\bar{\alpha}}$, the quadratic form $\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}^\top \mathbf{L}_0 \widehat{\boldsymbol{\mu}_{\bar{\alpha}}}$ will be upwardly biased by sampling error in $\beta_{\bar{\alpha}}^{(1)}$. To correct for this bias, we use the inverse Hessian (conditional on the best estimate of $p_{\bar{\alpha}}$) to get an approximate expression for the sampling variance ($S_{\bar{\alpha}}$) of $\beta_{\bar{\alpha}}^{(1)}$, and use this in order to get an improved estimate of the quadratic form as follows (see Supplementary information S5 for a general derivation):

$$\widehat{\boldsymbol{\mu}_{\bar{\alpha}}^\top \mathbf{L}_0 \boldsymbol{\mu}_{\bar{\alpha}}} = \widehat{\boldsymbol{\mu}_{\bar{\alpha}}}^\top \mathbf{L}_0 \widehat{\boldsymbol{\mu}_{\bar{\alpha}}} - \mathbf{X}^\top \mathbf{L}_0 \mathbf{X} S_{\bar{\alpha}} \tag{21}$$

where $\mathbf{X}$ is the fixed effect design matrix.

In addition, allele frequency changes will rarely be known without error and will most likely be estimated using a pool-seq approach. In such cases, it is necessary to add an additional covariance structure into the model that accommodates the

estimation error. In Supplementary information S6 we derive the covariance structure for projected allele-frequency change (in replicate m) as:

$$\overrightarrow{\boldsymbol{\mathcal{E}}}_m = 2\sigma_o^2 \mathbf{P} \left(\boldsymbol{\mathcal{L}}_{t_m,m} \circ \mathbf{Q}_{t_m,m} + \boldsymbol{\mathcal{L}}_{\tau_m,m} \circ \mathbf{Q}_{\tau_m,m} \right) \mathbf{P}^\top \tag{22}$$

where element ij of a $\mathbf{Q}$ matrix is the number of reads that span site i and j divided by the product of the coverages at the two sites. When $i = j$, $Q_{i,j}$ is simply the inverse of the coverage. When individuals have a constant probability of being sampled, σ_o^2 is expected to be one, but we treat it is a free parameter in order to capture any overdispersion (McCullagh and Nelder, 1989).

Comparison with Buffalo and Coop (2019)

To connect our work with Buffalo and Coop (2019) (henceforth 'B&C') it will be useful to express the covariance matrix $\mathbf{L}$ in terms of a diagonal matrix of standard deviations, $\mathbf{B}$, (half the square-root of the genetic diversities under random mating) and the correlation matrix, $\mathbf{R}$, such that $\mathbf{L} = \mathbf{BRB}$. We can then split the response to selection in generation t into two parts:

$$\begin{aligned} \mathbf{L}_t \boldsymbol{\alpha}_t = & \quad \mathbf{B}_t \mathbf{B}_t \boldsymbol{\alpha}_t \quad + \quad \mathbf{B}_t (\mathbf{R}_t - \mathbf{I}) \mathbf{B}_t \boldsymbol{\alpha}_t \\ = & \qquad \underset{S}{\Delta} \mathbf{p}_t \quad + \quad \underset{L}{\Delta} \mathbf{p}_t \end{aligned} \tag{23}$$

where the first term is due to direct selection at the loci, and the second term is due to linkage-disequilibria with other selected loci. It will also be useful to distinguish the additive *genetic* variance ($V_A(t)$) from the additive *genic* variance ($V_a(t)$) in generation t:

$$V_A(t) = \boldsymbol{\alpha}_t \mathbf{L}_t \boldsymbol{\alpha}_t^\top \tag{24}$$

versus

$$V_a(t) = \boldsymbol{\alpha}_t \mathbf{B}_t \mathbf{B}_t \boldsymbol{\alpha}_t^\top \tag{25}$$

with the distinction being that the additive *genetic* variance ($V_A(t)$) captures the contributions of both genetic diversities at individual loci and the linkage disequilibria between pairs of loci, while the additive *genic* variance ($V_a(t)$) captures the contributions of genetic diversities only.

We can also think about the additive genetic/genic covariance in fitness between time points t and τ for a population with genetic structure equal to that in generation t:

$$C_A(\tau \to t) = \boldsymbol{\alpha}_t \mathbf{L}_t \boldsymbol{\alpha}_\tau^\top \tag{26}$$

 and

$$C_a(\tau \to t) = \boldsymbol{\alpha}_t \mathbf{B}_t \mathbf{B}_t \boldsymbol{\alpha}_\tau^\top \tag{27}$$

⁴²¹ These will differ from $V_A(t)$ or $V_a(t)$ when the average effects at generation τ are
⁴²² different from those at generation t.

⁴²³ In the theory section above, we assumed $\mathbf{L}_t$ and $\boldsymbol{\alpha}_t$ were known and so the change
⁴²⁴ due to both direct and linked selection are fixed quantities, as is $V_A(t)$. In contrast,
⁴²⁵ B&C condition on $\mathbf{B}_t$ and $\boldsymbol{\alpha}_t$, and treat $\mathbf{R}_t$ as a random variable. Consequently,
⁴²⁶ in B&C, the change due to direct selection and $V_a(t)$ are fixed, as in our approach,
⁴²⁷ since genetic diversities and average effects are known. However, the change due to
⁴²⁸ indirect selection and $V_A(t)$ is random since the linkage-disequilibrium is unknown.
⁴²⁹ In our inference section we acknowledge that $\boldsymbol{\alpha}$ cannot be directly observed, but
⁴³⁰ develop a model for the distribution of α's, where estimates of $V_A(t)$ can be made
⁴³¹ after marginalising $\boldsymbol{\alpha}$. Similarly, in the approach of B&C the $\boldsymbol{\alpha}$'s and the elements
⁴³² of $\mathbf{B}$ for selected sites are also not directly observed, but they make a number of
⁴³³ strong assumptions that effectively allow the joint distribution of the average effects
⁴³⁴ and selected site diversities to be marginalised.

⁴³⁵

⁴³⁶ In Supplementary information S7 we work through the derivation of B&C using
⁴³⁷ our own notation and retaining a full multi-locus treatment. For easy comparison
⁴³⁸ with the present work, here we summarise both the explicit and implicit assumptions
⁴³⁹ underlying the general approach of B&C:

⁴⁴⁰ • A) Allele frequency changes are only measured over a single generation.

⁴⁴¹ • B) There is no direct selection on the loci for which allele frequency change is
⁴⁴² measured.

⁴⁴³ • C) The reference allele at a neutral locus is chosen arbitrarily.

⁴⁴⁴ • D) The signed linkage-disequilibrium between selected sites is zero, which pre-
⁴⁴⁵ cludes processes such as Hill-Robertson interference. Under this assumption
⁴⁴⁶ $V_A = V_a$.

⁴⁴⁷ • E) Changes in $\mathbf{R}$ are due to recombination alone - selection and drift are absent.

⁴⁴⁸ • F) Nongametic-phase linkage disequilibrium is absent.

⁴⁴⁹ • G) There is no relationship between the additive genic variation at a selected
⁴⁵⁰ site and its LD with neutral sites, as measured through $R_{t,ij}R_{\tau,ij}$ where i is a

neutral locus and j a selected locus. Note that since genetic diversity determines the additive genic variation and LD (even measured as a correlation) is constrained by the genetic diversities at the two loci, this is unlikely to be met (Sved and Hill, 2018).

- H) The average effects are constant, i.e. $\boldsymbol{\alpha}_t = \boldsymbol{\alpha}_\tau$ if estimates are to be interpreted as the additive genic variance (or additive genetic variance, if Assumption D is met) in generation τ. If they are not constant, it is the additive genic/genetic covariance between generations t and τ that is measured.

- I) The initial expected LD-structure between selected and neutral loci, $E[R^2_{t,ij}]$, is approximated from an expression for $E[L^2_{t,ij}]$, which is itself derived under mutation-recombination-drift balance (i.e. no selection). This assumption is partly relaxed in the Appendix by assuming the expected LD-structure between selected and neutral loci is equal to the observed LD-structure between all loci, not distinguishing between selected and neutral loci (Assumption I-b). However, in many cases the genetic diversity at selected loci will be less than that found at (or assumed for) neutral loci and so $E[R^2_{t,ij}]$ will likely be smaller than assumed (Sved and Hill, 2018).

- J) The ratio of genetic diversity in generation τ to genetic diversity in generation t ($\phi_{t,\tau}$), is constant across all selected loci. Under this assumption, dividing the estimate of $V_a(\tau)$ by $\phi_{t,\tau}$ gives an estimate of $V_a(t)$.

- K) The recombination rate between two sites g base pairs apart is given by $(1 - e^{-2g\bar{r}})/2$ (i.e. Haldane's (1919) mapping function) where $\bar{r}$ is the average crossover rate per site per generation.

- L) Neutral and selected loci are distributed uniformly and independently across the genome such that the distance between them has a triangular distribution.

- M) Since the genetic diversity at selected sites is not measured, it is assumed that $\phi_{t,\tau}$ is equal to the average ratio of genetic diversity in generation τ to genetic diversity in generation t across all neutral loci.

When applying the theory to data, the method of B&C requires two additional assumptions:

- Ib) The average LD between selected and neutral sites is equal to the average LD observed between all sites: see Assumption I) above.

- N) The (co)variance in allele frequency changes divided through by the average genetic diversity is a good approximation for the (co)variance of weighted allele frequency changes where the weights are the inverse of the square root of the genetic diversities.

- O) If allele frequencies are estimated from a sample, then the estimation error is binomially distributed.

Assumptions retained in our approach

Our approach works with diploid populations throughout, and assumes non-overlapping generations. While deriving our main theoretical result (Equation 4), we further assume an absence of meiotic drive and mutation. When extending our approach to practical situations and in our inference approach, we relax most of the assumptions made by B&C, but not all.

- Assumption E is only partly relaxed - the change in $\mathbf{R}$ does not assume a lack of drift but it does assume a lack of selection.

- We also make Assumption H if we choose to interpret $V_{\bar{A}}(0)$ as as an additive genetic variance (i.e. $V_A(0)$) rather than an additive genetic covariance over replicate/time-points.

- Assumptions E & H result in the unpredictable response to selection being zero. While we do not make this assumption, we do assume that the unpredictable responses to selection are independent across replicates. However, when applying the theory to data we assume that the within-replicate (residual) (co)variances are driven by drift only (rather than an unpredictable response to selection) and that the drift (co)variances are equal to their expected values (conditional on $\mathbf{L}_0$). In reality, the within-replicate covariances are likely larger, due to the unpredictable response to selection. This assumption might not be too severe if the unpredictable response to selection results in within-replicate (co)variances that are proportional to those under pure drift, although the residual variance may be higher than expected.

- In the absence of a recombination map, Assumption K may also be applied in our method.

- If $\mathbf{L}_0$ cannot be partitioned into gametic-phase and nongametic-phase contributions, assumption F might also be made by substituting $\mathbf{L}_0$ for $\mathbf{L}_0^{'}$ in $\tilde{\mathbf{L}}_0 \circ \mathbf{N}$ and $\tilde{\mathbf{L}}_0 \circ \mathbf{M}$.

- By relaxing some of the more extreme assumptions of B&C we are instead forced into making assumptions about the mean and covariance structure of the average effects of projected loci when making inferences.

Multilocus simulations

To validate our method and inference approach, we performed multilocus simulations using msprime (version 1.2.0) (Kelleher *et al.*, 2016) and SLiM (version 4.2.2) (Haller and Messer, 2023). The code for the simulations and the downstream analyses performed in R (version 4.3.3) is available on GitHub (https://github.com/manas-ga/Va_simulations) and includes the R library, Vw. We first describe the structure of the simulations in general terms and then discuss the specifics of our parameter choices in the context of the scaling used (see 'Simulation parameters' below). While these simulations do incorporate dominance effects, it is important to acknowledge that they do not include epistatic interactions between loci. They also assume random mating and non-overlapping generations. Furthermore, it is also important to note that genetic variation in fitness in our simulations was maintained, primarily, by drift-recombination-mutation-selection equilibrium, and our simulations do not capture other mechanisms of maintenance of genetic variation for fitness such as balancing selection or frequency-dependent selection.

We simulated a single, contiguous 1 million base-pair long genomic region. Our simulations had two distinct phases: the first phase simulated the history of an ancestral population and the second phase simulated a typical evolve and resequence experiment with independent replicate experimental populations derived from a base population drawn from the ancestral population.

Model for fitness

We used an additive model across loci for log absolute fitness, $log(W)$, as in Buffalo and Coop (2019). This choice was motivated by the fact that the variance in $log(W)$ is approximately equal to the variance in relative fitness (see Appendix 1 in Lynch and Walsh (1998)). Across-locus additivity implies that the genotypic value of log fitness for individual k is equal to $Y_k = \sum_{i=1}^{n_L} y_{k,i}$, where $y_{k,i}$ represents the genotypic contribution made by locus i to the log absolute fitness of individual k. Within loci, we employed the classical quantitative genetic fitness scheme (pp. 67 in Lynch and Walsh (1998)) adapted for *proportions* ($c_{k,i}$'s) (as opposed to counts) of reference alleles within individuals:

$$y_{k,i} = \begin{cases} 0 & \text{if } c_{k,i} = 0 \\ (1 + \kappa_i)\eta_i/2 & \text{if } c_{k,i} = 0.5 \\ \eta_i & \text{if } c_{k,i} = 1 \end{cases} \tag{28}$$

where κ determines the degree of dominance. Note that $\kappa = 0$ implies additivity and $\kappa = -1$ complete recessivity. The average effect for log absolute fitness at locus i is then given by $\eta_i^{(a)} = \eta_i[1 + \kappa_i(q_i - p_i)]$ (see Equation 4.10b in Lynch and Walsh (1998)). The average effects for (relative) fitness (α's) are well approximated by the $\eta^{(a)}$'s when they are small in magnitude, but higher-order approximations are required when the $\eta^{(a)}$'s are large (see Supplementary information S8). To obtain the log fitness of each individual we added a noise term drawn from a standard normal distribution (mean = 0, variance = 1) to the genotypic value. The absolute fitness of each individual was then obtained by exponentiating the individual's log fitness.

We sampled the η's from a distribution comprising a weighted mixture of three distributions: (1) a point mass at $\eta = 0$ representing neutral mutations, (2) a reflected gamma distribution with shape = 0.3 and scale = η_{scale} (typically 0.066, unless specified otherwise) representing deleterious mutations, and (3) a gamma distribution with shape = 0.3 and scale = η_{scale} (typically 0.066, unless specified otherwise) representing beneficial mutations. Thus, for both deleterious and beneficial mutations, the mean absolute η (i.e. $E[|\eta|]$) (scale × shape) was, typically, 0.02. The ratio of the frequency of beneficial to deleterious mutations, and η_{scale} varied among simulations. Below, we refer to the distribution of non-neutral η's (i.e. omitting mixture component (1) described above) as the 'distribution of fitness effects' (DFE), but note that this is in fact the distribution of non-zero effects on log fitness.

Phase 1 ('history phase'): Simulating the history of an ancestral population

We first used a neutral coalescent simulation implemented in msprime (Kelleher *et al.*, 2016) to construct genealogies for 2,500 diploid genomes (i.e. $N_e = 2,500$). To initialise a (non-equilibrium) set of selected loci, we then simulated mutations at a rate μ_{msp}, using the pyslim package to attach fitness effects (η) drawn randomly from the non-neutral part of the distribution described above. Since derived mutations are rare, and the DFE is predominantly deleterious, this will generate a positive relationship between the reference allele frequency and the fitness effects η's (and therefore the α's, i.e. the average effects for relative fitness), such that $\beta_{\bar{\alpha}}^{(1)} > 0$ but there should be no relationship between genetic diversities and the η's (i.e. $p_{\bar{\alpha}} = 0$). We implemented the fitness model described above by using SLiM's (Haller and

Messer, 2023) default recipe for polygenic selection (in the Wright-Fisher mode). SLiM code for implementing dominance effects in a quantitative genetic framework was adapted from Schaal *et al.* (2022). To reach mutation-selection-drift balance, we then let this population of 2,500 individuals evolve forward in time with selection for 25,000 generations. Non-neutral mutations drawn from the same DFE were allowed to occur in this period at a rate given by μ_{SLiM}. At generation 25,000, as the alleles reach mutation-selection-drift balance, we expect $\beta_{\bar{\alpha}}^{(1)}$ to have become more positive and $p_{\bar{\alpha}}$ to have become more negative, better reflecting a real population undergoing selection.

In generation 25,000, we sampled N_0 diploid individuals (typically 1000, unless specified otherwise) from this population, which then go on to become the base population in the next phase of the simulation. At this stage, we generated complete genomes for the N_0 individuals in the base population by using pyslim to add neutral mutations to the tree sequence recorded so far. To obtain our target number of loci, n_L (typically 65,000, unless specified otherwise), we set the neutral mutation rate to be $(n_L - n_{L_S})/g$ where n_{L_S} is the number of non-neutral segregating sites already present and g is the total branch length of the recorded tree sequence. We recorded the phased genotype of each parent at each locus, allowing us to construct $\mathbf{L}_0$ and $\mathbf{L}_0^{'}$.

Phase 2 ('experiment phase'): Simulating an evolve and resequence experiment

In the second phase of our simulations, again implemented in SLiM, we first allowed the base population to undergo one round of reproduction without selection to establish replicate experimental populations (typically 1,000 individuals in each of 10 replicates, unless specified otherwise). Next, we allowed each of these populations to evolve forward in time (typically three generations, unless specified otherwise) with selection as in the history phase. Since our goal was to estimate V_A in the base population, we restricted our analyses only to the set of loci segregating in the base population. Any new mutations in subsequent generations would, in all likelihood, occur at loci outside this set. Therefore, we did not simulate new mutations during the experiment phase. For each of the independent replicate populations, we recorded the genome-wide vector of allele frequencies in each generation of the experiment.

Simulation parameters

Varying the true levels of V_A

We varied the true (i.e. simulated) levels of V_A from *ca.* 0.01 to 0.1 by varying the number of non-neutral segregating sites (n_{L_S}) in the base population using a range of rates of non-neutral mutations in the history phase in both msprime (μ_{msp}) and SLiM (μ_{SLiM}). For example, in simulations with $E[|\eta|] = 0.02$ (i.e. $\eta_{scale} = 0.066$) and no beneficial mutations, we varied μ_{msp} between 5.56×10^{-9} and 5.56×10^{-8} and μ_{SLiM} between 5.56×10^{-7} and 5.56×10^{-6}, which resulted in n_{L_S} varying between *ca.* 10,000 and 64,000 in the base population (for other scenarios see Table 1). We set μ_{msp} to be an order of magnitude smaller than μ_{SLiM} in most simulations because otherwise all individuals would have had a fitness of zero at the end of the coalescent part of the history phase. This extreme mutation load arises because deleterious alleles, having previously evolved under neutrality, would segregate at fairly high frequencies. However, the choice of μ_{msp} is unlikely to significantly affect the composition of the base population in most simulations, since the non-neutral genetic diversity of the base population of the experiment phase was primarily determined by the drift-recombination-mutation-selection equilibrium reached over the course of the 25,000 generation long forward simulation, and therefore primarily dependent on μ_{SLiM}.

Number of segregating sites (n_L)

The requirement for handling large $n_L \times n_L$ matrices (e.g. $\mathbf{L}_0$, $\mathbf{L}_0^{'}$, $\mathbf{M}$, and $\mathbf{N}$, etc.) imposed an upper bound of around 70,000 on n_L to permit the analysis of simulations in parallel. Under most scenarios the entire target range of simulated V_A (i.e. approximately 0.01 to 0.1) could be achieved without n_{L_S} exceeding 65,000. Therefore, in general, we set n_L to 65,000 (but see 'Simplified simulations' below). However, under a small minority of scenarios – for example, at higher map lengths in the history phase (Figure 5B) or under strong dominance (Figure 3) – generating a larger V_A required significantly more than 65,000 non-neutral segregating sites. Consequently, in such cases, we had to discard simulations in which n_{L_S} was greater than 65,000 which generally happened at the higher mutation rates and, therefore, the higher levels of V_A.

Map length of the simulated region

There are two recombination-related parameters likely to influence the performance of our method: (1) the V_A per map length, and (2) the density of segregating sites per unit map length. Since our simulations were scaled down in an important way –

⁶⁴⁸ namely, they assumed that V_A was limited to only about 65,000 segregating sites as
⁶⁴⁹ opposed to millions of sites in real populations – it was not possible to simultaneously
⁶⁵⁰ set parameters (1) and (2) close to their realistic values. Therefore, we varied the
⁶⁵¹ effective map length from 0.001 to 2 morgans which spans scenarios where either the
⁶⁵² total map length of the simulated region is comparable to a typical *D. melanogaster*
⁶⁵³ autosome (∼0.5 morgan) or the density of segregating sites is comparable (∼0.065
⁶⁵⁴ morgan). Exact settings are detailed in the 'Reference parameters' section below.
⁶⁵⁵ The typical *D. melanogaster* map-length was determined from an effective crossover
⁶⁵⁶ rate of 10^{-8} (Comeron *et al.*, 2012; Wang *et al.*, 2023) and a 50Mb chromosome. The
⁶⁵⁷ typical site-density was determined under the assumption that there are one million
⁶⁵⁸ segregating sites per morgan, which is reasonably consistent with a recent evolve-
⁶⁵⁹ and-resequence study employing *D. melanogaster* (Bitter *et al.* (2024): 827,200 and
⁶⁶⁰ 893,465 sites on Chromosome 2 and 3 respectively).

⁶⁶¹ Since we only simulated a population of 2,500 individuals in the history phase,
⁶⁶² which is ∼500 times smaller than the estimate of N_e in *D. melanogaster* (Campos
⁶⁶³ *et al.*, 2013; Campos and Charlesworth, 2019), we scaled our map-length upwards by
⁶⁶⁴ a factor of 500 to achieve the target effective map-length described above (Campos
⁶⁶⁵ and Charlesworth (2019) but see Dabi and Schrider (2025) and Ferrari *et al.* (2025)
⁶⁶⁶ for potential issues). On the other hand, since the experiment phase simulates a
⁶⁶⁷ realistic evolve-and-resequence experiment with $N_e \approx 1,000$, this scaling was not
⁶⁶⁸ applied to map lengths in the experiment phase (see 'Reference parameters' below).
⁶⁶⁹ In what follows, the map-lengths implemented in the simulations are reported rather
⁶⁷⁰ than the target effective map-lengths.

Degree of dominance

⁶⁷² In the simulations with dominance effects (i.e. $\kappa \neq 0$) we always modelled delete-
⁶⁷³ rious alleles to be recessive. In practical terms, this meant that κ_i was set to be
⁶⁷⁴ positive when η_i was positive, and κ_i was set to be negative when η_i was negative.
⁶⁷⁵ For simplicity, in a given simulation, we assumed $|\kappa_i|$ to be the same across all loci,
⁶⁷⁶ although the actual dominance deviation, $d_i = \kappa_i \eta_i$ would be locus-specific. We sim-
⁶⁷⁷ ulated two different levels of $|\kappa|$: (1) $|\kappa| = 0.5$, and (2) $|\kappa| = 0.9$. These correspond
⁶⁷⁸ to $h = 0.25$ and $h = 0.05$, respectively, when the heterozygous fitness is $1 - hs$.
⁶⁷⁹ While $h = 0.25$ is consistent with the theoretical expectation from a fitness land-
⁶⁸⁰ scape based model (Manna *et al.*, 2011), $h = 0.05$, where every deleterious allele is
⁶⁸¹ almost completely recessive is probably extreme. Since full simulations with domi-
⁶⁸² nance were up to two orders of magnitude slower than their additive counterparts,
⁶⁸³ we switched on dominance effects only in the last 5,000 generations of the history

phase (i.e. from generation 20,001) as well as in the experiment phase. Relative to additive simulations, full simulations with dominance also required significantly more segregating sites to generate a given level of V_A. To generate sufficient V_A (i.e. between *ca.* 0.01 and 0.1) while keeping n_{L_S} below 67,500, we simulated deleterious mutations from a gamma distribution with a slightly larger mean effect on log fitness (i.e. $E[|\eta|] = 0.03$) in the history phase and set $|\kappa|$ to either 0.5 or 0.75. To test our method against extreme dominance (i.e. $|\kappa| = 0.9$), we performed additional full simulations in which we simulated the entire history phase under additivity, and switched on dominance effects only in the experiment phase. While the resulting genetic composition was far from the expected equilibrium, deleterious alleles were expected to be segregating at substantially lower frequencies than in the simplified simulations.

Reference parameters

Our aim was to investigate the sensitivity of our method to the following parameters: (1) map length in the history phase (0.5 morgan, 5 morgans, 50 morgans, 250 morgans), (2) map length in the experiment phase (0.01 morgan, 0.2 morgan, 2 morgans), (3) number of replicate populations in the experiment phase (3, 5, 10), (4) the population size of each of the replicate populations (100, 500, 1,000), (5) number of generations over which allele frequency changes were recorded in the experiment phase (1, 3, 5), (6) the ratio of the rates of beneficial to deleterious mutations (0%, 0.02 %, 2%), and (7) the mean of the gamma distribution from which non-neutral η's were sampled, $E[|\eta|]$ (0.02, 0.03, 0.06), and (8) the degree of dominance, κ (0, 0.5, 0.9).

Rather than vary all parameters in a fully factorial design, we selected a reference parameter set and explored the sensitivity of the method by changing each parameter in turn. The reference parameter set was (1) a map length of 0.5 morgan in the history phase (2) a map length of 2 morgans in the experiment phase, (3) 10 replicate populations in the experiment phase, (4) a population size of 1,000 in each of the replicate populations in the experiment phase, (5) 3 generations over which allele frequency changes were recorded in the experiment phase, (6) no beneficial mutations in the history phase, (7) $E[|\eta|] = 0.02$, and (8) complete additivity (i.e. $\kappa = 0$). The reference parameter set was chosen to best reflect evolve-and-resequence experiments using *D. melanogaster*.

Note that we varied $E[|\eta|]$ by using three different values (0.066, 0.09, and 0.2) for the scale of the gamma distribution (η_{scale}) from which the η's for non-neutral mutations were drawn while keeping the shape parameter fixed to 0.3. This meant

720 that the coefficient of variation was fixed to $1/\sqrt{0.3}$.

Simplified simulations

In addition to the full simulations described above, we performed a set of simplified,
proof-of-principle simulations to test the logic of our method. These simulations were
different from the full simulations in three ways:

(1) The history phase was highly abbreviated – the forward simulation imple-
mented in SLiM lasted only a single generation without any selection. This meant
that the ancestral population – from which replicate experimental population were
founded – had evolved entirely under neutrality.

(2) Rather than attach η's to the derived alleles generated with msprime, we
attached the η's randomly with respect to either of the two alleles segregating at the
locus. Amendments (1) and (2) generate a scenario where there is no relationship
between allele frequency and η: both $p_{\bar{\alpha}}$ and $\beta_{\bar{\alpha}}^{(1)}$ are expected to be zero.

(3) Given that the ancestral population had evolved without selection, these
simplified simulations had considerably higher genetic diversities compared to the full
simulations. This meant that fewer segregating non-neutral sites (n_{L_S}) were required
to achieve true levels of V_A between 0.01 and 0.1. Therefore, in these simulations,
we set n_L to be 3,000, much lower than the 65,000 used for the full simulations. We
achieved this by varying μ_{msp} between 3×10^{-9} and 2.35×10^{-8}. μ_{SLiM} was set to
0 in these simulations. This resulted in n_{L_S} at the end of the history phase being
between 160 and 1,902. We then added neutral mutations using a suitable rate as
described above such that n_L was expected to be 3,000.

Simulating pool-seq

To investigate the consequences of sampling noise around true allele frequencies in
real data, we reanalysed the reference sets for full and simplified simulations with
allele frequencies in the experiment phase obtained via simulated pool-seq. Specif-
ically, we investigated the sensitivity of our approach to the pool-seq coverage (i.e.
the expected number of reads spanning a given locus) and the degree of overdisper-
sion in the number of individuals sampled. We assumed that the number of reads
mapping to individual k followed a Poisson distribution with mean $\lambda_{x,k}$, which in
turn is sampled from a log-normal distribution with parameters μ_x and V_x. Vary-
ing V_x allowed us to control the degree of heterogeneity in the probabilities with
which individuals are sampled. For a given level of coverage and read length, we
determined the expected number of reads mapping to an individual ($E[\lambda_x]$). For a
given value of V_x and read-length, using the expression for the mean of a lognormal

distribution ($E[\lambda_x] = exp(\mu_x + V_x/2)$), we chose μ_x to achieve the target coverage: $\mu_x = log(E[\lambda_x]) - V_x/2$. We mapped reads to the genomes of various individuals in the population by sampling the starting positions for each read from a uniform distribution.

To select the length of the reads in these simulations, we first note that the expected number of segregating sites spanned by a 150 base-pair read in a recent evolve-and-resequence study (Bitter *et al.*, 2024) (827,200 segregating sites on chromosome 2 which is ∼50 Mb in length) is approximately 2.5. To achieve this, we fixed the read length to be 800 while analysing simplified simulations ($\sim$ 3,000 segregating sites) and 37 when analysing full simulations ($\sim$ 65,000 segregating sites). We used four different combinations of coverage and V_x: (1) coverage = 100x, $V_x = 0$; (2) coverage = 500x, $V_x = 0$; (3) coverage = 1000x, $V_x = 0$); (4) coverage = 1000x, $V_x = log(2)$. Note that $V_x = 0$ corresponds to a situation where individuals are sampled without overdispersion.

Comparison with B&C

To compare the precision and the accuracy of our approach to that of B&C, we performed additional simplified and full simulations. Although originally designed for the covariance in allele frequency change between multiple time points, B&C's method can be readily adapted to allele frequency changes recorded over a single generation in multiple independent evolutionary replicates (see Equation S59 and Buffalo and Coop (2020)). We set simulation parameters to their reference values (see above) with a few modifications. First, to make comparisons of biases clearer, we reduced noise by employing 50 replicate populations, and second, allele frequency change in the experiment phase was recorded over a single generation. We also ran these simulations at four different levels of map length in the history phase: 0.5 morgan, 5 morgans, 50 morgans, and 100 morgans.

To investigate the consequences of Assumptions B, Ib, and N, we implemented B&C's method using six different approaches. For clarity, we label the six approaches using a code (see Table 2) that indicates with superscripts whether an assumption is required ($+$) or not required (0) for an approach to work. To test Assumption B, we recorded allele frequency changes at either all segregating sites (B^+) or neutral sites only (B^0). To test Assumption Ib, we used either the average LD between all sites (Ib^+) or the average LD between selected sites with neutral sites (Ib^0). And finally, we either divided the (co)variance in allele frequency change throughout by twice the average genetic diversity (N^+; analogous to Equation 16 in B&C) or used the (co)variance of weighted allele frequency change, where the weights are the inverse

791 of the square root of twice the genetic diversity (N^0).

Results

Simplified simulations

We begin by discussing results from the simplified simulations in which we expect
there to be no relationship between allele frequencies and α's. Under our reference
parameter set (no dominance, map length in history $= 0.5$ morgan, map length in the
experiment $= 2$ morgans), our method provided precise and unbiased estimates of V_A
throughout the simulated range (0.01– 0.1) (Figure 1a). Furthermore, the estimates
of $p_{\bar{\alpha}}$ and $\beta_{\bar{\alpha}}^{(1)}$ were centred around 0 as expected (Figure 1b-c), and the residual vari-
ance was marginally above 1 (Figure 1d). Next, we investigated how our estimates of
V_A were affected by changing (1) map length of the genomic region being simulated,
(2) population size of each replicate population in the evolve and resequence experi-
ment, (3) number of replicate populations, and (4) number of generations over which
allele frequency changes were recorded in the experiment. Estimates of V_A remained
unbiased as the map length in the experiment became smaller, although estimates
became noisier, particularly at higher values of V_A (Figure 2a). As expected, es-
timates also became noisier as the number of individuals, replicate populations, or
generations became smaller (Figure 2b-d). However, estimates appeared upwardly
biased when the number of replicate populations or generations was small (Figure
2c-d). Estimates of V_A from simplified simulations incorporating dominance effects
in the fitness model were generally precise, but exhibited a small upward bias when
dominance effects were large in magnitude (Figure 3a). As the degree of dominance
increased, the estimates of $p_{\bar{\alpha}}$ and $\beta_{\bar{\alpha}}^{(1)}$ became increasingly more negative and the
residual variance became larger (Figure 3b-d).

Full simulations

In our full simulations we let the ancestral population evolve forward in time with
selection for 25,000 generations before simulating the experiment. In these simu-
lations, we expect the population to be at drift-recombination-mutation-selection
equilibrium, such that p_α is negative and $\beta_{\bar{\alpha}}^{(1)}$ is positive. Results from our refer-
ence set (no dominance, map length in history $= 0.5$ morgan, map length in the
experiment $= 2$ morgans) suggest that not only does our method provide precise and
unbiased estimates of V_A (Figure 4a), but it does so by correctly estimating the signs
of p_α and $\beta_{\bar{\alpha}}^{(1)}$ (Figure 4b-c). As before, estimates of V_A became slightly noisier at

shorter map lengths during the experiment (Figure 5a). The quality of our estimates of V_A was not significantly affected by adding beneficial mutations during the history phase (Figure 5c). On the other hand, our estimates were downwardly biased at larger map lengths in the history phase (Figure 5b, also see Supplementary Figure 1) and when non-neutral loci had on average larger fitness effects ($E[|\eta|]$) (i.e. when η_{scale} was larger) (Figure 5d). The bias at larger $E[|\eta|]$ was likely driven by the loss of additive genic variance at large-effect loci during the experiment (Supplementary Figure 2a-d).

When we simulated the history phase under additivity and switched on dominance only in the experiment phase, we obtained fairly reliable estimates of V_A (Supplementary Figure 4). However, switching on dominance in the last 5,000 generations of the history phase resulted in estimates that were significantly upwardly biased, especially when dominance effects were strong (Figure 3e). However, at the relatively low map length in the history phase (0.5 morgan) used in these simulations, the additive genetic variance for fitness was substantially lower than the additive genic variance (Supplementary Figure 5). This suggested the build up of unusually high negative linkage disequilibria between recessive deleterious alleles, generating pseudo-overdominance (Ohta and Kimura, 1970; Abu-Awad and Waller, 2023), a phenomenon typically observed in regions of extremely low recombination (Salson *et al.*, 2025). At higher levels of map length in the history phase (5 morgans), our method provided unbiased estimates of v_A (Figure 3f).

Finally, we re-analysed our standard set of full simulations in two different ways (Figure 5e-f). First, to accommodate cases in which phased genomes, and therefore $\mathbf{L}_0^{'}$, are unavailable, we assumed that $\mathbf{L}_0^{''}$ was 0 and set $\mathbf{L}_0^{'} = \mathbf{L}_0$. In other words, we assumed that $\tilde{\mathbf{L}}_0 = \mathbf{L}_0$. Our analyses suggest that this assumption leads to a slight downward bias in our estimates of V_A (Figure 5e). Second, to account for instances where N_E in the experiment phase is unknown, we replaced our estimate of N_E in the experiment phase by the number of individuals. Our analyses suggest that this does not affect our estimates of V_A in any way (Figure 5f), with the faster than expected drift being absorbed by an increased residual variance (Supplementary Figure 3): estimates of N_E can be obtained by dividing the assumed value of N_E by the estimated residual variance.

Simulating pool-seq

When we incorporated the expected covariance structure due to pool-seq sampling in our models, the performance of our method was practically unaffected by obtaining allele frequencies in the experiment phase using simulated pool-seq in the simplified

simulations (Figure 6A), although a modest negative bias was observed at low (100x) coverage in the full simulations (Supplementary Figure 6). Additionally, both in the simplified and the full simulations, the degree of overdispersion in the number of reads mapping to an individual did not adversely affect our estimates of V_A. Along expected lines (see Supplementary information S6), omitting the covariance structure of pool-seq sampling from our models led to estimates of V_A that were significantly downwardly biased, with virtually no V_A detected at 100x coverage (Figure 6B).

Comparison with B&C

In both simplified and full simulations, all implementations of B&C's method resulted in a strong upward bias at low recombination rates in the history phase (Figure 7 and Figure 8). This was likely a consequence of Assumption G being violated; the correlation between the contribution of a selected site j to V_a and the persistent association of this site with neutral alleles ($\sum_{\mathcal{N}_i}(R_{i_m j_m} R_{i_n j_n})$) was considerably higher than 0 when the map length in the history phase was low (panel h of Figure 7 and Figure 8). In contrast, the estimates of V_A provided by our method were unbiased and had far greater precision compared to any of the six implementations of B&C's method in the simplified simulations (Figure 7g), although there was a small upward bias at low recombination rates in the full simulations (Figure 8g). Due to the difficulty of distinguishing selected from unselected sites, B&C suggested that the average LD between all sites could be used instead of the LD between selected and unselected sites (Assumption Ib) and indeed this assumption seems to be well justified (panels d & f in Figure 7 and Figure 8 are very similar). However, an inability to distinguish selected from unselected sites would likely mean that B&C's method is applied to allele frequency change data from *all* segregating sites, contravening Assumption B. This results in considerable overestimation of V_A in simplified simulations (Figure 7a-b) but not in full simulations (Figure 8a-b). Under all scenarios the bias in the method of B&C can be reduced (but not eliminated) by relaxing Assumption N (compare panels b, d, and f in Figure 7-Figure 8 to panels a, c and e). Under assumption N, it is assumed that the (co)variance in allele frequency change weighted by the square root of twice the genetic diversities can be approximated by the (co)variances in allele frequency change divided through by twice the average diversities. However, this approximation is not required.

Discussion

In this paper we estimate the additive genetic variance for relative fitness (V_A) directly from the change in the genetic composition of a population caused by selection. Assuming only the absence of meiotic drive and ignoring mutation, we show that V_A can be conveniently expressed as a function of the genome-wide genetic diversity matrix $\mathbf{L}$, and the vector of genome-wide expected allele frequency change due to selection, $E[\Delta \mathbf{p}]$. In our inference approach, we describe how a linear mixed model can be employed to estimate $E[\Delta \mathbf{p}]$ via independent evolutionary replicates derived from the same base population with a known $\mathbf{L}$ – a common feature of evolve-and-resequence studies. Unlike alternative methods (Buffalo and Coop, 2019), this allows us to obtain estimates of V_A that are largely robust to the underlying genetic properties of the population and the genetic architecture of fitness. Moreover, the underlying modelling framework not only allows V_A to be estimated, but allows inferences about the relationship between effect sizes and allele frequency.

Although $E[\Delta p]$'s at individual loci cannot be usefully estimated, for our purposes it is sufficient to estimate their distribution as parameterised through the mean vector $\boldsymbol{\mu_{\bar{\alpha}}}$ and the (co)variance matrix $\boldsymbol{V_{\bar{\alpha}}}$ for the distribution of the average effects for fitness, the α's. Our inference approach uses a low-dimensional (3-parameter), but biologically sensible, model for the means and (co)variances. It is superficially surprising that such a simple model for an n_L (the number of loci) dimensional distribution can produce accurate results. However, in a typical dataset one expects the number of individuals, N, to be far smaller than the number of loci, n_L. Therefore, the non-null subspace of $\mathbf{L}$ has only N dimensions, and we can work with allele frequency changes projected into this reduced space. This provides a route to understanding how the distribution of α's can be estimated from selected changes in allele frequency that must be negligible compared to the impact of drift: the projection defines '*chunks*' of genome, and it is the frequency changes in these chunks, rather than individual alleles, that are tracked. Since the aggregate fitness effects of alleles across chunks will be more substantial, they can be more easily detected. Moreover, since these aggregate effects may involve a large number of loci, they will, thanks to the central limit theorem, tend to normality and, conditional on the projection, converge in distribution. Since the projection is defined by $\mathbf{L}$, a model of the α's that conditions on aspects of $\mathbf{L}$ ($p - q$ and pq in our case) is expected to be sufficient.

The results of our simulations demonstrate that our approach provides usefully precise and consistent estimates of V_A over a wide range of parameter combinations and experimental designs. In the simplified simulations, in which the relationship between allele frequencies and α's was expected to be absent, the model requires

only a single parameter: the variance in average effects, $\sigma_{\bar{\alpha}}^2$. If such a model is assumed, this is a relatively straightforward inference problem and the method of Buffalo and Coop (2019) can give accurate estimates if selected and neutral sites can be distinguished, and certain patterns of recombination and linkage-disequilibrium hold. Although our approach does not assume such a model, it can accurately infer V_A and also the lack of relationship between allele frequencies and α's, with the two parameters that determine the relationship between allele frequency and α all centered on their null expectations (Figure 1). However, in reality, independence between allele frequency and α seems implausible at mutation-selection-drift balance, and the elevated contribution of high-diversity loci to V_A under the simple, but unrealistic, scenario may result in greater power to detect V_A.

At mutation-selection-drift balance a negative relationship is expected between α's and allele frequencies (Charlesworth and Charlesworth, 2010). Although it could be argued that the strength of this relationship will be reduced in experimental evolution studies, which generally expose populations to novel environmental stressors (such as a laboratory environment, pathogens (Basu *et al.*, 2024), extreme population densities (Joshi and Mueller, 1996) and temperatures (Singh *et al.*, 2015; Hsu *et al.*, 2024), desiccation (Gibbs *et al.*, 1997), malnutrition (Kawecki *et al.*, 2021), or toxic substances (Godinho *et al.*, 2024; Xiao *et al.*, 2019)) it seems unlikely that no relationship would persist. Even under the more realistic scenario of our full simulations, in which the base population had evolved under selection for 25,000 generations, our approach provided reliable estimates of V_A that were remarkably robust to the details of the distribution of fitness effects, such as the relative frequency of beneficial mutations, or properties of the population such as the recombination rate. Furthermore, it did so by correctly inferring a negative relationship between fitness effects of alleles and genetic diversity, and a positive relationship between fitness effects and $p - q$, as expected under mutation-selection balance.

An important assumption of our method is that any consistent change in the α's is vanishingly small in the time-frame over which allele frequency changes are recorded. However, when fitness-causing alleles act non-additively, α's depend on allele frequency and so are expected to change as allele frequencies change. In spite of this, in the simplified simulations where allele frequency change was expected to be large, the performance of the method was minimally affected when allowing dominance at all loci. Moreover, given the form $\alpha_i \approx \eta_i[1 - \kappa_i(p_i - q_i)]$ (where the product $\eta_i \kappa_i$ is always positive in our simulations), the method correctly inferred that the two parameters describing the relationship between the α's and allele frequencies would become increasingly negative with the degree of dominance (Figure 3). Surprisingly, when we allowed dominance effects in the history phase and so allele frequencies

and effects had come to an equilibrium we found that our method overestimated V_A to varying degrees depending on how recessive the deleterious allele was. However, this was not because of a failure of the infinitesimal approximation in the experimental phase, but because substantial negative LD had built up during the history phase and our model of $V_{\bar{\alpha}}$ failed to capture this: the additive genetic variance was roughly half the additive genic variance. It seems that our simulations were resulting in substantial pseudo-overdominance (Abu-Awad and Waller, 2023) - a phenomenon characteristic of regions of very low recombination (Salson et al., 2025). When we performed identical simulations at higher recombination rates in the history phase, this problem was not observed.

Another key assumption of our method is that any consistent, selection-induced changes in $\mathbf{L}$ are negligible, which then allows us to model the expected evolution of $\mathbf{L}$ due to the action of drift and recombination. Selection-induced changes in $\mathbf{L}$ may be safely ignored if the genetic architecture of fitness is sufficiently polygenic such that selection coefficients associated with each locus are vanishingly small. It is important to note that this assumption was fully relaxed in all of our simulations, in which selection was free to influence the evolution of $\mathbf{L}$ along with drift and recombination. Despite this, the performance of our method was virtually unaffected, even in the simplified simulations where selection-induced changes in $\mathbf{L}$ were expected to be relatively large.

While our method generally performs well, some biases were observed. In terms of experimental design, small upward biases are evident when power is low - either because there are few replicates and/or allele frequency change is only calculated over a single generation. In terms of genetic architecture, our approach marginally underestimates V_A when the fitness effects of new non-neutral mutations are large. This downward bias seems to arise because there is a large contribution of rare highly deleterious variants to V_A and these get lost during the experimental phase. Nevertheless, even when 25% of V_A was lost during the course of the experiment the impact on estimates was rather minor. Similarly, modest downward biases were observed when the recombination rate in the history phase was increased, but the source of this bias has been harder to diagnose. Increasing the recombination rate resulted in an increase in the number of segregating selected sites and their genetic diversity, and a steeper relationship between α and genetic diversity, consistent with a reduction in the effects of background selection (Charlesworth et al., 1993) on weakly selected sites (Stephan et al., 1999). While it is not clear why this causes downward bias in the estimates, it is also unclear whether real populations would ever have such extreme genetic architecture where the bulk of the additive genetic variance for fitness is caused by highly deleterious variants segregating at very low frequencies. In

order to make the forward simulations manageable we were working with considerably smaller population and genome sizes than are typical of real populations. It is not clear, however, whether the standard rescaling of mutations rates, recombination rates and selection coefficients to accommodate this downsizing results in genetic architectures that would be typical of larger populations and larger genomes (Dabi and Schrider, 2025).

While our method performs well when applied to simulated data, application to real-world data would involve overcoming a number of challenges. First, we require genome-wide allele frequency change data from multiple independent evolutionary replicates – although this should be readily available for most evolve-and-resequence experiments using approaches such as pool-seq. An important point of consideration, therefore, is the minimum pool-seq coverage that the method requires. For our sim-plified simulations, there was little deterioration in model performance as coverage dropped or overdispersion increased, suggesting modest coverage (100x) should be sufficient if the appropriate covariance structure is used. While this might be surpris-ing, the relevant parameter is probably not coverage *per se*, but the number of reads overlapping at least one segregating site per unit map length, as this determines how accurately the frequency of a 'chunk' of genome can be measured. Using a similar argument, Tilk *et al.* (2019) demonstrated that supplementing pool-seq sampling with haplotype inference tools can result in nearly 500x 'effective' coverage at 10x empirical coverage. Given the ancestral haplotype structure is a core requirement of our method, it may be possible that even greater precision could be achieved by also using such tools. However, this could come with the risk that estimation errors are correlated across replicates and be mistaken for patterns of selection. For our full simulations, a moderate downward bias was observed at 100x coverage. While this may suggest that our method demands relatively high coverage ($\geq$500x), it is worth noting that we only implemented an approximate (i.e. diagonal) covariance structure for pool-seq sampling while analysing our simulations due to computational limita-tions. Incorporating the full covariance structure should lead to improved estimates.

Second, we require the genetic diversity matrix $\mathbf{L}$ in the base population from which the replicates are derived. This is not always the case for evolve-and-resequence studies, in which base populations are often split into replicate baseline popula-tions long before the experiment, or when newer selection regimes are derived mid-experiment (Burke *et al.*, 2010; Singh *et al.*, 2015; Gupta *et al.*, 2016; Robinson *et al.*, 2023). Furthermore, our method requires that sufficient individuals from the base population are individually sequenced to estimate $\mathbf{L}$, or – even better – its gametic phase ($\mathbf{L}'$) and non-gametic phase ($\mathbf{L}''$) components, although the required phasing should become more readily available with long-read sequencing. Third, to predict

how **L** evolves, we require an estimate of the recombination probability between all pairs of segregating sites, such as a recombination map for the population. In reality, recombination maps are likely to have been derived from other populations, in which recombination patterns may differ (Johnston, 2024). Although recombination maps can be approximated, for example by using Haldane's mapping function, it is not clear how sensitive the method is to errors in the recombination map. Fourth, our method requires the mean number of generations over which allele frequency changes are calculated, which may be hard to infer with overlapping generations. Since allele-frequency change will be roughly proportional to the number of generations that have elapsed (n_g), and V_A is quadratic in allele-frequency change, estimates of V_A might be out by a factor $(n_g/\widehat{n_g})^2$, where $\widehat{n_g}$ is the assumed number of generations. Fifth, our method uses effective population size to predict how **L** evolves (N_e) and to derive expressions for the drift (co)variance (N_E). Although reliable estimates of both effective population sizes may be hard to obtain, this is unlikely to be a major issue because: (i) $\mathbf{L}'$ decays with a rate roughly proportional to $1 - 1/N_e$ (likely making it insensitive to errors in estimating N_e) and (ii) our simulations suggest that using the wrong N_E does not adversely affect our estimates of V_A because the misspecification is absorbed by the residual variance of the model. Sixth, our simulations minimise the unpredictable response to selection by modelling a constant environment. However, with selection coefficients that vary in time or across replicates, the unpredictable response to selection will be greater. Indeed in outdoor mesocosms of *D. melanogaster*, Bitter *et al.* (2024) report that allele frequency changes can switch signs over time-points separated by a matter of weeks in spite of exhibiting highly concordant evolution between replicates. It is not clear to what degree this will affect inferences. Finally, selection can often act in different ways in different contexts such as space (Whitlock, 2015; Delph, 2018), time, and between the two sexes (Schenkel *et al.*, 2018). Our approach captures the effects of selection averaged over all these different contexts. Specifically, if loci have different fitness effects in males and females, we effectively estimate $(V_{A,f} + V_{A,m} + 2COV_{A,mf})/4$, where $V_{A,f}$ and $V_{A,m}$ are the additive genetic variances for relative fitness in females and males respectively, and $COV_{A,mf}$ is the intersexual additive genetic covariance for relative fitness. For these reasons, when applied to replicate populations, our estimates of V_A are perhaps best thought of as the additive genetic *covariance* in fitness between replicates. Although rarely made explicit, estimates from wild systems should be interpreted in the same way: the additive genetic covariance between the environments in which relatives live (Vehviläinen *et al.*, 2008).

Analysing data from wild populations is likely to entail further challenges. For example, in natural populations, allele frequency change from immigration may be

consequential, and without some modification is likely to be mistaken for allele frequency change caused by natural selection (Simon and Coop, 2024). Furthermore, replicate populations are unlikely to be available for natural systems – with some exceptions such as Trinidadian guppies (Reznick and Bryga, 1996) – although with some modifications our method could be adapted to use allele frequency change data from multiple time points. However, a lack of individual-level sequences in the base population would mean that the estimates of $\mathbf{L}$ will likely be considerably noisier in natural populations.

While we accept that the data requirements for our method are steep, and the list of caveats appears long, we do think that information about V_A can be successfully leveraged from current evolve and resequence studies. Going forward, we hope this work will inform future evolve and resequence study design, and that estimates of V_A from a wide range of organisms and environments become available. Partitioning V_A into genomic features is an obvious next step, and in the future we hope to go beyond simply knowing the magnitude of V_A and start to understand its underlying causes.

Data availability statement

This study does not use any data. The code used for simulations and analyses is available in the following GitHub repository: https://github.com/manas-ga/Va_simulations.

Acknowledgments

The authors would like to thank Bill Hill, Brian Charlesworth, Peter Keightley, Konrad Lohse, Bruce Walsh, Ben Longdon and Vince Buffalo for their insightful comments, and The Argyle for hosting us at the early stages of this project. Computer simulations and analyses described here were performed on the AC3 computing cluster based at Ashworth Laboratories, and on Eddie, the University of Edinburgh's main high performance computing facility.

Study funding

This work was funded by a Natural Environment Research Council (NERC) grant (reference: NE/W001330/1). For the purpose of open access, the authors have

applied a Creative Commons Attribution (CC BY) license to any Author Accepted
Manuscript version arising from this submission.

Conflict of interest

The authors have no conflicts of interest to declare.

References

Abbott, J. K., and E. H. Morrow, 2011 Obtaining snapshots of genetic variation
  using hemiclonal analysis. Trends in Ecology & Evolution **26**: 359–368.

Abu-Awad, D., and D. Waller, 2023 Conditions for maintaining and eroding pseudo-
  overdominance and its contribution to inbreeding depression. Peer Community
  Journal **3**.

Barton, N. H., and L. Partridge, 2000 Measuring fitness by means of balancer chro-
  mosomes. Genetics Research **75**: 297–313.

Basu, A., K. Tekade, A. Singh, P. N. Das, and N. G. Prasad, 2024 Experimental
  evolution for improved postinfection survival selects for increased disease resistance
  in *Drosophila melanogaster*. Evolution **78**: 1831–1843.

Bitter, M., S. Berardi, H. Oken, A. Huynh, E. Lappo *et al.*, 2024 Continuously
  fluctuating selection reveals fine granularity of adaptation. Nature **634**: 389–396.

Bonnet, T., M. B. Morrissey, P. de Villemereuil, S. C. Alberts, P. Arcese *et al.*, 2022
  Genetic variance in fitness indicates rapid contemporary adaptive evolution in wild
  animals. Science **376**: 1012–1016.

Buffalo, V., and G. Coop, 2019 The linked selection signature of rapid adaptation in
  temporal genomic data. Genetics **213**: 1007–1045.

Buffalo, V., and G. Coop, 2020 Estimating the genome-wide contribution of selection
  to temporal allele frequency change. Proceedings of the National Academy of
  Sciences **117**: 20672–20680.

Bulmer, M. G., 1980 *The mathematical theory of quantitative genetics.*. Clarendon
  Press.

Burke, M. K., J. P. Dunham, P. Shahrestani, K. R. Thornton, M. R. Rose *et al.*, 2010 Genome-wide analysis of a long-term evolution experiment with *Drosophila*. Nature **467**: 587–590.

Burt, A., 1995 The evolution of fitness. Evolution **49**: 1–8.

Butler, D., B. R. Cullis, A. R. Gilmour, B. J. Gogel, and R. Thompson, 2023 *ASReml-R Reference Manual, Version 4.2*. Queensland DPI, Brisbane, Australia.

Campos, J. L., and B. Charlesworth, 2019 The effects on neutral variability of recurrent selective sweeps and background selection. Genetics **212**: 287–303.

Campos, J. L., K. Zeng, D. J. Parker, B. Charlesworth, and P. R. Haddrill, 2013 Codon usage bias and effective population sizes on the X chromosome versus the autosomes in *Drosophila melanogaster*. Molecular biology and evolution **30**: 811–823.

Charlesworth, B., and D. Charlesworth, 2010 *Elements of evolutionary genetics*, volume 42. Springer.

Charlesworth, B., M. Morgan, and D. Charlesworth, 1993 The effect of deleterious mutations on neutral molecular variation. Genetics **134**: 1289–1303.

Charmantier, A., and B. C. Sheldon, 2006 Testing genetic models of mate choice evolution in the wild. Trends in Ecology & Evolution **21**: 417–419.

Comeron, J. M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in *Drosophila melanogaster*. PLoS Genetics **8**: e1002905.

Coulson, T., T. Benton, P. Lundberg, S. Dall, B. Kendall *et al.*, 2006 Estimating individual contributions to population growth: evolutionary fitness in ecological time. Proceedings of the Royal Society B: Biological Sciences **273**: 547–555.

Dabi, A., and D. R. Schrider, 2025 Population size rescaling significantly biases outcomes of forward-in-time population genetic simulations. Genetics **229**: iyae180.

de Los Campos, G., D. Sorensen, and D. Gianola, 2015 Genomic heritability: what is it? PLoS Genetics **11**: e1005048.

Delph, L. F., 2018 The study of local adaptation: a thriving field of research. Journal of Heredity **109**: 1–2.

Edwards, A. W., 1994 The fundamental theorem of natural selection. Biological Reviews **69**: 443–474.

Ewens, W. J., 1989 An interpretation and proof of the fundamental theorem of natural selection. Theoretical population biology **36**: 167–180.

Ewens, W. J., 2024 The fundamental theorem of natural selection: the end of a story. Evolution **78**: 803–808.

Falconer, D., and T. Mackay, 1996 *Introduction to Quantitative Genetics, 4th edn*. Addison Wesley Longman, Harlow.

Ferrari, T., S. Feng, X. Zhang, and J. Mooney, 2025 Parameter scaling in population genetics simulations may introduce unintended background selection: Considerations for scaled simulation design. Genome Biology and Evolution **17**: evaf097.

Fisher, R. A., 1930 *The genetical theory of natural selection*. Oxford University Press.

Fisher, R. A., 1941 Average excess and average effect of a gene substitution. Annals of Eugenics **11**: 53–63.

Fisher, R. A., 1958 *The genetical theory of natural selection, 2nd revised edn*. Dover Publications, New York.

Foerster, K., T. Coulson, B. C. Sheldon, J. M. Pemberton, T. H. Clutton-Brock *et al.*, 2007 Sexually antagonistic genetic variation for fitness in red deer. Nature **447**: 1107–1110.

Fowler, K., C. Semple, N. H. Barton, and L. Partridge, 1997 Genetic variation for total fitness in *Drosophila melanogaster*. Proceedings of the Royal Society of London. Series B: Biological Sciences **264**: 191–199.

Frank, S. A., and M. Slatkin, 1992 Fisher's fundamental theorem of natural selection. Trends in Ecology & Evolution **7**: 92–95.

Gardner, M. P., K. Fowler, N. H. Barton, and L. Partridge, 2005 Genetic variation for total fitness in *Drosophila melanogaster*: complex yet replicable patterns. Genetics **169**: 1553–1571.

Gianola, D., G. de Los Campos, W. G. Hill, E. Manfredi, and R. Fernando, 2009 Additive genetic variability and the Bayesian alphabet. Genetics **183**: 347–363.

Gibbs, A. G., A. K. Chippindale, and M. R. Rose, 1997 Physiological mechanisms of evolved desiccation resistance in *Drosophila melanogaster*. Journal of Experimental Biology **200**: 1821–1832.

Godinho, D. P., I. Fragata, A. Majer, L. R. Rodrigues, and S. Magalhães, 2024 Limits to the adaptation of herbivorous spider mites to metal accumulation in homogeneous and heterogeneous environments. Journal of Evolutionary Biology : voae003.

Grafen, A., 2015 Biological fitness and the fundamental theorem of natural selection. The American Naturalist **186**: 1–14.

Gupta, V., S. Venkatesan, M. Chatterjee, Z. A. Syed, V. Nivsarkar *et al.*, 2016 No apparent cost of evolved immune response in *Drosophila melanogaster*. Evolution **70**: 934–943.

Haldane, J. B. S., 1919 The combination of linkage values, and the calculation of distances between the loci of linked factors. Journal of Genetics **8**: 299–309.

Haller, B. C., and P. W. Messer, 2023 SLiM 4: multispecies eco-evolutionary modeling. The American Naturalist **201**: E127–E139.

Hendry, A. P., D. J. Schoen, M. E. Wolak, and J. M. Reid, 2018 The contemporary evolution of fitness. Annual Review of Ecology, Evolution, and Systematics **49**: 457–476.

Hill, W., and A. Robertson, 1968 Linkage disequilibrium in finite populations. Theoretical and applied genetics **38**: 226–231.

Hsu, S.-K., W.-Y. Lai, J. Novak, F. Lehner, A. M. Jakšić *et al.*, 2024 Reproductive isolation arises during laboratory adaptation to a novel hot environment. Genome Biology **25**: 141.

Johnston, S. E., 2024 Understanding the genetic basis of variation in meiotic recombination: past, present, and future. Molecular Biology and Evolution **41**: msae112.

Joshi, A., and L. D. Mueller, 1996 Density-dependent natural selection in *Drosophila*: trade-offs between larval food acquisition and utilization. Evolutionary Ecology **10**: 463–474.

Kawecki, T. J., B. Erkosar, C. Dupuis, B. Hollis, R. C. Stillwell *et al.*, 2021 The genomic architecture of adaptation to larval malnutrition points to a trade-off with adult starvation resistance in *Drosophila*. Molecular Biology and Evolution **38**: 2732–2749.

Kelleher, J., A. M. Etheridge, and G. McVean, 2016 Efficient coalescent simulation and genealogical analysis for large sample sizes. PLoS computational biology **12**: e1004842.

Kirkpatrick, M., T. Johnson, and N. Barton, 2002 General models of multilocus evolution. Genetics **161**: 1727–1750.

Kojima, K., 1959 Role of epistasis and overdominance in stability of equilibria with selection. Proceedings of the National Academy of Sciences **45**: 984–989.

Kruuk, L. E., 2004 Estimating genetic parameters in natural populations using the 'animal model'. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences **359**: 873–890.

Kruuk, L. E., T. H. Clutton-Brock, J. Slate, J. M. Pemberton, S. Brotherstone *et al.*, 2000 Heritability of fitness in a wild mammal population. Proceedings of the National Academy of Sciences **97**: 698–703.

Kruuk, L. E., and J. D. Hadfield, 2007 How to separate genetic and environmental causes of similarity between relatives. Journal of evolutionary biology **20**: 1890–1903.

Kulbaba, M. W., S. N. Sheth, R. E. Pain, V. M. Eckhart, and R. G. Shaw, 2019 Additive genetic variance for lifetime fitness and the capacity for adaptation in an annual plant. Evolution **73**: 1746–1758.

Lee, J. J., and C. C. Chow, 2013 The causal meaning of Fisher's average effect. Genetics research **95**: 89–109.

Lynch, M., and B. Walsh, 1998 *Genetics and analysis of quantitative traits*. Sinauer Associates, Inc., Sunderland.

Manna, F., G. Martin, and T. Lenormand, 2011 Fitness landscapes: an alternative theory for the dominance of mutation. Genetics **189**: 923–937.

Martinossi-Allibert, I., U. Savković, M. Đorđević, G. Arnqvist, B. Stojković *et al.*, 2018 The consequences of sexual selection in well-adapted and maladapted populations of bean beetles. Evolution **72**: 518–530.

McCullagh, P., and J. A. Nelder, 1989 *Generalised Linear Models*. Monographs on statistics and applied probability. Chapman & Hall, Cambridge, 2^{nd} edition.

Ohta, T., and M. Kimura, 1970 Development of associative overdominance through linkage disequilibrium in finite populations. Genetics Research **16**: 165–177.

Price, G. R., 1970 Selection and covariance. Nature **227**: 520–521.

Price, G. R., 1972 Fisher's 'fundamental theorem' made clear. Annals of human genetics **36**: 129–140.

Punzalan, D., M. Delcourt, and H. Rundle, 2014 Comparing the intersex genetic correlation for fitness across novel environments in the fruit fly, *Drosophila serrata*. Heredity **112**: 143–148.

Queller, D. C., 2017 Fundamental theorems of evolution. The American Naturalist **189**: 345–353.

Reznick, D. N., and H. A. Bryga, 1996 Life-history evolution in guppies (*Poecilia reticulata*: Poeciliidae). V. genetic basis of parallelism in life histories. The American Naturalist **147**: 339–359.

Richards, F. S., 1961 A method of maximum-likelihood estimation. Journal of the Royal Statistical Society Series B: Statistical Methodology **23**: 469–475.

Robertson, A., 1966 A mathematical model of the culling process in dairy cattle. Animal Science **8**: 95–108.

Robinson, C. E., H. Thyagarajan, and A. K. Chippindale, 2023 Evolution of reproductive isolation in a long-term evolution experiment with *Drosophila melanogaster*: 30 years of divergent life-history selection. Evolution **77**: 1756–1768.

Ruzicka, F., M. S. Hill, T. M. Pennell, I. Flis, F. C. Ingleby *et al.*, 2019 Genome-wide sexually antagonistic variants reveal long-standing constraints on sexual dimorphism in fruit flies. PLoS biology **17**: e3000244.

Salson, M., M. Duranton, S. Huynh, C. Mariac, C. Tranchant-Dubreuil *et al.*, 2025 Interplay between large low-recombining regions and pseudo-overdominance in a plant genome. Nature Communications **16**: 6458.

Santiago, E., and A. Caballero, 1998 Effective size and polymorphism of linked neutral loci in populations under directional selection. Genetics **149**: 2105–2117.

Schaal, S. M., B. C. Haller, and K. E. Lotterhos, 2022 Inversion invasions: when the genetic basis of local adaptation is concentrated within inversions in the face of gene flow. Philosophical Transactions of the Royal Society B **377**: 20210200.

Schenkel, M. A., I. Pen, L. W. Beukeboom, and J.-C. Billeter, 2018 Making sense of intralocus and interlocus sexual conflict. Ecology and Evolution **8**: 13035–13050.

Searle, S. R., 2006 *Matrix algebra useful for statistics*. John Wiley & Sons.

Shaw, R., and F. Shaw, 2014 Quantitative genetic study of the adaptive process. Heredity **112**: 13–20.

Simon, A., and G. Coop, 2024 The contribution of gene flow, selection, and genetic drift to five thousand years of human allele frequency change. Proceedings of the National Academy of Sciences **121**: e2312377121.

Singh, A., A. Hasan, and A. F. Agrawal, 2023 An investigation of the sex-specific genetic architecture of fitness in *Drosophila melanogaster*. Evolution **77**: 2015–2028.

Singh, K., E. Kochar, and N. Prasad, 2015 Egg viability, mating frequency and male mating ability evolve in populations of *Drosophila melanogaster* selected for resistance to cold shock. PloS one **10**: e0129992.

Stephan, W., B. Charlesworth, and G. McVean, 1999 The effect of background selection at a single locus on weakly selected, partially linked variants. Genetics Research **73**: 133–146.

Sved, J. A., and W. G. Hill, 2018 One hundred years of linkage disequilibrium. Genetics **209**: 629–636.

Tilk, S., A. Bergland, A. Goodman, P. Schmidt, D. Petrov *et al.*, 2019 Accurate allele frequencies from ultra-low coverage pool-seq samples in evolve-and-resequence experiments. G3: Genes, Genomes, Genetics **9**: 4159–4168.

Vedder, O., J. Komdeur, M. van der Velde, E. Schut, and M. J. Magrath, 2011 Polygyny and extra-pair paternity enhance the opportunity for sexual selection in blue tits. Behavioral Ecology and Sociobiology **65**: 741–752.

Vehviläinen, H., A. Kause, C. Quinton, H. Koskinen, and T. Paananen, 2008 Survival of the currently fittest: genetics of rainbow trout survival across time and space. Genetics **180**: 507–516.

Walsh, B., 2022 How full is the evolutionary fuel tank? Science **376**: 920–921.

Wang, Y., P. McNeil, R. Abdulazeez, M. Pascual, S. E. Johnston *et al.*, 2023 Variation in mutation, recombination, and transposition rates in *Drosophila melanogaster* and *Drosophila simulans*. Genome Research **33**: 587–598.

Weir, B. S., and C. C. Cockerham, 1989 Complete characterization of disequilibrium at two loci. In M. Feldman, editor, *Mathematical Evolutionary Theory*. Princeton University Press Princeton, NJ, 86–110.

Whitlock, M. C., 2015 Modern approaches to local adaptation. The American Naturalist **186**: S1–S4.

Wright, S., 1938 Size of population and breeding structure in relation to evolution. Science **87**: 340–431.

Xiao, C., N. Bayat Fard, K. Brzezinski, R. M. Robertson, and A. K. Chippindale, 2019 Experimental evolution of response to anoxia in *Drosophila melanogaster*: recovery of locomotion following CO_2 or N_2 exposure. Journal of Experimental Biology **222**: jeb199521.

Yang, J., S. H. Lee, M. E. Goddard, and P. M. Visscher, 2011 GCTA: a tool for genome-wide complex trait analysis. The American Journal of Human Genetics **88**: 76–82.

Zeng, J., R. De Vlaming, Y. Wu, M. R. Robinson, L. R. Lloyd-Jones *et al.*, 2018 Signatures of negative selection in the genetic architecture of human complex traits. Nature genetics **50**: 746–753.

 # Figures

Figure 1: Results of simplified simulations (map length in the history phase = 0.5 morgan, map length in the experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations $(E[|\eta|]) = 0.02$, and no dominance (i.e. $\kappa = 0$)). (a) A scatter plot of estimates of V_A vs true values of V_A. The solid black line indicates the 1:1 line. The inference of V_A was obtained by modelling the mean and the (co)variance of the average effects for relative fitness as $\boldsymbol{\mu}_{\bar{\alpha}} = \beta_{\bar{\alpha}}^{(1)}(\mathbf{p}_0 - \mathbf{q}_0)$ and $\mathbf{V}_{\bar{\alpha}} = \sigma_{\bar{\alpha}}^2 \mathbf{L}_0^{p_{\bar{\alpha}}}$, respectively. (b)-(d) Histograms of the estimates of $p_{\bar{\alpha}}$, $\beta_{\bar{\alpha}}^{(1)}$, and the residual variance, respectively. The vertical red lines indicate null expectations (0 for $p_{\bar{\alpha}}$ and $\beta_{\bar{\alpha}}^{(1)}$, and 1 for the residual variance). The black dashed lines indicate the means of the respective distributions.

Simplified simulations

Figure 2: Scatter plots of estimates of V_A vs true values of V_A for simplified simulations at different levels of (a) map length in the experiment, (b) population size of each replicate population in the evolve and resequence experiment, (c) number of replicate populations, and (d) number of generations over which allele frequency changes were recorded in the experiment. In each case, other than the parameter to be varied, the other parameters were fixed at their default values (the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations ($E[|\eta|]$) = 0.02, map length in the history phase = 0.5 morgan, map length in the experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, and no dominance ($\kappa = 0$)). The solid black line indicates the 1:1 line. The coloured lines represent regression lines for estimates of V_A vs true values of V_A.

Figure 3: Results of simulations with map lengths in the experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, and different degrees of dominance: $\kappa = 0$ (green circles), $\kappa = 0.5$ (blue triangles), $\kappa = 0.75$ (magenta plus signs) and $\kappa = 0.9$ (grey squares). (a-d) are the results from the simplified simulations with the map length in the history phase = 0.5 morgan and the mean of the gamma distribution from which effect sizes for log absolute fitness are sampled for non-neutral mutations ($E[|\eta|]$) = 0.02. (a) A scatter plot of estimates of V_A vs true values of V_A. The solid black line indicates the 1:1 line. The coloured lines represent regression lines for estimates of V_A vs true values of V_A. (b-d) Estimates of model parameters $p_{\bar{\alpha}}$, $\beta_{\bar{\alpha}}^{(1)}$, and the residual variance, respectively. (e-f) Scatter plots of estimates of V_A vs true values from full simulations (with $E[|\eta|] = 0.03$) when the map length in the history phase was either 0.5 morgan (e) or 5 morgans (f).

Full simulations

Figure 4: Results of full simulations with a burn-in phase of 25,000 generations (map length in the history phase = 0.5 morgan, map length in the experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations $(E[|\eta|]) = 0.02$, and no dominance (i.e. $\kappa = 0$)). (a) A scatter plot of estimates of V_A vs true values of V_A. The solid black line indicates the 1:1 line. The inference of V_A was obtained by modelling the mean and the (co)variance of the average effects for relative fitness as $\boldsymbol{\mu}_{\bar{\alpha}} = \beta_{\bar{\alpha}}^{(1)}(\mathbf{p}_0 - \mathbf{q}_0)$ and $\mathbf{V}_{\bar{\alpha}} = \sigma_{\bar{\alpha}}^2 \mathbf{L}_0^{p_{\bar{\alpha}}}$, respectively. (b)-(d) Histograms of the estimates of $p_{\bar{\alpha}}$, $\beta_{\bar{\alpha}}^{(1)}$, and the residual variance, respectively. The vertical red lines indicate null expectations (0 for $p_{\bar{\alpha}}$ and $\beta_{\bar{\alpha}}^{(1)}$, and 1 for the residual variance). The black dashed lines indicate the means of the respective distributions.

Full simulations

Figure 5: Scatter plots of estimates of V_A vs true values of V_A for full simulations with a burn-in phase of 25,000 generations at different values of (a) map length in the experiment phase, (b) map length during the history phase, (c) the ratio of rates of beneficial and deleterious mutations in the history phase, and (d) the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations ($E[|\eta|]$). In each case, other than the parameter to be varied, the other parameters were fixed at their default values (map length in the history phase = 0.5 morgan, map length in the experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, $E[|\eta|] = 0.02$, and no dominance (i.e. $\kappa = 0$)). The solid black line indicates the 1:1 line. The coloured lines represent regression lines for estimates of V_A vs true values of V_A. The effect of analysing the standard set of simulations (Figure 4) (e) without partitioning $\tilde{\mathbf{L}}_0$ into its gametic ($\mathbf{L}_0'$) and non-gametic phase ($\mathbf{L}_0''$) components, and (f) using $N_E = N = 1,000$. Note that in (F), a large number of green circles are eclipsed by the blue triangles.

Figure 6: Scatter plots of estimates of V_A vs true values of V_A for simplified simulations (map length in the history phase $= 0.5$ morgan, map length in the experiment phase $= 2$ morgans, number of replicate populations $= 10$, population size $= 1,000$, number of generations $= 3$, the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations $(E[|\eta|]) = 0.02$, and no dominance (i.e. $\kappa = 0$)) using either exact allele frequencies in the experiment phase (green circles), or allele frequencies in the experiment phase obtained via simulated pool-seq implemented without any overdispersion in the number of reads mapping to an individual, at three different levels of coverage (expected number of reads mapping a segregating site: 1000x (magenta triangles), 500x (blue squares), and 100x (grey plus symbols)) as well as estimates obtained from simulated pool-seq implemented with overdispersion $(V_x = log(2))$ in the number of reads mapping to an individual, at 1000x coverage (orange empty boxes with crosses). Reads were modelled to be 800 base-pairs long. The solid black line indicates the 1:1 line. The coloured lines represent regression lines for estimates of V_A vs true values of V_A. Estimates of V_A were obtained by either incorporating (a) or omitting (b) the expected covariance structure due to pool-seq sampling in the models.

Figure 7: Estimates of V_A obtained using our method (g) and various applications of the B&C Approach (a–f) from simplified simulations with varying map lengths in the history phase: 0.5 morgan (green circles), 5 morgans (magenta triangles), 50 morgans (blue squares), and 100 morgans (grey plus symbols). The solid black line indicates the 1:1 line. Each application of B&C is defined in terms of which three assumptions (B, Ib, N) are required or not (designated with a '+' or '0' superscript, respectively). Assumption B is that allele frequencies are tracked at unselected loci only, Assumption Ib is that the average LD between all sites is equal to the average LD between selected and unselected sites, and Assumption N is that the covariance between projected allele frequency changes is equal to the covariance between allele frequencies changes scaled by the average projection (see Table 2 for details). Panel h is the correlation between the contribution of a selected sites to V_a and their persistent associations with neutral alleles (i.e. deviations from Assumption G in B&C)

Figure 8: Estimates of V_A obtained using our method (g) and various applications of the B&C Approach (a–f) from full simulations with varying map lengths in the history phase: 0.5 morgan (green circles), 5 morgans (magenta triangles), 50 morgans (blue squares), and 100 morgans (grey plus symbols). The solid black line indicates the 1:1 line. Each application of B&C is defined in terms of which three assumptions (B, Ib, N) are required or not (designated with a '+' or '0' superscript, respectively). Assumption B is that allele frequencies are tracked at unselected loci only, Assumption Ib is that the average LD between all sites is equal to the average LD between selected and unselected sites, and Assumption N is that the covariance between projected allele frequency changes is equal to the covariance between allele frequencies changes scaled by the average projection (see Table 2 for details). Panel h is the correlation between the contribution of a selected sites to V_a and their persistent associations with neutral alleles (i.e. deviations from Assumption G in B&C). For clarity of presentation, in (a), (c), and (e), we have restricted the range of the y axis between 0 and 0.35. As a consequence, points having an estimate of V_A above 0.35 have been excluded from the plot for 0.5 morgan (green circles).

$\|\kappa\|$	$E[\|\eta\|]$	$\mu_{ben} : \mu_{del}$	μ_{msp}	μ_{SLiM}	$n_{L_{\mathcal{S}}}$
0	0.02	0	$5.56 \times 10^{-8} - 5.56 \times 10^{-7}$	$5.56 \times 10^{-7} - 5.56 \times 10^{-6}$	$10058 - 63755$
0	0.02	0.0002	$5.56 \times 10^{-8} - 5.56 \times 10^{-7}$	$5.56 \times 10^{-7} - 5.56 \times 10^{-6}$	$9742 - 63797$
0	0.02	0.02	$2.0 \times 10^{-8} - 2.0 \times 10^{-7}$	$2.0 \times 10^{-7} - 2.0 \times 10^{-6}$	$3127 - 29999$
0	0.03	0	$3.6 \times 10^{-8} - 3.6 \times 10^{-7}$	$3.6 \times 10^{-7} - 3.6 \times 10^{-6}$	$6565 - 42097$
0	0.06	0	$1.8 \times 10^{-8} - 1.6 \times 10^{-7}$	$1.8 \times 10^{-7} - 1.6 \times 10^{-6}$	$2998 - 18011$
0.5	0.03	0	$7.2 \times 10^{-8} - 3.24 \times 10^{-7}$	$7.2 \times 10^{-7} - 3.24 \times 10^{-6}$	$13738 - 49299$
0.75	0.03	0	$7.5 \times 10^{-8} - 2.07 \times 10^{-7}$	$7.5 \times 10^{-7} - 2.07 \times 10^{-6}$	$16870 - 38695$

Table 1: The ranges for the rates for non-neutral mutations used in the history phase of the full simulations implemented in msprime (μ_{msp}) (the coalescent simulation) and SLiM (μ_{SLiM}) (the forward simulation of the history), along with the ranges for the resulting number of non-neutral sites segregating in the population at the end of the history phase ($n_{L_{\mathcal{S}}}$) in simulations with different values of the absolute degree of dominance ($\|\kappa\|$), the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations ($E[\|\eta\|]$), and the ratio of the rate of beneficial mutations to the rate of deleterious mutations in the history phase ($\mu_{ben} : \mu_{del}$). Note that the ranges for the number of selected loci, $n_{L_{\mathcal{S}}}$, are only shown for the simulations where the map length in the history phase was 0.5 morgan.

Approach	Δp	Average LD between	$Cov(\Delta\mathbf{p}_m, \Delta\mathbf{p}_n)$
$B^+Ib^+N^+$	all sites	all sites	divided throughout by $(\bar{p}_m\bar{q}_m + \bar{p}_n\bar{q}_n)/2$
$B^0Ib^+N^+$	neutral sites	all sites	divided throughout by $(\bar{p}_m\bar{q}_m + \bar{p}_n\bar{q}_n)/2$
$B^0Ib^0N^+$	neutral sites	selected & neutral sites	divided throughout by $(\bar{p}_m\bar{q}_m + \bar{p}_n\bar{q}_n)/2$
$B^+Ib^+N^0$	all sites	all sites	replaced by $Cov(\Delta\overrightarrow{\mathbf{p}_m}, \Delta\overrightarrow{\mathbf{p}_n})$
$B^0Ib^+N^0$	neutral sites	all sites	replaced by $Cov(\Delta\overrightarrow{\mathbf{p}_m}, \Delta\overrightarrow{\mathbf{p}_n})$
$B^0Ib^0N^0$	neutral sites	selected & neutral sites	replaced by $Cov(\Delta\overrightarrow{\mathbf{p}_m}, \Delta\overrightarrow{\mathbf{p}_n})$

Table 2: We applied B&C's method to our simulations using six different approaches. The requirement of Assumptions B, Ib, and N is indicated using superscripts ("$+$" when required and "0" when not required). We used allele frequency change ($\Delta\mathbf{p}$) at either all segregating sites (Assumption B required) or at neutral segregating sites only (Assumption B not required). We computed average LD using either the LD between all segregating sites (Assumption Ib required) or the LD between selected and neutral sites (Assumption Ib not required). We used either the (co)variance in $\Delta\mathbf{p}$ divided throughout by twice the average genetic diversity (Assumption N required) or the (co)variance in weighted $\Delta\mathbf{p}$ (i.e. $\Delta\overrightarrow{\mathbf{p}}$) where the weights are the square roots of twice the genetic diversity at each site (Assumption N not required).

Symbol	Description
$\mathbf{B}$	Diagonal matrix of standard deviations for the c's
$c_{k,i}$	Number of reference alleles at locus i for individual k divided by 2.
$C_A(t \to \tau)$	Additive genetic covariance for relative fitness between generation t and τ for a population with genetic structure equal to that in generation t.
$C_a(t \to \tau)$	Additive genic covariance for relative fitness between generation t and τ for a population with genetic structure equal to that in generation t.
$\mathbf{D}_2$	Diagonal matrix of square-rooted eigenvalues of $\boldsymbol{\mathcal{D}}$
$\boldsymbol{\mathcal{D}}_m$	Matrix that gives the covariance in allele frequency changes between generation t_m and τ_m in replicate m due to drift.
$\boldsymbol{\mathcal{E}}_m$	Matrix that gives the covariance in allele frequency change estimation errors in replicate m.
$\mathbf{F}_{t\tau}$	$\mathbf{W}_{t\tau}$ but with the diagonals set to zero.
g	Total branch length of the tree sequence recorded in the history phase of the simulations
$\mathbf{H}_{t\tau}$	$\mathbf{W}_{t\tau}$ but with the off-diagonals set to zero.
$\mathbf{L}_{t,m}$	Covariance matrix of c's at time t in replicate m.
$\mathbf{L}'_{t,m}$	Covariance matrix of c's at time t in replicate m due to being on the same gametic contribution.
$\mathbf{L}''_{t,m}$	Covariance matrix of c's at time t in replicate m due to being on different gametic contributions.
$\Delta\mathbf{L}'_{t,m}$	The stochastic change in $\mathbf{L}'$ between time zero and t in replicate m.
$\tilde{\mathbf{L}}_0$	Weighted sum of $\mathbf{L}'_0$ and $\mathbf{L}''_0$ with weights for element ij being 1 and $z_{0,ij}/\zeta_{0,ij}$ respectively.
$\boldsymbol{\mathcal{L}}_{t,m}$	A matrix that gives, when post-multiplied by $\boldsymbol{\alpha}$, the predictable change in allele frequency due to selection between generation t_m and $t_m + 1$ in replicate m.
$\boldsymbol{\mathcal{L}}_m$	A matrix that gives, when post-multiplied by $\boldsymbol{\alpha}$, the predictable change in allele frequency due to selection between generation t_m and τ_m in replicate m.
$\mathbf{M}_{t,m}$	Matrix of weights for $\tilde{\mathbf{L}}_0$ that gives the covariance in allele frequency changes due to drift between time $t > 0$ and $t + 1$ in replicate m.
$\mathbf{M}_m$	Matrix of weights for $\tilde{\mathbf{L}}_0$ that gives the covariance in allele frequency changes due to drift between time $t_m > 0$ and τ_m in replicate m
n_L	Number of loci
$n_{L_{\mathcal{S}}}$	Number of non-neutral segregating sites
$N_{t,m}$	Census population size at time t in replicate m

$N_{e_{t,m}}$	Variance effective population size at time t in replicate m
$N_{E_{t,m}}$	Variance effective population size at time t in replicate m ignoring the impact of linked-selection
$\mathbf{N}_{t,m}$	Matrix of weights for $\tilde{\mathbf{L}}_0$ that gives the expected $\mathbf{L}$ at time $t > 0$ in replicate m.
$\mathbf{N}_m$	Matrix of weights for $\tilde{\mathbf{L}}_0$ that gives the sum of the expected $\mathbf{L}$ from time $t_m > 0$ to τ_m in replicate m.
$\mathcal{N}$	Used as a subscript to indicate the set of neutral loci.
O_{ij}	Number of pool-seq reads spanning sites i and j. If $i = j$, O_{ij} is the coverage.
$p_{\bar{\alpha}}$	Parameter that takes $\mathbf{L}_0$ to some power
$\mathbf{p}_{t,m}$	Vector of reference allele frequencies at time t in replicate m.
$\Delta\mathbf{p}_{t,m}$	Vector of reference allele frequency changes between time t and $t+1$ in replicate m.
$\Delta\mathbf{p}_m$	Vector of reference allele frequency changes between time t_m and τ_m in replicate m.
$\mathbf{P}$	Projection matrix for allele frequencies.
$\mathbf{q}_{t,m}$	Vector of alternate allele frequencies at time t in replicate m.
$\mathbf{Q}_m$	Matrix that is proportional to the covariance in allele frequency change estimation errors in replicate m.
r_{ij}	Recombination rate between loci i and j.
$R_{t,ij}$	Correlation in allele count between locus i and j at time t (Note the use of the uppercase to distinguish from the recombination rate r).
$\mathbf{R}_+$	Matrix of recombination probabilities.
$\mathbf{R}_-$	Matrix of non-recombination probabilities.
$\mathbf{R}$	Correlation matrix of the c's.
$\mathbf{S}_{\bar{\alpha}}$	Sampling covariance matrix for the parameters of the regression on the mean average effects, $\boldsymbol{\beta}_{\bar{\alpha}}$.
$\mathcal{S}$	Used as a subscript to indicate the set of selected loci.
t_m	Time at which allele frequencies are first measured in replicate m.
$\mathbf{U}$	Eigenvectors of $\mathbf{L}_0$.
$\mathbf{U_L}$	Eigenvectors of $\mathbf{L}_0$ with non-zero eigenvalues.
$\mathbf{U}_2$	Eigenvectors of $\boldsymbol{\mathcal{D}}$
$\boldsymbol{\mathcal{U}}_m$	Matrix that gives the covariance in allele frequency changes between generation t_m and τ_m in replicate m due to the unpredictable response to selection.
$V_a(t)$	Additive genic variance for relative fitness at time t
$V_A(t)$	Additive genetic variance for relative fitness at time t

$V_{\bar{A}}(t)$	Additive genetic covariance between replicate/time-points for relative fitness in a population with genotypic composition equal to that at time t.
V_x	Variance of the normal distribution from which $log(\lambda_x)$'s are sampled
$\mathbf{V}_{\bar{\alpha}}$	Covariance matrix for the mean average effects.
w	Relative fitness
W	Absolute fitness
$\mathbf{W}_{t\tau}$	A matrix with the ij^{th} element equal to $R_{t,ji}R_{\tau,ki}(b_{\tau,i}/b_{t,i})$
$\mathbf{X}$	Design matrix for the regression of the mean average effects on $\mathbf{p}_0 - \mathbf{q}_0$.
$y_{k,i}$	Genotypic contribution made by locus i to the log absolute fitness ($log(W)$) of individual k
Y_k	Genotypic value of log fitness for individual k
$z_{t,ij}$	$(1 - r_{t,ij})(1 - \frac{1}{2N_{e_t}})$
$\boldsymbol{\alpha}_{t,m}$	Vector of average effects for relative fitness at time t in replicate m.
$\bar{\boldsymbol{\alpha}}$	Vector of mean average effects for relative fitness.
$\Delta\alpha_{t,m}$	Vector of deviations of the average effects for relative fitness at time t in replicate m from the global mean.
$\beta_{\bar{\alpha}}^{(1)}$	Slope of the regression of the mean average effects on $\mathbf{p}_0 - \mathbf{q}_0$.
$\zeta_{t,ij}$	$r_{t,ij}(1 - \frac{1}{2N_{e_t}})$
η_i	Difference between the genotypic contributions to $log(W)$ by the reference and non-reference homozygotes at locus i
$\eta_i^{(a)}$	Average effect for $log(W)$ at locus i
η_{scale}	Scale of the gamma distribution from which $\eta's$ were sampled in the history phase of the simulations
κ_i	Degree of dominance at locus i for log absolute fitness.
$\lambda_{x,k}$	Mean of the Poisson distribution from which the number of reads mapping to individual k are sampled while simulating pool-seq
μ_{msp}	Mutation rate for non-neutral mutations in the coalescent part of the history phase of the simulations
μ_{SLiM}	Mutation rate for non-neutral mutations in the forward part of the history phase of the simulations
μ_x	Mean of the normal distribution from which $log(\lambda_x)$'s are sampled
$\boldsymbol{\mu}_{\bar{\alpha}}$	Vector of expected values for the mean average effects.
$\sigma_{\bar{\alpha}}^2$	Proportionality constant that relates $\mathbf{V}_{\bar{\alpha}}$ to $\mathbf{L}_0^{p_{\bar{\alpha}}}$.
σ_o^2	Parameter for scaling sampling (co)variances for allele frequencies: values greater than 1 indicate overdispersion.
τ_m	Time at which allele frequencies are finally measured in replicate m.

$\phi_{t,\tau}$	The ratio of genetic diversity in generation τ to genetic diversity in generation t assumed constant across all selected loci.
$\rightarrow$	Used above a symbol to indicate it is on the projected space.
$\frown$	Used above a symbol to indicate an estimate.

Table 3: Notation

Supplementary information for 'Estimating the additive genetic variance for relative fitness from changes in allele frequency'

Manas Geeta Arun[1,*], Aidan Angus-Henry[1,2], Darren J. Obbard[1] and Jarrod D. Hadfield[1]

[1]Institute of Ecology and Evolution, The University of Edinburgh, Ashworth Laboratories Charlotte Auerbach Road, Edinburgh, EH9 3FL, United Kingdom.

[2]Charité - Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.

[*]Corresponding author: manas.geetaarun@ed.ac.uk

S1 The dynamics of L under drift and recombination.

As outlined in the section 'Extending our approach to practical situations' our goal is to infer the additive genetic variance for relative fitness in the base population $(V_{\bar{A}}(0))$ using genome-wide allele frequency changes between generations t and τ. In order to derive the expected allele frequency change due to selection over this time period, we first derive the expected dynamics of $\mathbf{L}$ under drift and recombination.

The matrix $\mathbf{L}$ is a covariance matrix whose elements are proportional to the genotypic linkage-disequilbria (off-diagonals) or variance in genotypic allele frequencies (diagonals) at the start of a generation, before selection has acted. We can decompose $\mathbf{L}$ into $\mathbf{L}'$ and $\mathbf{L}''$ following the notation of Buffalo and Coop (2019) where $\mathbf{L}'$ represents the (co)variances that arise due to alleles in the same gamete and $\mathbf{L}''$ represents the (co)variances that arise due to alleles in the different gametic contributions of a genotype. Note that in Santiago and Caballero (1998) the primes have a subtly different meaning after the initial generation, as the double prime in following generations designates gametic phase disequilibrium due to recombination and nongametic phase disequilibrium in the initial generation. The elements of $\mathbf{L}'$ and $\mathbf{L}''$ have direct correspondences with genetic diversities and additive measures of disequilibria. Under the notation of Weir and Cockerham (1989) (see also Bulmer (1980), Chapter 12), the diagonal elements of $\mathbf{L}'$ are half the gametic genetic diversities $(\pi_i = p_i(1 - p_i) = 2L'_{ii})$, and the off-diagonals are half the gametic-phase disequilibria $(D_{ij} = 2L'_{ij})$. Note that Weir and Cockerham (1989) uses π to denote $p_i(1 - p_i)$ rather than the more usual (and less natural) $2p_i(1 - p_i)$. The diagonal elements of

i

$\mathbf{L}''$ are half the additive coefficients of Hardy Weinberg disequilibria ($D_i = 2L''_{ii}$), and the off-diagonals are half the nongametic-phase disequilibria ($D_{i/j} = 2L''_{ij}$).

To see these correspondences, imagine two bi-allelic loci, i and j, with reference/alternate alleles A/a and B/b, respectively. There are four possible gametic haplotypes, and we can denote the frequency of haplotype AB as p_{AB} in the gametes and the frequency of allele A as p_A. The proportion of copies of the reference allele at locus i in a randomly chosen individual, c_i, can be decomposed into into the sum of maternal and paternal contribution: $c_i = m_i + f_i$ where m_i (or f_i) takes the value $1/2$ if the mother (or the father) contributed a reference allele and 0 if not. Then,

$$
\begin{aligned}
L_{ij} &= COV(c_i, c_j) \\
&= COV(m_i + f_i, m_j + f_j) \\
&= COV(m_i, m_j) + COV(f_i, f_j) + COV(m_i, f_j) + COV(f_i, m_j)
\end{aligned} \tag{S1}
$$

Assuming haplotype frequencies are identical in male and female gametes we get

$$
\begin{aligned}
L_{ij} &= (p_{AB} - p_A p_B)/4 + (p_{AB} - p_A p_B)/4 + (p_{A/B} - p_A p_B)/4 + (p_{A/B} - p_A p_B)/4 \\
&= (p_{AB} - p_A p_B)/2 + (p_{A/B} - p_A p_B)/2 \\
&= D_{ij}/2 + D_{i/j}/2
\end{aligned} \tag{S2}
$$

where $p_{A/B}$ is the frequency of zygotes that have an A from their mother and a B from their father, or vice versa. When $i = j$,

$$
\begin{aligned}
L_{ii} &= COV(c_i, c_i) \\
&= (p_A^2 - p_A)/2 + (p_{A/A} - p_A^2)/2 \\
&= \pi_i/2 + D_i/2
\end{aligned} \tag{S3}
$$

where $p_{A/A}$ is the frequency of A homozygotes. The term D_i is an alternative, additive, measure of deviation from Hardy-Weinberg Equilibrium than the more commonly used inbreeding coefficient, F.

To derive expressions for the dynamics of $\mathbf{L}'$ and $\mathbf{L}''$, note that $\mathbf{L}''$ is generated anew each generation and under random mating has zero expectation such that

$$
L''_{t+1,ij} = e''_{t+1,ij} \tag{S4}
$$

where $e''_{t+1,ij}$ is a stochastic term with mean zero and variance given in Weir (1996). The elements of $\mathbf{L}'$ under recombination and drift are (Hill and Robertson, 1968; Santiago and Caballero, 1998)

ii

$$L'_{t+1,ij} = \left(1 - \frac{1}{2N_{e_t}}\right)\left((1 - r_{t,ij})L'_{t,ij} + r_{t,ij}L''_{t,ij}\right) + e'_{t+1,ij}$$
$$= z_{t,ij}L'_{t,ij} + \zeta_{t,ij}L''_{t,ij} + e'_{t+1,ij} \tag{S5}$$

where $z_{t,ij} = (1 - r_{t,ij})(1 - \frac{1}{2N_{e_t}})$ and $\zeta_{t,ij} = r_{t,ij}(1 - \frac{1}{2N_{e_t}})$, with $r_{t,ij}$ being the recombination rate between locus i and j in generation t, and N_{e_t} the effective population size in generation t. In the absence of selection, the stochastic terms, $e'_{t+1,ij}$, have zero mean and are uncorrelated over time, with variances given by Ohta and Kimura (1969, Eq 25, although those variances must be divided by 4 here since we are working with the frequency of alleles in individuals rather than gametes). The covariances between, for example, $e'_{t+1,ij}$ and $e'_{t+1,kl}$ are not given in Ohta and Kimura (1969), and may be unknown. Using the above results we can derive a recursion for L' at some arbitrary time t:

$$L'_{1,ij} = z_{0,ij}L'_{0,ij} + \zeta_{0,ij}L''_{0,ij} + e'_{1,ij}$$

$$L'_{2,ij} = z_{1,ij}(z_{0,ij}L'_{0,ij} + \zeta_{0,ij}L''_{0,ij} + e'_{1,ij}) + \zeta_{1,ij}L''_{1,ij} + e'_{2,ij}$$

$$L'_{3,ij} = z_{2,ij}(z_{1,ij}(z_{0,ij}L'_{0,ij} + \zeta_{0,ij}L''_{0,ij} + e'_{1,ij}) + \zeta_{1,ij}L''_{1,ij} + e'_{2,ij}) + \zeta_{2,ij}L''_{2,ij} + e'_{3,ij}$$

$$L'_{3,ij} = z_{2,ij}z_{1,ij}z_{0,ij}L'_{0,ij} + z_{2,ij}z_{1,ij}\zeta_{0,ij}L''_{0,ij} + z_{2,ij}z_{1,ij}e'_{1,ij} + z_{2,ij}\zeta_{1,ij}L''_{1,ij} + z_{2,ij}e'_{2,ij} + \zeta_{2,ij}L''_{2,ij} + e'_{3,ij}$$

$$\cdots = \cdots$$

$$L'_{t,ij} = L'_{0,ij}\prod_{k=0}^{t-1}z_{k,ij} + \sum_{k=0}^{t-1}\zeta_{k,ij}L''_{k,ij}\prod_{u=k+1}^{t-1}z_{u,ij} + \sum_{k=1}^{t-1}e'_{k,ij}\prod_{u=k}^{t-1}z_{u,ij} + e'_{t,ij}$$

$$L'_{t,ij} = L'_{0,ij}\prod_{k=0}^{t-1}z_{k,ij} + \zeta_{0,ij}L''_{0,ij}\prod_{k=1}^{t-1}z_{k,ij} + \sum_{k=1}^{t-1}\zeta_{k,ij}L''_{k,ij}\prod_{u=k+1}^{t-1}z_{u,ij} + \sum_{k=1}^{t-1}e'_{k,ij}\prod_{u=k}^{t-1}z_{u,ij} + e'_{t,ij}$$

$$L'_{t,ij} = L'_{0,ij}\prod_{k=0}^{t-1}z_{k,ij} + \frac{\zeta_{0,ij}}{z_{0,ij}}L''_{0,ij}\prod_{k=0}^{t-1}z_{k,ij} + \sum_{k=1}^{t-1}\frac{\zeta_{k,ij}}{z_{k,ij}}L''_{k,ij}\prod_{u=k}^{t-1}z_{u,ij} + \sum_{k=1}^{t-1}e'_{k,ij}\prod_{u=k}^{t-1}z_{u,ij} + e'_{t,ij}$$

$$L'_{t,ij} = \left(L'_{0,ij} + \frac{\zeta_{0,ij}}{z_{0,ij}}L''_{0,ij}\right)\prod_{k=0}^{t-1}z_{k,ij} + \sum_{k=1}^{t-1}\left(e'_{k,ij} + \frac{\zeta_{k,ij}}{z_{k,ij}}L''_{k,ij}\right)\prod_{u=k}^{t-1}z_{u,ij} + e'_{t,ij} \tag{S6}$$

Having the matrix $\mathbf{N}_t$ with the ij^{th} element $\prod_{k=0}^{t-1}z_{k,ij}$, we have for $t > 0$

$$\mathbf{L}_t = \mathbf{N}_t \circ \tilde{\mathbf{L}}_0 + \Delta\mathbf{L}'_t + \mathbf{L}''_t \tag{S7}$$

where $\circ$ is the Hadamard (i.e. lement-wise) product. The first term is the expected $\mathbf{L}$ (and $\mathbf{L}'$) in generation t, conditional on the genotypic composition of the population

iii

in generation 0, where $\tilde{\mathbf{L}}_0$ is the sum of $\mathbf{L}_0^{'}$ and $\mathbf{L}_0^{''}$ with the elements of the latter weighted by $\zeta_{0,ij}/z_{0,ij} = r_{0,ij}/(1 - r_{0,ij})$. $\Delta\mathbf{L}_t^{'}$ is a matrix with elements equal to the sum of the stochastic terms involving e in Equation S6 and represents stochastic changes in $\mathbf{L}^{'}$ from generation 0 to generation t. $\mathbf{L}_t^{''}$ are the new nongametic-phase disequilibria that arise in generation t. Note that if replicate populations are initiated from the offspring of Generation 0, then the stochastic terms, $\Delta\mathbf{L}_t^{'}$ and $\mathbf{L}_t^{''}$, will be unique in each replicate although the deterministic part is shared. If recombination rates are constant in time $r_{t,ij} = r_{ij}$ then $\zeta_{0,ij}/z_{0,ij} = r_{ij}/(1 - r_{ij})$ and $\prod_{k=0}^{t-1} z_{k,ij} = (1 - r_{ij})^t \prod_{k=0}^{t-1}(1 - \frac{1}{2N_{e_k}})$. When population sizes are also constant then $\prod_{k=0}^{t-1} z_{k,ij} = (1 - r_{ij})^t(1 - \frac{1}{2N_e})^t$.

S2 The mean and (co)variance of allele frequency changes.

Given our model for the dynamics of $\mathbf{L}$ (See Supplementary information S1) we can derive the mean and (co)variance of the allele-frequency changes within a replicate. Here expectations and variances are taken with respect to the evolutionary process, which not only includes the effects of drift ($\underset{D}{\Delta}\mathbf{p}$) but also stochastic changes in $\mathbf{L}$ and $\boldsymbol{\alpha}$. The change in allele frequency in replicate m is

$$
\begin{aligned}
\Delta\mathbf{p}_m &= \sum_{t=t_m}^{\tau_m-1}\left(\mathbf{L}_t\boldsymbol{\alpha} + \underset{D}{\Delta}\mathbf{p}_{t,m}\right) \\
&= \sum_{t=t_m}^{\tau_m-1}\left((\mathbf{N}_{t,m}\circ\tilde{\mathbf{L}}_0)\bar{\boldsymbol{\alpha}} + \Delta\mathbf{L}_{t,m}\bar{\boldsymbol{\alpha}} + (\mathbf{N}_{t,m}\circ\tilde{\mathbf{L}}_0)\Delta\boldsymbol{\alpha}_{t,m} + \Delta\mathbf{L}_{t,m}\Delta\boldsymbol{\alpha}_{t,m} + \underset{D}{\Delta}\mathbf{p}_{t,m}\right)
\end{aligned}
$$
(S8)

where $\Delta\mathbf{L}_{t,m} = \Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m}$, $\bar{\boldsymbol{\alpha}}$ is a vector of mean average effects and $\Delta\boldsymbol{\alpha}_{t,m}$ a vector of deviations specific to time t and replicate m. By definition, drift is non-directional and so $E\left[\underset{D}{\Delta}\mathbf{p}_{t,m}\right] = \mathbf{0}$. We also assume that the unpredictable response to selection has zero expectation which requires that deviations in $\mathbf{L}$ and $\boldsymbol{\alpha}$ from their predicted means are zero on average ($E[\Delta\mathbf{L}] = \mathbf{0}$ and $E[\Delta\boldsymbol{\alpha}] = \mathbf{0}$) and there is no correlation between them ($COV(\Delta\mathbf{L}, \Delta\boldsymbol{\alpha}) = 0$). If the genetic architecture of fitness is sufficiently polygenic, selection-induced changes in $\mathbf{L}$ may be negligible. Similarly, since the expected change in $\mathbf{L}$ is slow, the induced change in $\boldsymbol{\alpha}$ in the presence of non-additive gene action should be minor. Making these assumptions we have:

$$
\begin{aligned}
E\left[\Delta\mathbf{p}_m\right] &= \sum_{t=t_m}^{\tau_m-1}(\mathbf{N}_{t,m}\circ\tilde{\mathbf{L}}_0)\bar{\boldsymbol{\alpha}} \\
&= \left(\tilde{\mathbf{L}}_0 \circ \sum_{t=t_m}^{\tau_m-1}\mathbf{N}_{t,m}\right)\bar{\boldsymbol{\alpha}} \\
&= \left(\tilde{\mathbf{L}}_0 \circ \mathbf{N}^{(m)}\right)\bar{\boldsymbol{\alpha}} \\
&= \boldsymbol{\mathcal{L}}_m\bar{\boldsymbol{\alpha}}
\end{aligned}
$$
(S9)

where $\mathbf{N}^{(m)} = \sum_{t=t_m}^{\tau_m-1}\mathbf{N}_{t,m}$ and $\boldsymbol{\mathcal{L}}_m = \tilde{\mathbf{L}}_0\circ\mathbf{N}^{(m)}$. The within-replicate (co)variances are more challenging to derive. The variance due to the unpredictable selection at time t in replicate m is

$$
\begin{aligned}
VAR\left(\underset{U}{\Delta}\mathbf{p}_{t,m}\right) &= VAR\left((\Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m})\bar{\boldsymbol{\alpha}} + \boldsymbol{\mathcal{L}}_{t,m}\Delta\boldsymbol{\alpha}_{t,m} + (\Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m})\Delta\boldsymbol{\alpha}_{t,m}\right) \\
&= VAR(\Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m})\bar{\boldsymbol{\alpha}}\bar{\boldsymbol{\alpha}}^\top + \boldsymbol{\mathcal{L}}_{t,m}VAR(\Delta\boldsymbol{\alpha}_{t,m})\boldsymbol{\mathcal{L}}_{t,m} + VAR(\Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m})VAR(\Delta\boldsymbol{\alpha}_{t,m})
\end{aligned}
$$
(S10)

In addition, covariances between the unpredictable response at different time points within a replicate will be non-zero because $\Delta\mathbf{L}'_t$ is the sum of stochastic

v

changes from generation 1 to t and so $COV(\Delta\mathbf{L}'_{t_1,m}, \Delta\mathbf{L}'_{t_2,m}) = VAR(\Delta\mathbf{L}'_{t_1,m})$ when $t_1 \le t_2$, although $COV(\mathbf{L}''_{t_1,m}, \mathbf{L}''_{t_2,m}) = 0$. This gives

$$COV\left(\underset{U}{\Delta}\mathbf{p}_{t_1,m}, \underset{U}{\Delta}\mathbf{p}_{t_2,m}\right) = VAR(\Delta\mathbf{L}'_{t,m})\bar{\boldsymbol{\alpha}}\bar{\boldsymbol{\alpha}}^\top \tag{S11}$$

when $t_1 \le t_2$. Consequently, the total variance in the unpredictable response to selection will be

$$VAR\left(\underset{U}{\Delta}\mathbf{p}_m\right) = \sum_{t=t_m}^{\tau_m-1}(2(\tau_m - t) + 1)VAR(\Delta\mathbf{L}'_{t,m})\bar{\boldsymbol{\alpha}}\bar{\boldsymbol{\alpha}}^\top$$
$$+ \sum_{t=t_m}^{\tau_m-1}\left(VAR(\mathbf{L}''_{t,m})\bar{\boldsymbol{\alpha}}\bar{\boldsymbol{\alpha}}^\top + \boldsymbol{\mathcal{L}}_{t,m}VAR(\Delta\boldsymbol{\alpha}_{t,m})\boldsymbol{\mathcal{L}}_{t,m} + VAR(\Delta\mathbf{L}'_{t,m} + \mathbf{L}''_{t,m})VAR(\Delta\boldsymbol{\alpha}_{t,m})\right) \tag{S12}$$

and cannot be simplified - we simply refer to it as $\boldsymbol{\mathcal{U}}_m$.

For the drift (co)variances, note that under random mating, haplotypes are drawn from a multinomial with $2N$ trials. Using AB, Ab, AB and aB to denote the *number* of each haplotypes in the gamete pool, the number of mn haplotypes has variance $2Np_{mn}(1 - p_{mn})$ and the covariance in the numbers of mn and op haplotypes is $-2Np_{mn}p_{op}$ where p_{mn} is the frequency of the mn halpotype in the gamete pool (i.e the parental haplotype frequencies modified by recombination). The covariance in the number of A and B alleles sampled, is therefore

$$
\begin{aligned}
COV(AB + Ab, AB + aB) =\ & COV(AB, AB) + COV(AB, aB) \\
& + COV(Ab, AB) + COV(Ab, aB) \\
=\ & 2N\left[p_{AB}(1 - p_{AB}) - p_{AB}p_{aB} - p_{Ab}p_{AB} - p_{Ab}p_{aB}\right] \\
=\ & 2N\left[p_{AB}(1 - p_{AB} - p_{aB}) - p_{Ab}(p_{AB} + p_{aB})\right] \\
=\ & 2N\left[p_{AB}(1 - p_B) - p_{Ab}p_B\right] \\
=\ & 2N\left[p_{AB} - (p_{Ab} + p_{AB})p_B\right] \\
=\ & 2N\left[p_{AB} - p_A p_B\right] \\
=\ & 4N\bar{L}'_{ij}
\end{aligned}
\tag{S13}
$$

where $\bar{L}'_{ij}$ is the gametic-phase linkage disequilibria that would be achieved in an infinite population. We can divide through by $(1/2N)^2$ to obtain the drift covariance in frequency (rather than counts) as $\bar{L}'_{ij}/N$ (i.e the drift (co)variances in a allele frequency from generation t to $t+1$ are proportional to the gametic-phase disequilibria in generation $t + 1$ that would be achieved in an infinite population conditional on the genotypic composition of the population in generation t). This recovers the well known result for the drift variance in allele frequency: $\bar{L}'_{ii}/N = p_i(1 - p_i)/2N$. We can replace the census population size, N, with the effective population size N_E, since this approximates the sampling of genotypes in non-idealised populations well

(Ethier and Nagylaki, 1980). However, note that N_E differs from N_e in that it does include the impact of linked selection since this is conditioned on in the expectation, $E[\Delta \mathbf{p}]$. In matrix terms (since $\bar{L}'_{t+1,ij} = L'_{t,ij}(1 - r_{ij}) + L''_{t,ij}r_{ij}$):

$$
\begin{aligned}
VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right) &= \tfrac{1}{N_{E_{t,m}}} \bar{\mathbf{L}}'_{t+1,m} \\
VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right) &= \tfrac{1}{N_{E_{t,m}}} \left[\mathbf{R}_- \circ \mathbf{L}'_{t,m} + \mathbf{R}_+ \circ \mathbf{L}''_{t,m}\right] \\
VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right) &= \tfrac{1}{N_{E_{t,m}}} \left[\mathbf{R}_- \circ \left(\mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0 + \Delta \mathbf{L}'_{t,m}\right) + \mathbf{R}_+ \circ \mathbf{L}''_{t,m}\right] \\
VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right) &= \tfrac{1}{N_{E_{t,m}}} \left[\mathbf{R}_- \circ \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0 + \mathbf{R}_- \circ \Delta \mathbf{L}'_{t,m} + \mathbf{R}_+ \circ \mathbf{L}''_{t,m}\right]
\end{aligned}
\tag{S14}
$$

where $\mathbf{R}_+$ and $\mathbf{R}_-$ are matrices with the ij^{th} elements being r_{ij} and $1 - r_{ij}$ respectively. The expected drift terms (averaged over the evolutionary stochasticity in $\mathbf{L}$) is therefore:

$$
E\left[VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right)\right] = \tfrac{1}{N_{E_{t,m}}} \left(\mathbf{R}_- \circ \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0\right)
\tag{S15}
$$

We define a new matrix $\mathbf{M}_{t,m}$ with the ij^{th} element being $(1 - r_{ij})/N_{E_{t,m}}$ $\left(\mathbf{M}_{t,m} = \tfrac{1}{N_{E_{t,m}}}\mathbf{R}_-\right)$ to give

$$
E\left[VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right)\right] = \mathbf{M}_{t,m} \circ \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0
\tag{S16}
$$

Since the drift terms are independent, this gives:

$$
\begin{aligned}
E\left[VAR\left(\underset{D}{\Delta}\mathbf{p}_m\right)\right] &= \sum_{t=t_m}^{\tau_m-1} E\left[VAR\left(\underset{D}{\Delta}\mathbf{p}_{t,m}\right)\right] \\
&= \sum_{t=t_m}^{\tau_m-1} \tfrac{1}{N_{E_{t,m}}} \left(\mathbf{R}_- \circ \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0\right) \\
&= \tilde{\mathbf{L}}_0 \circ \sum_{t=t_m}^{\tau_m-1} \mathbf{M}_{t,m} \circ \mathbf{N}_{t,m} \\
&= \tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)}
\end{aligned}
\tag{S17}
$$

where $\mathbf{M}^{(m)} = \sum_{t=t_m}^{\tau_m-1} \mathbf{M}_{t,m} \circ \mathbf{N}_{t,m}$ and is null when $t_m = 0$ and $\tau_m = 1$. Note the ij^{th} element of $\mathbf{M}_{t,m} \circ \mathbf{N}_{t,m} = (1 - r_{ij})^{t+1} \tfrac{1}{N_{E_t}} \prod_{k=0}^{t-1}(1 - \tfrac{1}{2N_{e_{k,m}}})$ which reduces to $(1 - r_{ij})^{t+1} \tfrac{1}{N_{e_m}}(1 - \tfrac{1}{2N_{e_m}})^t$ with constant population size and $N_e = N_E$. $\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)}$ is denoted as $\mathcal{D}_m$ in the main text. It is not clear how large the evolutionary variance in $VAR\left(\underset{D}{\Delta}\mathbf{p}_m\right)$ is, and whether this would need to be accommodated.

When the vector of average effects, $\boldsymbol{\alpha}$, is constant in time and across replicates, $V_A(0) = \boldsymbol{\alpha}\mathbf{L}_0\boldsymbol{\alpha}^\top = \bar{\boldsymbol{\alpha}}\mathbf{L}_0\bar{\boldsymbol{\alpha}}^\top$. However, when they do vary, this equality does not hold and so we use the notation $V_{\bar{A}}(0)$ to distinguish $\bar{\boldsymbol{\alpha}}\mathbf{L}_0\bar{\boldsymbol{\alpha}}^\top$ from $V_A(0)$. We claim that $V_{\bar{A}}(0)$ can be interpreted as the genetic covariance in fitness between time-replicate combinations for a population with a genotypic composition identical to the base population: it is the expected covariance in breeding value of an individual taken from the base population and (hypothetically) raised in two different replicates at two different times.

To see this, we switch from our original definition of $\boldsymbol{\alpha}_{t,m}$ as being equal to $\bar{\boldsymbol{\alpha}} + \Delta\boldsymbol{\alpha}_{t,m}$ and with some abuse of notation use the form $\bar{\boldsymbol{\alpha}} + \Delta\boldsymbol{\alpha}_t + \Delta\boldsymbol{\alpha}_m + \Delta\boldsymbol{\alpha}_{t,m}$ where $\Delta\boldsymbol{\alpha}_t$ is the deviation of the average effects (averaged over replicates) at time t and $\Delta\boldsymbol{\alpha}_m$ is the deviation of the average effects (averaged over time) in replicate m, leaving $\Delta\boldsymbol{\alpha}_{t,m}$ as the 'residual' deviation in $\boldsymbol{\alpha}$. By definition, the terms $\Delta\boldsymbol{\alpha}_t$, $\Delta\boldsymbol{\alpha}_m$ and $\Delta\boldsymbol{\alpha}_{t,m}$ have expectation zero, where the expectations are taken over time, replicates and all time-replicate combinations respectively. $C_{\alpha_{t_m},\alpha_{\tau_n}} = \boldsymbol{\alpha}_{t,m}^\top\mathbf{L}_0\boldsymbol{\alpha}_{\tau,n}$ is the additive genetic covariance between replicate m at time t and replicate n at time τ given the base population structure.

$$C_{\alpha_{t_m},\alpha_{\tau_n}} = \left(\bar{\boldsymbol{\alpha}}^\top + \Delta\boldsymbol{\alpha}_t^\top + \Delta\boldsymbol{\alpha}_m^\top + \Delta\boldsymbol{\alpha}_{t,m}^\top\right)\mathbf{L}_0\left(\bar{\boldsymbol{\alpha}} + \Delta\boldsymbol{\alpha}_\tau + \Delta\boldsymbol{\alpha}_n + \Delta\boldsymbol{\alpha}_{\tau,n}\right) \quad \text{(S18)}$$

If we take the expectation of this over pairs of replicates we get:

$$C_{\alpha_t,\alpha_\tau} = \left(\bar{\boldsymbol{\alpha}}^\top + \Delta\boldsymbol{\alpha}_t^\top\right)\mathbf{L}_0\left(\bar{\boldsymbol{\alpha}} + \Delta\boldsymbol{\alpha}_\tau\right) \quad \text{(S19)}$$

since the Δ terms, and the product of pairs of Δ terms when $n \neq m$, have zero expectation. Here, C_{α_t,α_τ} is the between-replicate covariance at time t and τ, and has expectation (with respect to time) $\bar{\boldsymbol{\alpha}}^\top\mathbf{L}_0\bar{\boldsymbol{\alpha}}$ when $t \neq \tau$ and $\bar{\boldsymbol{\alpha}}^\top\mathbf{L}_0\bar{\boldsymbol{\alpha}} + Tr(\mathbf{L}_0 VAR(\Delta\boldsymbol{\alpha}_t))$ when $t = \tau$ where $VAR(\Delta\boldsymbol{\alpha}_t)$ are the (co)variances of $\boldsymbol{\alpha}$ over time.

viii

S3 Treating $\boldsymbol{\alpha}$ as random

In standard quantitative genetic theory, $\boldsymbol{\alpha}$ is considered fixed and the genotypes of individuals in the population are considered random, leading to a distribution of breeding values with variance V_A. More recently, however, dense molecular markers have been used as covariates for predicting genetic values, and the associated coefficients have been treated as random (Meuwissen *et al.*, 2001). As noted by Gianola *et al.* (2009), this raises interpretational problems when trying to relate the variance parameters associated with the coefficients to the concept of V_A.

In our inference section we show that

$$\begin{aligned} E[V_A(0)] &= E[\boldsymbol{\alpha}^\top \mathbf{L}_0 \boldsymbol{\alpha}] \\ &= Tr(\mathbf{L}_0 \mathbf{V}_\alpha) + \boldsymbol{\mu}_\alpha^\top \mathbf{L}_0 \boldsymbol{\mu}_\alpha \end{aligned} \tag{S20}$$

where here we assume the average effects are constant across time/replicates for ease and $\boldsymbol{\mu}_\alpha$ and $\mathbf{V}_\alpha$ are the mean vector and covariance matrix for the average effects (see the Appendix in Gianola *et al.* (2009) also). Critically, we interpret this expectation as an average over the epistemic uncertainty in $\boldsymbol{\alpha}$: it is a posterior mean after marginalising the distribution of $\boldsymbol{\alpha}$ (but conditioning on $\boldsymbol{\mu}_\alpha$ and $\mathbf{V}_\alpha$):

$$E[V_A(0)|\boldsymbol{\mu}_\alpha, \mathbf{V}_\alpha, \mathbf{L}_0] \propto \int_{\boldsymbol{\alpha}} \boldsymbol{\alpha}^\top \mathbf{L}_0 \boldsymbol{\alpha} \cdot |\mathbf{J}| \cdot Pr(\boldsymbol{\alpha}|\boldsymbol{\mu}_\alpha, \mathbf{V}_\alpha, \mathbf{L}_0) d\boldsymbol{\alpha} \tag{S21}$$

where $\mathbf{J}$ is the Jacobian of the transform from $\boldsymbol{\alpha}$ to V_A and depends on $\mathbf{L}_0$.

For an inferential method we see no contradiction between treating $\boldsymbol{\alpha}$ as fixed when *defining* V_A but treating it as random when *estimating* V_A. However, when treating $\bar{\boldsymbol{\alpha}}$ as random it is important to ensure that the underlying model is invariant with respect to which allele is chosen as the reference. Quantities such as V_A and genomic best linear unbiased predictors (gBLUP) are insensitive to which allele at a locus is chosen as the reference allele and which allele is chosen as the alternate allele. To make this explicit, consider the diagonal 'assignment' matrix $\mathbf{A}$ for which the diagonal elements are either 1 (the fittest allele is the reference allele) or -1 (the fittest allele is the alternate allele). Under a particular assignment, $\boldsymbol{\alpha} = \mathbf{A}\boldsymbol{\alpha}_+$ and $\mathbf{L} = \mathbf{A}\mathbf{L}_+\mathbf{A}$, where the subscript $+$ indicates the quantity had the fitter of the two alleles been the reference allele at all loci. If we consider V_A conditional on a particular assignment we get (since $\mathbf{A}\mathbf{A} = \mathbf{I}$):

ix

$$\begin{aligned} V_A &= \boldsymbol{\alpha}^\top \mathbf{L} \boldsymbol{\alpha} \\ &= (\mathbf{A}\boldsymbol{\alpha}_+)^\top \mathbf{A}\mathbf{L}_+ \mathbf{A}\mathbf{A}\boldsymbol{\alpha}_+ \\ &= \boldsymbol{\alpha}_+^\top \mathbf{A}\mathbf{A}\mathbf{L}_+ \mathbf{A}\mathbf{A}\boldsymbol{\alpha}_+ \\ &= \boldsymbol{\alpha}_+^\top \mathbf{L}_+ \boldsymbol{\alpha}_+ \end{aligned} \tag{S22}$$

showing we get the same value of V_A irrespective of the assignment we choose. In contrast, quantities such as $E[\mathbf{L}\boldsymbol{\alpha}]$ are sensitive to the assignment, since they undergo a sign reversal under a different choice:

$$\begin{aligned} \mathbf{L}\boldsymbol{\alpha} &= \mathbf{A}\mathbf{L}_+ \mathbf{A}\mathbf{A}\boldsymbol{\alpha}_+ \\ &= \mathbf{A}\mathbf{L}_+ \boldsymbol{\alpha}_+ \end{aligned} \tag{S23}$$

It is tempting to use the argument that $E[\mathbf{L}\boldsymbol{\alpha}] = \mathbf{0}$ when the reference allele is chosen at random, since α is equally likely to be positive as negative. The logic behind this argument can be expressed mathematically as $E[\boldsymbol{\alpha}] = E[\mathbf{A}]E[\boldsymbol{\alpha}_+]$ since the reference allele is chosen at random and so $\mathbf{A}$ must be independent of $\boldsymbol{\alpha}_+$. Under this same assumption $E[\mathbf{A}] = \mathbf{0}$, since any diagonal element has an equal chance of being -1 or 1, such that $E[\mathbf{L}\boldsymbol{\alpha}] = 0$. However, this logic is incorrect. The argument envisages $\mathbf{A}$ as random, yet for any particular analysis $\mathbf{A}$ is no longer a random variable but fixed - a choice has been made as to which allele is the reference allele - even if there remains epistemic uncertainty as to whether the reference allele is the fitter of the two alleles.

When the reference allele is chosen arbitrarily, any sensible distribution for $\boldsymbol{\alpha}$ must induce the same distribution on $\boldsymbol{\alpha}_+$ regardless of the assignment. If $\boldsymbol{\mu}_{\alpha_+}$ and $\mathbf{V}_{\alpha_+}$ are the means and (co)variances of the average effects had all reference alleles been the fitter allele, then the distribution for a particular assignment becomes $\boldsymbol{\mu}_\alpha = \mathbf{A}\boldsymbol{\mu}_{\alpha_+}$ and $\mathbf{V}_\alpha = \mathbf{A}\mathbf{V}_{\alpha_+}\mathbf{A}$. Given $\mathbf{A}^{-1} = \mathbf{A}$ this implies $\boldsymbol{\mu}_{\alpha_+} = \mathbf{A}\boldsymbol{\mu}_\alpha$ and $\mathbf{V}_{\alpha_+} = \mathbf{A}\mathbf{V}_\alpha\mathbf{A}$. For, $\boldsymbol{\mu}_\alpha$ this implies that suitable models should be (weighted) sums of differences between invariant properties of the alleles such that the difference reverses sign when the reference and alternate allele are switched. This might be their (log) frequency, such that the model is $\beta(p - q)$, with β a parameter, or it might be the difference in derived vs ancestral coded as 1 vs -1, such that the model is 2β or -2β depending on whether the reference allele is derived or ancestral, respectively.

If $\mathbf{V}_{\bar{\alpha}}$ is assumed to be diagonal, all models are permissible since the square removes any sign. Since the multiplication of diagonal matrices is not affected by order we can see this directly:

$$\begin{aligned}
\mathbf{V}_{\alpha_+} &= \mathbf{A}\mathbf{V}_\alpha\mathbf{A} \\
\mathbf{V}_{\alpha_+} &= \mathbf{A}\mathbf{A}\mathbf{V}_\alpha \\
\mathbf{V}_{\alpha_+} &= \mathbf{V}_\alpha
\end{aligned} \tag{S24}$$

However, for non-diagonal matrices a suitable distribution must result in a sign reversal of all covariances at a locus when the reference and alternate alleles are switched. The most obvious way to achieve this is to allow $\mathbf{V}_\alpha$ to be proportional to $\mathbf{L}^p$ since

$$\begin{aligned}
\mathbf{L}^p &= (\mathbf{A}\mathbf{L}_+\mathbf{A})^p \\
&= \mathbf{A}\mathbf{L}_+\mathbf{A}...\mathbf{A}\mathbf{L}_+\mathbf{A} \\
&= \mathbf{A}\mathbf{L}_+^p\mathbf{A}
\end{aligned} \tag{S25}$$

where $\mathbf{L}_+$ is the linkage-disequilibrium matrix had the fittest allele been the reference allele at all loci. Under this assumption

$$\begin{aligned}
\mathbf{V}_{\alpha_+} &= \mathbf{A}\mathbf{V}_\alpha\mathbf{A} \\
\mathbf{V}_{\alpha_+} &\propto \mathbf{A}\mathbf{L}^p\mathbf{A} \\
\mathbf{V}_{\alpha_+} &\propto \mathbf{A}\mathbf{A}\mathbf{L}_+^p\mathbf{A}\mathbf{A} \\
\mathbf{V}_{\alpha_+} &\propto \mathbf{L}_+^p
\end{aligned} \tag{S26}$$

A similar model was proposed by Zeng *et al.* (2018), although there, $\mathbf{L}$ was treated as diagonal such that the variance of the average effects are assumed to be proportional to the genetic diversities to some power.

xi

S4 Projection matrices

Rather than working with original allele frequencies, we project them into a new reduced subspace defined by $\mathbf{L}_0$ and claim that $V_A(0)$ does not depend on allele frequency changes outside of this space. de Los Campos *et al.* (2015) (in Supplementary Methods I) show that additive genetic variances are rotationally invariant: they are invariant to arbitrary linear transformations of the genotypes. Have $\mathbf{U}$ be *all* eigenvectors of $\mathbf{L}_0$ and $\mathbf{D}$ a diagonal matrix with the eigenvalues of $\mathbf{L}_0$ square-rooted along the diagonal. If we use the transformation $\mathbf{U}^\top$, de Los Campos *et al.* (2015) show that the average effects in the projected space are $\mathbf{U}^{-\top}\boldsymbol{\alpha}$ and the covariance of the projected genotypes is $\mathbf{U}^\top\mathbf{L}_0\mathbf{U}$, and:

$$V_A(0) = \boldsymbol{\alpha}^\top\mathbf{U}^{-\top}\mathbf{U}^\top\mathbf{L}_0\mathbf{U}\mathbf{U}^{-1}\boldsymbol{\alpha} = \boldsymbol{\alpha}^\top\mathbf{L}_0\boldsymbol{\alpha} \tag{S27}$$

Since $\mathbf{L}_0 = \mathbf{U}\mathbf{D}\mathbf{D}\mathbf{U}^\top$,

$$V_A(0) = \boldsymbol{\alpha}^\top\mathbf{U}^{-\top}\mathbf{U}^\top\mathbf{U}\mathbf{D}\mathbf{D}\mathbf{U}^\top\mathbf{U}\mathbf{U}^{-1}\boldsymbol{\alpha} \tag{S28}$$

which reduces to

$$V_A(0) = \boldsymbol{\alpha}^\top\mathbf{U}^{-\top}\mathbf{D}\mathbf{D}\mathbf{U}^{-1}\boldsymbol{\alpha} \tag{S29}$$

since eigenvectors are unitary $\mathbf{U}^{-1} = \mathbf{U}^\top$ and so $\mathbf{U}\mathbf{U}^\top = \mathbf{I}$. The maximum rank of $\mathbf{L}_0$ will generally be the number of individuals in the base population and will be substantially less then the number of loci. Consequently some diagonal elements of $\mathbf{D}$ will be zero which will set all elements in the corresponding column of $\mathbf{U}^{-\top}\mathbf{D}$ to zero. Consequently we can simply define the projection matrix $\mathbf{U_L}$ with all columns associated with zero eigenvalues dropped without loss of information.

To show that our chosen projection $(\mathbf{P} = \mathbf{D}_2^{-1}\mathbf{U}_2^\top\mathbf{U_L}^\top)$ results in identical and independently distributed residuals, we need to show that the drift covariance matrix is an identity matrix under this projection. If we write down the eigendecomposition of $\mathbf{U_L}^\top(\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)})\mathbf{U_L} = \mathbf{U}_2\mathbf{D}_2\mathbf{D}_2\mathbf{U}_2^\top$, and use the unitary property of $\mathbf{U_L}$ to note that $\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)}$ can be expressed as $\mathbf{U_L}\mathbf{U}_2\mathbf{D}_2\mathbf{D}_2\mathbf{U}_2^\top\mathbf{U_L}^\top$. The drift covariance matrix under the projection is then:

$$
\begin{aligned}
VAR(\underset{D}{\Delta}\overrightarrow{\mathbf{p}}_m) \quad &= \mathbf{P}\,VAR(\underset{D}{\Delta}\mathbf{p}_m)\mathbf{P}^\top \\
&= \mathbf{P}(\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)})\mathbf{P}^\top \\
&= \mathbf{P}\mathbf{U_L}\mathbf{U}_2\mathbf{D}_2\mathbf{D}_2\mathbf{U}_2^\top\mathbf{U_L}^\top\mathbf{P}^\top \\
&= \mathbf{D}_2^{-1}\mathbf{U}_2^\top\mathbf{U_L}^\top\mathbf{U_L}\mathbf{U}_2\mathbf{D}_2\mathbf{D}_2\mathbf{U}_2^\top\mathbf{U_L}^\top\mathbf{U_L}\mathbf{U}_2\mathbf{D}_2^{-1} \\
&= \mathbf{I}
\end{aligned}
\tag{S30}
$$

xii

S5 Bias correction for $\boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0\boldsymbol{\mu}_{\bar{\alpha}}$

The expression for $E[V_{\bar{A}}(0)]$ involves a quadratic in the vector of expected average effects, $\boldsymbol{\mu}_{\bar{\alpha}}$: $\boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0\boldsymbol{\mu}_{\bar{\alpha}}$. Replacing $\boldsymbol{\mu}_{\bar{\alpha}}$ with an estimate of it, $\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}$, will generally result in the estimate of $V_{\bar{A}}(0)$ being upwardly biased even if $\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}$ is an unbiased estimate of $\boldsymbol{\mu}_{\bar{\alpha}}$. To understand this, and correct for it, have the linear model

$$\boldsymbol{\mu}_{\bar{\alpha}} = \mathbf{X}\boldsymbol{\beta}_{\bar{\alpha}} \tag{S31}$$

where $\mathbf{X}$ is the fixed-effect design matrix and $\boldsymbol{\beta}_{\bar{\alpha}}$ an associated parameter vector. In our analyses of simulated data, $\mathbf{X}$ only has one column, $\mathbf{p}_0 - \mathbf{q}_0$, and $\boldsymbol{\beta}_{\bar{\alpha}}$ is a scalar: $\beta_{\bar{\alpha}}^{(1)}$. We can have

$$\widehat{\boldsymbol{\beta}_{\bar{\alpha}}} = \boldsymbol{\beta}_{\bar{\alpha}} + \mathbf{m}_{\bar{\alpha}} \tag{S32}$$

where $\mathbf{m}_{\bar{\alpha}}$ is the vector of deviations of the estimates from their true value. The expected estimate of $\boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0\boldsymbol{\mu}_{\bar{\alpha}}$ is then

$$\begin{aligned} E[\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}^{\top}\mathbf{L}_0\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}] =\ & \boldsymbol{\beta}_{\bar{\alpha}}^{\top}\mathbf{X}^{\top}\mathbf{L}_0\mathbf{X}\boldsymbol{\beta}_{\bar{\alpha}} + E[\mathbf{m}_{\bar{\alpha}}^{\top}\mathbf{X}^{\top}\mathbf{L}_0\mathbf{X}\mathbf{m}_{\bar{\alpha}}] \\ =\ & \boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0\boldsymbol{\mu}_{\bar{\alpha}} + E[\mathbf{m}_{\bar{\alpha}}^{\top}\mathbf{X}^{\top}\mathbf{L}_0\mathbf{X}\mathbf{m}_{\bar{\alpha}}] \end{aligned} \tag{S33}$$

Here, the expectation is taken over the distribution of estimates, and it is assumed the estimates are unbiased (since then, $E[\mathbf{m}_{\bar{\alpha}}] = \mathbf{0}$, and so $E[\mathbf{m}_{\bar{\alpha}}\boldsymbol{\beta}_{\bar{\alpha}}^{\top}] = \mathbf{0}$). This same assumption implies

$$E[\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}^{\top}\mathbf{L}_0\widehat{\boldsymbol{\mu}_{\bar{\alpha}}}] =\ \boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0\boldsymbol{\mu}_{\bar{\alpha}} + Tr\left(\mathbf{X}^{\top}\mathbf{L}_0\mathbf{X}E[\mathbf{m}_{\bar{\alpha}}^{\top}\mathbf{m}_{\bar{\alpha}}]\right) \tag{S34}$$

Since $E[\mathbf{m}_{\bar{\alpha}}^{\top}\mathbf{m}_{\bar{\alpha}}]$ is a matrix of sampling (co)variances for the parameters, to get an improved estimate we can use the inverse Hessian to get an approximate $\mathbf{S}_{\bar{\alpha}} = E[\mathbf{m}_{\bar{\alpha}}^{\top}\mathbf{m}_{\bar{\alpha}}]$:

$$\widehat{\boldsymbol{\mu}_{\bar{\alpha}}^{\top}\mathbf{L}_0}\boldsymbol{\mu}_{\bar{\alpha}} = \widehat{\boldsymbol{\mu}_{\bar{\alpha}}}^{\top}\mathbf{L}_0\widehat{\boldsymbol{\mu}_{\bar{\alpha}}} - Tr\left(\mathbf{X}^{\top}\mathbf{L}_0\mathbf{X}\mathbf{S}_{\bar{\alpha}}\right) \tag{S35}$$

In addition, $Tr(\mathbf{L}_0\widehat{\mathbf{V}_{\bar{\alpha}}})$ is most likely an upwardly biased estimator of $Tr(\mathbf{L}_0\mathbf{V}_{\bar{\alpha}})$. However, we found no easy way to determine or correct for the degree of bias, and the simulation results suggest that the bias is likely to be small, at least when the number of replicates or generations is moderately large.

S6 Analysing allele frequency data from pool-seq

The main derivation of our method assumes allele frequency changes in each replicate are known without error, and the majority of our simulations were analysed with this being true. However, this assumption is unlikely to be met in reality and here we develop a method for incorporating uncertainty in allele frequencies estimated using pool-seq.

Assuming that all individuals have the same probability of being sampled, the covariance in reference allele number at locus i and j is equal to the number of reads that span both sites, O_{ij}, multiplied by the covariance in allele number across haplotypes $2L'_{i,j}$. To get the covariance in reference allele frequency we must divide this by the number of reads spanning site i (O_{ii}) and the number of reads spanning site j to get $2L'_{i,j}O_{ij}/(O_{ii}O_{jj})$. When $i = j$ this reduces to the familiar binomial variance for allele frequency at a single site $2L'_{i,i}O_{ii}/(O_{ii}O_{ii}) = pq/O_{ii}$. When analysing binomial data the variance often exceeds its expectation - a phenomenon known as overdisperion. In the context of pool-seq this will be mostly driven by individuals, rather than gametic contributions, varying in their probability of being sampled. In this instance, the exact form of the sampling (co)variances will depend in a complicated way on both $\mathbf{L}'$ and $\mathbf{L}''$, but here we assume that for sites close enough to be spanned by the same read $\mathbf{L}'$ dominates over $\mathbf{L}''$ and at a site pq dominates over $\mathbf{L}''$. We then simply accommodate any overdispersion using a parameter that scales the sampling (co)variances, σ_o^2, which is constant over loci. This approach is widely used when analysing binomial data (the qausibinomial approach - (McCullagh and Nelder, 1989)) and values greater than one indicate overdispersion. In what follows we also assume σ_o^2 is constant across replicate/time-points although this could be relaxed.

Having the matrix $\mathbf{Q}$ with the ij^{th} element equal to $O_{ij}/(O_{ii}O_{jj})$ the sampling covariance in allele frequency in replicate m at time t_m is equal to $2\sigma_o^2(\mathbf{L}'_{t_m,m} \circ \mathbf{Q}_{t_m,m})$. Since errors in allele frequency estimates are independent at different time points the expected sampling covariance in allele frequency change is proportional to:

$$
\begin{aligned}
\boldsymbol{\mathcal{E}}_m =&\ 2\sigma_o^2 E\left[\boldsymbol{L}'_{t_m,m} \circ \mathbf{Q}_{t_m,m} + \boldsymbol{L}'_{\tau_m,m} \circ \mathbf{Q}_{\tau_m,m}\right] \\
=&\ 2\sigma_o^2 E\left[(\boldsymbol{\mathcal{L}}_{t_m,m} + \Delta\boldsymbol{L}'_{t_m,m}) \circ \mathbf{Q}_{t_m,m} + (\boldsymbol{\mathcal{L}}_{\tau_m,m} + \Delta\boldsymbol{L}'_{\tau_m,m}) \circ \mathbf{Q}_{\tau_m,m}\right] \quad \text{(S36)} \\
=&\ 2\sigma_o^2 (\boldsymbol{\mathcal{L}}_{t_m,m} \circ \mathbf{Q}_{t_m,m} + \boldsymbol{\mathcal{L}}_{\tau_m,m} \circ \mathbf{Q}_{\tau_m,m})
\end{aligned}
$$

where the expectation is taken over evolutionary realisations of $\mathbf{L}'$ and depends on $E[\Delta\boldsymbol{L}'_{t,m}] = 0$. The expected sampling covariance in projected allele frequency

xiv

change is therefore:

$$\overrightarrow{\boldsymbol{\mathcal{E}}}_m = 2\sigma_o^2 \mathbf{P} \left(\boldsymbol{\mathcal{L}}_{t_m,m} \circ \mathbf{Q}_{t_m,m} + \boldsymbol{\mathcal{L}}_{\tau_m,m} \circ \mathbf{Q}_{\tau_m,m} \right) \mathbf{P}^\top \tag{S37}$$

$\overrightarrow{\boldsymbol{\mathcal{E}}}$ is a square dense matrix equal in dimension to the number of projected loci. Moreover, it will vary over replicates such that the complete covariance structure across all replicates is a large block-diagonal matrix with large blocks. Given the computational burden this entails we fitted approximate models to the simulated data where $\overrightarrow{\boldsymbol{\mathcal{E}}}$ was diagonalised. For the same reason, we also used a diagonalised $\mathbf{Q}$ matrix which only contains information on coverage rather than complete information on the number of reads overlapping pairs of sites. In addition we also assumed $\boldsymbol{\mathcal{L}}_{t_m,m} = \boldsymbol{\mathcal{L}}_{\tau_m,m} = \mathbf{L}'_0$ for all m. For real data we advocate not using these approximations.

Estimation error in allele frequency at a single-site is comparable to drift, with coverage O_{ii}, or effective coverage $\sigma_o^2 O_{ii}$, equivalent to N_E. Because of this it might be tempting to think that estimates of $V_{\bar{A}}(0)$ would remain unbiased if estimation error was ignored since the residual variance will soak up the excess variance when model fitting. The only advantage, then, of including $\overrightarrow{\boldsymbol{\mathcal{E}}}$ is to increase precision when there is heterogeneity in coverage across sites. However, this is not the case - estimates of $V_{\bar{A}}(0)$ will be downwardly biased if estimation error is ignored and this bias can be substantial unless reads are very long and/or coverage is high. The reason for this is that a major source of information on $V_{\bar{A}}(0)$ comes from correlated changes in allele frequency at loci that are in LD, and in particular correlations that are tighter than expected under drift. If reads only span single sites then the error covariance in allele frequency is zero since $O_{ij} = 0$ resulting in correlations that are weaker than under drift which partly destroys the signal of selection. If reads are longer, then the error covariance in allele frequency will come to resemble that caused by drift and any bias when ignoring the errors should be reduced. However, at very long read lengths the bias may even become positive as the sampling correlations between distant sites on the same chromosome will be higher than that under drift where it will be broken down by a round of recombination (Equation S14).

S7   Comparison with the method of Buffalo and Coop (2019)

To make clearer the distinction between the approach of B&C and our approach, here we explicitly express the expectations and covariances appearing in B&C as conditional on $\mathbf{B}$ and $\boldsymbol{\alpha}$. In the sections dealing with our theory and inference, the conditioning (on $\mathbf{L}_0$) is left implicit. B&C work with the quantity

$$COV(\Delta\mathbf{p}_t, \Delta\mathbf{p}_\tau^\top | \mathbf{B}, \boldsymbol{\alpha}) \tag{S38}$$

where τ is some generation after t (B&C actually only work with the diagonal elements of this matrix, but we retain the full multi-locus model here for generality). Importantly, when B&C estimate the additive genetic variance in fitness they assume that both $\Delta\mathbf{p}_t$ and $\Delta\mathbf{p}_\tau$ both represent allele frequency change over a single generation (Assumption A). As the change due to drift will be independent in different generations, we can rewrite B&C's covariance:

$$\begin{aligned}
COV(\Delta\mathbf{p}_t, \Delta\mathbf{p}_\tau^\top | \mathbf{B}, \boldsymbol{\alpha}) =&\ COV(\mathbf{L}_t\boldsymbol{\alpha}_t, \boldsymbol{\alpha}_\tau^\top\mathbf{L}_\tau | \mathbf{B}, \boldsymbol{\alpha}) \\
=&\ E\left[\mathbf{L}_t\boldsymbol{\alpha}_t\boldsymbol{\alpha}_\tau^\top\mathbf{L}_\tau | \mathbf{B}, \boldsymbol{\alpha}\right] - E\left[\mathbf{L}_t | \mathbf{B}, \boldsymbol{\alpha}\right]\boldsymbol{\alpha}_t\boldsymbol{\alpha}_\tau^\top E\left[\mathbf{L}_\tau | \mathbf{B}, \boldsymbol{\alpha}\right]
\end{aligned} \tag{S39}$$

Since the diagonal elements of $\mathbf{L}$ have to be positive, a sufficient, but not necessary, condition for the final term to be zero is that there is no direct selection on the loci (i.e. $\boldsymbol{\alpha} = \mathbf{0}$). It is not a necessary condition because the change caused by direct selection at all loci could be exactly balanced by the change caused by indirect selection at other loci, although we ignore this unlikely scenario. The assumption that $\boldsymbol{\alpha} = \mathbf{0}$ is achieved in B&C by assuming that sites can be partitioned into neutral and selected sites and that allele frequency change is only tracked at the neutral sites (Assumption B). To understand the consequences of this assumption we consider all loci are being followed, both a selected set ($\mathcal{S}$) and a neutral set ($\mathcal{N}$). Consequently,

$$\boldsymbol{\alpha} = \left[\begin{array}{c} \mathbf{0} \\ \boldsymbol{\alpha}_\mathcal{S} \end{array} \right] \tag{S40}$$

and so

$$\boldsymbol{\alpha}\boldsymbol{\alpha}^\top = \left[\begin{array}{cc} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \boldsymbol{\alpha}_\mathcal{S}\boldsymbol{\alpha}_\mathcal{S}^\top \end{array} \right] \tag{S41}$$

We can also partition $\mathbf{L}$

xvi

$$\mathbf{L} = \left[\begin{array}{cc} \mathbf{L}_{\mathcal{N}} & \mathbf{L}_{\mathcal{N},\mathcal{S}} \\ \mathbf{L}_{\mathcal{S},\mathcal{N}} & \mathbf{L}_{\mathcal{S}} \end{array} \right] \tag{S42}$$

and writing $\mathbf{L}_{\mathcal{N},\mathcal{S}} = \mathbf{B}_{\mathcal{N}}\mathbf{R}_{\mathcal{N},\mathcal{S}}\mathbf{B}_{\mathcal{S}}$, Equation S39 for neutral sites becomes

$$\begin{aligned} COV(\Delta\mathbf{p}_{\mathcal{N}_t}, \Delta\mathbf{p}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha}) = \quad & \mathbf{B}_{\mathcal{N}_t} E\left[\mathbf{R}_{\mathcal{N}_t,\mathcal{S}_t}\mathbf{B}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_\tau}^\top \mathbf{B}_{\mathcal{S}_\tau}\mathbf{R}_{\mathcal{S}_\tau,\mathcal{N}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{N}_\tau} \\ & +\mathbf{B}_{\mathcal{N}_t} E\left[\mathbf{R}_{\mathcal{N}_t,\mathcal{S}_t}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_\tau}^\top \mathbf{B}_{\mathcal{S}_\tau} E\left[\mathbf{R}_{\mathcal{S}_\tau,\mathcal{N}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{N}_\tau} \end{aligned} \tag{S43}$$

Under the conditioning of B&C, $\mathbf{R}_{\mathcal{S},\mathcal{N}_\tau}$ is a random variable. If we assume that the linkage-disequilibrium between the neutral alleles and the selected alleles has arbitrary sign, then $E[\mathbf{R}_{\mathcal{S},\mathcal{N}_\tau}|\mathbf{B},\boldsymbol{\alpha}] = \mathbf{0}$, which will be met if the reference allele is chosen arbitrarily (e.g. not based on minor allele frequency (Good, 2022)). Under this assumption (Assumption C) the final term in Equation S43 disappears to give:

$$COV(\Delta\mathbf{p}_{\mathcal{N}_t}, \Delta\mathbf{p}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha}) = \quad \mathbf{B}_{\mathcal{N}_t} E\left[\mathbf{R}_{\mathcal{N}_t,\mathcal{S}_t}\mathbf{B}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_\tau}^\top \mathbf{B}_{\mathcal{S}_\tau}\mathbf{R}_{\mathcal{S}_\tau,\mathcal{N}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{N}_\tau} \tag{S44}$$

As in our inference section, the vector of allele frequency changes could be transformed using matrix $\mathbf{P}$ ($\Delta\overrightarrow{\mathbf{p}} = \mathbf{P}\Delta\mathbf{p}$) and B&C use the projection $\mathbf{P} = \mathbf{B}_{\mathcal{N}_t}^{-1}$ (they actually multiply this by $\sqrt{2}$ - see Equation S54) which results in

$$\begin{aligned} COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t}, \Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha}) = \quad & E\left[\mathbf{R}_{\mathcal{N}_t,\mathcal{S}_t}\mathbf{B}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_\tau}^\top \mathbf{B}_{\mathcal{S}_\tau}\mathbf{R}_{\mathcal{S}_\tau,\mathcal{N}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{N}_\tau}\mathbf{B}_{\mathcal{N}_t}^{-1} \\ = \quad & \mathbf{B}_{\mathcal{N}_t}^{-1} E\left[\mathbf{R}_{\mathcal{N}_t,\mathcal{S}_t}\mathbf{B}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_t}\boldsymbol{\alpha}_{\mathcal{S}_\tau}^\top \mathbf{B}_{\mathcal{S}_\tau}\mathbf{R}_{\mathcal{S}_\tau,\mathcal{N}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{N}_\tau} \end{aligned} \tag{S45}$$

Further derivation in B&C considers the expected value of a diagonal element of this matrix: the average (over neutral loci) covariance in projected allele frequency change. However, it is perhaps easier to note that the trace of this matrix is equal to this average multiplied by the number of neutral loci, $n_{L_\mathcal{N}}$ (see below). Since the trace of an outer product is equal to the inner product we get:

$$\begin{aligned} Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t}, \Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha})\right) \quad & = E\left[\boldsymbol{\alpha}_{\mathcal{S}_t}^\top \mathbf{B}_{\mathcal{S}_t}\mathbf{R}_{\mathcal{S}_t,\mathcal{N}_t}\mathbf{B}_{\mathcal{N}_t}^{-1}\mathbf{B}_{\mathcal{N}_\tau}\mathbf{R}_{\mathcal{N}_\tau,\mathcal{S}_\tau}\mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right] \\ & = \boldsymbol{\alpha}_{\mathcal{S}_\tau}^\top \mathbf{B}_{\mathcal{S}_t} E\left[\mathbf{R}_{\mathcal{S}_t,\mathcal{N}_t}\mathbf{B}_{\mathcal{N}_t}^{-1}\mathbf{B}_{\mathcal{N}_\tau}\mathbf{R}_{\mathcal{N}_\tau,\mathcal{S}_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau} \end{aligned} \tag{S46}$$

The diagonal element j of $\mathbf{W}_{t\tau} = \mathbf{R}_{\mathcal{S}_t,\mathcal{N}_t}\mathbf{B}_{\mathcal{N}_t}^{-1}\mathbf{B}_{\mathcal{N}_\tau}\mathbf{R}_{\mathcal{N}_\tau,\mathcal{S}_\tau}$ is equal to the sum of selected locus j's $R_{t,ji}R_{\tau,ji}(b_{\tau,i}/b_{t,i})$ across all neutral loci i. The jk^{th} off-diagonal element is the sum of $R_{t,ji}R_{\tau,ki}(b_{\tau,i}/b_{t,i})$ for selected loci j and k across all neutral

xvii

loci i. If we write $\mathbf{W}_{t\tau} = \mathbf{H}_{t\tau} + (\mathbf{W}_{t\tau} - \mathbf{H}_{t\tau})$ where $\mathbf{H}_{t\tau}$ and $\mathbf{W}_{t\tau} - \mathbf{H}_{t\tau}$ are zero but for the diagonal and off-diagonal elements respectively, then Equation S46 becomes

$$Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t}, \Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top|\mathbf{B},\boldsymbol{\alpha})\right) \begin{aligned} &= \boldsymbol{\alpha}_{\mathcal{S}_t}^\top \mathbf{B}_{\mathcal{S}_t} E\left[\mathbf{H}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right] \mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau} \\ &+ \boldsymbol{\alpha}_{\mathcal{S}_t}^\top \mathbf{B}_{\mathcal{S}_t} E\left[\mathbf{W}_{t\tau} - \mathbf{H}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right] \mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau} \end{aligned} \tag{S47}$$

If we focus on a system with two selected loci, j and k, then the final term in Equation S47 is equal to:

$$\sum_j \sum_k \left(\alpha_{t,j}\alpha_{\tau,k} b_{j_t,j_t} b_{k_\tau,k_\tau} \sum_i \left(E\left[R_{t,ij}R_{\tau,ik}|\mathbf{B},\boldsymbol{\alpha}\right] b_{i_\tau}/b_{i_t} \right) + \alpha_{j_\tau}\alpha_{k_t} b_{j_\tau,j_\tau} b_{k_t,k_t} \sum_i \left(E\left[R_{\tau,ij}R_{t,ik}|\mathbf{B},\boldsymbol{\alpha}\right] b_{i_\tau}/b_{i_t} \right) \right). \tag{S48}$$

Under Hill-Robertson interference, if α_j and α_k have the same sign we expect them to be in negative LD with each other, and as a consequence have opposing patterns of LD with the neutral loci (i.e. if $\alpha_j\alpha_k > 0$ then we expect $E[R_{i,j}R_{i,k}] < 0$ and vice versa). Although the terms of these products are evaluated at different generations in Equation S48 (t and τ) we expect the terms to share sign in the same way, generating a negative expectation for the second term in Equation S47. However, assuming an absence of Hill-Robertson interference, or signed linkage-disequilibrium more generally (Assumption D), then the final term in Equation S47 can be dropped:

$$Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t}, \Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top)|\mathbf{B},\boldsymbol{\alpha}\right) \quad = \boldsymbol{\alpha}_{\mathcal{S}_t}^\top \mathbf{B}_{\mathcal{S}_t} E\left[\mathbf{H}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right] \mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau} \tag{S49}$$

Based on a deterministic model for changes in linkage-disequilibrium (Assumption E) and assuming nongametic-phase linkage-disequilibrium is absent (Assumption F), Equations 40-44 in B&C derive an expression from which $E\left[\mathbf{H}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]$ can be computed. Under the assumption that drift (or selection) does not alter the dynamics of $\mathbf{R}$ then (Equation 42 in B&C):

$$L_{j_\tau,i_\tau} = L_{j_t,i_t}\frac{b_{j_\tau}^2}{b_{j_t}^2}(1 - r(g_{j,i}))^{\tau-t} \tag{S50}$$

which implies

$$R_{j_\tau,i_\tau} = R_{j_t,i_t}\frac{b_{j_\tau}b_{i_t}}{b_{j_t}b_{i_\tau}}(1 - r(g_{j,i}))^{\tau-t} \tag{S51}$$

and so

$$E\left[H_{j_t,j_\tau}|\mathbf{B},\boldsymbol{\alpha}\right] \begin{aligned} &= \sum_i E\left[R_{j_t,i_t}R_{j_\tau,i_\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\frac{b_{i_\tau}}{b_{i_t}} \\ &= \frac{b_{j_\tau}}{b_{j_t}}\sum_i E\left[R_{j_t,i_t}^2|\mathbf{B},\boldsymbol{\alpha}\right](1 - r(g_{j,i}))^{\tau-t} \end{aligned} \tag{S52}$$

xviii

where $r(g_{j,i})$ is the recombination rate as a function of the distance $g_{i,j}$ between the two loci. Writing $\mathbf{F}_{t\tau}$ a diagonal matrix with the j^{th} element equal to $\sum_i E\left[R_{j_t,i_t}^2|\mathbf{B},\boldsymbol{\alpha}\right](1-r(g_{j,i}))^{\tau-t}$, then $E\left[\mathbf{H}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right] = E\left[\mathbf{F}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{S}_\tau}\mathbf{B}_{\mathcal{S}_t}^{-1}$ and so

$$
\begin{aligned}
Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t},\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top|\mathbf{B},\boldsymbol{\alpha})\right) &= \boldsymbol{\alpha}_{\mathcal{S}_t}^\top\mathbf{B}_{\mathcal{S}_t}E\left[\mathbf{F}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{S}_\tau}\mathbf{B}_{\mathcal{S}_t}^{-1}\mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau} \\
&= \boldsymbol{\alpha}_{\mathcal{S}_t}^\top E\left[\mathbf{F}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{S}_\tau}\mathbf{B}_{\mathcal{S}_\tau}\boldsymbol{\alpha}_{\mathcal{S}_\tau} \\
&= \boldsymbol{\alpha}_{\mathcal{S}_t}^\top E\left[\mathbf{F}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]\mathbf{B}_{\mathcal{S}_\tau}^2\boldsymbol{\alpha}_{\mathcal{S}_\tau}
\end{aligned}
\tag{S53}
$$

where (under random mating) $\mathbf{B}_{\mathcal{S}_\tau}^2$ is a diagonal matrix with elements proportional to the genetic diversities in generation τ. However, this is still hard to evaluate since $E\left[\mathbf{F}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]$ will vary over loci in a way that may depend on $\mathbf{B}$ and hence $\mathbf{B}_{\mathcal{S}_t}^2$. The method of B&C assumes that the sample correlation between the diagonal elements of $E\left[\mathbf{F}_{t\tau}|\mathbf{B},\boldsymbol{\alpha}\right]$ and the elements $\alpha_{j_t}\alpha_{j_\tau}b_{j_\tau}^2$ is zero (Assumption G). Under this assumption:

$$
\begin{aligned}
Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t},\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top|\mathbf{B},\boldsymbol{\alpha})\right) &= \frac{1}{n_{L_\mathcal{S}}}\boldsymbol{\alpha}_{\mathcal{S}_t}^\top\mathbf{B}_{\mathcal{S}_\tau}^2\boldsymbol{\alpha}_{\mathcal{S}_\tau}E\left[Tr(\mathbf{F}_{t\tau})|\mathbf{B},\boldsymbol{\alpha}\right] \\
&= \frac{1}{n_{L_\mathcal{S}}}C_a(t\to\tau)E\left[Tr(\mathbf{F}_{t\tau})|\mathbf{B},\boldsymbol{\alpha}\right]
\end{aligned}
\tag{S54}
$$

where $n_{L_\mathcal{S}}$ is the number of selected loci (note that in B&C the leading term is $\frac{1}{2n_{L_\mathcal{S}}}$ not $\frac{1}{n_{L_\mathcal{S}}}$ and this is because our projection matrices are only proportional by a factor $\sqrt{2}$ - see Equation S45). The method of B&C further assumes (Assumption H) that the average effects are constant in time such that $\boldsymbol{\alpha}_t = \boldsymbol{\alpha}_\tau$ then this reduces to

$$
Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t},\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top|\mathbf{B},\boldsymbol{\alpha})\right) = \frac{V_a(\tau)}{n_{L_\mathcal{S}}}E\left[Tr(\mathbf{F}_{t\tau})|\mathbf{B},\boldsymbol{\alpha}\right]
\tag{S55}
$$

Under Assumption H, Assumption G implies that the additive genic variance contributed by a selected locus is independent of its associations with neutral loci as measured by R_{i_t,j_t}^2. However, allele frequencies at selected loci enter the expression for the additive genic variance, and they also dictate the range of R_{i_t,j_t} and therefore the magnitude of R_{i_t,j_t}^2: when selected allele frequencies are small compared to frequencies at neutral alleles, R_{i_t,j_t} cannot cover the full range of -1 to 1 (Sved and Hill, 2018).

B&C approximate $\mathbf{R}_t$, and therefore $\mathbf{F}_{t\tau}$ under the assumption of mutation-drift-recombination equilibrium (Assumption I). Ohta and Kimura (1971) derived the expectation of $L_{j_t,i_t}L_{j_t,i_t} = L_{j_t,i_t}^2$ under this assumption, although the expectation of R_{j_t,i_t}^2 can only be approximated as the expectation of L_{j_t,i_t}^2 divided by the expectation

of the genetic diversities at the two loci, and is only accurate when the minor allele frequencies are greater than 10% (McVean, 2002) – and can be out by orders of magnitude when allele frequencies are extreme (Song and Song, 2007), as can be expected at loci under selection. Moreover, Ohta and Kimura (1971) derived the expectation $E[L^2_{j_t,i_t}]$ yet the B&C approach actually requires $E\left[L^2_{j_t,i_t}|\mathbf{B},\boldsymbol{\alpha}\right]$ which is considerably more challenging to compute (Good, 2022). In the Appendix B&C also relax, to some extent, Assumption I, where $Tr(\mathbf{F}_{t\tau})$ is calculated empirically using all loci, neutral and selected (Equation 55). In both cases - using the neutral expectation (Assumption I) or empirical LD (Assumption I-b) - the expected LD between selected and neutral loci will be overestimated since in reality selected alleles will be rarer than neutral alleles and so their LD, even measured as a correlation, will be reduced compared to that between neutral loci (Sved and Hill, 2018).

Equation S55 allows $V_a(\tau)$ to be estimated, but B&C aim to estimate $V_a(t)$. Under Assumption H, and assuming that the proportional change in genetic diversity at selected loci between generation t and τ is constant across loci (Assumption J: $\mathbf{B}^2_{\mathcal{S}_t}\mathbf{B}^{-2}_{\mathcal{S}_\tau} = \phi_{t,\tau}\mathbf{I}$), $V_a(t) = \phi_{t,\tau}V_a(\tau)$.

The above derivation assumes that the sites can be partitioned into selected and neutral sites and that the map positions of all sites are known. In practice, this will often be infeasible and so a number of additional assumptions are required. In the absence of map positions, Haldane's (1919) mapping function is assumed for $r(g)$ (Assumption K) but, since the selected loci and their physical position, g, are assumed unknown, a model is also required for g. B&C assume that selected and neutral loci are distributed uniformly and independently such that $g_{j,i}$ has a triangular distribution (Assumption L). Also, since the selected sites are unobserved, $\phi_{t,\tau}$ cannot be computed and so it is assumed that that $\phi_{t,\tau}$ is equal to the ratio of genetic diversity at generation τ to genetic diversity at generation t across all neutral loci (Assumption M).

In addition to the assumptions/approximations made when developing the theory, additional assumptions/approximations are made when making inferences from data. Rather than taking the average (over loci) covariance (over evolutionary replicates) the covariance over loci is taken. However, from the law of total covariance this is expected to yield the correct result under Assumptions B and C. In addition, rather than calculating the (co)variances in projected allele frequency change, it is assumed (Assumption N) that the (co)variances in actual allele frequency divided through by the average projection is a good approximation: $COV_L(\Delta p_t, \Delta p_\tau)/E[b^2_t]$ is a good approximation of $COV_L(\Delta p_t/b_t, \Delta p_\tau/b_t)$ where COV_L designates a covari-

ance over loci (Equation 16 B&C). The two quantities are only expected to agree under restrictive conditions (Bohrnstedt and Goldberger, 1969). However, it is not necessary to make Assumption N and relaxing it can reduce the bias in the estimator considerably. Finally, if there is measurement error in allele frequencies and the same allele frequency measurements are used to calculate change over adjacent intervals then this will generate downward bias in the estimated covariances. This is corrected for in B&C by assuming the sampling error in allele frequencies is binomial around the true value (Assumption O), although in practice non-binomial causes of overdispersion are likely.

To connect Equation S55 with the derivation in B&C more clearly, we can write

$$Tr\left(COV(\Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_t}, \Delta\overrightarrow{\mathbf{p}}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha})\right) = n_{L_\mathcal{N}} \overline{COV(\Delta\overrightarrow{p}_{\mathcal{N}_t}, \Delta\overrightarrow{p}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha})} \tag{S56}$$

where the overline denotes the sample mean for the $n_{L_\mathcal{N}}$ neutral loci. Similarly,

$$\begin{aligned} Tr(\mathbf{F}_{t\tau}) &= \sum_j \sum_i E\left[R^2_{j_t, i_t} | \mathbf{B}, \boldsymbol{\alpha}\right] (1 - r(g_{j,i}))^{\tau-t} \\ &= n_{L_\mathcal{N}} \sum_j E\left[R^2_{j_t, \mathcal{N}_t} | \mathbf{B}, \boldsymbol{\alpha}\right] (1 - r(g_{j,\mathcal{N}}))^{\tau-t} \end{aligned} \tag{S57}$$

Equation S55 can then be written as

$$\overline{COV(\Delta\overrightarrow{p}_{\mathcal{N}_t}, \Delta\overrightarrow{p}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha})} = \frac{V_a(\tau)}{n_{L_\mathcal{S}}} \overline{\sum_j E\left[R^2_{j_t, \mathcal{N}_t} | \mathbf{B}, \boldsymbol{\alpha}\right] (1 - r(g_{j,\mathcal{N}}))^{\tau-t}} \tag{S58}$$

which is Equation 8 in B&C averaged over neutral loci and can be further simplified to

$$\overline{COV(\Delta\overrightarrow{p}_{\mathcal{N}_t}, \Delta\overrightarrow{p}_{\mathcal{N}_\tau}^\top | \mathbf{B}, \boldsymbol{\alpha})} = V_a(\tau) \overline{E\left[R^2_{\mathcal{S}_t, \mathcal{N}_t} | \mathbf{B}, \boldsymbol{\alpha}\right] (1 - r(g_{\mathcal{S},\mathcal{N}}))^{\tau-t}} \tag{S59}$$

where the overline indicates the sample average of all the $n_{L_\mathcal{S}} n_{L_\mathcal{N}}$ pairwise comparisons between selected loci and neutral loci.

Extending this theory to allele frequency change measured in independent replicates, rather than at different time-points in a single population is straightforward (Buffalo and Coop, 2020). If allele frequency change is measured in two populations, m and n, both of which have been derived independently from the base population t generation in the past, Equation S59 can be expressed as:

$$\overline{COV(\Delta\overrightarrow{p}_{\mathcal{N}_m}, \Delta\overrightarrow{p}_{\mathcal{N}_n}^\top | \mathbf{B}, \boldsymbol{\alpha})} = V_a(m, n) \overline{E\left[R^2_{\mathcal{S}_0, \mathcal{N}_0} | \mathbf{B}, \boldsymbol{\alpha}\right] (1 - r(g_{\mathcal{S},\mathcal{N}}))^{2t}} \tag{S60}$$

assuming the average effects have stayed constant.

S8 N_E and average effects under a log-linear model

In our simulations, we model the logarithm of absolute fitness of an individual k using an additive model across loci:

$$log(W_k) = Y_k = \sum_{i=1}^{n_L} y_{k,i} + e_k$$

where $y_{k,i}$ is the genotypic contribution of locus i to the log absolute fitness of individual k, and the environmental deviations, e, have zero mean and standard deviation σ_e. We allow within-locus deviations from additivity using a fitness-scheme equivalent to the one employed by Lynch and Walsh (1998) (pp. 67) adapted for *proportions* of reference alleles as opposed to counts:

$$y_{k,i} = \begin{cases} 0 & \text{if } c_{k,i} = 0 \\ (1 + \kappa_i)\eta_i/2 & \text{if } c_{k,i} = 0.5 \\ \eta_i & \text{if } c_{k,i} = 1 \end{cases} \tag{S61}$$

Note that in the parameterisation of Falconer and Mackay (1996), the genotypic value of the heterozygotes is $(\eta_i + d_i)/2$ such that $d_i = \kappa_i\eta_i$.

We then draw $2N_{t+1}$ parents with replacement from a multinomial with the probability of k being a parent of an offspring proportional to its absolute fitness $W_k = exp(Y_k)$. Following Kojima (1959) we can define Fisher's average effect for relative fitness in terms of the partial derivative of mean fitness with respect to each $E[c] = p_0$ and then rescale by mean fitness:

$$\begin{aligned} \alpha_i &= \frac{1}{\bar{W}}\frac{\partial \bar{W}}{\partial p_{0_i}} \\ &= \frac{1}{E[exp(Y)]}\frac{\partial E[exp(Y)]}{\partial p_{0_i}} \\ &= \frac{\partial log(E[exp(Y)])}{\partial p_{0_i}} \end{aligned} \tag{S62}$$

Assuming Y to be normally distributed with mean μ_Y and variance V_Y allows us to use the expression for the expectation of a log-normal distribution to write:

$$\begin{aligned} \alpha_i &= \frac{\partial log(exp(\mu_Y + V_Y/2))}{\partial p_i} \\ &= \frac{\partial(\mu_Y + V_Y/2)}{\partial p_i} \end{aligned} \tag{S63}$$

It is worth highlighting that, so far, we have made no assumptions about the distribution of the contributions (y_i) made by individual loci to Y.

Next, we assume Hardy-Weinberg genotypic frequencies to write:

$$
\begin{aligned}
\mu_Y & = \sum_{i=1}^{n_L} E[y_i] \\
& = \sum_{i=1}^{n_L} [p_i^2 \eta_i + 2 p_i q_i (1 + \kappa_i) \eta_i / 2] \\
& = \sum_{i=1}^{n_L} [p_i \eta_i + p_i q_i \kappa_i \eta_i] \\
& = \sum_{i=1}^{n_L} \eta_i [p_i + p_i q_i \kappa_i]
\end{aligned}
\tag{S64}
$$

Differentiating with respect to p_i yields:

$$
\frac{\partial \mu_y}{\partial p_i} = \eta_i [1 + \kappa_i (q_i - p_i)]
\tag{S65}
$$

Assuming that all additive genetic variance is genic:

$$
\begin{aligned}
V_Y & = \sum_{i=1}^{n_L} VAR(y_i) + \sigma_e^2 \\
& = \sum_{i=1}^{n_L} (E[y_i^2] - E[y_i]^2) + \sigma_e^2 \\
& = \sum_{i=1}^{n_L} [p_i^2 \eta_i^2 + 2 p_i q_i (1 + \kappa_i)^2 \eta_i^2 / 4 - \eta_i^2 (p_i + p_i q_i \kappa_i)^2] + \sigma_e^2 \\
& = \sum_{i=1}^{n_L} [p_i q_i (1 + \kappa_i)^2 \eta_i^2 / 2 - \eta_i^2 (2 p_i^2 q_i \kappa_i + p_i^2 q_i^2 \kappa_i^2)] + \sigma_e^2 \\
& = \sum_{i=1}^{n_L} [\frac{p_i q_i}{2} (1 + \kappa_i)^2 \eta_i^2 - \eta_i^2 p_i^2 q_i \kappa_i (2 + q_i \kappa_i)] + \sigma_e^2
\end{aligned}
\tag{S66}
$$

Differentiating with respect to p_i:

$$
\begin{aligned}
\frac{\partial V_y}{\partial p_i} & = \frac{q_i - p_i}{2} (1 + \kappa_i)^2 \eta_i^2 - \eta_i^2 \kappa_i [(2 p_i q_i - p_i^2)(2 + \kappa_i q_i) - p_i^2 q_i \kappa_i] \\
& = \frac{q_i - p_i}{2} (1 + \kappa_i)^2 \eta_i^2 - p_i \eta_i^2 \kappa_i [(2 q_i - p_i)(2 + \kappa_i q_i) - p_i q_i \kappa_i]
\end{aligned}
\tag{S67}
$$

Substituting expressions for $\frac{\partial \mu_y}{\partial p_i}$ and $\frac{\partial V_y}{\partial p_i}$ yields:

$$
\begin{aligned}
\alpha_i & = \eta_i [1 + \kappa_i (q_i - p_i)] + \\
& \quad \frac{q_i - p_i}{4} (1 + \kappa_i)^2 \eta_i^2 - \frac{p_i \eta_i^2 \kappa_i}{2} [(2 q_i - p_i)(2 + \kappa_i q_i) - p_i q_i \kappa_i]
\end{aligned}
\tag{S68}
$$

In the parameterisation of Falconer and Mackay (1996), this translates to:

$$
\begin{aligned}
\alpha_i & = \eta_i + d_i (q_i - p_i) + \\
& \quad \frac{q_i - p_i}{4} (\eta_i + d_i)^2 - \frac{d_i p_i}{2} [(2 q_i - p_i)(2 \eta_i + q_i d_i) - p_i q_i d_i]
\end{aligned}
\tag{S69}
$$

Note that as a first approximation Equations S69 and S68 reduce to the classical quantitative genetic expressions for the average effect of gene substitution (Equation 7.5 in Falconer and Mackay (1996) and Equation 4.10b in Lynch and Walsh (1998)).
In order to calculate N_E under this multinomial log-linear model we need to know the variance in offspring number V_o caused by any environmental or non-additive genetic variance. Then $N_E = 4N/(2 + V_o)$ where N is the census population size (Wright, 1938). The expected number of offspring for parent k is $2 N_{t+1} p_k$, as given

xxiii

above, and the variance in the number offspring is $2N_{t+1}p_k(1-p_k)$. From the the law of total variance $V_o = 4N_{t+1}^2Var(p) + 2N_{t+1}E[p(1-p)]$. Since $E[p(1-p)] = E[p] - E[p^2]$ and $Var(p) = E[p^2] - E[p]^2$ then $E[p(1-p)] = E[p] - Var(p) + E[p]^2$. Since $E[p] = 1/N_t$ by definition,

$$
\begin{aligned}
V_o &= 4N_{t+1}^2Var(p) + 2N_{t+1}E[p(1-p)] \\
&= 4N_{t+1}^2Var(p) + \tfrac{2N_{t+1}}{N_t} - 2N_{t+1}Var(p) + \tfrac{2N_{t+1}}{N_t^2} \\
&= (4N_{t+1}^2 - 2N_{t+1})Var(p) + \tfrac{2N_{t+1}}{N_t} + \tfrac{2N_{t+1}}{N_t^2}
\end{aligned}
\tag{S70}
$$

From the properties of the log-normal (with zero mean) we know $E[P] = exp(\sigma_e^2/2)$ and $Var(P) = (exp(\sigma_e^2) - 1)exp(\sigma_e^2)$. Since $p = P/(N_tE[P])$

$$
\begin{aligned}
Var(p) &= Var(P)/(N_tE[P])^2 \\
&= (exp(\sigma_e^2) - 1)exp(\sigma_e^2)/(N_texp((\sigma_e^2)/2))^2 \\
&= (exp(\sigma_e^2) - 1)exp(\sigma_e^2)/(N_t^2exp(\sigma_e^2)) \\
&= (exp(\sigma_e^2) - 1)/N_t^2
\end{aligned}
\tag{S71}
$$

If N_{t+1} and N_t are large then Equation S70 simplifies to

$$
V_o = 4N_{t+1}^2Var(p) + 2N_{t+1}/N_t
\tag{S72}
$$

such that

$$
V_o = (4N_{t+1}^2/N_t^2)(exp(\sigma_e^2) - 1) + 2N_{t+1}/N_t
\tag{S73}
$$

If the population size is constant then this simplifies to $V_o = 4exp(\sigma_e^2) - 2$ and the variance effective population size, N_{E_t}, is $4N_t/(2+V_o) = N_t/exp(\sigma_e^2)$. Note however, that in these expressions using σ_e^2 only assumes the non-additive genetic variance is zero. When non-additive genetic variance is present (which there will be to a small degree even with additivity on the log-scale) V_o will be greater than predicted here and N_E will be consequently smaller.

 # S9 Workflow for implementing the method

Box 1. General workflow of the method

We use genome-wide allele frequency change data ($\Delta \mathbf{p}$) from independent evolutionary replicates derived from the same ancestral base population with a known linkage structure ($\mathbf{L}_0$) to infer the mean ($\boldsymbol{\mu}_{\bar{\alpha}}$) and the variance ($\mathbf{V}_{\bar{\alpha}}$) of the distribution of average effects for fitness. Our goal is to then estimate an expectation for the additive genetic variance for relative fitness by averaging over the distribution of average effects; i.e. $E[V_{\bar{A}}(0)] = Tr(\mathbf{L}_0 \mathbf{V}_{\bar{\alpha}}) + \boldsymbol{\mu}_{\bar{\alpha}}^{\top} \mathbf{L}_0 \boldsymbol{\mu}_{\bar{\alpha}}$. We use simple, yet biologically sensible models, $\boldsymbol{\mu}_{\bar{\alpha}} = \beta_{\bar{\alpha}}^{(1)}(\mathbf{p}_0 - \mathbf{q}_0)$ and $\mathbf{V}_{\bar{\alpha}} = \sigma_{\bar{\alpha}}^2 \mathbf{L}_0^{p_{\bar{\alpha}}}$, which essentially allows the average effect for fitness at a site to depend on allele frequency and genetic diversity. This model can be readily implemented as a linear mixed model for suitably projected allele frequency change treating locus effects as random (see below). This, effectively, allows us to partition the total allele frequency change into a component caused by predictable responses to direct and indirect selection (modelled using the random effects of locus) and a component caused by unpredictable responses to selection as well as genetic drift (absorbed by the model residuals).

Experimental design and data:

A typical experimental design permitting the implementation of our method would involve a base populations of N_0 individuals which are sequenced individually. A number of independent evolutionary replicate populations derived from the base populations are then allowed to evolve between generations t and τ after the base population. If the replicate populations are derived from the base population after just a single round of reproduction t would be 1.

The following data are compiled from the experiment:

$\mathbf{C_{g,0}}$: A matrix with $2N_0$ rows and n_L columns indicating the presence or absence (1 or 0) at each of the n_L segregating sites in each of the $2N_0$ gametic contributions in the base population. If phased genomes are unavailable, the $N_0 \times n_L$ matrix of the proportions of the copies of the (arbitrarily chosen) reference allele at each locus in each individual ($\mathbf{C_0}$) may be used.

$\Delta \mathbf{p}_m$: For each replicate m, a vector representing allele frequency change at each segregating locus between generations t and τ after the base populations. t and τ must be known.

$\mathbf{R}_-$: An $n_L \times n_L$ matrix with 1's on the diagonal and the probabilities of non-recombination between pairs of sites as off-diagonal elements. A recombination map, if available, may be used to construct this matrix.

Step 1: Compute $\mathbf{L}_0$, $\mathbf{L}_0^{'}$, $\mathbf{L}_0^{''}$, and $\tilde{\mathbf{L}}_0$

First, we compute the covariance of $\mathbf{C_0}$ over individuals to obtain $\mathbf{L}_0$, and the co-variance of $\mathbf{C_{g,0}}$ over gametic contributions to obtain the matrix of gametic phase disequilibria $(\mathbf{L}_0^{'})$. The matrix of the non-gametic phase disequilibria $(\mathbf{L}_0^{''})$ is then given by $\mathbf{L}_0 - \mathbf{L}_0^{'}$ allowing us to compute $\tilde{\mathbf{L}}_0$ as a weighted sum of $\mathbf{L}_0^{'}$ and $\mathbf{L}_0^{''}$, with the the ij^{th} element of the latter being weighted by $r_{ij}/(1 - r_{ij})$, where r_{ij} is the probability of recombination between the two loci. If phased genomes are unavailable, $\tilde{\mathbf{L}}_0$ may be set to $\mathbf{L}_0$ which assumes that the non-gametic phase disequilibria are negligible.

Step 2: Compute the predicted L in generation t in replicate m $(\boldsymbol{\mathcal{L}}_{t,m})$

Given $\tilde{\mathbf{L}}_0$ in the base population, the expected $\mathbf{L}$ in generation t in replicate m under the action of drift and recombination is given by $\boldsymbol{\mathcal{L}}_{t,m} = \mathbf{N}_{t,m} \circ \tilde{\mathbf{L}}_0$, where the ij^{th} element of $\mathbf{N}_{t,m}$ is $(1 - r_{i,j})^t \prod_{k=0}^{t-1}(1 - \frac{1}{2N_{e_{k,m}}})$, and $\circ$ indicates Hadamard (i.e. element-wise) product. We then sum the $\boldsymbol{\mathcal{L}}_{t,m}$'s for all the generations over which $\Delta\mathbf{p}_m$ is measured to obtain $\boldsymbol{\mathcal{L}}_m = \sum_{t=t_m}^{\tau_m - 1} \boldsymbol{\mathcal{L}}_{t,m}$. Note that the expected predicable change due to selection between generations t and τ is $E[\Delta\mathbf{p}_m] = \boldsymbol{\mathcal{L}}_m\boldsymbol{\mu}_{\bar{\alpha}}$ and the among-replicate covariance in allele frequency change is $COV(\Delta\mathbf{p}_m, \Delta\mathbf{p}_n^{\top}) = \boldsymbol{\mathcal{L}}_m\mathbf{V}_{\bar{\alpha}}\boldsymbol{\mathcal{L}}_n$.

Step 3: Compute drift covariance and the projection matrix

Instead of working with raw allele frequency changes, we work with *projected* allele frequency changes. The projection matrix $\mathbf{P}$ is chosen to be such that (1) the number of dimensions of the dataset are reduced to the minimum of N_0 and n_L (typically $N_0 << n_L$), and (2) the allele frequency changes due to drift are independent and identically distributed. First, we perform eigen-decomposition of $\mathbf{L}_0$ (or, equivalently, singular-value decomposition of $\mathbf{C_0}$) and store the eigenvectors corresponding to non-zero eigenvalues $(\mathbf{U_L})$. The (co)variance in allele frequency change due to drift in replicate m is $\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)}$ where $\mathbf{M}^{(m)} = \sum_{t=t_m}^{\tau_m - 1} \mathbf{M}_{t,m} \circ \mathbf{N}_{t,m}$ and the ij^{th} element of $\mathbf{M}_{t,m}$ is $(1 - r_{i,j})/N_{E_{t,m}}$. We then perform eigen-decomposition of the drift covariance expressed using the eigen-vectors of $\mathbf{L}_0$, i.e. $\mathbf{U_L}^{\top}(\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)})\mathbf{U_L}$. Let $\mathbf{U}_2$ be a matrix having the eigenvectors of this $\mathbf{U_L}^{\top}(\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)})\mathbf{U_L}$ as columns, and let $\mathbf{D}_2$ be a diagonal matrix of square-rooted eigenvalues of this $\mathbf{U_L}^{\top}(\tilde{\mathbf{L}}_0 \circ \mathbf{M}^{(m)})\mathbf{U_L}$. We can then calculate the projection matrix $\mathbf{P} = \mathbf{D}_2^{-1}\mathbf{U}_2^{\top}\mathbf{U_L}^{\top}$, and obtain the projected allele frequency change vector for replicate m as $\Delta\overrightarrow{\mathbf{p}}_m = \mathbf{P}\Delta\mathbf{p}_m$. Note that the mean and covariance of the projected allele frequency changes due to predictable selection are $E[\Delta\overrightarrow{\mathbf{p}}_m] = \mathbf{P}_m\boldsymbol{\mathcal{L}}_m\boldsymbol{\mu}_{\bar{\alpha}}$ and $COV(\Delta\overrightarrow{\mathbf{p}}_m, \Delta\overrightarrow{\mathbf{p}}_n) = \mathbf{P}_m\boldsymbol{\mathcal{L}}_m\mathbf{V}_{\bar{\alpha}}\boldsymbol{\mathcal{L}}_n\mathbf{P}_n^{\top}$.

Step 4: Preparing data for model

Next, using our models $\boldsymbol{\mu}_{\bar{\alpha}} = \beta_{\bar{\alpha}}^{(1)}(\mathbf{p}_0 - \mathbf{q}_0)$ and $\mathbf{V}_{\bar{\alpha}} = \sigma_{\bar{\alpha}}^2 \mathbf{L}_0^{p_{\bar{\alpha}}}$ we calculate the fixed effect covariate $\mathbf{P}\boldsymbol{\mathcal{L}}_m(\mathbf{p}_0 - \mathbf{q}_0)$, and the covariance structure of the locus effects as $\mathbf{V}_{m,n} \propto \mathbf{P}\boldsymbol{\mathcal{L}}_m \mathbf{L}_0^{p_{\bar{\alpha}}} \boldsymbol{\mathcal{L}}_n \mathbf{P}^{\top}$.

Step 5: Fit linear mixed models

For a given $p_{\bar{\alpha}}$, our goals is to fit a linear mixed model with the projected allele frequency change in all replicates $(\Delta \vec{\mathbf{p}})$ as the response variable and a fixed effect predictor given by $\mathbf{P}\boldsymbol{\mathcal{L}}_m(\mathbf{p}_0 - \mathbf{q}_0)$ with a coefficient $\beta^{(1)}$ to be estimated. We treat locus effects to be random with a covariance structure assumed proportional to $\mathbf{V}_{m,n} \propto \mathbf{P}\boldsymbol{\mathcal{L}}_m \mathbf{L}_0^{p_{\bar{\alpha}}} \boldsymbol{\mathcal{L}}_n \mathbf{P}^{\top}$, with a proportionality constant $\sigma_{\bar{\alpha}}^2$ to be estimated. We use the R function *'optim()'* to find $\widehat{p_{\bar{\alpha}}}$ that maximises the conditional likelihood of the above model. We then use $\widehat{p_{\bar{\alpha}}}$, fit the above model, and obtain $\widehat{\beta^{(1)}}$ and $\widehat{\sigma_{\bar{\alpha}}^2}$.

Step 6: Estimate $E[V_{\bar{A}}(0)]$

The final step is to use the estimates obtained in the previous step $(\widehat{p_{\bar{\alpha}}}, \widehat{\beta^{(1)}}, \text{ and } \widehat{\sigma_{\bar{\alpha}}^2})$ to obtain $E[\widehat{V_{\bar{A}}(0)}] = Tr(\widehat{\mathbf{L}_0 \mathbf{V}_{\bar{\alpha}}}) + \boldsymbol{\mu}_{\bar{\alpha}}^{\top} \widehat{\mathbf{L}_0 \boldsymbol{\mu}_{\bar{\alpha}}}$. We obtain $Tr(\widehat{\mathbf{L}_0 \mathbf{V}_{\bar{\alpha}}})$ as $\widehat{\sigma_{\bar{\alpha}}^2} Tr(\mathbf{L}_0 \mathbf{L}_0^{\widehat{p_{\bar{\alpha}}}})$, and bias corrected $\boldsymbol{\mu}_{\bar{\alpha}}^{\top} \widehat{\mathbf{L}_0 \boldsymbol{\mu}_{\bar{\alpha}}}$ as $\widehat{\beta^{(1)}}^2 (\mathbf{p}_0 - \mathbf{q}_0)^{\top} \mathbf{L}_0 (\mathbf{p}_0 - \mathbf{q}_0) - VAR(\widehat{\beta^{(1)}})(\mathbf{p}_0 - \mathbf{q}_0)^{\top} \mathbf{L}_0 (\mathbf{p}_0 - \mathbf{q}_0)$.

 # S10   Supplementary figures

Supplementary Figure 1: Results of full simulations with a burn-in phase of 25,000 generations (map length in the history phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, the mean of the gamma distribution from which effect sizes for log absolute fitness are sampled for non-neutral mutations $(E[|\eta|]) = 0.03$, at different levels of the map length in the history phase (0.5 morgan: green circles, 5 morgan: magenta triangles, 50 morgans: blue squares, and 250 morgans: grey plus symbols). The coloured solid lines represent regression lines for estimates of V_A vs true values of V_A.

Supplementary Figure 2: (a) Fraction of the total additive genic variance ($V_{\bar{a}}(0)$) lost during the experiment (averaged over all the replicate populations) plotted as a function of the mean of the gamma distribution from which effect sizes for log absolute fitness are sampled for non-neutral mutations ($E[|\eta|]$). Each boxplot represents 100 independent simulations (for the corresponding level of $E[|\eta|]$) shown in Figure 5D. (b) The error in our estimates of V_A plotted versus the total additive genic variance (V_a) lost during the experiment (averaged over all the replicate populations) for simulations where the $E[|\eta|]$ was either 0.02 (green circles), 0.03 (blue triangles), or 0.06 (grey squares). The solid line has an intercept of 0 and a slope of -1. (c) The contribution of each non-neutral locus (averaged over all the replicate populations) to V_a lost during the experiment plotted versus the absolute average effect for relative fitness ($|\alpha|$) for that locus for a simulation with $E[|\eta|] = 0.06$ that had the maximum V_a loss (grey squares) and a simulation with $E[|\eta|] = 0.02$ that had a comparable true initial V_A but the minimum V_a loss (green circles) in the experiment. The solid curve represents the additive genic variance lost when a singleton goes extinct. (d) The total V_a lost at groups of loci classified by dividing $|\alpha|$ into intervals of 0.01 for the simulation with the maximum V_a loss (grey) and the simulation with the minimum V_a loss (green) in the experiment. In the simulation with the maximum V_A loss – in which the mean $|\alpha|$ for non-neutral segregating loci was 0.0410 – more than 50% of the total loss in V_a was driven by just 0.96% loci whose $|\alpha|$ was greater than 0.3.

Supplementary Figure 3: Residual variance in the models analysing the results of full simulations shown in Figure 4 using either the number of individuals (1,000) instead of the N_E or the true N_E.

Full simulations
(with dominance in the experiment phase)

Supplementary Figure 4: Results of full simulations (map lengths in the history phase = 0.5 morgans and experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, the mean of the gamma distribution from which effect sizes for log absolute fitness are sampled for non-neutral mutations ($E[|\eta|]$) = 0.03), in which dominance effects were switched on only in the experiment phase using different degrees of dominance: $|\kappa| = 0$ (green circles), $|\kappa| = 0.5$ (blue triangles), and $|\kappa| = 0.9$ (grey squares). (a) A scatter plot of estimates of V_A vs true values of V_A. The solid black line indicates the 1:1 line. The coloured lines represent regression lines for estimates of V_A vs true values of V_A. The inference of V_A was obtained by modelling the mean and the (co)variance of the average effects for relative fitness as $\boldsymbol{\mu}_{\bar{\alpha}} = \beta_{\bar{\alpha}}^{(1)}(\mathbf{p}_0 - \mathbf{q}_0)$ and $\mathbf{V}_{\bar{\alpha}} = \sigma_{\bar{\alpha}}^2 \mathbf{L}_0^{p_{\bar{\alpha}}}$, respectively. (b)-(d) Histograms of the estimates of $p_{\bar{\alpha}}$, $\beta_{\bar{\alpha}}^{(1)}$, and the residual variance, respectively. The horizontal dashed lines indicate null expectations (0 for $p_{\bar{\alpha}}$ and $\beta_{\bar{\alpha}}^{(1)}$, and 1 for the residual variance).

xxxi

Supplementary Figure 5: The true additive genetic variance for relative fitness (V_A) plotted versus the true additive genic variance for relative fitness (V_a) in full simulations with a burn-in phase of 25,000 generations performed using a map length in the history phase equal to either 0.5 morgan (a) or 5 morgans (b) with dominance effects switched on in the last 5,000 generations of the history phase. Three different degrees of dominance were used: $|\kappa| = 0$ (green circles), $\kappa = 0.5$ (blue triangles), and $\kappa = 0.75$ (magenta plus symbols). The mean of the gamma distribution from which effect sizes for log absolute fitness are sampled for non-neutral mutations $(E[|\eta|])$ was 0.03. The solid black lines indicates the 1:1 line. The coloured lines represent regression lines for true V_A vs true true V_a.

Full simulations
(incorporating covariance due to pool-seq in the analyses)

Legend: ● Exact ▲ 1000x ■ 500x + 100x ⊠ 1000x-overdispersed

Supplementary Figure 6: Scatter plots of estimates of V_A vs true values of V_A for full simulations with a burn-in phase of 25,000 generations (map length in the history phase = 0.5 morgan, map length in the experiment phase = 2 morgans, number of replicate populations = 10, population size = 1,000, number of generations = 3, the mean of the gamma distribution from which effect sizes for log absolute fitness were sampled for non-neutral mutations ($E[|\eta|]) = 0.02$, and no dominance (i.e. $\kappa = 0$)) using either exact allele frequencies in the experiment phase (green circles), or allele frequencies in the experiment phase obtained via simulated pool-seq implemented without any overdispersion in the number of reads mapping to an individual, at three different levels of coverage (expected number of reads mapping a segregating site: 1000x (magenta triangles), 500x (blue squares), and 100x (grey plus symbols)), as well as estimates obtained from simulated pool-seq implemented with overdispersion ($V_x = log(2)$) in the number of reads mapping to an individual, at 1000x coverage (orange empty boxes with crosses). Reads were modelled to be 37 base-pairs long. The solid black line indicates the 1:1 line. The coloured lines represent regression lines for estimates of V_A vs true values of V_A. Estimates of V_A were obtained by incorporating the expected covariance structure due to pool-seq sampling in the models.

xxxiii

References

Bohrnstedt, G. W., and A. S. Goldberger, 1969 On exact covariance of products of random variables. Journal of the American Statistical Association **64**: 1439–1442.

Buffalo, V., and G. Coop, 2019 The linked selection signature of rapid adaptation in temporal genomic data. Genetics **213**: 1007–1045.

Buffalo, V., and G. Coop, 2020 Estimating the genome-wide contribution of selection to temporal allele frequency change. Proceedings of the National Academy of Sciences **117**: 20672–20680.

Bulmer, M. G., 1980 *The mathematical theory of quantitative genetics.*. Clarendon Press.

de Los Campos, G., D. Sorensen, and D. Gianola, 2015 Genomic heritability: what is it? PLoS Genetics **11**: e1005048.

Ethier, S. N., and T. Nagylaki, 1980 Diffusion approximations of Markov chains with two time scales and applications to population genetics. Advances in Applied Probability **12**: 14–49.

Falconer, D., and T. Mackay, 1996 *Introduction to Quantitative Genetics, 4th edn*. Addison Wesley Longman, Harlow.

Gianola, D., G. de Los Campos, W. G. Hill, E. Manfredi, and R. Fernando, 2009 Additive genetic variability and the Bayesian alphabet. Genetics **183**: 347–363.

Good, B. H., 2022 Linkage disequilibrium between rare mutations. Genetics **220**: iyac004.

Haldane, J. B. S., 1919 The combination of linkage values, and the calculation of distances between the loci of linked factors. Journal of Genetics **8**: 299–309.

Hill, W., and A. Robertson, 1968 Linkage disequilibrium in finite populations. Theoretical and applied genetics **38**: 226–231.

Kojima, K., 1959 Role of epistasis and overdominance in stability of equilibria with selection. Proceedings of the National Academy of Sciences **45**: 984–989.

Lynch, M., and B. Walsh, 1998 *Genetics and analysis of quantitative traits*. Sinauer Associates, Inc., Sunderland.

McCullagh, P., and J. A. Nelder, 1989 *Generalised Linear Models*. Monographs on statistics and applied probability. Chapman & Hall, Cambridge, 2^{nd} edition.

McVean, G. A., 2002 A genealogical interpretation of linkage disequilibrium. Genetics **162**: 987–991.

Meuwissen, T. H., B. J. Hayes, and M. Goddard, 2001 Prediction of total genetic value using genome-wide dense marker maps. genetics **157**: 1819–1829.

Ohta, T., and M. Kimura, 1969 Linkage disequilibrium due to random genetic drift. Genetics Research **13**: 47–55.

Ohta, T., and M. Kimura, 1971 Linkage disequilibrium between two segregating nucleotide sites under the steady flux of mutations in a finite population. Genetics **68**: 571–580.

Santiago, E., and A. Caballero, 1998 Effective size and polymorphism of linked neutral loci in populations under directional selection. Genetics **149**: 2105–2117.

Song, Y. S., and J. S. Song, 2007 Analytic computation of the expectation of the linkage disequilibrium coefficient r^2. Theoretical population biology **71**: 49–60.

Sved, J. A., and W. G. Hill, 2018 One hundred years of linkage disequilibrium. Genetics **209**: 629–636.

Weir, B. S., 1996 *Genetic data analysis II: methods for discrete population genetic data*. Sunderland.

Weir, B. S., and C. C. Cockerham, 1989 Complete characterization of disequilibrium at two loci. In M. Feldman, editor, *Mathematical Evolutionary Theory*. Princeton University Press Princeton, NJ, 86–110.

Wright, S., 1938 Size of population and breeding structure in relation to evolution. Science **87**: 340–431.

Zeng, J., R. De Vlaming, Y. Wu, M. R. Robinson, L. R. Lloyd-Jones *et al.*, 2018 Signatures of negative selection in the genetic architecture of human complex traits. Nature genetics **50**: 746–753.