Questioning the Evidence for Host-Symbiont Codiversification in Mycorrhizal Symbioses

Fantine Bodin[1], Hélène Morlon[1], Benoît Perez-Lamarque[2]

[1]Institut de Biologie de l'ENS (IBENS), École Normale Supérieure, CNRS, INSERM, Université PSL, 46 rue d'Ulm, 75005 Paris, France.

[2]Université de Toulouse, Toulouse INP, CNRS, IRD, CRBE, Toulouse, France

Contact: fantine.bodin@ens.psl.eu (ORCID: 0009-0004-4376-9503),

helene.morlon@bio.ens.psl.eu (ORCID: 0000-0002-3195-7521),

benoit.perez-lamarque@utoulouse.fr (ORCID: 0000-0001-7112-7197)

Key words

Abstract

A large majority of plants form mycorrhizal symbioses. The ecological importance of these mutualistic interactions has sparked interest in their long evolutionary history. By examining interaction networks and phylogenetic trees, several studies have suggested that plants codiversify with their associated mycorrhizal fungi. However, recent research has demonstrated that phylogenetic congruence (often interpreted as codiversification) has often been conflated with another pattern called cophylogenetic signal (*i.e.,* closely related plants interacting with closely related fungi and *vice-versa*), which may arise from different biological processes. We performed cophylogenetic analyses on 29 diverse mycorrhizal networks to reevaluate the evidence for codiversification in mycorrhizal symbioses. We found significant cophylogenetic signal but no phylogenetic congruence: closely related plants interact with closely related fungi, but their phylogenies do not match. Instead of codiversification, this finding suggests that plants and fungi undergo diffuse coevolution, likely driven by trait-matching, where compatible, evolutionarily conserved plant and fungal traits govern their interactions. Our work highlights the importance of appropriately interpreting the cophylogenetic methods used to study the macroevolution of plants and their mycorrhizal fungi. It suggests that previous evidence of codiversification in mycorrhizal symbioses actually detected cophylogenetic signal, indicating that there is no evidence of codiversification occurring during the evolution of mycorrhizal symbioses.

Main text

More than 90% of vascular plants form mycorrhizal symbioses (Brundrett and Tedersoo, 2018). These mutualistic associations are crucial for nutrition of most plants and their associated fungi and thus for the functioning of whole ecosystems (Smith and Read, 2008; Tedersoo et al., 2020; van der Heijden et al., 2015). Mycorrhizal symbioses originated more than 400 million years ago, likely facilitating the transition of plants from water to land (Remy et al., 1994; Strullu-Derrien et al., 2018). Over time, these associations evolved and diversified, leading to the recognition of several types of mycorrhizal symbioses (van der Heijden et al., 2015). The ancient origin and the ecological importance of these interactions have sparked interest in their assembly and evolution. In natural communities, mycorrhizal fungi typically interact with multiple plant species, rather than forming exclusive one-to-one associations (van der Heijden et al., 2015). Therefore, they shape complex interaction networks (Chagnon et al., 2012; Jacquemyn et al., 2015; Perez-Lamarque et al., 2022; Petrolli et al., 2022; Rimington et al., 2019; Sepp et al., 2019). By considering the structure of mycorrhizal networks and the phylogenetic trees of plants and their associated mycorrhizal fungi, it has been suggested that phylogenetic congruence can occur in mycorrhizal symbioses (Arifin et al., 2023; Merckx and Bidartondo, 2008; Van Galen et al., 2023). Whilst these studies discuss multiple processes by which these patterns could arise, one possibility is that plants and their associated mycorrhizal fungi could be undergoing simultaneous speciation events (*i.e.,* cospeciation), leading to codiversification (see Table 1 for definitions).

Testing for the occurrence of codiversification in host-symbiont associations is notoriously challenging, and has spurred the development of several cophylogenetic methods (Blasco-Costa et al., 2021; De Vienne et al., 2013; Dismukes et al., 2022; Page, 1994; Perez-Lamarque and Morlon, 2024; Ronquist, 2002). However, two distinct patterns – cophylogenetic signal and phylogenetic congruence – have often been conflated (Perez-Lamarque and Morlon, 2024) (Figure 1; Table 1). In the context of mycorrhizal symbioses, cophylogenetic signal indicates that closely related plants tend to interact with closely related mycorrhizal fungi and *vice-versa*. In contrast, phylogenetic congruence is a more stringent, specific case of cophylogenetic signal, occurring when the phylogenetic trees of plants and their associated mycorrhizal fungi closely mirror each other. Codiversification refers to this pattern of phylogenetic congruence combined with matching divergence times, reflecting parallel diversification histories in plants and their associated fungi (Suzuki et al., 2022). Distinguishing these two patterns of cophylogenetic signal and phylogenetic congruence is crucial because they may stem from different biological processes. A cophylogenetic signal may simply arise if present-day mycorrhizal interactions assemble according to the trait matching of evolutionarily conserved traits (Perez-Lamarque and Morlon, 2024). It can also be generated by past biogeographic events that have constrained the distributions of extant species. In contrast, phylogenetic

congruence arises from more stringent scenarios. For instance, codiversification can
result from repeated cospeciation events driven by phylogenetic tracking, or from
successive vicariance events that repeatedly affect both plants and their associated
mycorrhizal fungi (Althoff et al., 2014; De Vienne et al., 2013; Perez-Lamarque and
Morlon, 2024). Global-fit methods such as ParaFit or PACo (Balbuena et al., 2013;
Hutchinson et al., 2017; Legendre et al., 2002) can assess cophylogenetic signal, but only
event-based methods such as Jane or eMPRess (Conow et al., 2010; Santichaivekin et al.,
2021) can evaluate phylogenetic congruence.

Here, we reevaluate the evidence for codiversification in mycorrhizal symbioses in light of
recent cophylogenetic developments (Perez-Lamarque and Morlon, 2024), using an
extensive collection of mycorrhizal networks. We manually screened the literature and
selected 29 networks from 13 different publications, ensuring coverage of various spatial
and evolutionary scales, as well as all four main types of mycorrhizal symbioses
(arbuscular mycorrhizae, ectomycorrhizae, orchid and ericoid mycorrhizae). For each
mycorrhizal network, if the phylogenetic trees were not already available, we retrieved
representative DNA sequences for both plant and mycorrhizal fungi, using multiple genes
when possible (Supplementary Table 1). Sequences were aligned using MAFFT (Katoh,
2002) and refined using trimAl (Capella-Gutiérrez et al., 2009). Robust phylogenetic trees
were then reconstructed using IQ-TREE (Nguyen et al., 2015), constraining deep nodes by
a robust backbone tree when necessary (Supplementary Table 1). To assess
cophylogenetic signal, we applied the global-fit method PACo (Balbuena et al., 2013;
Hutchinson et al., 2017) to each mycorrhizal network. We then evaluated phylogenetic
congruence using the event-based model eMPRess (Santichaivekin et al., 2021). This tool
reconciles phylogenetic trees by fitting of model of phylogenetic tracking that allows for
events of cospeciation, transfer, loss, and duplication. eMPRess returns the number of
each event and assesses the significance of the reconciled scenario through
permutations. We considered phylogenetic congruence to be supported when the
reconciled scenario had a significantly low cost (p-value < 0.05) and included more
cospeciation events than transfer events (Dorrell et al., 2021; Groussin et al., 2017; Perez-
Lamarque and Morlon, 2024).

We detected a significant cophylogenetic signal in 13 out of 29 mycorrhizal networks
(Figure 2). This pattern was consistently observed across the different types of
mycorrhizal symbioses, regardless of the spatial and evolutionary scales (Supplementary
Table 1). This confirms that there is evidence of frequent cophylogenetic signal in
mycorrhizal associations. In contrast, phylogenetic congruence was absent in all 29
networks (Figure 2). Our results were consistent across phylogenetic reconstruction
methods (Supplementary Table 1), meaning that uncertainty in fungal phylogenies does
not explain the observed patterns. The fungal phylogenies of 4 of the 29 networks were
built with only the ITS region, which may poorly resolve deep evolutionary relationships,

but generally supports recent splits. Our results were robust despite uncertain deep phylogenetic resolution in some networks because recent splits play the major role in the detection of phylogenetic congruence, while incongruence at deeper nodes can often be reconciled by a few host-transfer events using eMPRess. Therefore, given the high diversity of mycorrhizal networks analyzed, the consistency of our results indicates that they can be generalized beyond the specific sets of species and systems included in our analyses. Our findings suggest that previous studies that proposed codiversification in mycorrhizal symbioses based on visual inspections or global-fit methods detected cophylogenetic signal rather than phylogenetic congruence, and that there is in fact no evidence of codiversification between plants and their associated mycorrhizal fungi.

The absence of codiversification in mycorrhizal symbioses may stem from the fact that most mycorrhizal partners do not engage in highly specialized or one-to-one associations, but instead participate in complex interaction networks characterized by high partner sharing (Arifin et al., 2023; Hayward and Horton, 2014; Öpik et al., 2009; Perez-Lamarque et al., 2022; Sepp et al., 2019; Toju et al., 2016; Van Galen et al., 2023; Zhao et al., 2021). Such generalized interactions reduce the likelihood of frequent cospeciation (Althoff et al. 2014; Poisot et al. 2015), although our results do not rule out the occurrence of rare cospeciation events that may sporadically happen in some mycorrhizal associations and whose signal may be blurred by other events. While our findings suggest that codiversification is uncommon in mycorrhizal networks, this result also does not rule out the role of coevolution (Brundrett 2002, van der Heijden 2015). The widespread presence of cophylogenetic signal suggests that closely related plants tend to interact with closely related fungi. At regional and global scales, this signal could be at least partially explained by the biogeography of plants and/or mycorrhizal fungi. Vicariance events and dispersal limitations may lead to allopatric speciations and geographically independent radiations in both plants and fungi. As a result, closely related plants and closely related fungi tend to be present in similar places (Tedersoo et al., 2014), generating cophylogenetic signal (Perez-Lamarque and Morlon, 2024). However, we also observed cophylogenetic signal in mycorrhizal networks at local scale within individual communities (Supplementary Table 1), which cannot be due to biogeography. Therefore, the cophylogenetic signal we observed is likely explained by the fact that mycorrhizal interactions require a minimal level of trait matching, and that the involved mycorrhizal traits are evolutionary conserved. In this scenario, diffuse coevolution between interacting plants and fungi drives the evolutionary conservatism of key mycorrhizal traits (Cairney, 2000), such as plant root architecture, fungal nutrient exchange efficiency, or molecular signaling between plants and fungi (Castanedo et al., 2025; Delaux and Schornack, 2021; Karandashov et al., 2004; van der Heijden et al., 2015).

Our study highlights the importance of careful interpretation of cophylogenetic methods,
and suggests that mycorrhizal symbionts did not codiversify with their host plants.
Evolutionary history nevertheless plays an important role in shaping plant-mycorrhizal
networks, as evidenced by the phylogenetic conservatism of interactions. Trait matching
and trait conservatism likely mediates this phylogenetical signal, suggesting that a deeper
understanding of the functional traits underlying mycorrhizal compatibility is key to
understand the evolution of mycorrhizal symbioses. Finally, synthesizing our analyses
with similar studies in other plant-associated interactions, such as pollination, seed
dispersal, or parasitic interactions (Asar et al., 2022; Jackson, 2004; Russo et al., 2018;
Taengon et al., 2025), would help clarifying whether codiversification has ever play a
significant role in the evolutionary history of land plants.
174

 Tables

 Table 1: Cophylogenetic definitions in the context of mycorrhizal symbioses. These
 definitions are following the terminology of Perez-Lamarque and Morlon (2024).

Terminology	Process or pattern	Definition
Codiversification/ Co-cladogenesis:	Pattern	Parallel diversification of plant and mycorrhizal fungal lineages, in which the two clades exhibit phylogenetic congruence and matching divergence times. It can be generated through processes of phylogenetic tracking or successive vicariance events repeatedly affecting both clades.
Coevolution:	Process	Reciprocal evolutionary changes driven by selective pressures between two interacting lineages (pairwise coevolution). When more than two lineages are involved, the process is referred to as **diffuse coevolution**.
Cophylogenetic signal:	Pattern	Tendency of closely related plant species to interact with closely related mycorrhizal fungal partners (and *vice-versa*).
Cospeciation:	Process	Concomitant speciation event in a plant lineage and its associated mycorrhizal fungal lineage.
Phylogenetic congruence:	Pattern	Strong similarity between the phylogenetic trees of interacting plant and mycorrhizal fungal clades in terms of topology and relative branch lengths. When the divergence times of the two clades also coincide, this pattern corresponds to codiversification.
Phylogenetic tracking:	Process	Repeated plant speciation events leading to subsequent speciations in their associated mycorrhizal fungal lineages (**cospeciations**), for example through vertical transmission that isolates fungal populations between diverging plant lineages. Fungal lineages may also undergo host transfers, intra-host duplications, or losses at low frequency. Phylogenetic tracking typically results in a pattern of codiversification.
Trait matching	Process	Association of plants and fungi because of the complementarity of their mycorrhizal traits (for

		example plant root architecture, fungal nutrient exchange efficiency, or molecular signaling between plants and fungi). When these traits tend to be evolutionarily conserved, trait matching typically results in cophylogenetic signal.
Tree topology:	Pattern	Branching structure of the different nodes of a phylogenetic tree that does not consider absolute branch lengths (*i.e.* evolutionary time).

Figures

Figure 1: Cophylogenetic signal *versus* phylogenetic congruence. a. Cophylogenetic signal: closely related plants tend to interact with closely related mycorrhizal fungi and *vice-versa* (this pattern can be generated by trait matching of evolutionarily conserved traits, as represented here with colors representing trait values, or by past biogeographic events leading to closely related species being present in similar places). **b.** Phylogenetic congruence: the phylogenetic trees of plants and their associated fungi mirror one another (this pattern can be generated by codiversification, *i.e.* concomitant diversification events happening in both plant and fungal interacting lineages; see Table 1 for a list of definitions of cophylogenetic patterns and processes).

Figure 2: Cophylogenetic analysis of 29 mycorrhizal networks. For each
mycorrhizal network, we report its eMPRess p-value as a function of the ratio of
cospeciation to transfer events inferred by eMPRess. The grey regions indicate
whether there is phylogenetic congruence: the dark grey area indicates significant
phylogenetic congruence (eMPRess p-value < 0.05 and cospeciation / transfer ratio >
1), whereas the light grey areas indicate no significant phylogenetic congruence
(eMPRess p-value > 0.05 and/or cospeciation / transfer ratio < 1). In addition, the
shape of each symbol reflects the significance of the cophylogenetic signal inferred
using PACo: a dot represents a network with significant cophylogenetic signal (PACo
p-value < 0.05), whereas a cross represents a network without cophylogenetic signal
(PACo p-value > 0.05). The size of each shape indicates the type of genetic data
available for reconstructing the fungal phylogenetic trees. Detailed information on
each network can be obtained in Supplementary Table 1.
208

Methods

Screening for available mycorrhizal networks

We manually screened the literature to compile as many mycorrhizal networks as possible (Supplementary Table 1). We targeted publications that provided freely accessible interaction networks, along with the phylogenetic trees or the DNA sequences of both plants and mycorrhizal fungi. We retained networks from natural habitats and with taxonomic delineations at the plant species level. Fungal operational taxonomic units (OTUs) were typically delineated using a 97% similarity threshold. We aimed to include all major types of mycorrhizal symbioses: arbuscular mycorrhizae (n=11), ectomycorrhizae (n=5), orchid mycorrhizae (n=7), and ericoid mycorrhizae (n=6). We also selected networks encompassing a broad range of spatial scales – from communities in specific habitats of La Réunion (Perez-Lamarque et al., 2022) to the global arbuscular mycorrhizal network between Glomeromycotina fungi and land plant species (Öpik et al., 2010; Perez-Lamarque et al., 2020) – and evolutionary scales – from a 14 million-year-old ectomycorrhizal association between Pisonieae and their thelephoroid fungal symbionts (Hayward and Horton, 2014) to more than 450 million years of diversification of all land plants forming arbuscular mycorrhizae. Our networks also encompass a diversity of sampling strategies: from sampling all the species present in a local community (for example Perez-Lamarque et al., 2022; Sepp et al., 2019), to intentionally targeting on some specific plant clades only (as in Arifin et al., 2023; Shefferson et al., 2010) or gathering all plant-fungal associations available at regional or global scale (for instance Öpik et al., 2009; Perez-Lamarque et al., 2020).

In total, we studied 29 networks from 13 different publications (Arifin et al., 2023; Hayward and Horton, 2014; Martos et al., 2012; Merckx and Bidartondo, 2008; Öpik et al., 2010, 2009; Otero et al., 2011; Perez-Lamarque et al., 2022; Perez-Lamarque et al., 2020; Sepp et al., 2019; Shefferson et al., 2010; Toju et al., 2016; Van Galen et al., 2023; Zhao et al., 2021).

Recovering phylogenetic trees

For some publications, robust phylogenetic trees were already publicly available (Supplementary Table 1). For the others, we retrieved DNA sequences from GenBank, aligned them with MAFFT (Katoh, 2002), trimmed them with trimAl (Capella-Gutiérrez et al., 2009), and then reconstructed the phylogenetic trees with IQ-TREE (Nguyen et al., 2015). When multiple sequences were available for a single species, we selected one representative (Supplementary Table 2). For the plants, we consistently used sequences from at least two genes, selected from trnL, ITS, matK, matR, 18S, and ATPa (Supplementary Table 1). When no genetic data were available in the original publications, we reconstructed the missing plant phylogenetic trees with V.PhyloMaker2 (Jin and Qian, 2022). For the fungal phylogenetic reconstructions, the available sequences primarily included the SSU rRNA gene and/or the ITS region (ITS1, ITS2, or the

full ITS1-5.8S-ITS2 fragment), sometimes in addition to other genes (28S, EF1 alpha,
TOR2, ML 4-5). We considered robust the phylogenies obtained with multiple genes.
While the SSU rRNA gene alone can support robust phylogenetic tree, the fungal ITS
region is often too variable, especially ITS1 or ITS2 alone, to reliably resolve deeper
evolutionary relationships (Perez-Lamarque et al., 2022; Schoch et al., 2012). Therefore,
to obtain robust phylogenetic trees when only sequences of partial ITS1 or ITS2 were
available, we informed the IQ-TREE reconstructions with a robust backbone tree built
from multiple-gene datasets (following Talavera et al., 2022). Specifically, we used the
backbone trees of Weiß et al. (2016) for Sebacinales, of Johnston et al. (2019) for
Helotiales, and of Zhang et al. (2025) for Cantharellales (Supplementary Table 1). This
backbone approach improves bootstrap branch support, raising values >70% for most
nodes. In total, 15 fungal phylogenetic trees were reconstructed using either multi-gene
datasets or a backbone-tree approach, 10 using the SSU rRNA gene, 3 using the full ITS
region, and 1 using only ITS1, as no additional sequences or backbone trees were
available for this mycorrhizal system (Supplementary Table 1).

Evaluating cophylogenetic signal
We assessed cophylogenetic signal, *i.e.* whether closely related plants tend to interact
with closely related fungi, using the global-fit method PACo (Balbuena et al., 2013;
Hutchinson et al., 2017). This method returns a R^2 and a p-value obtained by comparison
with permutations. We ran 10,000 permutations for each mycorrhizal network.
Additionally, we checked the results with the global-fit method parafit (Legendre et al.,
2002; Paradis et al., 2004). Both methods yielded similar results (Supplementary Table 1).

Evaluating phylogenetic congruence
We assessed phylogenetic congruence, *i.e.* whether the phylogenetic trees of plants and
mycorrhizal fungi tend to mirror one another, with the event-based model eMPRess
(Santichaivekin et al., 2021). eMPRess assumes a process of phylogenetic tracking and
tries to reconcile the plant and fungal phylogenetic trees by fitting, using maximum
parsimony, events of cospeciation, host transfer, duplication, and loss, with a given cost
associated to each event. We ran eMPRess with the following costs: 0 for a cospeciation
event, 4 for a duplication event, 1 for a loss event, and 1 for a transfer event. We obtained
similar results using different cost values (Supplementary Table 1). To assess the
significance of each eMPRess reconciliation, we performed 1,000 permutations of the
mycorrhizal associations. We considered a reconciliation as significant when the total
cost was lower than 95% of the costs obtained from the permuted data (eMPRess p-value
< 0.05) and when the number of cospeciation events was larger than the number of
transfer events (cospeciation / transfer ratio > 1; Perez-Lamarque & Morlon, 2024).
Indeed, transfer events can obscure cospeciation signals and generate a symbiont
phylogeny that no longer matches the host phylogeny.

Importantly, eMPRess, like most event-based methods, evaluates phylogenetic
congruence by considering only the topologies of the plant and fungal phylogenetic trees.
It does not take into account whether the divergence times of corresponding plant and
fungal nodes are concomitant to fit a cospeciation event. Thus, to conclude that
codiversification has occurred, both a significant phylogenetic congruence (as detected
by eMPRess) and coincident divergence times between the two clades are needed. In this
study, we did not evaluate the latter aspect, as most fungal phylogenies were not time-
calibrated.
As phylogenetic congruence can only exist if one-to-one host-symbiont interactions are
frequent, eMPRess cannot account for symbionts interacting with multiple hosts, like
most event-based approach (De Vienne et al., 2013; Dismukes et al., 2022). Therefore,
when a fungal species interacted with several plant species, we considered two
approaches (Perez-Lamarque & Morlon, 2024): (1) we only picked the most abundant
interaction (or one interaction at random when abundances were not available;
Supplementary Table 1), or (2) we randomly simulated bifurcating sub-trees for fungal
taxa with multiple plants, such that each tip in the fungal tree is associated with a single
plant tip (Satler et al., 2019). As the second approach did not change our findings, we only
reported the results of the first approach in the main text (Supplementary Table 1).
Finally, we verified that our results obtained with eMPRess were robust to the small size
of some networks and the phylogenetic uncertainty in the fungal trees. Regarding the
small sampling size, it has been demonstrated using simulations that eMPRess still
performs well when the number of codiversifying species in low (Perez-Lamarque et al.
2024). For the phylogenetic uncertainty, we ran eMPRess with different fungal
phylogenetic trees obtained with different reconstruction methods (ITS only or the
backbone approach) and obtained consistent results in all cases.
316

Acknowledgments

The authors acknowledge members of the 'Modeling biodiversity' lab at IBENS and Laura Van Galen for helpful discussions, as well as Alice Garcia for initiating the project with preliminary analyses.

The authors also acknowledge Arild Arifin and Zhongtao Zhao for sharing the interaction networks and phylogenetic trees of plants and mycorrhizal fungi they studied, as well as all other authors whose data were directly available in their publications.

Competing interests

The authors declare no conflict of interests.

Author contributions

FB, HM, and BPL designed the study, FB performed the analyses, FB and BPL wrote a first version of the manuscript and all authors contributed to the revisions.

Code availability

All the scripts for performing the analyses are available through the following link: https://github.com/fantine-bodin/Cophylogenetics_mycorrhizal-symbioses.git

References

Althoff, D.M., Segraves, K.A., Johnson, M.T.J., 2014. Testing for coevolutionary diversification: linking pattern with process. Trends Ecol. Evol. 29, 82–89. https://doi.org/10.1016/j.tree.2013.11.003

Arifin, A.R., Phillips, R.D., Linde, C.C., 2023. Strong phylogenetic congruence between *Tulasnella* fungi and their associated Drakaeinae orchids. J. Evol. Biol. 36, 221–237. https://doi.org/10.1111/jeb.14107

Asar, Y., Ho, S.Y.W., Sauquet, H., 2022. Early diversifications of angiosperms and their insect pollinators: were they unlinked? Trends Plant Sci. 27, 858–869. https://doi.org/10.1016/j.tplants.2022.04.004

Balbuena, J.A., Míguez-Lozano, R., Blasco-Costa, I., 2013. PACo: A novel Procrustes Application to Cophylogenetic analysis. PLoS ONE 8, e61048. https://doi.org/10.1371/journal.pone.0061048

Blasco-Costa, I., Hayward, A., Poulin, R., Balbuena, J.A., 2021. Next-generation cophylogeny: unravelling eco-evolutionary processes. Trends Ecol. Evol. 36, 907–918. https://doi.org/10.1016/j.tree.2021.06.006

Brundrett, M.C., Tedersoo, L., 2018. Evolutionary history of mycorrhizal symbioses and
global host plant diversity. New Phytol. 220, 1108–1115.
https://doi.org/10.1111/nph.14976
Cairney, J.W.G., 2000. Evolution of mycorrhiza systems. Naturwissenschaften 87, 467–
475. https://doi.org/10.1007/s001140050762
Capella-Gutiérrez, S., Silla-Martínez, J.M., Gabaldón, T., 2009. trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics 25, 1972–1973. https://doi.org/10.1093/bioinformatics/btp348
Castanedo, L., Bianconi, M., Pellen, T., Vernie, T., Lefort, C., Libourel, C., Le Ru, A.,
Rodde, N., Cauet, S., Callot, C., Forrest, L., Nishiyama, T., Wickell, D., Li, F.-W.,
Keller, J., Delaux, P.-M., 2025. Leafy liverwort genomes shed light on the evolution
of ericoid symbiosis. https://doi.org/10.1101/2025.04.01.646537
Chagnon, P., Bradley, R.L., Klironomos, J.N., 2012. Using ecological network theory to
evaluate the causes and consequences of arbuscular mycorrhizal community
structure. New Phytol. 194, 307–312. https://doi.org/10.1111/j.1469-
8137.2011.04044.x
Conow, C., Fielder, D., Ovadia, Y., Libeskind-Hadas, R., 2010. Jane: a new tool for the
cophylogeny reconstruction problem. Algorithms Mol. Biol. 5.
https://doi.org/10.1186/1748-7188-5-16
De Vienne, D.M., Refrégier, G., López-Villavicencio, M., Tellier, A., Hood, M.E., Giraud, T.,
2013. Cospeciation vs host-shift speciation: methods for testing, evidence from
natural associations and relation to coevolution. New Phytol. 198, 347–385.
https://doi.org/10.1111/nph.12150
Delaux, P.-M., Schornack, S., 2021. Plant evolution driven by interactions with symbiotic
and pathogenic microbes. Science 371, eaba6605.
https://doi.org/10.1126/science.aba6605
Dismukes, W., Braga, M.P., Hembry, D.H., Heath, T.A., Landis, M.J., 2022.
Cophylogenetic methods to untangle the evolutionary history of ecological
interactions. Annu. Rev. Ecol. Evol. Syst. 53, 275–298.
https://doi.org/10.1146/annurev-ecolsys-102320-112823
Dorrell, R.G., Villain, A., Perez-Lamarque, B., Audren De Kerdrel, G., McCallum, G.,
Watson, A.K., Ait-Mohamed, O., Alberti, A., Corre, E., Frischkorn, K.R., Pierella
Karlusich, J.J., Pelletier, E., Morlon, H., Bowler, C., Blanc, G., 2021. Phylogenomic
fingerprinting of tempo and functions of horizontal gene transfer within
ochrophytes. Proc. Natl. Acad. Sci. 118, e2009974118.
https://doi.org/10.1073/pnas.2009974118
Groussin, M., Mazel, F., Sanders, J.G., Smillie, C.S., Lavergne, S., Thuiller, W., Alm, E.J.,
2017. Unraveling the processes shaping mammalian gut microbiomes over
evolutionary time. Nat. Commun. 8, 14319.
https://doi.org/10.1038/ncomms14319
Hayward, J., Horton, T.R., 2014. Phylogenetic trait conservation in the partner choice of a
group of ectomycorrhizal trees. Mol. Ecol. 23, 4886–4898.
https://doi.org/10.1111/mec.12903
Hutchinson, M.C., Cagua, E.F., Balbuena, J.A., Stouffer, D.B., Poisot, T., 2017. paco:
implementing Procrustean Approach to Cophylogeny in R. Methods Ecol. Evol. 8,
932–940. https://doi.org/10.1111/2041-210X.12736

Jackson, A.P., 2004. A reconciliation analysis oh host switching in plant-fungal
symbioses. Evolution 58, 1909–1923. https://doi.org/10.1111/j.0014-
3820.2004.tb00479.x
Jacquemyn, H., Brys, R., Waud, M., Busschaert, P., Lievens, B., 2015. Mycorrhizal
networks and coexistence in species-rich orchid communities. New Phytol. 206,
1127–1134. https://doi.org/10.1111/nph.13281
Jin, Y., Qian, H., 2022. V.PhyloMaker2: An updated and enlarged R package that can
generate very large phylogenies for vascular plants. Plant Divers. 44, 335–339.
https://doi.org/10.1016/j.pld.2022.05.005
Johnston, P.R., Quijada, L., Smith, C.A., Baral, H.-O., Hosoya, T., Baschien, C., Pärtel, K.,
Zhuang, W.-Y., Haelewaters, D., Park, D., Carl, S., López-Giráldez, F., Wang, Z.,
Townsend, J.P., 2019. A multigene phylogeny toward a new phylogenetic
classification of Leotiomycetes. IMA Fungus 10. https://doi.org/10.1186/s43008-
019-0002-x
Karandashov, V., Nagy, R., Wegmüller, S., Amrhein, N., Bucher, M., 2004. Evolutionary
conservation of a phosphate transporter in the arbuscular mycorrhizal symbiosis.
Proc. Natl. Acad. Sci. 101, 6285–6290. https://doi.org/10.1073/pnas.0306074101
Katoh, K., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on
fast Fourier transform. Nucleic Acids Res. 30, 3059–3066.
https://doi.org/10.1093/nar/gkf436
Legendre, P., Desdevises, Y., Bazin, E., 2002. A statistical test for host–parasite
coevolution. Syst. Biol. 51, 217–234.
https://doi.org/10.1080/10635150252899734
Martos, F., Munoz, F., Pailler, T., Kottke, I., Gonneau, C., Selosse, M., 2012. The role of
epiphytism in architecture and evolutionary constraint within mycorrhizal
networks of tropical orchids. Mol. Ecol. 21, 5098–5109.
https://doi.org/10.1111/j.1365-294X.2012.05692.x
Merckx, V., Bidartondo, M.I., 2008. Breakdown and delayed cospeciation in the
arbuscular mycorrhizal mutualism. Proc. R. Soc. B Biol. Sci. 275, 1029–1035.
https://doi.org/10.1098/rspb.2007.1622
Nguyen, L.-T., Schmidt, H.A., Von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies.
Mol. Biol. Evol. 32, 268–274. https://doi.org/10.1093/molbev/msu300
Öpik, M., Metsis, M., Daniell, T.J., Zobel, M., Moora, M., 2009. Large-scale parallel 454
sequencing reveals host ecological group specificity of arbuscular mycorrhizal
fungi in a boreonemoral forest. New Phytol. 184, 424–437.
https://doi.org/10.1111/j.1469-8137.2009.02920.x
Öpik, M., Vanatoa, A., Vanatoa, E., Moora, M., Davison, J., Kalwij, J.M., Reier, Ü., Zobel,
M., 2010. The online database Maarj *AM* reveals global and ecosystemic
distribution patterns in arbuscular mycorrhizal fungi (Glomeromycota). New
Phytol. 188, 223–241. https://doi.org/10.1111/j.1469-8137.2010.03334.x
Otero, J.T., Thrall, P.H., Clements, M., Burdon, J.J., Miller, J.T., 2011. Codiversification of
orchids (Pterostylidinae) and their associated mycorrhizal fungi. Aust. J. Bot. 59,
480. https://doi.org/10.1071/BT11053
Page, R., 1994. Parallel phylogenies: reconstructing the history of host-parasite
assemblages. Cladistics 10, 155–173. https://doi.org/10.1006/clad.1994.1010

Paradis, E., Claude, J., Strimmer, K., 2004. APE: Analyses of Phylogenetics and Evolution
in R language. Bioinformatics 20, 289–290.
https://doi.org/10.1093/bioinformatics/btg412
Perez-Lamarque, B., Morlon, H., 2024. Distinguishing Cophylogenetic Signal from
Phylogenetic Congruence Clarifies the Interplay Between Evolutionary History
and Species Interactions. Syst. Biol. syae013.
https://doi.org/10.1093/sysbio/syae013
Perez-Lamarque, B., Öpik, M., Maliet, O., Afonso Silva, A.C., Selosse, M., Martos, F.,
Morlon, H., 2022. Analysing diversification dynamics using barcoding data: The
case of an obligate mycorrhizal symbiont. Mol. Ecol. 31, 3496–3512.
https://doi.org/10.1111/mec.16478
Perez-Lamarque, B., Petrolli, R., Strullu-Derrien, C., Strasberg, D., Morlon, H., Selosse,
M.-A., Martos, F., 2022. Structure and specialization of mycorrhizal networks in
phylogenetically diverse tropical communities. Environ. Microbiome 17, 38.
https://doi.org/10.1186/s40793-022-00434-0
Perez-Lamarque, B., Selosse, M., Öpik, M., Morlon, H., Martos, F., 2020. Cheating in
arbuscular mycorrhizal mutualism: a network and phylogenetic analysis of
mycoheterotrophy. New Phytol. 226, 1822–1835.
https://doi.org/10.1111/nph.16474
Petrolli, R., Zinger, L., Perez-Lamarque, B., Collobert, G., Griveau, C., Pailler, T., Selosse,
M., Martos, F., 2022. Spatial turnover of fungi and partner choice shape
mycorrhizal networks in epiphytic orchids. J. Ecol. 110, 2568–2584.
https://doi.org/10.1111/1365-2745.13986
Remy, W., Taylor, T.N., Hass, H., Kerp, H., 1994. Four hundred-million-year-old vesicular
arbuscular mycorrhizae. Proc. Natl. Acad. Sci. 91, 11841–11843.
https://doi.org/10.1073/pnas.91.25.11841
Rimington, W.R., Pressel, S., Duckett, J.G., Field, K.J., Bidartondo, M.I., 2019. Evolution
and networks in ancient and widespread symbioses between Mucoromycotina
and liverworts. Mycorrhiza 29, 551–565. https://doi.org/10.1007/s00572-019-
00918-x
Ronquist, F., 2002. Parsimony analysis of coevolving species associations.
Russo, L., Miller, A.D., Tooker, J., Bjornstad, O.N., Shea, K., 2018. Quantitative
evolutionary patterns in bipartite networks: Vicariance, phylogenetic tracking or
diffuse co-evolution? Methods Ecol. Evol. 9, 761–772.
https://doi.org/10.1111/2041-210X.12914
Santichaivekin, S., Yang, Q., Liu, J., Mawhorter, R., Jiang, J., Wesley, T., Wu, Y.-C.,
Libeskind-Hadas, R., 2021. eMPRess: a systematic cophylogeny reconciliation
tool. Bioinformatics 37, 2481–2482.
https://doi.org/10.1093/bioinformatics/btaa978
Satler, J.D., Herre, E.A., Jandér, K.C., Eaton, D.A.R., Machado, C.A., Heath, T.A., Nason,
J.D., 2019. Inferring processes of coevolutionary diversification in a community of
Panamanian strangler figs and associated pollinating wasps*. Evolution 73, 2295–
2311. https://doi.org/10.1111/evo.13809
Schoch, C.L., Seifert, K.A., Huhndorf, S., Robert, V., Spouge, J.L., Levesque, C.A., Chen,
W., Fungal Barcoding Consortium, Fungal Barcoding Consortium Author List,
Bolchacova, E., Voigt, K., Crous, P.W., Miller, A.N., Wingfield, M.J., Aime, M.C., An,
K.-D., Bai, F.-Y., Barreto, R.W., Begerow, D., Bergeron, M.-J., Blackwell, M.,

Boekhout, T., Bogale, M., Boonyuen, N., Burgaz, A.R., Buyck, B., Cai, L., Cai, Q., Cardinali, G., Chaverri, P., Coppins, B.J., Crespo, A., Cubas, P., Cummings, C., Damm, U., De Beer, Z.W., De Hoog, G.S., Del-Prado, R., Dentinger, B., Diéguez-Uribeondo, J., Divakar, P.K., Douglas, B., Dueñas, M., Duong, T.A., Eberhardt, U., Edwards, J.E., Elshahed, M.S., Fliegerova, K., Furtado, M., García, M.A., Ge, Z.-W., Griffith, G.W., Griffiths, K., Groenewald, J.Z., Groenewald, M., Grube, M., Gryzenhout, M., Guo, L.-D., Hagen, F., Hambleton, S., Hamelin, R.C., Hansen, K., Harrold, P., Heller, G., Herrera, C., Hirayama, K., Hirooka, Y., Ho, H.-M., Hoffmann, K., Hofstetter, V., Högnabba, F., Hollingsworth, P.M., Hong, S.-B., Hosaka, K., Houbraken, J., Hughes, K., Huhtinen, S., Hyde, K.D., James, T., Johnson, E.M., Johnson, J.E., Johnston, P.R., Jones, E.B.G., Kelly, L.J., Kirk, P.M., Knapp, D.G., Kõljalg, U., Kovács, G.M., Kurtzman, C.P., Landvik, S., Leavitt, S.D., Liggenstoffer, A.S., Liimatainen, K., Lombard, L., Luangsa-ard, J.J., Lumbsch, H.T., Maganti, H., Maharachchikumbura, S.S.N., Martin, M.P., May, T.W., McTaggart, A.R., Methven, A.S., Meyer, W., Moncalvo, J.-M., Mongkolsamrit, S., Nagy, L.G., Nilsson, R.H., Niskanen, T., Nyilasi, I., Okada, G., Okane, I., Olariaga, I., Otte, J., Papp, T., Park, D., Petkovits, T., Pino-Bodas, R., Quaedvlieg, W., Raja, H.A., Redecker, D., Rintoul, T.L., Ruibal, C., Sarmiento-Ramírez, J.M., Schmitt, I., Schüßler, A., Shearer, C., Sotome, K., Stefani, F.O.P., Stenroos, S., Stielow, B., Stockinger, H., Suetrong, S., Suh, S.-O., Sung, G.-H., Suzuki, M., Tanaka, K., Tedersoo, L., Telleria, M.T., Tretter, E., Untereiner, W.A., Urbina, H., Vágvölgyi, C., Vialle, A., Vu, T.D., Walther, G., Wang, Q.-M., Wang, Y., Weir, B.S., Weiß, M., White, M.M., Xu, J., Yahr, R., Yang, Z.L., Yurkov, A., Zamora, J.-C., Zhang, N., Zhuang, W.-Y., Schindel, D., 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for *Fungi*. Proc. Natl. Acad. Sci. 109, 6241–6246. https://doi.org/10.1073/pnas.1117018109

Sepp, S., Davison, J., Jairus, T., Vasar, M., Moora, M., Zobel, M., Öpik, M., 2019. Non-random association patterns in a plant–mycorrhizal fungal network reveal host–symbiont specificity. Mol. Ecol. 28, 365–378. https://doi.org/10.1111/mec.14924

Shefferson, R.P., Cowden, C.C., Mccormick, M.K., Yukawa, T., Ogura-Tsujita, Y., Hashimoto, T., 2010. Evolution of host breadth in broad interactions: mycorrhizal specificity in East Asian and North American rattlesnake plantains (Goodyera spp.) and their fungal hosts: MYCORRHIZAL HOST BREADTH EVOLUTION. Mol. Ecol. 19, 3008–3017. https://doi.org/10.1111/j.1365-294X.2010.04693.x

Smith, S.E., Read, D.J., 2008. Mycorrhizal symbiosis, 3rd ed. ed. Academic Press, Amsterdam Boston.

Strullu-Derrien, C., Selosse, M., Kenrick, P., Martin, F.M., 2018. The origin and evolution of mycorrhizal symbioses: from palaeomycology to phylogenomics. New Phytol. 220, 1012–1030. https://doi.org/10.1111/nph.15076

Suzuki, T.A., Fitzstevens, J.L., Schmidt, V.T., Enav, H., Huus, K.E., Mbong Ngwese, M., Grießhammer, A., Pfleiderer, A., Adegbite, B.R., Zinsou, J.F., Esen, M., Velavan, T.P., Adegnika, A.A., Song, L.H., Spector, T.D., Muehlbauer, A.L., Marchi, N., Kang, H., Maier, L., Blekhman, R., Ségurel, L., Ko, G., Youngblut, N.D., Kremsner, P., Ley, R.E., 2022. Codiversification of gut microbiota with humans. Science 377, 1328–1332. https://doi.org/10.1126/science.abm7759

Taengon, C., Feng, Y., Zhang, Y., Aluthwattha, S.T., Chen, J., Wang, G., 2025. Cospeciation is not the dominant driver of plant–pollinator codiversification in

specialized pollination systems. Integr. Zool. 20, 437–442.
https://doi.org/10.1111/1749-4877.12886
Talavera, G., Lukhtanov, V., Pierce, N.E., Vila, R., 2022. DNA Barcodes Combined with
Multilocus Data of Representative Taxa Can Generate Reliable Higher-Level
Phylogenies. Syst. Biol. 71, 382–395. https://doi.org/10.1093/sysbio/syab038
Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N.S., Wijesundera, R., Ruiz, L.V.,
Vasco-Palacios, A.M., Thu, P.Q., Suija, A., Smith, M.E., Sharp, C., Saluveer, E.,
Saitta, A., Rosas, M., Riit, T., Ratkowsky, D., Pritsch, K., Põldmaa, K., Piepenbring,
M., Phosri, C., Peterson, M., Parts, K., Pärtel, K., Otsing, E., Nouhra, E.,
Njouonkou, A.L., Nilsson, R.H., Morgado, L.N., Mayor, J., May, T.W., Majuakim, L.,
Lodge, D.J., Lee, S.S., Larsson, K.-H., Kohout, P., Hosaka, K., Hiiesalu, I., Henkel,
T.W., Harend, H., Guo, L., Greslebin, A., Grelet, G., Geml, J., Gates, G., Dunstan,
W., Dunk, C., Drenkhan, R., Dearnaley, J., De Kesel, A., Dang, T., Chen, X.,
Buegger, F., Brearley, F.Q., Bonito, G., Anslan, S., Abell, S., Abarenkov, K., 2014.
Global diversity and geography of soil fungi. Science 346, 1256688.
https://doi.org/10.1126/science.1256688
Tedersoo, L., Bahram, M., Zobel, M., 2020. How mycorrhizal associations drive plant
population and community biology. Science 367.
https://doi.org/10.1126/science.aba1223
Toju, H., Tanabe, A.S., Ishii, H.S., 2016. Ericaceous plant–fungus network in a harsh
alpine–subalpine environment. Mol. Ecol. 25, 3242–3257.
https://doi.org/10.1111/mec.13680
van der Heijden, M.G.A., Martin, F.M., Selosse, M., Sanders, I.R., 2015. Mycorrhizal
ecology and evolution: the past, the present, and the future. New Phytol. 205,
1406–1423. https://doi.org/10.1111/nph.13288
Van Galen, L.G., Orlovich, D.A., Lord, J.M., Nilsen, A.R., Dutoit, L., Larcombe, M.J., 2023.
Correlated evolution in an ectomycorrhizal host–symbiont system. New Phytol.
238, 1215–1229. https://doi.org/10.1111/nph.18802
Weiß, M., Waller, F., Zuccaro, A., Selosse, M., 2016. Sebacinales – one thousand and one
interactions with land plants. New Phytol. 211, 20–40.
https://doi.org/10.1111/nph.13977
Zhang, J., Gu, Z., Zhou, C., Zhou, H., 2025. Molecular phylogeny and morphology reveal
four new species in Hymenochaetales and one new species in Cantharellales
from Southwestern China. MycoKeys 115, 87–135.
https://doi.org/10.3897/mycokeys.115.142433
Zhao, Z., Li, X., Liu, M.F., Merckx, V.S.F.T., Saunders, R.M.K., Zhang, D., 2021. Specificity
of assemblage, not fungal partner species, explains mycorrhizal partnerships of
mycoheterotrophic *Burmannia* plants. ISME J. 15, 1614–1627.
https://doi.org/10.1038/s41396-020-00874-x
581
582

Supporting information

Supplementary Table 1: Summary table with characteristics of each mycorrhizal network and the corresponding results of the cophylogenetic analyses. The table reports, for each network: the spatial, evolutionary and taxonomic scales, the network size, the type of mycorrhizal associations, how we recovered phylogenetic trees of both plants and fungi, the results of the evaluation of cophylogenetic signal (with 2 methods: PACo and ParaFit), and the results of the evaluation of phylogenetic congruence (with the event-based method eMPRess with different cost values for each reconciliation event).

Supplementary Table 2: Data selection for the mycorrhizal networks from Arifin et al. (Arifin et al., 2023). Modification of Table S1 from Arifin et al. (2023). As different sequences of the same OTU were available, the representative sequence of each fungal OTU chosen to build the phylogenetic tree are highlighted in green.