Review: Estimates of livestock genetic diversity in the UK

C. M. Rochus [a], C. F. Price [b], I. Pocrnić [a]

[a] *Division of Quantitative Biology, Roslin Institute, Royal (Dick) School of Veterinary*

Studies, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG, UK

[b] *Rare Breeds Survival Trust, Stoneleigh Park, Kenilworth, CV8 2LG, UK*

Corresponding author: Christina Rochus. Email: crochus@ed.ac.uk

Abstract

The United Kingdom (UK) is home to many local and rare livestock breeds. The local

breed populations are highly adapted to specific environments in the UK and these

and other rare breeds provide solutions to niche needs. Rare breeds often contribute

more than expected to overall species genetic diversity, which is important because

genetic variation is needed for adapting to new challenges. It is therefore very

important from both UK and global perspectives to maintain genetic diversity of rare

livestock breeds in the UK, and to do this, we need to evaluate the monitoring of

genetic diversity to identify gaps in our knowledge and prioritise resources reserved

for conservation purposes. The objectives of this study were to survey the literature

to: (1) summarise genetic/genomic characterisation (effective population size (N_e)

and inbreeding) of domestic populations (livestock and equine) in the UK and Ireland;

(2) compare number of populations on the UK's Rare Breed Survival Trust (RBST)

watch list and the number of UK and Irish populations in the peer-reviewed literature

with inbreeding, genetic diversity and/or N_e estimates; and (3) compare annually

reported (census-based) estimates of N_e with (inbreeding and DNA-based) estimates

from peer-reviewed literature. We found a total of 37 publications with N_e or

inbreeding estimates for UK or Ireland livestock populations, published from 1975 to

2024. While many (42%) of the breeds on the RBST watchlist have been included in

publications, there are still many breeds, and a few species (turkey, duck and geese)

with no publicly available pedigree- or DNA-based genetic diversity measures. We

found census-based N_e estimates were, on average, higher than DNA-based

estimates, likely due to violated assumptions when estimating N_e with census-based

data because of the way livestock mating systems are designed. Most peer-reviewed

papers estimated genetic diversity measures using pedigree, microsatellite markers,

or SNP markers. To identify breed-unique variants responsible for adaptive traits in

rare breeds, more studies using whole-genome sequencing will be needed.

Altogether, we have summarised the genetic diversity estimates on UK livestock

populations, identifying gaps in knowledge.

Keywords: Genetic diversity, inbreeding, effective population size (N_e), rare breeds,

livestock, equine

Implications

Maintaining genetic diversity in livestock helps safeguard against future challenges to

our food production systems. We surveyed publicly available measures of genetic

diversity for farm animal populations in the UK (including government, Rare Breed

Survival Trust (RBST), and peer-reviewed scientific publications) to identify any gaps.

We found census-based estimates of genetic diversity, which are used to inform and

prioritise resources for conservation efforts, on average, are likely overestimating

available genetic diversity. Additionally, most farm animal breeds had only these

census-based estimates. Genetic diversity estimates could be improved using

pedigree or DNA information, more accurately identifying endangered breeds.

Introduction

Through domestication and the spread of animal-based agriculture, livestock

populations have become highly adapted to many niche environments (Zeder, 2008).

Those populations today are called local breeds and are often maintained by breed

associations for their unique qualities and nostalgia. These breeds can be described

as hardy, many of which can thrive on low-quality feed not appropriate for human

consumption and habitat in areas and weather conditions where other breeds cannot.

Rare breeds have lost their historic popularity because they have not been improved

through systematic breeding programmes, unlike contemporary commercial

populations that currently use genomic selection to inform mating decisions. Rare

breeds are therefore not as efficient, or economically viable as commercial

populations raised in intensive farming systems. However, there are several roles

that rare breeds fill in the modern world. In the UK and European Union (EU), niche

products, like cheese and meat, from local breeds are sold with a geographical

indicator (PDO – protected designation of origin, and PGI – protected geographical

indication) ('Geographical indications and quality schemes explained', n.d.;

'Protected geographical food and drink names: UK GI schemes', 2024). Local breeds

in the UK are used for conservation grazing to maintain numerous habitats

(grassland, heathland, wood pasture, coastal and floodplains, fen, scrub and scrub

mosaics, saltmarsh and sand dunes habitats) ('Graze with livestock to maintain and

improve habitats', n.d.) and are used in organic farming (Van Diepen et al., 2007).

Local breeds also play an intangible cultural role in their communities providing a

glimpse into the past, including domestication processes and the Neolithic transition

to agriculture, all the way up to modern times. Finally, in an ever-changing world,

these historic breeds can play an important role as reservoirs of genetic variation,

providing the variation needed for sustainable agriculture systems with animals that

can adapt to future challenges, which has been recognised by the UN ('Resolution

adopted by the General Assembly on 25 September 2015: 70/1. Transforming our

world: the 2030 Agenda for Sustainable Development', 2015; *Kumming-Montreal*

Global biodiversity framework: Conference of the parties to the convention on

biological diversity, 2022).

There are many other important roles for local UK breeds to play in modern society,

besides maintaining total livestock species genetic diversity. To highlight a few UK

examples, we present breeds with unique traits and their current and historic uses.

The North Ronaldsay sheep, found primarily in the Orkneys has adapted to primarily

eat seaweed, mainly composed of brown kelp, after they were mostly contained to

the shore in the 19th century to protect cultivated land from being grazed (RBST,

n.d.; Porter, 2002). Many modern breeds struggle with some functional traits as a

result of a focus on improving production traits (Rauw et al., 1998). In contrast, local

breeds are prized for these functional traits, and Shetland cattle are an example of

this with their excellent calving ease (Schwabe and Hall, 1989), they can be bred with

larger continental breeds without calving issues (RBST, n.d.). Local breeds can

provide unique animal products that cannot be provided by commercial breeds. One

of the first EU protected animal products was the Gloucestershire Old Spots pig

(RBST, n.d.). In the application for this status, it was indicated that texture, and

flavour of these pigs are distinguishable from commercial pigs (The Gloucestershire

Old Spots Pig Breeders' Club, 2006). There were some traditional functions of local

breeds that may not be in their current usage, but could inspire more sustainable

practices. Before the use of pesticides to control parasites Shetland geese were

traditionally grazed in pastures after sheep were grazed to manage parasites like

liver fluke (RBST, n.d.). Finally, many of the local breeds can thrive and forage in

any climate, like the Scots Grey, a traditional Scottish chicken breed traced back to

the 16th century (RBST, n.d.). This ability to thrive outdoors is particularly useful for

conservation grazing. Many iconic UK landscapes, like meadows and grasslands

that harbour a third of British flora, were created from traditional farming practices

(Peach, 2019). In the Great Trossachs Forest, grazing allows for a more diverse

environment for wild birds, and opportunities for less competitive grass species to

seed and germinate, among other benefits (The Great Trossachs Forest, 2025).

They use a combination of Luing cattle, Hebridean sheep and Scottish Blackface

sheep, each providing different grazing patterns to contribute to the park's diversity

(The Great Trossachs Forest, 2025).

Local breeds from the British Isles have played an outsized role in global commercial

agriculture today. UK breeds have had widespread popularity, for example, purebred

and crossbred Angus, Jersey, and Guernsey cattle can be found in many countries

around the world, many global commercial pig companies have developed lines

based on Large White and Yorkshire breeds, and the Leicester Longwool was widely

used in crossbreeding schemes to improve the carcass and fleece of sheep breeds

in Europe, Australia, Canada, New Zealand, and the US (Porter, 2002). The

Leicester Longwool was the product of a new type of breeding, "in-and-in", developed

by English farmer Robert Bakewell in the 18[th] century, where inbreeding was used to

fix traits in a population (Wykes, 2004). This is believed to be the first implementation

of systematic selective breeding, when in comparison, many famers to this point had

been randomly mating livestock (Wykes, 2004).

The UK government recognises the importance of maintaining the genetic diversity of

its local breeds and has several national and international obligations to conserve

rare breeds, including the 2011 England Biodiversity Strategy, the UN Food and

Agriculture Organisation's (FAO) "Global Plan of Action", the UN Convention on

Biological Diversity "Strategic Plan for Biodiversity 2011-2020", and for input to the

European Farm Animal Biodiversity Information System (EFABIS) and FAO Global

Information System (DAD-IS) (Beale, 2024). To fulfil these obligations, the UK's

Department for Environment, Food and Rural Affairs (DEFRA) monitors the number

of active sires and dams for livestock populations by breed to inform and prioritise

resources for conservation efforts. DEFRA collects this census data through

collaboration with breed organisations and the Rare Breed Survival Trust (RBST), a

charity that conserves and promotes native livestock and equine breeds in the UK

(RBST, n.d.).

Additionally, several research groups have studied the genetic diversity and

inbreeding of local and rare breeds, publishing their results in peer-reviewed

literature. These studies were standalone projects or part of global consortia (Bovine

HapMap (The Bovine HapMap Consortium et al., 2009), Sheep Hapmap , Adaptmap

(for goats) (Stella et al., 2018), etc.). The objectives of this study were to survey the

literature to: (1) summarise genetic/genomic characterisation (genetic diversity,

effective population size (N_e) and inbreeding) of domestic populations (livestock and

equine) in the UK and Ireland; (2) compare number of populations on the RBST list

and the number of UK and Irish populations in the peer-reviewed literature with

inbreeding, genetic diversity and/or N_e estimates; and (3) compare annually reported

(census-based) estimates of N_e with (DNA-based) estimates from peer-reviewed

literature. By systematically reviewing the peer-reviewed genetic diversity literature

for UK livestock populations, we can identify gaps, which will help with prioritising

research areas.

Material and methods

We used Web of Science (https://www.webofscience.com) to search (on 2025-01-13)

using the following search term:

(inbreeding OR diversity OR "effective population size") AND (cattle OR cow* OR

bull* OR pig* OR chicken* OR sheep* OR goat* OR horse* OR duck* OR turkey* OR

geese) AND ("UK" OR "United Kingdom" OR "Britain" OR "Scotland" OR "England"

OR "Wales" OR "Northern Ireland" OR "Ireland") AND ("genomic" or "genetic" or

"pedigree").

We searched for inbreeding, diversity and N_e estimates for all species identified on

the Rare Breed Survival Trust's (RBST) 2024-2025 watchlist for breeds in the UK

('RBST watchlist 2024', 2024). Irish populations were included due to the shared

border between Northern Ireland and Ireland, making some livestock and equine

populations to likely be well-connected. Values were extracted from articles, tables,

and figures (using WebPlotDigitizer (Rohatgi, 2024)). The RBST watchlist ('RBST

watchlist 2024', 2024) is annually published and used to prioritise breeds for

conservation by considering genetic diversity and number of breeding females.

Every year DEFRA collects census numbers (they can be number of animals

registered or an estimate of total number) from breed organisations and the RBST,

and publishes them (Beale, 2024). We have used the 2023 release of census and

effective population estimates (Beale, 2024). This data is used to monitor numbers of

animals over the years to help with prioritisation of conservation efforts. DEFRA uses

the following formula to estimate N_e:

184
$$N_e = \frac{4N_m N_f}{N_m + N_f}$$

Where N_m is the number of breeding males, and N_f is the number of breeding

females) (Wright, 1931). N_e is the number of individuals that would be needed for the

inbreeding estimate to be true in an idealised population (Falconer and Mackay,

1996). An idealised population has the following conditions: (1) mating is restricted to

a population with no migration, (2) generations are discrete, (3) number of breeding

individuals are the same for every generation, (4) mating is random and includes self-

fertilisation, (5) there is no selection, and (6) mutation is disregarded (Falconer and

Mackay, 1996). The effective population estimate is standardised and comparable

between populations, and can be used to estimate the rate of inbreeding (Wright,

1931). The UN FAO guidance indicates N_e should be 50 animals per generation, or

more, as an N_e of 50 corresponds to 1% inbreeding rate per generation (when

number of breeding males and females are equal) (FAO Commission on Genetic

Resources for Food and Agriculture, 2015).

Study design

We have summarised the peer-reviewed literature on genetic diversity in UK

livestock populations by counting the number of published papers per species,

population, data type (pedigree, mtDNA, microsatellite (SSR), single nucleotide

polymorphisms (SNP), or whole genome sequencing (WGS)) and number of

populations both in the literature and on the RBST 2024 watchlist.

Additionally, for all inbreeding- and DNA-based N_e estimates found for UK

populations in the literature, we compared this with census-based N_e estimates

published by DEFRA. To make the closest comparisons, we compared census-

based estimates from the same year of DNA sampling that was used in the DNA-

based estimates. If the year of sampling was not available, year of publication was

used as a proxy. If there was no census-based estimate for the year of sampling/year

of publication of DNA-based estimates, the closest year with an estimate was used. If

the sampling occurred over a period of years, census-based N_e estimates from those

years were averaged. We used a paired t-test to determine if census-based and

DNA-based N_e estimates were significantly different.

Results

Summary of literature search

In total, we have included 36 peer-reviewed publications and one report written for

the RBST, published between 1975 and 2024, all of which have estimates of genetic

diversity, inbreeding, diversity or N_e of UK or Irish livestock or equine populations.

The 37 publications are summarised in **Table 1**, including sources of data for each

study (original data from a total of 50 publications). **Figure 1** shows our workflow,

including total number of papers found through the Web of Science search, number

of papers removed due to not meeting certain criteria, and number of papers

included that were not identified in the Web of Science search.

Figure 1: Literature search terms and number of results from Web of Science

Summary of counts

Of the publications included in our analyses, there were 14 papers on equine

populations, 11 on cattle, ten on sheep, one on pig, one on goat and one on chicken.

Information on breeds studied, and data type are included in **Table 1**. The number of

populations in both the literature and the RBST 2024 watchlist are plotted by

conservation priority status in **Figure 2**. Notably, there were no goat, turkey, duck, or

geese populations on the RBST 2024 watchlist that have estimates of genetic

inbreeding, diversity or N_e in the literature.

Figure 2: Clustered stacked bar plot of the number of UK and Irish livestock

populations on the RSBT list with estimates of inbreeding, diversity or effective

population size (N_e) estimated using pedigree or molecular markers available ("yes")

or not ("no")

Comparing genomic-based diversity estimates with census data

Of the N_e estimates from genomic data with census estimates available, 43 (of 53)

census estimates were greater than inbreeding- or LD-based estimates. The paired t-

test for census-based and inbreeding or LD-based N_e was significant (p=0.0009).

The median difference between the two was 391 (maximum difference was 24,068).

All census-based N_e estimates with their corresponding inbreeding- or DNA-based N_e

estimates from DEFRA are plotted in **Figure 2**. DNA-based N_e were estimated from

linkage disequilibrium (LD) methods.

Figure 3: Inbreeding (F)- and linkage disequilibrium (LD)-based effective population

size (N_e) estimates. N_e of 50 are indicated on both axes with grey dashed lines.

Discussion

It is important to maintain rare breeds to adapt to future challenges, to serve niche

markets, for conservation grazing, and to maintain cultural identity and contribute to

sustainable agriculture. However, there are finite resources available for conservation

of these breeds, and they are not immediately profitable in the larger economy, so it

is important to identify gaps, and prioritise certain aspects of livestock conservation.

In this sense, we discuss: (1) work already done in these areas and compare census-

based N_e estimates with DNA-based estimates, (2) potential value of WGS in rare

breeds, (3) usage of N_e estimates to guide conservation strategies, (4) limitations of

our study design and (5) future actions.

N_e estimates

The well-established threshold of concern for population conservation is a N_e of 50

(FAO Commission on Genetic Resources for Food and Agriculture, 2015). Franklin

(1980) first proposed a short-term aim of a N_e of 50 for small populations (with

livestock populations particularly in mind) as this corresponds to a 1% rate of

inbreeding ($\Delta F = \frac{1}{2N_e}$) (Wright, 1931), which is considered a tolerable level of

inbreeding, considering the negative effect of inbreeding depression. This was part of

the 50/500 rule, where Franklin also proposed a long-term minimum value for N_e of

500 (where mutation offsets the loss of genetic variation through drift) (Franklin,

1980). There have been more recent discussions on updating the 50/500 rule, which

is needed as our knowledge on genome biology has evolved in the following 45

years, however there has been no clear consensus on changing the recommended

minimum N_e (Waples, 2025; Pérez-Pereira et al., 2022; Frankham et al., 2014;

Jamieson and Allendorf, 2012).

The N_e threshold of concern may need to be updated, but N_e estimates as a measure

of genetic diversity also needs to be revisited. We compared census-based and

inbreeding- or DNA-(LD-)based N_e estimates using annually reported census-based

estimates from DEFRA and inbreeding- or LD-based estimates from the literature.

Census-based N_e estimates were, on average, higher than inbreeding- or LD-based

estimates (with a median difference of 391), and ~11% (6 of 53) of paired estimates

had an inbreeding or LD-based N_e estimate < 50, with a corresponding census-based

N_e > 50. This could indicate that several populations without DNA-based N_e

estimates but with census-based estimates appear to have an N_e > 50 but may

actually need conservation prioritisation.

We have observed very different N_e estimates based on data and methodology.

Estimating and publishing census-based estimates has been valuable as a low-cost

method for estimating N_e and observing trends over time. However, inbreeding and

DNA-based methods are estimates based on the genome, the diversity of which we

are trying to conserve. The difference between estimates could be expected due to

the assumptions of the methods used. In the census-based methods, there is an

assumption of an idealised population, which does not reflect the reality of livestock

breeding. DNA-based methods for estimating N_e also have assumptions, for

example, LD-based methods assume LD is produced solely from the finite population

size, and non-random mating and population structure are absent (Waples, 2025).

Additionally, any method using microsatellite or SNP arrays will also contend with

ascertainment bias of those DNA markers, where those markers have been identified

usually using animals not from the population of interest – which is especially

problematic for monitoring rare breeds with unique variants that we are interested in

conserving.

The census-based N_e estimates in this study showed that more common breeds in

the UK had higher estimates. An extreme example of this is the Holstein-Friesian

breed. The UK census-based N_e estimate for 2023 was 13,154 (Beale, 2024).

Holstein-Friesian is a globally popular commercial breed, and populations in different

countries are well-connected to each other, particularly through similar genetics

(global use of artificial insemination with elite bulls' semen). Therefore, we would

expect N_e estimates across countries to be somewhat similar to the UK estimate. N_e

estimates from the Netherlands (one of the countries where this breed originated

from), Canada, France, Ireland, Italy, and USA range from 18 to 150 (Ablondi et al.,

2022; Makanjuola et al., 2020; Harmen P. Doekes et al., 2018; H.P. Doekes et al.,

2018; Rodríguez-Ramilo et al., 2015; Danchin-Burge et al., 2011; Mc Parland et al.,

2007a; Weigel, 2001). These estimates came from a variety of inbreeding- and DNA-

based methods (but not the census-based method). This highlights how estimating

and comparing N_e estimates can be challenging (even the difference between 18 and

150 has vastly different implications for conservation), and that the census-based UK

estimate is unrealistic. In livestock populations there is often higher selection intensity

in males (fewer active breeding males than active breeding females), which can be

accounted for in Wright's estimate (Wright, 1931). However, the definition of "active

breeding males" could contribute to unreasonable N_e estimates. For example, some

of the active breeding males (e.g., elite bulls) likely have thousands of offspring, while

many active males only have a very small number of offspring, and this variation in

family size is not accounted for in the census-based estimate.

Whole-genome sequencing (WGS)

While genotyping strategies are becoming more affordable, especially SNP arrays,

typically rare breeds are raised as a hobby, and it is not financially feasible to have

continual producer-investment in routine genotyping. For producers raising rare

breeds for conservation purposes, SNP arrays are not ideal because they likely won't

contain unique breed-specific variants (due to ascertainment bias), of which are the

main interest of conservation programmes. WGS would be better for conservation

studies, however, it is more expensive. Sequencing costs are decreasing, and there

are strategies to reduce those costs even further, including using low-pass

sequencing and imputation, but this requires an initial investment to establish high

coverage genotyping of a reference population to identify breed-unique variants.

Only one publication included in this study used 15X coverage Illumina 150 bp

paired-end WGS (Lindsay-McGee et al., 2023). Due to the study design which also

used available genotypes from populations genotyped with an Affymetrix array, only

400K SNPs remained for analyses. In contrast, to give an idea of the magnitude of

the number of variants we could expect from sequencing, from a publication in 2019

from the 1000 Bull Genome Project where a total of 429 animals from 14 breeds

were sequenced, they identified 29.1 million single nucleotide variants, and 1.7

million insertions-deletions (Hayes et al., 2014). We could expect that there is a lot of

unexplored data from the Lindsay-McGee et al. (2023) study, and that it would be

valuable for further investment in generating WGS data for other rare breeds in the

UK, allowing for information on breed-unique variants to be included in management

plans.

Conservation management

As we previously mentioned, monitoring and follow-up conservation strategies are

mainly based on the FAO guidance (N_e of 50 per generation, equivalent to 1%

inbreeding rate per generation (FAO Commission on Genetic Resources for Food

and Agriculture, 2015)). Current census-based estimates of N_e should be replaced by

inbreeding-based estimates of N_e. For the census-based estimates of N_e, DEFRA is

using the number of active sires and dams, which is known because pedigree

information is collected and available. An immediate improvement would be to use

pedigree information to estimate rates of inbreeding, which could then be used to

calculate annual inbreeding-based N_e estimates. In the long-term, developing a

national strategy to regularly collect DNA samples for breeds on the RBST watchlist,

would allow for DNA-based estimates of genetic diversity measures.

Having DNA-based contemporary N_e estimates is critical to prioritise

breeds/populations for conservation activities. Furthermore, having genomic

(especially WGS) data on these breeds would not only allow for better N_e estimates

but would also lead to a better understanding of the general population structure.

This would add value for cost-effective decision making, particularly linked to

choosing the most optimal conservation system (*in situ*, *ex situ in vivo*, *ex situ in*

vitro), optimising the number of individuals within the breed to keep in the chosen

conservation system, as well as opening the door for future activities (e.g., precise

introgression, genome editing, etc.). Conservation activities and systems based on

genomic data were systematically reviewed in Oldenbroek et al. (2017), the

algorithms for selection and optimisation of the number of individuals are available

and can be easily adapted for conservation purposes, e.g., Pocrnic et al., (2022); Yu

et al., (2014), and for a recent review of the genome editing concepts applied to

livestock populations, including conservation applications, see Clark, (2020). We

particularly mention genome editing technology as there is potential to introduce

breed-unique causal variants into other populations (Leroy et al., 2016). Alternatively,

variants could be introduced by genome editing to make local populations more

resistant to disease or improve their fertility (traits that may suffer from inbreeding

depression), only changing traits to improve the survival of the breed while

maintaining the breed standards/definitions. Finally, a more complex application of

genome editing that could be considered is surrogate sire technology. Surrogate

sires are made sterile by CRISPR-Cas9 editing of *NANOS2* in mammals (Park et al.,

2017) or *DDX4* in chicken (Taylor et al., 2017), and donor-derived stem cells are

transplanted into the gonads (Taylor et al., 2017). The surrogate sire could then

disseminate the donor's sperm through natural breeding. The potential to exploit

surrogate sire technology in commercial livestock breeding was discussed by

Gottardo et al. (2019), but similarly, we could use it in a conservation context to

disseminate local breed sperm. Finally, novel technologies like gene drives are

emerging and could find their application in conservation strategies to help spread

desired alleles in a population (e.g., Faber et al. (2021); Webber et al. (2015); Esvelt

et al. (2014)).

Limitations of this study

Many of the results included in this study either came from or benefitted from

research consortiums. These consortiums build capacity through the training of

individuals, building databases and sharing resources. However, from these

consortium publications there were several breeds of UK origin, that were not

included in this study because of uncertainty on where samples were sampled.

Additionally, UK origin breeds in other countries may no longer reflect the current UK

population anymore due to the use of crossbreeding. Therefore, we didn't include

populations unless we were certain that they were UK or Ireland samples and if there

was a proportion of samples included from another country, there was some

evidence that these populations were well-connected with the corresponding UK

population.

There are more peer-reviewed publications that analyse genotypes from UK livestock

populations than we have included in this publication. We tried to keep only those

publications with some genetic diversity measure, inbreeding or N_e estimate. This

means papers looking at population structure and signatures of selections, while

related, were not included. Sometimes the data from those studies were used in

other publications where the measures we were interested in were estimated, and we

tried to cite both the original source of data and the publication with the analyses of

interest.

From systematically reviewing literature for genetic diversity, inbreeding and N_e

estimates for UK and Ireland livestock and equine populations, one recommendation

for other researchers is clear: include detailed information on populations analysed,

including sampling year, number of herds sampled from and sampling criteria. This

additional information will be helpful for future studies (for publicly available data

eventually used in other studies) and can be used to inform conservation decisions.

Future actions

This study was designed to identify gaps in our knowledge of the genetic diversity of

livestock populations in the UK. Most breeds on the RBST watchlist do not have

DNA-based studies, and for those with estimates, many are outdated. Most notably

missing were any estimates on duck, geese, goat and turkey breeds. These

populations should be prioritised for generating genomic data, and they need to

remain on the RBST watchlist and DEFRA monitoring programmes, potentially even

reassessing conservation categories based on DNA-based metrics when available.

The creation of a centralised database for all UK livestock rare breeds, integrating

DEFRA, RBST and academic sources, would help with implementing a systematic

approach to monitor genetic diversity, and fund the generation of genomic data to

update diversity estimates. Publicly funded genetic estimates and resulting

publications should be open access and allow for data sharing. Through the

standardisation and sharing of data, joint conservation programmes for

transboundary breeds can be developed, and breed connectedness and genetic

diversity measures can be jointly assessed.

Ethics approval

Ethics approval was not needed for this study.

Data and model availability statement

All data used in this study are available in the supplemental information.

Declaration of generative AI and AI-assisted technologies in the writing
process

We declare that no generative AI or AI-assisted technologies were used in the writing
process.

Author ORCIDs

C. M. Rochus 0000-0002-7570-6847

I. Pocrnić 0000-0001-524-7428

Declaration of interest

460   We declare that we have no conflicts of interest.

Acknowledgements

462   C.M.R. and I.P. acknowledge support from the Biotechnology and Biological

463   Sciences Research Council (BBSRC) awarded to The Roslin Institute (The University

464   of Edinburgh) (BBS/E/RL/230001A and BBS/E/RL/230001C). For the purpose of

465   open access, the authors have applied a CC BY public copyright license to any

466   author accepted manuscript version arising from this submission.

Financial support statement

469   C.M.R. and I.P. acknowledge support from the Biotechnology and Biological

470   Sciences Research Council (BBSRC) awarded to The Roslin Institute (The University

471   of Edinburgh) (BBS/E/RL/230001A and BBS/E/RL/230001C). For the purpose of

472   open access, the authors have applied a CC BY public copyright license to any

473   author accepted manuscript version arising from this submission.

References

Ablondi, M., Sabbioni, A., Stocco, G., Cipolat-Gotet, C., Dadousis, C., Kaam, J.-T.V., Finocchiaro, R., Summer, A., 2022. Genetic Diversity in the Italian Holstein Dairy Cattle Based on Pedigree and SNP Data Prior and After Genomic Selection. Frontiers in Veterinary Science 8, 773985. doi:10.3389/fvets.2021.773985

Banos, G., 2023. Genomic analysis of the North Ronaldsay sheep. Rare Breeds Survival Trust (RBST).

Banos, G., Talenti, A., Chatziplis, D., Sánchez-Molano, E., 2022. Genomic analysis of the rare British Lop pig and identification of distinctive genomic markers. PLOS ONE 17, e0271053. doi:10.1371/journal.pone.0271053

Barbato, M., Orozco-Terwengel, P., Tapio, M., Bruford, M.W., 2015. SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data 6. doi:10.3389/fgene.2015.00109

Beale, S., 2024. Official Statistics UK farm animal genetic resources (FAnGR): breed inventory results. UK Department for Environment, Food & Rural Affairs. Retrieved on from https://www.gov.uk/government/statistics/uk-farm-animal-genetic-resources-fangr-breed-inventory-results

Beynon, S.E., Slavov, G.T., Farré, M., Sunduimijid, B., Waddams, K., Davies, B., Haresign, W., Kijas, J., MacLeod, I.M., Newbold, C.J., Davies, L., Larkin, D.M., 2015. Population structure and history of the Welsh sheep breeds determined by whole genome genotyping. BMC Genetics 16, 65. doi:10.1186/s12863-015-0216-x

Bodo, I., Alderson, L., Langlois, B. (Eds.), 2005. Conservation genetics of endangered horse breeds. EAAP.

Bradley, D.G., MacHugh, D.E., Cunningham, P., Loftus, R.T., 1996. Mitochondrial diversity and the origins of African and European cattle. Proceedings of the National Academy of Sciences 93, 5131–5135. doi:10.1073/pnas.93.10.5131

Brinkmann, L., Gerken, M., Riek, A., 2012. Adaptation strategies to seasonal changes in environmental conditions of a domesticated horse breed, the Shetland pony (*Equus ferus caballus*). Journal of Experimental Biology 215, 1061–1068. doi:10.1242/jeb.064832

Browett, S., McHugo, G., Richardson, I.W., Magee, D.A., Park, S.D.E., Fahey, A.G., Kearney, J.F., Correia, C.N., Randhawa, I.A.S., MacHugh, D.E., 2018. Genomic Characterisation of the Indigenous Irish Kerry Cattle Breed. Frontiers in Genetics 9, 51. doi:10.3389/fgene.2018.00051

Cao, Y.-H., Xu, S.-S., Shen, M., Chen, Z.-H., Gao, L., Lv, F.-H., Xie, X.-L., Wang, X.-H., Yang, H., Liu, C.-B., Zhou, P., Wan, P.-C., Zhang, Y.-S., Yang, J.-Q., Pi, W.-H., Hehua, Ee., Berry, D.P., Barbato, M., Esmailizadeh, A., Nosrati, M., Salehian-Dehkordi, H., Dehghani-Qanatqestani, M., Dotsev, A.V., Deniskova, T.E., Zinovieva, N.A., Brem, G., Štěpánek, O., Ciani, E., Weimann, C., Erhardt, G., Mwacharo, J.M., Ahbara, A., Han, J.-L., Hanotte, O., Miller, J.M., Sim, Z., Coltman, D., Kantanen, J., Bruford, M.W., Lenstra, J.A., Kijas, J., Li, M.-H., 2021. Historical Introgression from Wild Relatives Enhanced Climatic Adaptation and Resistance to Pneumonia in Sheep. Molecular Biology and Evolution 38, 838–855. doi:10.1093/molbev/msaa236

Clark, E.L., 2020. Breeding in an Era of Genome Editing. In Encyclopedia of Sustainability Science and Technology (ed. Meyers, R.A.). Springer New York, New York, NY, pp. 1–16. doi:10.1007/978-1-4939-2493-6_1122-1

Corbin, L.J., Blott, S.C., Swinburne, J.E., Vaudin, M., Bishop, S.C., Woolliams, J.A., 2010. Linkage disequilibrium and historical effective population size in the Thoroughbred horse. Animal Genetics 41, 8–15. doi:10.1111/j.1365-2052.2010.02092.x

Cunningham, E.P., Dooley, J.J., Splan, R.K., Bradley, D.G., 2001. Microsatellite diversity, pedigree relatedness and the contributions of founder lineages to thoroughbred horses. Animal Genetics 32, 360–364. doi:10.1046/j.1365-2052.2001.00785.x

Danchin-Burge, C., Hiemstra, S.J., Blackburn, H., 2011. Ex situ conservation of Holstein-Friesian cattle: Comparing the Dutch, French, and US germplasm collections. Journal of Dairy Science 94, 4100–4108. doi:10.3168/jds.2010-3957

Decker, J.E., Pires, J.C., Conant, G.C., McKay, S.D., Heaton, M.P., Chen, K., Cooper, A., Vilkki, J., Seabury, C.M., Caetano, A.R., Johnson, G.S., Brenneman, R.A., Hanotte, O., Eggert, L.S., Wiener, P., Kim, J.-J., Kim, K.S., Sonstegard, T.S., Van Tassell, C.P., Neibergs, H.L., McEwan, J.C., Brauning, R., Coutinho, L.L., Babar, M.E., Wilson, G.A., McClure, M.C., Rolf, M.M., Kim, J., Schnabel, R.D., Taylor, J.F., 2009. Resolving the evolution of extant and extinct ruminants with high-throughput phylogenomics. Proceedings of the National Academy of Sciences 106, 18644–18649. doi:10.1073/pnas.0904691106

Dell, A., Curry, M., Yarnell, K., Starbuck, G., Wilson, P.B., 2020. Genetic analysis of the endangered Cleveland Bay horse: A century of breeding characterised by pedigree and microsatellite data. PLOS ONE 15, e0240410. doi:10.1371/journal.pone.0240410

Doekes, H.P., Veerkamp, R.F., Bijma, P., Hiemstra, S.J., Windig, J., 2018. Value of the Dutch Holstein Friesian germplasm collection to increase genetic variability and improve genetic merit. Journal of Dairy Science 101, 10022–10033. doi:10.3168/jds.2018-15217

Doekes, Harmen P., Veerkamp, R.F., Bijma, P., Hiemstra, S.J., Windig, J.J., 2018. Trends in genome-wide and region-specific genetic diversity in the Dutch-Flemish Holstein–Friesian breeding program from 1986 to 2015. Genetics Selection Evolution 50, 15. doi:10.1186/s12711-018-0385-y

Esvelt, K.M., Smidler, A.L., Catteruccia, F., Church, G.M., 2014. Concerning RNA-guided gene drives for the alteration of wild populations. eLife 3, e03401. doi:10.7554/eLife.03401

Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. Molecular Ecology Resources 10, 564–567. doi:10.1111/j.1755-0998.2010.02847.x

Faber, N.R., McFarlane, G.R., Gaynor, R.C., Pocrnic, I., Whitelaw, C.B.A., Gorjanc, G., 2021. Novel combination of CRISPR-based gene drives eliminates resistance and localises spread. Scientific Reports 11, 3719. doi:10.1038/s41598-021-83239-4

Falconer, D.S., Mackay, T.F.C., 1996. Introduction to Quantitative Genetics. Pearson Education Limited, Harlow, England.

FAO Commission on Genetic Resources for Food and Agriculture, 2015. The second
report on the state of the world's animal genetic resorces for food and
agriculture.
Fox-Clipsham, L.Y., Brown, E.E., Carter, S.D., Swinburne, J.E., 2011. Population
screening of endangered horse breeds for the foal immunodeficiency
syndrome mutation. Veterinary Record 169, 655–655. doi:10.1136/vr.100235
Frankham, R., Bradshaw, C.J.A., Brook, B.W., 2014. Genetics in conservation
management: Revised recommendations for the 50/500 rules, Red List criteria
and population viability analyses. Biological Conservation 170, 56–63.
doi:10.1016/j.biocon.2013.12.036
Franklin, 1980. Evolutionary changes in small populations. In Conservation Biology.
An Evolutionary-Ecological Perspective. pp. 135–149.
Funk, S.M., Guedaoura, S., Juras, R., Raziq, A., Landolsi, F., Luís, C., Martínez,
A.M., Musa Mayaki, A., Mujica, F., Oom, M.D.M., Ouragh, L., Stranger, Y.,
Vega-Pla, J.L., Cothran, E.G., 2020. Major inconsistencies of inferred
population genetic structure estimated in a large set of domestic horse breeds
using microsatellites. Ecology and Evolution 10, 4261–4279.
doi:10.1002/ece3.6195
Geographical indications and quality schemes explained, n.d. . European
Commission. Retrieved on from
https://agriculture.ec.europa.eu/farming/geographical-indications-and-quality-
schemes/geographical-indications-and-quality-schemes-explained_en
Golden, B.L., Brinks, J.S., Bourdon, R.M., 1991. A performance programmed method
for computing inbreeding coefficients from large data sets for use in mixed-
model analyses. Journal of Animal Science 69, 3564–3573.
doi:10.2527/1991.6993564x
Gottardo, P., Gorjanc, G., Battagin, M., Gaynor, R.C., Jenko, J., Ros-Freixedes, R.,
Bruce A. Whitelaw, C., Mileham, A.J., Herring, W.O., Hickey, J.M., 2019. A
Strategy To Exploit Surrogate Sire Technology in Livestock Breeding
Programs. G3 Genes|Genomes|Genetics 9, 203–215.
doi:10.1534/g3.118.200890
Graze with livestock to maintain and improve habitats, n.d. . UK Department for
Environment, Food & Rural Affairs. Retrieved on from
https://defrafarming.blog.gov.uk/graze-with-livestock-to-maintain-and-improve-
habitats/
Grilz-Seger, G., Neuditschko, M., Ricard, A., Velie, B., Lindgren, G., Mesarič, M.,
Cotman, M., Horna, M., Dobretsberger, M., Brem, G., Druml, T., 2019.
Genome-Wide Homozygosity Patterns and Evidence for Selection in a Set of
European and Near Eastern Horse Breeds. Genes 10, 491.
doi:10.3390/genes10070491
Gu, J., Orr, N., Park, S.D., Katz, L.M., Sulimova, G., MacHugh, D.E., Hill, E.W., 2009.
A Genome Scan for Positive Selection in Thoroughbred Horses. PLoS ONE 4,
e5767. doi:10.1371/journal.pone.0005767
Gutiérrez, J.P., Goyache, F., 2005. A note on ENDOG: a computer program for
analysing pedigree information. Journal of Animal Breeding and Genetics 122,
172–176. doi:10.1111/j.1439-0388.2005.00512.x
Hayes, B.J., Macleod, I.M., Daetwyler, H.D., Bowman, P.J., Chamberlian, A.J., Jagt,
C.J.V., Capitan, A., Pausch, H., Stothard, P., Liao, X., Schrooten, C., Mullaart,
E., Fries, R., Guldbrandtsen, B., Lund, M.S., Boichard, D., Veerkamp, R.F.,
Vantassell, C.P., Gredler, B., Druet, T., Bagnato, A., Vilkki, J., Dekoning, D.J.,

Santus, E., Goddard, M.E., 2014. Genomic prediction from whole genome sequence in livestock: the 1000 Bull Genomes Project. In 10th World Congress of Genetics Applied to Livestock Production. Vancouver, Canada, p. 6.

Hayes, B.J., Visscher, P.M., McPartlan, H.C., Goddard, M.E., 2003. Novel Multilocus Measure of Linkage Disequilibrium to Estimate Past Effective Population Size. Genome Research 13, 635–643. doi:10.1101/gr.387103

Henderson, C.R., 1976. A Simple Method for Computing the Inverse of a Numerator Relationship Matrix Used in Prediction of Breeding Values. Biometrics 32, 69. doi:10.2307/2529339

Hill, E.W., Stoffel, M.A., McGivney, B.A., MacHugh, D.E., Pemberton, J.M., 2022. Inbreeding depression and the probability of racing in the Thoroughbred horse. Proceedings of the Royal Society B: Biological Sciences 289, 20220487. doi:10.1098/rspb.2022.0487

Hill, W.G., 1972. Effective size of populations with overlapping generations. Theoretical Population Biology 3, 278–289. doi:10.1016/0040-5809(72)90004-4

Jamieson, I.G., Allendorf, F.W., 2012. How does the 50/500 rule apply to MVPs? Trends in Ecology & Evolution 27, 578–584. doi:10.1016/j.tree.2012.07.001

Jorde, P.E., Ryman, N., 1995. Temporal allele frequency change and estimation of effective size in populations with overlapping generations. Genetics 139, 1077–1090. doi:10.1093/genetics/139.2.1077

Juras, R., Cothran, E.G., Klimas, R., 2003. Genetic Analysis of Three Lithuanian Native Horse Breeds. Acta Agriculturae Scandinavica, Section A - Animal Science 53, 180–185. doi:10.1080/09064700310012971

Kerr, E., Marr, M.M., Collins, L., Dubarry, K., Salavati, M., Scinto, A., Woolley, S., Clark, E.L., 2024. Analysis of genotyping data reveals the unique genetic diversity represented by the breeds of sheep native to the United Kingdom. BMC Genomic Data 25, 82. doi:10.1186/s12863-024-01265-3

Khanshour, A., Juras, R., Blackburn, R., Cothran, E.G., 2015. The Legend of the Canadian Horse: Genetic Diversity and Breed Origin. Journal of Heredity 106, 37–44. doi:10.1093/jhered/esu074

Khanshour, A.M., Juras, R., Cothran, E.G., 2013. Microsatellite analysis of genetic variability in Waler horses from Australia. Australian Journal of Zoology 61, 357. doi:10.1071/ZO13062

Kijas, J.W., Lenstra, J.A., Hayes, B., Boitard, S., Porto Neto, L.R., San Cristobal, M., Servin, B., McCulloch, R., Whan, V., Gietzen, K., Paiva, S., Barendse, W., Ciani, E., Raadsma, H., McEwan, J., Dalrymple, B., other members of the International Sheep Genomics Consortium, 2012. Genome-Wide Analysis of the World's Sheep Breeds Reveals High Levels of Historic Mixture and Strong Recent Selection. PLoS Biology 10, e1001258. doi:10.1371/journal.pbio.1001258

Kijas, J.W., Porto-Neto, L., Dominik, S., Reverter, A., Bunch, R., McCulloch, R., Hayes, B.J., Brauning, R., McEwan, J., the International Sheep Genomics Consortium, 2014. Linkage disequilibrium over short physical distances measured in sheep using a high-density SNP chip. Animal Genetics 45, 754–757. doi:10.1111/age.12197

Kumming-Montreal Global biodiversity framework: Conference of the parties to the convention on biological diversity (No. CBD/COP/15/L.25), 2022. . Convention on Biological Diversity, Montreal, Canada.

Leroy, G., Besbes, B., Boettcher, P., Hoffmann, I., Capitan, A., Baumung, R., 2016. Rare phenotypes in domestic animals: unique resources for multiple applications. Animal Genetics 47, 141–153. doi:10.1111/age.12393

Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. Bioinformatics 25, 1451–1452. doi:10.1093/bioinformatics/btp187

Lindsay-McGee, V., Sanchez-Molano, E., Banos, G., Clark, E.L., Piercy, R.J., Psifidi, A., 2023. Genetic characterisation of the Connemara pony and the Warmblood horse using a within-breed clustering approach. Genetics Selection Evolution 55, 60. doi:10.1186/s12711-023-00827-w

MacHugh, D.E., Loftus, R.T., Cunningham, P., Bradley, D.G., 1998. Genetic structure of seven European cattle breeds assessed using 20 microsatellite markers. Animal Genetics 29, 333–340. doi:10.1046/j.1365-2052.1998.295330.x

MacHugh, D.E., Shriver, M.D., Loftus, R.T., Cunningham, P., 1997. Microsatellite DNA Variation and the Evolution, Domestication and Phylogeography of Taurine and Zebu Cattle (Bos taurus and Bos indicus). Genetics 146, 1071–1086. doi:https://doi.org/10.1093/genetics/146.3.1071

MacHugh, D.E., Troy, C.S., McCormick, F., Olsaker, I., Eythórsdóttir, E., Bradle, D.G., 1999. Early medieval cattle remains from a Scandinavian settlement in Dublin: genetic analysis and comparison with extant breeds. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences 354, 99–109. doi:10.1098/rstb.1999.0363

Mahon, G.A.T., Cunningham, E.P., 1982. Inbreeding and the inheritance of fertility in the thoroughbred mare. Livestock Production Science 9, 743–754. doi:10.1016/0301-6226(82)90021-5

Makanjuola, B.O., Miglior, F., Abdalla, E.A., Maltecca, C., Schenkel, F.S., Baes, C.F., 2020. Effect of genomic selection on rate of inbreeding and coancestry and effective population size of Holstein and Jersey cattle populations. Journal of Dairy Science 103, 5183–5199. doi:10.3168/jds.2019-18013

Manunza, A., Ramirez-Diaz, J., Cozzi, P., Lazzari, B., Tosser-Klopp, G., Servin, B., Johansson, A.M., Grøva, L., Berg, P., Våge, D.I., Stella, A., 2023. Genetic diversity and historical demography of underutilised goat breeds in North-Western Europe. Scientific Reports 13, 20728. doi:10.1038/s41598-023-48005-8

Martin, I., 1975. A Genetic Analysis of the Galway Sheep Breed: 3. Level of Inbreeding in the Pedigree Galway Sheep Breed.

Mc Parland, S., Kearney, J.F., Rath, M., Berry, D.P., 2007a. Inbreeding Effects on Milk Production, Calving Performance, Fertility, and Conformation in Irish Holstein-Friesians. Journal of Dairy Science 90, 4411–4419. doi:10.3168/jds.2007-0227

Mc Parland, S., Kearney, J.F., Rath, M., Berry, D.P., 2007b. Inbreeding trends and pedigree analysis of Irish dairy and beef cattle populations. Journal of Animal Science 85, 322–331. doi:10.2527/jas.2006-367

McGahern, A.M., Edwards, C.J., Bower, M.A., Heffernan, A., Park, S.D.E., Brophy, P.O., Bradley, D.G., MacHugh, D.E., Hill, E.W., 2006. Mitochondrial DNA sequence diversity in extant Irish horse populations and in ancient horses. Animal Genetics 37, 498–502. doi:10.1111/j.1365-2052.2006.01506.x

McHugo, G., Browett, S., Randhawa, I., Howard, D., Mullen, M., Richardson, I., Park, S., Magee, D., Scraggs, E., Dover, M., Correia, C., Hanrahan, J., MacHugh,

D., 2019. A Population Genomics Analysis of the Native Irish Galway Sheep
Breed. FRONTIERS IN GENETICS 10. doi:10.3389/fgene.2019.00927

Meuwissen, T.H.E., Luo, Z., 1992. Computing inbreeding coefficients in large
populations. Genetics, selection, evolution 24, 305–313. doi:10.1186/1297-
9686-24-4-305

Nei, M., 1987. Molecular evolutionary genetics, 2. print. ed. Columbia Univ. Pr, New
York, NY.

Oldenbroek, J.K., 2017. Management of Animal Genetic Diversity. Wageningen
Academic Publishers, Wageningen, the Netherlands.

Orozco-terWengel, P., Barbato, M., Nicolazzi, E., Biscarini, F., Milanesi, M., Davies,
W., Williams, D., Stella, A., Ajmone-Marsan, P., Bruford, M.W., 2015.
Revisiting demographic processes in cattle with genome-wide population
genetic analysis. Frontiers in Genetics 6. doi:10.3389/fgene.2015.00191

Park, K.-E., Kaucher, A.V., Powell, A., Waqas, M.S., Sandmaier, S.E.S., Oatley, M.J.,
Park, C.-H., Tibary, A., Donovan, D.M., Blomberg, L.A., Lillico, S.G., Whitelaw,
C.B.A., Mileham, A., Telugu, B.P., Oatley, J.M., 2017. Generation of germline
ablated male pigs by CRISPR/Cas9 editing of the NANOS2 gene. Scientific
Reports 7, 40176. doi:10.1038/srep40176

Peach, J., 2019. The great British meadow. Royal Botanic Garden Kew. Retrieved on
from https://www.kew.org/read-and-watch/great-british-meadow

Pérez-Pereira, N., Wang, J., Quesada, H., Caballero, A., 2022. Prediction of the
minimum effective size of a population viable in the long term. Biodiversity and
Conservation 31, 2763–2780. doi:10.1007/s10531-022-02456-z

Peter, C., Bruford, M., Perez, T., Dalamitra, S., Hewitt, G., Erhardt, G., 2007. Genetic
diversity and subdivision of 57 European and Middle-Eastern sheep breeds.
Animal Genetics 38, 37–44. doi:10.1111/j.1365-2052.2007.01561.x

Petersen, J.L., Mickelson, J.R., Rendahl, A.K., Valberg, S.J., Andersson, L.S.,
Axelsson, J., Bailey, E., Bannasch, D., Binns, M.M., Borges, A.S., Brama, P.,
da Câmara Machado, A., Capomaccio, S., Cappelli, K., Cothran, E.G., Distl,
O., Fox-Clipsham, L., Graves, K.T., Guérin, G., Haase, B., Hasegawa, T.,
Hemmann, K., Hill, E.W., Leeb, T., Lindgren, G., Lohi, H., Lopes, M.S.,
McGivney, B.A., Mikko, S., Orr, N., Penedo, M.C.T., Piercy, R.J., Raekallio,
M., Rieder, S., Røed, K.H., Swinburne, J., Tozaki, T., Vaudin, M., Wade, C.M.,
McCue, M.E., 2013. Genome-Wide Analysis Reveals Selection for Important
Traits in Domestic Horse Breeds. PLoS Genetics 9.
doi:10.1371/journal.pgen.1003211

Pocrnic, I., Lindgren, F., Tolhurst, D., Herring, W.O., Gorjanc, G., 2022. Optimisation
of the core subset for the APY approximation of genomic relationships.
Genetics Selection Evolution 54, 76. doi:10.1186/s12711-022-00767-x

Porter, V. (Ed.), 2002. Mason's World Dictionary of Livestock Breeds, Types and
Varieties. CABI Publishing, Wallingford, Oxon, UK.

Protected geographical food and drink names: UK GI schemes, 2024. . UK
Government. Retrieved on from https://www.gov.uk/guidance/protected-
geographical-food-and-drink-names-uk-gi-schemes

Prystupa, J.M., Hind, P., Cothran, E.G., Plante, Y., 2012. Maternal Lineages in
Native Canadian Equine Populations and Their Relationship to the Nordic and
Mountain and Moorland Pony Breeds. Journal of Heredity 103, 380–390.
doi:10.1093/jhered/ess003

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D.,
Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J., Sham, P.C., 2007. PLINK:

A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. The American Journal of Human Genetics 81, 559–575. doi:10.1086/519795

Purfield, D.C., McParland, S., Wall, E., Berry, D.P., 2017. The distribution of runs of homozygosity and selection signatures in six commercial meat sheep breeds. PLOS ONE 12, e0176780. doi:10.1371/journal.pone.0176780

Rafter, P., McHugh, N., Pabiou, T., Berry, D.P., 2022. Inbreeding trends and genetic diversity in purebred sheep populations. animal 16, 100604. doi:10.1016/j.animal.2022.100604

Rauw, W.M., Kanis, E., Noordhuizen-Stassen, E.N., Grommers, F.J., 1998. Undesirable side effects of selection for high production efficiency in farm animals: a review. Livestock Production Science 56, 15–33. doi:10.1016/S0301-6226(98)00147-X

RBST, n.d. North Ronaldsay. Rare Breed Survival Trust. Retrieved on from https://www.rbst.org.uk/north-ronaldsay

RBST, n.d. Shetland. Rare Breed Survival Trust. Retrieved on from https://www.rbst.org.uk/shetland

RBST, n.d. Gloucestershire Old Spots. Rare Breed Survival Trust. Retrieved on from https://www.rbst.org.uk/gloucestershire-old-spots

RBST, n.d. Shetland. Rare Breed Survival Trust. Retrieved on from https://www.rbst.org.uk/shetland-geese

RBST, n.d. Scots Grey. Rare Breed Survival Trust. Retrieved on from https://www.rbst.org.uk/scots-grey

RBST, n.d. Introduction. Rare Breed Survival Trust. Retrieved on from https://www.rbst.org.uk/our-mission

RBST watchlist 2024, 2024. . Rare Breeds Survival Trust (RBST). Retrieved on from https://www.rbst.org.uk/news/rbst-watchlist-2024

Resolution adopted by the General Assembly on 25 September 2015: 70/1. Transforming our world: the 2030 Agenda for Sustainable Development, 2015.

Rodríguez-Ramilo, S.T., Fernández, J., Toro, M.A., Hernández, D., Villanueva, B., 2015. Genome-Wide Estimates of Coancestry, Inbreeding and Effective Population Size in the Spanish Holstein Population. PLOS ONE 10, e0124157. doi:10.1371/journal.pone.0124157

Rohatgi, A., 2024. WebPlotDigitizer.

Roughsedge, T., Brotherstone, S., Visscher, P.M., 1999. Quantifying genetic contributions to a dairy cattle population using pedigree analysis. Livestock Production Science 60, 359–369. doi:10.1016/S0301-6226(99)00106-2

Schwabe, A.E., Hall, S.J.G., 1989. Dystocia in nine British breeds of cattle and its relationship to dimensions of the dam and calf. Veterinary Record 125, 636–639. doi:10.1136/vr.125.26-27.636

Stella, A., Nicolazzi, E.L., Van Tassell, C.P., Rothschild, M.F., Colli, L., Rosen, B.D., Sonstegard, T.S., Crepaldi, P., Tosser-Klopp, G., Joost, S., 2018. AdaptMap: exploring goat diversity and adaptation. Genetics Selection Evolution 50, 61, s12711-018-0427–5. doi:10.1186/s12711-018-0427-5

Tajima, F., 1983. EVOLUTIONARY RELATIONSHIP OF DNA SEQUENCES IN FINITE POPULATIONS. Genetics 105, 437–460. doi:10.1093/genetics/105.2.437

Taylor, L., Carlson, D.F., Nandi, S., Sherman, A., Fahrenkrug, S.C., McGrew, M.J., 2017. Efficient TALEN-mediated gene targeting of chicken primordial germ cells. Development dev.145367. doi:10.1242/dev.145367

Tenesa, A., Navarro, P., Hayes, B., Duffy, D., Clarke, G., Goddard, M., Visscher, P.,
2007. Recent human effective population size estimated from linkage
disequilibrium. GENOME RESEARCH 17, 520–526. doi:10.1101/gr.6023607
The Bovine HapMap Consortium, Gibbs, R.A., Taylor, J.F., Van Tassell, C.P.,
Barendse, W., Eversole, K.A., Gill, C.A., Green, R.D., Hamernik, D.L., Kappes,
S.M., Lien, S., Matukumalli, L.K., McEwan, J.C., Nazareth, L.V., Schnabel,
R.D., Weinstock, G.M., Wheeler, D.A., Ajmone-Marsan, P., Boettcher, P.J.,
Caetano, A.R., Garcia, J.F., Hanotte, O., Mariani, P., Skow, L.C., Sonstegard,
T.S., Williams, J.L., Diallo, B., Hailemariam, L., Martinez, M.L., Morris, C.A.,
Silva, L.O.C., Spelman, R.J., Mulatu, W., Zhao, K., Abbey, C.A., Agaba, M.,
Araujo, F.R., Bunch, R.J., Burton, J., Gorni, C., Olivier, H., Harrison, B.E., Luff,
B., Machado, M.A., Mwakaya, J., Plastow, G., Sim, W., Smith, T., Thomas,
M.B., Valentini, A., Williams, P., Womack, J., Woolliams, J.A., Liu, Y., Qin, X.,
Worley, K.C., Gao, C., Jiang, H., Moore, S.S., Ren, Y., Song, X.-Z.,
Bustamante, C.D., Hernandez, R.D., Muzny, D.M., Patil, S., San Lucas, A.,
Fu, Q., Kent, M.P., Vega, R., Matukumalli, A., McWilliam, S., Sclep, G., Bryc,
K., Choi, J., Gao, H., Grefenstette, J.J., Murdoch, B., Stella, A., Villa-Angulo,
R., Wright, M., Aerts, J., Jann, O., Negrini, R., Goddard, M.E., Hayes, B.J.,
Bradley, D.G., Barbosa Da Silva, M., Lau, L.P.L., Liu, G.E., Lynn, D.J.,
Panzitta, F., Dodds, K.G., 2009. Genome-Wide Survey of SNP Variation
Uncovers the Genetic Structure of Cattle Breeds. Science 324, 528–532.
doi:10.1126/science.1167936
The Gloucestershire Old Spots Pig Breeders' Club, 2006. Application for registration
of a TSG council regulation (EC) No 509/2006 'Traditionally farmed
Gloucestershire Old Spots Pork' EC No UK/007-0024 TSG.
The Great Trossachs Forest, 2025. Conservation grazing. The Great Trossachs
Forest. Retrieved on from
https://www.thegreattrossachsforest.co.uk/conservation-grazing
Van Diepen, P., McLean, B., Frost, D., 2007. Livestock breeds and organic farming
systems.
Wall, E., Brotherstone, S., Kearney, J.F., Woolliams, J.A., Coffey, M.P., 2005. Impact
of Nonadditive Genetic Effects in the Estimation of Breeding Values for
Fertility and Correlated Traits. Journal of Dairy Science 88, 376–385.
doi:10.3168/jds.S0022-0302(05)72697-7
Waples, R.S., 2025. The Idiot's Guide to Effective Population Size. Molecular
Ecology e17670. doi:10.1111/mec.17670
Webber, B.L., Raghu, S., Edwards, O.R., 2015. Is CRISPR-based gene drive a
biocontrol silver bullet or global conservation threat? Proceedings of the
National Academy of Sciences 112, 10565–10567.
doi:10.1073/pnas.1514258112
Weigel, K.A., 2001. Controlling Inbreeding in Modern Breeding Programs. Journal of
Dairy Science 84, E177–E184. doi:10.3168/jds.S0022-0302(01)70213-5
Weir, B.S., Cockerham, C.C., 1984. Estimating F-Statistics for the Analysis of
Population Structure. Evolution 38, 1358–1370.
Weir, B.S., Hill, W.G., 1980. EFFECT OF MATING STRUCTURE ON VARIATION IN
LINKAGE DISEQUILIBRIUM. Genetics 95, 477–488.
doi:10.1093/genetics/95.2.477
Wiener, P., Burton, D., Williams, J.L., 2004. Breed relationships and definition in
British cattle: a genetic analysis. Heredity 93, 597–602.
doi:10.1038/sj.hdy.6800566

Wilkinson, S., Wiener, P., Teverson, D., Haley, C.S., Hocking, P.M., 2012.
Characterization of the genetic diversity, structure and admixture of British
chicken breeds. Animal Genetics 43, 552–563. doi:10.1111/j.1365-
2052.2011.02296.x
Williams, J.L., Hall, S.J.G., Del Corvo, M., Ballingall, K.T., Colli, L., Ajmone Marsan,
P., Biscarini, F., 2015. Inbreeding and purging at the genomic Level: the
Chillingham cattle reveal extensive, non-random SNP heterozygosity. Animal
Genetics 47, 19–27. doi:10.1111/age.12376
Winton, C.L., Hegarty, M.J., McMahon, R., Slavov, G.T., McEwan, N.R., Davies-
Morel, M.C.G., Morgan, C.M., Powell, W., Nash, D.M., 2013. Genetic diversity
and phylogenetic analysis of native mountain ponies of Britain and Ireland
reveals a novel rare population. Ecology and Evolution 3, 934–947.
doi:10.1002/ece3.507
Winton, C.L., McMahon, R., Hegarty, M.J., McEwan, N.R., Davies-Morel, M.C.G.,
Morgan, C., Nash, D.M., 2020. Genetic diversity within and between British
and Irish breeds: The maternal and paternal history of native ponies. Ecology
and Evolution 10, 1352–1367. doi:10.1002/ece3.5989
Wright, S., 1950. THE GENETICAL STRUCTURE OF POPULATIONS. Annals of
Eugenics 15, 323–354. doi:10.1111/j.1469-1809.1949.tb02451.x
Wright, S., 1931. Evolution in Mendelian populations. Genetics 16, 97–159.
doi:10.1093/genetics/16.2.97
Wright, S., McPhee, H.C., 1925. An approximate method of calculating coefficients of
inbreeding and relationship from livestock pedigrees. Journal of Agricultural
Research 31, 377–383.
Wykes, D.L., 2004. Robert Bakewell (1725–1795) of Dishley: farmer and livestock
improver. The Agricultural History Review 52.
Yu, X., Woolliams, J.A., Meuwissen, T.H., 2014. Prioritizing animals for dense
genotyping in order to impute missing genotypes of sparsely genotyped
animals. Genetics Selection Evolution 46, 46. doi:10.1186/1297-9686-46-46
Zeder, M. a, 2008. Domestication and early agriculture in the Mediterranean Basin:
Origins, diffusion, and impact. Proceedings of the National Academy of
Sciences of the United States of America 105, 11597–11604.
doi:10.1073/pnas.0801317105
901

 Table 1: Publications characterising genetic diversity of UK and/or Irish livestock and

 equine populations

Species	Populations	Data	Methods	Publication	Data from
Cattle	Aberdeen Angus	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	3
		30 SSR	H_O, H_E	4	
	Angus	Pedigree	F_{PED}	5	
	Ancient Irish samples	mtDNA	ND	2	
	Ayrshire	30 SSR	H_O, H_E	4	
	Belted Galloway	50K SNP	N_{e-LD}, F_{ROH}, H_O	6	7
	Charolais	Pedigree	F_{PED}	5	
	Chillingham	777K SNP	F_{IS}, F_{ROH}	8	
		777K SNP	F_{IS}	9	
	Devon	50K SNP	H_O	6	7
	Dexter	30 SSR	H_O, H_E	4	
		50K SNP	N_{e-LD}, F_{ROH}, H_O	6	7
	English Longhorn	50K SNP	H_O	6	7
	Galloway	50K SNP	H_O	6	7
	Guernsey	30 SSR	H_O, H_E	4	
		50K SNP	F_{ROH}, H_O	6	7 & 10
	Hereford	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	3
		30 SSR	H_O, H_E	4	
		Pedigree	N_{e-F}, F_{PED}	5	
	Highland	30 SSR	H_O, H_E	4	
		50K SNP	H_O	6	7
	Holstein-Friesian	Pedigree	F_{PED}	11	
		Pedigree	F_{PED}	12	
		Pedigree	N_{e-F}, F_{PED}	7 & 13	
	Friesian	30 SSR	H_O, H_E	4	
	Jersey	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	3
		30 SSR	H_O, H_E	4	
	Kerry	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	
		50K SNP	N_{e-LD}, F_{ROH}, H_O	6	This study, 1 & 14
	Limousin	Pedigree	F_{PED}	5	
	Simmental	Pedigree	N_{e-F}, F_{PED}	5	
	South Devon	50K SNP	H_O	6	7
	Sussex	50K SNP	H_O	6	7
	Welsh Black	50K SNP	H_O	6	7
	White Park	777K SNP	F_{IS}	9	
		50K SNP	H_O	6	7
Chicken	Appenzeller	30 SSR	F_{IS}, H_O, H_E	15	
	Araucana	30 SSR	F_{IS}, H_O, H_E	15	
	Brahma	30 SSR	F_{IS}, H_O, H_E	15	
	Buff Orpington	30 SSR	F_{IS}, H_O, H_E	15	
	Cochin	30 SSR	F_{IS}, H_O, H_E	15	
	Croad Langshan	30 SSR	F_{IS}, H_O, H_E	15	
	Derbyshire Redcap	30 SSR	F_{IS}, H_O, H_E	15	
	Dorking	30 SSR	F_{IS}, H_O, H_E	15	

	Hamburgh	30 SSR	F_{IS}, H_O, H_E	15	
	Indian Game	30 SSR	F_{IS}, H_O, H_E	15	
	Ixworth	30 SSR	F_{IS}, H_O, H_E	15	
	Leghorn (coloured)	30 SSR	F_{IS}, H_O, H_E	15	
	Light Sussex	30 SSR	F_{IS}, H_O, H_E	15	
	Lincolnshire Buff	30 SSR	F_{IS}, H_O, H_E	15	
	Maran	30 SSR	F_{IS}, H_O, H_E	15	
	Marsh Daisy	30 SSR	F_{IS}, H_O, H_E	15	
	Norfolk Grey	30 SSR	F_{IS}, H_O, H_E	15	
	Old English Pheasant Fowl	30 SSR	F_{IS}, H_O, H_E	15	
	Rhode Island Red	30 SSR	F_{IS}, H_O, H_E	15	
	Scots Dumpy	30 SSR	F_{IS}, H_O, H_E	15	
	Scots Grey	30 SSR	F_{IS}, H_O, H_E	15	
	Silkie	30 SSR	F_{IS}, H_O, H_E	15	
	Spanish	30 SSR	F_{IS}, H_O, H_E	15	
	Sussex	30 SSR	F_{IS}, H_O, H_E	15	
Equine	Carneddau pony	50K SNP	H_O, H_E	16	
		17 SSR	H_O, H_E	16	
		mtDNA	ND	16	
		17 SSR	F_{IS}, H_O, H_E	17	16
		mtDNA	ND	17	16
	Cleveland Bay horse	10 SSR	H_O	18	
		15 SSR	$N_{e\text{-}F}$, H_O, H_E	19	This study & 20-22
		Pedigree	$N_{e\text{-}F}$, F_{PED}	23	
		15 SSR	H_O, H_E	23	
	Clydesdale horse	10 SSR	H_O	18	
		15 SSR	$N_{e\text{-}F}$, H_O, H_E	19	This study & 20-22
	Connemara pony	394 SSR	H_O	24	
		50K SNP	H_O, H_E	16	
		17 SSR	H_O, H_E	16	
		mtDNA	ND	16	
		17 SSR	F_{IS}, H_O, H_E	17	16
		mtDNA	ND	17	16
		400K SNP	F_{ROH}	25	
	Dales pony	10 SSR	H_O	18	
		15 SSR	H_O, H_E	23	26
		15 SSR	$N_{e\text{-}F}$, H_O, H_E	19	This study & 20-22
	Dartmoor pony	17 SSR	F_{IS}, H_O, H_E	17	
		mtDNA	ND	17	
	Eriskay pony	10 SSR	H_O	18	
		15 SSR	$N_{e\text{-}F}$, H_O, H_E	19	This study & 20-22
	Exmoor pony	10 SSR	H_O	18	
		50K SNP	$N_{e\text{-}LD}$, F_{IS}	27	
		600K SNP	F_{ROH}	28	
		15 SSR	H_O, H_E	23	29
		15 SSR	$N_{e\text{-}F}$, H_O, H_E	19	This study & 20-22
	Fell pony	10 SSR	H_O	18	
		50K SNP	$N_{e\text{-}LD}$, F_{IS}	27	
		17 SSR	F_{IS}, H_O, H_E	17	
		mtDNA	ND	17	
		15 SSR	H_O, H_E	23	26
		15 SSR	$N_{e\text{-}F}$, H_O, H_E	19	This study & 20-22

Breed	Marker	Parameters		Reference
Hackney horse	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Hackney pony	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Highland pony	10 SSR	H_O	18	
	50K SNP	H_O, H_E	16	
	17 SSR	H_O, H_E	16	
	mtDNA	ND	16	
	17 SSR	F_{IS}, H_O, H_E	17	16
	mtDNA	ND	17	16
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Irish Draught horse	10 SSR	H_O	18	
	15 SSR	H_O, H_E	23	30
Kerry Bog pony	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
New Forest pony	50K SNP	N_{e-LD}, F_{IS}	27	
	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Shetland pony	13 SSR	H_O, H_E	31	
	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
	15 SSR	H_O, H_E	23	32
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Shire horse	10 SSR	H_O	18	
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Suffolk Punch horse	10 SSR	H_O	18	
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Thoroughbred horse	Pedigree	F_{PED}	33	
	13 SSR	F_{IS}, H_O, H_E	31	
	Pedigree	F_{PED}	31	
	10 SSR	H_O	18	
	15 SSR	H_O, H_E	34	
	394 SSR	H_O	24	
	30K SNP	N_{e-LD}	35	
	50K SNP	N_{e-LD}, F_{IS}	27	
	15 SSR	H_O, H_E	23	24
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
	30K SNP	F_{ROH}	36	
Warmblood horse	400K SNP	F_{ROH}	25	
	400K SNP	F_{ROH}	25	
Welsh pony	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Section A Welsh pony	50K SNP	H_O, H_E	16	
	17 SSR	H_O, H_E	16	
	mtDNA	ND	16	
	17 SSR	H_O, H_E	17	16
	mtDNA	ND	17	16
Section B Welsh pony	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
Section C Welsh pony	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
Section D Welsh pony	50K SNP	H_O, H_E	16	
	17 SSR	H_O, H_E	16	
	mtDNA	ND	16	

		17 SSR	F_{IS}, H_O, H_E	17	16
		mtDNA	ND	17	16
Goat	Bilberry	50K SNP	F_{ROH}, H_O, H_E	37	38
	Old Irish Goat	50K SNP	F_{ROH}, H_O, H_E	37	38
	Traditional Aran	50K SNP	F_{ROH}, H_O, H_E	37	38
Pig	British Lop	60K SNP	N_{e-F}, N_{e-LD}, F_{PED}, F_{ROH}	39	
Sheep	Badger Faced Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Balwen	50K SNP	N_{e-LD}, F_{IS}	40	
	Belclare	50K SNP	N_{e-LD}, F_{ROH}	41	
		600K SNP	F_{IS}, H_O, H_E	42	
		Pedigree	N_{e-F}, F_{PED}	43	
	Beltex	50K SNP	N_{e-LD}, F_{ROH}	41	
	Beulah	50K SNP	N_{e-LD}, F_{IS}	40	
	Black Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Blue-faced Leicester	600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
	Border Leicester	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
		50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
		600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	46
		50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	Boreray	50K SNP	F_{IS}, H_E	44	
	Brecknock Hill Cheviot	50K SNP	N_{e-LD}, F_{IS}	40	
	Charolais	50K SNP	N_{e-LD}, F_{ROH}	41	
		Pedigree	N_{e-F}, F_{PED}	43	
	Cheviot (commercial)	50K SNP	F_{ROH}, H_O	47	
	Clun Forest	50K SNP	N_{e-LD}, F_{IS}	40	
	Dolgellau Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Dorset Horn	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
		50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
	EasyCare	50K SNP	F_{ROH}, H_O	47	
	Exmoor Horn	31 SSR	N_{e-F}, H_O, H_E	48	
	Galway	Pedigree	F_{PED}	49	
		50K SNP	N_{e-LD}, F_{IS}, H_E	44	
		50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	
		Pedigree	N_{e-F}, F_{PED}	43	
	Greyface Dartmoor	50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	Hill Flock Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Hill Radnor	50K SNP	N_{e-LD}, F_{IS}	40	
	Improved Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Kerry Blackface	600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
	Kerry Hill	50K SNP	N_{e-LD}, F_{IS}	40	
	Lincoln Longwool	50K SNP	F_{ROH}, H_O	47	
	Llandovery White Faced	50K SNP	N_{e-LD}, F_{IS}	40	
	Llanwenog	50K SNP	N_{e-LD}, F_{IS}	40	
	Lleyn	50K SNP	N_{e-LD}, F_{IS}	40	
		600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
		50K SNP	F_{ROH}, H_O	47	
	Norfolk Horn	50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	North Ronaldsay	50K SNP	N_{e-LD}, F_{ROH}	50	
	Oxford Down	50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	Poll Dorset	50K SNP	F_{ROH}, H_O	47	
	Ryeland	50K SNP	F_{ROH}, H_O	47	
	Scottish Blackface	31 SSR	N_{e-F}, H_O, H_E	48	
		50K SNP	N_{e-LD}, F_{IS}, H_E	44	

	50K SNP	$N_{e\text{-LD}}$, F_{IS}, F_{ROH}, H_O	45	44
	50K SNP	$N_{e\text{-F}}$, F_{ROH}, H_O	47	
Soay	50K SNP	$N_{e\text{-LD}}$, F_{IS}, H_E	44	
	50K SNP	$N_{e\text{-LD}}$, F_{IS}, F_{ROH}, H_O	45	44
South Wales Welsh Mountain	50K SNP	$N_{e\text{-LD}}$, F_{IS}	40	
Southdown sheep	600K SNP	F_{IS}, H_O, H_E	42	
Suffolk	50K SNP	$N_{e\text{-LD}}$, F_{IS}, H_E	44	
	50K SNP	$N_{e\text{-LD}}$, F_{ROH}	41	
	50K SNP	$N_{e\text{-LD}}$, F_{IS}, F_{ROH}, H_O	45	
	Pedigree	$N_{e\text{-F}}$, F_{PED}	43	
	50K SNP	F_{ROH}, H_O	47	
Swaledale	31 SSR	$N_{e\text{-F}}$, H_O, H_E	48	
Talybont Welsh Mountain	50K SNP	$N_{e\text{-LD}}$, F_{IS}	40	
Texel	50K SNP	$N_{e\text{-LD}}$, F_{IS}, H_E	44	
	50K SNP	$N_{e\text{-LD}}$, F_{ROH}	41	
	Pedigree	$N_{e\text{-F}}$, F_{PED}	43	
Tregaron Welsh Mountain	50K SNP	$N_{e\text{-LD}}$, F_{IS}	40	
Vendeen	50K SNP	$N_{e\text{-LD}}$, F_{ROH}	41	
	Pedigree	$N_{e\text{-F}}$, F_{PED}	43	
Waterford Blackface	600K SNP	$N_{e\text{-LD}}$, F_{IS}, H_O, H_E	42	
Welsh Hardy Speckled Faced	50K SNP	$N_{e\text{-LD}}$, F_{IS}	40	
Welsh Mountain	31 SSR	$N_{e\text{-F}}$, H_O, H_E	48	
Whiteface Woodland	50K SNP	F_{ROH}, H_O	47	
Wicklow Cheviot	600K SNP	$N_{e\text{-LD}}$, F_{IS}, H_O, H_E	42	
Wiltshire Horn	50K SNP	$N_{e\text{-LD}}$, F_{IS}, H_E	44	
	50K SNP	$N_{e\text{-LD}}$, F_{IS}, F_{ROH}, H_O	45	44
	50K SNP	$N_{e\text{-F}}$, F_{ROH}, H_O	47	

Observed heterozygosity (H_O); expected heterozygosity (H_E); nucleotide diversity (ND) using (Tajima, 1983), (Arlequin v.3.5 (Excoffier and Lischer, 2010; Nei, 1987)), or (DnaSP (Librado and Rozas, 2009; Nei, 1987)); pedigree-based inbreeding coefficient (F_{PED}) using (Meuwissen and Luo, 1992), (Henderson, 1976), (Wright and McPhee, 1925), (Gutiérrez and Goyache, 2005; Wright, 1931), (ABTK (Golden et al., 1991; Henderson, 1976)), or an unknown pedigree-based method; inbreeding coefficient (F_{IS}) using (Wright, 1950), (PLINK (Purcell et al., 2007; Wright, 1950)), or an unknown method likely based on (Wright, 1950); runs of homozygosity inbreeding coefficient (F_{ROH}); LD-based effective population size ($N_{e\text{-LD}}$) using (Barbato et al., 2015), (Tenesa et al., 2007), (Weir and Hill, 1980), (Hayes et al., 2003), or (Kijas et al., 2012); F-based effective population size ($N_{e\text{-F}}$), with F estimates (Hill, 1972), (Weir and Cockerham, 1984), (ENDOG (Gutiérrez and Goyache, 2005; Wright, 1950)), (Jorde and Ryman, 1995).

(1) MacHugh et al., 1998; (2) MacHugh et al., 1999; (3) Bradley et al., 1996; (4) Wiener et al., 2004; (5) Mc Parland et al., 2007; (6) Browett et al., 2018; (7) Decker et al., 2009; (8) Williams et al., 2016; (9) (Orozco-terWengel et al., 2015); (10) The Bovine HapMap Consortium et al., 2009; (11) Roughsedge et al., 1999; (12) Wall et al., 2005; (13) Mc Parland et al., 2007; (14) MacHugh et al., 1997; (15) Wilkinson et al., 2012; (16) Winton et al., 2013; (17) Winton et al., 2020; (18) Alderson (Bodo et al., 2005; (19) Funk et al., 2020; (20) Khanshour et al., 2013; (21) Khanshour et al., 2015; (22) Juras et al., 2003; (23) Dell et al., 2020; (24) Gu et al., 2009; (25) Lindsay-McGee et al., 2023; (26) Fox-Clipsham et al., 2011; (27) Petersen et al., 2013; (28) Grilz-Seger et al., 2019; (29) Prystupa et al., 2012; (30) McGahern et al., 2006; (31) Cunningham et al., 2001; (32) Brinkmann et

al., 2012; (33) Mahon and Cunningham, 1982; (34) Cunningham (Bodo et al., 2005); (35) Corbin et al., 2010; (36) Hill et al.,

2022; (37) Manunza et al., 2023; (38) Stella et al., 2018; (39) Banos et al., 2022; (40) Beynon et al., 2015; (41) Purfield et al.,

2017; (42) Cao et al., 2021; (43) Rafter et al., 2022; (44) Kijas et al., 2012; (45) McHugo et al., 2019; (46) Kijas et al., 2014; (47)

Kerr et al., 2024; (48) Peter et al., 2007; (49) Martin, 1975; (50) Banos, 2023.

Supplementary file 1: Supplementary_file_1.xlsx – Effective population size,

inbreeding, observed and expected heterozygosity, and nucleotide diversity of UK

and Irish livestock and equine populations