Review: Assessing available genetic diversity estimates of rare breeds of

livestock

C. M. Rochus [a], C. F. Price [b], I. Pocrnić [a]

[a] *Division of Quantitative Biology, Roslin Institute, Royal (Dick) School of Veterinary*

Studies, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG,

United Kingdom

[b] *Rare Breeds Survival Trust, Stoneleigh Park, Kenilworth, CV8 2LG,* United Kingdom

Corresponding author: Christina Rochus. Email: crochus@ed.ac.uk

Abstract

The United Kingdom is home to many local and rare livestock breeds. The local

breed populations are highly adapted to specific environments in the United Kingdom

and these and other rare breeds provide solutions to niche needs. Rare breeds often

contribute more than expected to overall species genetic diversity, which is important

because genetic variation is needed for adapting to new challenges. It is therefore

very important from both United Kingdom and global perspectives to maintain genetic

diversity of rare livestock breeds in the United Kingdom, and to do this, we need to

evaluate the monitoring of genetic diversity to identify gaps in our knowledge and

prioritise resources reserved for conservation purposes. The objectives of this study

were to survey the literature to: (1) summarise genetic/genomic characterisation

(effective population size (N_e) and inbreeding) of domestic populations (livestock and

equine) in the United Kingdom and Ireland; (2) compare number of populations on

the United Kingdom's Rare Breeds Survival Trust watch list and the number of United

Kingdom and Irish populations in the peer-reviewed literature with inbreeding, genetic

diversity and/or N_e estimates; and (3) compare annually reported (census-based)

estimates of N_e with (inbreeding- and DNA-based) estimates from peer-reviewed

literature. We found a total of 37 publications with N_e or inbreeding estimates for

United Kingdom or Ireland livestock populations, published from 1975 to 2024. While

many (42%) of the breeds on the Rare Breeds Survival Trust watchlist have been

included in publications, there are still many breeds, and a few species (turkey, duck

and geese) with no publicly available pedigree- or DNA-based genetic diversity

measures. We found census-based N_e estimates were, on average, higher than

DNA-based estimates, likely due to violated assumptions when estimating N_e with

census-based data because of the way livestock mating systems are designed. Most

peer-reviewed papers estimated genetic diversity measures using pedigree,

microsatellite markers, or single nucleotide polymorphisms (**SNP**) markers. To

identify breed-unique variants responsible for adaptive traits in rare breeds, more

studies using whole-genome sequencing will be needed. Altogether, we have

summarised genetic diversity estimates of United Kingdom livestock populations,

identifying gaps in knowledge.

Keywords: Inbreeding, effective population size, census data, genomic data,

conservation

Implications

Maintaining genetic diversity in livestock helps safeguard against future challenges to

our food production systems. We surveyed publicly available measures of genetic

diversity for farm animal populations in the United Kingdom (including government,

Rare Breeds Survival Trust, and peer-reviewed scientific publications) to identify any

gaps. We found census-based estimates of genetic diversity, which are used to

inform and prioritise resources for conservation efforts, on average, are likely

overestimating available genetic diversity. Additionally, most farm animal breeds had

only these census-based estimates. Genetic diversity estimates could be improved

using pedigree or DNA information, more accurately identifying endangered breeds.

Introduction

Livestock populations in the United Kingdom originated in centres of domestication in

Southwest Asia, East Asia, and Mesoamerica, with the first domesticated species

arriving to the British Isles during the Neolithic transition to agriculture (Larson et al.,

2014; Whittle et al., 2011). Through this process, livestock populations have become

highly adapted to many niche environments (Zeder, 2008). Those populations today

are called local breeds and are often maintained by breed associations for their

unique qualities and nostalgia. These breeds can be described as hardy, many of

which can thrive on low-quality feed not appropriate for human consumption and

habitat in areas and weather conditions where other breeds cannot. Rare breeds

have lost their historic popularity because commercial breeds are more productive

(Canali, 2006). As a consequence of decreasing population sizes, rare breed

societies do not have the resources to support breeding programmes, unlike

contemporary commercial populations that can use genomic selection to inform

mating decisions (DEFRA, 2021). However, there are several roles that rare breeds

fill in the modern world. In the United Kingdom and European Union, niche products,

like cheese and meat, from local breeds are sold with a geographical indicator

(protected designation of origin, and protected geographical indication)

('Geographical indications and quality schemes explained', n.d.; 'Protected

geographical food and drink names: UK GI schemes', 2024). Local breeds in the

United Kingdom are used for conservation grazing to maintain numerous habitats

(grassland, heathland, wood pasture, coastal and floodplains, fen, scrub and scrub

mosaics, saltmarsh and sand dunes habitats) ('Graze with livestock to maintain and

improve habitats', n.d.) and are used in organic farming (Van Diepen et al., 2007).

Local breeds also play an intangible cultural role in their communities providing a

glimpse into the past, including domestication processes and the Neolithic transition

to agriculture, all the way up to modern times. Finally, in an ever-changing world,

these historic breeds can play an important role as reservoirs of genetic variation,

providing the variation needed for sustainable agriculture systems with animals that

can adapt to future challenges, which has been recognised by the United Nations

(UN environment programme, 2022; United Nations General Assembly, 2015).

There are many other important roles for local breeds to play in modern society,

besides maintaining total livestock species genetic diversity. To highlight a few

examples from the United Kingdom, we present breeds with unique traits and their

current and historic uses.   The North Ronaldsay sheep, found primarily in the

Orkneys has adapted to primarily eat seaweed, mainly composed of brown kelp, after

they were mostly contained to the shore in the 19th century to protect cultivated land

from being grazed (RBST, n.d.; Porter, 2002). Many modern breeds struggle with

some functional traits as a result of a focus on improving production traits (Rauw et

al., 1998). In contrast, local breeds are prized for these functional traits, and

Shetland cattle are an example of this with their excellent calving ease (Schwabe and

Hall, 1989), they can be bred with larger continental breeds without calving issues

(RBST, n.d.). Local breeds can provide unique animal products that cannot be

provided by commercial breeds. One of the first protected animal products in the

European Union was the Gloucestershire Old Spots pig (RBST, n.d.). In the

application for this status, it was indicated that texture, and flavour of these pigs are

distinguishable from commercial pigs (The Gloucestershire Old Spots Pig Breeders'

Club, 2006). There were some traditional functions of local breeds that may not be in

their current usage, but could inspire more sustainable practices. Before the use of

pesticides to control parasites Shetland geese were traditionally grazed in pastures

after sheep were grazed to manage parasites like liver fluke (RBST, n.d.). Finally,

many of the local breeds can thrive and forage in any climate, like the Scots Grey, a

traditional Scottish chicken breed traced back to the 16th century (RBST, n.d.). This

ability to thrive outdoors is particularly useful for conservation grazing. Many iconic

landscapes in the United Kingdom, like meadows and grasslands that harbour a third

of British flora, were created from traditional farming practices (Peach, 2019). In the

Great Trossachs Forest, grazing allows for a more diverse environment for wild birds,

and opportunities for less competitive grass species to seed and germinate, among

other benefits (The Great Trossachs Forest, 2025). They use a combination of Luing

cattle, Hebridean sheep and Scottish Blackface sheep, each providing different

grazing patterns to contribute to the park's diversity (The Great Trossachs Forest,

2025).

Local breeds from the British Isles have played an outsized role in global commercial

agriculture today. Breeds from the United Kingdom have had widespread popularity,

for example, purebred and crossbred Angus, Jersey, and Guernsey cattle can be

found in many countries around the world, many global commercial pig companies

have developed lines based on Large White and Yorkshire breeds, and the Leicester

Longwool was widely used in crossbreeding schemes to improve the carcass and

fleece of sheep breeds in Europe, Australia, Canada, New Zealand, and the US

(Porter, 2002). The Leicester Longwool was the product of a new type of breeding,

"in-and-in", developed by English farmer Robert Bakewell in the 18th century, where

inbreeding was used to fix traits in a population (Wykes, 2004). This is believed to be

the first implementation of systematic selective breeding, when in comparison, many

famers to this point had been randomly mating livestock (Wykes, 2004).

The United Kingdom government recognises the importance of maintaining the

genetic diversity of its local breeds and has several national and international

obligations to conserve rare breeds, including the 2011 England Biodiversity

Strategy, the United Nation's Food and Agriculture Organisation's "Global Plan of

Action", the United Nation's Convention on Biological Diversity "Strategic Plan for

Biodiversity 2011-2020", and for input to the European Farm Animal Biodiversity

Information System and Food and Agriculture Organisation's Global Information

System (Beale, 2024). To fulfil these obligations, the Department for Environment,

Food and Rural Affairs in the United Kingdom monitors the number of active sires

and dams for livestock populations by breed to inform and prioritise resources for

conservation efforts (with data for cattle, equine, sheep, goat and pig publically

available as far back as 2000, and for poultry available from 2021) (Beale, 2024).

The Department for Environment, Food and Rural Affairs collects this census data

through collaboration with breed organisations and the Rare Breeds Survival Trust, a

charity established in 1973 that conserves and promotes native livestock and equine

breeds in the United Kingdom (RBST, n.d.). Census data is used to estimate

effective population size (N_e), which is the number of individuals in an idealised

population that would generate the same rate of inbreeding as found in the

population of interest (Falconer and Mackay, 1996), allowing for interpretation and

comparison of N_e values across populations and species.

Additionally, several research groups have studied the genetic diversity and

inbreeding of local and rare breeds, publishing their results in peer-reviewed

literature. These studies were standalone projects or part of global consortia (Bovine

HapMap (The Bovine HapMap Consortium et al., 2009), Sheep Hapmap, Adaptmap

(for goats) (Stella et al., 2018), etc.). The objectives of this study were to survey the

literature to: (1) summarise genetic/genomic characterisation (genetic diversity, N_e

and inbreeding) of domestic populations (livestock and equine) in the United

Kingdom and Ireland; (2) compare number of populations on the Rare Breeds

Survival Trust list and the number of populations in the United Kingdom and Ireland

in the peer-reviewed literature with inbreeding, genetic diversity and/or N_e estimates;

and (3) compare annually reported (census-based) estimates of N_e with (DNA-based)

estimates from peer-reviewed literature. By systematically reviewing the peer-

reviewed genetic diversity literature for livestock populations in the United Kingdom,

we can identify gaps, which will help with prioritising research areas.

Material and methods

We used Web of Science (https://www.webofscience.com) to search (on 2025-01-13)

using the following search term:

(inbreeding OR diversity OR "effective population size") AND (cattle OR cow* OR

bull* OR pig* OR chicken* OR sheep* OR goat* OR horse* OR duck* OR turkey* OR

geese) AND ("UK" OR "United Kingdom" OR "Britain" OR "Scotland" OR "England"

OR "Wales" OR "Northern Ireland" OR "Ireland") AND ("genomic" or "genetic" or

"pedigree").

We searched for inbreeding, diversity and N_e estimates for all species identified on

the Rare Breeds Survival Trust's 2024-2025 watchlist for breeds in the United

Kingdom ('RBST watchlist 2024', 2024). Irish populations were included due to the

shared border between Northern Ireland and Ireland, making some livestock and

equine populations to likely be well-connected. Values were extracted from articles,

tables, and figures (using WebPlotDigitizer (Rohatgi, 2024)). The Rare Breeds

Survival Trust watchlist ('RBST watchlist 2024', 2024) is annually published and used

to prioritise breeds for conservation by considering genetic diversity and number of

breeding females.

Every year the Department for Environment, Food and Rural Affairs collects census

numbers (they can be number of animals registered or an estimate of total number)

from breed organisations and the Rare Breeds Survival Trust, and publishes them

(Beale, 2024). We have used the 2023 release of census and effective population

estimates (Beale, 2024). This data is used to monitor numbers of animals over the

years to help with prioritisation of conservation efforts. The Department for

Environment, Food and Rural Affairs uses the following formula to estimate N_e:

$$N_e = \frac{4N_m N_f}{N_m + N_f}$$

Where N_m is the number of breeding males, and N_f is the number of breeding

females) (Wright, 1931). N_e is the number of individuals that would be needed for the

inbreeding estimate to be true in an idealised population (Falconer and Mackay,

1996). An idealised population has the following conditions: (1) mating is restricted to

a population with no migration, (2) generations are discrete, (3) number of breeding

individuals are the same for every generation, (4) mating is random and includes self-

fertilisation, (5) there is no selection, and (6) mutation is disregarded (Falconer and

Mackay, 1996). The effective population estimate is standardised and comparable

between populations, and can be used to estimate the rate of inbreeding (Wright,

1931). The United Nation's Food and Agriculture Organisation's guidance indicates

N_e should be 50 animals per generation, or more, as an N_e of 50 corresponds to 1%

inbreeding rate per generation (when number of breeding males and females are

equal) (FAO Commission on Genetic Resources for Food and Agriculture, 2015).

Study design

We have summarised the peer-reviewed literature on genetic diversity in livestock

populations in the United Kingdom by counting the number of published papers per

species, population, data type (pedigree, mitochondrial DNA, microsatellite, single

nucleotide polymorphisms (**SNP**), or whole genome sequencing) and number of

populations both in the literature and on the Rare Breeds Survival Trust 2024

watchlist.

Additionally, for all inbreeding- and DNA-based N_e estimates found for populations in

the United Kingdom in the literature, we compared this with census-based N_e

estimates published by the Department for Environment, Food and Rural Affairs. To

make the closest comparisons, we compared census-based estimates from the same

year of DNA sampling that was used in the DNA-based estimates. If the year of

sampling was not available, year of publication was used as a proxy. If there was no

census-based estimate for the year of sampling/year of publication of DNA-based

estimates, the closest year with an estimate was used. If the sampling occurred over

a period of years, census-based N_e estimates from those years were averaged. We

used a paired t-test to determine if census-based and DNA-based N_e estimates were

significantly different.

Results

Summary of literature search

In total, we have included 36 peer-reviewed publications and one report written for

the Rare Breeds Survival Trust, published between 1975 and 2024, all of which have

estimates of genetic diversity, including inbreeding coefficients, nucleotide diversity

or N_e of livestock or equine populations in the United Kingdom or Ireland (see

Supplementary Table S1 for information on publications and diversity estimates

included in this study). The 37 publications are summarised in **Table 1**, including

sources of data for each study (original data from a total of 50 publications). **Figure 1**

shows our workflow, including total number of papers found through the Web of

Science search, number of papers removed due to not meeting certain criteria, and

number of papers included that were not identified in the Web of Science search.

Summary of counts

Of the publications included in our analyses, there were 14 papers on equine

populations, 11 on cattle, ten on sheep, one on pig, one on goat and one on chicken.

Information on breeds studied, and data type are included in **Table 1**. The number of

populations in both the literature and the Rare Breeds Survival Trust 2024 watchlist

are plotted by conservation priority status in **Figure 2**. Notably, there were no goat,

turkey, duck, or geese populations on the Rare Breeds Survival Trust 2024 watchlist

that have estimates of genetic inbreeding, diversity or N_e in the literature.

Comparing genomic-based diversity estimates with census data

Of the N_e estimates from genomic data with census estimates available, 43 (of 53)

census estimates were greater than inbreeding- or linkage disequilibrium (**LD**)-based

estimates. The paired t-test for census-based and inbreeding or LD-based N_e was

significant (p=0.0009). The median difference between the two was 391 (maximum

difference was 24,068). All census-based N_e estimates with their corresponding

inbreeding- or LD-based N_e estimates from the Department for Environment, Food

and Rural Affairs are plotted in **Figure 3**.

Discussion

It is important to maintain rare breeds to adapt to future challenges, to serve niche

markets, for conservation grazing, and to maintain cultural identity and contribute to

sustainable agriculture. However, there are finite resources available for conservation

of these breeds, and they are not immediately profitable in the larger economy, so it

is important to identify gaps, and prioritise certain aspects of livestock conservation.

In this sense, we discuss: (1) work already done in these areas and compare census-

based N_e estimates with DNA-based estimates, (2) potential value of whole genome

sequence data for rare breeds, (3) usage of N_e estimates to guide conservation

strategies, (4) limitations of our study design and (5) future actions.

Effective population size estimates

The well-established threshold of concern for population conservation is a N_e of 50

(FAO Commission on Genetic Resources for Food and Agriculture, 2015). Franklin

(1980) first proposed a short-term aim of a N_e of 50 for small populations (with

livestock populations particularly in mind) as this corresponds to a 1% rate of

inbreeding ($\Delta F = \frac{1}{2N_e}$) (Wright, 1931), which is considered a tolerable level of

inbreeding, considering the negative effect of inbreeding depression. This was part of

the 50/500 rule, where Franklin also proposed a long-term minimum value for N_e of

500 (where mutation offsets the loss of genetic variation through drift) (Franklin,

1980). There have been more recent discussions on updating the 50/500 rule, which

is needed as our knowledge on genome biology has evolved in the following 45

years, however there has been no clear consensus on changing the recommended

minimum N_e (Waples, 2025; Pérez-Pereira et al., 2022; Frankham et al., 2014;

Jamieson and Allendorf, 2012).

The N_e threshold of concern may need to be updated, but N_e estimates as a measure

of genetic diversity also needs to be revisited. We compared census-based and

inbreeding- or LD-based N_e estimates using annually reported census-based

estimates from the Department for Environment, Food and Rural Affairs and

inbreeding- or LD-based estimates from the literature. Census-based N_e estimates

were, on average, higher than inbreeding- or LD-based estimates (with a median

difference of 391), and ~11% (6 of 53) of paired estimates had an inbreeding or LD-

based N_e estimate < 50, with a corresponding census-based N_e > 50. This could

indicate that several populations without LD-based N_e estimates but with census-

based estimates appear to have an N_e > 50 but may actually need conservation

prioritisation.

We have observed very different N_e estimates based on data and methodology.

Estimating and publishing census-based estimates has been valuable as a low-cost

method for estimating N_e and observing trends over time. However, inbreeding and

LD-based methods are estimates based on the genome, the diversity of which we

are trying to conserve. The difference between estimates could be expected due to

the assumptions of the methods used. In the census-based methods, there is an

assumption of an idealised population, which does not reflect the reality of livestock

breeding. LD-based methods for estimating N_e also have assumptions, for example,

an assumption is that LD is produced solely from the finite population size, and non-

random mating and population structure are absent (Waples, 2025). Additionally, any

method using microsatellite or SNP arrays will also contend with ascertainment bias

of those DNA markers, where those markers have been identified usually using

animals not from the population of interest – which is especially problematic for

monitoring rare breeds with unique variants that we are interested in conserving.

The census-based N_e estimates in this study showed that more common breeds in

the United Kingdom had higher estimates. An extreme example of this is the

Holstein-Friesian breed. The United Kingdom census-based N_e estimate for 2023

was 13 154 (Beale, 2024). Holstein-Friesian is a globally popular commercial breed,

and populations in different countries are well-connected to each other, particularly

through similar genetics (global use of artificial insemination with elite bulls' semen).

Therefore, we would expect N_e estimates across countries to be somewhat similar to

the United Kingdom estimate. N_e estimates from the Netherlands (one of the

countries where this breed originated from), Canada, France, Ireland, Italy, and the

United States of America range from 18 to 150 (Ablondi et al., 2022; Makanjuola et

al., 2020; Harmen P. Doekes et al., 2018; H.P. Doekes et al., 2018; Rodríguez-

Ramilo et al., 2015; Danchin-Burge et al., 2011; Mc Parland et al., 2007a; Weigel,

2001). These estimates came from a variety of inbreeding- and LD-based methods

(but not the census-based method). This highlights how estimating and comparing N_e

estimates can be challenging (even the difference between 18 and 150 has vastly

different implications for conservation), and that the census-based United Kingdom

estimate is unrealistic. The footnotes of Table 1 demonstrate the wide range of

diversity estimates, and even methodologies within diversity estimate type, all with

certain assumptions and strengths, rendering meaningful comparisions and

intrepretation of results challenging. In the case of census-based N_e estimates for

livestock populations there is often higher selection intensity in males (fewer active

breeding males than active breeding females), which can be accounted for in

Wright's estimate (Wright, 1931). However, the definition of "active breeding males"

could contribute to unreasonable N_e estimates. For example, some of the active

breeding males (e.g., elite bulls) likely have thousands of offspring, while many active

males only have a very small number of offspring, and this variation in family size is

not accounted for in the census-based estimate.

Whole-genome sequencing

While genotyping strategies are becoming more affordable, especially SNP arrays,

typically rare breeds are raised as a hobby, and it is not financially feasible to have

continual producer-investment in routine genotyping. For producers raising rare

breeds for conservation purposes, SNP arrays are not ideal because they likely will

not contain unique breed-specific variants (due to ascertainment bias, as seen in

Geibel et al. (2021)), which are the main interest of conservation programmes. Next

generation sequencing technologies would be better for conservation studies,

however, they can be more expensive. Whole genome sequencing costs are

decreasing, and there are strategies to reduce those costs even further, including

using low-pass sequencing and imputation, but this requires an initial investment to

establish high coverage genotyping of a reference population to identify breed-unique

variants.

Only one publication included in this study used 15X coverage Illumina 150 base pair

paired-end whole genome sequence data (Lindsay-McGee et al., 2023). Due to the

study design which also used available genotypes from populations genotyped with

an Affymetrix array, only 400K SNPs remained for analyses. In contrast, to give an

idea of the magnitude of the number of variants we could expect from sequencing,

from a publication in 2019 from the 1000 Bull Genome Project where a total of 429

animals from 14 breeds were sequenced, they identified 29.1 million single

nucleotide variants, and 1.7 million insertions-deletions (Hayes et al., 2014). We

could expect that there is a lot of unexplored data from the Lindsay-McGee et al.

(2023) study, and that it would be valuable for further investment in generating whole

genome sequence data for other rare breeds in the United Kingdom, allowing for

information on breed-unique variants to be included in management plans.

Conservation management

As we previously mentioned, monitoring and follow-up conservation strategies are

mainly based on the Food and Agriculture Organisation's guidance (N_e of 50 per

generation, equivalent to 1% inbreeding rate per generation (FAO Commission on

Genetic Resources for Food and Agriculture, 2015)). Current census-based

estimates of N_e should be replaced by inbreeding-based estimates of N_e. For the

census-based estimates of N_e, the Department for Environment, Food and Rural

Affairs is using the number of active sires and dams, which is known because

pedigree information is collected and available. An immediate improvement would

be to use pedigree information to estimate rates of inbreeding, which could then be

used to calculate annual inbreeding-based N_e estimates. In the long-term,

developing a national strategy to regularly collect DNA samples for breeds on the

Rare Breeds Survival Trust watchlist, would allow for DNA-based estimates of

genetic diversity measures.

Having DNA-based contemporary N_e estimates is critical to prioritise

breeds/populations for conservation activities. Furthermore, having genomic

(especially whole genome sequence) data on these breeds would not only allow for

better N_e estimates but would also lead to a better understanding of the general

population structure. This would add value for cost-effective decision making,

particularly linked to choosing the most optimal conservation system (*in situ*, *ex situ*

in vivo, *ex situ in vitro*), optimising the number of individuals within the breed to keep

in the chosen conservation system, as well as opening the door for future activities

(e.g., precise introgression, genome editing, etc.). Conservation activities and

systems based on genomic data were systematically reviewed in Oldenbroek et al.

(2017), the algorithms for selection and optimisation of the number of individuals are

available and can be easily adapted for conservation purposes, e.g., Pocrnic et al.,

(2022); Yu et al., (2014), and for a recent review of the genome editing concepts

applied to livestock populations, including conservation applications, see Clark,

(2020). We particularly mention genome editing technology as there is potential to

introduce breed-unique causal variants into other populations (Leroy et al., 2016).

Alternatively, variants could be introduced by genome editing to make local

populations more resistant to disease or improve their fertility (traits that may suffer

from inbreeding depression), only changing traits to improve the survival of the breed

while maintaining the breed standards/definitions. Finally, a more complex

application of genome editing that could be considered is surrogate sire technology.

Surrogate sires are made sterile by CRISPR-Cas9 editing of *NANOS2* in mammals

(Park et al., 2017) or *DDX4* in chicken (Taylor et al., 2017), and donor-derived stem

cells are transplanted into the gonads (Taylor et al., 2017). The surrogate sire could

then disseminate the donor's sperm through natural breeding. The potential to exploit

surrogate sire technology in commercial livestock breeding was discussed by

Gottardo et al. (2019), but similarly, we could use it in a conservation context to

disseminate local breed sperm. Finally, novel technologies like gene drives are

emerging and could find their application in conservation strategies to help spread

desired alleles in a population (e.g., Faber et al. (2021); Webber et al. (2015); Esvelt

et al. (2014)).

Limitations of this study

Many of the results included in this study either came from or benefitted from

international research consortiums. These consortiums build capacity through the

training of individuals, building databases and sharing resources. However, from

these consortium publications there were several breeds that originated in the United

Kingdom, that were not included in the current study because of uncertainty of

sampling locations. Additionally, breeds originating in the United Kingdom with

populations in other countries, may no longer reflect the current United Kingdom

population anymore due to the use of crossbreeding. Therefore, we did not include

populations unless we were certain that they were United Kingdom or Ireland

samples and if there was a proportion of samples included from another country,

there was some evidence that these populations were well-connected with the

corresponding population in the United Kingdom.

There are more peer-reviewed publications that analyse genotypes from livestock

populations in the United Kingdom than we have included in this publication. We tried

to keep only those publications with some genetic diversity measure, inbreeding or

N_e estimate. This means papers looking at population structure and signatures of

selections, while related, were not included. Sometimes the data from those studies

were used in other publications where the measures we were interested in were

estimated, and we tried to cite both the original source of data and the publication

with the analyses of interest.

From systematically reviewing literature for genetic diversity, inbreeding and N_e

estimates for livestock and equine populations in the United Kingdom and Ireland,

one recommendation for other researchers is clear: include detailed information on

populations analysed, including sampling year, number of herds sampled from and

sampling criteria. This additional information will be helpful for future studies (for

publicly available data eventually used in other studies) and can be used to inform

conservation decisions.

Future actions

This study was designed to identify gaps in our knowledge of the genetic diversity of

livestock populations in the United Kingdom. Most breeds on the Rare Breeds

Survival Trust watchlist do not have DNA-based studies, and for those with

estimates, many are outdated. Most notably missing were any estimates on duck,

geese, goat and turkey breeds. These populations should be prioritised for

generating genomic data, and they need to remain on the Rare Breeds Survival Trust

watchlist and the Department for Environment, Food and Rural Affairs monitoring

programmes, potentially even reassessing conservation categories based on DNA-

based metrics when available. The creation of a centralised database for all rare

livestock breeds in the United Kingdom, integrating the Department for Environment,

Food and Rural Affairs, Rare Breeds Survival Trust and academic sources, would

help with implementing a systematic approach to monitor genetic diversity, and fund

the generation of genomic data to update diversity estimates. Publicly funded

genetic estimates and resulting publications should be open access and allow for

data sharing. Through the standardisation and sharing of data, joint conservation

programmes for transboundary breeds can be developed, and breed connectedness

and genetic diversity measures can be jointly assessed.

Ethics approval

Ethics approval was not needed for this study.

Data and model availability statement

The data were not deposited in an official repository. All data are available in the

supplemental information.

Declaration of generative AI and AI-assisted technologies in the writing

process

We declare that no generative AI or AI-assisted technologies were used in the writing

process.

Author ORCIDs

C. M. Rochus 0000-0002-7570-6847

I. Pocrnić 0000-0001-524-7428

479 Declaration of interest

480 We declare that we have no conflicts of interest.

481 Acknowledgements

487 Financial support statement

488 C.M.R. and I.P. acknowledge support from the Biotechnology and Biological

489 Sciences Research Council (BBSRC) awarded to The Roslin Institute (The University

490 of Edinburgh) (BBS/E/RL/230001A and BBS/E/RL/230001C).

References

Ablondi, M., Sabbioni, A., Stocco, G., Cipolat-Gotet, C., Dadousis, C., Kaam, J.-T.V., Finocchiaro, R., Summer, A., 2022. Genetic Diversity in the Italian Holstein Dairy Cattle Based on Pedigree and SNP Data Prior and After Genomic Selection. Frontiers in Veterinary Science 8, 773985. doi:10.3389/fvets.2021.773985

Banos, G., 2023. Genomic analysis of the North Ronaldsay sheep. Rare Breeds Survival Trust (RBST), Kenilworth, United Kingdom. Retrieved on 8 September 2025 from https://www.rbst.org.uk/Handlers/Download.ashx?IDMF=d71f4977-b219-479f-be8b-b70702988263

Banos, G., Talenti, A., Chatziplis, D., Sánchez-Molano, E., 2022. Genomic analysis of the rare British Lop pig and identification of distinctive genomic markers. PLOS ONE 17, e0271053. doi:10.1371/journal.pone.0271053

Barbato, M., Orozco-Terwengel, P., Tapio, M., Bruford, M.W., 2015. SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. Frontiers in Genetics 6, 109. doi:10.3389/fgene.2015.00109

Beale, S., 2024. Official Statistics UK farm animal genetic resources (FAnGR): breed inventory results. UK Department for Environment, Food & Rural Affairs. Retrieved on 8 September 2025 from https://www.gov.uk/government/statistics/uk-farm-animal-genetic-resources-fangr-breed-inventory-results

Beynon, S.E., Slavov, G.T., Farré, M., Sunduimijid, B., Waddams, K., Davies, B., Haresign, W., Kijas, J., MacLeod, I.M., Newbold, C.J., Davies, L., Larkin, D.M., 2015. Population structure and history of the Welsh sheep breeds determined by whole genome genotyping. BMC Genetics 16, 65. doi:10.1186/s12863-015-0216-x

Bodo, I., Alderson, L., Langlois, B. (Eds), 2005. Conservation genetics of endangered horse breeds. EAAP Scientific Series, Wageningen Academic Publishers, Wageningen, Netherlands. Retrieved on 8 September 2025 from https://brill.com/edcollbook/title/68563?language=en&srsltid=AfmBOorAnLyai-phwpa_e2vwhxyGbIRGn70_5iUyppsgFkkea8d1VgKS

Bradley, D.G., MacHugh, D.E., Cunningham, P., Loftus, R.T., 1996. Mitochondrial diversity and the origins of African and European cattle. Proceedings of the National Academy of Sciences 93, 5131–5135. doi:10.1073/pnas.93.10.5131

Brinkmann, L., Gerken, M., Riek, A., 2012. Adaptation strategies to seasonal changes in environmental conditions of a domesticated horse breed, the Shetland pony (*Equus ferus caballus*). Journal of Experimental Biology 215, 1061–1068. doi:10.1242/jeb.064832

Browett, S., McHugo, G., Richardson, I.W., Magee, D.A., Park, S.D.E., Fahey, A.G., Kearney, J.F., Correia, C.N., Randhawa, I.A.S., MacHugh, D.E., 2018. Genomic Characterisation of the Indigenous Irish Kerry Cattle Breed. Frontiers in Genetics 9, 51. doi:10.3389/fgene.2018.00051

Canali, G., 2006. Common agricultural policy reform and its effects on sheep and goat market and rare breeds conservation. Small Ruminant Research 62, 207–213. doi:10.1016/j.smallrumres.2005.08.021

Cao, Y.-H., Xu, S.-S., Shen, M., Chen, Z.-H., Gao, L., Lv, F.-H., Xie, X.-L., Wang, X.-H., Yang, H., Liu, C.-B., Zhou, P., Wan, P.-C., Zhang, Y.-S., Yang, J.-Q., Pi,

W.-H., Hehua, Ee., Berry, D.P., Barbato, M., Esmailizadeh, A., Nosrati, M.,
Salehian-Dehkordi, H., Dehghani-Qanatqestani, M., Dotsev, A.V., Deniskova,
T.E., Zinovieva, N.A., Brem, G., Štěpánek, O., Ciani, E., Weimann, C.,
Erhardt, G., Mwacharo, J.M., Ahbara, A., Han, J.-L., Hanotte, O., Miller, J.M.,
Sim, Z., Coltman, D., Kantanen, J., Bruford, M.W., Lenstra, J.A., Kijas, J., Li,
M.-H., 2021. Historical Introgression from Wild Relatives Enhanced Climatic
Adaptation and Resistance to Pneumonia in Sheep. Molecular Biology and
Evolution 38, 838–855. doi:10.1093/molbev/msaa236

Clark, E.L., 2020. Breeding in an Era of Genome Editing. In Encyclopedia of
Sustainability Science and Technology (ed. Meyers, R.A.). Springer New York,
New York, NY, pp. 1–16. doi:10.1007/978-1-4939-2493-6_1122-1

Corbin, L.J., Blott, S.C., Swinburne, J.E., Vaudin, M., Bishop, S.C., Woolliams, J.A.,
2010. Linkage disequilibrium and historical effective population size in the
Thoroughbred horse. Animal Genetics 41, 8–15. doi:10.1111/j.1365-
2052.2010.02092.x

Cunningham, E.P., Dooley, J.J., Splan, R.K., Bradley, D.G., 2001. Microsatellite
diversity, pedigree relatedness and the contributions of founder lineages to
thoroughbred horses. Animal Genetics 32, 360–364. doi:10.1046/j.1365-
2052.2001.00785.x

Danchin-Burge, C., Hiemstra, S.J., Blackburn, H., 2011. Ex situ conservation of
Holstein-Friesian cattle: Comparing the Dutch, French, and US germplasm
collections. Journal of Dairy Science 94, 4100–4108. doi:10.3168/jds.2010-
3957

Decker, J.E., Pires, J.C., Conant, G.C., McKay, S.D., Heaton, M.P., Chen, K.,
Cooper, A., Vilkki, J., Seabury, C.M., Caetano, A.R., Johnson, G.S.,
Brenneman, R.A., Hanotte, O., Eggert, L.S., Wiener, P., Kim, J.-J., Kim, K.S.,
Sonstegard, T.S., Van Tassell, C.P., Neibergs, H.L., McEwan, J.C., Brauning,
R., Coutinho, L.L., Babar, M.E., Wilson, G.A., McClure, M.C., Rolf, M.M., Kim,
J., Schnabel, R.D., Taylor, J.F., 2009. Resolving the evolution of extant and
extinct ruminants with high-throughput phylogenomics. Proceedings of the
National Academy of Sciences 106, 18644–18649.
doi:10.1073/pnas.0904691106

DEFRA, 2021. Policy paper: Native livestock breeds: reducing extinction risk.
Department for Environment, Food & Rural Affairs. Retrieved on 8 September
2025 from https://www.gov.uk/government/publications/native-livestock-
breeds-reducing-extinction-risk/native-livestock-breeds-reducing-extinction-risk

Dell, A., Curry, M., Yarnell, K., Starbuck, G., Wilson, P.B., 2020. Genetic analysis of
the endangered Cleveland Bay horse: A century of breeding characterised by
pedigree and microsatellite data. PLOS ONE 15, e0240410.
doi:10.1371/journal.pone.0240410

Doekes, H.P., Veerkamp, R.F., Bijma, P., Hiemstra, S.J., Windig, J., 2018. Value of
the Dutch Holstein Friesian germplasm collection to increase genetic
variability and improve genetic merit. Journal of Dairy Science 101, 10022–
10033. doi:10.3168/jds.2018-15217

Doekes, Harmen P., Veerkamp, R.F., Bijma, P., Hiemstra, S.J., Windig, J.J., 2018.
Trends in genome-wide and region-specific genetic diversity in the Dutch-
Flemish Holstein–Friesian breeding program from 1986 to 2015. Genetics
Selection Evolution 50, 15. doi:10.1186/s12711-018-0385-y

Esvelt, K.M., Smidler, A.L., Catteruccia, F., Church, G.M., 2014. Concerning RNA-
589   guided gene drives for the alteration of wild populations. eLife 3, e03401.
590   doi:10.7554/eLife.03401

591   Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs
592   to perform population genetics analyses under Linux and Windows. Molecular
593   Ecology Resources 10, 564–567. doi:10.1111/j.1755-0998.2010.02847.x

594   Faber, N.R., McFarlane, G.R., Gaynor, R.C., Pocrnic, I., Whitelaw, C.B.A., Gorjanc,
595   G., 2021. Novel combination of CRISPR-based gene drives eliminates
596   resistance and localises spread. Scientific Reports 11, 3719.
597   doi:10.1038/s41598-021-83239-4

598   Falconer, D.S., Mackay, T.F.C., 1996. Introduction to Quantitative Genetics. Pearson
599   Education Limited, Harlow, England.

600   FAO Commission on Genetic Resources for Food and Agriculture, 2015. The second
601   report on the state of the world's animal genetic resorces for food and
602   agriculture. FAO, Rome, Italy.

603   Fox-Clipsham, L.Y., Brown, E.E., Carter, S.D., Swinburne, J.E., 2011. Population
604   screening of endangered horse breeds for the foal immunodeficiency
605   syndrome mutation. Veterinary Record 169, 655–655. doi:10.1136/vr.100235

606   Frankham, R., Bradshaw, C.J.A., Brook, B.W., 2014. Genetics in conservation
607   management: Revised recommendations for the 50/500 rules, Red List criteria
608   and population viability analyses. Biological Conservation 170, 56–63.
609   doi:10.1016/j.biocon.2013.12.036

610   Franklin, 1980. Evolutionary changes in small populations. In Conservation Biology.
611   An Evolutionary-Ecological Perspective. Sinauer Associates, Sunderland, MA,
612   USA, pp. 135–149.

613   Funk, S.M., Guedaoura, S., Juras, R., Raziq, A., Landolsi, F., Luís, C., Martínez,
614   A.M., Musa Mayaki, A., Mujica, F., Oom, M.D.M., Ouragh, L., Stranger, Y.,
615   Vega-Pla, J.L., Cothran, E.G., 2020. Major inconsistencies of inferred
616   population genetic structure estimated in a large set of domestic horse breeds
617   using microsatellites. Ecology and Evolution 10, 4261–4279.
618   doi:10.1002/ece3.6195

619   Geibel, J., Reimer, C., Weigend, S., Weigend, A., Pook, T., Simianer, H., 2021. How
620   array design creates SNP ascertainment bias. PLOS ONE 16, e0245178.
621   doi:10.1371/journal.pone.0245178

622   Geographical indications and quality schemes explained, n.d. . European
623   Commission. Retrieved on 8 September 2025 from
624   https://agriculture.ec.europa.eu/farming/geographical-indications-and-quality-
625   schemes/geographical-indications-and-quality-schemes-explained_en

626   Golden, B.L., Brinks, J.S., Bourdon, R.M., 1991. A performance programmed method
627   for computing inbreeding coefficients from large data sets for use in mixed-
628   model analyses. Journal of Animal Science 69, 3564–3573.
629   doi:10.2527/1991.6993564x

630   Gottardo, P., Gorjanc, G., Battagin, M., Gaynor, R.C., Jenko, J., Ros-Freixedes, R.,
631   Bruce A. Whitelaw, C., Mileham, A.J., Herring, W.O., Hickey, J.M., 2019. A
632   Strategy To Exploit Surrogate Sire Technology in Livestock Breeding
633   Programs. G3 Genes|Genomes|Genetics 9, 203–215.
634   doi:10.1534/g3.118.200890

635   Graze with livestock to maintain and improve habitats, n.d.. UK Department for
636   Environment, Food & Rural Affairs. Retrieved on 8 September 2025 from

https://defrafarming.blog.gov.uk/graze-with-livestock-to-maintain-and-improve-
habitats/

Grilz-Seger, G., Neuditschko, M., Ricard, A., Velie, B., Lindgren, G., Mesarič, M.,
Cotman, M., Horna, M., Dobretsberger, M., Brem, G., Druml, T., 2019.
Genome-Wide Homozygosity Patterns and Evidence for Selection in a Set of
European and Near Eastern Horse Breeds. Genes 10, 491.
doi:10.3390/genes10070491

Gu, J., Orr, N., Park, S.D., Katz, L.M., Sulimova, G., MacHugh, D.E., Hill, E.W., 2009.
A Genome Scan for Positive Selection in Thoroughbred Horses. PLoS ONE 4,
e5767. doi:10.1371/journal.pone.0005767

Gutiérrez, J.P., Goyache, F., 2005. A note on ENDOG: a computer program for
analysing pedigree information. Journal of Animal Breeding and Genetics 122,
172–176. doi:10.1111/j.1439-0388.2005.00512.x

Hayes, B.J., Macleod, I.M., Daetwyler, H.D., Bowman, P.J., Chamberlian, A.J., Jagt,
C.J.V., Capitan, A., Pausch, H., Stothard, P., Liao, X., Schrooten, C., Mullaart,
E., Fries, R., Guldbrandtsen, B., Lund, M.S., Boichard, D., Veerkamp, R.F.,
Vantassell, C.P., Gredler, B., Druet, T., Bagnato, A., Vilkki, J., Dekoning, D.J.,
Santus, E., Goddard, M.E., 2014. Genomic prediction from whole genome
sequence in livestock: the 1000 Bull Genomes Project. In Proceedings of the
12th World Congress on Genetics Applied to Livestock Production
(WCGALP), 17-22 August 2014, Vancouver, Canada. Retrieved on 8
September 2025 from https://hal.science/hal-01193911v1/document

Hayes, B.J., Visscher, P.M., McPartlan, H.C., Goddard, M.E., 2003. Novel Multilocus
Measure of Linkage Disequilibrium to Estimate Past Effective Population Size.
Genome Research 13, 635–643. doi:10.1101/gr.387103

Henderson, C.R., 1976. A Simple Method for Computing the Inverse of a Numerator
Relationship Matrix Used in Prediction of Breeding Values. Biometrics 32, 69.
doi:10.2307/2529339

Hill, E.W., Stoffel, M.A., McGivney, B.A., MacHugh, D.E., Pemberton, J.M., 2022.
Inbreeding depression and the probability of racing in the Thoroughbred
horse. Proceedings of the Royal Society B: Biological Sciences 289,
20220487. doi:10.1098/rspb.2022.0487

Hill, W.G., 1972. Effective size of populations with overlapping generations.
Theoretical Population Biology 3, 278–289. doi:10.1016/0040-5809(72)90004-
4

Jamieson, I.G., Allendorf, F.W., 2012. How does the 50/500 rule apply to MVPs?
Trends in Ecology & Evolution 27, 578–584. doi:10.1016/j.tree.2012.07.001

Jorde, P.E., Ryman, N., 1995. Temporal allele frequency change and estimation of
effective size in populations with overlapping generations. Genetics 139,
1077–1090. doi:10.1093/genetics/139.2.1077

Juras, R., Cothran, E.G., Klimas, R., 2003. Genetic Analysis of Three Lithuanian
Native Horse Breeds. Acta Agriculturae Scandinavica, Section A - Animal
Science 53, 180–185. doi:10.1080/09064700310012971

Kerr, E., Marr, M.M., Collins, L., Dubarry, K., Salavati, M., Scinto, A., Woolley, S.,
Clark, E.L., 2024. Analysis of genotyping data reveals the unique genetic
diversity represented by the breeds of sheep native to the United Kingdom.
BMC Genomic Data 25, 82. doi:10.1186/s12863-024-01265-3

Khanshour, A., Juras, R., Blackburn, R., Cothran, E.G., 2015. The Legend of the
Canadian Horse: Genetic Diversity and Breed Origin. Journal of Heredity 106,
37–44. doi:10.1093/jhered/esu074

Khanshour, A.M., Juras, R., Cothran, E.G., 2013. Microsatellite analysis of genetic
variability in Waler horses from Australia. Australian Journal of Zoology 61,
357. doi:10.1071/ZO13062
Kijas, J.W., Lenstra, J.A., Hayes, B., Boitard, S., Porto Neto, L.R., San Cristobal, M.,
Servin, B., McCulloch, R., Whan, V., Gietzen, K., Paiva, S., Barendse, W.,
Ciani, E., Raadsma, H., McEwan, J., Dalrymple, B., other members of the
International Sheep Genomics Consortium, 2012. Genome-Wide Analysis of
the World's Sheep Breeds Reveals High Levels of Historic Mixture and Strong
Recent Selection. PLoS Biology 10, e1001258.
doi:10.1371/journal.pbio.1001258
Kijas, J.W., Porto-Neto, L., Dominik, S., Reverter, A., Bunch, R., McCulloch, R.,
Hayes, B.J., Brauning, R., McEwan, J., the International Sheep Genomics
Consortium, 2014. Linkage disequilibrium over short physical distances
measured in sheep using a high-density SNP chip. Animal Genetics 45, 754–
757. doi:10.1111/age.12197
Larson, G., Piperno, D.R., Allaby, R.G., Purugganan, M.D., Andersson, L., Arroyo-
Kalin, M., Barton, L., Climer Vigueira, C., Denham, T., Dobney, K., Doust,
A.N., Gepts, P., Gilbert, M.T.P., Gremillion, K.J., Lucas, L., Lukens, L.,
Marshall, F.B., Olsen, K.M., Pires, J.C., Richerson, P.J., Rubio de Casas, R.,
Sanjur, O.I., Thomas, M.G., Fuller, D.Q., 2014. Current perspectives and the
future of domestication studies. Proceedings of the National Academy of
Sciences 111, 6139–6146. doi:10.1073/pnas.1323964111
Leroy, G., Besbes, B., Boettcher, P., Hoffmann, I., Capitan, A., Baumung, R., 2016.
Rare phenotypes in domestic animals: unique resources for multiple
applications. Animal Genetics 47, 141–153. doi:10.1111/age.12393
Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of
DNA polymorphism data. Bioinformatics 25, 1451–1452.
doi:10.1093/bioinformatics/btp187
Lindsay-McGee, V., Sanchez-Molano, E., Banos, G., Clark, E.L., Piercy, R.J., Psifidi,
A., 2023. Genetic characterisation of the Connemara pony and the
Warmblood horse using a within-breed clustering approach. Genetics
Selection Evolution 55, 60. doi:10.1186/s12711-023-00827-w
MacHugh, D.E., Loftus, R.T., Cunningham, P., Bradley, D.G., 1998. Genetic structure
of seven European cattle breeds assessed using 20 microsatellite markers.
Animal Genetics 29, 333–340. doi:10.1046/j.1365-2052.1998.295330.x
MacHugh, D.E., Shriver, M.D., Loftus, R.T., Cunningham, P., 1997. Microsatellite
DNA Variation and the Evolution, Domestication and Phylogeography of
Taurine and Zebu Cattle (Bos taurus and Bos indicus). Genetics 146, 1071–
1086. doi:https://doi.org/10.1093/genetics/146.3.1071
MacHugh, D.E., Troy, C.S., McCormick, F., Olsaker, I., Eythórsdóttir, E., Bradle,
D.G., 1999. Early medieval cattle remains from a Scandinavian settlement in
Dublin: genetic analysis and comparison with extant breeds. Philosophical
Transactions of the Royal Society of London. Series B: Biological Sciences
354, 99–109. doi:10.1098/rstb.1999.0363
Mahon, G.A.T., Cunningham, E.P., 1982. Inbreeding and the inheritance of fertility in
the thoroughbred mare. Livestock Production Science 9, 743–754.
doi:10.1016/0301-6226(82)90021-5
Makanjuola, B.O., Miglior, F., Abdalla, E.A., Maltecca, C., Schenkel, F.S., Baes, C.F.,
2020. Effect of genomic selection on rate of inbreeding and coancestry and

effective population size of Holstein and Jersey cattle populations. Journal of Dairy Science 103, 5183–5199. doi:10.3168/jds.2019-18013

Manunza, A., Ramirez-Diaz, J., Cozzi, P., Lazzari, B., Tosser-Klopp, G., Servin, B., Johansson, A.M., Grøva, L., Berg, P., Våge, D.I., Stella, A., 2023. Genetic diversity and historical demography of underutilised goat breeds in North-Western Europe. Scientific Reports 13, 20728. doi:10.1038/s41598-023-48005-8

Martin, I., 1975. A Genetic Analysis of the Galway Sheep Breed: 3. Level of Inbreeding in the Pedigree Galway Sheep Breed. Irish Journal of Agricultural Research 14, 269–274.

Mc Parland, S., Kearney, J.F., Rath, M., Berry, D.P., 2007a. Inbreeding Effects on Milk Production, Calving Performance, Fertility, and Conformation in Irish Holstein-Friesians. Journal of Dairy Science 90, 4411–4419. doi:10.3168/jds.2007-0227

Mc Parland, S., Kearney, J.F., Rath, M., Berry, D.P., 2007b. Inbreeding trends and pedigree analysis of Irish dairy and beef cattle populations. Journal of Animal Science 85, 322–331. doi:10.2527/jas.2006-367

McGahern, A.M., Edwards, C.J., Bower, M.A., Heffernan, A., Park, S.D.E., Brophy, P.O., Bradley, D.G., MacHugh, D.E., Hill, E.W., 2006. Mitochondrial DNA sequence diversity in extant Irish horse populations and in ancient horses. Animal Genetics 37, 498–502. doi:10.1111/j.1365-2052.2006.01506.x

McHugo, G., Browett, S., Randhawa, I., Howard, D., Mullen, M., Richardson, I., Park, S., Magee, D., Scraggs, E., Dover, M., Correia, C., Hanrahan, J., MacHugh, D., 2019. A Population Genomics Analysis of the Native Irish Galway Sheep Breed. Frontiers in Genetics 10, 927. doi:10.3389/fgene.2019.00927

Meuwissen, T.H.E., Luo, Z., 1992. Computing inbreeding coefficients in large populations. Genetics, selection, evolution 24, 305–313. doi:10.1186/1297-9686-24-4-305

Nei, M., 1987. Molecular evolutionary genetics, 2. print. ed. Columbia Univ. Pr, New York, NY.

Oldenbroek, J.K., 2017. Management of Animal Genetic Diversity. Wageningen Academic Publishers, Wageningen, the Netherlands.

Orozco-terWengel, P., Barbato, M., Nicolazzi, E., Biscarini, F., Milanesi, M., Davies, W., Williams, D., Stella, A., Ajmone-Marsan, P., Bruford, M.W., 2015. Revisiting demographic processes in cattle with genome-wide population genetic analysis. Frontiers in Genetics 6, 191. doi:10.3389/fgene.2015.00191

Park, K.-E., Kaucher, A.V., Powell, A., Waqas, M.S., Sandmaier, S.E.S., Oatley, M.J., Park, C.-H., Tibary, A., Donovan, D.M., Blomberg, L.A., Lillico, S.G., Whitelaw, C.B.A., Mileham, A., Telugu, B.P., Oatley, J.M., 2017. Generation of germline ablated male pigs by CRISPR/Cas9 editing of the NANOS2 gene. Scientific Reports 7, 40176. doi:10.1038/srep40176

Peach, J., 2019. The great British meadow. Royal Botanic Garden Kew. Retrieved on 8 September 2025 from https://www.kew.org/read-and-watch/great-british-meadow

Pérez-Pereira, N., Wang, J., Quesada, H., Caballero, A., 2022. Prediction of the minimum effective size of a population viable in the long term. Biodiversity and Conservation 31, 2763–2780. doi:10.1007/s10531-022-02456-z

Peter, C., Bruford, M., Perez, T., Dalamitra, S., Hewitt, G., Erhardt, G., 2007. Genetic diversity and subdivision of 57 European and Middle-Eastern sheep breeds. Animal Genetics 38, 37–44. doi:10.1111/j.1365-2052.2007.01561.x

Petersen, J.L., Mickelson, J.R., Rendahl, A.K., Valberg, S.J., Andersson, L.S.,
Axelsson, J., Bailey, E., Bannasch, D., Binns, M.M., Borges, A.S., Brama, P.,
da Câmara Machado, A., Capomaccio, S., Cappelli, K., Cothran, E.G., Distl,
O., Fox-Clipsham, L., Graves, K.T., Guérin, G., Haase, B., Hasegawa, T.,
Hemmann, K., Hill, E.W., Leeb, T., Lindgren, G., Lohi, H., Lopes, M.S.,
McGivney, B.A., Mikko, S., Orr, N., Penedo, M.C.T., Piercy, R.J., Raekallio,
M., Rieder, S., Røed, K.H., Swinburne, J., Tozaki, T., Vaudin, M., Wade, C.M.,
McCue, M.E., 2013. Genome-Wide Analysis Reveals Selection for Important
Traits in Domestic Horse Breeds. PLoS Genetics 9, e54997.
doi:10.1371/journal.pgen.1003211
Pocrnic, I., Lindgren, F., Tolhurst, D., Herring, W.O., Gorjanc, G., 2022. Optimisation
of the core subset for the APY approximation of genomic relationships.
Genetics Selection Evolution 54, 76. doi:10.1186/s12711-022-00767-x
Porter, V. (Ed.), 2002. Mason's World Dictionary of Livestock Breeds, Types and
Varieties. CABI Publishing, Wallingford, Oxon, UK.
Protected geographical food and drink names: UK GI schemes, 2024. UK
Government. Retrieved on 8 September 2025 from
https://www.gov.uk/guidance/protected-geographical-food-and-drink-names-
uk-gi-schemes
Prystupa, J.M., Hind, P., Cothran, E.G., Plante, Y., 2012. Maternal Lineages in
Native Canadian Equine Populations and Their Relationship to the Nordic and
Mountain and Moorland Pony Breeds. Journal of Heredity 103, 380–390.
doi:10.1093/jhered/ess003
Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D.,
Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J., Sham, P.C., 2007. PLINK:
A Tool Set for Whole-Genome Association and Population-Based Linkage
Analyses. The American Journal of Human Genetics 81, 559–575.
doi:10.1086/519795
Purfield, D.C., McParland, S., Wall, E., Berry, D.P., 2017. The distribution of runs of
homozygosity and selection signatures in six commercial meat sheep breeds.
PLOS ONE 12, e0176780. doi:10.1371/journal.pone.0176780
Rafter, P., McHugh, N., Pabiou, T., Berry, D.P., 2022. Inbreeding trends and genetic
diversity in purebred sheep populations. animal 16, 100604.
doi:10.1016/j.animal.2022.100604
Rauw, W.M., Kanis, E., Noordhuizen-Stassen, E.N., Grommers, F.J., 1998.
Undesirable side effects of selection for high production efficiency in farm
animals: a review. Livestock Production Science 56, 15–33.
doi:10.1016/S0301-6226(98)00147-X
RBST, n.d. North Ronaldsay. Rare Breeds Survival Trust. Retrieved on 8 September
2025a from https://www.rbst.org.uk/north-ronaldsay
RBST, n.d. Shetland. Rare Breeds Survival Trust. Retrieved on 8 September 2025b
from https://www.rbst.org.uk/shetland
RBST, n.d. Gloucestershire Old Spots. Rare Breeds Survival Trust. Retrieved on 8
September 2025c from https://www.rbst.org.uk/gloucestershire-old-spots
RBST, n.d. Shetland. Rare Breeds Survival Trust. Retrieved on 8 September 2025d
from https://www.rbst.org.uk/shetland-geese
RBST, n.d. Scots Grey. Rare Breeds Survival Trust. Retrieved on 8 September
2025e from https://www.rbst.org.uk/scots-grey
RBST, n.d. Introduction. Rare Breeds Survival Trust. Retrieved on 8 September
2025f from https://www.rbst.org.uk/our-mission

RBST watchlist 2024, 2024. Rare Breeds Survival Trust (RBST). Retrieved on 8
September 2025 from https://www.rbst.org.uk/news/rbst-watchlist-2024
Rodríguez-Ramilo, S.T., Fernández, J., Toro, M.A., Hernández, D., Villanueva, B.,
2015. Genome-Wide Estimates of Coancestry, Inbreeding and Effective
Population Size in the Spanish Holstein Population. PLOS ONE 10,
e0124157. doi:10.1371/journal.pone.0124157
Rohatgi, A., 2024. WebPlotDigitizer, v4. Retrieved on 8 September 2025 from
https://automeris.io
Roughsedge, T., Brotherstone, S., Visscher, P.M., 1999. Quantifying genetic
contributions to a dairy cattle population using pedigree analysis. Livestock
Production Science 60, 359–369. doi:10.1016/S0301-6226(99)00106-2
Schwabe, A.E., Hall, S.J.G., 1989. Dystocia in nine British breeds of cattle and its
relationship to dimensions of the dam and calf. Veterinary Record 125, 636–
639. doi:10.1136/vr.125.26-27.636
Stella, A., Nicolazzi, E.L., Van Tassell, C.P., Rothschild, M.F., Colli, L., Rosen, B.D.,
Sonstegard, T.S., Crepaldi, P., Tosser-Klopp, G., Joost, S., 2018. AdaptMap:
exploring goat diversity and adaptation. Genetics Selection Evolution 50, 61,
s12711-018-0427–5. doi:10.1186/s12711-018-0427-5
Tajima, F., 1983. EVOLUTIONARY RELATIONSHIP OF DNA SEQUENCES IN
FINITE POPULATIONS. Genetics 105, 437–460.
doi:10.1093/genetics/105.2.437
Taylor, L., Carlson, D.F., Nandi, S., Sherman, A., Fahrenkrug, S.C., McGrew, M.J.,
2017. Efficient TALEN-mediated gene targeting of chicken primordial germ
cells. Development 144, 928–934. doi:10.1242/dev.145367
Tenesa, A., Navarro, P., Hayes, B., Duffy, D., Clarke, G., Goddard, M., Visscher, P.,
2007. Recent human effective population size estimated from linkage
disequilibrium. GENOME RESEARCH 17, 520–526. doi:10.1101/gr.6023607
The Bovine HapMap Consortium, Gibbs, R.A., Taylor, J.F., Van Tassell, C.P.,
Barendse, W., Eversole, K.A., Gill, C.A., Green, R.D., Hamernik, D.L., Kappes,
S.M., Lien, S., Matukumalli, L.K., McEwan, J.C., Nazareth, L.V., Schnabel,
R.D., Weinstock, G.M., Wheeler, D.A., Ajmone-Marsan, P., Boettcher, P.J.,
Caetano, A.R., Garcia, J.F., Hanotte, O., Mariani, P., Skow, L.C., Sonstegard,
T.S., Williams, J.L., Diallo, B., Hailemariam, L., Martinez, M.L., Morris, C.A.,
Silva, L.O.C., Spelman, R.J., Mulatu, W., Zhao, K., Abbey, C.A., Agaba, M.,
Araujo, F.R., Bunch, R.J., Burton, J., Gorni, C., Olivier, H., Harrison, B.E., Luff,
B., Machado, M.A., Mwakaya, J., Plastow, G., Sim, W., Smith, T., Thomas,
M.B., Valentini, A., Williams, P., Womack, J., Woolliams, J.A., Liu, Y., Qin, X.,
Worley, K.C., Gao, C., Jiang, H., Moore, S.S., Ren, Y., Song, X.-Z.,
Bustamante, C.D., Hernandez, R.D., Muzny, D.M., Patil, S., San Lucas, A.,
Fu, Q., Kent, M.P., Vega, R., Matukumalli, A., McWilliam, S., Sclep, G., Bryc,
K., Choi, J., Gao, H., Grefenstette, J.J., Murdoch, B., Stella, A., Villa-Angulo,
R., Wright, M., Aerts, J., Jann, O., Negrini, R., Goddard, M.E., Hayes, B.J.,
Bradley, D.G., Barbosa Da Silva, M., Lau, L.P.L., Liu, G.E., Lynn, D.J.,
Panzitta, F., Dodds, K.G., 2009. Genome-Wide Survey of SNP Variation
Uncovers the Genetic Structure of Cattle Breeds. Science 324, 528–532.
doi:10.1126/science.1167936
The Gloucestershire Old Spots Pig Breeders' Club, 2006. Application for registration
of a TSG council regulation (EC) No 509/2006 'Traditionally farmed
Gloucestershire Old Spots Pork' EC No UK/007-0024 TSG. Retrieved on 8
September 2025 from

https://assets.publishing.service.gov.uk/media/5fd36838d3bf7f306291b5a3/pf
n-traditionally-farmed-gloucs-old-spots-pork.pdf
The Great Trossachs Forest, 2025. Conservation grazing. The Great Trossachs
Forest. Retrieved on 8 September 2025 from
https://www.thegreattrossachsforest.co.uk/conservation-grazing
UN environment programme, 2022. Kumming-Montreal Global biodiversity
framework (No. CBD/COP/15/L.25). Conference of the parties to the
convention on biological diversity, Convention on Biological Diversity,
Montreal, Canada. Retrieved on 8 September 2025 from
https://www.cbd.int/doc/c/e6d3/cd1d/daf663719a03902a9b116c34/cop-15-l-
25-en.pdf
United Nations General Assembly, 2015. Resolution adopted by the General
Assembly on 25 September 2015: 70/1. Transforming our world: the 2030
Agenda for Sustainable Development. Retrieved on 8 September 2025 from
https://docs.un.org/en/A/RES/70/1
Van Diepen, P., McLean, B., Frost, D., 2007. Livestock breeds and organic farming
systems. ADAS Wales and Organic Centre Wales. Retrieved on 8 September
2025 from https://orgprints.org/id/eprint/10822/1/breeds07.pdf
Wall, E., Brotherstone, S., Kearney, J.F., Woolliams, J.A., Coffey, M.P., 2005. Impact
of Nonadditive Genetic Effects in the Estimation of Breeding Values for
Fertility and Correlated Traits. Journal of Dairy Science 88, 376–385.
doi:10.3168/jds.S0022-0302(05)72697-7
Waples, R.S., 2025. The Idiot's Guide to Effective Population Size. Molecular
Ecology e17670. doi:10.1111/mec.17670
Webber, B.L., Raghu, S., Edwards, O.R., 2015. Is CRISPR-based gene drive a
biocontrol silver bullet or global conservation threat? Proceedings of the
National Academy of Sciences 112, 10565–10567.
doi:10.1073/pnas.1514258112
Weigel, K.A., 2001. Controlling Inbreeding in Modern Breeding Programs. Journal of
Dairy Science 84, E177–E184. doi:10.3168/jds.S0022-0302(01)70213-5
Weir, B.S., Cockerham, C.C., 1984. Estimating F-Statistics for the Analysis of
Population Structure. Evolution 38, 1358–1370.
Weir, B.S., Hill, W.G., 1980. EFFECT OF MATING STRUCTURE ON VARIATION IN
LINKAGE DISEQUILIBRIUM. Genetics 95, 477–488.
doi:10.1093/genetics/95.2.477
Whittle, A.W.R., Healy, F., Bayliss, A., Allen, M.J., 2011. Gathering time: dating the
early Neolithic enclosures of southern Britain and Ireland. Oxbow Books,
Oxford.
Wiener, P., Burton, D., Williams, J.L., 2004. Breed relationships and definition in
British cattle: a genetic analysis. Heredity 93, 597–602.
doi:10.1038/sj.hdy.6800566
Wilkinson, S., Wiener, P., Teverson, D., Haley, C.S., Hocking, P.M., 2012.
Characterization of the genetic diversity, structure and admixture of British
chicken breeds. Animal Genetics 43, 552–563. doi:10.1111/j.1365-
2052.2011.02296.x
Williams, J.L., Hall, S.J.G., Del Corvo, M., Ballingall, K.T., Colli, L., Ajmone Marsan,
P., Biscarini, F., 2015. Inbreeding and purging at the genomic Level: the
Chillingham cattle reveal extensive, non‐random SNP heterozygosity. Animal
Genetics 47, 19–27. doi:10.1111/age.12376

Winton, C.L., Hegarty, M.J., McMahon, R., Slavov, G.T., McEwan, N.R.,
936   Davies‑Morel, M.C.G., Morgan, C.M., Powell, W., Nash, D.M., 2013. Genetic
937   diversity and phylogenetic analysis of native mountain ponies of Britain and
938   Ireland reveals a novel rare population. Ecology and Evolution 3, 934–947.
939   doi:10.1002/ece3.507
940   Winton, C.L., McMahon, R., Hegarty, M.J., McEwan, N.R., Davies‑Morel, M.C.G.,
941   Morgan, C., Nash, D.M., 2020. Genetic diversity within and between British
942   and Irish breeds: The maternal and paternal history of native ponies. Ecology
943   and Evolution 10, 1352–1367. doi:10.1002/ece3.5989
944   Wright, S., 1950. THE GENETICAL STRUCTURE OF POPULATIONS. Annals of
945   Eugenics 15, 323–354. doi:10.1111/j.1469-1809.1949.tb02451.x
946   Wright, S., 1931. Evolution in Mendelian populations. Genetics 16, 97–159.
947   doi:10.1093/genetics/16.2.97
948   Wright, S., McPhee, H.C., 1925. An approximate method of calculating coefficients of
949   inbreeding and relationship from livestock pedigrees. Journal of Agricultural
950   Research 31, 377–383.
951   Wykes, D.L., 2004. Robert Bakewell (1725–1795) of Dishley: farmer and livestock
952   improver. The Agricultural History Review 52, 38–55.
953   Yu, X., Woolliams, J.A., Meuwissen, T.H., 2014. Prioritizing animals for dense
954   genotyping in order to impute missing genotypes of sparsely genotyped
955   animals. Genetics Selection Evolution 46, 46. doi:10.1186/1297-9686-46-46
956   Zeder, M. a, 2008. Domestication and early agriculture in the Mediterranean Basin:
957   Origins, diffusion, and impact. Proceedings of the National Academy of
958   Sciences of the United States of America 105, 11597–11604.
959   doi:10.1073/pnas.0801317105
960
961

 Tables

 Table 1: Publications characterising genetic diversity of United Kingdom and/or Irish

 livestock and equine populations

Species	Populations	Data	Methods	Publication	Data from
Cattle	Aberdeen Angus	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	3
		30 SSR	H_O, H_E	4	
	Angus	Pedigree	F_{PED}	5	
	Ancient Irish samples	mtDNA	ND	2	
	Ayrshire	30 SSR	H_O, H_E	4	
	Belted Galloway	50K SNP	N_{e-LD}, F_{ROH}, H_O	6	7
	Charolais	Pedigree	F_{PED}	5	
	Chillingham	777K SNP	F_{IS}, F_{ROH}	8	
		777K SNP	F_{IS}	9	
	Devon	50K SNP	H_O	6	7
	Dexter	30 SSR	H_O, H_E	4	
		50K SNP	N_{e-LD}, F_{ROH}, H_O	6	7
	English Longhorn	50K SNP	H_O	6	7
	Galloway	50K SNP	H_O	6	7
	Guernsey	30 SSR	H_O, H_E	4	
		50K SNP	F_{ROH}, H_O	6	7 & 10
	Hereford	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	3
		30 SSR	H_O, H_E	4	
		Pedigree	N_{e-F}, F_{PED}	5	
	Highland	30 SSR	H_O, H_E	4	
		50K SNP	H_O	6	7
	Holstein-Friesian	Pedigree	F_{PED}	11	
		Pedigree	F_{PED}	12	
		Pedigree	N_{e-F}, F_{PED}	7 & 13	
	Friesian	30 SSR	H_O, H_E	4	
	Jersey	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	3
		30 SSR	H_O, H_E	4	
	Kerry	20 SSR	H_O, H_E	1	
		mtDNA	ND	2	
		50K SNP	N_{e-LD}, F_{ROH}, H_O	6	This study, 1 & 14
	Limousin	Pedigree	F_{PED}	5	
	Simmental	Pedigree	N_{e-F}, F_{PED}	5	
	South Devon	50K SNP	H_O	6	7
	Sussex	50K SNP	H_O	6	7
	Welsh Black	50K SNP	H_O	6	7
	White Park	777K SNP	F_{IS}	9	
		50K SNP	H_O	6	7
Chicken	Appenzeller	30 SSR	F_{IS}, H_O, H_E	15	
	Araucana	30 SSR	F_{IS}, H_O, H_E	15	
	Brahma	30 SSR	F_{IS}, H_O, H_E	15	
	Buff Orpington	30 SSR	F_{IS}, H_O, H_E	15	
	Cochin	30 SSR	F_{IS}, H_O, H_E	15	
	Croad Langshan	30 SSR	F_{IS}, H_O, H_E	15	

	Derbyshire Redcap	30 SSR	F_{IS}, H_O, H_E	15	
	Dorking	30 SSR	F_{IS}, H_O, H_E	15	
	Hamburgh	30 SSR	F_{IS}, H_O, H_E	15	
	Indian Game	30 SSR	F_{IS}, H_O, H_E	15	
	Ixworth	30 SSR	F_{IS}, H_O, H_E	15	
	Leghorn (coloured)	30 SSR	F_{IS}, H_O, H_E	15	
	Light Sussex	30 SSR	F_{IS}, H_O, H_E	15	
	Lincolnshire Buff	30 SSR	F_{IS}, H_O, H_E	15	
	Maran	30 SSR	F_{IS}, H_O, H_E	15	
	Marsh Daisy	30 SSR	F_{IS}, H_O, H_E	15	
	Norfolk Grey	30 SSR	F_{IS}, H_O, H_E	15	
	Old English Pheasant Fowl	30 SSR	F_{IS}, H_O, H_E	15	
	Rhode Island Red	30 SSR	F_{IS}, H_O, H_E	15	
	Scots Dumpy	30 SSR	F_{IS}, H_O, H_E	15	
	Scots Grey	30 SSR	F_{IS}, H_O, H_E	15	
	Silkie	30 SSR	F_{IS}, H_O, H_E	15	
	Spanish	30 SSR	F_{IS}, H_O, H_E	15	
	Sussex	30 SSR	F_{IS}, H_O, H_E	15	
Equine	Carneddau pony	50K SNP	H_O, H_E	16	
		17 SSR	H_O, H_E	16	
		mtDNA	ND	16	
		17 SSR	F_{IS}, H_O, H_E	17	16
		mtDNA	ND	17	16
	Cleveland Bay horse	10 SSR	H_O	18	
		15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
		Pedigree	N_{e-F}, F_{PED}	23	
		15 SSR	H_O, H_E	23	
	Clydesdale horse	10 SSR	H_O	18	
		15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
	Connemara pony	394 SSR	H_O	24	
		50K SNP	H_O, H_E	16	
		17 SSR	H_O, H_E	16	
		mtDNA	ND	16	
		17 SSR	F_{IS}, H_O, H_E	17	16
		mtDNA	ND	17	16
		400K SNP	F_{ROH}	25	
	Dales pony	10 SSR	H_O	18	
		15 SSR	H_O, H_E	23	26
		15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
	Dartmoor pony	17 SSR	F_{IS}, H_O, H_E	17	
		mtDNA	ND	17	
	Eriskay pony	10 SSR	H_O	18	
		15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
	Exmoor pony	10 SSR	H_O	18	
		50K SNP	N_{e-LD}, F_{IS}	27	
		600K SNP	F_{ROH}	28	
		15 SSR	H_O, H_E	23	29
		15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
	Fell pony	10 SSR	H_O	18	
		50K SNP	N_{e-LD}, F_{IS}	27	
		17 SSR	F_{IS}, H_O, H_E	17	

Breed	Marker	Parameters	Ref	Notes
	mtDNA	ND	17	
	15 SSR	H_O, H_E	23	26
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Hackney horse	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Hackney pony	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Highland pony	10 SSR	H_O	18	
	50K SNP	H_O, H_E	16	
	17 SSR	H_O, H_E	16	
	mtDNA	ND	16	
	17 SSR	F_{IS}, H_O, H_E	17	16
	mtDNA	ND	17	16
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Irish Draught horse	10 SSR	H_O	18	
	15 SSR	H_O, H_E	23	30
Kerry Bog pony	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
New Forest pony	50K SNP	N_{e-LD}, F_{IS}	27	
	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Shetland pony	13 SSR	H_O, H_E	31	
	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
	15 SSR	H_O, H_E	23	32
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Shire horse	10 SSR	H_O	18	
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Suffolk Punch horse	10 SSR	H_O	18	
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Thoroughbred horse	Pedigree	F_{PED}	33	
	13 SSR	F_{IS}, H_O, H_E	31	
	Pedigree	F_{PED}	31	
	10 SSR	H_O	18	
	15 SSR	H_O, H_E	34	
	394 SSR	H_O	24	
	30K SNP	N_{e-LD}	35	
	50K SNP	N_{e-LD}, F_{IS}	27	
	15 SSR	H_O, H_E	23	24
	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
	30K SNP	F_{ROH}	36	
Warmblood horse	400K SNP	F_{ROH}	25	
	400K SNP	F_{ROH}	25	
Welsh pony	15 SSR	N_{e-F}, H_O, H_E	19	This study & 20-22
Section A Welsh pony	50K SNP	H_O, H_E	16	
	17 SSR	H_O, H_E	16	
	mtDNA	ND	16	
	17 SSR	H_O, H_E	17	16
	mtDNA	ND	17	16
Section B Welsh pony	17 SSR	F_{IS}, H_O, H_E	17	
	mtDNA	ND	17	
Section C Welsh pony	17 SSR	F_{IS}, H_O, H_E	17	

		mtDNA	ND	17	
	Section D Welsh pony	50K SNP	H_O, H_E	16	
		17 SSR	H_O, H_E	16	
		mtDNA	ND	16	
		17 SSR	F_{IS}, H_O, H_E	17	16
		mtDNA	ND	17	16
Goat	Bilberry	50K SNP	F_{ROH}, H_O, H_E	37	38
	Old Irish Goat	50K SNP	F_{ROH}, H_O, H_E	37	38
	Traditional Aran	50K SNP	F_{ROH}, H_O, H_E	37	38
Pig	British Lop	60K SNP	N_{e-F}, N_{e-LD}, F_{PED}, F_{ROH}	39	
Sheep	Badger Faced Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Balwen	50K SNP	N_{e-LD}, F_{IS}	40	
	Belclare	50K SNP	N_{e-LD}, F_{ROH}	41	
		600K SNP	F_{IS}, H_O, H_E	42	
		Pedigree	N_{e-F}, F_{PED}	43	
	Beltex	50K SNP	N_{e-LD}, F_{ROH}	41	
	Beulah	50K SNP	N_{e-LD}, F_{IS}	40	
	Black Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Blue-faced Leicester	600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
	Border Leicester	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
		50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
		600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	46
		50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	Boreray	50K SNP	F_{IS}, H_E	44	
	Brecknock Hill Cheviot	50K SNP	N_{e-LD}, F_{IS}	40	
	Charolais	50K SNP	N_{e-LD}, F_{ROH}	41	
		Pedigree	N_{e-F}, F_{PED}	43	
	Cheviot (commercial)	50K SNP	F_{ROH}, H_O	47	
	Clun Forest	50K SNP	N_{e-LD}, F_{IS}	40	
	Dolgellau Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Dorset Horn	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
		50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
	EasyCare	50K SNP	F_{ROH}, H_O	47	
	Exmoor Horn	31 SSR	N_{e-F}, H_O, H_E	48	
	Galway	Pedigree	F_{PED}	49	
		50K SNP	N_{e-LD}, F_{IS}, H_E	44	
		50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	
		Pedigree	N_{e-F}, F_{PED}	43	
	Greyface Dartmoor	50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	Hill Flock Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Hill Radnor	50K SNP	N_{e-LD}, F_{IS}	40	
	Improved Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
	Kerry Blackface	600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
	Kerry Hill	50K SNP	N_{e-LD}, F_{IS}	40	
	Lincoln Longwool	50K SNP	F_{ROH}, H_O	47	
	Llandovery White Faced	50K SNP	N_{e-LD}, F_{IS}	40	
	Llanwenog	50K SNP	N_{e-LD}, F_{IS}	40	
	Lleyn	50K SNP	N_{e-LD}, F_{IS}	40	
		600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
		50K SNP	F_{ROH}, H_O	47	
	Norfolk Horn	50K SNP	N_{e-F}, F_{ROH}, H_O	47	
	North Ronaldsay	50K SNP	N_{e-LD}, F_{ROH}	50	
	Oxford Down	50K SNP	N_{e-F}, F_{ROH}, H_O	47	

Breed	Marker	Metrics	Ref	Ref
Poll Dorset	50K SNP	F_{ROH}, H_O	47	
Ryeland	50K SNP	F_{ROH}, H_O	47	
Scottish Blackface	31 SSR	N_{e-F}, H_O, H_E	48	
	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
	50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
	50K SNP	N_{e-F}, F_{ROH}, H_O	47	
Soay	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
	50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
South Wales Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
Southdown sheep	600K SNP	FI_S, H_O, H_E	42	
Suffolk	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
	50K SNP	N_{e-LD}, F_{ROH}	41	
	50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	
	Pedigree	N_{e-F}, F_{PED}	43	
	50K SNP	F_{ROH}, H_O	47	
Swaledale	31 SSR	N_{e-F}, H_O, H_E	48	
Talybont Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
Texel	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
	50K SNP	N_{e-LD}, F_{ROH}	41	
	Pedigree	N_{e-F}, F_{PED}	43	
Tregaron Welsh Mountain	50K SNP	N_{e-LD}, F_{IS}	40	
Vendeen	50K SNP	N_{e-LD}, F_{ROH}	41	
	Pedigree	N_{e-F}, F_{PED}	43	
Waterford Blackface	600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
Welsh Hardy Speckled Faced	50K SNP	N_{e-LD}, F_{IS}	40	
Welsh Mountain	31 SSR	N_{e-F}, H_O, H_E	48	
Whiteface Woodland	50K SNP	F_{ROH}, H_O	47	
Wicklow Cheviot	600K SNP	N_{e-LD}, F_{IS}, H_O, H_E	42	
Wiltshire Horn	50K SNP	N_{e-LD}, F_{IS}, H_E	44	
	50K SNP	N_{e-LD}, F_{IS}, F_{ROH}, H_O	45	44
	50K SNP	N_{e-F}, F_{ROH}, H_O	47	

Abbreviations: H_O = Observed heterozygosity; H_E = expected heterozygosity; ND = nucleotide diversity using (Tajima, 1983),

(Arlequin v.3.5 (Excoffier and Lischer, 2010; Nei, 1987)), or (DnaSP (Librado and Rozas, 2009; Nei, 1987)); F_{PED} = pedigree-

based inbreeding coefficient using (Meuwissen and Luo, 1992), (Henderson, 1976), (Wright and McPhee, 1925), (Gutiérrez and

Goyache, 2005; Wright, 1931), (ABTK (Golden et al., 1991; Henderson, 1976)), or an unknown pedigree-based method; F_{IS} =

inbreeding coefficient using (Wright, 1950), (PLINK (Purcell et al., 2007; Wright, 1950)), or an unknown method likely based on

(Wright, 1950); F_{ROH} = runs of homozygosity inbreeding coefficient; N_{e-LD} = Linkage disequilibrium (LD)-based effective

population size using (Barbato et al., 2015), (Tenesa et al., 2007), (Weir and Hill, 1980), (Hayes et al., 2003), or (Kijas et al.,

2012); N_{e-F} = Inbreeding (F)-based effective population size, with F estimates from (Hill, 1972), (Weir and Cockerham, 1984),

(ENDOG (Gutiérrez and Goyache, 2005; Wright, 1950)), (Jorde and Ryman, 1995); SSR = single sequence repeats, also known

as microsatellites; SNP = single nucleotide polymorphism; mtDNA = mitochondrial DNA.

[1] MacHugh et al., 1998

[2] MacHugh et al., 1999

[3] Bradley et al., 1996

980	[4] Wiener et al., 2004
981	[5] Mc Parland et al., 2007
982	[6] Browett et al., 2018
983	[7] Decker et al., 2009
984	[8] Williams et al., 2016
985	[9] Orozco-terWengel et al., 2015
986	[10] The Bovine HapMap Consortium et al., 2009
987	[11] Roughsedge et al., 1999
988	[12] Wall et al., 2005
989	[13] Mc Parland et al., 2007
990	[14] MacHugh et al., 1997
991	[15] Wilkinson et al., 2012
992	[16] Winton et al., 2013
993	[17] Winton et al., 2020
994	[18] Alderson (Bodo et al., 2005)
995	[19] Funk et al., 2020
996	[20] Khanshour et al., 2013
997	[21] Khanshour et al., 2015
998	[22] Juras et al., 2003
999	[23] Dell et al., 2020
1000	[24] Gu et al., 2009
1001	[25] Lindsay-McGee et al., 2023
1002	[26] Fox-Clipsham et al., 2011
1003	[27] Petersen et al., 2013
1004	[28] Grilz-Seger et al., 2019
1005	[29] Prystupa et al., 2012
1006	[30] McGahern et al., 2006
1007	[31] Cunningham et al., 2001
1008	[32] Brinkmann et al., 2012
1009	[33] Mahon and Cunningham, 1982
1010	[34] Cunningham (Bodo et al., 2005)
1011	[35] Corbin et al., 2010
1012	[36] Hill et al., 2022
1013	[37] Manunza et al., 2023
1014	[38] Stella et al., 2018
1015	[39] Banos et al., 2022
1016	[40] Beynon et al., 2015
1017	[41] Purfield et al., 2017

[42] Cao et al., 2021

[43] Rafter et al., 2022

[44] Kijas et al., 2012

[45] McHugo et al., 2019

[46] Kijas et al., 2014

[47] Kerr et al., 2024

[48] Peter et al., 2007

[49] Martin, 1975

[50] Banos, 2023

1027

Figures

Figure 1: Literature search terms and number of results from Web of Science, for a

search of genetic diversity measures of cattle, chicken, duck, equine, geese, goat,

pig, sheep, and turkey populations in the United Kingdom and Ireland.

Figure 2: Clustered stacked bar plot of the number of United Kingdom and Irish

livestock populations on the Rare Breeds Survival Trust's watchlist with estimates of

inbreeding, diversity or effective population size estimated using pedigree or

molecular markers available ("yes") or not ("no").

Figure 3: Inbreeding (F)- and linkage disequilibrium (LD)-based effective population

size (N_e) estimates of cattle, equine, pig, and sheep breeds. N_e of 50 are indicated on

both axes with grey dashed lines.

1039

1040 **Supplementary files**

1041 **Supplementary Table S1:** Supplementary_file_1.xlsx – Effective population size,

1042 inbreeding, observed and expected heterozygosity, and nucleotide diversity of United

1043 Kingdom and Irish livestock and equine populations

3379 — (inbreeding OR diversity OR "effective population size") AND (cattle OR cow* OR bull* OR pig* OR chicken* OR sheep* OR goat* OR horse* OR duck* OR turkey* OR geese) AND ("UK" OR "United Kingdom" OR "Britain" OR "Scotland" OR "England" OR "Wales" OR "Northern Ireland" OR "Ireland") AND ("genomic" or "genetic" or "pedigree")

384 — Excluded (NOT)
infection OR tropic* OR Africa* OR China OR Chinese OR simulation OR virus OR phage OR strain* OR isolate* OR crop* OR MHC OR zoo* OR zebra* OR pigeon* OR orchid* OR Ethiopia* OR disease OR parasite* OR mice OR bee* OR primate* OR dog* OR plant* OR MOET OR butterfly OR butterflies

23 — Checked for correct species, population sampled in the UK or Ireland, estimated effective population size and/or inbreeding

37 — Added
Papers that were missed that we were aware of
Papers that were original sources of data from the search results
Papers from hand searches