Parkinson's-Linked Synaptojanin 1 Loss in Dopamine Neurons Triggers Synaptic

Degeneration and Striatal TH Interneuron Compensation

Bandhan Mukherjee[1,#], Youneng Lin[1,#], Man Liu[1,&], Zhaoyan Li[1, &], Xiaoxing Cao[1], and Mian

Cao[1,2]*

1. Programme in Neuroscience and Behavioural Disorders, Duke-NUS Medical School,

Singapore

2. Department of Physiology, National University of Singapore, Singapore

#,&: These authors contributed equally

* Corresponding Author

Contact Information: mian.cao@duke-nus.edu.sg

Abstract

Synaptic dysfunction is increasingly recognized as an early hallmark of Parkinson's disease (PD). Synaptojanin 1 (SJ1), a synaptic phosphoinositide phosphatase essential for synaptic vesicle recycling, is genetically linked to early-onset Parkinsonism (EOP). While germline SJ1 knockout mice are perinatal lethal, SJ1-R258Q knock-in mice recapitulate EOP-like symptoms and exhibit selective dystrophy of nigrostriatal dopamine (DA) terminals. However, these whole-body mutants limit understanding of SJ1's role in the DA system. Here, we generated DA neuron-specific SJ1 conditional KO (SJ1-DA cKO) mice. Complete SJ1 loss caused severe synaptic dystrophy throughout the striatum, indicating a cell-autonomous role of SJ1 in both substantia nigra and ventral tegmental area DA subtypes. Surprisingly, despite profound synaptic degeneration and DA deficiency, SJ1-DA cKO mice showed no overt motor deficits. Instead, we observed a robust induction of striatal tyrosine hydroxylase-positive interneurons (iTHINs), which expressed multiple DA markers and formed new connections with degenerating DA terminals, suggesting a potential local compensation. In contrast, acute SJ1 deletion in adult/aged DA neurons caused similar terminal pathology but limited iTHINs induction. Together, our findings reveal a critical role for SJ1 in maintaining DA synaptic function and uncover an adaptive striatal response to DA loss, offering insights into compensatory mechanisms relevant to PD pathogenesis.

Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disorder, primarily affecting movement. The pathological hallmarks of PD include the selective loss of dopamine (DA) neurons in the substantia nigra pars compacta (SNc) and DA deficiency in the striatum.

Although over 90% of PD cases are sporadic, recent genetic studies have identified more than 20 risk genes and 90 genomic risk loci associated with both familial and sporadic PD[1]. Several of these genes regulate endocytic trafficking at presynaptic terminals, suggesting that impaired synaptic vesicle (SV) recycling is an early pathological feature of PD[2]. A major SV recycling pathway involves clathrin-mediated endocytosis (CME), in which Synaptojanin 1 (SJ1), a lipid phosphatase, plays a crucial role[3]. SJ1 facilitates clathrin uncoating by dephosphorylating $PI(4,5)P_2$ on the plasma membrane and releasing clathrin adaptors. In its absence, clathrin-coated vesicles (CCVs) accumulate at synapses, as demonstrated in multiple model organisms[4-7].

A homozygous missense mutation (R258Q) in SJ1 has been previously identified in early-onset Parkinsonism (EOP), establishing SJ1 as the PARK20 gene[8,9]. This mutation inactivates the Sac phosphatase domain, resulting in partial loss-of-function (LOF). Since then, additional SJ1 mutations have been linked to Parkinsonism and epilepsy, most of which are homozygous missense or nonsense mutations[2,10]. A strong genotype-phenotype correlation exists, where nonsense mutations causing premature truncation in the homozygous state led to severe neurodegeneration, seizures, and early mortality[11]. In contrast, homozygous or compound heterozygous missense mutations are associated with milder phenotypes, including Parkinsonism and seizure susceptibility[10,12].

More recently, rare heterozygous SJ1 variants have been implicated in EOP, and Sac domain variants have been linked to sporadic PD[13]. Additionally, a novel expression quantitative trait locus (eQTL) connection between rs10847864 and SJ1 was identified, highlighting convergence between familial and GWAS studies[14]. Furthermore, SJ1 expression is reduced in blood and postmortem PD brain tissues, suggesting it may serve as a biomarker for sporadic PD[15].

Multiple mouse models have been developed to study SJ1 LOF. Complete SJ1 loss in constitutive homozygous knockout (KO) mice results in early mortality and severe epilepsy, mirroring human patients with homozygous nonsense mutations[11,16]. Aged SJ1-haploinsufficient (+/-) mice show PD-like features, including alpha-synuclein accumulation,

autophagy defects, and DA terminal degeneration[17]. They also exhibit impaired astrocytic

glucose sensing and lysosomal dysfunction in midbrain neurons[18,19].

Our previously generated SJ1 knock-in (KI) mice carrying the R258Q mutation (SJ1-KI[RQ])

exhibit neurological symptoms reminiscent of EOP, including early-onset epilepsy and motor

coordination deficits[20]. These mice display generalized synaptic endocytic defects and

selective synaptic degeneration in a subset of DA neurons in nigrostriatal pathway, a hallmark

of PD pathology[20]. Additionally, we demonstrated a synergistic interaction between SJ1 and

auxilin (PARK19), another PD-linked clathrin uncoating protein[21]. Auxilin-KO mice exhibit

dystrophic nigrostriatal DA terminals similar to SJ1-KI[RQ] mice. In double mutant (Aux-KO/SJ1-

KI[RQ]) mice, lifespan is dramatically reduced, and synaptic endocytic defects and striatal

dystrophic changes are exacerbated, further supporting the vulnerability of nigrostriatal

neurons to synaptic endocytic impairments[21].

Synergistic effects of SJ1 with other PD-linked proteins have also been demonstrated in

double mutant mouse models. INPP5F/Sac2, a PI4P phosphatase involved in endosomal

trafficking, has been identified as a PD risk gene through GWAS[22]. Sac2-KO/ SJ1-KI[RQ] mice

show shortened lifespan, more severe endocytic defects, and earlier DA terminal dystrophy[23].

In addition, combining SJ1 heterozygosity with LRRK2 G2019S worsens DA release deficits

and motor impairments in aged mice[24].

While these mouse models have provided valuable insights into SJ1 function in PD, they are

all constitutive whole-body mutants, leading to complex behavioral phenotypes and even early

death due to global SJ1 loss in the brain. To better delineate SJ1's role in DA neurons in PD,

we generated novel DA neuron-specific SJ1 conditional KO models (both genetic and acute

deletions) in this study. Complete SJ1 loss in DA neurons results in severe DA terminal

dystrophy and DA deficiency throughout the striatum, but surprisingly, no overt motor deficits.

Instead, we observed a robust compensatory increase in striatal tyrosine hydroxylase-positive

interneurons (THINs). These induced THINs acquire DA-like properties and form new

connections with DA terminals, potentially compensating for DA loss. Our study not only

reinforces the critical role of SJ1 in DA synaptic transmission but also reveals a novel

compensatory mechanism, which if pharmacologically enhanced, might delay or mitigate PD.

Results

Generation of DA-specific SJ1 conditional knockout (cKO) mice genetically

Given that conventional SJ1 KO mice exhibit perinatal lethality within 24 hours of birth, we
generated SJ1 flx/flx mice by inserting loxP sites flanking exon 4 of *Synj1* gene using
CRISPR/Cas9. These mice were then crossed with the DAT-IRES-Cre line to create DA-
specific SJ1 conditional KO mice (DAT-Cre;SJ1 flx/flx), hereafter referred to as SJ1 cKO[DAT]
(Figures 1A and S1A).

To confirm successful recombination, we performed immunohistochemistry (IHC) on fixed
frozen brain sections using antibodies against SJ1 and DA marker tyrosine hydroxylase (TH).
In SJ1 cKO[DAT] mice, SJ1 protein was selectively depleted in TH-positive DA neurons in the
SNc and ventral tegmental area (VTA), while SJ1 expression in TH-negative neurons and non-
DA brain regions, such as the hippocampus, remained intact (Figure 1B). These findings
confirm efficient and specific KO of SJ1 in DA neurons. SJ1 cKO[DAT] mice survived into
adulthood without seizures, suggesting that epilepsy reported in SJ1-KI[RQ] mice and
constitutive SJ1 KO mice is likely due to SJ1 loss in other brain regions, such as the
hippocampus and cortex[16,20].

DA Terminal Pathology and DA Deficiency in the SJ1 cKO[DAT] Striatum

To investigate DA-specific pathology in SJ1 cKO[DAT] mice, we performed IHC for DA-related
markers in both midbrain (DA soma) and striatum (DA terminals). In the midbrain, TH-positive
DA neurons appeared morphologically intact, with no significant loss of DA neurons in either
the SNc or VTA at 2-3 months old and 6-8 months old (Figure S1B). However, in the striatum,
we observed numerous dopamine transporter (DAT) positive dystrophic clusters in SJ1
cKO[DAT] but not in the control (Figures 1C and 1D). In line with our previous study[20,21], this
striatal pathology emerged as early as three weeks after birth, progressively increasing in size
and number, and peaking at 2–3 months of age (Figure S1E). Importantly, SJ1 cKO[DAT] mice
exhibited this pathology throughout the striatum, including the dorsal and ventral striatum
(Figures 1C and S1C). This differs from the previously reported SJ1-KI[RQ] mice[20], where
dystrophic DA terminals were primarily restricted to the dorsal striatum (nigrostriatal pathway)
(Figure S1D). This widespread pathology observed in the striatum suggests that the complete
loss of SJ1 impacted both SNc and VTA DA neurons. Together, the results from SJ1-KI[RQ] and
SJ1 cKO[DAT] mice indicate that the loss of SJ1 in DA neurons is sufficient to cause DA terminal
dystrophy in a cell-autonomous and gene-dosage dependent manner.

As shown in Figures 1E and S2A, these dystrophic clusters were immunopositive for several
other DA markers, including TH, aromatic L-amino acid decarboxylase (AADC), vesicular
monoamine transporter 2 (vMAT2), and synaptic vesicle glycoprotein 2C (SV2C). Additional
analyses revealed partial colocalization of these DAT-positive clusters with general synaptic

markers such as synaptotagmin-1 (Syt1) and α-synuclein (Figures S2B and S2C). In contrast, the pathological form of α-synuclein (pSer129-α-syn) was not detected in these clusters (Figure S2D). Moreover, these dystrophic clusters lacked colocalization with D2 dopamine receptors (D2R) or the medium spiny neuron (MSN) marker Darpp32, indicating that the pathology is restricted to DA terminals (Figures S2E and S2F). Additionally, immunostaining with microglial marker, ionized calcium-binding adapter molecule (Iba1), and astrocyte marker, glial fibrillary acidic protein (GFAP), did not show any obvious gliosis and inflammation in the striatum (Figures S2G and S2H). Interestingly, these clusters were negative for clathrin-dependent endocytic proteins either (e.g., clathrin light chain, endophilin 1, and amphiphysin 1), suggesting a clathrin-independent mechanism may underlie DA terminal dystrophy (Figures S3A-S3D). Indeed, electron microscopy (EM) of the SJ1 cKO[DAT] striatum revealed multilayered, whirl-like membrane structures, similar to those previously observed in SJ1-KI[RQ] and Auxilin-KO mutant mouse models (Figure 1F)[21]. These membrane accumulations correlate with DA marker protein clusters and likely reflect abnormal plasma membrane invaginations at DA terminals. They envelop various organelles, including SVs, mitochondria, autophagosomes, and endo/lysosomes; however, clathrin-coated vesicle (CCV) accumulation was not observed (Figures 1F and S4A-S4B).

We further performed western blot analysis to assess potential changes in DA-related and synaptic/endocytic protein levels in striatal tissue. SJ1 cKO[DAT] mice showed a significant reduction in DAT and vMAT2, while TH and AADC levels remained unchanged, indicating impaired presynaptic DA transmission (Figures 2A and 2B). Levels of clathrin-associated endocytic proteins—including SJ1, BIN1, endophilin 1, dynamin, and AP2—were unaffected (Figures S5A and S5B).

To assess DA metabolism and homeostasis, we conducted high-performance liquid chromatography (HPLC) on striatal extracts. DA levels were significantly reduced in SJ1 cKO[DAT] mice at both 2–3 and 6–8 months of age (Figure 2C), indicating striatal DA deficiency. Concurrently, elevated levels of DA metabolites DOPAC (not significant) and HVA suggest increased DA degradation, likely resulting from impaired SV recycling, reduced vesicular sequestration, and cytosolic DA buildup (Figure 2C).

No Apparent Motor Coordination Deficits in SJ1 cKO[DAT] Mice

The strong DA terminal pathology and DA deficiency prompted us to examine the motor function of SJ1 cKO[DAT] mice. In the open field test, the 2-3 months old SJ1 cKO[DAT] mice performed equally well compared to control mice, while 6-8 months old SJ1 cKO[DAT] mice showed reduced total distance and center distance, suggesting their locomotion or exploratory

activity is gradually reduced during aging (Figures 2E and 2F). We further performed more specific motor coordination tests: rotarod test and balanced beam test. Surprisingly, unlike SJ1-KI[RQ] mice which showed strong motor deficits[20], we did not observe obvious motor coordination deficit in SJ1 cKO[DAT] mice when tested for rotarod and balance beam at either 2-3 months or 6-8 months old (Figures 2G and 2H). These results indicate either the DA terminal pathology observed is not severe enough to impair basic motor function, or compensatory mechanisms may have mitigated DA deficiency.

Locally Adaptive induced TH-positive Interneurons (iTHINs) in SJ1 cKO[DAT] Striatum

In addition to the TH/DAT-positive dystrophic clusters in DA terminals, we observed a marked increase in TH-positive neurons within the striatum of SJ1 cKO[DAT] mice (Figure 3A). Striatal TH-positive neurons have been reported in rodents[25-27], primates[28,29], and humans[30-32].. Previous studies using TH-EGFP mice identified them as TH-positive interneurons (THINs), typically undetectable by IHC in wild-type (WT) mice due to low TH expression masked by dense DA axons[33]. In SJ1 cKO[DAT] mice, however, the strong TH protein signal led us to define them as induced TH-positive interneurons (iTHINs). These iTHINs appeared in both dorsal and ventral striatum (Figure S6A), reaching ~200 per coronal section (Figure 3B). Their emergence paralleled DA terminal dystrophy—first detectable at 3 weeks, peaking at 2–3 months, and persisting into adulthood and aging (Figure S6B). Given that SJ1 deletion is restricted to midbrain DA neurons, this suggests a robust and sustained local adaptation in the striatum.

In the striatum, 95% of neurons are MSNs. In addition, there are several other striatal interneurons, including neuropeptide Y (NPY), parvalburmin (PV), Calretinin-positive GABA interneurons and choline acetyltransferase (ChAT)-positive cholinergic interneurons[34]. To determine the identity of iTHINs, we performed double staining of TH with specific markers for different neuronal types. As shown in Figure S7A-S7C, iTHINs were not positive for NPY, PV, and ChAT, suggesting they represent a distinct subtype of interneurons. We observed few iTHINs occasionally expressed Calretinin or Darpp32 (Figures S7D and S7E), which is consistent with previous studies and suggest heterogeneity within this population[35,36]. In addition, we don't see obvious changes of other neuronal types we examined in terms of their number, distribution and marker expression levels. Together, these results further support the specificity of iTHINs induction in response to DA terminal pathology.

DA-Like Properties of iTHINs in SJ1 cKO[DAT] Striatum

To investigate whether iTHINs adopt DA-like properties to compensate for DA deficiency, we first assessed their expression of key biosynthetic enzymes. Importantly, in addition to TH, which is the first enzyme to convert L-tyrosine to L-DOPA, we also observed co-expression of the second enzyme AADC, which further converts L-DOPA to DA. All iTHINs were positive for both TH and AADC, suggesting they can potentially produce DA (Figure 3C).

We next examined the expression of DA transporters: vMAT2 and DAT, which are required for vesicular packaging and reuptake of DA, respectively. Both markers colocalized with TH-positive clusters at dystrophic DA terminals (Figures 3D and 3E) but were absent from iTHINs somata—although DAT gene promoter activity was detected in these cells (see Figure 5E).

Given the known heterogeneity of striatal THINs[34,37], we explored whether iTHINs exhibit subregional adaptations linked to specific midbrain DA neuron subtypes. We first examined aldehyde dehydrogenase 1A1 (ALDH1A1), a marker of A9 SNc DA neurons—the population most vulnerable in PD[38]. As expected, ALDH1A1 labeled ventral-tier SNc neurons in both control and SJ1 cKO[DAT] mice (Figure S8A). In the striatum, ALDH1A1-positive fibers aggregated with TH/DAT-positive clusters in the dorsal region of SJ1 cKO[DAT] mice (Figures 3F and S8B), indicating dystrophic changes in A9-derived terminals. Notably, iTHINs in this region also expressed ALDH1A1 protein, whereas those in the ventral striatum did not, suggesting region-specific adaptation (Figures 3G and S8B). We also examined Calbindin, a marker of A10 VTA DA neurons[39]. While Calbindin labeled a subset of TH-positive neurons in the VTA (Figure S8C), it was not expressed in any iTHINs (Figure S8D). Together, these results show that a subset of iTHINs—particularly in the dorsal striatum—exhibit A9-like molecular feature, potentially reflecting a circuit-specific compensatory response to selective nigrostriatal terminal pathology.

A Second Genetic SJ1 cKO Model (SJ1 cKO[TH]) Recapitulates DA Terminal Pathology and iTHINs Adaptation

To validate our findings, we generated an independent cell type-specific KO model by crossing SJ1 flx/flx mice with the TH-IRES-Cre line, obtaining SJ1 cKO[TH] mice (Figure 4A). While this Cre line also targets TH-positive neurons in other regions (e.g., norepinephrine neurons in the locus coeruleus), our analysis focused on DA neurons in the ventral midbrain and striatum.

As shown in Figure 4B, SJ1 was selectively and completely depleted in TH-positive DA neurons of the SNc and VTA. No obvious DA soma loss was observed in the midbrain of 2-3 months old mice. However, DA terminals in the striatum displayed the same hallmark pathology seen in SJ1 cKO[DAT] mice—early-onset (1–2 months) dystrophic clusters positive

for DA markers including TH, AADC, DAT, vMAT2, SV2C, and ALDH1A1 (Figures 4C–4F). These clusters were broadly distributed across the striatum, with ALDH1A1 selectively labeling clusters in the dorsal region (Figure 4G), further confirming both SNc and VTA DA neurons were affected due to complete loss of SJ1. Importantly, iTHINs were again robustly induced in the striatum, paralleling DA terminal pathology. These iTHINs co-expressed TH and AADC, with dorsal striatal subsets also expressing ALDH1A1 (Figures 4G and 4H), indicating their DA-like phenotype.

Together, results from both SJ1 cKO^{DAT} and cKOTH models demonstrate that SJ1 loss in DA neurons induces cell-autonomous synaptic pathology, which in turn triggers a secondary, non-cell-autonomous compensatory response via local iTHINs.

iTHINs Derive from a pre-existing pool of WT THINs and activate DAT gene expression in response to SJ1 loss

To test whether striatal iTHINs in SJ1 cKO models originate from a pre-existing population of THINs, we genetically labeled original THINs by crossing Ai9 reporter mice (Cre-dependent tdTomato expression) with SJ1 cKOTH mice, generating Ai9;SJ1 cKOTH animals (Figure 5A). This enables tdTomato (tdT) expression in all TH gene expressing cells, including DA neurons in the midbrain and THINs in the striatum. As expected, tdT-expressing neurons in the midbrain colocalized with TH immunostaining (Figure 5B). In the striatum, we observed numerous tdT-expressing neurons in both Ai9;TH-Cre (control) and Ai9;SJ1 cKOTH mice, representing the total THINs (Figure 5C). Consistent with prior studies[27], these tdT-expressing striatal neurons in control mice were not positive for anti-TH staining, indicating that they lacked TH protein (Figure 5C). In contrast, all iTHINs (TH-positive) in SJ1 cKOTH mice colocalized with tdT, comprising approximately 30% of total tdT-expressing neurons (Figures 5C and 5D). This supports the idea that a subset of WT THINs were induced to express DA-related proteins following SJ1 loss.

We further examined the Cre-dependent tdT labeling pattern in Ai9;SJ1 cKO^{DAT} mice with its respective control, Ai9;DAT-Cre (Figure 5A). As expected, DAT-Cre drove tdT labeling in midbrain DA soma and striatal axons in both control and mutant mice (Figures 5E and 5F). In control striatum (Ai9;DAT-Cre), only DA axons were labeled, confirming that WT THINs do not normally express DAT gene. However, in Ai9;SJ1 cKO^{DAT} mice, we unexpectedly observed tdT-expressing cell bodies in the striatum (Figure 5F). Importantly, almost all tdT-expressing neurons were TH-positive, indicating that DAT gene promoter is activated in iTHINs (Figure 5G). Despite this, DAT protein was not detected, possibly due to post-transcriptional

suppression. These findings reinforce the DA-like identity of iTHINs and reveal their
transcriptional adaptability to DA neuron dysfunction.

iTHINs Establish Novel Connectivity with DA Terminals in SJ1 cKO Striatum

To further investigate potential interactions between DA neurons and iTHINs, we employed
an AAV-based strategy to locally label DA neurons in the ventral midbrain using a Cre-
dependent system. We unilaterally injected AAV5-flex-tdTomato into the ventral midbrain of
SJ1 cKOTH, SJ1 cKO^{DAT} and respective Cre line controls, TH-Cre and DAT-Cre (Figure 6A).
Four weeks post-injection, we performed TH immunostaining on midbrain and striatum
sections to visualize DA neurons and their projections.

In both SJ1 cKO and control mice, tdT expression was detected in DA soma (midbrain) and
DA fibers (striatum), confirming successful AAV-mediated expression and diffusion throughout
DA neurons (Figures 6B-6C). Surprisingly, in SJ1 cKOTH and SJ1 cKO^{DAT} mice, we observed
a subset of iTHINs (anti-TH positive) also strongly expressed tdT in the striatum (Figure 6C).
Particularly in SJ1 cKOTH mice, despite TH-Cre driving recombination in all THINs, tdT
expression was exclusively found in iTHINs (Figures 6C and 6D). In contrast, when AAV5-
flex-tdTomato was directly injected into the striatum of SJ1 cKOTH mice, both iTHINs and
original WT THINs (anti-TH negative) were randomly labeled (Figures S9A and S9B 2nd row).
As in other controls, striatal AAV injection labeled WT THINs in TH-Cre but not DAT-Cre mice,
and labeled iTHINs specifically in SJ1 cKO^{DAT} mice (Figures S9B). These results suggest the
presence of a newly formed and specific connection between DA neurons and iTHINs, which
may allow long-distance material transfer, such as AAV axonal transport and inter-neuronal
spreading.

To further trace the connectivity between DA neurons and iTHINs, we used a Cre-independent
AAV strategy by injecting AAV9-TH-EGFP into the midbrain of B6 control and SJ1 cKO mice
(Figure 6E). In both SJ1 cKOTH and SJ1 cKO^{DAT} mice, EGFP expression was observed in TH-
positive DA soma within the SNc and VTA, as well as in their DA fibers projecting to the
striatum (Figures 6F and 6G). Similar to AAV5-flex-tdTomato, only a subset of local TH-
positive iTHINs expressed EGFP in the SJ1 cKO striatum but not B6 control (Figure 6G and
6H). To confirm the selectivity of this connection, we repeated the AAV9-TH-EGFP midbrain
labeling in Ai9;SJ1 cKOTH mice, in which all striatal THINs expressed tdT (Figure S9C). As
shown in Figure S9D, EGFP expression was exclusively detected in tdT/TH-positive double
positive iTHINs but absent in tdT-positive/TH-negative original WT THINs, reinforcing the
notion of selective connectivity between DA neurons and iTHINs.

As a third approach, we injected AAV-hSyn-mCherry into the ventral midbrain of SJ1 cKO
mouse models (Figure 6I). As expected, the pan-neuronal hSyn promoter drove mCherry
expression in different neurons near the injection site, including a subset of TH-positive DA
neurons in the midbrain of both SJ1 cKOTH and SJ1 cKO^{DAT} (Figure 6J). Interestingly, in the
striatum of SJ1 cKO—but not WT mice, we observed mCherry-expressing neurons that were
exclusively iTHINs (anti-TH positive), suggesting specific connectivity and material transfer
(Figures 6K and 6L). In contrast, direct injection of AAV-hSyn-mCherry into WT striatum
resulted in random labeling of various local neurons, including Darpp32-positive MSNs and
other neuronal types (Figures S9E and S9F). This result further ruled out the possibility of
non-specific AAV leakage from dystrophic DA terminals in SJ1 cKO striatum.

Together, these three independent approaches provide compelling evidence that DA neurons
and iTHINs establish a selective and novel connection following SJ1 loss in the DA system.

Upregulation of DA markers in iTHINs distinguishes SJ1 cKO from acute DA lesion mouse models

Previous studies using neurotoxin-induced PD rodent and non-human primate models have
shown the increase of THINs in the striatum following DA depletion[40-43]. To further characterize
the iTHINs observed in our genetic SJ1 cKO models, we directly compared them with two
acute DA lesion models: (1) the 6-OHDA model and (2) a caspase-3 (Casp3)-mediated
ablation model. In the 6-OHDA paradigm, B6 mice received a unilateral injection of the
neurotoxin into the midbrain, and brains were collected for IHC analysis two weeks post-
injection (Figure 7A). For the Casp3 lesion model, a Cre-dependent AAV-taCasp3-TEVp was
unilaterally injected into the midbrain of DAT-Cre mice, and brain tissue analyzed one-month
post-injection (Figure 7C). In both models, we observed clear evidence of DA neuron
degeneration in the midbrain and a near-complete loss of DA projections in the striatum
(Figures 7B and 7D). Following the depletion of DA fibers, a population of local TH-positive
neurons emerged in the striatum (Figure 7E), consistent with previously reported "THINs"[33] .
However, compared to the striatum of SJ1 cKO mice, 1) the TH protein expression level is
much weaker in these neurons (Figures 7E and 7F); 2) they did not express other DA-related
markers, including AADC and ALDH1A1 (Figure 7G and 7H); 3) they did not exhibit
endogenous DAT promoter activity (under the Ai9;DAT-Cre background, Figure S10A),
indicating that iTHINs in SJ1 DA cKO striatum acquire more DA-like features. In Table S1, we
summarized the differences between iTHINs in SJ1 DA cKO mice and WT THINs from these
two neurotoxin models.

In summary, various labeling strategies revealed that about 30% of the original WT THINs were specifically induced to express DA-like features (TH/AADC protein and DAT promoter activity) following DA-specific SJ1 cKO. In these iTHINs, a subset in the dorsal striatum also expressed the A9 DA-specific marker ALDH1A1, matching their nigrostriatal DA input from the midbrain. Furthermore, iTHINs appear to form new and specific connections with dystrophic DA terminals, further supporting a role in adaptive compensation for DA deficiency (Figure S10B).

Acute SJ1 KO in Adulthood Triggers Similar DA Terminal Pathology with limited Developmental Compensation from iTHINs

As described above, iTHINs in genetic SJ1 cKO mice begin to appear at 2-3 weeks of age, peaking at 2-3 months, suggesting that this is an adaptation during early development stage. To bypass this developmental adaption, we generated acute SJ1 KO mice in DA neurons by unilaterally injecting AAV-TH-Cre virus into the ventral midbrain of adult (2-month) or aged (12-month) SJ1 flx/flx mice (Figure 8A). To assess the potential non-specific toxic effect of AAV-Cre expression in DA neurons[44,45], we performed parallel injections in WT mice and found no loss of DA somata or fibers (Figures S11A and S11B).

Following AAV injection, we confirmed successful acute KO of SJ1 by examining brain tissue sections 3-4 months post injection, observing efficient depletion of SJ1 in most TH-positive DA neurons on the ipsilateral midbrain (Figure 8B). Immunostaining results revealed numerous TH/DAT-positive clusters throughout the striatum, resembling the DA terminal pathology observed in genetic SJ1 cKO mice (Figures 8C and 8D). This confirms the synaptic defects across the entire DA population projecting to the striatum due to the complete loss of SJ1. Importantly, even though the number and density of DA terminal clusters in the striatum were comparable to those observed in genetic SJ1 cKO mice, very few TH-positive iTHINs were detected---about 10% of the number observed in genetic SJ1 cKO mice (Figures 8E and 8F). While the DA pathology resulting from SJ1 acute KO was comparable in adult and aged mice, our findings suggest that the secondary compensatory response from iTHINs is primarily induced during early development to compensate for DA terminal defects. Future longitudinal studies are needed to determine whether acute SJ1 loss leads to progressive DA neurodegeneration and motor deficits via axonal die-back.

Discussion

In this study, we examined the consequences of SJ1 loss in DA neurons using newly generated DA-specific SJ1 cKO mice. We identified two major cellular outcomes: (1) a primary

defect marked by severe dystrophy of DA terminals, with accumulation of DA markers and

excess membranes in both dorsal and ventral striatum; and (2) a localized compensatory

response involving a robust increase in DA-like striatal TH-positive interneurons (iTHINs) that

communicate with DA terminals. Comparative analysis with acute adult SJ1 deletion and other

lesion models indicates that this iTHINs induction is specific to early developmental loss of

SJ1. This adaptive striatal plasticity may serve as an early neuroprotective mechanism to

maintain DA homeostasis or delay terminal degeneration, offering a potential therapeutic

strategy to preserve DA function in PD.

Cell-Autonomous, Clathrin-Independent Role of SJ1 in the Neuromodulatory System

SJ1 has been extensively studied in excitatory and inhibitory synapses, where it plays a key

role in clathrin uncoating during clathrin-mediated SV endocytosis and recycling. Constitutive

whole-body SJ1 KO mice die postnatally with severe epilepsy, suggesting imbalanced

excitatory/inhibitory transmission during development[16]. While DA-specific SJ1 cKO mice

survive to adulthood without seizures, our findings show that DA terminals from both the SNc

and VTA undergo significant dystrophic changes, indicating a cell-autonomous role for SJ1 in

regulating DA synaptic function. Surprisingly, no accumulation of endocytic proteins involved

in CME or CCVs were detected in these dystrophic terminals. As neuromodulators, DA

neurons release transmitters via "volume transmission," a slower process that does not rely

on precise synaptic contact sites[46]. This contrasts with the fast "synaptic transmission" seen

in excitatory and inhibitory synapses, which primarily recycle SVs through CME. Given these

differences, it is plausible that DA synapses utilize alternative mechanisms for SV recycling,

such as bulk endocytosis [47].

In addition to DA, another neuromodulator norepinephrine neurons are also vulnerable in PD,

likely due to shared features such as pace-making firing, extensive axonal arborization, and

accumulation of cytosolic oxidized catecholamines. Future studies using our SJ1 conditional

KO model could determine whether SJ1 similarly regulates SV recycling in norepinephrine

neurons and how its dysfunction contributes to their selective degeneration in PD.

Gene Dosage-Dependent SJ1 Loss Correlates with the Severity of DA pathology

There is a clear genotype-phenotype correlation in SJ1-linked PD cases, which is mirrored in

mouse models carrying different SJ1 mutations. DA pathology in these models also reflects

SJ1 gene dosage. The SJ1-KI[RQ] model shows selective degeneration of nigrostriatal terminals,

while the SJ1 cKO model exhibits more widespread DA terminal dystrophy. This indicates that

both A9 and A10 DA neurons are susceptible to impaired SJ1-mediated SV recycling, with A9

SNc neurons being particularly vulnerable, even with partial SJ1 loss. A similar synergistic effect is seen in the Auxilin KO/SJ1-KI[RQ] double mutant, which displays more severe DA terminal pathology than either single mutant[21]. More recently, SJ1 haploinsufficiency has been linked to abnormal mesolimbic signaling and mild motivational deficits[48], suggesting a broader role for SJ1 in VTA DA function that can be further explored using our SJ1 DA cKO mice. Together, these findings highlight impaired SV recycling as a key driver of DA dysfunction, with partial impairment leading to selective degeneration of SNc DA terminals as a hallmark of PD pathogenesis.

Synaptic Transmission Defects in DA Terminals Are Insufficient to Trigger Motor Dysfunction

SJ1 DA-specific cKO mice do not exhibit obvious motor deficits, despite showing more severe terminal pathology and DA deficiency than SJ1-KI[RQ] mice. This surprising finding suggests several possibilities: (1) motor symptoms in SJ1-KI[RQ] mice may arise from combined SJ1 dysfunction across multiple brain regions, not just DA neurons; (2) the dystrophic DA terminal changes observed in SJ1 mutant mice may represent a prodromal phase of PD that does not progress to full clinical manifestations within the mouse lifespan; and (3) local compensatory mechanisms within the striatum may preserve basic motor function despite significant synaptic dysfunction.

Our findings are consistent with recent studies using DA-specific cKO mouse models of two important synaptic transmission proteins: the Ca^{2+} sensor Syt1 and the release site organizer RIM1/2[49,50]. In both cKO models, although activity- or action potential-induced phasic DA release is significantly abolished, the mice do not exhibit obvious motor defects[49,50]. Furthermore, early compensatory changes, such as receptor adaptation and enhanced DA synthesis, packaging, and innervation, have been observed in the striatum of Syt1 DA-cKO mice[49]. These results suggest that DA-dependent motor functions can be sustained by basal extracellular DA levels and striatal compensatory plasticity.

Taken together, the converging results from three DA-specific cKO models with impaired synaptic transmission support the idea that motor symptoms of PD only emerge after extensive DA denervation. These models may therefore capture key aspects of the pre-symptomatic or early compensatory stages of PD.

Adaptive Compensation of iTHINs in SJ1 DA cKO Striatum

DA-like Features of iTHINs

A prominent adaptive change in SJ1 DA cKO mice is a robust increase in iTHINs, accounting

for ~30% of the original total THINs population in the striatum. A much milder induction of

iTHINs was also observed in SJ1-KI[RQ] and Auxilin KO/SJ1-KI[RQ] double mutants, suggesting

that this response is specific to endocytic defects and modulated by gene dosage[21].

Neurotoxin-induced PD models (6-OHDA, MPTP) in rodents and non-human primates have

shown increased THINs in the striatum following DA depletion, suggesting a potential

compensatory or protective role[40-43]. However, it remains debatable whether these THINs

qualify as bona fide DA neurons capable of expressing key DA markers (e.g., AADC, DAT)

and producing/releasing DA[33,43,51].

In our study, when compared with acute 6-OHDA and Casp3 lesion models, iTHINs in SJ1

cKO mice displayed distinct DA-like features: strong co-expression of TH and AADC,

activation of the DAT gene promoter, and expression of the SNc DA neuron marker ALDH1A1

in a subset of dorsal striatal iTHINS. While SV-related markers like vMAT2 were not detected,

it's worth nothing that non-vesicle mediated somato-dendritic DA release in the midbrain has

been proposed previously[52]. Future biochemical and physiological studies are needed to

determine iTHINs capacity for DA production and release. Alternatively, iTHINs may retain

their GABAergic interneuron identity and contribute to striatal homeostasis by modulating local

circuitry[33].

Crosstalk Between iTHINs and Dystrophic DA Terminals in SJ1 cKO Striatum

Our observations further support long-distance communication between DA neurons and

iTHINs. First, a subset of iTHINs expressing A9 SNc DA marker ALDH1A1, are enriched in

the dorsal striatum, the primary target of ALDH1A1-positive DA fibers. This indicates a region-

specific and circuit-dependent compensatory response to DA terminal degeneration. Second,

AAV-based tracing from SJ1 KO DA neurons leads to selective co-labeling of striatal iTHINs,

suggesting material transfer or spreading from degenerating DA terminals to iTHINs. The

underlying mechanisms such as through newly formed synapses or non-synaptic mechanisms

such as exosomes or tunneling nanotubes should be explored in future studies[53,54].

What signal do striatal THINs detect to trigger their conversion? While extracellular DA levels

were previously shown to negatively regulate THINs number[55], acute DA lesion models with

severe DA depletion do not induce iTHINs. This indicates that other factors—likely released

from DA fibers—are required to initiate and sustain DA-like features in iTHINs. Notably, acute

SJ1 loss in adult or aged DA neurons significantly reduced the number of iTHINs, suggesting

this signal diminishes with age. Identifying this signal could open new therapeutic avenues for circuit-level compensation in PD.

Furthermore, prior studies suggest THINs form connections with other striatal neurons[27,34], raising the possibility that iTHINs may also engage in local crosstalk—an area that can be further explored using single-nucleus RNA sequencing and spatial transcriptomics.

Methods and Materials

Animals

Mice were housed in SPF rooms with a 12-hour light/dark cycle. All experiments involving animals were conducted according to the protocol approved by SingHealth Institutional Animal Care and Use Committee (IACUC).

SJ1-KI[RQ] mouse line was generated as described before and a gift from Prof Pietro De Camili (Yale University)[20]. SJ1 flx/flx (C57BL/6J-Synj1[em1cyagen]; S-CKO-00357, Cyagen) was generated by inserting loxP sites flanking exon 4 of Synj1 using CRISPR/Cas9. These mice were crossed with DAT-IRES-Cre (C001024, Cyagen) and TH-IRES-Cre knock in line (generated from Prof Luis de Lecea Lab, Standford University; kindly provided by Dr Sarah Luo, IMCB-A*STAR) to obtain double heterozygous, +/DAT-Cre;SJ1 +/flx and +/TH-Cre;SJ1 +/flx mice, respectively. To create DA-specific SJ1 conditional KO mice, double heterozygous mice were crossed with homozygous SJ1 flx/flx, obtaining +/DAT-Cre;SJ1 flx/flx and +/TH-Cre; SJ1 flx/fox mice, referred to as SJ1 cKO[DAT] and SJ1 cKO[TH], respectively. Appropriate littermate or age-matched controls (SJ1 flx/flx or +/DAT-Cre;SJ1 +/flx or +/TH-Cre;SJ1 +/flx) were used in all experiments. All mice used in the experiments with Cre-dependant promoter were heterozygous, simply referred as DAT-Cre and TH-Cre in the manuscript.

Ai9 mice (B6.Cg-Gt(ROSA)26Sortm9(CAG-tdTomato)Hze/J; RRID:IMSR_JAX:007909) were kindly provided by Prof Enrico Petretto (Duke-NUS) for specific labeling following Cre-mediated recombination. Homozygous Ai9 mice were first crossed with homozygous SJ1 flx/flx to obtain double heterozygous, +/Ai9;SJ1 +/flx mice. Subsequent models involving Ai9 were all heterozygous Ai9, hence simply referred as Ai9 in the manuscript. Ai9;SJ1 +/flx mice were then crossed with SJ1 cKO[DAT] and SJ1 cKO[TH] to obtain Ai9;SJ1 cKO[DAT] and Ai9;SJ1 cKO[TH], respectively. Appropriate littermate or age-matched controls were used in all experiments.

For SJ1 acute KO mice model, SJ1 flx/flx mice were employed. Briefly, mice were stereotactic injected AAV.rTH.PI.Cre.SV40 (1ul) (Addgene-#107788) unilaterally in right midbrain (AP 3.3

mm, ML 1.2 mm, DV 4.3 mm) after anesthetized by isoflurane (5% induction, 2% maintenance at 0.5-0.8 L/min oxygen or air flow). After 3 to 4 month of injection, mice brain tissues were obtained for IHC.

For 6-hydroxydopamine (6-OHDA; H4381, Sigma-Aldrich) lesion model, B6 mice (C57BL/6J; RRID:IMSR_JAX:000664) were treated with desipramine (D3900, Sigma-Aldrich; i.p. 25 mg/kg) 30 min prior to 6-OHDA infusions to prevent noradrenergic neurons. 6-OHDA (2.88 ug, dissolved in 0.2% ascorbate/saline solution) was unilaterally pressure injected into the right midbrain (AP 3.3 mm, ML 1.2 mm, DV 4.3 mm). Mice brain tissues were harvested for IHC after 2 weeks of 6-OHDA injection.

For Casp3 ablation model, Cre-dependent virus, pAAV-flex-taCasp3-TEVp (1ul) (Addgene-#45580) were unilaterally injected in Cre heterozygous mice right midbrain (AP 3.3 mm, ML 1.2 mm, DV 4.3 mm) after anesthetizing. Mice brain tissues were harvested for IHC after one month of virus injection.

AAV Labelling Surgery

Animals were anesthetized using isoflurane (5% induction, 2% maintenance at 0 .5-0.8 L/min oxygen or air flow) and secured in a stereotaxic apparatus. For virus injection, pAAV-FLEX-tdTomato (1ul) (Addgene-#28306), rAAV-mTH-EGFP (BrainCase-#BC-0319) (0.5ul) or pAAV-hSyn-mCherry (Addgene-#114472) (1ul) was infused into the midbrain (targeting the VTA and SNc) and the striatum. The coordinates for the midbrain were AP 3.3 mm, ML 1.2 mm, DV 4.3 mm. For striatal infusion, the coordinates were AP -0.3 mm, ML -2.0 mm, DV -2.5 and -4.0 mm.

Infusions were conducted using a Hamilton syringe (Hamilton-Syringe, NEUROS, 7002, 2UL. The infusion rate for all surgery was 100nl/min, unless stated otherwise. Syringe was left in place for an additional 10 min before withdrawal to prevent retrograde mass flow. All injection procedures were performed on a heating pad (37°C) to maintain body temperature. Following surgery, mice recovered in another clean cage on a heating pad until they were awake.

Antibodies

The antibodies used in this study were obtained from commercial sources as stated: mouse anti-α-synuclein (610786, RRID: AB_2748880), mouse anti-AP2 (611350, RRID: AB_398872) and mouse anti-Dynamin (610245, RRID: AB_397640) from BD Biosciences; goat anti-ChAT

(AB144P, RRID: AB_2079751), mouse anti-clathrin light chain (AB9884, RRID: AB_992745), rat anti-DAT (MAB369, RRID: AB_2190413), mouse anti-SV2C (MABN367, RRID: AB_2905667) and rabbit anti-TH (AB152, RRID: AB_390204) from Merck Millipore; mouse anti-β-actin (sc-47778, RRID: AB_2714189) and mouse anti-BIN1 99D (sc-13575, RRID:AB_626753) from Santa Cruz Biotechnology; rabbit anti-ALDH1A1 (HPA002123, RRID: AB_1844722), rabbit anti-GFAP (ZRB2383, RRID: AB_2905668) and rabbit anti-SJ1 (HPA011916, RRID: AB_1857692) from Sigma-Aldrich; rabbit anti-Endophilin 1 (159002, RID: AB_887757), mouse-anti Calretinin (214111, RRID: AB_2619904), mouse anti-TH (213211, RRID: AB_2636901), rabbit anti-Amphiphysin 1 (120002, RRID: AB_887690), rabbit anti-AADC (369003, RRID: AB_2620131), guinea pig-anti Calbindin D28k (214005, RRID: AB_2619902), mouse anti-Parvalbumin (195011, RRID: AB_2619883) and rabbit anti-D2R (376 203, RRID: AB_2636918) from Synaptic Systems; rabbit anti-VMAT2 (MSFR106400, RRID: AB_2571857) from Nittobo Medical; rabbit anti-DARPP-32 (2306, RRID: AB_823479) and rabbit anti-Neuropeptide Y (11976, RRID: AB_2716286) from Cell Signaling Technology; rabbit anti-Iba1 (019-19741, RRID: AB_839504) from FUJIFILM Wako Pure Chemical Corporation; goat anti-ALDH1A1 (AF5869_RRID: AB_2044597) from R&D Systems; rabbit anti-alpha synuclein phospho S129 (EP1536Y, RRID:AB_765074) from abcam; mouse anti-GAPDH (60004-1, RRID: AB_2107436) and rabbit anti-GAPDH (10494-1-AP, RRID: AB_2263076) from Proteintech.

Secondary antibodies used were all purchased from commercial sources as stated: donkey anti-mouse IgG (H + L) Alexa Fluor Plus 488 (A32766, RRID: AB_2762823), donkey anti-mouse IgG (H + L) Alexa Fluor 594 (A21203, RRID: AB_2535789), donkey anti-mouse IgG (H + L) Alexa Fluor Plus 647 (A32787, RRID: AB_2762830), donkey anti-rabbit IgG (H + L) Alexa Fluor Plus 488 (A32790, RRID: AB_2762833), donkey anti-rabbit IgG (H + L) Alexa Fluor Plus 594 (A32754, RRID: AB_2762827), donkey anti-rabbit IgG (H + L) Alexa Fluor 647 (A32795, RRID: AB_2762835), donkey anti-rat IgG (H + L) Alexa Fluor Plus 488 (A48269, RRID: AB_2893137), donkey anti-rat IgG (H + L) Alexa Fluor 594 (A21209, RRID: AB_2535795), donkey anti-goat IgG (H + L) Alexa Fluor 488 (A11055, RRID: AB_2534102), donkey anti-goat IgG (H + L) Alexa Fluor 594 (A32758, RRID: AB_2762828), goat anti-guinea pig IgG (H + L) Alexa Fluor 488 (A11073, RRID: AB_2534117) and goat anti-guinea pig IgG (H + L) Alexa Fluor 647 (A21450, RRID: AB_2535867) from Life Technologies; goat anti-mouse IgG light chain HRP (AP200P, RRID:AB_805324) and mouse anti-rabbit IgG light chain HRP (MAB201P, RRID:AB_827270) from Merck; IRDye 800CW donkey anti-rabbit IgG (926-32213, RRID: AB_621848), IRDye 800CW donkey anti-mouse IgG (926-32212, RRID: AB_621847), IRDye 680RD donkey anti-mouse IgG (926-68072, RRID: AB_10953628) and IRDye 800CW goat anti-rat (926-32219, RRID: AB_1850025) from LI-COR Biosciences.

Brain Histology and Immunofluorescence

Mice were anesthetized with a Ketamin/Xylazine anesthetic cocktail injection, perfused transcardially with ice-cold 4% paraformaldehyde in 1 X phosphate-buffered saline (PBS) and the brains were kept in the same fixative overnight at 4°C. On the next day, brains were transferred to 30% sucrose in 1 X PBS and kept overnight at 4°C on top of a roller. Brains were then embedded in OCT (Tissue-Tek) and freeze in liquid nitrogen-cooled 2-methylbutane (Isopentane). Coronal or sagittal sections of 20 μm thickness were cut with a cryostat. The sections were either mounted on adhesive slide, SuperFrost® Plus (VWR) or collected as floating sections in 1 X PBS in a 24-well plate. Sections were then blocked in 5% BSA and 0.1% Triton X-100 in 1 X PBS for 1 hour at room temperature. After blocking, sections were incubated with primary antibodies and kept overnight at 4°C. Subsequently, sections were washed 3 times for 10 min with 0.1% Triton X-100 in 1 X PBS, then incubated with Alexa-conjugated secondary antibodies for 1 hour at room temperature. Finally, the sections were mounted with Fluoromount-G (Invitrogen) and sealed with nail polish. The floating sections were mounted onto slides after all the staining procedures. Images were acquired with Andor BC43 (Oxford Instruments) with 10x, 20x and 40x oil objective or a spinning disk system (Gataca Systems) based on an inverted microscope (Nikon Ti2-E; Nikon) equipped with a confocal spinning head (CSU-W, Yokogawa) and a Plan-Apo 20x and 40x oil objective.

For stereological analysis of TH-positive DA neurons in the midbrain, 30 μm coronal section were collected incubated with 0.1% H_2O_2 for 5 min to quench endogenous peroxidase activity and subsequently incubated with buffer containing 5% BSA, and 0.1% Triton X-100 in 1X PBS for 1 hour at room temperature. Next, sections were incubated overnight at 4°C with anti-TH primary antibody (1:1000). The sections were then washed with 1X PBS three times for 10 min and incubated with a biotinylated anti-rabbit secondary antibody (Vector Laboratories, USA) for 1 hour at room temperature, later followed by incubation with Avidin/Biotin complex (ABC) reagent (Vector Laboratories, USA) at room temperature. Finally, immunoreactivity was revealed by incubation with diaminobenzidine (DAB). Stereological analysis was performed using the optical fractionator probe in the Stereo Investigator (MBF Bioscience) software. A total of 7-9 coronal sections of every fourth sections were collected across the midbrain and used for counting. The SN and VTA regions were outlined based on the Allen mouse brain atlas and counting was performed using 63x oil lens. The parameters used were guard zone of 2 μm, dissector height of 13 μm and counting frame of 200 μm x 200 μm.

Electron Microscopy

2 months old mouse was anesthetized and fixed by transcardial perfusion with 4%
paraformaldehyde and 0.125% glutaraldehyde (EMS, Hatfield, PA) in 0.1 M PB buffer. Brain
was removed and dissected in small pieces (1 × 1 × 1 mm^3) and further incubated in 2.5%
glutaraldehyde in 0.1 M PB buffer at 4 °C overnight. The samples were then washed for 5 mins
in 1 X PBS for three times. The samples were post-fixed with 1% osmium tetroxide for 1 hour
and washed with distilled water. They are then dehydrated through an ethanol series (25%,
50%, 75%, 95% and 100%) and absolute acetone, before infiltrating with araldite
resin/acetone mixtures. On the next day, samples were further infiltrated in pure resin at 40°C
to 50°C, then embedded in resin and polymerised at 60°C for 24 hours. The embedded sample
was sectioned into 100nm slices with a Leica UC6 ultramicrotome and stained with lead citrate.
Images were captured with JEOL JEM-1400Flash TEM, operating at 100kV.

Motor Behavioral Tests

613 SJ1 cKO^{DAT} and mice between 2-8 months old were used for balance beam and rotarod tests.
614 The mice were divided into two age groups: 2-3 months old (n= 13 for Ctrl, n=15 for SJ1
615 cKO^{DAT}) and 6-8 months old (n=9 for Ctrl and n=10 for SJ1 cKO^{DAT}). Both males and females
616 were used for the behavioral assays. Both tests were conducted during the light period. The
617 mice were allowed to be acclimatized to the test for at least 30 min before each test.

Balance Beam Test

619 A narrow beam (10 mm width) was suspended 15 cm above a soft padding. Mice were placed
620 on end of the beam and during their trip towards the other end, the number of missteps (paw
621 slips) was recorded. The test was conducted thrice for each mouse and the average number
622 of missteps was calculated.

Rotarod Test

624 Mice were placed on a rod which is rotating at 4 rpm with an acceleration to 40 rpm within 6
625 min. The duration of the mice staying on the rod is measured three times at 10 min intervals
626 and the average 'latency to fall' was determined for each mouse.

Open Field Test

628 Mice were placed in a chamber measured 40 cm (length) x 40 cm (width) x 30 cm (height)
629 and recorded for 15-minute. Recorded videos were analyzed with TopScan video tracking
630 software. The mice path tracking and time spent in the defined area were analyzed. The mice
631 used for two age groups: 2-3 months old (n= 12 for Ctrl, n= 14 for SJ1 cKO^{DAT}) and 6-8 months
632 old (n= 6 for Ctrl and n= 7 for SJ1 cKO^{DAT}).

HPLC

Striatum tissues dissected from mice brain were homogenized in 0.5 N perchloric acid with 100 µM of deferoxamine mesylate and 100 µM of glutathione. The homogenized samples were further sonicated, centrifuged and the supernatants were filtered using 0.1 µm PVDF centrifugal filters before collecting the filtrates for HPLC analysis. A reversed-phase UltiMate 3000 HPLC system (Thermo Fisher Scientific) with an electrochemical detector and a reversed-phase column (Vydac Denali, C18, 4.6 x 250 mm, 5 µm particle size, 120 Å pore size) were used to run the samples. The HPLC run was performed at a flow rate of 0.5 ml per minute with a mobile phase containing 1.3% sodium acetate, 0.01% EDTA (pH8.0), 0.5% sodium 1-heptanesulfonate, 7% acetonitrile (v/v) and 2% methanol (v/v), adjusted to pH 4.0 with acetic acid. All buffers used for HPLC analysis were double filtered through 0.2 µM nylon membranes. Dopamine, DOPAC and HVA levels in the samples was identified by retention time of their respective standard and quantified by measuring the area under the peak using the software ChromeleonTM 7.2 Chromatography Data System (Thermo Fisher Scientific). The areas under the peaks were normalized to their respective tissue weight.

Immunoblotting

Striatum tissues were homogenized in buffer containing 20 mM Tris-HCl (pH 7.4), 150 mM NaCl, 2 mM EDTA and cOmpleteTM EDTA-free Protease Inhibitor Cocktail (Roche, USA). The homogenized samples were centrifuged at 700 g for 10 min to obtain the post-nuclear supernatant (PNS). Protein concentration was determined by the Pierce BCA Protein Assay Kit. SDS-PAGE and western blotting were performed by standard procedures. The membranes were blotted with primary antibodies and then IRDye® Secondary Antibodies (LI-COR, USA) or HRP light chain secondary antibodies. The protein bands were detected using Odyssey ® CLx imaging system or Bio-Rad ChemiDoc Touch Imaging System, and its intensity was quantified via Image Studio. A minimum of three independent sets of samples were used for quantification.

Quantification of DAT Immunoreactivity Clusters and TH+ Striatal Interneurons

The quantification of DAT-positive clustering in striatum was conducted using Fiji software (Version ImageJ 2.160/1.54p/Java 1.8.0_322 (64-bit), RRID: SCR_002285). For each mouse, 5 random sampling sites with a size of 300 µm x 300 µm were selected respectively for both

dorsal and ventral striatum. At least three mice from each genotype and each age group were collected for this purpose. The threshold setting for the clusters was set to quantify puncta bigger than 5 μm^2 to ensure normal positive axonal terminals were not counted in. The average size of clusters and average number of clusters was counted automatically with Fiji plugin "Analyse Particles" based on all five random sampling sites for each mouse.

Quantification of TH-positive and tdT-expressing cells was done using Fiji software with full stitched half coronal section. Image acquisition involved tiling of the entire section with either 10x or 20x objective lens on Andor BC43. Cell Counter plugin in Fiji software was used to count visible cell bodies across the striatum (outlined based on Allen mouse brain atlas). Cell count from individual channels was used for percentage ratio quantification.

Quantification for TH-positive cells intensity was done using Fiji software. A consistent threshold was applied across all images to isolate TH-positive signals. The 8-bit figure type was used to measure the mean density of the TH signal. Background fluorescence was determined from adjacent areas without specific staining. All measurements were performed under identical settings to ensure consistency across experimental groups. For each mouse, 6 random TH-positive cells were selected from both dorsal and ventral striatum. At least three mice from each condition were collected for this purpose.

Statistical Analysis

All statistical analysis was performed using GraphPad Prism (Ver 10.4.1, RRID: SCR_002798). All graphs were plotted on GraphPad Prism. Data are presented as mean ± SEM unless otherwise stated. Statistical significance was determined using the Student's unpaired t test for the comparison of two independent groups or ANOVA with Tukey's Honest Significant Difference post hoc test for multiple group comparisons. Data with p-values <0.05, <0.01, <0.001, and <0.0001 are represented by asterisks *, **, ***, and ****.

Acknowledgment

We thank Prof Zhang Suchun's lab (Duke-NUS) and Asst Prof Jun Nishiyama's lab (Duke-NUS) for providing some reagents for the experiments. We thank Prof Pietro De Camili (Yale University), Prof Shirish Shenolikar (Duke University) and Prof Wang Hongyan (Duke-NUS) for reading the manuscript and providing comments. We thank Dr Sarah Luo Xinwei (IMCB-A*STAR) and Prof Enrico Petretto (Duke-NUS) for kindly providing TH-IRES-Cre mouse line and Ai9 mouse line, respectively. We also thank Prof Pietro De Camili for gifting SJ1-KI^{RQ}

mouse line. We thank Asst Prof Wong Peiyan (Duke-NUS) from animal behaviour core for
training. We thank Electron Microscopy Unit (NUS) for EM preparation and imaging. Select
portions of this manuscript were edited for clarity and grammar using the ChatGPT language
model (OpenAI). Scientific content and interpretations were solely formulated by the authors.
This research was supported in part by grant from Ministry of Education Academic Research
Fund (AcRF) Tier 2 Singapore (MOE-T2EP30122-0009) and Tier 1 (FY2024-MOET1-0001).

Declaration of interest

The authors declare no competing interests.

Author Contributions

B.M., Y.L., M.L., Z.L. and M.C. designed the experiments. Y.L., M.L. and Z.L. performed
sample preparations, histology experiments and microscopy imaging. Y.L. and X.C performed
HPLC and analysis. Y.L. preformed western blotting and analysis. Y.L. and M.C. performed
sample preparations for EM. B.M., Y.L., M.L. and X.C. performed behavioural experiments
and analysis. B.M., M.L. and Z.L. performed stereotaxic injections. Y.L. and M.L. performed
quantification and statistical analysis. M.L. and X.C. performed mice genotyping. B.M., Y.L.,
M.L., Z.L. and M.C. wrote the paper. All authors read and approved the final manuscript.

Availability of data and materials

The datasets generated and/or analyzed in this study are available from the corresponding
author Mian Cao on request.

Figure Legends

Figure 1. SJ1 conditional KO (SJ1 cKO^{DAT}) in DA neurons exhibit dystrophic DA axon terminals in the striatum

(A) Generation of SJ1 cKO^{DAT} mice by crossing SJ1 flx/flx mice with DAT-IRES-Cre line to create DA-specific SJ1 conditional KO mice (DAT-Cre;SJ1 flx/flx), hereafter referred to as SJ1 cKO^{DAT}. Appropriate littermate control (SJ1 flx/flx or DATCre;SJ1+/flx) were used in all experiments.

(B) Double staining of TH and SJ1 to show SJ1 specifically KO in midbrain DA neurons. Green arrows: TH-positive DA neurons; White arrows: TH-negative midbrain neurons. SJ1 expression in hippocampus shows that other regions were not affected. Scale bar: 20 μm.

(C) DAT staining of 2 months old Ctrl and SJ1 cKO^{DAT} shows clustering phenotype in dorsal (DS) and ventral striatum (VS). Scale bar: 20 μm.

(D) Quantification for DAT clusters at 2-3 months and 6-8 months of age. The average number of clusters is counted in five randomly selected, 300 × 300-μm regions of interest (ROIs). Data are represented as mean ± SEM (**p < 0.01, and ***p < 0.001, and ****p < 0.0001 by one-way ANOVA with post-hoc Tukey's test). n=3 mice per genotype.

(E) Double staining of DAT with TH, AADC, vMAT2 and SV2C showed colocalization in the dorsal striatum (DS) of SJ1 cKO^{DAT} mice. Scale bar: 20 μm.

(F) Representative EM micrographs of nerve terminals from the ventral striatum (VS) of SJ1 cKO^{DAT}. Red arrows: "Whirl"-like multilayered membranes; Synaptic vesicles (SVs), endosomes (endo), autophagosome (AP) and mitochondria (mt) were also labeled in the images. Scale bar: 500 nm.

Figure 2. DA deficiency in SJ1 cKO^{DAT} striatum with no apparent motor deficit

(A) Representative blots for DA markers, including DAT, vMAT2, TH and AADC from 2-3 months old Ctrl and SJ1 cKO^{DAT} half brain striatal homogenates.

(B) Quantification of proteins level shown in (A). Significant reduction of DAT and vMAT2 in SJ1 cKO^{DAT} after protein levels normalized to the level of actin. Data are represented as mean ± SEM (**p < 0.01, and ***p < 0.001 with Student's unpaired t-test). n=3-5 mice per genotype.

(C), (D) HPLC analysis of SJ1 cKO^{DAT} mice showed an average of 50% reduction in striatal DA levels for (C) 2-3 months old and (D) 6-8 months old. Data are represented as mean ± SEM

(*p < 0.05, **p < 0.01 by one-way ANOVA with Student's unpaired t-test). n=5 for Ctrl and 2-

3M SJ1 cKO^{DAT}; n=6 for Ctrl and 6-8M SJ1 cKO^{DAT}.

(E) Performance of SJ1 cKO^{DAT} mice in open field test. No significant difference in locomotor

activity at 2-3 months old. Significant p-value for total distance and center distance at 6-8

months old indicate reduced exploratory activity. Data are represented as mean ± SEM

(**p < 0.01 with Student's unpaired t-test). n numbers used for each age group of the Ctrl and

SJ1 cKO^{DAT} mice (2-3M: 12 and 14; 6-8M: 6 and 7).

(F) Representative tracing of Ctrl and SJ1 cKO^{DAT} mice at 2-3 months and 6-8 months old

recorded for 15 minutes. The center square box indicates the "center zone".

(G) Performance of two age groups of SJ1 cKO^{DAT} mice on the accelerated rotarod. Data are

represented as mean ± SEM. n numbers used for each age group of the Ctrl and SJ1 cKO^{DAT}

mice (2–3M: 13 and 15; 6–8 M: 9 and 10).

(H) Quantification for the number of missteps during the balance beam test for two age groups

of SJ1 cKO^{DAT} mice. Data are represented as mean ± SEM. n numbers used for each age

group of the Ctrl and SJ1 cKO^{DAT} mice (2–3M: 13 and 15; 6–8M: 9 and 10).

Figure 3. Robust appearance of induced TH-positive interneurons (iTHINs) with DA-like

characteristics in the SJ1 cKO^{DAT} striatum

(A) Numerous induced TH-positive interneurons (iTHINs) observed in the striatum of SJ1

cKO^{DAT} mice but not in Ctrl. Scale bar: 50 µm.

(B) Quantification of iTHINs in 2-3 months and 6-8 months age group. Number of iTHINs were

quantified on single coronal striatum section of one hemisphere. Data are represented as

mean ± SEM (**p < 0.01). n=3 mice per genotype.

(C) Representative image of iTHINs expressing AADC in the striatum of 2 months old SJ1

cKO^{DAT} mice compared to Ctrl. White arrows: iTHINs. Asterisk: clusters. Scale bar: 20 µm.

(D) Double staining of vMAT2 and TH shows vMAT2 is clustered with TH at DA terminals but

not expressed in iTHINs in 2 months old SJ1 cKO^{DAT} striatum. White arrows: iTHINs. Asterisk:

clusters. Scale bar: 20 µm.

(E) Double staining of DAT and TH shows DAT is clustered with TH at DA terminals but not

expressed in iTHINs in 2 months old SJ1 cKO^{DAT} striatum. White arrows: iTHINs. Asterisk:

clusters. Scale bar: 20 µm.

(F), (G) Double staining of ALDH1A1 and TH shows ALDH1A1 is detected in both dystrophic
DA terminals and iTHINs in SJ1 cKO^{DAT} (F) dorsal but not (G) ventral striatum. White arrows:
iTHINs. Asterisk: clusters. Scale bar: 50 μm.

Figure 4. SJ1 conditional KO using TH-IRES-Cre (SJ1 cKOTH) phenocopies SJ1 cKO^{DAT}
pathological characteristics

(A) Generation of SJ1 cKOTH mice by crossing SJ1 flx/flx mice with TH-IRES-Cre line to create
TH-specific SJ1 conditional KO mice (TH-Cre;SJ1 flx/flx), hereafter referred to as SJ1 cKOTH.
Appropriate littermate controls (SJ1 flx/flx or TH-Cre;SJ1 +/flx) were used in all experiments.

(B) Double staining of TH and SJ1 showed specific loss of SJ1 in DA neurons. SJ1 expression
in hippocampus shows that other regions were not affected. Scale bar: 20 μm.

(C) DAT staining of 2 months Ctrl and SJ1 cKOTH showed similar clustering phenotype in
dorsal and ventral striatum. Scale bar: 20 μm.

(D) Double staining of vMAT2 with SV2C showed co-aggregation in the striatum of SJ1 cKOTH
mice. Scale bar: 20 μm.

(E) Double staining of DAT and TH showed DAT is clustered with TH at DA terminals (asterisk),
but not expressed in iTHINs (white arrows) in the striatum of SJ1 cKOTH mice. Scale bar:
20 μm.

(F) Double staining of AADC and TH showed AADC is clustered with TH at DA terminals
(asterisk), and also co-expressed in iTHINs (white arrows). Scale bar: 20 μm.

(G), (H) Double staining of ALDH1A1 and TH shows ALDH1A1 is detected in both dystrophic
DA terminals (asterik) and iTHINs (white arrows) in SJ1 cKO^{DAT} dorsal (G) but not ventral (H)
striatum. Scale bar: 50 μm.

Figure 5. iTHINs emerge from original WT THINs and induce DAT gene expression after
SJ1 loss

(A) Simplified schematic showing the generation of Ai9-tdT labeled SJ1 cKOTH and SJ1 cKO^{DAT}
mice, referred to as Ai9;SJ1 cKOTH and Ai9;SJ1 cKO^{DAT}, respectively. Ai9-tdT reporter driven
by CAG promoter expresses DA-specific tdT signal after Cre-mediated recombination by
either TH- or DAT-IRES-Cre line.

(B) Representative images of tdT expression and immunostaining of TH in the midbrain DA neurons of 2 months old Ai9;TH-Cre (Ctrl) and Ai9;SJ1 cKO^TH. Scale bar: 100 µm.

(C) Representative images of tdT expression and immunostaining of TH in the striatum of 2 months old Ai9;TH-Cre (Ctrl) and Ai9;SJ1 cKO^TH. Blue arrows: tdT expressing THINs driven by TH-Cre recombination (negative for TH immunostaining). White arrows: iTHINs expressing tdT signal (positive for TH immunostaining). Scale bar: 50 µm

(D) Percentage of TH-Cre driven tdT expressing neurons positive for TH immunostaining shown in (C). In average, 34.11% of total tdT expressing neurons were positive for TH immunostaining (iTHINs). The number of neurons was quantified on a single stitched coronal striatum section for both tdT expressing neurons and tdT/TH-positive neurons. Data are represented as mean ± SEM. n=3 mice per genotype.

(E) Representative images of tdT expression and TH immunostaining of DA neurons in the midbrain of 2 months old Ai9;DAT-Cre (Ctrl) and Ai9;SJ1cKO^DAT. Scale bar: 100 µm.

(F) Representative images of tdT expression and immunostaining of TH and DAT in the striatum of Ai9;DAT-Cre (Ctrl) and Ai9;SJ1cKO^DAT. tdT expressing striatal neurons were positive for TH but not DAT immunostaining, indicating DAT gene promoter is activated in iTHINs but DAT protein is undetected. White arrows: all tdT-expressing THINs driven by DAT-Cre that were positive for TH immunostaining. Scale bar: 50 µm.

(G) Percentage of DAT-Cre driven tdT expressing neurons positive for TH immunostaining shown in (F). Majority of tdT expressing neurons were iTHINs. The number of neurons was quantified on a single stitched coronal striatum section for both tdT expressing neurons and tdT/TH-positive neurons. Data are represented as mean ± SEM. n=3 mice per genotype.

Figure 6. iTHINs establish specific and new connection with dystrophic DA fibers in SJ1 DA cKO striatum

(A) Schematic showing experiment procedure and timeline. pAAV-FLEX-tdTomato (Cre-dependent) was injected unilaterally into the ventral midbrain of SJ1 cKO^TH, SJ1 cKO^DAT mice and Cre mouse controls at 2-3 months old. The animal brains were collected 2 weeks post injection for IHC.

(B) Representative images of TH immunostaining and tdT expression in the injected midbrain DA neurons of both SJ1 cKO^TH and SJ1 cKO^DAT mice. Scale bar: 100 µm.

(C) Representative images of TH immunostaining in the striatum. While tdT injection in TH-Cre and DAT-Cre mice only labeled some DA fibers, tdT injection in SJ1 cKOTH and SJ1 cKO^{DAT} labeled both DA fibers and several iTHINs. White arrows: iTHINs double positive for tdT signal and TH immunostaining. Scale bar: 50 µm.

(D Percentage of tdT expressing neurons positive for anti-TH staining in both SJ1 cKOTH and SJ1 cKO^{DAT}. The number of neurons was quantified on a single stitched coronal striatal section for both tdT and TH/tdT-positive neurons. Data are represented as mean ± SEM. n=3 mice per genotype.

(E) Schematic showing experiment procedure and timeline. rAAV-mTH-EGFP was unilaterally injected into the midbrain of B6, SJ1 cKOTH and SJ1 cKO^{DAT} mice at 2-3 months old. The animal brains were collected 2 weeks post injection for IHC.

(F) Representative images of TH immunostaining with EGFP expression in the injected midbrain DA neurons of both SJ1 cKOTH and SJ1 cKO^{DAT} mice. Scale bar: 100 µm.

(G) Representative images of TH immunostaining and EGFP expression in the striatum. No EGFP expressing THINs observed in B6 striatum. Striatum of injected SJ1 cKOTH and SJ1 cKO^{DAT} mice show EGFP specifically expressed in iTHINs. White arrows: iTHINs double positive for EGFP and anti-TH staining. Scale bar: 50 µm.

(H) Percentage of EGFP expressing neurons positive for anti-TH staining in both SJ1 cKOTH and SJ1 cKO^{DAT}. The number of neurons was quantified on a single stitched coronal striatal section for both EGFP neurons and TH/EGFP-positive neurons. Data are represented as mean ± SEM. n=3 mice per genotype.

(I) Schematic showing experiment procedure and timeline. pAAV-hSyn-mCherry was injected unilaterally in the midbrain of B6, SJ1 cKOTH and SJ1 cKO^{DAT} mice at 2-3 months old. The animal brains were collected 2 weeks post injection for IHC.

(J) Representative images of TH immunostaining and mCherry expression in the injected midbrain DA neurons of both SJ1 cKOTH and SJ1 cKO^{DAT} mice. Scale bar: 100 µm.

(K) Representative images of TH immunostaining and mCherry expression in the striatum of injected mice. No mCherry labeled THINs in B6 striatum. Striatum of injected SJ1 cKOTH and SJ1 cKO^{DAT} show mCherry specifically expressed in iTHINs. White arrows: iTHINs double positive for mCherry and anti-TH staining. Scale bar: 50 µm.

(L) Percentage of mCherry expressing neurons positive for anti-TH staining in both SJ1 cKOTH and SJ1 cKO^{DAT}. The number of neurons was quantified on a single stitched coronal striatal

section for both mCherry and TH/mCherry-positive neurons. Data are represented as
mean ± SEM. n=3 mice per genotype.

Figure 7. THINs appearing in 6-OHDA and Casp3 lesion mice are distinct from iTHINs
observed in SJ1 cKO mice

(A) Schematic showing the midbrain of 2-3 months old B6 mice unilaterally injected with 6-
OHDA and collected 2 weeks post injection for IHC.

(B) Representative coronal section shows the extent of 6-OHDA lesion in the striatum and
midbrain DA neurons of B6 mice. Scale bar: 500 μm.

(C) Schematic showing the midbrain of 2-3 months old DAT-Cre mice unilaterally injected with
pAAV-flex-taCasp3-TEVp (Cre-dependent) and collected 1 month post injection for IHC.

(D) Representative coronal section shows the extent of Casp3-mediated lesion in the striatum
and midbrain DA neurons of DAT-Cre mice. Scale bar: 500 μm.

(E) Representative images of TH-positive neurons (white arrows) observed in the striatum of
SJ1 cKO^{DAT} and lesioned hemisphere of 6-OHDA and Casp3 injected mice. Exposure time
and threshold were set the same for anti-TH imaging in all three conditions. Scale bar: 20 μm.

(F) The expression of anti-TH neurons observed in 6-OHDA and Casp3 lesioned models is
significantly lower than iTHINs observed in SJ1 cKO^{DAT} striatum. Data are represented as
mean ± SEM (****$p < 0.0001$). n=3 mice per genotype.

(G) Unlike iTHINs in SJ1 cKO^{DAT} mice, AADC is undetectable in either 6-OHDA or Casp3
lesioned striatum. Scale bar: 20 μm.

(H) Unlike iTHINs in SJ1 cKO^{DAT} mice, ALDH1A1 is undetectable in either 6-OHDA or Casp3
lesioned striatum. Scale bar: 20 μm.

Figure 8. Acute SJ1 KO in DA neurons of adult and aged mice triggers similar DA
terminal pathology but fewer iTHINs

(A) Schematic showing unilateral stereotaxic injection of AAV9-TH-Cre virus into the ventral
midbrain of 2 months (adult) or 12 months (aged) SJ1 flx/flx mice to obtain SJ1 acute KO in
DA neurons. Animal brains were collected 3-4 months post injection for IHC.

(B) TH and SJ1 immunostaining of the adult and aged SJ1 acute KO mice showed SJ1 specific
KO in the injected midbrains. Scale bar: 100 μm.

(C) DAT immunostaining of the adult and aged SJ1 acute KO mice shows similar clustering

phenotype in dorsal (DS) and ventral striatum (VS) as observed in SJ1 cKO^{DAT} mice. Scale

bar: 20 µm.

(D) No significant difference of the number of DAT clusters between SJ1 cKO^{DAT}, adult and

aged SJ1 acute KO. The average number of clusters is counted in five randomly selected,

300 × 300-µm regions of interest (ROIs). Data are represented as mean ± SEM. n = 3-4 mice

per genotype.

(E) Representative images of striatum show comparable TH/DAT-positive clusters but much

lesser iTHINs in the injected hemisphere of adult and aged SJ1 acute KO mice when

compared to SJ1 cKO^{DAT}. White arrows: iTHINs. Scale bar: 50 µm.

(F) Number of iTHINs in the adult and aged SJ1 acute KO is significantly lower than SJ1

cKO^{DAT} mice. Number of iTHINs was quantified in stitched coronal section of the injected

striatum. Data are represented as mean ± SEM (**$p < 0.01$). n=3-4 mice per genotype.

Supplementary Figure Legends

Supplementary Figure 1. Further Characterization of DA system in SJ1 cKO^{DAT} mice

(A) Generation strategy to obtain SJ1 cKO^{DAT} mouse model. Heterozygous DAT-IRES-Cre line was crossed with homozygous SJ1 flx/flx to obtain double heterozygous DAT-Cre;SJ1 +/flx. Double heterozygous were further crossed with each other to obtain SJ1 cKO^{DAT} and control mice.

(B) Stereological analysis of TH-positive DA cell bodies in the SN and VTA region of Ctrl and SJ1 cKO^{DAT} mice at 2-3 months and 6-8 months age group. There is no significant loss of DA neurons in both ages. The estimated cell numbers in one hemisphere are shown. Data are represented as mean ± SEM (Student's unpaired t-test). n=3 mice per genotype.

(C), (D) Tiling of a single coronal striatum section labelled for DAT in 2 months (C) SJ1 cKO^{DAT} and (D) SJ1-KI^{RQ} mice. Unlike SJ1-KI^{RQ}, DAT clusters were observed in the entire striatum of SJ1 cKO^{DAT} mice. Scale bar: 500 µm.

(E) Representative images of DAT immunostaining in the striatum of SJ1 cKO^{DAT} at 3 weeks, 2 months, 6 months and 12 months age group. Scale bar: 50 µm.

Supplementary Figure 2. Other markers examined in the SJ1 cKO^{DAT} striatum

(A) Immunostaining of vMAT2 with SV2C showed colocalization for clusters in the striatum of 2 months SJ1 cKO^{DAT} mice. Scale bar: 20 µm.

(B), (C) Immunostaining of DAT with (B) Syt1 and (C) alpha-Syn showed partial colocalization with DAT-positive clusters in the striatum of 2 months SJ1 cKO^{DAT} mice. Scale bar: 20 µm.

(D) Immunostaining of DAT with pSer129-α-syn showed no misfolded alpha-Synuclein accumulated in DAT-positive clusters in the striatum of 2 months SJ1 cKO^{DAT} mice. Scale bar: 20 µm.

(E), (F) Immunostaining of DAT with (E) D2R and (F) Darpp32 showed no colocalization with DAT-positive clusters in the striatum of 2 months SJ1 cKO^{DAT} mice, suggesting there is no effect on postsynaptic neurons. Scale bar: 20 µm.

(G), (H) Immunostaining of DAT with (G) GFAP and (H) Iba1 showed similar glia cell pattern in both Ctrl and SJ1 cKO^{DAT} mice at 2 months old, suggesting there is no gliosis and inflammation in the striatum. Scale bar: 20 µm.

Supplementary Figure 3. Endocytic markers examined in the SJ1 cKO^{DAT} striatum

(A)-(D) Double staining of DAT with (A) SJ1, (B) CLC, (C) amphiphysin 1 and (D) endophilin
1 showed no colocalization of these clathrin-related endocytic proteins with DAT-positive
clusters in the striatum of 2 months old SJ1 cKO^{DAT} mice. Scale bar: 20 µm.

Supplementary Figure 4. More EM images of whirl-like membrane accumulation at DA
terminals in SJ1 cKO^{DAT} striatum.

(A), (B) More representative EM images to show the abnormal lipid membrane accumulation
which is correlated with DAT/TH-positive protein clusters in 2 months old SJ1 cKO^{DAT} striatum.
Abnormal whirl-like multilayered membrane (red arrows) surrounding the synaptic vesicles
(SVs) and endosomes (endo). Mitochondria (mt) also labeled in the terminals. Scale bar:
200 nm and 500nm.

Supplementary Figure 5. Western blot analysis of endocytic markers in the SJ1 cKO^{DAT}
striatum

(A) Representative blots for endocytic markers, including SJ1, BIN1, Endophilin 1, Dynamin 1
and AP2 from 2-3 months old Ctrl and SJ1 cKO^{DAT} half brain striatal homogenates. The protein
expression levels of these endocytic factors remain unchanged.

(B) Quantification of proteins level shown in (A). Protein levels were normalized to the level of
actin. n=3-5 mice per genotype.

Supplementary Figure 6. Distribution of iTHINs in whole striatum of SJ1 cKO^{DAT}

(A) Tiling of a single coronal striatum section with TH immunostaining in 2 months old SJ1
cKO^{DAT} mice. Insets: TH-positive clusters and iTHINs were both observed throughout the
striatum, including dorsal striatum (DS), ventral striatum (VS) and olfactory tubercle (OT).
Scale bar: 500 µm.

(B) Representative images of TH immunostaining in the striatum of SJ1 cKO^{DAT} at 3 weeks, 2
months, 6 months and 12 months age group. TH-positive iTHINs (white arrows) and clusters
(asterisk) were labeled. Scale bar: 50 µm.

Supplementary Figure 7. iTHINs are a distinct group of striatal interneurons

(A)-(C) Immunostaining of TH with (A) NPY, (B) PV and (C) ChAT in SJ1 cKO^{DAT} striatum showed that iTHINs is a distinct group of neurons from other striatal interneurons. Scale bar: 20 μm.

(D), (E) Very few iTHINs in the striatum of SJ1 cKO^{DAT} occasionally co-express (D) Calretinin and (E) Darpp32, suggesting heterogeneity within iTHINs subpopulation. Scale bar: 20 μm.

Supplementary Figure 8. A subpopulation of iTHINs in SJ1 cKO^{DAT} dorsal striatum expresses ALDH1A1

(A) Immunostaining of TH and ALDH1A1 in the midbrain of 2 months SJ1 cKO^{DAT} mice shows the ventral tier DA neurons expressing ALDH1A1. Scale bar: 20 μm.

(B) Tiling of a single coronal striatum section immunostained for TH and ALDH1A1 of 2 months old Ctrl and SJ1 cKO^{DAT} mice. TH/ALDH1A1-positive iTHINs were mainly distributed in dorsal striatum (DS). Scale bar: 500 μm

(C) Immunostaining of TH and Calbindin in the midbrain of 2 months SJ1 cKO^{DAT} mice shows some VTA DA neurons expressing Calbindin. Scale bar: 20 μm.

(D) Immunostaining shows that iTHINs in the striatum of 2 months SJ1 cKO^{DAT} mice were negative for Calbindin. Scale bar: 50 μm.

Supplementary Figure 9. Control AAV labeling for the specific DA-iTHINs connection

(A) Schematic showing experiment procedure and timeline. pAAV-FLEX-tdTomato (Cre-dependent) was injected unilaterally into the striatum of TH-Cre, DAT-Cre, SJ1 cKOTH and SJ1 cKO^{DAT} mice at 2-3 months old. Animal brains were collected for IHC 2 weeks post injection.

(B) Representative images of tdT expression and TH immunostaining in the injected striatum of SJ1 cKOTH and SJ1 cKO^{DAT} with their respective Cre line control, TH-Cre and DAT-Cre mice. In TH-Cre, tdT labeled WT THINs (red arrows) that were negative for anti-TH. In SJ1 cKOTH, TH immunostaining labeled all iTHINs (blue arrows), some tdT-expressing WT THINs (red arrow) and tdT expressing iTHINs (white arrow) were observed. No tdT-expressing neurons observed in DAT-Cre injected striatum. While TH immunostaining labeled all iTHINs (blue arrow), tdT specifically labeled some iTHINs (white arrow) in SJ1 cKO^{DAT} mice. Scale bar: 50 μm.

(C) Schematic showing experiment procedure and timeline. rAAV-mTH-EGFP was unilaterally

in the midbrain of B6, Ai9;TH-Cre and Ai9;SJ1 cKOTH mice at 2-3 months old. Animal brains

were collected 2 weeks post injection for IHC.

(D) Representative images of EGFP expression, tdT expression and TH immunostaining in

the striatum of injected mice. EGFP was not expressed in WT THINs (red arrows) in injected

Ai9;TH-Cre mice. However, EGFP was expressed in TH/tdT-positive iTHINS (white arrows)

but not TH-negative/tdT-positive WT THINs (red arrows) in Ai9;SJ1 cKOTH mice. This suggests

EGFP labelling is specific to iTHINs in SJ1 cKO mice. Scale bar: 50 μm.

(E) Schematic showing experiment procedure and timeline. pAAV-hSyn-mCherry was

unilaterally injected into the striatum of B6 mice at 2-3 months old. Animal brains were

collected 2 weeks post injection for IHC.

(F) Representative images of mCherry expression and Darpp32 immunostaining in the

striatum of injected mice. mCherry-positive neurons were randomly observed in the striatum

of injected mice, with a subset of them positive for Darpp32 (white arrows). Scale bar: 50 μm.

1029

Supplementary Figure 10. Controls for lesioned models and Summary of iTHINs

(A) Representative striatum images of Ai9;DAT-Cre mice unilaterally injected with Casp3 (Cre-

mediated) in the midbrain. THINs appearing in the striatum of lesioned model (white arrows)

were not expressing tdT, indicating DAT gene promoter is not activated in lesioned model

THINs. Scale bar: 200 μm.

(B) Illustration summary of the new features of THINs induced in the SJ1 DA cKO striatum.

1036

1037 **Supplementary Figure 11. Controls for SJ1 Acute KO model**

1038 (A), (B) Representative images of (A) adult B6 mice (2 months old) and (B) aged B6 mice (12

1039 months old) unilaterally injected with AAV-TH-Cre virus show no loss of DA somata in the

1040 midbrain or DA fibers in the striatum, validating the safety and non-toxicity of AAV TH-Cre used.

1041 Scale bar: 200 μm.

1042

1043 **Supplementary Table 1. Comparison of THINs observed in WT, genetic SJ1 DA cKO and**

1044 **acute DA lesion mouse models**

1045   Immunostaining against TH, AADC, ALDH1A1 and DAT, and Cre-dependent labelling were

1046   performed to examine the gene/protein expression of these DA markers in THINs and iTHINs.

1047

1048

1049

1050

References

1. Bandres-Ciga, S., Diez-Fairen, M., Kim, J.J., and Singleton, A.B. (2020). Genetics of Parkinson's disease: An introspection of its journey towards precision medicine. Neurobiol Dis *137*, 104782. 10.1016/j.nbd.2020.104782.
2. Ng, X.Y., and Cao, M. (2024). Dysfunction of synaptic endocytic trafficking in Parkinson's disease. Neural Regen Res *19*, 2649-2660. 10.4103/NRR.NRR-D-23-01624.
3. Saheki, Y., and De Camilli, P. (2012). Synaptic vesicle endocytosis. Cold Spring Harbor perspectives in biology *4*, a005645. 10.1101/cshperspect.a005645.
4. Milosevic, I., Giovedi, S., Lou, X., Raimondi, A., Collesi, C., Shen, H., Paradise, S., O'Toole, E., Ferguson, S., Cremona, O., and De Camilli, P. (2011). Recruitment of Endophilin to Clathrin-Coated Pit Necks Is Required for Efficient Vesicle Uncoating after Fission. Neuron *72*, 587-601. 10.1016/j.neuron.2011.08.029.
5. Verstreken, P., Koh, T.-W., Schulze, K.L., Zhai, R.G., Hiesinger, P.R., Zhou, Y., Mehta, S.Q., Cao, Y., Roos, J., and Bellen, H.J. (2003). Synaptojanin Is Recruited by Endophilin to Promote Synaptic Vesicle Uncoating. Neuron *40*, 733-748. 10.1016/S0896-6273(03)00644-5.
6. Schuske, K.R., Richmond, J.E., Matthies, D.S., Davis, W.S., Runz, S., Rube, D.A., van der Bliek, A.M., and Jorgensen, E.M. (2003). Endophilin Is Required for Synaptic Vesicle Endocytosis by Localizing Synaptojanin. Neuron *40*, 749-762. 10.1016/S0896-6273(03)00667-6.
7. Hayashi, M., Raimondi, A., O'Toole, E., Paradise, S., Collesi, C., Cremona, O., Ferguson, S.M., and De Camilli, P. (2008). Cell- and stimulus-dependent heterogeneity of synaptic vesicle endocytic recycling mechanisms revealed by studies of dynamin 1-null neurons. Proc Natl Acad Sci U S A *105*, 2175-2180. 10.1073/pnas.0712171105.
8. Krebs, C.E., Karkheiran, S., Powell, J.C., Cao, M., Makarov, V., Darvish, H., Di Paolo, G., Walker, R.H., Shahidi, G.A., Buxbaum, J.D., et al. (2013). The Sac1 domain of SYNJ1 identified mutated in a family with early-onset progressive Parkinsonism with generalized seizures. Hum Mutat *34*, 1200-1207. 10.1002/humu.22372.
9. Quadri, M., Fang, M., Picillo, M., Olgiati, S., Breedveld, G.J., Graafland, J., Wu, B., Xu, F., Erro, R., Amboni, M., et al. (2013). Mutation in the SYNJ1 gene associated with autosomal recessive, early-onset Parkinsonism. Hum Mutat *34*, 1208-1215. 10.1002/humu.22373.
10. Lesage, S., Mangone, G., Tesson, C., Bertrand, H., Benmahdjoub, M., Kesraoui, S., Arezki, M., Singleton, A., Corvol, J.C., and Brice, A. (2021). Clinical Variability of SYNJ1-Associated Early-Onset Parkinsonism. Front Neurol *12*, 648457. 10.3389/fneur.2021.648457.
11. Dyment, D.A., Smith, A.C., Humphreys, P., Schwartzentruber, J., Beaulieu, C.L., Consortium, F.C., Bulman, D.E., Majewski, J., Woulfe, J., Michaud, J., and Boycott, K.M. (2015). Homozygous nonsense mutation in SYNJ1 associated with intractable epilepsy and tau pathology. Neurobiology of aging *36*, 1222 e1221-1225. 10.1016/j.neurobiolaging.2014.09.005.
12. Rauschendorf, M.A., Jost, M., Stock, F., Zimmer, A., Rosler, B., Rijntjes, M., Piroth, T., Coenen, V.A., Reinacher, P.C., Meyer, P.T., et al. (2017). Novel compound heterozygous synaptojanin-1 mutation causes l-dopa-responsive dystonia-parkinsonism syndrome. Mov Disord *32*, 478-480. 10.1002/mds.26876.
13. Senkevich, K., Parlar, S.C., Chantereault, C., Yu, E., Ahmad, J., Ruskey, J.A., Asayesh, F., Spiegelman, D., Waters, C., Monchi, O., et al. (2024). Are rare heterozygous SYNJ1 variants associated with Parkinson's disease? NPJ Parkinsons Dis *10*, 201. 10.1038/s41531-024-00809-9.

14. Farrow, S.L., Schierding, W., Gokuladhas, S., Golovina, E., Fadason, T., Cooper, A.A., and O'Sullivan, J.M. (2022). Establishing gene regulatory networks from Parkinson's disease risk loci. Brain *145*, 2422-2435. 10.1093/brain/awac022.

15. Gao, X., Huang, Z., Feng, C., Guan, C., Li, R., Xie, H., Chen, J., Li, M., Que, R., Deng, B., et al. (2020). Multimodal analysis of gene expression from postmortem brains and blood identifies synaptic vesicle trafficking genes to be associated with Parkinson's disease. Brief Bioinform. 10.1093/bib/bbaa244.

16. Cremona, O., Di Paolo, G., Wenk, M.R., Luthi, A., Kim, W.T., Takei, K., Daniell, L., Nemoto, Y., Shears, S.B., Flavell, R.A., et al. (1999). Essential role of phosphoinositide metabolism in synaptic vesicle recycling. Cell *99*, 179-188. S0092-8674(00)81649-9 [pii].

17. Pan, P.Y., Sheehan, P., Wang, Q., Zhu, X., Zhang, Y., Choi, I., Li, X., Saenz, J., Zhu, J., Wang, J., et al. (2020). Synj1 haploinsufficiency causes dopamine neuron vulnerability and alpha-synuclein accumulation in mice. Hum Mol Genet *29*, 2300-2312. 10.1093/hmg/ddaa080.

18. Zhu, X., Prakash, S.S., McAuliffe, G., and Pan, P.Y. (2023). Synaptojanin1 Modifies Endolysosomal Parameters in Cultured Ventral Midbrain Neurons. eneuro *10*. 10.1523/ENEURO.0426-22.2023.

19. Pan, P.-Y., Zhu, J., Rizvi, A., Zhu, X., Tanaka, H., and Dreyfus, C.F. (2021). Synaptojanin1 deficiency upregulates basal autophagosome formation in astrocytes. Journal of Biological Chemistry *297*. 10.1016/j.jbc.2021.100873.

20. Cao, M., Wu, Y., Ashrafi, G., McCartney, A.J., Wheeler, H., Bushong, E.A., Boassa, D., Ellisman, M.H., Ryan, T.A., and De Camilli, P. (2017). Parkinson Sac Domain Mutation in Synaptojanin 1 Impairs Clathrin Uncoating at Synapses and Triggers Dystrophic Changes in Dopaminergic Axons. Neuron *93*, 882-896 e885. 10.1016/j.neuron.2017.01.019.

21. Ng, X.Y., Wu, Y., Lin, Y., Yaqoob, S.M., Greene, L.E., De Camilli, P., Cao, M., and (2023). Mutations in Parkinsonism-linked endocytic proteins synaptojanin1 and auxilin have synergistic effects on dopaminergic axonal pathology. npj Parkinson's Disease *9*. 10.1038/s41531-023-00465-5.

22. Nalls, M.A., Pankratz, N., Lill, C.M., Do, C.B., Hernandez, D.G., Saad, M., DeStefano, A.L., Kara, E., Bras, J., Sharma, M., et al. (2014). Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. Nature Genetics *46*, 989-993. 10.1038/ng.3043.

23. Cao, M., Park, D., Wu, Y., and De Camilli, P. (2020). Absence of Sac2/INPP5F enhances the phenotype of a Parkinson's disease mutation of synaptojanin 1. Proc Natl Acad Sci U S A *117*, 12428-12434. 10.1073/pnas.2004335117.

24. Pan, P.Y., Li, X., Wang, J., Powell, J., Wang, Q., Zhang, Y., Chen, Z., Wicinski, B., Hof, P., Ryan, T.A., and Yue, Z. (2017). Parkinson's Disease-Associated LRRK2 Hyperactive Kinase Mutant Disrupts Synaptic Vesicle Trafficking in Ventral Midbrain Neurons. J Neurosci *37*, 11366-11376. 10.1523/JNEUROSCI.0964-17.2017.

25. Tashiro, Y., Sugimoto, T., Hattori, T., Uemura, Y., Nagatsu, I., Kikuchi, H., and Mizuno, N. (1989). Tyrosine hydroxylase-like immunoreactive neurons in the striatum of the rat. Neuroscience Letters *97*, 6-10. doi.org/10.1016/0304-3940(89)90130-4.

26. Nakahara, T., Yamamoto, T., Endo, K., and Kayama, H. (2001). Neuronal ectopic expression of tyrosine hydroxylase in the mouse striatum by combined administration of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and 3-nitropropionic acid. Neuroscience *108*, 601-610. doi.org/10.1016/S0306-4522(01)00441-9.

27. Ibáñez-Sandoval, O., Tecuapetla, F., Unal, B., Shah, F., Koós, T., and Tepper, J.M. (2010). Electrophysiological and Morphological Characteristics and Synaptic Connectivity of Tyrosine Hydroxylase-Expressing Neurons in Adult Mouse Striatum. The Journal of Neuroscience *30*, 6999-7016. 10.1523/JNEUROSCI.5996-09.2010.

28. Dubach, M., Schmidt, R., Kunkel, D., Bowden, D.M., Martin, R., and German, D.C. (1987). Primate neostriatal neurons containing tyrosine hydroxylase:

Immunohistochemical evidence. Neuroscience Letters *75*, 205-210.
doi.org/10.1016/0304-3940(87)90298-9.

29.   Betarbet, R., Turner, R., Chockkan, V., DeLong, M.R., Allers, K.A., Walters, J., Levey,
A.I., and Greenamyre, J.T. (1997). Dopaminergic neurons intrinsic to the primate
striatum. J Neurosci *17*, 6761-6768. 10.1523/JNEUROSCI.17-17-06761.1997.

30.   Porritt, M.J., Batchelor, P.E., Hughes, A.J., Kalnins, R., Donnan, G.A., and Howells,
D.W. (2000). New dopaminergic neurons in Parkinson's disease striatum. Lancet
356, 44-45. 10.1016/S0140-6736(00)02437-5.

31.   Cossette, M., Lecomte, F., and Parent, A. (2005). Morphology and distribution of
dopaminergic neurons intrinsic to the human striatum. J Chem Neuroanat *29*, 1-11.
10.1016/j.jchemneu.2004.08.007.

32.   Huot, P., Levesque, M., and Parent, A. (2006). The fate of striatal dopaminergic
neurons in Parkinson's disease and Huntington's chorea. Brain *130*, 222-232.
10.1093/brain/awl332.

33.   Xenias, H.S., Ibáñez-Sandoval, O., Koós, T., and Tepper, J.M. (2015). Are Striatal
Tyrosine Hydroxylase Interneurons Dopaminergic? The Journal of Neuroscience *35*,
6584-6599. 10.1523/JNEUROSCI.0195-15.2015.

34.   Tepper, J.M., Koós, T., Ibanez-Sandoval, O., Tecuapetla, F., Faust, T.W., and Assous,
M. (2018). Heterogeneity and Diversity of Striatal GABAergic Interneurons: Update
2018. Frontiers in Neuroanatomy *12*, 91. 10.3389/fnana.2018.00091.

35.   Darmopil, S., Muñetón‐Gómez, V.C., De Ceballos, M.L., Bernson, M., and Moratalla,
R. (2008). Tyrosine hydroxylase cells appearing in the mouse striatum after
dopamine denervation are likely to be projection neurones regulated by l‐DOPA.
European Journal of Neuroscience *27*, 580-592. 10.1111/j.1460-9568.2008.06040.x.

36.   Masuda, M., Miura, M., Inoue, R., Imanishi, M., Saino‐Saito, S., Takada, M.,
Kobayashi, K., and Aosaki, T. (2011). Postnatal development of tyrosine hydroxylase
mRNA‐expressing neurons in mouse neostriatum. European Journal of
Neuroscience *34*, 1355-1367. 10.1111/j.1460-9568.2011.07873.x.

37.   Mao, M., Nair, A., and Augustine, G.J. (2019). A Novel Type of Neuron Within the
Dorsal Striatum. Frontiers in Neural Circuits *13*, 32. 10.3389/fncir.2019.00032.

38.   Cai, H., Liu, G., Sun, L., and Ding, J. (2014). Aldehyde Dehydrogenase 1 making
molecular inroads into the differential vulnerability of nigrostriatal dopaminergic
neuron subtypes in Parkinson's disease. Translational Neurodegeneration *3*, 27.
10.1186/2047-9158-3-27.

39.   Anderegg, A., Poulin, J.-F., and Awatramani, R. (2015). Molecular heterogeneity of
midbrain dopaminergic neurons – Moving toward single cell resolution. FEBS Letters
589, 3714-3726. doi.org/10.1016/j.febslet.2015.10.022.

40.   Palfi, S., Leventhal, L., Chu, Y., Ma, S.Y., Emborg, M., Bakay, R., Déglon, N.,
Hantraye, P., Aebischer, P., and Kordower, J.H. (2002). Lentivirally Delivered Glial
Cell Line-Derived Neurotrophic Factor Increases the Number of Striatal
Dopaminergic Neurons in Primate Models of Nigrostriatal Degeneration. The Journal
of Neuroscience *22*, 4942. 10.1523/JNEUROSCI.22-12-04942.2002.

41.   Tande, D., Hoglinger, G., Debeir, T., Freundlieb, N., Hirsch, E.C., and Francois, C.
(2006). New striatal dopamine neurons in MPTP-treated macaques result from a
phenotypic shift and not neurogenesis. Brain *129*, 1194-1200. 10.1093/brain/awl041.

42.   Ünal, B., Shah, F., Kothari, J., and Tepper, J.M. (2015). Anatomical and
electrophysiological changes in striatal TH interneurons after loss of the nigrostriatal
dopaminergic pathway. Brain Structure and Function *220*, 331-349. 10.1007/s00429-
013-0658-8.

43.   Troshev, D., Bannikova, A., Blokhin, V., Pavlova, E., Kolacheva, A., and Ugrumov, M.
(2023). Compensatory Processes in Striatal Neurons Expressing the Tyrosine
Hydroxylase Gene in Transgenic Mice in a Model of Parkinson's Disease.
International Journal of Molecular Sciences *24*, 16245. 10.3390/ijms242216245.

44. Erben, L., Welday, J.P., Murphy, R., and Buonanno, A. (2022). Toxic and Phenotypic
Effects of AAV_Cre Used to Transduce Mesencephalic Dopaminergic Neurons.
International Journal of Molecular Sciences *23*. 10.3390/ijms23169462.
45. Rezai Amin, S., Gruszczynski, C., Guiard, B.P., Callebert, J., Launay, J.-M., Louis, F.,
Betancur, C., Vialou, V., and Gautron, S. (2019). Viral vector-mediated Cre
recombinase expression in substantia nigra induces lesions of the nigrostriatal
pathway associated with perturbations of dopamine-related behaviors and hallmarks
of programmed cell death. Journal of Neurochemistry *150*, 330-340.
doi.org/10.1111/jnc.14684.
46. Özçete, Ö.D., Banerjee, A., and Kaeser, P.S. (2024). Mechanisms of
neuromodulatory volume transmission. Molecular Psychiatry *29*, 3680-3693.
10.1038/s41380-024-02608-3.
47. Chanaday, N.L., Cousin, M.A., Milosevic, I., Watanabe, S., and Morgan, J.R. (2019).
The Synaptic Vesicle Cycle Revisited: New Insights into the Modes and Mechanisms.
The Journal of Neuroscience *39*, 8209. 10.1523/JNEUROSCI.1158-19.2019.
48. Mejaes, J.I., Saenz, J., O'Brien, C., Pizzano, C.M., Pan, P.Y., and Barker, D.J. (2024).
Haploinsufficiency of the Parkinson's disease gene synaptojanin1 is associated with
abnormal responses to psychomotor stimulants and mesolimbic dopamine signaling.
Front Behav Neurosci *18*, 1359225. 10.3389/fnbeh.2024.1359225.
49. Delignat-Lavaud, B., Kano, J., Ducrot, C., Masse, I., Mukherjee, S., Giguere, N.,
Moquin, L., Levesque, C., Burke, S., Denis, R., et al. (2023). Synaptotagmin-1-
dependent phasic axonal dopamine release is dispensable for basic motor behaviors
in mice. Nat Commun *14*, 4120. 10.1038/s41467-023-39805-7.
50. Cai, X., Liu, C., Tsutsui-Kimura, I., Lee, J.-H., Guo, C., Banerjee, A., Lee, J., Amo, R.,
Xie, Y., Patriarchi, T., et al. (2024). Dopamine dynamics are dispensable for
movement but promote reward responses. Nature *635*, 406-414. 10.1038/s41586-
024-08038-z.
51. Troshev, D., Bannikova, A., Blokhin, V., Kolacheva, A., Pronina, T., and Ugrumov, M.
(2022). Striatal Neurons Partially Expressing a Dopaminergic Phenotype: Functional
Significance and Regulation. International Journal of Molecular Sciences *23*, 11054.
10.3390/ijms231911054.
52. Rice, M.E., Patel, J.C., and Cragg, S.J. (2011). Dopamine release in the basal
ganglia. Neuroscience *198*, 112-137. 10.1016/j.neuroscience.2011.08.066.
53. Ramírez-Jarquín, U.N., Sharma, M., Shahani, N., Li, Y., Boregowda, S., and
Subramaniam, S. (2022). Rhes protein transits from neuron to neuron and facilitates
mutant huntingtin spreading in the brain. Science Advances *8*, eabm3877.
10.1126/sciadv.abm3877.
54. Rufino-Ramos, D., Leandro, K., Perdigão, P.R.L., O'Brien, K., Pinto, M.M., Santana,
M.M., van Solinge, T.S., Mahjoum, S., Breakefield, X.O., Breyne, K., and Pereira de
Almeida, L. (2023). Extracellular communication between brain cells through
functional transfer of Cre mRNA mediated by extracellular vesicles. Molecular
Therapy *31*, 2220-2239. 10.1016/j.ymthe.2023.05.012.
55. Busceti, C.L., Bucci, D., Molinaro, G., Di Pietro, P., Zangrandi, L., Gradini, R.,
Moratalla, R., Battaglia, G., Bruno, V., Nicoletti, F., and Fornai, F. (2012). Lack or
Inhibition of Dopaminergic Stimulation Induces a Development Increase of Striatal
Tyrosine Hydroxylase-Positive Interneurons. PloS one *7*, e44025.
10.1371/journal.pone.0044025.

1257

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

A

B

Supplementary Figure 5

A

TH

SJ1 cKO^DAT

DS

DS

VS

VS

OT

OT

2 months

B

| 3 weeks | 2 months | 6 months | 12 months |

TH

SJ1 cKO^DAT

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9

A

TH   tdT   Merge

6-OHDA

Casp3$^{DAT-Cre}$

Ai9;DAT-Cre mice

B

△ WT THINs (TH gene+/TH-)

▲ iTHINs (TH+/AADC+/DAT gene+/ALDH1A1+)

▲ iTHINs (TH+/AADC+/DAT gene+/ALDH1A1-)

DMS

DLS

NAc

OT

MSN / other INs

DA-iTHINs communication

Dystrophic DA terminals

SNc   VTA

SJ1 DA cKO

A9 SNc DA

A10 VTA DA

Supplementary Figure 10

A

AAV-TH-CRE injected in adult B6 mice

B

AAV-TH-CRE injected in aged B6 mice

Supplementary Figure 11

Mice Model		TH	AADC	ALDH1A1	DAT
Ai9;TH-Cre	WT THINs	- (*TH gene+)	-	-	-
SJ1 cKO^{DAT}	iTHINs	+++	+++	+++ (*subregion)	- (*DAT gene+)
SJ1 cKOTH	iTHINs	+++	+++	+++ (*subregion)	-
6-OHDA model		+	-	-	-
Casp3 model		+	-	-	-

+++ positive and high expression;

+ positive and mild expression;

- negative/cannot be detected with immunostaining;

*remarks

Table S1. Immunostaining of THINs observed in genetic and lesioned mice models