Tissue-Specific Iron Levels Modulate Lipid Peroxidation and the FLASH Radiotherapy Effect

Running title: Interplay of Iron and Lipid Peroxidation in the FLASH effect

Nuria Vilaplana-Lopera[1*], Jiyoung Kim[1*], Gilyeong Nam[2], Iain D. C. Tullis[1], Salome Paillas[1], Jia-Ling
Ruan[1], Yanyan Jiang[1], Sohee Park[3], Pei Ju Lee[4], Tianxu Hou[1], Ellie Walker[1], Ammar Abu-Halawa[1],
Mark A. Hill[1], Changhoon Choi[3], Ik Jae Lee[2], Youngtae Jeong[5], Samira Lakhal-Littleton[6], Chee Kin
Then[7,8], Shing-Chuan Shen[4], Amato J. Giaccia[1], Kristoffer Petersson[1], Eui Jung Moon[1#]

[1]Department of Oncology, University of Oxford, Oxford, United Kingdom
[2]Department of Radiation Oncology, Severance Hospital, Yonsei University Medical School, Seoul,
Republic of Korea
[3]Department of Radiation Oncology, Samsung Hospital, Seoul, Republic of Korea
[4]Graduate Institute of Medical Sciences, College of Medicine, Taipei Medical University, Taipei,
Taiwan
[5]Department of New Biology, Daegu Gyeongbuk Institute of Science and Technology (DGIST), Daegu,
Republic of Korea.
[6]Department of Physiology, Anatomy & Genetics, University of Oxford, Oxford, United Kingdom
[7]Department of Radiation Oncology, Shuang Ho Hospital, Taipei Medical University, New Taipei City,
Taiwan
[8]Graduate Institute of Clinical Medicine, College of Medicine, Taipei Medical University, Taipei,
Taiwan
21
#Corresponding author
Eui Jung Moon
Old Road Campus Research Building
Roosevelt Drive, Headington, OX3 7DQ
United Kingdom
ejung.moon@oncology.ox.ac.uk
28

*N.V.L. and J.K. are co-first authors.

Abstract

Iron is vital to living cells, playing a key role in cellular respiration, DNA synthesis, and various

metabolic functions. Importantly, cancer cells have a higher dependency on iron compared to normal

cells to support their rapid growth and survival. Due to this fact, tumors are more vulnerable to

ferroptosis, an iron-dependent form of regulated cell death. Radiation therapy (RT), a standard treatment

for many cancer patients, is known to induce ferroptosis. Ultra-high dose rate FLASH RT offers an

improved therapeutic window by minimizing damage to normal tissues while preserving tumor control.

However, the precise biological mechanisms behind the protective effects of FLASH RT on normal

tissues remain unclear. In this study, we propose that variations in lipid peroxidation and ferroptosis,

driven by intrinsic differences in iron levels between normal and cancerous tissues, contribute to this

effect. Our findings show that FLASH RT increases lipid peroxidation and induces ferroptosis in tumor

cells but does not significantly elevate lipid peroxidation in normal tissues compared to conventional

RT. To determine whether raising iron levels in normal tissues could abrogate the protective effects of

FLASH, mice were fed a high-iron diet before RT. A high-iron diet before and after RT reversed the

protective effect of FLASH, resulting in increased intestinal damage and lipid peroxidation. This

suggests that baseline iron levels and iron-driven lipid peroxidation are critical factors in mediating the

protective outcomes of FLASH RT. Overall, our study sheds light on the role of iron in modulating RT

responses and provides new mechanistic insights into how FLASH RT influences normal and cancerous

tissues.

Introduction

Iron is a vital element that plays a fundamental role in maintaining essential cellular functions such as oxygen transport, energy production, hematopoiesis, and DNA replication (1). Its biological importance becomes particularly essential in cancer, where tumor cells exhibit a strong dependence on iron for survival. Therefore, elevated levels of iron and ferritin, the primary intracellular iron storage protein, are frequently observed in the tissues and serum of cancer patients (2-5). This iron dependency opens a critical therapeutic window through ferroptosis, an iron-dependent form of regulated cell death characterized by excessive lipid peroxidation (6).

Radiation therapy (RT), a standard care for many cancer patients to achieve local control and to treat metastatic burden, has been shown to promote ferroptosis by inducing lipid peroxidation, thereby enhancing tumor cell death (7). Furthermore, combining RT with ferroptosis inducers such as RSL3, erastin, and sulfasalazine, has been shown to increase tumor radiosensitivity and overcome radioresistance (7-9). While RT works by damaging tumor DNA either directly or indirectly via the generation of reactive oxygen species (ROS), ferroptosis operates independently of DNA damage pathways, making it a promising approach for developing therapeutic strategies to increase RT response (10).

FLASH RT uses ultra-high dose rates to increase the therapeutic window of RT (11). Specifically, FLASH RT results in less normal tissue toxicity while maintaining tumor control rates comparable to conventional RT. Normal tissue sparing after FLASH RT is known as the FLASH effect and has been attributed in part to the interplay between oxygen depletion and ROS production (11-14). Although studies have demonstrated that FLASH RT reduces tissue oxygen concentrations both in normal tissues and tumors, oxygen depletion may not be the only mechanism accounting for normal tissue sparing, as the observed reduction in oxygen could be considered insufficient to induce tissue protection (15, 16). It has also been suggested that radical-radical recombination as well as immune modulation are potential mechanisms of FLASH RT (14, 17-19). Recently, the contribution of lipids and lipid peroxidation was proposed as the key link between FLASH and ferroptosis, an iron-dependent regulated cell death pathway (20-22). However, the biological mechanisms by which FLASH RT alters lipid peroxidation and potentially ferroptosis remain unclear. Therefore, this study investigated how lipid peroxidation

and ferroptosis are differentially modulated after FLASH RT in tumor and normal tissues and whether

varying iron levels in these tissues plays a role in the FLASH effect.

Materials and Methods

Cell lines and cell culture

A549 and MDA-MB-231 were purchased from ATCC and maintained at 37 °C in a humidified

incubator with 5% CO_2 in Dulbecco's Modified Eagle Medium supplemented with 10% fetal bovine

serum (F7524, Sigma Aldrich) and 1% antibiotic/antimycotic (15240-062, Gibco). Cells were routinely

tested for mycoplasma.

siRNA transfection

ONTARGETplus SMARTpool siRNA targeting TFRC (Horizon Discovery) was transfected into A549

and MDA-MB-231 cells according to the manufacturer's instructions. Briefly, 2×10^5 A549 and MDA-

MB-231 cells were seeded in 6-well plates. After 24 hours of incubation, the cells were washed and

cultured in 1.6 mL of antibiotic-free DMEM supplemented with 10% FBS. A siRNA solution was

prepared by diluting 10 µL of 20 µM TFRC-targeting siRNA in 190 µL of Opti-MEM (31985070,

GIBCO), while 7.5 µL of RNAiMax (13778030, Thermo Fisher Scientific) was diluted in 192.5 µL of

Opti-MEM. The two diluted components were then combined and incubated at room temperature for

15 minutes before being added to the cells. Cells were incubated overnight before being seeded for

further experiments.

Intracellular iron measurement

Intracellular iron levels were measured using the FerroOrange (F374, Dojindo) probe according to the

manufacturer's instructions. Briefly, cells were irradiated and incubated for the specified times before

being washed with PBS and incubated with a 1 µM FerroOrange solution in the serum-free phenol red-

free media (31053028, Gibco) for 30 minutes at 37 ℃. Fluorescence (Ex 531/40 – Em 629/53) was

directly imaged after the incubation using the Celigo Image Cytometer (Nexcelom).

Reagents

Ferr-1 (ab142829, Abcam) was dissolved in DMSO at 10 mM concentration and diluted in cell culture

media to the specified concentrations. Ammonium iron (II) sulfate (09719, Merck) was dissolved in

deionized water at 100 mM concentration and diluted in cell culture media to specified concentrations.

Deferoxamine (D9533, Sigma Aldrich) was dissolved in DMSO at 100 mM concentration and diluted

in cell culture media to the specified concentrations.

RNA sequencing

RNA samples were extracted from MDA-MB-231 cells 24 hours after being irradiated with 10 Gy in a

^{137}Cs irradiator for sequencing as previously described[23]. Briefly, RNA was extracted using the

RNeasy Mini Kit (74104, Qiagen) following the manufacturer's protocol. Library preparation,

sequencing (paired-end 150 bp sequencing, 20 million reads), and bioinformatic analysis were

performed by Novogene Co. (Cambridge, UK) using the Illumina NovaSeq platform. The genes that

significantly altered by 10 Gy RT were analyzed by Enrichr[24-26]. The Gene Set Enrichment Analysis

(GSEA) was performed using gene sets obtained from the Molecular Signatures Database (MSigDB

v2024.1.Hs) [27]. Raw gene count data were normalized to transcripts per million (TPM) prior to GSEA

analysis. Analyses were conducted using the GSEA software (v4.3.3) from the Broad Institute. Gene

sets with nominal P-value < 0.05 and False Discovery Rate (FDR) $< 25\%$ were considered significantly

enriched. Additionally, the heatmap was generated using Python (3.11.8).

Radiation

RT was performed in the Radiation Biophysics facility at the University of Oxford. Cells were exposed

to γ-rays from a ^{137}Cs sealed source irradiator (GSR D1, Gamma-Service Medical, Leipzig, Germany;

dose rate of 1.7 Gy/min). For alpha particle treatment, cells were irradiated with 3.3 MeV alpha particles

(LET = 121 keV μm^{-1}) using the Oxford ^{238}Pu irradiator [28].

FLASH radiation

Conventional and FLASH RT was performed using an electron linear accelerator which delivers

electrons of 6 MeV nominal energy in a circular horizontal beam collimated to 5 cm in diameter,

previously described with more details (29, 30). *In vitro* experiments were performed in T12.5 flasks

or 35 mm cell culture dishes irradiated one by one in a vertical position in the central part of the beam

(Fig. S4). For *in vivo* experiments, the mice were anesthetized with isoflurane supplemented with ~55%

oxygen (90% oxygen evenly mixed with air), then placed in an immobilization cradle and irradiated in

an upright position. A 6 mm brass collimator with a 20 x 20 mm^2 or a 33×30 mm^2 aperture was used

to further define the RT field covering the whole thorax or the whole abdominal area respectively

(Supplementary Fig. 5B). RT was delivered with 3.5 µs electron pulses with a repetition rate of 25 or

300 Hz (25 Hz for conventional and 300 Hz for FLASH). Dose rates were 0.1 Gy/s for conventional

and > 2 kGy/s for FLASH, with a dose-per-pulse of ~4 mGy and 5 Gy, respectively. Doses ranging

from 5 to 20 Gy were given for the *in vivo* and *in vitro* experiments.

The prescribed doses were verified during delivery with pieces of Gafchromic EBT-XD film (Ashland

Inc, Covington, KY) placed at the surface or between (abdominal RT) two pieces of Perspex

representing a mouse phantom positioned exactly as the mice in the beam path (30). The films were

scanned (Epson Perfection v850 Pro, Seiko Epson Corporation, Nagano, Japan), 24 hours post-RT, and

the red channel was analyzed for a 20×20 mm^2 central part of the exposed film, using ImageJ (version

1.52a, Wayne Rasband). The film had previously been calibrated in a clinical 6 MeV electron beam

from a Varian Truebeam (Varian Medical Systems Inc, Palo Alto, CA) linear accelerator at the

Churchill Hospital site in Oxford, UK. Online verification of the delivered dose was achieved using an

Advanced Markus ionization chamber (PTW-Freiburg GmbH, Freiburg, Germany) positioned in the

beam (corrected for ion recombination) in the beam energy monitor and collimator system (not

disturbing the collimated part of the beam), as well as a toroidal beam charge monitor (31, 32). The

beam energy monitor was used to verify that the electron beam energy was consistently 6 MeV, for

both FLASH and conventional RT (29). The overall uncertainty in dosimetry was estimated to be 4%

(including a measured output variation of within 2%).

Proton RT

Proton beam RT was given as previously described(33). After placing cell dishes at the midpoint of a

10 cm wide spread-out-Bragg-peak (SOBP, a 230 MeV proton beam generated by a proton therapy

system (Sumitomo Heavy Industries, Ltd., Niihama, Japan) was given at a dose rate of 2.14 Gy/min at

the Samsung Proton Therapy Center in Seoul, South Korea.

Lipid peroxidation

Lipid peroxidation was assessed using the BODIPY 581/591 C11 (D3861, Invitrogen) probe according

to the manufacturer's instructions. Briefly, cells were irradiated and incubated for the specified times

before incubation with a 2 μM BODIPY 581/591 C11 solution in PBS for 30 minutes at 37 ºC. The

cells were detached and washed with PBS before FITC fluorescence intensity (oxidized BODIPY

emission) was measured using a CytoFLEX (Beckman Coulter) flow cytometer. FlowJo software was

used to assess the geometric mean of the intensity of oxidized BODIPY 581/591 C11.

Clonogenic assay

For [137]Cs IR clonogenic assays, single cells were plated in 6-well plates and allowed to settle for 24

hours before RT. For FLASH vs. conventional irradiated clonogenic assays, cells were seeded in T12.5

flasks and treated with the specified concentrations of Ferr-1 for 24 hours before RT. Prior to RT, the

flasks were positioned vertically. Following RT, the cells were detached, counted, and replated as single

cells in 6-well plates. Seeding densities were optimized for each cell line and RT dose. Colonies were

cultured for 8 to 12 days before being stained with a 20% methanol/0.5% crystal violet solution.

Colonies were then counted to determine plating efficiency (PE = colony number/number of cells

plated), and the surviving fraction (SF) was calculated using the formula: SF = (number of colonies

formed after treatment) / (number of cells seeded $\times$ PE), following the methodology of a previous study

(34).

Patient data

In total, 748 genes in the Cancer Gene Consensus were acquired from the Catalogue Of Somatic

Mutations In Cancer (COSMIC) and analyzed by DAVID functional analysis to identify the enriched

pathways (35, 36). Tissue microarrays were purchased from Biomax (BR1008b – Breast cancer,

LC10011b – Lung cancer). *TFRC* expression data from the TCGA PanCancer Atlas was obtained from

cBioPortal, including mRNA expression z-scores (log RNA Seq V2 RSEM) normalized relative to

normal samples, along with corresponding clinical data (37). *TFRC* expression across different tumor

types was visualized using TCGA datasets through The University of ALabama at Birmingham CANcer

data analysis Portal (UALCAN) (38).

In vivo experiments

All animal experiments were performed according to the guidelines of the United Kingdom Home

Office and the University of Oxford under the project licenses PP8415318 and PP4558762. Female

BALB/cAnNCrl mice were purchased from Charles River (6-8 weeks, strain code 028) and housed in

individually ventilated cages containing no more than six mice per cage, in a 12/12-hour light/dark

cycle. For normal mouse lung studies, lung tissues were collected 24 hours or 7 days after RT either at

conventional or FLASH dose rates. To determine iron accumulation in the upper intestine, mice were

fed with either control diet with 200 ppm iron (TD. 08713, Envigo) or a high-iron diet with 5000 ppm

iron (TD140464, Envigo) for 24, and 48 hours (39). Intestinal tissues were collected at the specified

time points to assess the correlation between iron diet and RT effects. For RT studies, mice were fed

either a control diet or a high-iron diet for evaluation. After 24 hours, they were either kept as controls

or exposed to conventional or FLASH dose rate RT. Post-RT, mice were monitored and weighed daily.

One group of mice returned to a control diet, and the other group of mice remained on a high-iron diet.

Seventy-two hours after RT, mice were sacrificed to collect blood samples and intestinal tissues.

For tissue processing, lung tissues were inflated with 10% formalin and kept in formalin for 24 hours

before storing in 70% ethanol. Intestinal samples were flushed with PBS, fixed in formalin for 24 hours.

Then they were cut with micro scissors to expose the lumen. The tissues were then rolled from the

posterior end to reveal the inner lumen and stored in 70% ethanol until being processed and embedded

in paraffin. Sections of 6 μm thickness were prepared for immunohistochemistry analysis using

microtome.

Mouse tissues

H460 and Calu6 xenograft tumor tissues as well as normal mouse lung and intestine tissues treated with conventional, or FLASH RT were kindly gifted by the Petersson group. Briefly, H460 and Calu6 cells were subcutaneously injected into athymic nude mice. When the tumor size reached 100 mm^3, mice were randomized into three groups: control, conventional RT, and FLASH RT (15 Gy for H460, 20 Gy for Calu6). When the tumor size reached 800 mm³ or 50 days post-RT, the mice were sacrificed to collect tumor tissues. The harvested tissues were fixed in formalin, then stored in 70% ethanol until further processing and paraffin embedding. Six µm sections were cut and used for immunohistochemistry.

Immunohistochemistry

Paraffin-embedded tissues were sectioned and stained as previously described (23, 40). Briefly, after incubation in a 65 °C oven for 30 minutes, slides were deparaffinized by immersing in Histo-Clear (National Diagnostics, HS-200) twice for 3 minutes followed by rehydration for 3 minutes in 100 %, 70 %, and 50 % ethanol. The slides were washed in deionized water for 5 minutes before antigen retrieval was performed in 1 x Citrate buffer (10 x Citrate buffer (pH 6.0), Sigma #C9999; diluted with deionized water and 0.05 % Tween-20) which was heated to 110 °C in a pressure cooker for 3 minutes. Then, the slides were cooled at room temperature for 20 minutes. Endogenous peroxidase and phosphatase activities were blocked by Dual Endogenous Enzyme Block (Agilent Technologies, #S2003) for 20 minutes. The slides were blocked with 10 % goat serum for 30 minutes and incubated with an anti-4-HNE antibody (Thermo Fisher, #BS-6313R or abcam, #ab48506) diluted in 1:100 or 1:200 with antibody diluent (abcam, #ab64211) at 4 °C overnight. After washing three times with PBS/0.05 % Tween-20, the secondary antibody with HRP-labelled polymers was added to slides (Agilent Technologies; #K4003 or #K4001) and incubated at room temperature for 20 minutes. Slides were then washed three times with PBS/0.05% Tween-20 and incubated with DAB (3,3′-Diaminobenzidine) (#K3468) to visualize the HRP signals. The slides were counterstained with hematoxylin (1.09249.0500, Sigma Aldrich) and mounted with aqueous mounting medium (1.08562.0050, Merck).

Tissue iron was stained using a Prussian Blue staining kit (ab150674, Abcam) according to the

manufacturer's protocol, with a peroxide-DAB enhancing step. Briefly, slides were deparaffinized and

incubated with Prussian Blue solution for 1 hour at room temperature before washing 3 times with deionized

water. If no blue prussiate was visible, slides were incubated with a 0.3% hydrogen peroxide and 0.01M

sodium azide methanol solution for 30 minutes at room temperature. Slides were then washed 4 times with

PBS, incubated with DAB until tissues turned brown. Slides were washed with 3 changes of deionized

water before counterstaining with nuclear fast red for 4 minutes at room temperature. Slides were

dehydrated in 50%, 70%, and 100% ethanol and 2 changes of 100% xylene before mounting with DPX

(06522, Sigma Aldrich).

Hematoxylin and eosin (H&E) staining was performed by deparaffinizing tissues followed by

incubation with 25% Harris' hematoxylin (CODE) diluted in water for 3 minutes. Slides were dipped

10 times in acid alcohol (1% hydrochloric acid in 70% ethanol) and then washed with 10 dips in tap

water. Slides were dipped 10 times into Scott's tap water and then washed with 10 dips in tap water.

Slides were dipped 10 times in 0.5% lithium carbonate and then washed with running tap water. Slides

were then dipped 10 times in 50%, 70%, and 90% ethanol. Slides were then incubated in alcoholic eosin

for 1 minute before being dipped 10 times 90% ethanol and repeated with fresh 90% ethanol. Slides

were then dehydrated in 100% ethanol and 2 changes of 100% xylene before mounting with DPX.

All slides were imaged with an Aperio Scanscope CS digital pathology scanner (Leica Biosystems

Imaging, Inc.) at 40X or 20X magnification.

For 4-HNE staining, three representative fields were imaged per slide at 10x or 20 x magnification.

DAB staining intensity was analyzed using FIJI (Image-J-based open-source software). Images were

chosen from the irradiated lung lobe and upper intestine areas where iron is mainly accumulated.

Positively stained areas were extracted using color deconvolution with the brightness and contrast

optimized for each set of images and the values were kept consistent across all images from the same

set. Positively stained areas were converted into binary images, and the integrated density values were

calculated.

The amount of iron in the upper intestines or microarray cores were assessed using FIJI. First, positively

stained iron areas were selected and extracted by using color threshold, converted into binary images

and positive areas were measured. Similarly, total tissue areas were selected and extracted using the color threshold, converted into binary and measured. The percentage of iron in each tissue/core was determined using the iron positive and total tissue areas.

Three representative fields of H&E stained images 1 mm in size were randomized and the remaining crypts were manually counted (Fig. S5). The percentage of remaining crypts was determined by comparing to the untreated and non-irradiated control.

Ferritin ELISA

Mouse serum was collected by centrifugation of mouse blood at 3000 rpm for 10 minutes. Serum ferritin levels were measured by using a commercially available mouse ferritin ELISA kit according to the manufacturer's protocol (Abcam, #ab157713).

Statistical analysis

The data are presented as the mean, the standard deviation, and the individual observations of at least three replicates. The data were analyzed using GraphPad Prism software (GraphPad Prism 10, Dotmatics) as indicated in each figure. Briefly, for comparisons between two groups, we performed two-tailed Student's *t*-tests. Nonparametric Kolmogorov-Smirnov test was used for tissue microarray data. To compare multiple groups, we used one-way ANOVA for single-condition experiments and two-way ANOVA for experiments involving multiple conditions. A *P* value of <0.05 was considered statistically significant.

Study approval

This study involved mouse studies. Experimental procedures were carried out under a project license issued by the UK Home Office under the UK Animals (Scientific Procedures) Act of 1986.

Results

Iron is essential for tumor survival

Iron staining in tissue microarrays (TMA) of breast and lung cancer patients using Prussian Blue with

DAB (3,3′-Diaminobenzidine) indicated that iron was highly elevated in both lung (Fig. 1A) and breast

(Fig. 1B) cancer samples compared to corresponding normal tissues. The functional enrichment analysis

of 748 genes from the Catalogue of Somatic Mutations in Cancer (COSMIC) Cancer Gene Consensus

further confirmed that cancer driving genes involved in iron metabolism or homeostasis including "4

iron 4 sulfur cluster binding", "response to iron ion", and "cellular iron ion binding" were significantly

amplified, supporting the crucial needs of iron in cancer (Fig. 1C) (35). Specifically, transferrin receptor

(TFRC), a receptor facilitating iron transport, was included in COSMIC. Expression of TFRC was

significantly correlated with poor overall survival in all cancer types (Fig. 1D) and its expression was

highly upregulated in the majority of cancer types (Fig. 1E). The inhibition of *TFRC*, which inhibited

iron uptake, significantly decreased cell survival in both lung (A549) (Fig. 1F) and breast (MDA-MB-

231) cancer cells (Fig. 1G), indicating that iron is critical for tumor cell survival.

RT induces lipid peroxidation and ferroptosis

RT induces cell death primarily through the induction of DNA damage. Recently, ferroptosis has been

identified as a contributor to RT-induced cell death that occurs independently of DNA damage (7-10).

RNA sequencing was performed in MDA-MB-231, a human breast cancer cell line, treated with 0 or

10 Gy of RT to determine changes in gene expression induced by RT. Cells were irradiated with γ-rays

from a ^{137}Cs irradiator 24 hours before RNA extraction and transcriptomic analysis. Kyoto

Encyclopedia of Genes and Genomes (KEGG) enrichment analysis using Enrichr identified significant

changes in genes associated with the cell cycle, DNA replication, p53 pathway, and DNA replication,

which are hallmarks of cells exposed to RT-induced DNA damage (24-26) (Fig. 2A). Additionally, RT

also affected genes involved in ferroptosis (Fig. 2A, Fig. S1A) in line with previous studies (7-9). Gene

Set Enrichment Analysis (GSEA) further identified a strong association between RT exposure and

significant alterations in ferroptosis-related genes (27) (Fig. 2B, Fig. S1B). To assess the biological

significance of the changes in gene expression, the effect of RT on lipid peroxidation and ferroptosis

was examined in MDA-MB-231 (human breast cancer) and A549 (human lung cancer) cell lines *in*

vitro. We first detected lipid peroxidation 24 hours after exposure to 4 Gy of RT that increased in a

dose-dependent manner up to 10 Gy (Fig. 2C). There was also a time-dependent increase in lipid

peroxidation when irradiating cells at 10 Gy with a statistically significant enhancement after 4 hours

of RT (Fig. 2D) and a further time-dependent increase up to 72 hours (Fig. 2E). Since increases in lipid

peroxidation can lead to ferroptosis, clonogenic survival was assessed using ferrostatin-1 (Ferr-1), a

selective inhibitor of ferroptosis. We found that Ferr-1 significantly enhanced clonogenic survival in

both cell lines following RT, confirming that RT induced ferroptosis contributes to the decreased

survival of these tumor cells (Fig. 2F).

Iron enhances RT sensitivity and ferroptosis

Since iron is central to lipid peroxidation, increasing iron levels by adding ammonium iron (II) sulfate

also increased lipid peroxidation significantly after 10 Gy RT in both A549 (Fig. 3A) and MDA-MB-

231 (Fig. 3B). The treatment of Ferr-1 abrogated the increased lipid peroxidation induced by iron. These

results were supported by decreased clonogenic survival upon ammonium iron (II) sulfate treatment.

The radiosensitizing effect of iron was rescued by Ferr-1 treatment, suggesting that iron

supplementation indeed leads to RT-induced ferroptosis (Fig. 3C and 3D). Consistent with a previous

report (9), iron depletion using deferoxamine (DFO), an iron chelator, also reduced lipid peroxidation

(Fig. 3E and 3F) and RT-induced tumor cell death (Fig. 3G and 3H). Overall, these data highlight the

critical role that iron plays in tumor cell survival and ferroptosis in response to RT.

FLASH RT induces ferroptosis in tumor cells

FLASH, an ultra-high dose rate RT, has been shown to result in less lipid peroxidation in micelles and

liposomes (20). To investigate whether FLASH RT also reduces lipid peroxidation in tumor cells, A549

and MDA-MB-231 were treated with electrons at conventional (6 MeV, 0.1 Gy/s) or FLASH RT (6

MeV, > 2 kGy/s) dose rates. The RT dose used in these experiments was 10 Gy, as the FLASH effect

is larger at higher RT doses (≥ 10 Gy) (13, 14, 41, 42). Interestingly, FLASH RT induced lipid

peroxidation in both cancer cell lines, and it was comparable to levels found with conventional RT (Fig.

4A and 4B). Furthermore, tumor cells treated with doses of RT ranging from 5 to 20 Gy exhibited a

dose-dependent increase in lipid peroxidation both by FLASH RT and conventional RT dose rates (Fig,

4C and 4D). Clonogenic assays using Ferr-1 confirmed that Ferr-1 increased cell clonogenicity in both

conventional and FLASH RT treated cells, indicating that ferroptosis contributes to cell death after

conventional and FLASH RT in cancer cells (Fig. 4E and 4F). In addition to dose rate differences, we

also investigated how different RT sources affected RT-induced lipid peroxidation using alpha particles

and proton (Fig. S2), indicating that various RT modalities increased lipid peroxidation.

FLASH RT induces lipid peroxidation in tumors but not in normal tissues

The tumor microenvironment consists of complex interactions between the tumor and its surrounding

counterparts, including fibroblasts, blood vessels, and low oxygen levels (43). To determine whether

the effect of FLASH RT on lipid peroxidation could be detected *in vivo*, tumor tissues were collected

from mouse xenograft models of two different human lung cancer cell lines (H460 and Calu6) for

staining with 4-hydroxynonenal (4-HNE), a classical marker for lipid peroxidation. Consistent with the

in vitro observations, both conventional and FLASH RT (15 Gy or 20 Gy) significantly increased 4-

HNE staining in these tumor tissues (Fig. 5A).

In contrast to tumors, we did not observe similar effects between conventional and FLASH RT in

normal mouse lung tissues, supporting previous observation of reduced lipid peroxidation (20, 22).

Conventional RT treatment of the whole thorax of BALB/c mice at 10 Gy significantly increased lipid

peroxidation in lung tissues at 24 hours and 7 days post-treatment. In contrast, the same dose of FLASH

RT resulted in markedly decreased lipid peroxidation compared to conventional RT across all tested

conditions. (Fig. 5B). These data suggest that reduced lipid peroxidation levels induced by FLASH RT

in normal tissue compared to tumors may contribute to the reduced normal tissue damage observed with

FLASH RT.

Increasing iron levels in normal tissue abolishes FLASH sparing effect

To investigate why the reduction in lipid peroxidation observed with FLASH is not evident in tumour

tissues, we hypothesized that intrinsic differences in iron levels between normal and tumour tissues may

account for this variation. in lipid peroxidation found after FLASH RT compared to conventional RT

(21). To test our hypothesis, we increased iron levels specifically in the normal intestine of BALB/c

mice. The intestinal tissue is highly sensitive to RT-induced toxicities due to the presence of rapidly dividing cells, particularly in the crypts and endothelium. However, multiple studies have shown that FLASH RT significantly reduces intestinal damage compared to conventional RT (30, 44, 45). Since iron absorption primarily occurs in the small intestine, particularly in the duodenum and proximal jejunum, mice were fed a high-iron diet (5000 ppm iron) to elevate iron levels locally without disrupting systemic iron homeostasis (46). Iron accumulation in the small intestine was confirmed as early as 24 and 48 hours after initiating the high-iron diet (Fig. S3A).

To examine whether high iron levels at the time of RT affect RT responses, mice were fed with a high-iron diet (HI) for 24 hours and treated with RT at conventional or FLASH RT dose rates, then returned to the control diet (HI 24-hour) or remained on a high-iron diet for an additional 72 hours (HI 96-hour) to determine the effect of iron before or after the RT (Fig. 6A). Serum ferritin levels suggested that the iron diet did not cause any systemic changes in iron levels (Fig. S3B). Consistent with normal lung tissues, conventional RT at both 8 and 10 Gy increased lipid peroxidation (4-HNE), with less induction with FLASH RT in the control diet group (200 ppm iron) (Fig. 6B and 6C). Similarly, lipid peroxidation was not significantly different after FLASH RT in mice fed an HI for 24 hours compared to non-irradiated mice, while conventional RT still showed high induction. However, the HI diet before and after RT (HI 96-hour) abrogated the FLASH RT effect with a significant increase in lipid peroxidation relative to the non-irradiated group and showed no significant difference from the conventional RT group.

These results were further supported by quantifying the number of crypts after RT and FLASH RT (Fig. 6D and 6E). Consistent with previous reports (30, 44, 45), the conventional RT groups exhibited a significantly lower number of remaining crypts compared to the FLASH RT groups following treatment either at 8 or 10 Gy (8 Gy: 33.4 % Conv vs. 54 % FLASH, 10 Gy: 15.9 % Conv vs. 36 % FLASH). A high-iron diet for 24 hours (HI 24-hour) did not alter this trend of intestinal damage. Notably, in the HI 96-hour mice, the difference between conventional and FLASH RT groups was no longer significant. The prolonged high iron diet exposure resulted in a reduced number of surviving crypts even in the FLASH RT groups, eliminating the FLASH sparing effect observed in other conditions. Taking together,

our findings suggest that iron availability plays a crucial role in RT-induced lipid peroxidation and

contributes to the reduced normal tissue damage observed with FLASH RT.

Discussion

Iron is essential in cell survival and biogenesis, as well as in rapidly proliferating tumor cells (2).

However, if homeostatic levels of iron become excessive, cells will die due to ROS production through

the Fenton reaction. In ferroptosis, iron is crucial as it initiates cell death by facilitating ROS production,

leading to lipid peroxidation (6). The present study evidenced the crucial role of iron in cancer cells and

their dependency on iron transport for survival. We also confirmed that excess iron increased the

radiosensitization of tumor cells through ferroptosis, while iron chelation protected against RT-induced

ferroptosis. Given our observation of significantly elevated iron levels in lung and breast cancer tumors

compared to normal tissues, we hypothesized that baseline tissue iron levels may critically influence

the reduction of normal tissue toxicity following FLASH radiotherapy.

Approximately 50% of cancer patients undergo RT during their cancer treatment. Advances in RT

techniques have resulted in more precise and focused treatments (47, 48). However, the effectiveness

of RT is often limited by the radioresistance of tumors and normal tissue toxicities, particularly for

tumors located in sensitive areas such as the abdomen or bone marrow. FLASH RT, introduced in 2014

by Fauvadon *et al*. shows promise in enhancing the therapeutic window of RT by decreasing normal

tissue toxicity while maintaining tumor control (49). Studies by multiple research groups have further

validated FLASH effects using various *in vitro* and *in vivo* models (50). Moreover, the successful

application of FLASH dose rates to cancer patients in several clinical trials is promising for its

translation into the clinic (51-54).

The most widely proposed mechanism for decreased FLASH RT normal tissue toxicity is oxygen

depletion (11-14, 50). Extensive *in vitro* and *in vivo* studies have shown that the sparing effect seen at

ultra-high dose rates varies with variable oxygen, supporting the crucial role of oxygen in response to

FLASH RT (13-16, 55). However, some of these studies also indicated that FLASH-induced hypoxia

alone was not sufficient to achieve normal tissue sparing (15, 16). To fully account for decreased normal

tissue toxicity, multiple studies have proposed other potential FLASH effects, including radical-radical

recombination, decreased DNA damage, conservation of the stem cell niche, and immune responses

(14, 17-19, 44, 45, 56-60). While these studies suggest that the FLASH effect may arise from the

complex interactions of chemical and biological responses within the tissue microenvironment, the

underlying cause of the differential effects on tumor and normal tissues remains incompletely elucidated.

In this study, we demonstrate that FLASH RT induced lipid peroxidation and ferroptosis, an iron

dependent regulated cell death pathway, based on our *in vitro* clonogenic survival (6, 21). Furthermore,

we showed that these effects are attributed to differences in iron levels in normal and tumor tissues.

Supporting our hypothesis, a previous study by Froidevaux *et al*. showed that FLASH RT did not

induce lipid peroxidation, an essential step for ferroptosis, compared to conventional RT, using micelles

and liposomes (20). However, this study did not differentiate FLASH effects between tumor vs normal

tissues, and did not account for the complexity of the tissue microenvironment. In the present study, we

compared the effects of FLASH RT on lipid peroxidation *in vitro* and *in vivo* utilizing several cancer

cells, xenograft tumors, and normal tissues. In cancer cells and tumors, we found increased lipid

peroxidation compared to non-irradiated controls after conventional and FLASH RT. Moreover, this

increase in lipid peroxidation corresponded to an increase in ferroptosis *in vitro*. In contrast, we

observed significantly reduced lipid peroxidation in normal tissues after FLASH RT compared to

conventional RT, in line with the previous observations regarding FLASH effects on lipid peroxidation

(20, 57). Interestingly, we also demonstrated that a variety of RT types (including protons and alpha

particles) are capable of inducing lipid peroxidation in cancer cells despite qualitative and quantitative

differences in DNA damage they cause. Since the FLASH effect has been seen in studies using proton

beams as well as heavier ion beams (helium and carbon beams), our data further support the role of

lipid peroxidation for the FLASH effect (16, 45, 58, 61-67).

To test our hypothesis regarding the role of tissue iron levels in the FLASH effect, we selectively

increased iron levels in the intestines without altering systemic iron homeostasis. In the control diet

group, the normal tissue protection by FLASH RT was evidenced by decreased lipid peroxidation and

more regenerating crypts compared to conventional RT. Interestingly, elevating iron levels only at the

time of RT was not sufficient to reverse the FLASH effect, since rapid iron turnover immediately

decreased normal tissue iron levels after withdrawal of the high-iron diet. However, when high iron

levels were sustained for another 72 hours after IR, the FLASH effect was abolished in the upper

intestines, indicating that sufficient iron availability during and after RT is necessary to induce lipid

peroxidation leading to tissue damage, which reflects underlying biological processes.

The kinetic model proposed by Spitz *et al*. suggests that the limited Fenton type reaction in normal

tissues compared to tumors contributes to the FLASH effect (68). Our study supports this model by

demonstrating the variation in intrinsic iron levels between tumors and normal tissues and highlighting

the necessity of the prolonged iron presence for FLASH-induced damage. Future studies exploring the

benefits or disadvantages of FLASH effects on iron-rich tissues (such as the liver or bone marrow),

focusing on lipid peroxidation and ferroptosis, will be required. In conclusion, the present study

highlights that iron, lipid peroxidation and ferroptosis contribute to the FLASH effect, furthering our

understanding of the mechanisms underlying reduced toxicity in normal tissue after FLASH RT.

Moreover, our study could also be used as a rationale as to which tissues or organs would benefit from

FLASH RT.

Acknowledgements and funding resources

We acknowledge our use of the gene set enrichment analysis, GSEA software, and Molecular Signature

Database (MSigDB) (Subramanian, Tamayo, et al. (2005), PNAS 102, 15545-

15550, http://www.broad.mit.edu/gsea/). Fig. 5A was created in https://BioRender.com. This work was

supported by Medical Research Council (MRC) Program grant (MC_UU_00001/11), Ministry of

Health and Welfare in Korea (RS-2023-00266627), National Institute of Health PO1 grant

(5P01CA257904-03), Gray Trust (AN9050), Cancer Research UK—RadNet (C6078/A28736),

MRC (MR/X006611/1), PhD students Study Abroad Program, National Science and Technology

Council (NSTC), Taiwan (114-2917-I-038 -001), and John Fell Fund (AND00180).

Author contributions

Study Conception & Design: N.V.L., M.H., C.C., S.L., K.P., and E.J.M.; Performed Experiment and

Data Collection: N.V.L., J.K, G.N., I.D.C.T., S.P., J.R, Y.J., S.P., P.J.L., T.H., E.W., A.A, M.H., C.C.,

C.K.T., K.P., and E.J.M.; Performed Data Analysis: N.V.L., J.K, G.N., S.P., P.J.L., T.H., E.W., M.H.,

C.C., and E.J.M.; Interpretation of data analysis: N.V.L., J.K, M.H., C.C., S.C.S., I.J.L., Y.T.J., A.J.G.,

K.P., and E.J.M. ;Writing the first draft: N.V.L. and E.J.M.; Figures Design: N.V.L., J.K., and E.J.M.

Supervision: A.J.G., K.P., and E.J.M; Funding acquisition: I.J.L., S.C.S., A.J.G., K.P., and E.J.M. All

authors contributed to manuscript editing and revision

Declaration of Interests

The authors have declared that no conflict of interest exists.

References

1.   Federico G, Carrillo F, Dapporto F, Chiariello M, Santoro M, Bellelli R, et al. NCOA4 links

iron bioavailability to DNA metabolism. Cell Rep. 2022;40(7):111207.

2.   Torti SV, Manz DH, Paul BT, Blanchette-Farra N, Torti FM. Iron and Cancer. Annu Rev

Nutr. 2018;38:97-125.

3.   Hann HW, Stahlhut MW, Menduke H. Iron enhances tumor growth. Observation on

spontaneous mammary tumors in mice. Cancer. 1991;68(11):2407-10.

4.   Radulescu S, Brookes MJ, Salgueiro P, Ridgway RA, McGhee E, Anderson K, et al. Luminal

iron levels govern intestinal tumorigenesis after Apc loss in vivo. Cell Rep. 2012;2(2):270-82.

5.   Sukiennicki GM, Marciniak W, Muszynska M, Baszuk P, Gupta S, Bialkowska K, et al. Iron

levels, genes involved in iron metabolism and antioxidative processes and lung cancer incidence.

PLoS One. 2019;14(1):e0208610.

6.   Dixon SJ, Lemberg KM, Lamprecht MR, Skouta R, Zaitsev EM, Gleason CE, et al.

Ferroptosis: an iron-dependent form of nonapoptotic cell death. Cell. 2012;149(5):1060-72.

7.   Lei G, Zhang Y, Koppula P, Liu X, Zhang J, Lin SH, et al. The role of ferroptosis in ionizing

radiation-induced cell death and tumor suppression. Cell Res. 2020;30(2):146-62.

8.   Lang X, Green MD, Wang W, Yu J, Choi JE, Jiang L, et al. Radiotherapy and

Immunotherapy Promote Tumoral Lipid Oxidation and Ferroptosis via Synergistic Repression of

SLC7A11. Cancer Discov. 2019;9(12):1673-85.

9.   Ye LF, Chaudhary KR, Zandkarimi F, Harken AD, Kinslow CJ, Upadhyayula PS, et al.

Radiation-Induced Lipid Peroxidation Triggers Ferroptosis and Synergizes with Ferroptosis Inducers.

ACS Chem Biol. 2020;15(2):469-84.

10.   E. Hall AG. Radiation Biology for Radiologist. 8th Edition ed: Wolters Kluwer Health; 2018.

11.   Wilson JD, Hammond EM, Higgins GS, Petersson K. Ultra-High Dose Rate (FLASH)

Radiotherapy: Silver Bullet or Fool's Gold? Front Oncol. 2019;9:1563.

12.   Moon EJ, Petersson K, Olcina MM. The importance of hypoxia in radiotherapy for the

immune response, metastatic potential and FLASH-RT. Int J Radiat Biol. 2022;98(3):439-51.

13. Adrian G, Konradsson E, Lempart M, Back S, Ceberg C, Petersson K. The FLASH effect

depends on oxygen concentration. Br J Radiol. 2020;93(1106):20190702.

14. Montay-Gruel P, Acharya MM, Petersson K, Alikhani L, Yakkala C, Allen BD, et al. Long-

term neurocognitive benefits of FLASH radiotherapy driven by reduced reactive oxygen species. Proc

Natl Acad Sci U S A. 2019;116(22):10943-51.

15. Cao X, Zhang R, Esipova TV, Allu SR, Ashraf R, Rahman M, et al. Quantification of Oxygen

Depletion During FLASH Irradiation In Vitro and In Vivo. Int J Radiat Oncol Biol Phys. 2021.

16. El Khatib M, Van Slyke AL, Velalopoulou A, Kim MM, Shoniyozov K, Allu SR, et al.

Ultrafast Tracking of Oxygen Dynamics During Proton FLASH. Int J Radiat Oncol Biol Phys.

2022;113(3):624-34.

17. Jin JY, Gu A, Wang W, Oleinick NL, Machtay M, Spring Kong FM. Ultra-high dose rate

effect on circulating immune cells: A potential mechanism for FLASH effect? Radiother Oncol.

2020;149:55-62.

18. Zhu H, Xie D, Wang Y, Huang R, Chen X, Yang Y, et al. Comparison of intratumor and local

immune response between MV X-ray FLASH and conventional radiotherapies. Clin Transl Radiat

Oncol. 2023;38:138-46.

19. Kim YE, Gwak SH, Hong BJ, Oh JM, Choi HS, Kim MS, et al. Effects of Ultra-high doserate

FLASH Irradiation on the Tumor Microenvironment in Lewis Lung Carcinoma: Role of Myosin Light

Chain. Int J Radiat Oncol Biol Phys. 2021;109(5):1440-53.

20. Pascal Froidevaux VG, Claude Bailat, Walter Reiner Geyer, François Bochud, Marie-

Catherine Vozenin. FLASH irradiation does not induce lipid peroxidation in lipids micelles and

liposomes. Radiation Physics and Chemistry. 2023;205.

21. Vilaplana-Lopera N, Abu-Halawa A, Walker E, Kim J, Moon EJ. Ferroptosis, a key to

unravel the enigma of the FLASH effect? Br J Radiol. 2022;95(1140):20220825.

22. Grilj V, Paisley R, Sprengers K, Geyer WR, Bailat C, Bochud F, et al. Average dose rate is

the primary determinant of lipid peroxidation in liposome membranes exposed to pulsed electron

FLASH beam. Radiation Physics and Chemistry. 2024;222.

23.   Moon EJ, Mello SS, Li CG, Chi JT, Thakkar K, Kirkland JG, et al. The HIF target MAFF

promotes tumor invasion and metastasis through IL11 and STAT3 signaling. Nat Commun.

2021;12(1):4308.

24.   Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr: interactive and

collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics. 2013;14:128.

25.   Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a

comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res.

2016;44(W1):W90-7.

26.   Xie Z, Bailey A, Kuleshov MV, Clarke DJB, Evangelista JE, Jenkins SL, et al. Gene Set

Knowledge Discovery with Enrichr. Curr Protoc. 2021;1(3):e90.

27.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set

enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.

Proc Natl Acad Sci U S A. 2005;102(43):15545-50.

28.   Tracy BL, Stevens DL, Goodhead DT, Hill MA. Variation in RBE for Survival of V79-4

Cells as a Function of Alpha-Particle (Helium Ion) Energy. Radiat Res. 2015;184(1):33-45.

29.   Berne A, Petersson K, Tullis IDC, Newman RG, Vojnovic B. Monitoring electron energies

during FLASH irradiations. Phys Med Biol. 2021;66(4):045015.

30.   Ruan JL, Lee C, Wouters S, Tullis IDC, Verslegers M, Mysara M, et al. Irradiation at Ultra-

High (FLASH) Dose Rates Reduces Acute Normal Tissue Toxicity in the Mouse Gastrointestinal

System. Int J Radiat Oncol Biol Phys. 2021;111(5):1250-61.

31.   Petersson K, Jaccard M, Germond JF, Buchillier T, Bochud F, Bourhis J, et al. High dose-per-

pulse electron beam dosimetry - A model to correct for the ion recombination in the Advanced

Markus ionization chamber. Med Phys. 2017;44(3):1157-67.

32.   Vojnovic B, Tullis IDC, Newman RG, Petersson K. Monitoring beam charge during FLASH

irradiations. Frontiers in Physics. 2023;11.

33.   Choi C, Lee GH, Son A, Yoo GS, Yu JI, Park HC. Downregulation of Mcl-1 by Panobinostat

Potentiates Proton Beam Therapy in Hepatocellular Carcinoma Cells. Cells. 2021;10(3).

34.   Franken NA, Rodermond HM, Stap J, Haveman J, van Bree C. Clonogenic assay of cells in

vitro. Nat Protoc. 2006;1(5):2315-9.

35.   Sondka Z, Bamford S, Cole CG, Ward SA, Dunham I, Forbes SA. The COSMIC Cancer

Gene Census: describing genetic dysfunction across all human cancers. Nat Rev Cancer.

2018;18(11):696-705.

36.   Huang DW, Sherman BT, Tan Q, Kir J, Liu D, Bryant D, et al. DAVID Bioinformatics

Resources: expanded annotation database and novel algorithms to better extract biology from large

gene lists. Nucleic Acids Res. 2007;35(Web Server issue):W169-75.

37.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer

genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer

Discov. 2012;2(5):401-4.

38.   Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Ponce-Rodriguez I,

Chakravarthi B, et al. UALCAN: A Portal for Facilitating Tumor Subgroup Gene Expression and

Survival Analyses. Neoplasia. 2017;19(8):649-58.

39.   Lakhal-Littleton S, Wolna M, Chung YJ, Christian HC, Heather LC, Brescia M, et al. An

essential cell-autonomous role for hepcidin in cardiac iron homeostasis. Elife. 2016;5.

40.   Jiang Y, Martin J, Alkadhimi M, Shigemori K, Kinchesh P, Gilchrist S, et al. Olaparib

increases the therapeutic index of hemithoracic irradiation compared with hemithoracic irradiation

alone in a mouse lung cancer model. Br J Cancer. 2021;124(11):1809-19.

41.   Vozenin MC, Hendry JH, Limoli CL. Biological Benefits of Ultra-high Dose Rate FLASH

Radiotherapy: Sleeping Beauty Awoken. Clin Oncol (R Coll Radiol). 2019;31(7):407-15.

42.   Adrian G, Konradsson E, Beyer S, Wittrup A, Butterworth KT, McMahon SJ, et al. Cancer

Cells Can Exhibit a Sparing FLASH Effect at Low Doses Under Normoxic In Vitro-Conditions. Front

Oncol. 2021;11:686142.

43.   Anderson NM, Simon MC. The tumor microenvironment. Curr Biol. 2020;30(16):R921-R5.

44.   Levy K, Natarajan S, Wang J, Chow S, Eggold JT, Loo PE, et al. Abdominal FLASH

irradiation reduces radiation-induced gastrointestinal toxicity for the treatment of ovarian cancer in

mice. Sci Rep. 2020;10(1):21600.

45.   Kim MM, Verginadis, II, Goia D, Haertter A, Shoniyozov K, Zou W, et al. Comparison of

FLASH Proton Entrance and the Spread-Out Bragg Peak Dose Regions in the Sparing of Mouse

Intestinal Crypts and in a Pancreatic Tumor Model. Cancers (Basel). 2021;13(16).

46.   J. F. Collins SRLF, X. Wang, G. J. Anderson. Mechanisms and Regulation of Intestinal Iron

Transport. In: Said HM, editor. Physiology of the Gastrointestinal Tract. 6th edition ed: Academic

Press; 2018. p. 1451-83.

47.   Bernier J, Hall EJ, Giaccia A. Radiation oncology: a century of achievements. Nat Rev

Cancer. 2004;4(9):737-47.

48.   Delaney G, Jacob S, Featherstone C, Barton M. The role of radiotherapy in cancer treatment:

estimating optimal utilization from a review of evidence-based clinical guidelines. Cancer.

2005;104(6):1129-37.

49.   Favaudon V, Caplier L, Monceau V, Pouzoulet F, Sayarath M, Fouillade C, et al. Ultrahigh

dose-rate FLASH irradiation increases the differential response between normal and tumor tissue in

mice. Sci Transl Med. 2014;6(245):245ra93.

50.   Limoli CL, Vozenin MC. Reinventing Radiobiology in the Light of FLASH Radiotherapy.

Annu Rev Cancer Biol. 2023;7:1-21.

51.   Kinj R, Gaide O, Jeanneret-Sozzi W, Dafni U, Viguet-Carrin S, Sagittario E, et al.

Randomized phase II selection trial of FLASH and conventional radiotherapy for patients with

localized cutaneous squamous cell carcinoma or basal cell carcinoma: A study protocol. Clin Transl

Radiat Oncol. 2024;45:100743.

52.   Mascia AE, Daugherty EC, Zhang Y, Lee E, Xiao Z, Sertorio M, et al. Proton FLASH

Radiotherapy for the Treatment of Symptomatic Bone Metastases: The FAST-01 Nonrandomized

Trial. JAMA Oncol. 2023;9(1):62-9.

53.   Bourhis J, Sozzi WJ, Jorge PG, Gaide O, Bailat C, Duclos F, et al. Treatment of a first patient

with FLASH-radiotherapy. Radiother Oncol. 2019;139:18-22.

54.   Gaide O, Herrera F, Jeanneret Sozzi W, Goncalves Jorge P, Kinj R, Bailat C, et al.

Comparison of ultra-high versus conventional dose rate radiotherapy in a patient with cutaneous

lymphoma. Radiother Oncol. 2022;174:87-91.

55.   Ha B, Liang K, Liu C, Melemenidis S, Manjappa R, Viswanathan V, et al. Real-time optical

oximetry during FLASH radiotherapy using a phosphorescent nanoprobe. Radiother Oncol.

2022;176:239-43.

56.   Eggold JT, Chow S, Melemenidis S, Wang J, Natarajan S, Loo PE, et al. Abdominopelvic

FLASH Irradiation Improves PD-1 Immune Checkpoint Inhibition in Preclinical Models of Ovarian

Cancer. Mol Cancer Ther. 2022;21(2):371-81.

57.   Zhu H, Xie D, Yang Y, Huang S, Gao X, Peng Y, et al. Radioprotective effect of X-ray

abdominal FLASH irradiation: Adaptation to oxidative damage and inflammatory response may be

benefiting factors. Med Phys. 2022;49(7):4812-22.

58.   Velalopoulou A, Karagounis IV, Cramer GM, Kim MM, Skoufos G, Goia D, et al. FLASH

Proton Radiotherapy Spares Normal Epithelial and Mesenchymal Tissues While Preserving Sarcoma

Response. Cancer Res. 2021;81(18):4808-21.

59.   Fouillade C, Curras-Alonso S, Giuranno L, Quelennec E, Heinrich S, Bonnet-Boissinot S, et

al. FLASH Irradiation Spares Lung Progenitor Cells and Limits the Incidence of Radio-induced

Senescence. Clin Cancer Res. 2020;26(6):1497-506.

60.   Labarbe R, Hotoiu L, Barbier J, Favaudon V. A physicochemical model of reaction kinetics

supports peroxyl radical recombination as the main determinant of the FLASH effect. Radiother

Oncol. 2020;153:303-10.

61.   Kristensen L, Poulsen PR, Kanouta E, Rohrer S, Ankjaergaard C, Andersen CE, et al. Spread-

out Bragg peak FLASH: quantifying normal tissue toxicity in a murine model. Front Oncol.

2024;14:1427667.

62.   Daugherty EC, Zhang Y, Xiao Z, Mascia AE, Sertorio M, Woo J, et al. FLASH radiotherapy

for the treatment of symptomatic bone metastases in the thorax (FAST-02): protocol for a prospective

study of a novel radiotherapy approach. Radiat Oncol. 2024;19(1):34.

63.   Ni H, Reitman ZJ, Zou W, Akhtar MN, Paul R, Huang M, et al. FLASH radiation reprograms

lipid metabolism and macrophage immunity and sensitizes medulloblastoma to CAR-T cell therapy.

Nat Cancer. 2025.

64. Chowdhury P, Velalopoulou A, Verginadis, II, Morcos G, Loo PE, Kim MM, et al. Proton

FLASH Radiotherapy Ameliorates Radiation-induced Salivary Gland Dysfunction and Oral Mucositis

and Increases Survival in a Mouse Model of Head and Neck Cancer. Mol Cancer Ther.

2024;23(6):877-89.

65. Verginadis, II, Velalopoulou A, Kim MM, Kim K, Paraskevaidis I, Bell B, et al. FLASH

proton reirradiation, with or without hypofractionation, reduces chronic toxicity in the normal murine

intestine, skin, and bone. Radiother Oncol. 2025;205:110744.

66. Tinganelli W, Weber U, Puspitasari A, Simoniello P, Abdollahi A, Oppermann J, et al.

FLASH with carbon ions: Tumor control, normal tissue sparing, and distal metastasis in a mouse

osteosarcoma model. Radiother Oncol. 2022;175:185-90.

67. Dokic I, Moustafa M, Tessonnier T, Meister S, Ciamarone F, Akbarpour M, et al. Ultra-High

Dose Rate Helium Ion Beams: Minimizing Brain Tissue Damage while Preserving Tumor Control.

Mol Cancer Ther. 2024.

68. Spitz DR, Buettner GR, Petronek MS, St-Aubin JJ, Flynn RT, Waldron TJ, et al. An

integrated physico-chemical approach for explaining the differential impact of FLASH versus

conventional dose rate irradiation on cancer and normal tissue responses. Radiother Oncol.

2019;139:23-7.

Figure legends

Fig 1. Iron is essential for tumor survival

(A-B) Tissue microarray slides from breast and lung cancer patients were analyzed to measure iron levels using Prussian blue. Compared to normal lung (A) or breast tissues (B), iron levels were significantly higher in lung and breast cancer tissues (Lung cancer; n = 20 for normal tissues, n = 32 for lung adenocarcinoma tissues, Breast cancer; n = 11 for normal tissues, n = 90 for cancer tissues). (C) The functional enrichment analysis of 748 genes from the Catalogue of Somatic Mutations in Cancer (COSMIC) Cancer Gene Consensus showed significantly regulated pathways of cancer driving genes "4 iron 4 sulfur cluster binding", "response to iron ion", and "cellular iron ion binding". (D) Analysis of TCGA PanCancer Atlas data revealed that cancer patients exhibiting higher *TFRC* expression in tumor tissues compared to normal tissues had significantly poorer overall survival outcomes (Low *TFRC*: n=2014, High *TFRC*: n=3257; log-rank test, $p < 0.0001$) (E) TCGA data showing *TFRC* mRNA expression in normal and various cancer patient tissues were visualized using UALCAN (F-G) The role of iron in tumor cell survival was determined by knocking down *TFRC*, an iron transporter. Compared to the siRNA against scrambled sequence, the inhibition of *TFRC* expression significantly decreased intracellular iron and cell survival measured by clonogenic assay in both A549 (F) and MDA-MB-231 (G) cells. Statistical tests were performed by nonparametric Kolmogorov-Smirnov test (A-B), log-rank (C), and upaired *t*-test (F-G).

Fig 2. RT induces lipid peroxidation and ferroptosis in cancer cells in a dose and time-dependent manner. (A) RNA sequencing was performed after treating MDA-MB-231 cells with 0 or 10 Gy of RT. Pathways analysis by Enrichr identified that genes involved in the "Cell Cycle", "p53 signaling pathway", "DNA replication" and "Ferroptosis" were significantly altered by RT. (B) GSEA enrichment further confirmed significant alterations of genes involved in ferroptosis after RT. (C-E) Lipid peroxidation was measured using C11 BODIPY in A549 and MDA-MB-231 24 hours after RT at varying doses from 2 to 10 Gy (C) as well as at different time points after 10 Gy of RT (D-E). (F) Clonogenic survival was determined after RT with or without pre-treatment of 20 µM (A549, I) or 4 µM (MDA-MB-231, J) Ferr-1. RT was given using a [137]Cs irradiator. Error bars indicate standard

deviation (SD) (n = 3 per group). Statistical tests were performed by One-way ANOVA with Dunnett's

multiple comparisons test (C-E) and unpaired *t*-test (F).

Fig 3. Iron enhances RT sensitivity and ferroptosis.

The role of iron in lipid peroxidation and cell survival was determined by measuring lipid peroxidation

using C11 BODIPY and clonogenic assay, respectively. Adding a suboptimal dose of iron (50 μM and

5 μM of ammonium iron (II) sulfate in A549 and MDA-MB-231 respectively) increased lipid

peroxidation (A-B) and radiosensitization (C-D) in both cell lines. Treatment with Ferr-1 (20 μM and

4 μM in A549 and MDA-MB-231, respectively) decreased lipid peroxidation induced by RT (A-B) and

reversed radiosensitization (C-D) indicating that ferroptosis was further induced by external iron. (E-H)

Inhibition of iron availability using deferoxamine (DFO) (200 nM and 100 nM in A549 and MDA-MB-

231, respectively) decreased lipid peroxidation induced by RT (E-F) and increased radioresistance (G-

H) of A549 and MDA-MB-231 cells. (A-H, n = 3/group) Error bars indicate standard deviation (SD).

Two-way ANOVA with Tukey's multiple comparisons test (A-B), One-way ANOVA with Tukey's

multiple comparisons test (C-D, G-H), and unpaired *t*-test (E-F). For C-D and G-H, 6 Gy conditions

were compared.

Fig 4. FLASH RT induces similar levels of lipid peroxidation and ferroptosis with conventional

RT in cancer cells. (A-B) A549 (A) and MDA-MB-231 (B) cells were irradiated at 10 Gy with

conventional or FLASH dose rates with an electron linear accelerator. A significant increase in lipid

peroxidation was observed 24 hours after RT using C11 BODIPY in all treatment groups. (C-D) A

significant dose-dependent induction of lipid peroxidation was observed after conventional and FLASH

RT. (E-F) Clonogenic survival was determined after conventional or FLASH RT with or without pre-

treatment of 20 μM (A549) or 4 μM (MDA-MB-231) Ferr-1. Error bars indicate standard deviation

(SD) (n = 3 per group). Statistical tests were performed by One-way ANOVA with Dunnett's multiple

comparisons test (A-B), Two-way ANOVA with Tukey's multiple comparisons test (C-D), and One-

way ANOVA with Tukey's multiple comparisons test (E-F).

F**ig 5. FLASH RT induces lipid peroxidation in mouse xenograft tumors while sparing normal**

lung tissues. (A-B) Lung tumor tissues (H460 and Calu6) from subcutaneous mouse xenograft models

were stained with 4-HNE, a lipid peroxidation marker. After conventional and FLASH RT (15 Gy for

H460 and 20 Gy for Calu6), there was a significant increase in 4-HNE staining intensity (n = 4 for H460,

n = 6 for Calu6 0 Gy and FLASH RT, n = 8 for Calu6 conventional RT). (C-D) BALB/c mouse lung

tissues were stained with 4-HNE after 10 Gy of conventional or FLASH RT. Lipid peroxidation was

significantly enhanced 24 hours and 7 days after conventional RT. However, FLASH RT did not change

lipid peroxidation (n = 6 per group at each time point). Error bars indicate standard deviation (SD).

Statistical tests were performed by One-way ANOVA (A-B) or Two-way ANOVA (C-D) with Tukey's

multiple comparisons test.

Fig 6. Increasing iron levels in normal tissue reverts FLASH sparing effect. (A) BALB/c mice were

fed with high iron diet (5000 ppm iron) for 24 hours to increase iron levels in normal intestine. After

RT treatment at 8 or 10 Gy, mice were returned to a normal diet or stayed on a high-iron diet for another

72 hours. (HI: High iron diet) (B-C) Lipid peroxidation, which was stained with 4-HNE was highly

increased by both 8 (B) and 10 Gy (C) conventional RT in the control or high iron diet groups both for

24 hours and 96 hours. FLASH RT at 8 or 10 Gy did not increase lipid peroxidation as significantly as

conventional RT with the control or 24-hour high-iron diet. However, a prolonged high-iron diet even

after RT treatment eliminated the FLASH effect on lipid peroxidation. (D-E) Tissue damage in the

upper intestines was determined by H&E staining. The presence of remaining intestinal crypts (arrow

heads) was counted and analyzed. The number of remaining crypts was significantly decreased after 8

(D) and 10 Gy of conventional RT in the control or high-iron diet groups for 24 hours and 96 hours.

FLASH RT-induced damages were significantly lower than conventional RT in the control or 24-hour

high-iron diet mice. However, a prolonged high iron diet, diminished the sparing observed in FLASH

irradiated tissues since FLASH RT induced similar tissue damage to conventional RT. (B - E, for 8 Gy,

n = 8 in every condition; for 10 Gy, control diet n = 12, HI diet 24 hours or 96 hours n = 6) Error bars

indicate standard deviation (SD). Statistical test was performed by Two-way ANOVA with Tukey's

multiple comparisons.

Fig. 1

(A) Lung cancer

(B) Breast cancer

(C) 748 genes from
The Cancer Gene Consensus

GO Term	p-value
4 iron, 4 sulfur cluster binding	0.008
Response to iron ion	0.017
cellular iron ion homeostasis	0.03

(D)

TCGA PanCancer Atals

(E)

Expression of TFRC across TCGA cancers (with tumor and normal samples)

(F)

A549   A549

(G)

MDA-MB-231   MDA-MB-231

Fig. 2

Fig. 3

Fig. 4

(A) A549

(B) MDA-MB-231

(C) A549

(D) MDA-MB-231

(E) A549

(F) MDA-MB-231

Fig. 5

(A)

	Non-treated	Conventional	FLASH

H460 100μm

Calu6 100μm

H460

Lipid peroxidation (4-HNE)

p<0.0001
p=0.0056

0Gy   15Gy Conv   15Gy FLASH

Calu6

Lipid peroxidation (4-HNE)

p=0.019
p=0.0002

0Gy   20Gy Conv   20Gy FLASH

(B)

	Non-treated	10Gy Conv	10Gy FLASH

24h

7d 100μm

Lipid peroxidation (4-HNE)

ns
p=0.0114
p=0.0016

ns
p=0.0072
p<0.0001

☐ 0Gy
▨ 10Gy Conv
▨ 10Gy FLASH

24 hr   7 days

Fig. 6