IN-VIVO EVIDENCE FOR INCREASED TAU DEPOSITION IN TEMPORAL LOBE EPILEPSY

Raúl R. Cruces[1], Jack Lam[1], Thaera Arafat[1], Jessica Royer[1], Judy Chen[1], Ella Sahlas[1], Arielle Dascal[1], Daniel Mendelson[1], Raluca Pana[2], Robert Hopewell[1], Chris Hung-Hsin Hsiao[1], Gassan Massarweh[1], Jean-Paul Soucy[1], Pedro Rosa-Neto[1], Marie-Christine Guiot[2], Sylvia Villeneuve[3], Lorenzo Caciagli[4,5], Matthias J. Koepp[4], Andrea Bernasconi[1], Neda Bernasconi[1], Boris C. Bernhardt[1*].*

1. McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University, Montreal, Québec, Canada.
2. Montreal Neurological Institute, McGill University Health Centre (MUHC) and other McGill University
3. StoP-AD Centre, Douglas Mental Health University Institute, McGill University, Montréal, QC, Canada
4. UCL Queen Square Institute of Neurology, University College London, London WC1N 3BG, UK
5. Department of Neurology, Inselspital, Sleep-Wake-Epilepsy Center, Bern University Hospital, University of Bern, Bern, Switzerland

joint senior authors

Corresponding author
Raul R. Cruces, MD PhD
raul.rodriguez-cruces@mcgill.ca

Boris C. Bernhardt, PhD
boris.bernhardt@mcgill.ca

GLOSSARY:
TLE = temporal lobe epilepsy. **PET** = positron emission tomography. **MRI** = magnetic resonance imaging. **NI** = neuroinflammation

KEYWORDS:
Temporal lobe epilepsy | [^{18}F]MK-6240 | tauopathy | neuroimaging | neuroinflammation

SHORT TITLE:
Increased *in-vivo* tau in TLE

ABSTRACT

Temporal lobe epilepsy (TLE), the most common pharmaco-resistant epilepsy in adults, has

been linked to structural brain changes extending beyond the mesiotemporal areas. While not

traditionally viewed as a neurodegenerative disorder, recent *ex-vivo* studies have shown

elevated levels of misfolded tau protein in TLE. This study investigated tau deposition in TLE

patients using the *in-vivo* PET tracer [18F]MK-6240 combined with ex vivo tau

immunohistochemistry (IHC). We studied 28 TLE patients and 28 healthy controls to assess

tau uptake and its relationship with brain connectivity, clinical variables, and cognitive function

alongside post-surgical tissue from a subset of patients. Compared to controls, TLE patients

exhibited markedly increased [18F]MK-6240 uptake in bilateral superior and medial temporal

regions and the parietal cortex, with tau accumulation following regional functional and

structural connectivity and cognitive impairment. IHC revealed phosphorylated tau in 6/7

cases, including subpial, neuritic, and neuronal inclusions. These findings suggest that tau

accumulation contributes to cognitive decline observed in TLE, supporting a potential role of

tau in epilepsy-related neurodegeneration.

1 INTRODUCTION

Mounting evidence suggests that temporal lobe epilepsy (TLE), the most common pharmaco-
resistant epilepsy in adults, is not a static condition, but one that is associated with progressive
changes extending beyond the mesiotemporal disease epicenter[1–4]. Although the exact causes
of disease progression remain unknown, several lines of evidence suggest similar mechanisms
as in Alzheimer's disease (AD), an established neurodegenerative condition. First, cognitive
profiles in middle-aged pharmaco-resistant epilepsy patients resemble those of mild cognitive
impairment, which can be linked to AD[5]. Furthermore, epidemiological studies have indicated
an elevated risk for pharmaco-resistant epilepsy patients to be diagnosed with AD, and *vice*
versa[6,7]. Across all age groups, patients with AD show a 3-fold higher risk of suffering from
epilepsy[8,9]. Finally, a recent analysis of *ex-vivo* specimens in operated TLE patients has shown
highly elevated levels of misfolded tau protein in operated TLE patients[10], a hallmark of AD
pathology[11]. Such findings complement data from animal models of TLE, which demonstrate
that seizure activity triggers tau hyperphosphorylation and amyloidogenic pathways[12,13].

Whole-brain patterns of increases of phosphorylated tau in epilepsy remain unknown due to
limits imposed by the scope of resection. Also, while *ex-vivo* tau increases in TLE were related
to postoperative cognitive decline[14], correlations with measures of preoperative cognitive
function remain incompletely understood. These considerations underscore the need for *in-vivo*
analyses of tau in TLE to evaluate its potential role as both a modulator of cognitive function
and a predictor of clinical outcomes. Research in AD and other neurodegenerative conditions
has benefitted from the introduction of second-generation PET tracers that are sensitive and
specific to pathological deposits. In particular, the [^{18}F]-MK-6240 tracer (*henceforth* MK-
6240) has shown high affinity and selectivity to intracellular neurofibrillary tangles[15–17]. In
contrast, while there is an extensive PET literature in epilepsy on presurgical localization of
hypometabolism via [^{18}F]fluorodeoxyglucose[18,19], PET studies in epilepsy have so far not
systematically investigated tau tracers[7]. As such, the whole brain distribution of tau in epileptic
patients remains unknown. Unlike research in AD, where tau uptake patterns are increasingly
understood and contextualized with respect to patterns of brain connectivity[1–3,20–22], there have
not been any systematic assessments of potential network-level effects underlying the
accumulation of these pathological deposits in TLE.

The objective of our study was to investigate the distribution and prevalence of tau aggregates
in TLE using *in-vivo* MK-6240 PET. Tau mapping was complemented with network decoding
models informed by multimodal MRI investigations in the same participants that tested for
spatial associations of tau uptake in patients with patterns of global and local connectivity.
Moreover, linear models examined associations with cognitive function assessed outside the
scanner and clinical measures from the patients presurgical examination.

RESULTS

We recruited 28 TLE patients (14 Females, age=36±12) and 28 healthy volunteers (14 Females,

age=33±7). with no significant differences in age and sex distribution. The mean onset of

epilepsy in TLE patients was 21±14 years, the mean duration of epilepsy was 15.3±11.5 years,

and 29% had hippocampal atrophy (volume or asymmetry less than 1.5 standard deviations

from controls). Further clinical features are listed in **Table 1**. Analyses are referenced to seizure

onset and reported separately for the ipsilateral and contralateral hemispheres.

Widespread increases in tau uptake in TLE

Mean cortical MK-6240 SUVR was increased in patients with TLE relative to controls, both

ipsilateral (t=3.16, p=<0.002) and contralateral (t=3.25, p=0.002) to the seizure focus. The

individualized MK-6240 SUVR histogram analysis across the cortical surface of all subjects

(**Figure 1A**) shows that there is a high prevalence of TLE patients with high MK-6240 uptake,

which is reflected in the mean group uptake (**Figure 1B**). Indeed, 75% of TLE patients but no

controls demonstrated elevated mean SUVR above 1.5 (97.5th percentile based on controls,

Table 2), with a spatial distribution generally resembling the statistical effects. This pattern

was more pronounced in the ipsilateral hemisphere but showed a similar bilateral distribution

(**Supplementary Figure 1 and Table 2**).

Surface-wide analysis of the group difference of cortical MK-6240 SUVR indicated bilateral

high uptake in patients relative to controls, particularly in superior-posterior temporal and

parietal and occipital cortex (**Figure 1C**). Statistical comparisons were based on two-tailed

mixed-effects models controlling for age and sex, with significance assessed using random

field theory (p_{RFT}<0.025). The strongest effects were localized to the ipsilateral frontal

operculum and the posterior cortex, including the anterior and inferior parietal lobe (postcental,

supramarginal and angular gyrus) and the lateral temporal (superior temporal gyrus) and

occipital cortices. In the contralateral hemisphere, a similar pattern was observed, sparing the

occipital region, but with a larger involvement of the temporal lobe, encompassing the posterior

section of the superior and middle temporal gyri, and the premotor cortex.

In significant clusters derived from the main group- difference analysis, the overall effect was

similar in both hemispheres but slightly higher ipsilateral to seizure onset (ipsilateral: t=5.04,

p<0.001; contralateral: t=4.78, p<0.001, **Figure 1D**). No significant differences were observed

in the subcortical structures analyzed; however we found a trend of higher MK-6240 SUVR

uptake in the ipsilateral hippocampus (t=-2.00, uncorrected p =0.049, **Table 3**).

To assess whether MK-6240 increases were seen even after correcting for grey matter atrophy,

a common feature in TLE[23,24], we re-ran the mixed effects model controlling vertex-wise for

regional grey matter thickness estimates and our results were largely unaffected

(**Supplementary Figure 2A-B**). Additionally, we evaluated the robustness of our results using

different reference areas to calculate the MK-6240 SUVR and confirmed high consistency

(**Supplementary Figure 3**).

We evaluated the effects of sex on MK-6240 uptake to investigate potential sex-related

differences in tau pathology. Our findings indicated that females with TLE overall exhibited

an elevated Tau burden relative to their male counterparts (**Supplementary Figure 4**).

A subgroup of 7 healthy controls and 13 TLE patients had longitudinal tau PET data available
(mean difference between scan 1 and 2=18.75 months, minimum=12 months, maximum 42
months). While there was a non-significant trend for subtle increases in TLE in the ipsilateral
temporal lobe at uncorrected thresholds (**Supplementary Figure 2C**), MK-6240 uptake
showed no significant longitudinal change in either healthy controls or patients, and no group
differences in rate of change were observed (**Figure 1E**).

7
Figure 1. A) Subject-Wise distribution of cortical MK-6240 uptake. *Density histograms illustrate the distribution*
of global cortical MK-6240 uptake for individual participants in each group. Red squares indicate regions where
SUVR exceeds the abnormality threshold (SUVR > 97.5th percentile). Brain images represent individual
participants from each group with progressively increasing MK-6240 values. ***B) Group-wise mean tau MK-6240***
uptake. *Cortical phosphorylated tau deposits were assessed with MK-6240 and visualized as standardized uptake*
value ratios (SUVR). The average MK-6240 uptake is shown for healthy controls and TLE patients. ***C) Group***
differences in MK-6240 uptake. *Group comparisons between patients and controls were performed using a*
mixed-effects model, adjusting for age and sex, and corrected for multiple comparisons using random field theory

1   (*p*RFT *< 0.025*). *Significant regions are outlined in white.* **D) Mean SUVR values for significant regions** *obtained*
2   *from the group differences analysis. Violin plots are shown by hemisphere and group.* **E) Longitudinal**
3   **trajectories of MK-6240 uptake** *by group between the PET scans 1 and 2. Lines correspond to the longitudinal*
4   *participants, and transparent points are the cross-sectional subjects.*
5

Association with demographic, clinical, and cognitive measures

Neither age MK-6240 SUVR (r = −0.03, p = 0.80) nor disease duration were significantly
associated with MK-6240 SUVR in patients (r = −0.08, p = 0.68) in clusters of between-group
differences. At the surface level, MK-6240 SUVR was associated with age in the ipsilateral
medial-to-anterior cingulate gyrus and inferior frontal lobe, and with disease duration in the
contralateral inferior parietal region and the antero-inferior portion of the occipital lobe.
Notably, the latter duration association to tau remained consistent even after controlling for
age. Age at seizure onset, contralateral hippocampal volume, interictal epileptiform discharge
frequency and generalize tonic clonic seizure frequency were not correlated to MK-6240
SUVR, neither in clusters of between-group findings nor at the vertex-level (**Supplementary**
figure 5).

When studying associations to hippocampal volume, we noted that lower ipsilateral
hippocampal volume related to higher MK-6240 SUVR in patients in clusters of findings (r =
−0.28, p = 0.04; **Figure 2A-left**). Furthermore, lower ipsilateral hippocampal volumes were
associated with higher MK-6240 SUVR in the contralateral posterior superior parietal lobe
(**Figure 2A-right**).

We also studied associations between clusters of tau PET measures and cognitive variables,
notably scores on the EpiTrack[25], as well as scores on previously validated tasks tapping into
episodic and semantic memory[26] (**Figure 2B-left**). In line with prior results[27], our TLE patients
showed lower performance on the *EpiTrack* (r = -0.29, p = 0.03) and *episodic memory task* (r
= −0.32, p = 0.03), indicating an association between increased Tau burden and lower cognitive
performance. In contrast, their performance did not differ from controls for the *semantic*
memory task (*p*=0.09, p=0.57). At the surface level, cognitive performance measures had
distinct associations with MK-6240 SUVR, lower EpiTrack scores showed only a trend toward
higher MK-6240 SUVR, whereas reduced episodic memory performance exhibited a negative
association with MK-6240 SUVR (**Figure 2B-right**). This pattern involved the ipsilateral
angular gyrus and bilateral occipital lobes. No associations were observed for semantic
memory (**Supplementary figure 5**).

Figure 2. A) Tau MK-6240 SUVR and clinical relationships. *Scatterplots show an association of mean MK-6240 SUVR in regions that showed significant between-group differences (see Figure 1) with clinical variables. Corresponding cortical projections visualize the regional effects of these variables on MK-6240 SUVR. Dark red on the brain maps indicate regions where age and duration are associated with increased tau deposition, as measured by t-statistics.* **B) Tau MK-6240 SUVR and behavioral relationships.** *Scatterplots display the relationship of mean MK-6240 SUVR in regions of significant-between group differences with behavioral measures. Brain maps illustrate the corresponding regional effects, dark blue indicating areas where lower cognitive performance is associated with higher MK-6240 SUVR. Mixed-effects models were applied to assess the association between MK-6240 SUVR and these features, corrected for multiple comparisons with random field theory (p<0.025, two tails). Significant regions after multiple comparisons are outlined in white in the surface-based model and an asterisk (*) indicates significant results in the analyses.*

Additionally, we evaluate the impact of all available clinical features on MK-6240 uptake using
a linear mixed-effects model. No significant clinical predictor evaluated in the model impacted
MK-6240 uptake except for a larger patient–control difference in females than in males, and
an association between a higher number of antiseizure medications and lower MK-6240 SUVR.
We also assessed subcortical and hippocampal MK-6240 SUVR and did not observe any
significant between group differences (**Supplementary Figure 6, Supplementary Table 3**).
Moreover, we did not observe any significant correlations between subcortical MK-6240
SUVR and clinical or behavioral measures.
9

Association to network architecture

We assessed the susceptibility of globally highly interconnected regions to tauopathy using
group-based network decoding models (**Figure 3**). Vertex-wise measures of mean connectivity
and neighborhood-weighted connectivity informed by MK-6240 were derived from group-
average structural and functional connectomes and correlated with vertex-wise *t*-statistics
obtained from the group-difference mixed-effects model.

This analysis revealed a marginal association at trend levels between MK-6240 SUVR and
highly connected regions defined by mean functional connectivity ($r = 0.44$, $p_{spin} = 0.11$, **Figure**
3A-left). Associations with mean structural connectivity were non-significant ($r = 0.11$, $p_{spin} =$
0.29, **Figure 3B-left**).

In contrast, associations between tau uptake differences and neighborhood-weighted
connectivity showed strong correlations for both functional ($r = 0.51$, p_{spin} 0.02, **Figure 3A-**
right) and structural networks ($r = 0.52$, $p_{spin} < 0.001$, **Figure 3B-right**).

Figure 3. A) Relationship between MK-6240 group differences and functional connectivity. The left panel displays a scatterplot of mean node values from the functional connectome against between-group differences in MK-6240 tau uptake (see Figure 1). The right panel visualizes neighboring regions of each node determined by functional connectivity, with their association to tau between- group differences. B) Relationship between MK-6240 group differences and structural connectivity, derived from diffusion MRI tractography. The left panel shows a logarithmic scatter plot of mean node values against group differences. The right panel highlights neighboring regions based on structural connectivity, which show a strong trend of increased tau deposition. The central image depicts how functional (upper) and structural (lower) connectivity informed by MK-6240 group differences was used to generate neighbor-weighted MK-6240 estimates. Points are color-coded based on MK-6240 t-values, with higher t-values indicating greater tau deposits in TLE compared to controls. The red line represents non-linear trends in the data. These findings suggest that both functional and structural network properties significantly reflected the spatial distribution of MK-6240 increases in TLE, with network hubs and their connected regions being particularly vulnerable.

Direct effect of Tau on memory surpasses network-mediated influence

In a final integrative analysis, we evaluated the effects of MK-6240 on cognitive performance
and the potential modulation of the structural connectome using structural equation modeling
(SEM). In the SEM analysis (**Figure 4),** mean MK-6240 SUVR from significant clusters was
significantly associated with structural neighborhood-weighted connectivity (β = 0.38, p =
0.002), but not with structural connectome efficiency. Higher MK-6240 was directly and
significantly associated with lower EpiTrack performance (β = −0.49, p = 0.009) and lower
episodic memory performance (β = −0.55, p = 0.004), but not with semantic memory (β =
−0.14, p = 0.56). Paths from structural network measures to memory outcomes were not
significant, and none of the indirect (mediation) effects reached significance. The total effect
of MK-6240 on episodic memory was significant, indicating a predominantly direct
relationship rather than mediation.

13
14 ***Figure 4.*** *Structural equation model showing relationships between mean significant MK-6240, structural*
15 *network measures (SC weighted neighbors and SC efficiency), and behavioral outcomes. Values represent*
16 *standardized path coefficients. Significant paths (p < 0.05) are shown in blue for positive effects and orange for*
17 *negative effects, the rest is gray. Dotted lines represent covariance between features. Indirect effects via semantic*
18 *network measures were tested but were not significant.*

Heterogeneous AT8 Tau depositions in post-surgical TLE tissue
Neuropathological evaluation of AT8-immunoreactive phosphorylated tau (p-tau) revealed a
spectrum of variable p-tau deposition in 5/6 cases for which adequate specimen was available
(**Supplementary Table 5**). The most frequent finding was focal subpial immunoreactivity,
characterized by a thin layer of positivity at the cortical margin, which was particularly
prominent in cases diagnosed with focal cortical dysplasia (FCD Type II and III).
Intracytoplasmic neuronal inclusions were identified in several instances, most notably in a
case with a hippocampal cavernoma, where perilesional neurons exhibited definite
intracytoplasmic staining alongside strong "tuffed-like" pathology in adjacent areas. In a case
of FCD-II the morphology was more complex, presenting as strong neuritic staining within the
hippocampal CA4 region and focal, strong longitudinal striated patterns. One case had less
suitable specimen, but overall exhibited negative findings, with only negligible dot-like
staining at fragment edges (**Supplementary Figure 9**).
14

1 DISCUSSION

This study represents novel *in-vivo* evidence of elevated phosphorylated tau, measured by
[18F]MK-6240 PET, in young adults with TLE, and demonstrates association with markers of
network architecture as well as clinical and cognitive measures. Comparing patients to controls,
increased tau accumulation in TLE was primarily observed bilateral temporo-posterior regions,
and we could furthermore demonstrate that MK-6240 uptake was associated with measures of
functional and structural connectivity, indicating that highly interconnected network hubs and
their neighboring regions are particularly vulnerable to tau deposition. Notably, females with
TLE exhibited a higher tau burden than their male counterparts, suggesting potential sex-
related differences in tau pathology. Cognitive analyses showed that higher tau burden was
directly associated with lower performance on the Epitrack and impaired episodic memory.
Mediation analyses did not support indirect effects of structural network measures on cognitive
outcomes, indicating that the relationship between tau burden and memory impairment was
predominantly direct rather than mediated by network properties. Although longitudinal
changes were modest, these findings suggest that tau pathology may contribute to disease-
related network disruption and cognitive impairment in drug-resistant TLE. These findings
suggest that tau accumulation may contribute to cognitive decline, supporting the potential role
of tau in neurodegeneration in pharmaco-resistant epilepsy *in-vivo*.

While so far unexplored in epilepsy research, the [18F]MK-6240 PET tracer has been repeatedly
applied to map disease processes in established neurodegenerative conditions, in particular
AD[28–30]. In those assessments, MK-6240 has been shown to generally describe the typical
Braak pattern of pathological spread away from an entorhinal/mesiotemporal disease epicentre
to involve bilateral, large scale networks[31]. Several assessments in AD have also shown that
tau PET changes have associations with overall cognitive impairment and atrophy, both at the
time of scanning[32,33] and prospectively[34]. These findings have been complemented with several
histopathological and electron microscopic validation experiments[35], suggesting that MK-6240
does scale with the overall deposition of hyperphosphorylated tau. A key strength of the current
study is its focus on young and middle-aged individuals, an underexplored population in tau
imaging research. Despite the younger age of the cohort, we observed a robust increase in MK-
6240 binding in our patients. The observed pattern of tau deposition, involving the superior
and middle temporal regions, as well as parietal-occipital areas, aligns in part with earlier *ex-
vivo* observations of increased abnormalities in the ipsilateral temporal lobe relative to the
epileptogenic focus[3,10,36–38], and confirms recent CSF-biomarker findings suggesting increased
tau in TLE[39]. These results provide *in-vivo* evidence suggesting that TLE is associated with an
elevated risk of tau-associated pathophysiological processes[39,40]. Crucially, we observed robust
increases in tau deposition in TLE patients relative to controls after adjusting for age and sex,
with findings remaining significant after correction for multiple comparisons. Prevalence
mapping provided evidence for slightly greater tau burden ipsilateral to the seizure focus,
whereas group-level effects were largely bilateral. This pattern was not directly aligned with
ipsilateral temporo-limbic connectivity and may therefore reflect a more complex pathological
process that cannot be fully explained with respect to a unilateral mesiotemporal epicentre.

The current sample was composed out of equal proportions of females and males in both the patient and control cohorts. This allowed us to systematically assess interactions between group and sex, which pointed towards more marked between-group differences in females compared to males. Indeed, while both female and male patients demonstrated common increases in tau uptake in central and posterior divisions, female patients additionally displayed increases in tau compared to female controls in anterior midline, insular, and temporo-anterior divisions. In the context of patients diagnosed with AD and individuals with a familiar risk for it, sex effects have increasingly been recognized. This is also motivated by a twofold higher lifetime risk for developing AD in women, which cannot be explained by higher life expectancy alone[41]. Prior studies have indicated higher tau accumulation in both the brain and plasma in older females compared to males that are both cognitively normal and those that are at risk for developing AD. Increased tau in females is generally not accompanied by an elevated increase in amyloid, suggesting either independent mechanisms or an elevated effect of pre-existing amyloid on tau accumulation in females[42,43]. While the evidence for sex-differences in longitudinal tau effects remains inconclusive, there is work that overall suggests faster cognitive decline in females with significant tau accumulation compared to their male counterparts[44]. It has been suggested that sex hormones and sex chromosomes interact with various disease mechanisms during aging, affecting inflammatory, metabolic, and autophagy processes[45]. These complex interactions affect disease progression between the sexes in classical neurodegenerative conditions and may furthermore interact with disease related factors and reported sex differences in epilepsy[46].

Pathological validation of PET findings through post-operative immunohistochemistry is essential for confirming in-vivo observations. Post-surgical neuropathological analysis using the AT8 antibody revealed p-tau deposition in cases with cortical MK-6240 positivity, providing direct histopathological support for our PET findings. The presence of p-tau in drug-resistant epilepsy suggests a significant overlap between chronic seizure activity and neurodegenerative processes. Several studies have shown that p-tau accumulation correlates with duration of refractory epilepsy and cognitive decline[10,14,36,38], appearing across a diverse spectrum of surgical etiologies including FCD, hippocampal sclerosis, and acquired lesions[47]. Our findings align with these observations, particularly in FCD-II and III cases where p-tau was frequently identified in cortico-amygdalohippocampectomy resections.

An important consideration when interpreting both PET and histopathological findings is the structural heterogeneity of tau fibrils across tauopathies, as disease-specific conformations influence tracer binding properties. In ex-vivo studies, both 3R and 4R isoforms were confirmed in positive cases, yet many epilepsy cases show minimal or absent p-tau, suggesting age-related vulnerability and involvement of multiple distinct mechanisms. This variability underscores the complexity of tau pathology in epilepsy. While the PET signal most likely reflects overall increases in tau burden, findings may also co-localize with inflammatory cascades, given reports linking cortical atrophy patterns to microglial activation in TLE[48,49]. Growing evidence further suggests that neuroinflammation may accelerate tau phosphorylation and misfolding, creating a self-perpetuating cycle of neuronal dysfunction[50,51].

Collectively, our findings support the hypothesis that epilepsy may act as a "secondary tauopathy," where seizure-related metabolic stress activates protein aggregation pathways similar to primary neurodegenerative diseases, explaining p-tau detection even in younger

individuals[37]. The convergence of histopathological and global cortical MK-6240 PET findings highlights the heterogeneity of neuropathological mechanisms underlying TLE. Future studies clarifying tau conformation, post-translational modifications, and tracer binding properties will be essential for enhancing the biological specificity of these observations.

Previous PET studies in TLE have primarily focused on functional rather than pathological markers, with FDG-PET commonly used to map regional hypometabolism for seizure lateralization, surgical planning, and outcome prediction[52–54]. Cross-sectional studies have reported moderate associations between epilepsy duration and FDG-PET hypometabolism, mainly in temporal lobe regions[55,56]. In contrast, longitudinal FDG-PET evidence remains limited and inconclusive. A study of 16 adults with TLE undergoing repeated FDG-PET scans found no significant association between scan interval and the rate of change in regional FDG-PET uptake, although patients with longer disease duration showed greater evidence of progressive decline[57]. Similar findings have been reported in pediatric non-lesional epilepsy, where persistent seizures were associated with expansion of the hypometabolic cortex over time, whereas seizure reduction was linked to decreased hypometabolic extent[58].

Connectome decoding models have previously been applied to describe network effects in the expression of specific imaging phenotypes, in both neurodevelopmental and neurodegenerative conditions[20,59–62]. In our study, we explored global as well as local connectivity associations to patient-related MK-6240 increases. While the former approach aimed to test an overall susceptibility of highly interconnected hub regions to tauopathy, the latter assessed a more local association between tau uptake and a region's connectivity patterns. Notably, while TLE-related increases in tau showed only marginal associations with global functional connectivity, stronger associations emerged with local connectivity patterns, both structurally and functionally. These findings suggest a subtle vulnerability of polysynaptic functional hubs, alongside a more pronounced influence of local network architecture. The selective susceptibility of highly connected cortical hubs in TLE has been described in studies of atrophy, and our findings extend this vulnerability to tau pathology[63]. In neurodegeneration research, local connectivity has been linked to the mechanisms of regional propagation, in which tau spreads through anatomically connected networks to affect an increasing set of brain regions[31,64,65].

In clusters of between-group differences, we did not find an overall association between MK-6240 and either disease duration or age. However, surface-based analyses revealed that longer disease duration was associated with increased MK-6240 SUVR in contralateral parieto-occipital regions, while age was associated with increased SUVR in anterior ipsilateral inferior cingulate and inferior frontal regions. These spatially distributed effects may reflect a broader disease process involving interconnected networks rather than localized effects primarily limited to the ipsilateral temporal lobe[14]. Increasing the sample size with ideally longer longitudinal follow up times and an older average age of included patients may boost sensitivity to detect progressive tau uptake in TLE in future work and might help to clarify the temporal dynamics of tau accumulation and its potential role in neurodegenerative cascades associated with epilepsy. This expanded design help to identify clinically relevant subgroups that show the most marked progressive increases in tau, and to thus establish whether tau increases are

pre-existing or whether they cumulatively increase throughout the TLE disease course. In AD,
prior work has reported an increase in tau misfolding depositions over periods longer than five
years and in individuals older than 65 years[66,67]. We did not systematically genotype our
participants to explore potential associations between *in-vivo* tau uptake and genetic
background. A prior meta-analysis of 46 studies[68] found that the APOE ε4 allele[69] was
associated with increased epilepsy risk, earlier onset, worse cognition, and a history of febrile
convulsions, while the APOE ε2 allele was linked to reduced risk; APOE ε3 showed no
association[70]. The APOE gene was more strongly related to temporal lobe, drug-refractory, and
late-onset epilepsy, with no link between ε4 and psychiatric symptoms or seizure-free
outcomes after surgery[8].

In AD, increased tau burden is associated with reduced hippocampal volume, likely reflecting
neuronal loss driven by abnormal tau deposition and network disconnection from memory
systems[71–73]. Hippocampal atrophy has also been linked to high amyloid-β burden in tau-
positive cases and to cognitive decline[74]. In our study, we observed an association between
increased cortical MK-6240 uptake and reduced ipsilateral hippocampal volume. This
relationship may reflect network-level effects that impact the hippocampus together with large-
scale systems. In several prior studies, associations between hippocampal pathology and
network level compromise have been demonstrated[75–77], suggesting a coupling of pathological
processes in TLE occurring the mesiotemporal lobe and at large scale. However, detailed
analyses of these mechanisms beyond the correlational assessment conducted here are beyond
the scope of the present manuscript, and further studies are needed to clarify the role of
hippocampal tau pathology in cognition and network disconnection.

Our findings also demonstrated significant associations between MK-6240 uptake and poorer
performance in executive function and episodic memory, which again support a potential role
of phosphorylated tau in progressive, multisystem dysfunction in TLE. These findings suggest
a vulnerability of cognitive networks governing episodic memory and executive function to
tau-related pathology in TLE. The preservation of semantic memory in our patients and the
absence of a correlation with MK-6240 may, on the other hand, suggest compensatory
mechanisms similar to those in AD, where episodic memory and executive function are more
markedly affected than semantic memory and conceptual knowledge[3,78]. In future work, the
incorporation of other measures probing into additional mechanisms of memory dysfunctions,
as well as those tapping into processes mediated by other systems (*e.g.,* parieto-frontal control
processes) may be valuable.
Together, these findings suggest that elevated phosphorylated tau in young adults with TLE
may represent a candidate mechanism contributing to large-scale network dysfunction. As tau
imaging in epilepsy advances, integrating larger, longitudinal, and multimodal datasets across
diverse age ranges will be critical to clarifying disease trajectories and individual variability.
Ultimately, this work lays the groundwork for a network-based molecular framework of
epilepsy, which can be refined as more high-resolution *in-vivo* and *post-mortem* data become
available.

METHODS

Participants

We recruited 28 TLE patients (14 Females, mean±SD age=33±8 years) and 28 healthy
volunteers with a similar age and sex distribution (14 Females, mean±SD age=37±12 years)
between October 2019 and July 2025. Diagnosis of TLE and lateralization of seizure foci were
determined by the criteria of the International League Against Epilepsy (ILAE), based on a
comprehensive examination that included clinical history, neurological examination, review of
medical records, video-electroencephalography (EEG) recordings, and clinical MRI. As in
earlier work[75], hippocampal atrophy was determined as volumes or left-right asymmetry
beyond 1.5 standard deviations of the corresponding mean of healthy controls: 8 (29%) patients
had unilateral hippocampal atrophy ipsilateral to the focus and 19 (71%) had normal
hippocampal volume. Mean±SD age at seizure onset was 21±14 years and mean±SD duration
of epilepsy was 15±11 years. Healthy controls were recruited via advertisement; all denied a
history of neurological, psychiatric, or significant medical comorbidities, including traumatic
brain injury. A subgroup in both cohorts had longitudinal acquisition (28/13 patients and 28/7
controls) with a minimum follow up time of 12 months (median=15 months, range=12 months
– 42 months). Interictal epileptiform discharge frequency (IEDs) was obtained either from the
epilepsy monitoring unit or from the long-term monitoring (EEG). Generalized tonic-clonic
seizures frequency (GTCS) was obtained through retrospective clinical data review.

Standard protocol approvals, registrations, and patient consents

The Ethics Committee of the Montreal Neurological Institute and Hospital (MNI) approved the
study (#2018-4148). Written informed consent, including a statement for open sharing of
collected research data, was obtained from all participants.

PET Image acquisition and processing

Tau PET scanning was performed on a Siemens high-resolution research tomograph (HRRT)
at the McConnell Brain Imaging Center of the MNI, using $[^{18}F]$-MK-6240 tracer[17], which
shows low off-target binding[16]. The tracer was injected intravenously, and images were
acquired after 90 minutes for approximately 30 minutes. A previous validation study in our
center has shown that scans 90-110 minutes post-injection may optimally balance duration and
signal. In TLE, mean injected MK-6240 activity was 226.1±22.6 MBq/L. In controls, it was
237.2±21.5 MBq/L. Images were reconstructed with an ordered subset expectation
maximization algorithm. A short transmission scan (11 minutes) with a rotating Cesium137
source was used for attenuation correction. Images underwent correction for dead time, decay,
as well as random and scattered coincidences. Dynamic MK-6240 images (4 in total) were co-
registered and averaged. Average MK-6240 uptake was registered to the T1-weighted MRI
(T1w) using a novel affine, edge-based approach with a cross-correlation cost function and
histogram matching (**Supplementary Figure 7A**). Next, it was partial volume corrected using
a probabilistic mask of the grey matter. We calculated standardized uptake value ratio (SUVR)

images using the cerebellar grey matter as a reference region[79]. To assess reproducibility in the
mixed effects models, we additionally tested the use of the brainstem and a composite reference
region[80] (**Supplementary Figure 3**). The SUVR volume was registered to the medial cortical
surface derived from T1-weighted MRI, resulting in surface-based MK-6240 maps
(**Supplementary Figure 7B**). Surface-based MK-6240 data were blurred with a 10-mm
FWHM surface-based kernel, and participant-specific data were re-sampled to the fsLR-32k
standard template surface[81]. The following subcortical structures were automatically
segmented on the T1w image for each hemisphere: thalamus, caudate nucleus, putamen, globus
pallidus, nucleus accumbens, as well as the hippocampus After calculating the mean value of
all the voxels of each structure, we computed SUVR values based on the same reference region
as for the other structures.

Cognitive assessments
Cognitive assessment included the EpiTrack, a 15-min screening tool primarily targeting
evaluate executive function, designed to monitor cognitive side effects of anti-seizure
medication and seizures[79]. Episodic and semantic memory were assessed using previously
validated tasks. While the episodic memory task involved learning/recalling object pairs across
encoding and retrieval phases, the semantic memory task required identifying conceptually
related object pairs from visual prompts[80,81]. For logistic reasons, the cognitive tests were
performed at the time of the MRI scanning (13 ± 15 months apart from the PET scanning across
participants).

Multimodal MRI acquisition and processing
MRI scans were also obtained at the McConnell Brain Imaging Centre, using a 3T Siemens
Magnetom Prisma-Fit scanner equipped with a 64-channel head coil. Participants underwent a
T1w structural scan, followed by multi-shell diffusion weighted imaging (DWI) and resting-
state functional MRI (rs-fMRI). T1w images were acquired using a 3D-MPRAGE sequence
(0.8 mm isovoxels). Diffusion-weighted imaging data were collected using a multi-band
accelerated 2D spin-echo EPI sequence (1.6 mm isovoxels) with three b-value shells (300, 700,
2000 s/mm²) and corresponding diffusion directions (10, 40, 90), along with phase-reversed b0
images for distortion correction. The rs-fMRI scans were acquired with a 7-min multiband 2D-
BOLD EPI sequence (3 mm isovoxels, TR = 600 ms), with participants instructed to keep their
eyes open and fixate on a cross. For detailed parameters, see the MICs database
(http://n2t.net/ark:/70798/d72xnk2wd397j190qv)[82].
Multimodal MRI data processing and feature fusion leveraged micapipe v0.2.3
(http://micapipe.readthedocs.io), with details described elsewhere[83]. In brief, T1w data were
processed for intensity abnormalities, followed by automated segmentation of subcortical
structures using FSL FIRST[84], the generation of probabilistic tissue estimates using FSL
FAST[85], and cortical surface segmentations using FreeSurfer 6.0[86]. Surfaces were manually
quality controlled and corrected if necessary.
The rs-fMRI data underwent motion correction, distortion correction, denoising, bandpass
filtering, surface mapping, and surface registration. We computed pairwise Product-moment

correlation coefficients between vertices to generate a subject-level functional connectivity
(FC) matrix. Resulting correlation coefficients were normalized via Fisher r-to-z
transformation.
DWI preprocessing involved motion correction, denoising, and correction for inhomogeneity
and geometric distortions. Anatomically constrained tractography was performed using tissue
segmentations derived from each participant's T1w images that were co-registered to native
DWI space. Multi-shell and multi-tissue response functions were estimated, followed by
constrained spherical deconvolution and intensity normalization. A tractogram with 40M
streamlines was generated and refined using SIFT2[87]. Structural connectome (SC) weights
were defined as the filtered streamline counts between cortical vertices using the fsLR-5k
standard surface. A secondary SC was generated using streamline edge lengths as weights.
These two connectomes were combined to create a weighted SC (SCw) that integrates
connectivity strength and distance information, such that the connection with high streamline
counts but short lengths are penalized relative to longer connections.

Data Normalization
A novel PET-to-MRI registration was performed in two steps. First, edge images were
generated from the T1w and the average MK-6240 PET map. Next, the edge image of the MK-
6240 average map was registered to the T1w edge image using a cross-correlation cost function
with histogram matching. (**Supplementary figure 7A**). Individual quality control was
performed for each registration to ensure optimal results. For DWI and rs-fMRI, a two-step
registration process was applied. First, a modality-agnostic, deep learning-based
segmentation[88] was used to generate labeled images for each modality. These labeled images
were then registered to the T1w-derived label map using affine registration, followed by
nonlinear registration to ensure alignment in the subject's native T1w space.
The resulting transformations were applied to map PET data onto the subject's cortical surface,
reconstructed from the native T1w image. This native surface was subsequently resampled to
the fsLR standard surface meshes at two different resolutions ~32,000 and ~5,000 vertices per
hemisphere. The fsLR-32k surface was used to sample quantitative values from MK-6240,
while the fsLR-5k surface was used for constructing both structural and functional
connectomes derived from DWI and rs-fMRI.

Statistical analysis
Analyses were performed using R and BrainStat[89], an open-access toolbox for brain-wide
statistical analysis (http://brainstat.readthedocs.io). In patients, prior to analysis, cortical
features were sorted into ipsilateral/contralateral relative to the seizure focus.

a) Mapping of tau uptake in TLE. We compared MK-6240 vertex-wise SUVR between
individuals with TLE and healthy controls using a surface-based linear mixed effects model,
controlling for age, sex, and participant as a random effect. Separate surface-based models were
applied to control for cortical thickness and to compare males with TLE versus male controls,
and females with TLE versus female controls. An additional interaction model was used to

assess the combined effects of group and sex on tau uptake. Here and below, surface-based
analyses were corrected for multiple comparisons using random field theory (two-tailed), with
a significance threshold set at $p < 0.025$. As in the cortical analysis, we compared MK-6240
uptake in subcortical ROI between TLE patients and healthy controls using mixed effects
models. Models included age as a covariate, a sex by group interaction, and a random intercept
for each participant. Statistical significance was assessed using a two-tailed threshold of $p <$
0.025, with false discovery rate correction for multiple comparisons.
8
In addition to examining between-group differences, a binary tau probability map was created
by thresholding the MK-6240 PET data in every individual. The threshold was defined as the
97.5th percentile (1.56 SUVR) of values in the healthy control population (**Table 2**). Vertices
exceeding this threshold were assigned a value of 1, while all other vertices were set to 0,
resulting in a binary map indicating regions of elevated tau deposition. The group difference in
mean MK-6240 SUVR values from significant clusters (MK-6240$_{sig}$) in both hemispheres was
assessed using a Welch two sample t-test.
Longitudinal changes in MK-6240 were assessed using linear-mixed-effects models. One
model evaluated the change in MK-6240 uptake over time from the mean values extracted from
significant regions. This model was fitted using restricted maximum likelihood (REML), and
t-tests for fixed effects were calculated using Satterthwaite's approximation for degrees of
freedom. Separate estimated slopes and 95% confidence intervals were obtained for each group
using the Kenward-Roger method. Model assumptions were checked via inspection of scaled
residuals, and between-subject variability was captured by the participant-level random
intercept. A second model evaluated surface-based longitudinal changes in MK-6240 uptake
comparing TLE and healthy controls. This model was adjusted for age and sex and corrected
for multiple comparisons using random field theory ($p<0.025$).
26
b) Relationship to clinical and cognitive variables. Correlation analyses were initially
performed using mean MK-6240 SUVR values extracted from significant clusters in both
hemispheres identified in the group-difference analysis (**Figure 2, left column**). These findings
were complemented with surface-based linear models using vertex-wise MK-6240 SUVR
values (Figure 2, right column). These analyses included only cross-sectional data from the
first time point. We examined associations to age, disease duration, age at onset,
ipsilateral/contralateral hippocampal volume, and cognitive measures. Unless otherwise
specified, these analyses controlled for age and sex. In addition to the analyses of Figure 2 we
evaluated the relationships between MK-6240 and age at onset of the epilepsy and the
contralateral hippocampal volume, but no significant correlations were observed
(**Supplementary figure 5**).
38
c) Network contextualization. We correlated the t-statical map from the above between-group
comparison in MK-6240 between patients and controls with network metrics derived from
group-level SC and FC. Specifically, network metrics were mean connectivity and
neighborhood-weighted connectivity[90]. SCs and FCs were modeled as non-thresholded,
weighted graphs. Each subject's connectome was represented by a vertex-wise adjacency
matrix based on either structural or functional connectivity. Group-level connectomes were

then constructed by averaging edge weights across individuals, separately for DWI and rs-fMRI data.

From the group-level weighted connectivity graphs, two primary network measures were derived: mean connectivity and neighborhood-weighted connectivity, the latter also referred to as Neighborhood Deformation Estimates[90]. These metrics were used as dependent variables in the group-level correlation analyses, contextualizing regional differences in MK-6240 findings (t-statistical map) within the broader network architecture.

To compute neighborhood-weighted connectivity, we applied the following procedure. For each cortical node (D_i), the feature values of its connected neighbors were extracted and averaged (dj), weighted by the strength of their connection (NC_{ij}). Self-connections (i.e., diagonal elements of the connectivity matrix, $j{\neq}i$) and the medial wall vertices were excluded from the computation to avoid circular influence. The neighborhood-weighted value for a given node was thus a degree-normalized, connectivity-weighted average of its neighbors' feature values, such as the t-statistical map of MK-6240 difference. See **Equation 1**:

$$D_i = \frac{1}{N_i} \sum_{j=1, j\neq i}^{N_i} d_j \times NC_{ij}$$

This method quantifies the relation between the original nodal measurements, for example regional PET uptake, and the connectivity-weighted average of these same measurements across each node's network-defined neighborhood. In doing so, it captures the extent to which regional signal patterns are spatially structured by the underlying network topology. Detailed implementation and supporting scripts are available in our project repository: *https://github.com/MICA-MNI/2025_in-vivo_tauPET-mk6240_TLE/blob/main/scripts/Fig-2_network_contextualization.ipynb*

To correct for multiple comparisons and account for the spatial autocorrelation inherent in surface-based data, non-parametric spin permutation testing was used (using 10,000 permutations)[91,92]. Significance was assessed relative to these null models. Medial wall vertices were excluded from all calculations.

d) Structural equation modeling. SEM examined whether graph-theoretical measures of structural brain networks (see below) were associated with the relationship between MK-6240sig SUVR and three memory outcomes (EpiTrack, episodic memory, and semantic memory). Models were estimated using maximum likelihood with non-parametric bootstrapping (10,000 resamples) to obtain robust standard errors and confidence intervals, particularly for indirect effects. Model fit was summarized using the root mean square error of approximation and the standardized root mean square residual, and standardized path coefficients are reported.

Graph-theoretical metrics were derived from individual fsLR-5k structural and functional weighted connectomes[93]. For each participant, we computed mean strength, mean neighborhood-weighted connectivity, mean characteristic path length, weighted global efficiency[94], and weighted clustering coefficient[95,96]. Functional connectomes were Fisher-z

transformed, absolute values were taken, normalized to the maximum weight, and thresholded
to retain the top 20% of connections[97,98]. Structural connectomes were weighted by edge length
to penalize short-distance connections with high connectivity and subsequently normalized to
the maximum weight. Individual mean neighborhood-weighted connectivity was calculated
using each subject's weighted structural connectome (NC_{ij}) with MK-6240 uptake values from
connected cortical neighbors (*di*, see **Equation 1**). The medial wall was removed from all
connectomes. To explore associations among variables, correlograms were generated including
MK-6240sig, behavioral measures, and network metrics. Due to high collinearity among
network measures, only structural weighted global efficiency and structural mean
neighborhood-weighted connectivity were retained for SEM analyses. Functional network
metrics were excluded because they showed no significant associations with behavioral
outcomes (**Supplementary Figure 8**).

Three preliminary SEMs were first estimated separately for each cognitive outcome (EpiTrack,
episodic memory, semantic memory). Based on these results, a single multivariate SEM was
specified with MK-6240sig as the predictor, structural weighted global efficiency, and mean
neighborhood-weighted connectivity as mediators, and the three memory measures as
correlated outcomes. The predictor was specified to directly predict both mediators (paths *a1*
and *a2*), which were allowed to covary. Each outcome was then regressed on both mediators
and on MK-6240 signal, estimating mediator-to-outcome (*b*) paths and direct (*c'*) effects.
Residual covariances were specified among outcomes to account for shared unexplained
variance. Indirect effects were calculated as the product of *a* and *b* paths for each mediator and
outcome, and total indirect effects were obtained by summing mediator-specific indirect
effects. Total effects were computed as the sum of direct and indirect effects. All parameters
were estimated simultaneously, and full model estimates are reported in **Table 4**.
25
Tau immnunohistochemistry
Brain tissue samples from the surgical resections of the temporal neocortex, parahippocampal
regions, hippocampus, and amygdala were processed according to standardized clinical
procedures at the Montreal Neurological Institute. Samples were fixed in 10% neutral buffered
formalin and embedded in paraffin (FFPE). Immunohistochemistry (IHC) was carried on a
Ventana Discovery Ultra automated staining platform, following manufacturer instructions.
The sections were pre-treated with CC1, and the primary monoclonal antibody AT8
(recognizing phosphorylated tau at Ser202/Thr205; Thermo Fisher Scientific; MN1020) used
at 1:1200 dilution.
Whole-slide imaging was performed using a Hamamatsu NanoZoomer S210 digital scanner
(C13239-01). All digitized images were reviewed and interpreted by an expert
neuropathologist to characterize the distribution and morphology of phosphorylated tau
inclusions.

DATA AVAILABILITY
All the data used in the study is hosted on OSF (https://osf.io/ct3gw).
3
4 **CODE AVAILABILITY**
5 All processing workflow scripts and the code for analyses and statistics are openly available
6 on GitHub (https://github.com/MICA-MNI/2025_in-vivo_tauPET-mk6240_TLE).
7

ACKNOWLEDGEMENTS
We thank the McConnell Brain Imaging Centre (BIC) PET and MRI units of the MNI, and the
study participants for supporting our research. RRC received support from the Fonds de la
Recherche du Québec – Santé (FRQ-S), the Montreal Neurological Institute Jeanne Timmins
Costello Fellowship, and the Healthy Brains, Healthy Lives – Entrepreneur Postdoc
Fellowship. JR received support from the Canadian Open Neuroscience Platform (CONP) and
Canadian Institutes of Health Research (CIHR). MK is supported by the Wellcome Trust
(221934/Z/20/Z). LC acknowledges previous support from Brain Research UK (Award
14181). BCB acknowledges support from CIHR (FDN-154298, PJT-174995), SickKids
Foundation (NI17-039), Natural Sciences and Engineering Research Council (NSERC;
Discovery-1304413), Azrieli Center for Autism Research of the Montreal Neurological
Institute (ACAR), BrainCanada, FRQ-S, the Helmholtz International BigBrain Analytics and
Learning Laboratory (Hiball), the Canada Research Chairs Program (CRC), and the Centre of
Excellence in Epilepsy at the Neuro (CEEN).

1   REFERENCES

1.   Pitkänen, A. & Sutula, T. P. Is epilepsy a progressive disorder? Prospects for new therapeutic approaches in temporal-
lobe epilepsy. *Lancet Neurol.* **1**, 173–181 (2002).
2.   Galovic, M. *et al.* Progressive Cortical Thinning in Patients With Focal Epilepsy. *JAMA Neurol* **76**, 1230–1239 (2019).
3.   Tai, X. Y. *et al.* Review: Neurodegenerative processes in temporal lobe epilepsy with hippocampal sclerosis: Clinical,
pathological and neuroimaging evidence. *Neuropathol. Appl. Neurobiol.* **44**, 70–90 (2018).
4.   Caciagli, L. *et al.* A meta-analysis on progressive atrophy in intractable temporal lobe epilepsy: Time is brain? *Neurology*
89, 506–516 (2017).
5.   Boyle, P. A., Wilson, R. S., Aggarwal, N. T., Tang, Y. & Bennett, D. A. Mild cognitive impairment. *Neurology* **67**, 441–
445 (2006).
6.   Noebels, J. A perfect storm: Converging paths of epilepsy and Alzheimer's dementia intersect in the hippocampal
formation. *Epilepsia* **52 Suppl 1**, 39–46 (2011).
7.   Lam, A. D. *et al.* Association of Seizure Foci and Location of Tau and Amyloid Deposition and Brain Atrophy in Patients
With Alzheimer Disease and Seizures. *Neurology* **103**, e209920 (2024).
8.   Irizarry, M. C. *et al.* Incidence of new-onset seizures in mild to moderate Alzheimer disease. *Arch. Neurol.* **69**, 368–372
(2012).
9.   Amatniek, J. C. *et al.* Incidence and predictors of seizures in patients with Alzheimer's disease. *Epilepsia* **47**, 867–872
(2006).
10. Tai, X. Y. *et al.* Hyperphosphorylated tau in patients with refractory epilepsy correlates with cognitive decline: a study
of temporal lobe resections. *Brain* **139**, 2441–2455 (2016).
11. Salemme, S. *et al.* The prognosis of mild cognitive impairment: A systematic review and meta-analysis. *Alzheimers
Dement (Amst)* **17**, e70074 (2025).
12. Canet, G. *et al.* Seizure activity triggers tau hyperphosphorylation and amyloidogenic pathways. *Epilepsia* **63**, 919–935
(2022).
13. Alves, M., Kenny, A., de Leo, G., Beamer, E. H. & Engel, T. Tau Phosphorylation in a Mouse Model of Temporal Lobe
Epilepsy. *Front. Aging Neurosci.* **11**, 490080 (2019).
14. Thom, M. & Koepp, M. Tau protein in drug-resistant epilepsy and cognitive decline. in *Inflammation and Epilepsy: New
Vistas* 149–184 (Springer International Publishing, Cham, 2021).
15. Malarte, M.-L., Nordberg, A. & Lemoine, L. Characterization of MK6240, a tau PET tracer, in autopsy brain tissue from
Alzheimer's disease cases. *Eur. J. Nucl. Med. Mol. Imaging* **48**, 1093–1102 (2021).
16. Aguero, C. *et al.* Autoradiography validation of novel tau PET tracer [F-18]-MK-6240 on human postmortem brain tissue.
Acta Neuropathologica Communications **7**, 1–15 (2019).
17. Hopewell, R. *et al.* A simplified radiosynthesis of [F]MK-6240 for tau PET imaging. *J Labelled Comp Radiopharm* **62**,
109–114 (2019).
18. Knowlton, R. C. *et al.* In vivo hippocampal glucose metabolism in mesial temporal lobe epilepsy. *Neurology* **57**, 1184–
1190 (2001).
19. Juhász, C. *et al.* Glucose and [11C]flumazenil positron emission tomography abnormalities of thalamic nuclei in temporal
lobe epilepsy. *Neurology* **53**, 2037–2045 (1999).
20. Seeley, W. W., Crawford, R. K., Zhou, J., Miller, B. L. & Greicius, M. D. Neurodegenerative diseases target large-scale
human brain networks. *Neuron* **62**, 42–52 (2009).
21. de Bruin, H. *et al.* Connectivity as a universal predictor of tau progression in atypical Alzheimer's disease. *Brain* **148**,
3893–3912 (2025).
22. Ottoy, J. *et al.* Tau follows principal axes of functional and structural brain organization in Alzheimer's disease. *Nat
Commun* **15**, 5031 (2024).
23. Kerestes, R. *et al.* Patterns of subregional cerebellar atrophy across epilepsy syndromes: An ENIGMA-Epilepsy study.
Epilepsia **65**, 1072–1091 (2024).
24. Whelan, C. D., Haaker, J. G. & Sisodiya, S. M. *Structural Brain Abnormalities in the Common Epilepsies Assessed in a
Worldwide ENIGMA Study*. (2018).
25. Lutz, M. T. & Helmstaedter, C. EpiTrack: tracking cognitive side effects of medication on attention and executive
functions in patients with epilepsy. *Epilepsy Behav* **7**, 708–714 (2005).
26. Cabalo, D. G. *et al.* Differential reorganization of episodic and semantic memory systems in epilepsy-related
mesiotemporal pathology. *Brain* **147**, 3918–3932 (2024).
27. Tavakol, S. *et al.* Differential relational memory impairment in temporal lobe epilepsy. *Epilepsy Behav* **155**, 109722
(2024).
28. Betthauser, T. J. *et al.* In Vivo Characterization and Quantification of Neurofibrillary Tau PET Radioligand 18F-MK-
6240 in Humans from Alzheimer Disease Dementia to Young Controls. *Journal of Nuclear Medicine* **60**, 93–99 (2019).
29. Pascoal, T. A. *et al.* 18F-MK-6240 PET for early and late detection of neurofibrillary tangles. *Brain* **143**, 2818–2830
(2020).
30. Kreisl, W. C. *et al.* Patterns of tau pathology identified with 18F-MK-6240 PET imaging. *Alzheimer's & Dementia* **18**,
272–282 (2022).
31. Pascoal, T. A. *et al.* Longitudinal 18F-MK-6240 tau tangles accumulation follows Braak stages. *Brain* **144**, 3517–3528
(2021).
32. Malpetti, M., La Joie, R. & Rabinovici, G. D. Tau Beats Amyloid in Predicting Brain Atrophy in Alzheimer Disease:
Implications for Prognosis and Clinical Trials. *Journal of Nuclear Medicine* **63**, 830–832 (2022).
33. Ossenkoppele, R. *et al.* Accuracy of Tau Positron Emission Tomography as a Prognostic Marker in Preclinical and
Prodromal Alzheimer Disease: A Head-to-Head Comparison Against Amyloid Positron Emission Tomography and

Magnetic Resonance Imaging. *JAMA Neurol* **78**, 961–971 (2021).

34. La Joie, R. *et al.* Prospective longitudinal atrophy in Alzheimer's disease correlates with the intensity and topography of baseline tau-PET. *Sci. Transl. Med.* **12**, (2020).

35. Kunach, P. *et al.* Cryo-EM structure of Alzheimer's disease tau filaments with PET ligand MK-6240. *Nat Commun* **15**, 8497 (2024).

36. Gourmaud, S. *et al.* Alzheimer-like amyloid and tau alterations associated with cognitive deficit in temporal lobe epilepsy. *Brain* **143**, 191–209 (2020).

37. Smith, K. M. *et al.* Tau deposition in young adults with drug-resistant focal epilepsy. *Epilepsia* **60**, 2398–2403 (2019).

38. Thom, M. *et al.* Neurofibrillary tangle pathology and Braak staging in chronic epilepsy in relation to traumatic brain injury and hippocampal sclerosis: a post-mortem study. *Brain* **134**, 2969–2981 (2011).

39. Ballerini, A. *et al.* Late-onset temporal lobe epilepsy: insights from brain atrophy and Alzheimer's disease biomarkers. *Brain* **148**, 185–198 (2025).

40. Bernasconi, N. Is epilepsy a curable neurodegenerative disease? *Brain* **139**, 2336–2337 (2016).

41. Snyder, H. M. *et al.* Sex biology contributions to vulnerability to Alzheimer's disease: A think tank convened by the Women's Alzheimer's Research Initiative. *Alzheimers Dement* **12**, 1186–1196 (2016).

42. Wisch, J. K. *et al.* Sex-related Differences in Tau Positron Emission Tomography (PET) and the Effects of Hormone Therapy (HT). *Alzheimer Dis Assoc Disord* **35**, 164–168 (2021).

43. Ferretti, M. T., Dimech, A. S. & Chadha, A. S. *Sex and Gender Differences in Alzheimer's Disease*. (Academic Press, 2021).

44. Zeighami, Y., Tremblay, C. & Dadar, M. Sex differences in vulnerability to tau pathology: Impact on cognitive decline. *Alzheimers Dement* **21**, e70634 (2025).

45. Lopez-Lee, C., Torres, E. R. S., Carling, G. & Gan, L. Mechanisms of sex differences in Alzheimer's disease. *Neuron* **112**, 1208–1221 (2024).

46. Hopping, L., Kyriakopoulos, P. & Bui, E. Sex and gender differences in epilepsy. *Int Rev Neurobiol* **164**, 235–276 (2022).

47. Mrzyglod, A. *et al.* Patterns of phosphorylated tau accumulation in a spectrum of acquired and developmental brain lesions associated with refractory epilepsy. *Epilepsia* (2025) doi:10.1111/epi.18418.

48. Altmann, A. *et al.* A systems-level analysis highlights microglial activation as a modifying factor in common epilepsies. *Neuropathol Appl Neurobiol* **48**, e12758 (2022).

49. Dickstein, L. P. *et al.* Neuroinflammation in neocortical epilepsy measured by PET imaging of translocator protein. *Epilepsia* **60**, 1248–1254 (2019).

50. Langworth-Green, C. *et al.* Chronic effects of inflammation on tauopathies. *Lancet Neurol* **22**, 430–442 (2023).

51. Toscano, E. C. B. *et al.* Hyperphosphorylated Tau in Mesial Temporal Lobe Epilepsy: a Neuropathological and Cognitive Study. *Mol Neurobiol* **60**, 2174–2185 (2023).

52. O'Brien, T. J. *et al.* The utility of a 3-dimensional, large-field-of-view, sodium iodide crystal--based PET scanner in the presurgical evaluation of partial epilepsy. *J Nucl Med* **42**, 1158–1165 (2001).

53. Theodore, W. H. *et al.* [18F]fluorodeoxyglucose positron emission tomography in refractory complex partial seizures. *Ann Neurol* **14**, 429–437 (1983).

54. Vinton, A. B. *et al.* The extent of resection of FDG-PET hypometabolism relates to outcome of temporal lobectomy. *Brain* **130**, 548–560 (2007).

55. Griffith, H. R. *et al.* Preoperative FDG-PET temporal lobe hypometabolism and verbal memory after temporal lobectomy. *Neurology* **54**, 1161–1165 (2000).

56. Courtney, M. R. *et al.* F-FDG-PET hypometabolism as a predictor of favourable outcome in epilepsy surgery: protocol for a systematic review and meta-analysis. *BMJ Open* **12**, e065440 (2022).

57. Sharpe, C. *et al.* Longitudinal changes of focal cortical glucose hypometabolism in adults with chronic drug resistant temporal lobe epilepsy. *Brain Imaging Behav* **15**, 2795–2803 (2021).

58. Benedek, K., Juhász, C., Chugani, D. C., Muzik, O. & Chugani, H. T. Longitudinal changes in cortical glucose hypometabolism in children with intractable epilepsy. *J Child Neurol* **21**, 26–31 (2006).

59. Zhou, J., Gennatas, E. D., Kramer, J. H., Miller, B. L. & Seeley, W. W. Predicting regional neurodegeneration from the healthy brain functional connectome. *Neuron* **73**, 1216–1227 (2012).

60. Crossley, N. A. *et al.* The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain* **137**, 2382–2395 (2014).

61. Royer, J. *et al.* Epilepsy and brain network hubs. *Epilepsia* **63**, 537–550 (2022).

62. Georgiadis, F. *et al.* Connectome architecture shapes large-scale cortical alterations in schizophrenia: a worldwide ENIGMA study. *Mol Psychiatry* **29**, 1869–1881 (2024).

63. Larivière, S. *et al.* Network-based atrophy modeling in the common epilepsies: A worldwide ENIGMA study. *Sci Adv* **6**, (2020).

64. Liu, L. *et al.* Trans-synaptic spread of tau pathology in vivo. *PLoS One* **7**, e31302 (2012).

65. Braak, H., Alafuzoff, I., Arzberger, T., Kretzschmar, H. & Del Tredici, K. Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol* **112**, 389–404 (2006).

66. Harrison, T. M. *et al.* Longitudinal tau accumulation and atrophy in aging and alzheimer disease. *Ann Neurol* **85**, 229–240 (2019).

67. Pan, N., Liu, S., Ge, X., Zheng, Y. & for Alzheimer's Disease Neuroimaging Initiative. Association of hippocampal atrophy with tau pathology of temporal regions in preclinical Alzheimer's disease. *J Alzheimers Dis* 13872877251314785 (2025).

68. Liang, Y. *et al.* Association of apolipoprotein E genotypes with epilepsy risk: A systematic review and meta-analysis. *Epilepsy Behav* **98**, 27–35 (2019).

69. Chen, Y. *et al.* Seizure frequency, APOE ε4, and cognitive function in older people with epilepsy. *Acta Epileptol* **7**, 30

(2025).

70. Liu, S., He, Z., Shi, W. & Li, J. The association between APOE gene polymorphisms and the risk, characteristics, and prognosis of epilepsy: A systematic review and meta-analysis. *Epilepsy Behav* **160**, 110070 (2024).

71. de Souza, L. C. *et al.* CSF tau markers are correlated with hippocampal volume in Alzheimer's disease. *Neurobiol Aging* **33**, 1253–1257 (2012).

72. Aschenbrenner, A. J., Gordon, B. A., Benzinger, T. L. S., Morris, J. C. & Hassenstab, J. J. Influence of tau PET, amyloid PET, and hippocampal volume on cognition in Alzheimer disease. *Neurology* **91**, e859–e866 (2018).

73. Harrison, T. M. *et al.* Tau deposition is associated with functional isolation of the hippocampus in aging. *Nat Commun* **10**, 4900 (2019).

74. Pan, F.-F. *et al.* Associations of hippocampal volumes, brain hypometabolism, and plasma NfL with amyloid, tau, and cognitive decline. *Alzheimers Dement* **21**, e70005 (2025).

75. Bernhardt, B. C. *et al.* Temporal lobe epilepsy: Hippocampal pathology modulates connectome topology and controllability. *Neurology* **92**, e2209–e2220 (2019).

76. Bernhardt, B. C. *et al.* The spectrum of structural and functional imaging abnormalities in temporal lobe epilepsy. *Ann Neurol* **80**, 142–153 (2016).

77. Larivière, S. *et al.* Brain Networks for Cortical Atrophy and Responsive Neurostimulation in Temporal Lobe Epilepsy. *JAMA Neurol* **81**, 1199–1209 (2024).

78. Simons, J. S., Graham, K. S. & Hodges, J. R. Perceptual and semantic contributions to episodic memory: evidence from semantic dementia and Alzheimer's disease. *J. Mem. Lang.* **47**, 197–213 (2002).

79. Pascoal, T. A. *et al.* In vivo quantification of neurofibrillary tangles with [F]MK-6240. *Alzheimers. Res. Ther.* **10**, 74 (2018).

80. Landau, S. M. *et al.* Measurement of longitudinal β-amyloid change with 18F-florbetapir PET and standardized uptake value ratios. *J Nucl Med* **56**, 567–574 (2015).

81. Van Essen, D. C., Glasser, M. F., Dierker, D. L., Harwell, J. & Coalson, T. Parcellations and hemispheric asymmetries of human cerebral cortex analyzed on surface-based atlases. *Cereb. Cortex* **22**, 2241–2262 (2012).

82. Royer, J. *et al.* An Open MRI Dataset For Multiscale Neuroscience. *Sci Data* **9**, 569 (2022).

83. Cruces, R. R. *et al.* Micapipe: A pipeline for multimodal neuroimaging and connectome analysis. *Neuroimage* **263**, 119612 (2022).

84. Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans. Med. Imaging* **20**, 45–57 (2001).

85. Woolrich, M. W. *et al.* Bayesian analysis of neuroimaging data in FSL. *Neuroimage* **45**, S173–86 (2009).

86. Fischl, B. FreeSurfer. *Neuroimage* **62**, 774–781 (2012).

87. Tournier, J.-D. *et al.* MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. *Neuroimage* **202**, 116137 (2019).

88. Billot, B. *et al.* SynthSeg: Segmentation of brain MRI scans of any contrast and resolution without retraining. *Med Image Anal* **86**, 102789 (2023).

89. Larivière, S. *et al.* BrainStat: A toolbox for brain-wide statistics and multimodal feature associations. *Neuroimage* **266**, 119807 (2023).

90. Shafiei, G. *et al.* Spatial Patterning of Tissue Volume Loss in Schizophrenia Reflects Brain Network Architecture. *Biol Psychiatry* **87**, 727–735 (2020).

91. Vos de Wael, R. *et al.* BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Commun Biol* **3**, 103 (2020).

92. Alexander-Bloch, A. F. *et al.* On testing for spatial correspondence between maps of human brain structure and function. *Neuroimage* **178**, 540–551 (2018).

93. Rubinov, M. & Sporns, O. Complex network measures of brain connectivity: uses and interpretations. *Neuroimage* **52**, 1059–1069 (2010).

94. Barrat, A., Barthélemy, M., Pastor-Satorras, R. & Vespignani, A. The architecture of complex weighted networks. *Proc Natl Acad Sci U S A* **101**, 3747–3752 (2004).

95. Opsahl, T. & Panzarasa, P. Clustering in weighted networks. *Soc. Networks* **31**, 155–163 (2009).

96. Wasserman, S. & Faust, K. *Social Network Analysis: Methods and Applications.* (Cambridge University Press, 1994).

97. Luppi, A. I. *et al.* Systematic evaluation of fMRI data-processing pipelines for consistent functional connectomics. *Nat Commun* **15**, 4745 (2024).

98. Zuo, X.-N., Biswal, B. B. & Poldrack, R. A. *Reliability and Reproducibility in Functional Connectomics.* (Frontiers Media SA, 2019).

AUTHOR INFORMATION
Authors and affiliations
McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University,
Montreal, Québec, Canada.
*Raúl R. Cruces, Jack Lam**, *Thaera Arafat, Jessica Royer, Judy Chen, Ella Sahlas, Arielle*
Dascal, Daniel Mendelson, Robert Hopewell, Chris Hung-Hsin Hsiao, Gassan Massarweh,
Jean-Paul Soucy, Pedro Rosa-Neto, Andrea Bernasconi, Neda Bernasconi, Boris C.*
*Bernhardt**
* joint senior authors
10
11 Montreal Neurological Institute, McGill University Health Centre (MUHC) and other McGill
12 University
13 *Raluca Pana, Marie-Christine Guiot*
14
15 StoP-AD Centre, Douglas Mental Health University Institute, McGill University, Montréal,
16 QC, Canada
17 *Sylvia Villeneuve*
18
19 UCL Queen Square Institute of Neurology, University College London, London WC1N 3BG,
20 UK
21 *Matthias Koepp*
22
23 Department of Neurology, Inselspital, Sleep-Wake-Epilepsy Center, Bern University Hospital,
24 University of Bern, Bern, Switzerland
25 *Lorenzo Caciagli*
26
27 Peter O'Donnell Jr. Brain Institute (OBI), University of Texas Southwestern Medical Centre (UTSW),
28 Dallas, TX, USA
29 *Pedro Rosa-Neto*
30
31 **Corresponding authors**
32 Raul Cruces, MD PhD, Boris C. Bernhardt, PhD

1   **TABLE 1. CLINICAL FEATURES**

Participant	Sex	Age	Handedness	Language	Epilepsy classification	Lateralization	Origin	Onset	Duration	Hippocampal atrophy	GTCSF	IED	Surgery	ENGEL
HC20	F	22	R	fr	focal	R	TLE	10	12	no	2	2	no	
HC10	F	25	R	en	focal	R	mTLE	17	8	yes	0.25	4	yes	IVB
HC19	F	28	R	fr	focal	R	mTLE	0.5	27.5	no	1	3	yes	IA
HC21	F	29	R	en	focal	R	TLE	23	6	no	0	3	no	
HC24	F	31	R	en	focal	R	mTLE	6	25	no	12	3	no	
HC02	F	33	L	en	focal	L	TLE	24	9	yes	130	5	no	
HC01	F	35	R	en	focal	L	mTLE	27	8	no	4	3	yes	IA
HC25	F	35	R	fr	focal	L	TLE	31	4	no	2	3	no	
HC15	F	41	R	fr	focal	R	TLE	39	2	no	0	5	no	
HC09	F	45	R	fr	focal	BL	mTLE	7	38	no	0	3	no	
HC26	F	47	R	fr	focal	L	TLE	12	35	yes	2	5	no	
HC28	F	50	R	fr	focal	R	TLE	32	18	yes	0.22	5	no	
HC27	F	61	R	en	focal	L	TLE	20	41	no	0	NA	no	
HC14	F	65	R	en	focal	L	TLE	41	24	yes	0	3	no	
HC04	M	25	R	en	focal	R	mTLE	2	23	no	0	3	no	
HC05	M	26	R	en	focal	L	mTLE	18	8	no	2.4	0	no	
HC17	M	27	R	fr	focal	L	TLE	25	2	no	2	3	no	
HC22	M	27	R	fr	focal	BL	TLE	8	19	yes	1	0	no	
HC11	M	28	R	en	focal	L	mTLE	26	2	yes	0	3	yes	IA
HC07	M	29	R	fr	focal	R	TLE	2	27	no	1.5	5	no	
HC16	M	29	L	en	focal	L	TLE	6	23	no	0	5	no	
HC13	M	30	R	en	focal	-	TLE	24	6	no	0	3	no	
HC23	M	30	R	en	focal	L	TLE	25	5	no	48	1	no	
HC08	M	33	R	en	focal	R	TLE	18	15	no	0	3	yes	IA
HC12	M	40	R	fr	focal	R	mTLE	24	16	no	0	1	yes	IA
HC06	M	44	R	fr	focal	R	mTLE	28	16	no	0	5	yes	IA
HC03	M	52	R	en	focal	L	mTLE	50	2	yes	0	NA	no	
HC18	M	62	R	fr	focal	L	mTLE	55	7	yes	0	3	no	

F: Female.   M: Male
HA: Hippocampal atrophy of the ipsilateral hippocampus. Defined as a hippocampal volume more than 2 standard deviations below the mean
of controls, normalized by intracranial volume.
Fr: French.   En: English
Epilepsy classification: focal, multifocal, generalized.
L: Left; R: Right; BL: Bilateral
mTLE: Mesial temporal lobe epilepsy
TLE: Temporal lobe epilepsy
IED: Interictal epileptiform discharge frequency from EMU or LTMO.
GTCSF: generalized tonic-clonic seizures frequency per year.
IED was coded on an ordinal 5-point scale (1-5), with increasing values reflecting higher frequency. For example, none was coded as 0, rare
as 1, and frequent as 5.

1 **TABLE 2. SUVR THRESHOLDS**

Group	97.5 th	99.0 th
Healthy controls	*1.56*	*1.65*
All subjects	1.69	1.82

2
3 ***Supplementary Table 2.*** *SUVR thresholds. Percentiles for SUVR values based on the mean cortical uptake of all*
4 *vertices by group. Values represent standardized uptake value ratios (SUVR). We used the 97.5th percentile*
5 *threshold based on healthy controls in the current manuscript.*
6

A | High-probability areas of increase MK6240 by group

7
8 **Supplementary Figure 1.** *Cortical map of abnormal MK-6240 SUVR probability.* Regions with SUVR values
9 greater than 1.5 (percentile 97.5th based on controls) were identified and visualized for each group. Dark red
10 represents areas with higher prevalence of abnormal SUVR uptake, light red indicates lower prevalence. The
11 maximum MK-6240 SUVR probability is in 75% of the TLE population.

TABLE 3. MEAN SUBCORTICAL MK-6240 SUVR VALUES

Structure	Patient	Healthy	T-statistic	p-value
Ipsilateral thalamus	0.67±0.11	0.66±0.11	0.673	0.5
Ipsilateral caudate	0.63±0.12	0.62±0.12	0.845	0.4
Ipsilateral putamen	0.69±0.14	0.66±0.12	1.34	0.2
Ipsilateral pallidus	0.64±0.14	0.64±0.13	0.895	0.4
Ipsilateral amygdala	0.63±0.13	0.61±0.09	0.972	0.3
Ipsilateral hippocampus	0.71±0.12	0.71±0.11	0.412	0.7
Ipsilateral accumbens	0.59±0.14	0.60±0.17	0.411	0.7
Contralateral thalamus	0.69±0.10	0.65±0.10	1.27	0.2
Contralateral caudate	0.63±0.11	0.59±0.11	0.87	0.4
Contralateral putamen	0.68±0.12	0.63±0.13	1.5	0.14
Contralateral pallidus	0.62±0.14	0.57±0.13	0.718	0.5
Contralateral amygdala	0.64±0.10	0.60±0.11	1.28	0.2
Contralateral hippocampus	0.69±0.11	0.69±0.10	-0.37	0.7
Contralateral accumbens	0.57±0.13	0.54±0.14	0.292	0.8

1 Mean±SD

2 Welch Two Sample t-test, **uncorrected** p-values for multiple comparisons.

A | Group differences in thickness

B | Group differences in MK6240 controlled by thickness

C | Longitudinal changes: Patients more than Controls

Supplementary Figure 2. A) Impact of cortical thickness on between-group differences in MK-6240 uptake. Group difference in cortical thickness. The comparison was evaluated using a mixed effects model, controlling for age and sex, and corrected for multiple comparisons with random field theory (p<0.025). **B)** MK-6240 uptake comparing TLE and healthy controls after controlling for cortical thickness. **C)** Surface-based longitudinal changes in MK-6240 uptake comparing TLE and healthy controls. No cortical regions showed statistically significant longitudinal changes or group differences in the longitudinal slopes. Mixed-effects models were adjusted for age and sex and corrected for multiple comparisons using random field theory (p<0.025). Significant regions, when present, are outlined in white.

1
Supplementary Figure 3. *Consistency of MK-6240 quantification across reference regions.* ***A)*** *Scatterplot of the*
raw mean MK-6240 values in TLE patients and controls across different reference regions. ***B)*** *Correlation of t-*
statistics from between-group difference maps across SUVR reference regions (brainstem, cerebellar gray matter,
and a composite region). The top-right scatterplots show vertex-wise t-statistic comparisons for each pair of
reference choices, along with Pearson correlation coefficients (r), demonstrating high consistency across methods
(r ≥ 0.96). Blue-shaded brain maps illustrate the corresponding reference regions. The bottom-left matrix displays
Spearman's rho values between comparisons. ***C)*** *Group comparisons between patients and controls for each*
reference region. The same mixed-effects model used in the main analyses was applied here, adjusting for age
and sex, and corrected for multiple comparisons using random field theory (pRFT < 0.025). Significant cortical
regions are outlined in white. Below the surfaces the scatter plots display mean SUVR values for significant
regions in each hemisphere by group. Collectively, these results indicate strong agreement in regional t-statistics
across the reference region choices used for MK-6240 SUVR computation.
14

1
2   ***Supplementary Figure 4.*** *Sex differences in MK-6240 uptake. The comparison was evaluated using a mixed*
3   *effects model, controlling for age, and corrected for multiple comparisons with random field theory (p<0.025).*
4   ***A)*** *Group comparisons between only female patients and female controls.* ***B)*** *Group comparisons between only*
5   *male patients and male controls.* ***C)*** *Group comparisons between patients and controls with an interaction of*
6   *group by sex, where higher T-values represent where females have higher uptake than males.*

Supplementary Figure 5. *Tau MK-6240 SUVR and onset age, hippocampal volume, IEDs, GTCS and spatial task. Scatterplots display the relationship of mean significant MK-6240 SUVR with measures not explored in the main manuscript. For the spatial task only 12 out of 28 TLE patients had available data. Brain maps illustrate the corresponding regional effects, dark blue indicating areas where lower values of the corresponding measures are associated with higher MK-6240 SUVR. Mixed-effects models were used to assess these association, with correction for multiple comparisons with random field theory (p<0.025, two tails). No association reached statistical significance.*

1
Supplementary Figure 6. *Group differences and correlations of subcortical and hippocampal MK-6240 uptake.*
The left panel shows the t-statistics from mixed-effects models examining group differences in each subcortical
mean significant MK-6240 SUVR, including an interaction of group by sex and covariates: age, disease duration,
and ipsilateral hippocampal volume (I.hippocampus). The right panel shows Spearman correlations between
subcortical mean significant MK-6240 SUVRs and the same clinical and behavioral variables evaluated in the
cortical MK-6240 analysis. P-values were corrected for multiple comparisons using the Bonferroni-Holm
method; no results reached significance.

A | PET to T1w registration

B | Main processing workflow

Supplementary Figure 7. *PET image processing workflow. **A)** Edge-based registration between PET MK-6240 and T1-weighted (T1w) MRI. Edge detection was applied to both PET and T1w images, and the resulting edge maps were affine-registered using a cross-correlation cost function with histogram matching. **B)** Workflow of co-registration between the structural T1w MRI and the MK-6240 PET image. Normalization by the cerebellar mean SUVR and mapping to the mid-thickness surface of each subject. The right image shows an example of one subject MK-6240 map projected on a standardized surface, fsLR-32k.*

A | Correlogram of MK-6240, behavior and connectome derived metrics

B | Individual structural equations models

1
Supplementary Figure 8. *Impact of the connectome derived metric on the SUVR uptake and behavior.* ***A)***
Correlogram of the connectome derived metrics, behavioral scores and mean MK-6240 from significant clusters.
Orange square represents the correlations between behavioral scores and network derive metrics from structural
and functional connectomes (neighborhood-weighted connectivity, strength, cluster coefficient, efficiency and
characteristic path length). The purple rectangle in the center represents the correlations between network
derived metrics from functional connectomes. The blue rectangle in the bottom-right represents the correlations
between network derived metrics from structural connectomes.
B) *Individual Structural equation models showing relationships between mean significant MK-6240, structural*
network measures (SC weighted neighbors and SC efficiency), and behavioral outcomes. Values represent
standardized path coefficients. Significant paths (p < 0.05) are shown in blue for positive effects and red for
negative effects, the rest is gray. Indirect effects via semantic network measures were tested but were not
significant.

TABLE 4. **MAIN STRUCTURAL EQUATION MODEL SUMMARY**

Regressions	Estimate	Std.Err	z-value	P(>|z|)	Std.lv	Std.all
SC neighbours ~						
MK-6240 (a1)	***0.17***	***0.056***	***3.051***	***0.002***	***0.17***	***0.378***
SC efficiency ~						
MK-6240 (a2)	0.053	0.164	0.321	0.748	0.053	0.045
EpiTrack ~						
SC neighbours (b11)	9.945	8.821	1.127	0.26	9.945	0.534
SC efficiency (b12)	-1.92	3.31	-0.58	0.562	-1.92	-0.267
MK-6240 (c1)	***-4.08***	***1.557***	***-2.62***	***0.009***	***-4.08***	***-0.486***
Episodic ~						
SC neighbours (b21)	9.769	9.801	0.997	0.319	9.769	0.586
SC efficiency (b22)	-3.656	3.481	-1.05	0.294	-3.656	-0.567
MK-6240 (c2)	***-4.148***	***1.456***	***-2.848***	***0.004***	***-4.148***	***-0.552***
Semantic ~						
SC neighbours (b31)	8.348	6.701	1.246	0.213	8.348	0.567
SC efficiency (b32)	-2.158	2.075	-1.04	0.298	-2.158	-0.379
MK-6240 (c3)	-0.921	1.559	-0.591	0.555	-0.921	-0.139

Covariances	Estimate	Std.Err	z-value	P(>|z|)	Std.lv	Std.all
SC neighbours ~~						
SC efficiency	***0.011***	***0.002***	***6.041***	***>0.001***	***0.011***	***0.954***
EpiTrack ~~						
Episodic	***0.457***	***0.174***	***2.624***	***0.009***	***0.457***	***0.373***
Semantic	***0.577***	***0.194***	***2.979***	***0.003***	***0.577***	***0.501***
Episodic ~~						
Semantic	0.194	0.134	1.446	0.148	0.194	0.189

Variances	Estimate	Std.Err	z-value	P(>|z|)	Std.lv	Std.all
SC neighbours	0.004	0.001	6.145	>0.001	0.004	0.857
SC efficiency	0.031	0.005	6.029	>0.001	0.031	0.998
EpiTrack	1.37	0.337	4.065	>0.001	1.37	0.848
Episodic	1.094	0.173	6.328	>0.001	1.094	0.844
Semantic	0.969	0.4	2.425	0.015	0.969	0.957

Regressions	Estimate	Std.Err	z-value	P(>|z|)	Std.lv	Std.all
Indirect EpiTrack > SC neighbours	1.694	1.595	1.062	0.288	1.694	0.202
Indirect EpiTrack > SC efficiency	-0.101	0.633	-0.16	0.873	-0.101	-0.012
Indirect EpiTrack > Total	1.593	1.312	1.214	0.225	1.593	0.19
Indirect Episodic > SC neighbours	1.664	1.844	0.903	0.367	1.664	0.221
Indirect Episodic > SC efficiency	-0.193	0.894	-0.216	0.829	-0.193	-0.026
Indirect Episodic > Total	1.471	1.505	0.978	0.328	1.471	0.196
Indirect Semantic > SC neighbours	1.422	1.202	1.183	0.237	1.422	0.214
Indirect Semantic > SC efficiency	-0.114	0.526	-0.216	0.829	-0.114	-0.017
Indirect Semantic > Total	1.308	1.003	1.305	0.192	1.308	0.197
Total EpiTrack	-2.487	1.339	-1.857	0.063	-2.487	-0.296
Total Episodic	***-2.677***	***0.988***	***-2.71***	***0.007***	***-2.677***	***-0.356***
Total Semantic	0.388	1.166	0.332	0.74	0.388	0.058

1   TABLE 5. POST-SURGICAL TLE CASES

ID	Surgical Procedure	Histology	AT8+	Neuropathology
S1	Right CAH	FCD-II/HS	yes	Focal subpial and few intracytoplasmic neuronal inclusions. Weak subpial staining with negative neurons.
S2	Right AH	FCD-IIA/HS	yes	Subpial positivity and rare isolated neurons with weak staining.
S3	Right fusiform and parahippocampal resection	FCD-?	no	Neurons negative; dot-like edge staining only.
S4	Left CAH	FCD-IIA	yes	Weak subpial stain and rare neuronal positivity. Otherwise, negative with focal background
S5	Right CAH	FCD-IIIA/HS2	yes	Weak subpial staining is present, with a few neurons showing intracytoplasmic positivity, alongside strong neuritic staining in CA4 and focal longitudinal striated staining.
S6	Left amygdalar resection	Cavernoma	yes	Definite intracytoplasmic neuronal positivity around cavernoma; strong tufted-like staining in focal areas

2   *CAH: cortico amygdalo hippocampectomy*
3   *AH: amygdalo hippocampectomy*

Supplementary Figure 9. *Cortical MK-6240 PET and AT8 immunohistochemistry in post-surgical tissue.* Six cases are presented. For each case, cortical MK-6240 SUVR z-scores relative to controls are displayed at the top of each panel (range: −4 to 4). Cortical surface projections are shown in the following order (*left to right*): left lateral, left superior, right superior, and right lateral views (*top row*), followed by left medial, left inferior, right inferior, and right medial views (*bottom row*). Representative post-surgical tissue sections stained with AT8 immunohistochemistry are shown beneath each cortical surface. Panels correspond to individual cases labeled A–F, with multiple sections per case. (A1–A3, B1–B2, C1–C2, D1–D2, E1–E5, F1–F3).