Genome landscape and genetic architecture of recombination in domestic goats (*Capra Hircus*).

Alice Etourneau[1*], Rachel Rupp[1], Bertrand Servin[1*]

[1]GenPhySE, Université de Toulouse, INRAE, ENVT, 31320 Castanet-Tolosan, France

*Corresponding Authors

E-mail addresses:

AE: alice.etourneau@inrae.fr

RR: rachel.rupp@inrae.fr

BS: bertrand.servin@inrae.fr

Abstract

Background

Recombination is a fundamental biological process, both in participating to the creation of viable gametes and as a driver of genetic diversity. Characterising recombination is therefore of strong interest in breeding populations. In this study, we used ~50K genotyped data and pedigree from two French populations (Alpine and Saanen) of domestic goats (*Capra hircus*) to build sex-specific recombination maps, and to explore the genetic basis of two recombination phenotypes: genome-wide recombination rate (GRR) and intra-chromosomal shuffling.

Results

Sex-specific recombination maps showed higher recombination in males than females for both breeds (Alpine autosomal map size = 35.1M in males and 30.5M in females; Saanen map size = 34.0M in males and 29.0M in females). Heterochiasmy is particularly notable on small chromosomes, and at both ends of the chromosomes. Yet, no difference in shuffling has been found between populations. Genetic parameters on recombination phenotypes could only be estimated in males, due to lack of data in females. Both phenotypes are significantly heritable (h^2=0.12 (0.03) for GRR and h^2=0.034 (0.015) for shuffling, when pooling breeds). GWAS on male GRR identified several significant loci, including *RNF212*, *RNF212B* and *SSH1*, the last one being a novel locus for this phenotype. Correlation of SNP effects between breeds is low for both recombination phenotypes (0.25 for GRR and 0.04 for shuffling), indicating different genetic determinants in the two breeds.

Conclusions

This study contributes to the understanding of recombination evolution in ruminants, both between breeds and species.

Background

Recombination is a fundamental biological process in eukaryotes. It is necessary for the proper segregation of chromosomes during meiosis [1] and is also a great driver of genetic diversity by creating new combinations of alleles and avoiding the accumulation of deleterious mutations over time [2,3]. Studying recombination can therefore be important both to understand functional reproduction in a species, and to study evolutionary drivers of genetic diversity.

Of the different approaches to study recombination, pedigree-based studies, as opposed to population studies for instance, have several advantages: first they provide a direct measure of meiotic recombination, essentially unaffected by evolutionary pressures, through the transmission of gametes in the pedigree and second, they allow to assign crossovers to specific individuals and therefore estimate the genetic basis of the recombination process. Pioneered in humans [4,5], the increased availability of cheap genotyping tools has led to a large number of recombination studies, especially in livestock. Indeed, the implementation of genotyping routines within genomic selection programs put at the disposal of researchers both the tools and the data to study recombination in pedigrees. As such, multiple studies were conducted in cattle [6–11], but also in sheep [12], chicken [13], pigs [14–17] and Atlantic salmon [18]. Apart from livestock, recombination in several natural populations with available pedigrees have also been studied: Soay sheep [19], red deer [20,21] and wild house sparrow [22].

This approach of characterising recombination through the use of genotypes and pedigree data allows to estimate recombination rates and build recombination maps of a species. Results have shown that heterochiasmy, or sex-differences in recombination, is widespread in many clades, but its sign varies: in most species like pigs [14–16], humans [4], deer [20], chicken [13] or Atlantic salmon [18], females recombine more than males; while in other species like cattle [8–

11] or sheep [19], males show higher recombination than females. [22] and [23] illustrate this

inconsistent heterochiasmy in birds and fishes, respectively.

This high diversity in recombination features shows the relevance of studying the

recombination process itself. One way to study this process is through genetics, which allows

to estimate heritability and find associated QTLs for phenotypes such as recombination rate

and, more recently, intra-chromosomal shuffling [24]. The genetic control of recombination

rate, despite showing common basis between species, was also shown to be very diverse, even

between several populations or the two sexes of the same species. Focusing only on ruminants,

heritability for phenotypes of recombination intensity varies between 0.04 in Norwegian Red

cattle [11] and 0.22 in Dutch Holstein-Friesian cattle [6]. But variations in heritability can be

observed between species and, most notably, sexes within a species [9,12,19,21]. Identified

QTL for these phenotypes can also vary between populations and species. Cattle [6,8–11],

sheep [12,19] and red deer [21] all show a signal in the region of the genes

RNF212B/HEI10/REC8/REC114, four genes in very close proximity to each other in these

three species. In cattle [6,9] and sheep [12,19], associations with the *RNF212* gene have also

been found. Yet despite these similarities, genes like MSH4 or PRDM9 have only been detected

in cattle among ruminants, although this is possibly due to the increased statistical power in this

species. When studying genetic control within sexes and breeds separately in a species, the

differences are even starker. In sheep, [19] found a QTL at RNF212 only in Soay females, and

not in males. Yet [12] actually found a QTL for this same gene in males from another

population, the Lacaune. In cattle, [8] identified thirteen loci associated with recombination

intensity, and only three of them were shared between sexes. In [9]'s bovine population, four

of their ten detected QTLs are sex-specific. And once again, [11] found sex differences in

Norwegian red dairy cattle.

In this context of strong genetic diversity in very related groups, we present a new study on

another ruminant species: the goat. In goats, the last genetic map was built with a linkage

methodology in 1998 [25]. At the time, its primary goal was still to understand the organisation

of genes between one another, and the map was never updated since. Genome-wide

recombination rate was estimated through the use of cytogenetics techniques by counting

MLH1 foci [26,27] but never confirmed using other methods.

In 2010, the first reference genome assembly was built for the domestic goat [28] and in 2013,

an international goat genome consortium was created to acquire more genomic data and

investigate genetic variations in the species. This lead to the development of a 50k SNP chip

[29] and later on, the creation of a sequence-level SNP panel of more than a thousand goats

from all over the world [30].

In France, a goat breeding program for two breeds, Alpine and Saanen, has been in place since

the 1960s, its main goal being to improve milk production for the purpose of cheese-making

with the progressive inclusion of health and udder morphology characteristics. Since 2014 and

the availability of the 50k SNP chip, selection candidates are now systematically genotyped.

Furthermore, despite the fact that the breeding program is male-oriented, females have started

to be genotyped as well in the last few years, both as part of the breeding program and within

experimental programs. Therefore, we took advantage of genotyping and pedigree data

available in the two Alpine and Saanen breeds to study recombination in this ruminant species

using a pedigree-based approach, for the first time to our knowledge. In an opportunity to

examine heterochiasmy and breed differences in this new species, we built updated, sex-specific

recombination maps, developing novel methodologies to quantitatively measure sex differences

in recombination patterns and take into account uncertainty about crossover detection. In

addition, we also studied the genetic basis of two recombination phenotypes in both available

breeds, by estimating genetic parameters and performing genome-wide association studies.

Methods

Study population and genotype data

In this work, we exploited pedigree and genotyping data of 7,588 goats born between 2008 and 2019. They amount to 3,169 males and 4,419 females from both the Alpine (5,051 samples) and Saanen (2,537 samples) breeds. The animals were genotyped with versions 1 and 2 of the Illumina Inc. GoatSNP50 chip. Only autosomal SNPs and markers common to both arrays were kept, resulting in 53,347 SNPs.

Quality control of physical marker ordering

Errors in the physical ordering of markers can lead to detect spurious recombination events. Recombination studies therefore require a robust marker ordering. Here, this ordering was built from two independent physical maps of the SNPs: the reference genome assembly ARS1 and the Radiation Hybrid (RH) map of [31]. We decided to keep only SNP markers that have a consistent ordering on both physical maps. This was done by identifying, for each chromosome, the longest increasing subsequence (LIS)[32] in the mapping of the RH order to the assembly order. This requires knowing the order of the SNP markers on both maps. For the reference genome ordering, genomic positions of the 53,347 v1 autosomal markers of the GoatSNP50 Illumina chip were obtained from the ARS1 assembly as described on the International Goat Genome website (http://www.goatgenome.org/projects.html). For the RH map, additional analysis was required as it was built using SNP array of different species (ovine and bovine SNPs herein called external SNPs). We mapped the external SNPs to the ARS1 reference genome and then projected the position of each caprine SNP on the RH map from the positions of its two flanking external SNPs. For goat markers located at the extremities of the

chromosome, which had only one flanking external SNP and could not be projected on the RH

map, we assumed the reference genome order was correct (but see below for further data quality

control steps). **Figure S1 (See Additional File 1**) illustrates this quality control step on two

chromosomes. After this step, 54,013 markers were kept.

Furthermore, an initial detection of crossovers on the genome (using methods detailed below)

revealed 7 genomic regions (123 SNP markers in total) that showed extremely high

recombination rates (See example on chromosome 1 in **Additional File 1, Figure S2**). Detailed

analysis of the synteny conservation with the human genome in these regions showed that they

corresponded systematically to conserved synteny breakpoints. As these events were most

probably some remaining unidentified assembly errors, markers involved were discarded in

downstream analysis. For a list of these markers, see **Additional File 2, Table S1**.

Genotype quality control

Standard data cleaning procedures were applied using PLINK 1.9 [33]. Animals and SNPs with

call-rate below 99% were removed (**See Additional File 1, Figure S3**). Monomorphic SNPs in

the dataset were removed. No MAF filter was applied at this step as even a small MAF may be

informative for crossover detection in families polymorphic for these markers. Markers with

strong departures from Hardy-Weinberg genotype proportions ($p < 10^{-7}$) within each breed

were removed. Note that families with a sire having more than 50 offspring were not used to

compute the Hardy-Weinberg equilibrium (HWE) test as fathers carrying rare alleles in such

designs can have a great impact on testing for HWE.

Mendelian errors were tested using the *mendel* command of the Yapp software version 0.3 [34].

It evaluated the number of incoherent markers between parent and offspring and automatically

corrected the kinship relationships in the pedigree which were beyond the threshold fixed by

the software.

After these quality control steps, 46,081 markers and 7,457 animals, including 4,520 Alpine

and 2,937 Saanen, were left.

Data phasing and crossover detection

Genotype phasing and crossover detection were done using Yapp version 0.4. First, the *phase*

command was used in order to phase genotypes. In order to reconstruct the paternally and

maternally inherited haplotypes of each individual (i.e. to infer the phase of heterozygous loci),

Yapp uses a set of discrete steps. First, using all available genotypes, it identifies the

unambiguous transmissions from parents to offspring, *e.g.* the alleles transmitted by a

homozygous parent or received by a homozygous offspring are easily phased. This leads to a

set of partially resolved phases. Second, for each parent in the pedigree, from oldest to newest,

it improves the inference of parental phases by combining all partially resolved transmitted

haplotypes using the algorithm of [35]. Third, after this first pass in the pedigree, the transmitted

haplotypes from parent to offspring are improved using a Hidden Markov Model along the

whole chromosome similar to the approach of [36] and segregation indicators are inferred.

Then, the *recomb* command was used among a total of 6,434 meioses to detect crossovers. A

crossover in the parent meiosis is defined as a change in phase observed in the offspring, on the

inherited chromosome from the parent. Crossovers are detected between two heterozygous

SNPs and can only be detected between the first and the last phased, heterozygous SNPs.

Meioses and crossovers were quality-controlled to ensure that only truly informative data was

retained. All meioses with a parent or offspring phased at less than 90% of markers were

removed. Furthermore, meioses with a parent that itself had none of its parents genotyped and

only one offspring, were removed, as it is impossible to detect crossovers in such a setting.

Distance between two crossovers on the same chromosome was calculated for every meiosis.

As crossovers are detected within an interval, distance between crossovers was calculated as

the distance between the closest interval boundaries. All double-crossovers closer than 5 Mb

from one another were considered false-positives due to phasing errors and both were removed

(**See Additional File 1, Figure S4**). After filtering, the dataset contained 6,204 meioses and

204,727 crossovers.

Recombination maps

Recombination maps were constructed by estimating recombination rates in non-overlapping

intervals of one megabase (Mb) along the genome. The methodology for estimating

recombination rates was adapted from [12]. For a given genomic interval i and a meiosis m,

the count of the number of crossovers Y_{mi} is modelled using a Poisson distribution whose

parameter (measured in Morgans) depends on the recombination rate c_i (in centiMorgans per

Mb, cM/Mb) assumed constant in the interval and across meioses, and the length of the genomic

interval l_{mi} (in Mb) that can depend on the meiosis (see below).

Then, the expected number of crossovers in an interval is $0.01c_i l_{mi}$ and Y_{mi} is modelled as:

$$Y_{mi} \mid c_i \sim Poisson(0.01c_i l_{mi}) \qquad (\,1\,).$$

194   As in [12], we define a prior on c_i as:

$$c_i \sim \Gamma(\alpha, \beta) \quad (\,2\,).$$

196   As in [12], we used an empirical Bayes approach to set α and β from the genome-wide

197   distribution of Maximum Likelihood Estimates of recombination rates ($\alpha = 3.02$ and $\beta =$

198   2.32, see **Additional File 1, Figure S5**). Given the conjugacy of Poisson and Gamma

199   distributions, the posterior distribution for c_i is then:

$$c_i \mid Y_{.i} \sim \Gamma(\alpha + \sum_m Y_{mi}, \beta + 0.01 \sum_m l_{mi}) \qquad (\,3\,).$$

201   Stemming from the Bayesian model $(\,3\,)$, the expected value for the Gamma law was

202   used to estimate the chromosomic recombination rate:

203
$$\widehat{c_{chr}} = \frac{\alpha + \sum_m Y_{chr,m}}{\beta + \sum_m l_{chr,m}} \quad (4),$$

204   With α and β the prior parameters and m a meiosis. c_{chr} was estimated for both breeds and

205   sexes, and a model explaining c_{chr} by the log of the chromosome size was fitted for all

206   populations.

Meiosis Informativity

In the equations above, the length of the interval is allowed to depend on the interval and the

meiosis considered. This is to account for the variation in the informativity of the data for the

detection of crossovers. First, because samples are genotyped on a 50k SNP chip, no phasing

information is available for the tips of chromosome before and after the first and the last SNP

of the chromosome, which makes detecting crossovers impossible in these regions. Second, for

a given meiosis, only SNPs that are heterozygous and phased in the parent are informative for

crossover detection. Hence for a particular meiosis, this can further reduce the portion of the

genome where crossovers can be detected (**Fig. 1**).

[Figure 1 here]

Accounting for uncertainty in crossover locations

Crossovers are not determined at a specific position but resolved within genome regions that

vary in size depending on the parent and offspring genotypes. The average length of these

intervals in our data is 1,4Mb with a 95% range of 0.1Mb to 10.2Mb (**See Additional File 1,**

Figure S6). To account for this uncertainty in the position of crossovers, we used a Monte Carlo

approach as in [12]: we first sampled crossover positions uniformly in the detection intervals

and then the recombination rate from its posterior (Equation 4). This was done 100 times,

providing for each interval 100 samples from the posterior distribution of recombination rate.

Let us name $E(c_i|Y_{\cdot ik}) = b_{ik}$ the expected value of posterior k for genomic interval i. The

expected value of all combined posteriors for interval i is therefore:

$$\hat{c_i} = E(c_i|Y_{\cdot i \cdot}) = E_k\big(E(c_i|Y_{\cdot ik})\big) = E_k(b_{ik}) \qquad (5),$$

And according to posterior variance law:

$$Var(\hat{c_i}) = E\big(Var_k(c_i|Y_{\cdot ik})\big) = Var(c_i) - Var_k(E(c_i|Y_{\cdot ik}) = Var(c_i) - Var_k(b_{ik}) \qquad (6).$$

Thus, recombination rate c for genomic interval i was estimated as the mean of the 100

posteriors' expected values.

Assessing sex differences in local recombination rates

The Bayesian model described above is interesting in that it allows to account naturally for the

uncertainties in the data and provides estimates of recombination rates even in regions with

little information. Using this approach, we could in principle evaluate the sex differences in

recombination rates from the posterior samples obtained by applying the model separately in

both sexes. However, when interrogating the difference in local recombination rates between

sexes, the model does not allow to separate the effect of differences in rates and those of

differences in uncertainty. Therefore, the two sexes could be deemed to have different rates due

to the amount of data in each sex being unbalanced (for example female estimates are typically

more influenced by the prior than male estimates). To deal with this issue, we derived a

permutation scheme that can separate both effects.

Our measure of heterochiasmy in an interval i is:

$$\hat{d_i} = log2(\widehat{c_{Mi}}) - log2(\widehat{c_{Fi}}) \qquad (7).$$

That is, the log-ratio of male to female recombination (in base 2 for easier interpretation).

To account for the difference in data informativity between sexes, we permuted the sex of the

parents 1,000 times to obtain a sample of $\mathbf{d_i}$ under the null of no difference between sexes. As

permutations do not change the number of male and female meioses, they allow to account for

difference in informativity between sexes. The resulting distribution of $\mathbf{d_i}$ in the permutation

samples were very close to normally distributed (**See Additional File 1, Figure S7**), so in each

interval the null distribution of $\mathbf{d_i}$ can be described by a normal distribution with parameters set

at the sample mean and variance. This in turn allows to derive p-values for a difference between

sexes in each interval. Finally, to account for multiple testing across intervals, the False

Discovery Rate (FDR) approach of [37] using the R "qvalue" package was applied on the p-

values and the significance threshold set at 0.05.

Recombination phenotypes

The crossover dataset described above offers the opportunity to study the individual variation

in some aspects of recombination (i.e. recombination phenotypes) and interrogate their genetic

architectures. However, because we had very little data in females this was only possible to do

in males. Two phenotypes were studied: the individual Genome-wide Recombination Rate

(GRR) and the intra-chromosomal shuffling.

The intensity of recombination in a meiosis can be measured by summing the crossover count

on all autosomes (Autosomal crossover count (ACC)). However, as explained above, the

informativity of marker data varies between meioses and can influence the value of ACC. To

take this into account, rather than the ACC as a measure of recombination intensity we use the

genome-wide recombination rate (GRR), defined as the ACC divided by the informative

portion of the genome (in Mb).

The other phenotype we studied is a measure of the efficacy of a meiosis to create new allelic

combinations in the offspring: the rate of intra-chromosomal shuffling $\bar{r}_{intra}$ [24]. It is defined

as the probability that two randomly chosen loci uncouple their alleles during meiosis and is

calculated as:

$$\overline{r}_{intra} = \sum_{k=1}^{n} 2p_k(1 - p_k)L_k^2 \qquad (\,8\,),$$

Where for chromosome k, p_k is the proportion of SNPs inherited from one parent of the focal

animal (or grandparent of the offspring), L_k is the chromosome's proportion in the autosomal

genome, and n is the number of autosomes.

276 **Heritabilities**

Heritabilities of the GRR and $\overline{r}_{intra}$ phenotypes were estimated using all males with a

calculated phenotype, i.e. 520 parents accounting for 5,707 meioses (**Table 1** for detail).

To estimate heritability, variance components were estimated by restricted maximum likelihood

(REML) using the Wombat software [38]. An animal model including a permanent

environment effect was fitted:

$$\mathbf{y} \sim \mathbf{X\beta} + \mathbf{Za} + \mathbf{Zu} + \mathbf{e} \qquad (\,9\,).$$

$\mathbf{y}$ is the vector of observations (GRR or $\overline{r}_{intra}$); $\boldsymbol{\beta}$ is the vector of fixed effects, $\mathbf{a}$ is the vector

of direct animal (additive) genetic effects; $\mathbf{u}$ is the vector of animal permanent environment

effect; $\mathbf{e}$ is the vector of residual effects; $\mathbf{a} \sim N(0, \mathbf{A}\sigma_A^2)$, $u \sim N(0, \mathbf{I}\sigma_B^2)$ and $\mathbf{X}, \mathbf{Z}$ are the

incidence matrixes. The pedigree matrix $\mathbf{A}$ was computed on five generations of the genealogy.

The model was applied to all breeds separately or to the entire population at once. A fixed effect

for breed was therefore added where necessary.

Once the variance components had been estimated, values were incorporated into the model as

fixed values to obtain the Estimated Breeding Values (EBV) for each phenotype. Due to the

absence of differences in estimated heritability between breeds (see Results), EBVs were

estimated by pooling samples from both breeds. Note that although we did not fit a model for

female recombination phenotypes, EBVs can still be predicted for females as they are connected

to males in the pedigree.

Genome-wide association studies

Genotype Imputation

As an imputation reference panel, we used the whole-genome sequence VarGoats data

presented in [30] and retrieved from https://www.goatgenome.org/home.html. This data

consisted of 1,372 goats out of which 252 European samples closely related to our study

samples were used (from the Alpine, Fossés, Lorraine, Poitevine, Provençale, Pyrénées, Rove,

Savoie and Saanen breeds).

SNP filtering was then applied on the resulting VCF based on several criteria, using bcftools

(http://www.htslib.org/doc/1.1/bcftools.html). First, we removed SNPs having discordant

reference alleles between the SNP array manifest and the resequencing data, and SNPs showing

strictly less than two counts of the minor allele across all selected samples or with a proportion

of missing genotypes greater than 10%. Then, SNPs were filtered based on the distribution of

the following GATK 3.6 [39] variant-calling quality parameters: DP, FS, VQSLOD (**See**

Additional File 1, Figure S8). The chosen threshold for SNP removal were DP < 10,000; DP

> 25,000; FS > 50; VQSLOD < 10 and VQSLOD > 25. The resulting number of SNPs

remaining was 22,741,231 out of which 43,358 SNPs are present on the V1 GoatSNP50

Illumina chip. This corresponds to removing 5.6% of the chip SNPs.

The Beagle4 software [40] was used to impute the few missing genotypes of the sequence panel

and correct the called genotypes by using the genotype likelihood (*gl*) method, a modelscale of

1.5 [41], an effective population size of 1000 and the recombination maps estimated using a

pedigree-based approach *(see above)*. To assess the quality of the resulting panel genotypes, we

leveraged information on 91 animals for which both sequencing and SNP genotyping data were

available: we computed R² and genotype concordance before and after imputation. Both metrics

were highly improved by the imputation procedure: mean R2 increased from 0.928 to 0.990

and mean genotype concordance from 0.974 to 0.997. (**See Additional File 1, Figures S9 &**

S10). Phasing of the reference panel was done using ShapeIt4 [42]. As some animals from the

reference panel were previously phased using pedigree information, their phased haplotypes for

the SNP chip markers were used as scaffold reference for phasing with ShapeIt4. The resulting

VCF file, phased and imputed, was used as reference panel for imputation.

Shapeit4 was also run on the phased VCF of 7,457 samples produced by Yapp from the SNP

array data to solve the few unphased genotypes and impute the few remaining missing

genotypes. The resulting fully phased 50k chip data was imputed to sequence using the Impute5

software [43]. Recombination maps were provided to increase power and N_e was set at 1000.

Impute5 provides a measure of confidence for all imputed markers (imputation R2) which was

used in later steps to filter SNPs *(see below)*.

Statistical Analysis

GWAS were performed separately in each breed using the predicted EBV for male GRR and

$\overline{r}_{intra}$ as phenotypes: 589 samples were used in Alpine and 254 in Saanen (Table 1). SNPs with

MAF<5% and imputation R2<0.3 were removed from the analysis leaving N=15,988,092 and

15,734,503 SNPs in Alpine and Saanen respectively. A univariate linear mixed model was used

for testing SNP association with the phenotype using the Gemma software [44], with the "leave

one chromosome out" (LOCO) strategy (*i.e.* the genetic relationship matrix (GRM) used to

analyse a chromosome is computed on all other chromosomes except this one).

Significance was assessed using the ash method [45] implemented in the "ashr" R package, by

calculating s-values for each SNP. In the same way as for q-values [37], thresholding s-values

at level α gives a set of significant features so that a proportion of α is expected to be false-

positives. While [37]'s FDR gives indication on the confidence that effects are different from

zero, [45]'s False Sign Rate (FSR) gives indication on the confidence of the *sign* of effects. As

this method takes into account the β estimates as well as their standard errors, the resulting s-

values can induce a reranking of SNPs compared to methods only taking p-values into account.

All SNPs with s-value<0.01 were declared significant and SNPs with s-value<0.05 were

declared suggestive.

Correlations of SNP effects between breeds were measured by the correlation of β estimates

for SNPs that were significant in at least one of the two breeds, and using standard errors as

weights.

Annotation of GWAS loci

A GWAS locus was defined from pairwise Linkage Disequilibrium (R2) between all significant

SNPs on a given chromosome. R2 calculation was done using the PLINK 1.9 software.

Significant SNPs were considered to belong to the same peak if their R2 was equal or superior

to 0.2. For the lead SNP of each peak, a high LD area was defined as follows: an area whose

boundaries correspond to the SNP furthest away in R2>0.6 from the lead variant, on both sides

of the lead variant. We aimed at evaluating the evidence for association of all genes present in

Ensembl (gff3 release 111). As this annotation did not include two major genes associated with

the recombination process, *RNF212* and *PRDM9*, we manually added them to the annotation

file after blasting their bovine gene sequence on ARS1 (**See Additional File 3, Tables S2 &**

S3).

For all genes, we evaluated the following criteria: if the gene is within 2Mb of the lead variant;

if it is the closest gene to the right, to the left or both; if it is the closest gene to another

significant SNP of the peak and if it belongs within the "R2>0.6" area of the peak. Based on

this we calculated a "candidate score", ranging from 0 to 5, as the number of criteria satisfied

by the gene.

Results

Recombination landscape

After quality control, genotype data on 7,457 animals led to the detection of 204,727 crossovers

in 6,204 meioses from 843 parents. As a large portion of the dataset comes from a dairy

selection program, most genotyped parents were males. As males also typically have more

genotyped offspring than females, there are 10 times more male crossovers than female ones in

the recombination data (Table 1). The software used to detect crossovers provides the

information on the parts of the genome that are informative for the detection of crossovers (See

Methods). Overall, all meioses had more than 94% of the genome informative for crossovers

(**See Additional File 1, Figure S11**). Females had slightly more informative meioses than

males (mean=0.99% vs 0.98%). Indeed, although females are usually less well connected in the

pedigree than males, the ones that passed our quality control criteria for meioses (**see Methods**)

tended to be better connected than the average male.

Table 1: Summary table of data after all genotype and crossover quality control steps.

Breed	Sex	Headcount	Parents	Meiosis count	Crossover count
Alpine	M	1,739	327	3,562	120,057
	F	2,781	262	386	11,329
Saanen	M	1,339	193	2,145	70,307
	F	1,598	61	111	3,050
Total		7,457	843	6,204	204,727

Chromosome-wide rates

A linear model was used to test differences in genetic map size between populations. Breed and

sex parameters were found significant (sex effect p-value = 10^{-61}; breed effect p-value =

10^{-13}; interaction p-value = 0.52). The size of the genetic map was estimated at 35.1M for

Alpine males, 34.0M for Saanen males, 30.5M for Alpine females and 29.0M for Saanen

females (**See Additional File 1, Figure S12**). Therefore, the genetic map is slightly shorter in

Saanen than in Alpine and much shorter in females than males.

The mean recombination rate of each chromosome was estimated with the Bayesian model, and

a linear model depending on the log size of the chromosomes was fit on the estimates (**Fig. 2a**).

Again, a significant difference between breeds was found (p-value = 0.01) but, as the effect is

very small, the figure shows breed-averaged estimates to avoid over-plotting. A significant

difference between sexes is also observed (p-value $\approx 10^{-48}$): males overall have a higher

chromosomic recombination rate than females, and the recombination rate is greater on small

chromosomes than big chromosomes for both sexes. Overall, the chromosomic recombination

rate has a log relation with the chromosome size. It is observed that the difference in

recombination between the sexes is greater on small chromosomes than on big chromosomes,

on which recombination rates are almost alike.

Chromosome landscape was calculated from 1Mb resolution Bayesian estimates of

recombination rate: all positions in Mb were converted to positions relative to the centromere,

and recombination rates of all chromosomes were studied depending on their position to the

centromere. Considering that all goat autosomes are acrocentric, the beginning of all

chromosomes (or centromeric tip) is located at 0%, and the end (or telomeric tip) is located at

100%. The results are presented in **Fig. 2b**. Females have a much flatter recombination pattern

along the chromosomes compared to males, and both sexes seem to recombine as much in the

middle of chromosomes.

Local recombination rates

Recombination rates were estimated locally on 1Mb intervals on all autosomes (**See Additional**

File 4, Table S4). Differences between sexes were estimated as the log ratio of male to female

recombination rate and significance was assessed using its empirical null distribution calculated

from permutations (see Methods). Recombination maps for each chromosome are available in

Additional File 5 (**Figures S18 to S46**), and **Fig. 2c** illustrates this inference on chromosome

25. It exhibits the general trend of the sex differences in recombination patterns where males

recombine significantly more than females at the tips of the chromosomes. As all goat

autosomes are acrocentric, this means that male recombination rate is higher not only at the

telomeric end, but also at the centromeric end. A total of 871 intervals of the genome are

significant for sex differences (out of 2477 intervals). Of the significant intervals, 559 (64% of

the significant intervals) show higher recombination rate in males and are localized for the most

part at both ends of chromosomes. The 312 remaining significant intervals (36% of the

significant intervals) show higher recombination rate in females and are located in the middle

of chromosomes. For a list of all significant intervals, see **Additional File 6, Table S5**.

[Figure 2 here]

Genetic parameters and architecture of recombination phenotypes

Phenotype distributions

Two phenotypes were used to characterize the genetic basis of recombination in the goat: the

individual Genome-wide Recombination Rate (GRR), a measure of recombination *intensity* and

intra-chromosomal shuffling ($\overline{r}_{intra}$) a measure of the *distribution* of crossovers on

chromosomes. This is because crossovers located in the middle of a chromosome lead to create

more novel allelic combinations and therefore increase intra-chromosomal shuffling while

crossovers located at the tips of a chromosome decrease intra-chromosomal shuffling.

GRR is defined as the per meiosis crossover count divided by the length of informative genome.

The average value of GRR over all meioses is 1.39 cM per Mb. The distributions of GRR for

different breeds and sexes are shown in **Figure S13** (**See Additional File 1**).

Intra-chromosomal shuffling $\bar{r}_{intra}$ was defined as stated in Material & Methods. The average

value of $\bar{r}_{intra}$ is 0.011. The distributions of $\bar{r}_{intra}$ between breeds and sexes are shown in

Figure S14 (See Additional File 1). A linear model was fitted to estimate differences between

populations. A representation of the estimates can be found in **Figure S15 (See Additional File**

1). No significant difference was found between sexes (p-value = 0.12) but Alpine goats had

significantly higher $\bar{r}_{intra}$ (p-value = 0.03). Furthermore, the interaction between breed and sex

was found significant (female Alpine $+4.01 \times 10^{-4}$; p-value = 0.03), meaning that in average,

Alpine females shuffled more than Alpine males, but Saanen females shuffled less than Saanen

males.

Heritability estimates

Heritabilities of GRR and $\bar{r}_{intra}$ were estimated using an animal model taking into account

repeated measures, only for the male phenotype. GRR heritability was estimated at 0.12 ± 0.03

and was consistent in both breeds. $\bar{r}_{intra}$ heritability was estimated at 0.034 ± 0.015 when both

breeds were analysed together. No significant permanent environment effect was detected for

any phenotype (**Table 2**).

Table 2: Variance components and heritability (SE) estimates for male genome-wide

recombination rate and intra-chromosomal shuffling in a population of 843 Alpine and

Saanen goats.

Phenotype	Population	σ_A^2	σ_B^2	σ_E^2	τ^2	h^2
GRR	All	0.50 (0.12)	0.05 (0.09)	3.52 (0.07)	0.01 (0.02)	0.12 (0.03) [a]
	Alpine	0.47 (0.16)	0.11 (0.13)	3.57 (0.09)	0.03 (0.03)	0.11 (0.04) [a]
	Saanen	0.47 (0.17)	0.00 (0.13)	3.42 (0.11)	0.00 (0.03)	0.12 (0.04) [a]
$\bar{r}_{intra}$	All	0.002 (0.001)	0.000 (0.001)	0.052 (0.001)	0.000 (0.01)	0.034 (0.015) [a]
	Alpine	0.002 (0.001)	0.000 (0.001)	0.053 (0.001)	0.004 (0.019)	0.033 (0.019)

	Saanen	0.001 (0.001)	0.000 (0.001)	0.051 (0.002)	0.000 (0.021)	0.027 (0.020)

450 Heritability was considered significant when (h²±2SE) did not include 0. τ^2 = permanent environment effect; [a] =

451 significant heritability. With σ_A^2 the genetic variance; σ_B^2 the permanent environment variance and σ_E^2 the residual

452 variance.

Genome-Wide Associations

The EBVs predicted from the animal model were used as phenotypes for the GWAS on GRR

and $\bar{r}_{intra}$. GWAS results using imputed genotypes to the sequence level are illustrated in **Fig.**

3 for Alpine and Saanen, on both phenotypes.

The analyses on GRR revealed 15 significant QTLs in Alpine and 4 suggestive QTLs in Saanen.

For Alpine, there were one QTL on chromosomes 1, 6, 7, 13 and 29; three QTLs on

chromosomes 5 and 10; as well as two QTLs on chromosomes 9 and 17. For Saanen, there were

one suggestive QTL on chromosomes 2, 4, 6 and 14. There is no overlap between the significant

QTLs in Alpine and the suggestive QTLs in Saanen. Correlation of SNP effects between the

two breeds for this phenotype was calculated using standard errors as weights. It amounted to

0.25.

The analyses on $\bar{r}_{intra}$ revealed 7 significant QTLs in Alpine and 41 suggestive QTLs in

Saanen. Correlation of $\hat{\beta}$ between the two breeds is 0.04. For Alpine, there were one QTL on

chromosomes 1, 9 and 19; as well as two QTLs on chromosomes 8 and 10. The lead variant on

chromosome 9 is 5kb away from the closest significant QTL for GRR GWAS, and the one on

chromosome 10 is 268bp away from its respective significant QTL for GRR. Correlation of the

$\hat{\beta}$ between the GWAS for GRR and $\bar{r}_{intra}$ is 0.89. For Saanen, there were five QTLs on

chromosome 1; two QTLs on chromosomes 2, 4, 6, 8, 14, 17, 18, 19, 20, 21 and 24; one QTL

on chromosomes 3, 5, 7, 9, 11, 13, 22 and 28; as well as three QTLs on chromosome 10 and

26. Despite some QTLs being on the same chromosomes as for GRR, no suggestive region is

within 2Mb of suggestive QTLs for GRR. Correlation between the two GWAS in Saanen is

0.92.

[Figure 3 here]

Considering the GWAS on GRR, the most significant QTL in Alpine is located on chromosome

17 at around 7Mb (lead variant s-value=0.00236) and is illustrated in **Fig. 4a**. In this region, 58

candidate genes can be found, including *SSH1* which scores 5/5. The significant SNPs are

located directly within the *SSH1* gene, in an intronic region. The lead variant has a MAF of 0.49

in the population. The QTL illustrated in **Fig. 4b** contains significant SNPs in high LD in a

region of 300kb and is rich in genes. It is the second most significant QTL (lead variant s-

value=0.00238). Our standardised method for ranking candidate genes identifies 86 genes with

a score of at least 1/5. The highest scoring (3/5) is *CARMIL3* because of its physical proximity

to the lead variant. Among the candidates for this region are two genes previously associated

with recombination: *RNF212B* (rank 17, score 1/5) and *REC8* (rank 51, score 1/5). The QTL

illustrated in **Fig. 4c** is located at the end of chromosome 6 and is the 6th most significant of the

genome. It contains 36 candidate genes including *SPON2* (rank 1, score 4/5) and *RNF212* (rank

2, score 3/5).

In Saanen, the most suggestive QTL can be found on chromosome 6 at around 12.5Mb (lead

variant s-value=0.0326) and is illustrated in **Fig. 4d**. Several significant SNPs of the area fall

within the *ANK2* gene which scores at 4/5 as a result. Eight other candidate genes can be found

in the region. Other notable genes in both breeds can be found in Table 3. For exhaustive results

for GWAS candidate genes in both breeds, see **Additional File 7**, **Tables 6 & 7**.

Table 3: Genes scoring above 4/5 for the GWAS on EBVs of male GRR in Alpine or Saanen.

Breed	QTL of interest (chromosome: position of the lead variant)	Candidate gene (score/5)
Alpine	1:156523831	KCNH8 (5/5)

	5:94902996	GRIN2B (5/5)
	5:106271734	B4GALNT3 (5/5)
	6:117176129	SPON2 (4/5)
	7:105398227	SNAP47 (4/5)
	9:53226533	THEMIS (5/5)
	10:1686345	EFCAB11 (5/5)
	13:17395272	CCNY (4/5)
	17:6965601	SSH1 (5/5)
	17:7730110	GLTP (5/5)
	29:42944888	FERMT3 (5/5)
	2:29863833	TNS1 (5/5)
Saanen	6:12545797	ANK2 (4/5)
	14:91668757	B3GAT2 (5/5)

Considering the GWAS on $\overline{r}_{intra}$, the most significant QTL is the first signal of chromosome

8 in Alpine (lead variant s-value=0.0015). The gene *PTPRD*, which scores 4/5 for this region,

has no known link with the recombination process. Another significant QTL can be found for

this phenotype and breed at the end of chromosome 1 (**Fig. 4e**). The highest scoring candidate

genes are *ERG* (4/5), *ETS2* (3/5) and *KCNJ15* (2/5), but none of them seem to be involved in

recombination either. The region is 6Mb away from *PRDM9*. The most significant QTL for

$\overline{r}_{intra}$ in Saanen can be found on chromosome 4 (lead variant s-value=0.0026). The gene *AVL9*

scores 5/5 for this region but does not seem to be involved in the recombination process. For

exhaustive candidate genes results for GWAS on $\overline{r}_{intra}$, see **Additional File 7, Tables 8 & 9**.

[Figure 4 here]

Discussion

This study is a detailed analysis of the recombination process and its genetic architecture in a

new species: the goat (*Capra Hircus*). It provides precise estimates of recombination rates along

the genome, using a new statistical inference approach that accounts for different levels of

uncertainty in crossover data. This analysis revealed that the genetic length of the ARS1 goat

genome assembly is 35.1M (Alpine males) corresponding to a genome-wide recombination rate

of 1.42cM/Mb. However, recombination was found to be highly variable along the genome,

with rates varying from 0.21 to 10.32cM/Mb. An important feature of the recombination

landscape is the large heterochiasmy in some genomic regions, mostly located at chromosome

ends. This is consistent with patterns of recombination in a closely related species, the sheep

Ovis aries [12,19].

In a second part of this study, we characterized the genetic architecture of recombination

intensity and localisation through the study of two recombination phenotypes, GRR and $\overline{r}_{intra}$.

This analysis revealed low but significant genetic components to these traits and identified

several QTLs for both traits, some of which match loci found in other species while others are

new. Overall, our study adds to the growing knowledge on recombination and will contribute

to understanding the evolution of recombination, a fundamental trait for individual fitness and

adaptive potential of populations. We now discuss the importance of our findings and the future

research questions it suggests.

Studying recombination from livestock breeding programmes data

Livestock breeding populations are good models to study the genetic architecture of the

recombination process as they offer large datasets of genotyped individuals related through well

characterised pedigrees. An important caveat of these datasets however lies in the unbalanced

genotyping between sexes. In ruminant programs of genomic selection such as the one used

here, male selection candidates are typically all genotyped while females are much less likely

to be, although this tends to change with the decrease in genotyping costs. In our dataset, the

number of dams was very small. We still managed to correctly characterise heterochiasmy

along the genome, but the number of dams was too small to study the genetic architecture of

recombination in this sex. We expect the number of genotyped females to grow in the future

which will allow studying genetic determinants of heterochiasmy in this species. However, it

should be noted that offspring size will still typically be smaller for females, so that taking into

account uncertainty as we did here will remain relevant.

Recombination landscape of the goat genome

Our study employs a similar strategy as previous studies in animal populations [8,10–12,16,18–

20,22,46], but we aimed at better accounting for the uncertainty in crossover data. The first

source of uncertainty arises from the unprecise localisation of crossovers due to the genotyping

density, the second by the homozygosity of individuals in the pedigree, and the third concerns

only female recombination rate estimation, as our available female data is much scarcer. To

tackle the first problem, [12] proposed a random sampling approach of crossover locations to

average recombination rate estimation over crossover uncertainty. Here, we used the same

strategy but extended it further to better identify genomic regions with significant differences

in recombination rates between sexes. To tackle the second problem, we made use of an

information provided by the software used to phase individuals (Yapp) that explicitly identifies

regions of the genome where meioses are uninformative for crossover detection. In order to

integrate both sources of uncertainty, we leveraged the Bayesian Gamma-Poisson model of [12]

that allows to incorporate this information naturally. One caveat of this model is that it is much

harder to implement if one wants to add a hierarchical level to account for differences between

groups (*e.g.* sex) but with non-null covariances. Recent developments around Poisson-Log-

Normal models [47] could be used for this purpose (see e.g. [48] in the context of recombination

rate inference) but this will require further developments, in particular to better account for the

longitudinal correlation of recombination rates along the genome. Finally, we proposed a

permutation method to assess the significance of heterochiasmy along chromosomes. This

method allows to draw conclusions from a lowly informative dataset in terms of sex differences,

which could not have been confidently untangled otherwise. We believe it could be useful in

future studies on heterochiasmy.

The average recombination rate of the goat genome (1.42cM/Mb varying from 1.12 to

1.71cM/Mb among chromosomes) is very similar to other ruminants (e.g. 1.5cM/Mb in sheep

[12], 0.98cM/Mb in cattle [46]). A striking feature of its landscape is a strong heterochiasmy

that we could reveal even if the number of female meioses available in our datasets was

relatively low. This heterochiasmy is mostly concentrated in a few megabases at the

chromosome ends, irrespective of the total chromosome size. As a consequence, the difference

in average recombination rate between sexes is smaller for large chromosomes than small ones

(**Fig. 2a**). The biological causes underlying this heterochiasmy are still unknown. Further

research combining genetics studies with cytogenetics and functional studies will most likely

be required to unravel them.

Patterns of recombination and heterochiasmy in the goat match those found in sheep [12,19,48]

and cattle [8,10,11] although maybe to a lesser extent for this last species. Given that these three

species have highly colinear genomes, it seems possible to make a quantitative comparison of

recombination patterns along ruminant chromosomes to reveal how recombination evolved

with speciation.

Genetic parameters and architecture of recombination phenotypes

We studied two different phenotypes: the Genome-Wide Recombination Rate (GRR) which is

a measure of recombination intensity, and intra-chromosomal shuffling ($\overline{r}_{intra}$) which depends

on recombination intensity too but also measures differences in the localisation of crossovers.

As expected, we observed a high correlation between the two phenotypes (0.62). However,

despite strong correlation of the GWAS results on GRR and $\overline{r}_{intra}$, their genetic architectures

were found only partially overlapping, illustrating that they are probably capturing different

underlying biological processes (for example those involved in Double Strand Break (DSB)

localisation from those involved in DSB repair).

Heritability for male GRR was estimated at 0.12 with similar results for each breed, and similar

to those found in other ruminant species for this phenotype (ACC or GRR): *e.g.* 0.22 for Dutch

Holstein-Friesan males in cattle [6], 0.13 in several cattle breeds [9], 0.12 for Soay sheep [19]

and 0.23 for Lacaune sheep [12].

The heritability for male $\overline{r}_{intra}$ was estimated at 0.034 and consistent between the two breeds,

although only slightly significantly different from zero (SE=0.015). Heritability for this

phenotype seems to be smaller than in other species as [18] estimated it at 0.11 in salmon, a

species with *PRDM9* hotspots, and [22] estimated it at 0.14 in sparrows, a species without

PRDM9 hotspots.

The animal models used for estimating genetic parameters also allow to predict the additive

genetic values (EBVs) of every animal in the pedigree. These EBVs on male GRR and $\overline{r}_{intra}$

were used as phenotypes for our GWAS. This permits to have only one measure of phenotype

for each individual, and therefore to run analyses using the fast computing procedures required

for whole genome sequence imputed datasets. Using EBVs as phenotypes for GWAS is

sometimes discouraged for different reasons. First, to avoid too much shrinkage in the

phenotypes: ideally all animals should have either own and/or direct offspring phenotypes.

Here, out of the 843 individuals in the GWAS, only 323 (38%) did not have any own

phenotypes. These were all females, as they obviously cannot be directly measured for male

recombination phenotypes. Among these females, only 43 (5% of all animals) were not directly

related to a male with own information for these phenotypes, *i.e.* had a coefficient of

relationship smaller than ½ with measured samples. So overall, our dataset meets this first

requirement. Another caveat of using EBVs as phenotype is that their precisions can sometimes

vary widely between individuals, especially when there are large differences in their number of

repeated measures. In our case, the posterior variances of EBVs ranged from 2.6 to 7.1, *i.e.* a

factor smaller than 3 between the most and least precise predictions. Also, although precisions

of females EBVs tended to be smaller than males their distribution was unimodal (**See**

Additional File 1, Figures S16 & S17). Given this homogeneity in the precision of EBVs, we

did not deem deregression, a procedure that comes with its own caveats, necessary. Overall, for

this dataset we believe that phenotypic records are sufficiently informative and homogeneously

distributed among animals to use EBVs as phenotypes. Finally, we note that we used the

pedigree relationship matrix to derive EBVs while the GRM was used in the GWAS, in addition

to using the LOCO methodology, avoiding potential convergence issues (see **Methods**).

GWAS results for male GRR and $\bar{r}_{intra}$ showed low genetic correlation between Alpine and

Saanen. Despite identical heritability results, no QTL were found in common between the two

breeds. Besides, the correlation of the β for the SNPs that are significant in at least one breed

is only 0.25 for GRR and 0.04 for $\bar{r}_{intra}$. This suggests strong differences in the underlying

genetic determinants of recombination between these two breeds, although they are both

European breeds originating from the Alps and closely related in a worldwide diversity panel

[49]. Despite this relatively close genetic proximity, those breeds have been shown to have

different genetic architecture for other traits. For example, [50] found genetic heterogeneity in

the determinism of dairy-related traits between the two breeds, despite similar breeding goals.

Our results add to the literature that suggests that genome-wide recombination landscapes tend

to be mostly shared between populations while the underlying genetic determinants of the

process can be quite different, *e.g.* [12] in sheep, [15,16] in pigs, although results in cattle tend

to show they are more conserved [10].

Candidate genes associated with QTL loci in the GWAS were prioritised by implementing a

scoring methodology using criteria based on physical and genetic distance to significant SNPs.

This methodology was developed in order to provide an objective measure for the investigation

of candidate genes underlying GWAS peaks and quickly highlight strong candidates in regions

rich in annotations. For instance, 13Mb on chromosome 5 contain three different peaks in rapid

succession and 141 genes are annotated in the region. The scoring allowed to quickly identify

the most promising genes and order them all depending on their physical and genetic proximity

to each peak. Genes involved in meiosis in other species were searched for in a second step to

identify functional candidates at QTLs among the positional candidates.

The GWAS results for GRR in Alpine goats show a signal towards the end of chromosome 6,

in very close proximity to the *RNF212* gene. *RNF212* has been found to be associated with

GRR in humans [51], cattle [6,9] and sheep [12,19]. It has also been proved that *RNF212* is

involved in and is necessary for meiosis in mammals by playing a central role to the designation

of crossover sites and the formation of crossover-specific recombination complexes [52,53].

Yet it only scores second candidate for the signal on chromosome 6 because it does not directly

contain the significant SNPs. Of course, this is still consistent with a regulatory variant affecting

RNF212 expression.

One signal with significant SNPs as far as 330kb apart can be found at the end of chromosome

10 between 79.9 and 80.3Mb in the Alpine breed (**Fig. 4b**). The genes *RNF212B* and *REC8* are

in the region. They were previously suspected to be responsible for signals in sheep [12,19] and

cattle [6,9] but the studies had difficulty attributing the signal to one of the two genes due to a

lack of resolution. Among our candidates, *RNF212B* ranks 17[th] and *REC8* ranks 51[st], as the

region is very rich in annotated genes. Functionally, both these genes make very good

candidates. On the one hand, *RNF212B* is a homolog of *RNF212*, this gene's involvement in

recombination being described just above, and the two proteins *rnf212* and *rnf212b* have been

showed to interact with each other and to be necessary for their stability during the

recombination process [54]. On the other hand, REC8 is also known to be involved in meiotic

recombination [55] through DSB formation [56] as well as in implementing homologue bias

during DSB repairs [57–59]. But the gene richness of the region (62 genes in 690kb), their

relative physical distance to the lead variant and the fact that, surprisingly, the main part of the

signal is located in an area of 250kb completely devoid of annotated protein coding genes,

makes discriminating the two candidates, or even determining if they are good candidates, quite

difficult.

Among significant QTLs, the most significant signal was found in Alpine on chromosome 17

around 7Mb (**Fig. 4a**). The signal falls directly within the gene *SSH1*, which scored 5/5

according to our ranking method. This gene belongs to the slingshot phosphatase family,

constituted of three genes [60]. Investigation of the GTEx database [61,62] showed that it is

expressed in testis and automatic annotations [63,64] indicate it may be involved in chromatin

remodelling mechanisms. Furthermore, its paralog *SSH2* showed protein-enrichment in a study

aiming to predict mouse meiosis-essential genes [65] and is known to be involved in later

spermatogenesis [66]. But current knowledge of these genes' functions as well as the strong

signal found for a QTL within *SSH1* suggest this family may be involved in recombination.

When finding QTLs of interest at the end of chromosome 1 in Alpine, both for GRR and $\overline{r}_{intra}$,

one could have assumed that these signals were associated with *PRDM9*, known regulator of

recombination hotspots [67,68] and located at the very end of chromosome 1 in goats (**See**

Additional File 3, **Table 1**). Therefore, it was surprising to see that *PRDM9* actually scores

very low for the GRR signal (1/5 score) due to its physical distance from the significant QTL

and is even too far from the $\overline{r}_{intra}$ signal to gain any score point. It could be explained that no

association was found between *PRDM9* and GRR because GRR is a phenotype of intensity of

recombination, and *PRDM9* regulates the location of recombination. But $\overline{r}_{intra}$ is a

recombination phenotype linked with the position of crossovers and the association was even

more expected, especially considering that [17] found a clear signal for *PRDM9* in their own

GWAS on pig $\overline{r}_{intra}$ phenotype. Furthermore, the third highest ranking candidate gene in goats

for $\overline{r}_{intra}$ is *KCNJ15* (2/5 score and **Fig. 4e**). Although this gene seems to have no link with

recombination, an association for this gene with GRR has previously been found in sheep [12].

Furthermore, [6] found signal for the genes *KCNJ2* and *KCNJ16*, paralogs of *KCNJ15*, in their

GWAS for GRR. Thus, further investigation on the involvement of *KCNJ15* in recombination

may be relevant.

Conclusions

In this study, we have found that recombination patterns are very alike between the Alpine and

Saanen breeds, but heterochiasmy is strongly present, manifested at the chromosome ends. The

recombination pattern we observed is well conserved with closely related species like sheep

and cattle. We also showed that male GRR and male $\overline{r}_{intra}$ are heritable and that this value is

conserved in the two breeds. Heritability for GRR is concordant with results found in sheep and

cattle, but goat heritability for $\overline{r}_{intra}$ seems to be lower than in the other species in which it was

estimated. GWAS results showed low correlation between the two breeds, for both phenotypes,

and it seems that despite similar recombination patterns and same heritability for recombination

traits, the underlying genetic architecture of recombination is different between Alpine and

Saanen. Candidate genes found in GWAS are partially in common with what was previously

found in literature, with some novel genes. This supports the hypothesis that genetic control for

recombination is well conserved among many lowly-related species, but also that the genes

involved in this control can evolve quickly.

Declarations

702   **Ethics approval and consent to participate**

703   Not applicable

Consent for publication

Not applicable

Availability of data and materials

Initial genotypes and pedigree were provided by Capgenes and INRAE. Their access is restricted by the provider. Crossover data and informativity data, derived from these initial datasets, as well as genotypes of the phenotyped parents and scripts for analysis are available at https://doi.org/10.57745/HPYLLT. Sequence data for the imputation reference panel can be obtained from the VarGoats consortium: https://www.goatgenome.org/vargoats.html.

Competing interests

The authors declare that they have no competing interests.

Funding

Genotype data from the genomic selection programme have received funding from: the French Genomcap and Phenofinlait programmes (ANR, Apis-Gène, CASDAR, FranceAgriMer, France Génétique Elevage, the French Ministry of Agriculture Agrifood, and Forestry), CASDAR (Genovicap et Maxi'male). Genotype data from the INRAE research database have received funding from: UE 7th framework programme ("3SR" grant #245140), Region Centre-Val de Loire (CAPRIMAM grant) and APISGENE (ACTIVE GOAT grant). AE was funded by the INRAE Animal Genetics and Ecology and Biodiversity divisions.

Authors' contributions

All authors participated in conceptualizing this study and to the data curation and analyses. All authors developed methodology for this study and administered the project. RR provided the initial dataset as resources for the project. BS and AE did programming and software development to produce results. BS and RR supervised the project. All authors validated the

727 results. AE and BS did visualisation work through diverse presentations and wrote the original

728 draft. All authors reviewed, edited and approved the final manuscript.

Acknowledgements

Many numerical analyses were performed on the genobioinfo bioinformatics platform Toulouse

Midi-Pyrénées (Bioinfo Genotoul). We are thankful to Capgenes and the INRAE P3R unit

(https://doi.org/10.15454/1.5483259352597417E12) for providing access to SNP array

genotypes, and to the VarGoats consortium for early access to whole genome sequencing data.

We would like to thank Susan Johnston, Tristan Mary-Huard and Leopoldo Sanchez Rodriguez

for fruitful discussions during the production of these results.

References

1. Koehler KE, Hawley RS, Sherman S, Hassold T. Recombination and nondisjunction in
humans and flies. Human Molecular Genetics. 1996;5:1495–504.

2. Muller HJ. THE RELATION OF RECOMBINATION TO MUTATIONAL ADVANCE.
Mutat Res. 1964;106:2–9.

3. Crow JF, Kimura M. Evolution in Sexual and Asexual Populations. The American Naturalist.
1965;99:439–50.

4. Kong A, Gudbjartsson DF, Sainz J, Jonsdottir GM, Gudjonsson SA, Richardsson B, et al. A
high-resolution recombination map of the human genome. Nat Genet. 2002;31:241–7.

5. Coop G, Wen X, Ober C, Pritchard JK, Przeworski M. High-Resolution Mapping of
Crossovers Reveals Extensive Variation in Fine-Scale Recombination Patterns Among
Humans. Science. 2008;319:1395–8.

6. Sandor C, Li W, Coppieters W, Druet T, Charlier C, Georges M. Genetic Variants in REC8,
RNF212, and PRDM9 Influence Male Recombination in Cattle. Paigen K, editor. PLoS Genet.
2012;8:e1002854.

7. Weng Z-Q, Saatchi M, Schnabel RD, Taylor JF, Garrick DJ. Recombination locations and
rates in beef cattle assessed from parent-offspring pairs. Genet Sel Evol. 2014;46:34.

8. Ma L, O'Connell JR, VanRaden PM, Shen B, Padhi A, Sun C, et al. Cattle Sex-Specific
Recombination and Genetic Control from a Large Pedigree Analysis. Auton AJ, editor. PLoS
Genet. 2015;11:e1005387.

9. Kadri NK, Harland C, Faux P, Cambisano N, Karim L, Coppieters W, et al. Coding and
noncoding variants in *HFM1* , *MLH3* , *MSH4* , *MSH5* , *RNF212* , and *RNF212B* affect
recombination rate in cattle. Genome Res. 2016;26:1323–32.

10. Shen B, Jiang J, Seroussi E, Liu GE, Ma L. Characterization of recombination features and
the genetic basis in multiple cattle breeds. BMC Genomics. 2018;19:304.

11. Brekke C, Johnston SE, Gjuvsland AB, Berg P. Variation and genetic control of individual
recombination rates in Norwegian Red dairy cattle. Journal of Dairy Science. 2023;106:1130–
41.

12. Petit M, Astruc J-M, Sarry J, Drouilhet L, Fabre S, Moreno CR, et al. Variation in
Recombination Rate and Its Genetic Determinism in Sheep Populations. Genetics.
2017;207:767–84.

13. Weng Z, Wolc A, Su H, Fernando RL, Dekkers JCM, Arango J, et al. Identification of
recombination hotspots and quantitative trait loci for recombination rate in layer chickens. J
Animal Sci Biotechnol. 2019;10:1–13.

14. Johnsson M, Whalen A, Ros-Freixedes R, Gorjanc G, Chen C-Y, Herring WO, et al. Genetic
variation in recombination rate in the pig. Genet Sel Evol. 2021;53:54.

15. Lozada-Soto EA, Maltecca C, Wackel H, Flowers W, Gray K, He Y, et al. Evidence for
recombination variability in purebred swine populations. J Anim Breed Genet. 2021;138:259–
73.

16. Brekke C, Berg P, Gjuvsland AB, Johnston SE. Recombination rates in pigs differ between
breeds, sexes and individuals, and are associated with the RNF212, SYCP2, PRDM7, MEI1
and MSH4 loci. Genet Sel Evol. 2022;54:33.

17. Brekke C, Gjuvsland AB, Berg P, Johnston SE. Independent genetic basis of meiotic
crossover positioning and interference in domestic pigs. Sci Rep. 2025;15:9260.

18. Brekke C, Johnston SE, Knutsen TM, Berg P. Genetic architecture of individual meiotic
crossover rate and distribution in Atlantic Salmon. Sci Rep. 2023;13:20481.

19. Johnston SE, Bérénos C, Slate J, Pemberton JM. Conserved Genetic Architecture
Underlying Individual Recombination Rate Variation in a Wild Population of Soay Sheep (
Ovis aries). Genetics. 2016;203:583–98.

20. Johnston SE, Huisman J, Ellis PA, Pemberton JM. A High-Density Linkage Map Reveals
Sexual Dimorphism in Recombination Landscapes in Red Deer (Cervus elaphus). G3
(Bethesda). 2017;7:12.

21. Johnston SE, Huisman J, Pemberton JM. A Genomic Region Containing REC8 and
RNF212B Is Associated with Individual Recombination Rate Variation in a Wild Population
of Red Deer. G3 (Bethesda). 2018;8:12.

22. McAuley JB, Servin B, Burnett HA, Brekke C, Peters L, Hagen IJ, et al. The Genetic
Architecture of Recombination Rates is Polygenic and Differs Between the Sexes in Wild
House Sparrows (Passer domesticus). Molecular Biology and Evolution. 2024;41:msae179.

23. Cooney CR, Mank JE, Wright AE. Constraint and divergence in the evolution of male and
female recombination rates in fishes. Evolution. 2021;75:2857–66.

24. Veller C, Kleckner N, Nowak MA. A rigorous measure of genome-wide genetic shuffling
that takes into account crossover positions and Mendel's second law. Proceedings of the
National Academy of Sciences. 2019;116:1659–68.

25. Schibler L, Vaiman D, Oustry A, Giraud-Delville C, Cribiu EP. Comparative Gene
Mapping: A Fine-Scale Survey of Chromosome Rearrangements between Ruminants and
Humans. Genome Res. 1998;8:901–15.

26. Fröhlich J, Vozdova M, Kubickova S, Cernohorska H, Sebestova H, Rubes J. Variation of
Meiotic Recombination Rates and MLH1 Foci Distribution in Spermatocytes of Cattle, Sheep
and Goats. Cytogenet Genome Res. 2015;146:211–21.

27. Muñoz-Fuentes V, Marcet-Ortega M, Alkorta-Aranburu G, Linde Forsberg C, Morrell JM,
Manzano-Piedras E, et al. Strong Artificial Selection in Domestic Mammals Did Not Result in
an Increased Recombination Rate. Mol Biol Evol. 2015;32:510–23.

28. Dong Y, Xie M, Jiang Y, Xiao N, Du X, Zhang W, et al. Sequencing and automated whole-
genome optical mapping of the genome of a domestic goat (Capra hircus). Nat Biotechnol.
2013;31:135–41.

29. Tosser-Klopp G, Bardou P, Bouchez O, Cabau C, Crooijmans R, Dong Y, et al. Design and
Characterization of a 52K SNP Chip for Goats. Liu Z, editor. PLoS ONE. 2014;9:e86227.

30. Denoyelle L, Talouarn E, Bardou P, Colli L, Alberti A, Danchin C, et al. VarGoats project:
a dataset of 1159 whole-genome sequences to dissect Capra hircus global diversity. Genet Sel
Evol. 2021;53:86.

31. Du X, Servin B, Womack JE, Cao J, Yu M, Dong Y, et al. An update of the goat genome
assembly using dense radiation hybrid maps allows detailed analysis of evolutionary
rearrangements in Bovidae. BMC Genomics. 2014;15:625.

32. Schensted, C. Longest increasing and decreasing subsequences. Can J Math. 1961;13:179–
91.

33. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. GigaSci. 2015;4:7.

34. Servin B. Yapp: a suite of tools to analyze genomic data in pedigrees [Internet]. INRAE;
2024. Available from: https://yapp.readthedocs.io

35. Favier A, Elsen J-M, de Givry S, Legarra A. Optimal haplotype reconstruction in half-sib
families. In: Dovier A, Dal Palù A, Will S, editors. Edinburgh, UK: EasyChair; 2010 [cited
2022 May 5]. p. 27–37. Available from: https://easychair.org/publications/paper/SpJ

36. Druet T, Georges M. LINKPHASE3: an improved pedigree-based phasing algorithm robust
to genotyping and map errors. Bioinformatics. 2015;31:1677–9.

37. Storey JD, Tibshirani R. Statistical significance for genomewide studies. Proc Natl Acad
Sci USA. 2003;100:9440–5.

38. Meyer K. WOMBAT—A tool for mixed model analyses in quantitative genetics by
restricted maximum likelihood (REML). J Zhejiang Univ - Sci B. 2007;8:815–21.

39. Van der Auwera G, O'Connor BD. Genomics in the Cloud: Using Docker, GATK, and
WDL in Terra (1st edition). O'Reilly Media; 2020.

40. Browning BL, Browning SR. Genotype Imputation with Millions of Reference Samples.
The American Journal of Human Genetics. 2016;98:116–26.

41. Pook T, Mayer M, Geibel J, Weigend S, Cavero D, Schoen CC, et al. Improving Imputation
Quality in BEAGLE for Crop and Livestock Data. G3 Genes|Genomes|Genetics. 2020;10:177–
88.

42. Delaneau O, Zagury J-F, Robinson MR, Marchini JL, Dermitzakis ET. Accurate, scalable
and integrative haplotype estimation. Nat Commun. 2019;10:5436.

43. Rubinacci S, Delaneau O, Marchini J. Genotype imputation using the Positional Burrows
Wheeler Transform. Scheet P, editor. PLoS Genet. 2020;16:e1009049.

44. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association studies.
Nat Genet. 2012;44:821–4.

45. Stephens M. False discovery rates: a new deal. Biostat. 2016;kxw041.

46. Qanbari S, Wittenburg D. Male recombination map of the autosomal genome in German
Holstein. Genet Sel Evol. 2020;52:73.

47. Chiquet J, Mariadassou M, Robin S. The Poisson-Lognormal Model as a Versatile
Framework for the Joint Analysis of Species Abundances. Front Ecol Evol. 2021;9:588292.

48. Vassilief H, Id Bella M, Hazard D, Tortereau F, Faraut T, Johnston SE, et al. 202. Sex
differences in recombination maps are associated with differential hotspot usage in sheep.
Proceedings of 12th World Congress on Genetics Applied to Livestock Production (WCGALP)
[Internet]. Rotterdam, the Netherlands: Wageningen Academic Publishers; 2022 [cited 2025
Feb 12]. p. 861–4. Available from: https://www.wageningenacademic.com/doi/10.3920/978-
90-8686-940-4_202

49. Colli L, Milanesi M, Talenti A, Bertolini F, Chen M, Crisà A, et al. Genome-wide SNP
profiling of worldwide goat populations reveals strong partitioning of diversity and highlights
post-domestication migration routes. Genet Sel Evol. 2018;50:58.

50. Martin P, Palhière I, Maroteau C, Clément V, David I, Klopp GT, et al. Genome-wide
association mapping for type and mammary health traits in French dairy goats identifies a
pleiotropic region on chromosome 19 in the Saanen breed. Journal of Dairy Science.
2018;101:5214–26.

51. Kong A, Thorleifsson G, Frigge ML, Masson G, Gudbjartsson DF, Villemoes R, et al.
Common and low-frequency variants associated with genome-wide recombination rate. Nat
Genet. 2014;46:11–6.

52. Reynolds A, Qiao H, Yang Y, Chen JK, Jackson N, Biswas K, et al. RNF212 is a dosage-
sensitive regulator of crossing-over during mammalian meiosis. Nat Genet. 2013;45:269–78.

53. Qiao H, Prasada Rao HBD, Yang Y, Fong JH, Cloutier JM, Deacon DC, et al. Antagonistic
roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. Nat
Genet. 2014;46:194–9.

54. Ito M, Yun Y, Kulkarni DS, Lee S, Sandhu S, Nuñez B, et al. Distinct and interdependent
functions of three RING proteins regulate recombination during mammalian meiosis.
Proceedings of the National Academy of Sciences. 2025;122:e2412961121.

55. Parisi S, McKay MJ, Molnar M, Thompson MA, van der Spek PJ, van Drunen-
Schoenmaker E, et al. Rec8p, a Meiotic Recombination and Sister Chromatid Cohesion
Phosphoprotein of the Rad21p Family Conserved from Fission Yeast to Humans. Mol Cell
Biol. 1999;19:3515–28.

56. Kugou K, Fukuda T, Yamada S, Ito M, Sasanuma H, Mori S, et al. Rec8 guides canonical
Spo11 distribution along yeast meiotic chromosomes. Mol Biol Cell. 2009;20:3064–76.

57. Klein F, Mahr P, Galova M, Buonomo SB, Michaelis C, Nairz K, et al. A central role for
cohesins in sister chromatid cohesion, formation of axial elements, and recombination during
yeast meiosis. Cell. 1999;98:91–103.

58. Brar GA, Hochwagen A, Ee LS, Amon A. The Multiple Roles of Cohesin in Meiotic
Chromosome Morphogenesis and Pairing. MBoC. 2009;20:1030–47.

59. Kim KP, Weiner BM, Zhang L, Jordan A, Dekker J, Kleckner N. Sister cohesion and
structural axis components mediate homolog bias of meiotic recombination. Cell.
2010;143:924–37.

60. Patterson KI, Brummer T, O'brien PM, Daly RJ. Dual-specificity phosphatases: critical
regulators with diverse cellular targets. Biochemical Journal. 2009;418:475–89.

61. Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, et al. The Genotype-Tissue
Expression (GTEx) project. Nat Genet. 2013;45:580–5.

62. GTEx Portal - SSH1 gene page [Internet]. GTEx Portal. [cited 2025 Aug 14]. Available
from: https://www.gtexportal.org/home/gene/SSH1

63. SSH1 Gene - GeneCards | SSH1 Protein | SSH1 Antibody [Internet]. GeneCards. [cited
2025   Aug   14].   Available   from: https://www.genecards.org/cgi-
bin/carddisp.pl?gene=SSH1&keywords=SSH1

64. Seal RL, Braschi B, Gray K, Jones TEM, Tweedie S, Haim-Vilmovsky L, et al.
Genenames.org: the HGNC resources in 2023. Nucleic Acids Res. 2023;51:D1003–9.

65. Fang K, Li Q, Wei Y, Zhou C, Guo W, Shen J, et al. Prediction and Validation of Mouse
Meiosis-Essential Genes Based on Spermatogenesis Proteome Dynamics. Molecular & Cellular
Proteomics. 2021;20:100014.

66. Xu K, Su X, Fang K, Lv Y, Huang T, Li M, et al. The Slingshot phosphatase 2 is required
for acrosome biogenesis during spermatogenesis in mice. Yan W, Harper DM, Hammoud S,
editors. eLife. 2023;12:e83129.

67. Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M, et al. PRDM9 Is a Major
Determinant of Meiotic Recombination Hotspots in Humans and Mice. Science. 2010;327:836–
40.

68. Baker Z, Schumer M, Haba Y, Bashkirova L, Holland C, Rosenthal GG, et al. Repeated
losses of PRDM9-directed recombination despite the conservation of PRDM9 across
vertebrates. de Massy B, editor. eLife. 2017;6:e24133.

Figures

Figure 1: Illustration of the notion of interval informativity.

Figure 2: Estimated recombination rates for males and females at several scales.

(a): Chromosomic recombination rate depending on chromosome size. (b): Bayesian estimates
of the chromosome landscape of recombination rate depending on distance to centromere. All
29 chromosomes are acrocentric. Confidence intervals at 99%. (c): Recombination map on
chromosome 25 at 1Mb resolution. Confidence interval under H_0 estimated using sex
permutations.

Figure 3: Manhattan plots of genome-wide association studies (GWAS) for male recombination
phenotypes after imputation to sequence level.

GRR and $\overline{r}_{intra}$ stand respectively for EBVs on male GRR and EBVs on male $\overline{r}_{intra}$.

Figure 4: Zoom on GWAS Manhattan plots for regions of interest in both populations and
phenotypes.

Additional Files

Additional File 1 Figure S1

File format: PDF

Title: Position of (b)ovine (top) and goat (bottom) chip SNPs on both the RH map and the ARS1 assembly, on chromosomes 2 and 18.

Description: Goat SNPs have been filtered using the longest increasing subsequence algorithm, and only remaining SNPs have been represented.

Additional File 1 Figure S2

File format: PDF

Title: Crossover distribution on the 35-45 Mb area of chromosome 1.

Description: Red lines = position of conserved synteny breaks.

Additional File 1 Figure S3

File format: PDF

Title: Additive SNP and sample call-rate cumulative distributions.

Description: Left: SNP call-rate; right: sample call-rate. Black line: chosen threshold=0.01.

Additional File 1 Figure S4

File format: PDF

Title: Distribution of the distance between two crossovers per chromosome per meiosis.
Description: Red line: chosen threshold=5Mb

Additional File 1 Figure S5

File format: PDF

Title: Density distribution of the Maximum Likelihood estimates of recombination rates on 1Mb intervals of the genome.

Description: Black: gamma law; extracted parameters α=3.08 & β=2.20.

Additional File 1 Figure S6

File format: PDF

Title: Distribution of crossover detection interval sizes.

Additional File 1 Figure S7

File format: PDF

Title: Distribution of all permuted recombination rate log ratio for all genomic intervals.

Additional File 1 Figure S8

File format: PDF

Title: SNP distribution of sequencing and variant calling quality parameters.

959 Description: A: DP parameter; B: FS parameter; C: VQSLOD parameter.

Additional File 1 Figure S9

961 File format: PDF

962 Title: R² evolution according to allele frequency.

963 Description: (A) Before genotype imputation. Mean R²=0.928. (B) After genotype imputation.
964 Mean R²=0.990.

Additional File 1 Figure S10

966 File format: PDF

967 Title: Genotype concordance according to allele frequency.
968 Description: (A) Before genotype imputation. Mean genotype concordance=0.974. (B) After
969 genotype imputation. Mean genotype concordance=0.997.

Additional File 1 Figure S11

971 File format: PDF

972 Title: Distribution the proportion of the autosomal genome informative for crossovers in parents
973 (M=males vs F=females).

Additional File 1 Figure S12

975 File format: PDF

976 Title: Comparison of recombination map size estimates between populations.

977 Description: Confidence interval $= \mu \pm 2 \times standard\ error$.

Additional File 1 Figure S13

979 File format: PDF

980 Title: Distribution of genome-wide recombination rates (GRR) in meioses.

Additional File 1 Figure S14

982 File format: PDF

983 Title: Distribution of intra-chromosomal shuffling in meioses.

Additional File 1 Figure S15

985 File format: PDF

986 Title: Comparison of intra-chromosomal shuffling estimates between populations.

987 Description: Confidence interval $= \mu \pm 2 \times standard\ error$.

Additional File 1 Figure S16

Additional File 1 Figure S17

Additional File 2 Table S1

File format: XLS

Title: SNPs within synteny breakpoints

Description: File listing all 123 SNPs removed from dataset after identifying synteny
breakpoints with human genome on ARS1 assembly.

Additional File 3 Table S2

File format: XLS

Title: BLAST results for the bovine gene sequence RNF212 on ARS1 assembly.

Additional File 3 Table S3

File format: XLS

Title: BLAST results for the bovine gene sequence PRDM9 on ARS1 assembly.

Additional File 4 Table S4

File format: XLS

Title: Sex-specific estimated recombination rates for each 1Mb interval of the autosomal goat
genome.

Additional File 5 Figure S18

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 1, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S19

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 2, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S20

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 3, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S21

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 4, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S22

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 5, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S23

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 6, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S24

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 7, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S25

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 8, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom
panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found
significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S26

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 9, at 1Mb
resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S27

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 10, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S28

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 11, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S29

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 12, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S30

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 13, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S31

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 14, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S32

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 15, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S33

Title: Estimated recombination rates for males and females on chromosome 16, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S34

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 17, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S35

Title: Estimated recombination rates for males and females on chromosome 18, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S36

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 19, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S37

Title: Estimated recombination rates for males and females on chromosome 20, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S38

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 21, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S39

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 22, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S40

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 23, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S41

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 24, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S42

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 25, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S43

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 26, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S44

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 27, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S45

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 28, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 5 Figure S46

File format: PDF

Title: Estimated recombination rates for males and females on chromosome 29, at 1Mb resolution.

Description: Log ratio between male and female recombination rates were plotted in the bottom panel. Confidence interval of log ratio under null hypothesis appears in grey. Intervals found significant for sex differences are marked with a red dot on the bottom panel.

Additional File 6 Table S5

File format: XLS

Title: Significant 1Mb intervals for sex differences in recombination rate.

Additional File 7 Table S6

File format: PDF

Title: Candidate genes for GWAS on EBVs for male genome-wide recombination rate in Alpine goats.

Description: All genes scoring at least 1 as candidates are showed. For each QTL of interest, genes have been sorted from higher scoring to lower scoring. Manual annotation has been added

on genes previously found in GWAS for recombination phenotypes in ruminants, referencing the corresponding papers.

Additional File 7 Table S7

File format: PDF

Title: Candidate genes for GWAS on EBVs for male genome-wide recombination rate in Saanen goats.

Description: All genes scoring at least 1 as candidates are showed. For each QTL of interest, genes have been sorted from higher scoring to lower scoring. Manual annotation has been added on genes previously found in GWAS for recombination phenotypes in ruminants, referencing the corresponding papers.

Additional File 7 Table S8

File format: PDF

Title: Candidate genes for GWAS on EBVs for male intra-chromosomal shuffling in Alpine goats.

Description: All genes scoring at least 1 as candidates are showed. For each QTL of interest, genes have been sorted from higher scoring to lower scoring. Manual annotation has been added on genes previously found in GWAS for recombination phenotypes in ruminants, referencing the corresponding papers.

Additional File 7 Table S9

File format: PDF

Title: Candidate genes for GWAS on EBVs for male intra-chromosomal shuffling in Saanen goats.

Description: All genes scoring at least 1 as candidates are showed. For each QTL of interest, genes have been sorted from higher scoring to lower scoring. Manual annotation has been added on genes previously found in GWAS for recombination phenotypes in ruminants, referencing the corresponding papers.

informative interval = I_{mi}

interval on which crossovers can be detected

chromosome sequence

1Mb interval