DNA methylation networks during pig fetal development: a joint fused ridge estimation approach

Karolina M. Wachala[1¤,2*], Jani de Vos[3¤a,4], Ole Madsen[4], Martijn F.L. Derks[4†], Carel F.W. Peeters[2†]

1 Interfaculty Bioinformatics Unit, University of Bern, Bern, Switzerland

2 Mathematical and Statistical Methods group (Biometris), Wageningen University & Research, Wageningen, Netherlands

3 Research and Technology Centre, Hendrix Genetics, Boxmeer, Netherlands

4 Animal Breeding and Genomics group, Wageningen University & Research, Wageningen, Netherlands

¤Current address Karolina M. Wachala

¤a Current address Jani de Vos

†These authors contributed equally to this work.

* Corresponding author: karolina.wachala@unibe.ch

Abstract

Although an organism's genetic information is predominantly identical among most of its cell types, the epigenome regulates the expression of the genome in a cell type- and context-dependent manner. In mammals, DNA methylation in regulatory regions, such as promoters, primarily regulates gene expression by inducing transcriptional inactivation. With genome-wide approaches came the realization that DNA methylation patterns underlying mammalian development are considerably more dynamic than previously recognized. This realization highlights the need for methodological approaches capable of capturing this phenomenon. In this study, we investigated the feasibility of modeling DNA methylation networks by jointly estimating regularized precision matrices from time- and tissue-specific omics data derived from the pig genome. For that, we utilized RNA-Seq and RRBS data that span seven pig tissues at

three developmental stages: early organogenesis, late organogenesis, and newborn. Our analysis focused on 61, 48, and 74 genes — differentially expressed across developmental stages and CpG-methylated in promoter regions, from endoderm-, mesoderm-, and ectoderm-derived tissues, respectively. Using a joint fused ridge approach, we were able to *borrow* information across tissues and time points, enabling more robust network inference. This analytical framework advances exploratory methods for studying organism development using pig as a model species. Our results highlight the importance of fetal-maternal immunity and the circulatory system in early development, and shed light on dynamic interactions across tissues, organ systems, and germ layers. We anticipate that this flexible framework can be extended to other omics data and species, facilitating future research.

Author summary

This study explores how gene activity is controlled during pig fetal development using networks. The epigenome can turn genes on and off, depending on, e.g. the cell's function and the organism's growth stage. This process includes chemical changes to DNA, such as methylation. We focused on DNA methylation patterns during pig fetal development by analyzing samples from several organs at three stages of fetal growth. We combined gene activity data with DNA methylation data by using a network-based method that allows us to visualize and study how genes interact across different tissues and developmental stages. We discovered that the immune and circulatory systems are critical during early development. We have also observed complex interactions across tissues and organ systems. Our study provides a flexible toolbox with clear, step-by-step explanations. This analytical framework can be used to explore developmental patterns at the genomic scale in other species and can be adapted to different types of biological data.

Introduction

Significance of swine in biomedical research and livestock systems

The pig is a crucial large animal model species in various fields, such as xenotransplantation and surgical training [1]. It is used to study disease conditions owing to its anatomical and physiological similarities to humans [2]. With swine genomics recognized for its importance in biomedical research, efforts to develop complete swine (epi)genome maps have accelerated [3, 4]. Furthermore, since porcine genome sequences became crucial for discovering molecular genetic variants, they have enabled research into the genetic control of quantitative traits such as growth, reproduction, and robustness, for example, resilience to stressors, which are economically and societally desirable in livestock systems [5, 6]. The interest in genetic control of desirable traits is not surprising as pigs hold significance as a crucial global source of meat production [7]. Genomic selection, which utilizes genomic information to predict genetic potential of individual animals for desirable traits, is widely used in animal breeding with the aim of genetic improvements of livestock [8, 9]. However, as DNA methylation, among other functional epigenetic data, is pivotal to understand the regulatory mechanisms guiding pig development to e.g., predict genomic potential, efforts to characterize epigenetic mechanisms underlying porcine development are accelerating [10].

Exploring DNA methylation and its role in swine development

Epigenetics is a bridge between an organism's genotype and phenotype. Epigenetic modifications influence the phenotype through chemical changes altering gene expression without changing the base-pairing of the DNA sequence itself [11]. Mammals, with pigs among them, exhibit three main processes of epigenetic regulation: (i) DNA methylation, (ii) Histone modifications, and (iii) Coding and non-coding RNAs [12]. Epigenomics relates to the analysis of these processes across numerous genes in a cell or an organism [12]. DNA methylation is a crucial epigenetic modification of gene regulation essential for mammalian development, and has been the most extensively investigated process of epigenetic regulation since its discovery in 1948 [13]. DNA

methylation controls the transcriptional activity of genes by altering the chromosomal   30
structure, DNA conformation, and DNA stability [14]. It occurs via the covalent   31
transfer of the methyl group to the C-5 position of the cytosine ring of eukaryotic DNA,   32
regulated by a family of DNA methyltransferases (DNMTs), namely DNMT3a and   33
DNMT3b [15, 16]. The DNMT1 enzyme ensures the maintenance of DNA methylation.   34
Upon establishing DNA methylation patterns, DNMT enzymes aid in continuation of   35
said patterns into subsequent cellular generations. Although DNA methylation is   36
usually erased during zygote formation, it is subsequently re-established in the embryo   37
upon implementation [15]. The formation of 5-methylcytosine resulting from DNA   38
methylation is essential for normal embryonic development in mammals, as it is vital for   39
processes such as genomic imprinting, maintenance of X chromosome inactivation,   40
chromosomal maintenance and genomic stability [12, 13, 15].   41

In the majority of mammalian somatic cells, the covalent transfer of methyl group   42
occurs predominantly at the 5'—C—phosphate—G—3' (CpG) dinucleotides, with most   43
CpG sites in the genome fully methylated [17]. Additionally, gene promoters contain   44
CpG islands, which are CpG-rich regions primarily regulating gene transcription [18].   45
Gene promoters are short segments of DNA (100–1,000 base pairs) where gene   46
transcription is initiated, and methylation was revealed to regulate gene expression by   47
switching-off transcription of the corresponding gene [19, 20]. However, the genome's   48
promoters remain generally unmethylated [21]. Nevertheless, owing to advancements in   49
the genome-scale mapping of methylation, DNA methylation can be studied in various   50
genomic contexts, from transcriptional start sites (TSSs) through gene bodies to   51
regulatory elements, gene promoters and enhancers [19, 22]. Although traditionally   52
DNA methylation is primarily associated with CpG islands, demethylation seems to   53
occur predominantly in DNA regions that are not CpG islands, which has led to the   54
recognition of DNA methylation as a vastly dynamic process [22]. The dynamic nature   55
of DNA methylation highlights the complexity of this epigenetic regulatory process, its   56
potential impact on gene expression, and, consequently, tissue growth and   57
morphogenesis during organism development.   58

Dynamic epigenomic switches in swine: Integrating network approaches

The GENE-SWitCH project, working in cooperation with the Functional Annotation of Animal Genomes (FAANG) Consortium, aimed to uncover the dynamic epigenetic switches in the functional genome of swine [23]. Since gene expression, transcripts and regulatory regions differ in a time- and tissue-specific manner, a static view of the genome and current genome annotation data sets may restrain a deeper understanding of gene expression and its function in mammalian development [24]. Consequently, a statistical approach allowing for the analysis of these complex and dynamic data sets is needed. However, epigenomic data are high-dimensional, as the variable dimension (genes) exceeds the observation dimension (biological replicates) [25]. The high dimensionality of the dataset leads to a so-called *curse of dimensionality* [26]. As the number of dimensions in a dataset increases, the mathematical problem space increases exponentially [27]. Therefore, the curse of dimensionality increases the computational and mathematical complexity of elucidating the epigenomics mechanisms that govern organism development [26]. In recent years, researchers developed numerous computational methods for conquering the curse of dimensionality [26]. One such method is ridge shrinkage [28]. Ridge shrinkage promotes more stable and reliable estimates of model parameters [29]. Therefore, it is applicable in omics data analyses. Next to ridge regularization, networks have gained popularity as a visually appealing and interpretable approach for the analysis of molecular mechanisms measured in thousands of variables, e.g., genes [30].

However, combining data from multiple datasets, e.g., pig tissues at developmental stages in a network, might dilute the unique biology of each sample, leading to an oversimplified representation of the dynamic epigenomics landscape during organism development. There is a need for a visually appealing and interpretable modeling approach, not only to extract DNA methylation networks and visualize how they change during organism development, but to separate signal from noise efficiently in high-dimensional DNA methylation datasets. Therefore, we aim to apply a network modeling approach utilizing data fusion over time and tissue types. Fusion will allow us to *borrow* strength from variables across DNA methylation datasets, helping to improve

the stability and accuracy of generated networks, by promoting the detection of significant edges with higher precision [31]. To decide where fusion penalties should be applied, we will utilize results from hierarchical clustering of RNA sequencing (RNA-Seq) data. Moreover, using ridge shrinkage and fusion, we will extract the parameters describing network topology jointly for all tissues originating from the same somatic germ layer. Furthermore, we will visualize differential networks, capturing how DNA methylation dynamics change between developmental stages for each tissue. Thereafter, we will analyze the extracted networks. With all these steps combined our networks will more accurately capture the intricate biological context of the GENE-SWitCH Reduced Representation Bisulphite Sequencing (RRBS) datasets, allowing us to gain insights into DNA methylation dynamics driving pig development.

Materials and methods

Initial datasets

All the data used in this study were collected prior to our research by the GENE-SWitCH consortium funded by the European Union's Horizon 2020 Research and Innovation Programme. The RRBS and RNA-Seq data we used are publicly available through the FAANG data portal. RRBS uses methylation-sensitive restriction enzyme digestion on the genomic DNA [32]. The fragmented genomic DNA is then treated with bisulphite and sequenced [32]. After treatment of fragmented genomic DNA with bisulphite, unmethylated cytosine residues are converted to uracil and subsequently to thymine during polymerase chain reaction (PCR) amplification, while 5-methylcytosine residues remain unaffected [33,34]. GENE-SWitCH has collected RRBS and RNA-Seq data from seven porcine tissues representative of the three somatic germ layers:

   I. Endoderm – liver, lung, small intestine;

  II. Mesoderm – kidney, skeletal muscle from hindlimb;

 III. Ectoderm – hindbrain, skin.

Moreover, the data stem from three developmental stages: 30 days post-fertilization (dpf) or early organogenesis, 70dpf or late organogenesis, and from newborn piglets

(NB). Additionally, at 30dpf, pooling of samples (biological replicates) coming from multiple fetuses was performed to ensure enough genetic material will be available for sequencing. RNA-Seq data collection was followed up by the normalization of read counts and differential expression analysis. The RRBS data, RNA-Seq data, and the differential expression analysis results provided by the GENE-SWitCH were the starting point for our research.

Data preparation

The GENE-SWitCH project has utilized RRBS to study genome-wide DNA methylation of swine at a single-nucleotide resolution. Since RRBS enriches for CpG-dense regions while using restriction enzymes cutting at specific sites for fragmentation, it does not cover non-CpG areas, and CpGs in areas without the enzyme restriction site [32, 35]. Therefore, we focused on methylation in the CpG-rich promoter regions to ensure proper coverage. However, the annotation of promoter regions in CpG-methylated genes is not trivial, as for meaningful inference the most precise promoter region specification is desirable. The results from chromatin immunoprecipitation followed by sequencing (ChIP-Seq) served us to that end. This technology uses antibodies specific to a DNA-binding protein of interest to identify regulatory regions of the genome [36]. In the case of promoter regions, the H3 lysine 4 trimethylation (H3K4me3) is a core histone mark used to identify active promoter regions [36]. By investigating the distribution of histone modification signals around TSSs we identified the number of base pairs (bp) upstream and downstream of TSSs where promoters are most likely to lie. We specified that the read distribution peak is around 200bp upstream, and 50bp downstream of gene to the TSS. Moreover, we extracted methylation values only for the significantly differentially expressed genes (DEGs) between developmental stages, as obtained by GENE-SWitCH [24]. We wanted to retain all DEGs between developmental stages; therefore we did not filter on the fold change. Moreover, we excluded genes located on the X chromosome from the analyses, to mitigate potential biases related to sex differences in gene expression and methylation. We extracted the methylation values separately for each tissue at developmental stage, resulting in 21 files representing 7 pig tissues at 3 developmental stages.

Prior to exploratory analysis, we processed the data further. First, since CpG [148] methylation might occur at multiple positions in the promoter region of a gene, while [149] numerous genes have alternative promoters aiding pre-mRNA splicing [37], we recorded [150] the median of replicate methylation values per gene; other data, including genes for [151] which all the methylation values were 0, was omitted. Excluding genes that exhibited [152] zero methylation values in all replicates ensured the presence of between-sample [153] variation. We thereafter transposed the resulting data frames, with replicates arranged [154] in rows, and genes arranged in columns. Secondly, we normalized (mean centered and [155] divided by the standard deviation) the columns of the numeric matrices obtained to [156] normalize the range of methylation values, which is a crucial step when implementing [157] algorithms that calculate distances between data. [158]

Gaussian Graphical Models [159]

We aimed to visualize networks representing relationships among variables, where each [160] variable corresponds to the methylation value of a CpG methylated at a promoter DEG [161] between developmental stages. Network science evolved from graph theory, a [162] mathematical branch exploring relationships between pairs of objects [38]. Therefore, a [163] network is just a collection of elements called nodes, linked together by their interactions, [164] known as edges [39]. As the presence of an edge between nodes in the network indicates [165] a conditional dependency between the variables, we aimed to extract, visualize and [166] analyze Gaussian Graphical Models (GGMs) using the `rags2ridges` R package [40, 41]. [167] GGM is a type of graphical model that captures pairwise relationships among variables, [168] based on the assumption that the joint distribution of these variables follows a [169] multivariate normal distribution [40]. When we assume multi-normality, the nodes in [170] the network represent variables that follow a multivariate normal distribution. In this [171] context, conditional independence between nodes is equivalent to the corresponding [172] partial correlation zero, indicating no relationship, and therefore the absence of an edge. [173]

Probabilistic graphs, which are networks, can depict such conditional dependencies. [174] These graphs consist of interconnected elements, known as nodes, representing variables, [175] while the connections between nodes, termed edges, illustrate their interactions [38]. In [176] the case of conditional (in)dependence, nodes represent variables, whereas edges [177]

represent their conditional dependencies. If X ⊥ Y | Z Z holds true, the edge between X   178
and Y variable will be excluded from the graph, whereas conditional dependence is   179
illustrated by the presence of an edge, as illustrated in Fig 1.   180

Fig 1. Undirected conditional independence graph. X and Y variables are conditionally independent conditioned on Z, meaning that once Z is given, knowing the value of X is superfluous concerning the value of Y, and vice versa. Since X ⊥ Y | Z holds true, the edge between X and Y variable is excluded from the graph

Consequently, partial correlation can measure conditional (in)dependence [42], which   181
can be formulated as   182

$$X \perp Y \mid Z \leftrightarrow p_{XY \cdot Z} = 0 \tag{1}$$

for conditional independence and   183

$$X \not\perp Y \mid Z \leftrightarrow p_{XY \cdot Z} \neq 0 \tag{2}$$

in the case of conditional dependence, where $p_{XY \cdot Z}$ is a partial correlation coefficient   184
denoting the relationship between two random variables, X and Y, conditioned on a set   185
of random variables Z which possibly explain this association [40]. The parameters of   186
GGMs, i.e., the partial correlations, can be estimated by inverting the sample covariance   187
matrix Σ to obtain the precision matrix Ω. Let us consider a covariance matrix   188

$$\Sigma = \begin{array}{c} \\ x \\ y \\ z \end{array}\begin{array}{ccc} x & y & z \\ \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix} \end{array}, \tag{3}$$

with the inverse covariance matrix (precision matrix) given by

$$\Omega = \Sigma^{-1} = \begin{array}{c} \\ x \\ y \\ z \end{array}\begin{array}{ccc} x & y & z \\ \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ -1 & -1 & 3 \end{pmatrix} \end{array}, \tag{4}$$

such that $\Sigma \times \Omega$ equals the identity matrix $\mathbf{I}$:

$$\Sigma \times \Omega = I = \begin{array}{c} \\ x \\ y \\ z \end{array}\begin{array}{ccc} x & y & z \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{array}. \tag{5}$$

The zeros in the precision matrix indicate conditional independence of two variables – X

and Y, conditioned on the third variable Z (or a set of variables). Correlation is a

normalized covariance [43]

$$\rho_{XY} = \frac{\text{cov}(X, Y)}{\sigma_X \sigma_Y}, \tag{6}$$

and the inversion of the covariance matrix yields a partial correlation structure between

all the variables. The partial correlation coefficient $\rho_{x_i x_j \cdot \text{rest}}$ between X_i and X_j

variables of a partial correlation matrix, given all other variables, is associated with the

precision matrix $\Omega = \omega_{ij} = \Sigma^{-1}$ such that

$$\rho_{x_i x_j \cdot \text{rest}} = -\frac{\omega_{ij}}{\sqrt{\omega_{ii}\omega_{jj}}}, \tag{7}$$

where a partial correlation matrix entry $\omega_{ij} = 0$ indicates a zero partial correlation

(absence of an edge in the network) [40].

Nonparanormal transformation

However, GGMs assume that the modeled variables follow a multivariate normal distribution, i.e., are jointly Gaussian such that for a $\mathbf{X} \in \mathbb{R}^{n \times p}$ where the rows X_i, $i = 1, \ldots, n$, are independently drawn from the same p-variate Gaussian distribution $\mathcal{N}(\mu, \Sigma)$ [44]. This assumption is not met when modeling RRBS data, since the violation of the normality assumption is frequently observed in methylation data analysis [45]. This violation originates from the fact that methylation values are bounded between 0 and 1, and their variance is usually smaller near the boundaries than near the middle of the interval, implying the violation of the homoscedasticity assumption [46]. Therefore, methylation values follow a scaled beta distribution. If not accounted for, the distribution of the project data would limit the detection of true positive edges in the networks. Consequently, gene methylation data from each tissue at developmental stage had to be transformed prior to network modeling.

The need for a multivariate Gaussian distribution between model variables implies a requirement for normality in the marginal distributions and linearity in all relationships between variables [47]. To achieve that, we transformed the RRBS data by estimating the Gaussian copula to relax the assumption of normality [48]. A copula is a specific instance of a multivariate cumulative distribution function characterized by having uniform marginal distributions, such that in a 2D ($p = 2$) setting, the marginal distributions are

$$X_1 \sim \mathcal{N}(0, 1) \quad \text{and} \quad X_2 \sim \mathcal{N}(0, 1). \tag{8}$$

[49] proposed the nonparanormal (NPN) transformation, a special case of a Gaussian copula, to incorporate non-Gaussian data to overcome the limiting assumption of multivariate normal data. In a 2D setting, the nonparanormal transformation aims to establish a set of monotonic functions f_1, f_2 for a 2-dimensional random variable $X = (X_1, X_2)$ such that the set of functions is multivariate normal distributed and $X \sim \mathrm{NPN}(\mu, \Sigma, f)$. This transformation preserves the conditional (in)dependence structure between the initial, non-Gaussian variables since each function f_i depends solely on the X_i variable. Therefore, the distribution of $f(X)$ has the same factorization as the distribution of X [49]. With NPN transformation the uniform marginal

distributions (equation 8) are transformed by a set of functions such that $f_1(X_1) = u_1$ [229]

and $f_2(X_2) = u_2$, with the corresponding Gaussian copula [230]

$$C(u_1, \ldots, u_p) = \Phi_{\mu,\Sigma}(\Phi^{-1}(u_1), \ldots, \Phi^{-1}(u_p)), \qquad (9)$$

where Φ^{-1} denotes the inverse normal standard cumulative distribution function (CDF), [231]

and Φ_2 denotes the CDF of multivariate distribution $N\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}\right)$ [49]. Here, [232]

we provided an explanation for a 2D setting; however, the same operation could be [233]

easily extended to higher dimensions, since the estimation of a nonparanormal [234]

transformation is computationally efficient, requiring only one pass of $X \in \mathbb{R}^{n \times p}$ [50]. [235]

We implemented the nonparanormal transformation using the `huge` R package, which [236]

estimates the Gaussian copula by transforming the variables marginally by applying [237]

smooth, i.e., continuously differentiable, functions [50]. We conducted the NPN [238]

transformation via the shrunken empirical cumulative distribution function (ECDF), [239]

where the parameter values are shrunken during transformation. We performed this [240]

transformation of methylation values separately for each tissue at developmental stage. [241]

This shrunken estimate helps to reduce the impact of extreme observations. Therefore, [242]

it can be effective in mitigating the impact of extreme values in situations where data [243]

values are constrained within a specific range, e.g., methylation values between 0 and 1. [244]

In this setting, we consider values approaching 0 or 1 to be extreme. [245]

Hierarchical clustering of RNA-Seq data [246]

After transforming methylation values separately for each tissue at developmental stage, [247]

we performed hierarchical clustering of genes and samples using RNA-Seq data. Our [248]

goal was to visualize the correlation-based relationships between samples by utilizing [249]

gene expression data, to facilitate the construction of penalty matrices guiding joint [250]

fused ridge network extraction from methylation data. To do so, we extracted unique [251]

Ensembl gene IDs for each tissue-developmental stage combination. These genes were [252]

then matched with a TPM (transcript per million) file provided by GENE-SWitCH to [253]

extract the count-based RNA-Seq transcript expression levels of only the genes that [254]

were CpG methylated at the promoter region and differentially expressed between [255]

30-70dpf and 70dpf-NB. Furthermore, we filtered out genes located on the X ~256~
chromosome, ultimately obtaining expression values of 2718 unique genes present in all ~257~
tissues at developmental stages for clustering. TPM accounts for sequencing depth, ~258~
which is the total number of reads obtained from an RNA-Seq run, and gene length ~259~
[51]. Nevertheless, the expression data needed pre-processing. All TPM values were ~260~
log-transformed, such that $TPM = \log_2(TPM + 1)$, to ensure that they are non-zero ~261~
allowing normalization prior to clustering. Moreover, the RNA-Seq data were scaled per ~262~
gene. After log transformation, the distribution of TPM-based gene expression ~263~
approached a normal distribution, whereas scaling brought the variables to a comparable ~264~
scale. Correlation-based hierarchical clustering followed. The pairwise similarity ~265~
between genes was computed as the Pearson correlation since the expression data were ~266~
approximately normally distributed. However, since scaling was performed on genes, not ~267~
samples, we computed the pairwise similarity between samples as the nonparametric ~268~
Spearman correlation. Both genes and samples were clustered using complete linkage to ~269~
create compact, distinct clusters. Finally, we implemented the average silhouette width ~270~
(AWS) method to choose the optimal number of sample clusters [52]. A silhouette ~271~
represents each cluster by comparing cluster tightness and separation, illustrating which ~272~
data points are positioned within the assigned cluster and which lie between the clusters ~273~
[53]. AWS is calculated for each data point by measuring its intra-cluster and ~274~
inter-cluster distance, with high values indicating better clustering [52]. The optimal ~275~
number of clusters was chosen as k = 16, the location of the maximum, indicating tight, ~276~
well-separated clusters. We subsequently generated a heatmap illustrating the ~277~
hierarchical clustering of genes and samples based on the preprocessed RNA-Seq data. ~278~

Network extraction ~279~

In preparation for network extraction, all the methylation matrices obtained after: (i) ~280~
Taking a median of methylation values per gene; (ii) Removing genes with no ~281~
between-sample variation; (iii) Scaling methylation values per gene; and (iv) ~282~
Implementing NPN transformation, were used to obtain an intersection of genes that ~283~
were present in all seven tissues and three developmental stages. However, since only ~284~
one gene was shared between all the matrices, germ-layer-specific intersections of genes ~285~

were obtained to allow for network modeling of methylation dynamic guiding pig fetal   286

development. Thereafter, the methylation matrices were filtered to contain methylation   287

values of only the genes present in the intersection. In Table 1 we provide an overview   288

of the 21 matrices   289

[3 tissues × 3 developmental stages + 2(2 tissues × 3 developmental stages)] obtained   290

for further analysis.   291

Table 1. An overview of the germ layer-specific methylation matrices obtained for
downstream analyses

	Liver	Lung	Intestine	Kidney	Muscle	Hindbrain	Skin
Endoderm	X	X	X				
Mesoderm				X	X		
Ectoderm						X	X

The methylation matrices consisted of: (i) $n = 3, p = 61$ for endoderm-specific matrices;   292

(ii) $n = 3, p = 48$ for mesoderm-specific matrices; and (iii) $n = 3, p = 74$ for   293

ectoderm-specific matrices. In Supplementary File 1, Tables S1-S3, we provide an   294

overview of all germ layer-specific genes that we used for network modeling. We will   295

now outline the procedure to obtain a joint estimation of germ layer- and class-specific   296

regularized precision matrices.   297

First, let y_{ig} denote a realization of a p-dimensional random variable (methylation   298

value of a gene), for $i = 1, \ldots, n_g$ independent observations (biological replicates)   299

corresponding to $g = 1, \ldots, G$ non-overlapping classes (tissues at developmental stages),   300

drawn from a multivariate normal distribution $Y_g \sim N_p(0, \Sigma_g)$. The parameters of the   301

multivariate Gaussian – sample mean vector and sample precision matrix $S_g^{-1} = \Omega_g$ can   302

be derived through maximum likelihood estimation (MLE). The multivariate normal   303

log-likelihood parameterized in terms of the precision matrix for the joint $\sum_{g=1}^{G} n_g$ data   304

observations is defined by   305

$$L(\{\Omega_g\}; \{S_g\}) \propto \sum_g n_g \left(\ln |\Omega_g| - \mathrm{tr}(S_g \Omega_g)\right), \qquad (10)$$

where $\{\Omega_g\}$ and $\{S_g\}$ denote sets $\{\Omega_g\}_{g=1}^G$ and $\{S_g\}_{g=1}^G$ respectively, and the maximum   306

likelihood estimate $\hat{\Omega}_g^{ML} = S_g^{-1}$ when $n_g > p$ [41,54]. To derive the parameters of the   307

multivariate Gaussian through MLE we want to invert the sample covariance matrix S_g 308

to obtain the precision matrix Ω_g for each class g [41]. However, it is not possible with   309

variable dimension (p) exceeding the observation dimension (n) [41]. In such a case, the covariance matrix does not have a full rank since the!re are not enough observations to span the information of the variable dimension, and thus, it cannot be inverted. Since the variable dimension (genes) exceeds the observation dimension (biological replicates) in our data, a joint fused ridge estimation will allow the sample covariance matrix inversion, while retaining class-specific information [41]. Therefore, we seek to derive the MLE adapting formula in 10 to include the fused l_2 given by

$$-\sum_{g} \frac{\lambda_{gg}}{2} \|\Omega_g - T_g\|_2^2 + \sum_{(g_1,g_2)} \frac{\lambda_{g_1 g_2}}{4} \|(\Omega_{g_1} - T_{g_1}) - (\Omega_{g_2} - T_{g_2})\|_2^2 \qquad (11)$$

where T_g represent class-specific target matrices that can serve to incorporate prior information of the network structure by integrating known relationships or dependencies between variables into the matrix, λ_{gg} represent class-specific ridge penalty parameters controlling the rate of shrinkage of the unbiased, high variance Ω_g towards the biased, low variance T_g, and $\lambda_{g_1 g_2}$ denote pair-specific fusion penalty parameters controlling the extent of entry-wise similarities between $(\Omega_g - T_{g_1})$ and $(\Omega_g - T_{g_2})$ to be retained [41, 54]. Considering the fused ridge penalty in 11, the maximizing argument for a single data class g is defined by [54] as:

$$\hat{\Omega}_g(\Lambda, \{\Omega_{g'}\}_{g' \neq g}) =$$

$$\left[\lambda_g I_p + \frac{1}{4}(S_g - \lambda_g T_g)^2 \right]^{1/2}$$

$$+ \frac{1}{2}(S_g - \lambda_g T_g)^{-1}, \quad (12)$$

where

$$S_g = S_g - \sum_{g' \neq g} \frac{\lambda_{g'g}}{n_g}(\Omega_{g'} - T_{g'}), \quad T_g = T_g, \qquad (13)$$

and

$$\lambda_g = \frac{\lambda_{g\bullet}}{n_g}, \qquad (14)$$

with $\lambda_{g\bullet}$ denoting the sum of entries in the gth row or column of the penalty matrix Λ. The penalty matrix Λ can store the class-specific (ridge) and pair-specific (fusion) penalties with λ_{gg} along the diagonal and $\lambda_{g_1 g_2}$ off-diagonal [41]. Since GGMs are graphs representing complex systems, e.g., methylation values of DEGs on a tissue- and time-scale, that partially share a common structure across data classes while still retaining class-specific features, setting uniform class-specific ridge penalties and uniform pair-specific fusion penalties is deemed restrictive [54, 55]. The fused l_2-penalized maximum likelihood approach allows simultaneous estimation of multiple precision matrices from high-dimensional data classes. In the analysis of a factorial design, such as tissues at different developmental stages, a more precise regulation of the individual values of λ_{gg} and $\lambda_{g_1 g_2}$ is suitable for the analysis [54]. It can be achieved by, e.g., informing the penalty matrix Λ with RNA-Seq hierarchical clustering results.

Therefore, we utilized the hierarchical clustering results to inform three separate penalty structures, such that each unique data class (tissue at developmental stage) was assigned a ridge penalty, assuming considerable differences in class-specific methylation patterns. Furthermore, fusion penalties were free to be estimated for data classes clustering the closest together. Figure providing visualization of hierarchical clustering results will follow in the Hierarchical clustering of RNA-Seq data section of the Results. Here we provide an overview of the penalty structures defined for the endoderm:

$$\Lambda^{En} = \begin{array}{c} \\ \textbf{L30} \\ \textbf{L70} \\ \textbf{LNB} \\ \textbf{Lu30} \\ \textbf{Lu70} \\ \textbf{LuNB} \\ \textbf{I30} \\ \textbf{I70} \\ \textbf{INB} \end{array} \begin{array}{c} \begin{array}{ccccccccc} \textbf{L30} & \textbf{L70} & \textbf{LNB} & \textbf{Lu30} & \textbf{Lu70} & \textbf{LuNB} & \textbf{I30} & \textbf{I70} & \textbf{INB} \end{array} \\ \left[\begin{array}{ccccccccc} \lambda_{L30} & \lambda_{L30L70} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \lambda_{L30L70} & \lambda_{L70} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \lambda_{LNB} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \lambda_{Lu30} & 0 & 0 & \lambda_{Lu30I30} & 0 & 0 \\ 0 & 0 & 0 & 0 & \lambda_{Lu70} & \lambda_{Lu70LuNB} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \lambda_{Lu70LuNB} & \lambda_{LuNB} & 0 & 0 & 0 \\ 0 & 0 & 0 & \lambda_{Lu30I30} & 0 & 0 & \lambda_{I30} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \lambda_{I70} & \lambda_{I70INB} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \lambda_{I70INB} & \lambda_{INB} \end{array} \right] \end{array},$$

mesoderm:

$$
\Lambda^{Me} = \begin{array}{c} \\ \textbf{K30} \\ \textbf{K70} \\ \textbf{KNB} \\ \textbf{M30} \\ \textbf{M70} \\ \textbf{MNB} \end{array}
\begin{array}{cccccc}
\textbf{K30} & \textbf{K70} & \textbf{KNB} & \textbf{M30} & \textbf{M70} & \textbf{MNB} \\
\left[\begin{array}{cccccc}
\lambda_{K30} & \lambda_{K30K70} & 0 & 0 & 0 & 0 \\
\lambda_{K30K70} & \lambda_{K70} & 0 & 0 & 0 & 0 \\
0 & 0 & \lambda_{KNB} & 0 & 0 & 0 \\
0 & 0 & 0 & \lambda_{M30} & 0 & 0 \\
0 & 0 & 0 & 0 & \lambda_{M70} & \lambda_{M70MNB} \\
0 & 0 & 0 & 0 & \lambda_{M70MNB} & \lambda_{MNB}
\end{array} \right]
\end{array},
$$

and ectoderm:

$$
\Lambda^{Ec} = \begin{array}{c} \\ \textbf{H30} \\ \textbf{H70} \\ \textbf{HNB} \\ \textbf{S30} \\ \textbf{S70} \\ \textbf{SNB} \end{array}
\begin{array}{cccccc}
\textbf{H30} & \textbf{H70} & \textbf{HNB} & \textbf{S30} & \textbf{S70} & \textbf{SNB} \\
\left[\begin{array}{cccccc}
\lambda_{H30} & \lambda_{H30H70} & 0 & 0 & 0 & 0 \\
\lambda_{H30H70} & \lambda_{H70} & 0 & 0 & 0 & 0 \\
0 & 0 & \lambda_{HNB} & 0 & 0 & 0 \\
0 & 0 & 0 & \lambda_{S30} & 0 & 0 \\
0 & 0 & 0 & 0 & \lambda_{S70} & \lambda_{S70SNB} \\
0 & 0 & 0 & 0 & \lambda_{S70SNB} & \lambda_{SNB}
\end{array} \right]
\end{array}.
$$

The `optPenalty.fused()` function from the `rags2ridges` R package determined each germ layer's optimal ridge and fusion penalty parameters [41]. It utilized leave one out cross validation (LOOCV) of the negative fused log-likelihood score and employed the Nelder-Mead algorithm for function minimization [41,56]. Thereafter, we used local false discovery rate (lFDR) with a 0.99 threshold to sparsify the regularized partial correlation matrices. The lFDR is the empirical posterior probability that an edge is null, given the observed partial correlations [41]. An edge will be retained if its posterior probability of being present is larger than or equal to the specified threshold. In our case, $(1 - \text{lFDR}) \geq 0.99$. In Table 2, Table 3 and Table 4, we summarize the

numbers and percentages of the edges retained for network visualization for each tissue   358

at developmental stage.   359

Table 2. Number and percentage of edges retained for each tissue at developmental stage originating from endoderm

	Liver			Lung			Intestine		
	30dpf	70dpf	NB	30dpf	70dpf	NB	30dpf	70dpf	NB
No.	1271	1304	0	0	1679	1762	1292	0	1294
%	69.45	71.26	0	0	91.75	96.28	70.6	0	70.71

Table 3. Number and percentage of edges retained for each tissue at developmental stage originating from mesoderm

	Muscle			Kidney		
	30dpf	70dpf	NB	30dpf	70dpf	NB
No.	0	0	744	756	741	0
%	0	0	65.96	67.02	65.69	0

Table 4. Number and percentage of edges retained for each tissue at developmental stage originating from ectoderm

	Hindbrain			Skin		
	30dpf	70dpf	NB	30dpf	70dpf	NB
No.	1966	0	0	0	2425	1731
%	72.79	0	0	0	89.78	64.09

Network visualization   360

Network visualization of time-dependent, tissue-specific methylation patterns followed   361

the extraction of regularized and sparsified partial correlation matrices. We used the   362

`Union()` function to visualize class-specific networks in the same coordinates between   363

developmental stages for each tissue [41]. This allowed for the detection of differential   364

connections between class-specific networks. Therefore, we obtained unions of 30-70dpf   365

and 70dpf-NB germ-layer specific, sparsified partial correlation matrices for each tissue,   366

sub-setting them to features containing nonzero row (column entries) so that the feature   367

was connected in at least one of the time-specific networks. Using the `DiffGraph()`   368

function, we visualized two differential networks (30-70dpf and 70dpf-NB) for each   369

tissue, with the `Ugraph()` 30dpf networks determining the coordinates of 70dpf   370

networks, whereas 70dpf networks determined the coordinates of NB networks [41]. The Fruchterman-Reingold (FR) algorithm was used to place the nodes, minimizing the number of crossing edges while ensuring that all edges were approximately equal in length [57]. To enhance the visual comparison of time-dependent differential networks, we indicated nodes (genes) to be either up-regulated (red) or down-regulated (green) between developmental stages. We extracted this information from DGE analysis results obtained by the GENE-SWitCH.

Network analysis

Network analysis followed the visualizations of time- and tissue-dependent methylation patterns. First, we detected communities in our networks with the `cluster_fluid_communities()` function from the `igraph` R package [58,59]. Communities in the network are nodes that are more densely connected with each other than with other nodes in the network [60]. While extracting this structural information of a network, the fluid community detection algorithm allows nodes to belong to multiple communities simultaneously [61]. Based on network topologies, we set two communities to be found for each network to capture distinct groups of nodes with cohesive interactions while maintaining graph interpretability. Second, network statistics were derived using the `GGMnetworkStats.fused()` function for each differential network [41]. In particular, we focused on genes with high betweenness centrality, indicating that these nodes are particularly important for the information flow in the networks. We did not specify a universal threshold, but rather focused on capturing a trend in the range of high betweenness centrality values in each network. We thereafter converted the gene IDs of genes with high betweennesss centrality from the Ensembl Sscrofa11.1 to Entrez IDs of human orthologs with `g:Profiler` [62]. Next, we conducted gene set enrichment analysis with `Enrichr` [63–65]. As a background, we used the single cell gene expression profiles from Developmental Single Cell Atlas of Gene RegulaTion and Expression (DESCARTES) ~ 4 million single cells from >110 samples representing 15 fetal organs (72–129 days post fertilization) [66]. This would allow us to evaluate if we are getting biologically feasible results for tissue-specific methylation networks obtained with the joint fused ridge approach we

implemented. Thereafter, we submitted the converted gene sets with high betweennesss   401
centrality for analysis with the ChIP-X Enrichment Analysis Version 3 (`ChEA3`) tool, to   402
predict transcription factors (TFs) associated with the obtained methylation networks   403
[67]. `ChEA3` computes the overlap between the submitted gene set and `ChEA3` libraries of   404
the TF target gene set (human and mouse) with Fisher's Exact Test, using a   405
background of 20000 genes [67]. Benjamini-Hochberg correction is thereafter used to   406
compute FDRs for each gene set in the library separately [67]. The correction is   407
followed by assigning a rank to each gene set, such that 1 indicates a gene set having   408
the lowest corrected p value from the Fisher's Exact Test [67]. TFs are proteins   409
regulating gene expression by binding to DNA sequence, and DNA methylation at   410
promoters is thought to repress TFs binding [21, 68]. However, the interplay between   411
gene expression, DNA methylation and TFs is even more complex, as it was recently   412
proposed that TF binding can also inhibit DNA methylation [69]. Therefore, adding   413
this additional layer of complexity to our network analysis, could provide is more   414
complete insights into methylation dynamics driving pig fetal development.   415

Results   416

Hierarchical clustering of RNA-Seq data   417

We performed hierarchical clustering of RNA-Seq data to obtain insights into gene   418
expression patterns underlying tissue diversification during pig fetal development.   419
Similarities in expression profiles between samples were of interest, and we identified 16   420
optimal clusters with AWS (see Supplementary File 1). In Fig 2 we present the results   421
from hierarchical clustering of the gene expression of CpG methylated at promoter   422
regions DEGs between developmental stages, and samples representative of all seven pig   423
tissues at three developmental stages. We excluded genes located on the X chromosome   424
from the analysis. Moreover, we utilized the results from hierarchical clustering to   425
decide where we should apply fusion during regularization for network extraction,   426
therefore encoding prior, biologically meaningful information in the penalty matrices.   427
Liver, kidney, and hindbrain samples were clustered by tissue, whereas lung, ileum,   428
muscle, and skin clustered by developmental stage. Additionally, the following samples   429

Fig 2. Hierarchical clustering results of TPM (Transcripts Per Million) gene expression data of all samples. Genes are clustered in rows, and samples are in columns. The row z-score corresponds to the fact that gene expression data has been mean-centered and scaled, as $z = \frac{(y-\mu)}{sd}$. Stage 1 denotes 30 days post-fertilization, stage 2 denotes 70 days post-fertilization, and stage 3 denotes newborn. There are four replicates for each tissue at developmental stage. Liver, kidney and hindbrain cluster by tissue, meaning that all samples coming from the same tissue cluster together

originating from the same germ layer clustered more closely together than with other samples: (i) Liver 30dpf – Liver 70dpf; (ii) Lung 30dpf – Ileum 30dpf; (iii) Lung 70dpf – Lung NB; (iv) Ileum 70dpf – Ileum NB; (v) Kidney 30dpf – Kidney 70dpf; (vi) Muscle 70dpf – Muscle NB; (vii) Hindbrain 30dpf – Hindbrain 70dpf; and (viii) Skin 70dpf – Skin NB. This information has been reflected in the penalty matrices, separate for each embryological layer, as shown in the Network extraction section of the Materials and methods.

Network extraction

Here, we present the results from jointly estimating multiple regularized partial correlation matrices under optimal penalty matrices informed by hierarchical clustering results. The optimal ridge and fusion penalty parameters estimated for each embryological layer are demonstrated. The optimal penalties were determined via LOOCV for all genes present in germ layer-specific intersections. The optimal penalties for the 61 genes found in all endoderm samples were

$$
\Lambda^{En} = \begin{array}{c} \\ \text{L30} \\ \text{L70} \\ \text{LNB} \\ \text{Lu30} \\ \text{Lu70} \\ \text{LuNB} \\ \text{I30} \\ \text{I70} \\ \text{INB} \end{array}
\begin{bmatrix}
\overset{\text{L30}}{2.34\times10^2} & \overset{\text{L70}}{5.16\times10^{-20}} & \overset{\text{LNB}}{0} & \overset{\text{Lu30}}{0} & \overset{\text{Lu70}}{0} & \overset{\text{LuNB}}{0} & \overset{\text{I30}}{0} & \overset{\text{I70}}{0} & \overset{\text{INB}}{0} \\
5.16\times10^{-20} & 3.83\times10^2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 2.37\times10^{11} & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 2.37\times10^8 & 0 & 0 & 1.68\times10^{-21} & 0 & 0 \\
0 & 0 & 0 & 0 & 1.40\times10^2 & 8.54\times10^{-17} & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 8.54\times10^{-17} & 2.39 & 0 & 0 & 0 \\
0 & 0 & 0 & 1.68\times10^{-21} & 0 & 0 & 9.16\times10^2 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1.13\times10^7 & 3.21\times10^{-25} \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 3.21\times10^{-25} & 6.91\times10^2
\end{bmatrix},
$$

for the 48 genes present in all mesoderm samples:

$$
\Lambda^{Me} = \begin{array}{c} \\ \text{K30} \\ \text{K70} \\ \text{KNB} \\ \text{M30} \\ \text{M70} \\ \text{MNB} \end{array}
\begin{bmatrix}
\overset{\text{K30}}{2.56} & \overset{\text{K70}}{6.95\times10^{-23}} & \overset{\text{KNB}}{0} & \overset{\text{M30}}{0} & \overset{\text{M70}}{0} & \overset{\text{MNB}}{0} \\
6.95\times10^{-23} & 2.17\times10^6 & 0 & 0 & 0 & 0 \\
0 & 0 & 2.08\times10^2 & 0 & 0 & 0 \\
0 & 0 & 0 & 2.56\times10^2 & 0 & 0 \\
0 & 0 & 0 & 0 & 2.28\times10^2 & 1.83\times10^{-20} \\
0 & 0 & 0 & 0 & 1.83\times10^{-20} & 2.42\times10^7
\end{bmatrix},
$$

and the 74 genes present in all ectoderm samples:

$$
\Lambda^{Ec} = \begin{array}{c} \\ \text{H30} \\ \text{H70} \\ \text{HNB} \\ \text{S30} \\ \text{S70} \\ \text{SNB} \end{array}
\begin{bmatrix}
\overset{\text{H30}}{2.07\times10^2} & \overset{\text{H70}}{1.23\times10^{-19}} & \overset{\text{HNB}}{0} & \overset{\text{S30}}{0} & \overset{\text{S70}}{0} & \overset{\text{SNB}}{0} \\
1.23\times10^{-19} & 6.21\times10^9 & 0 & 0 & 0 & 0 \\
0 & 0 & 3.49\times10^7 & 0 & 0 & 0 \\
0 & 0 & 0 & 1.42\times10^7 & 0 & 0 \\
0 & 0 & 0 & 0 & 6.86\times10^2 & 2.69\times10^{-22} \\
0 & 0 & 0 & 0 & 2.69\times10^{-22} & 1.18\times10^2
\end{bmatrix}.
$$

The penalty values for every germ layer-specific network extraction indicated strong differences in class-specific partial correlation matrices, favoring class-specific regularization over retaining entry-wise similarities between data classes. Moreover, we used spectral condition number plots to assess the conditioning of class-specific partial correlation matrices estimated under the optimal ridge penalties. An example plot can be found in Supplementary File 1.

Network visualization

Following the optimal penalties extraction, conditioning assessment and support

determination with a lFDR threshold of 0.99, we visualized the differential networks.

The following networks were empty: Liver NB, Lung 30dpf, Intestine 70dpf, Kidney NB,

Muscle 30dpf, Muscle 70dpf, Hindbrain 70dpf, Hindbrain NB, Skin 30dpf. Therefore,

the differential visualizations could not be obtained for these tissues at developmental

stages. Here, in Fig 3, we provide an example of a network obtained for liver between

early organogenesis and late organogenesis. We present all the rest visualized

differential networks in Supplementary File 1.

Fig 3. Differential network between Liver stage 30dpf and Liver stage 70dpf. Red nodes indicate up-regulated genes and green down-regulated genes. Edges unique to class Liver 30dpf are visualized in pink, while edges unique to class Liver 70dpf are in orange. Solid edges correspond to positive partial correlations, whereas dashed edges indicate negatively weighted partial correlations

The Lung 70dpf-NB and Skin 70dpf-NB networks showed a sparse *hairball* structure. Liver 30-70dpf and Skin 70dpf-NB were more densely connected than other differential networks extracted. Furthermore, there were noticeably more edges unique to late organogenesis (70dpf) in the skin network, while in the lung network edges unique to the newborn stage (NB).

Network analysis

Continuing with network analysis, based on network topologies we decided to detect two fluid communities for each tissue developmental stage combination. In Fig 4 we provide an example of community detection visualization obtained for the Liver 30dpf network. The rest of the networks visualizing fluid community structure can be found in Supplementary File 1.

Vertices

1: ENSSSCG00000000906
2: ENSSSCG00000001906
3: ENSSSCG00000002882
4: ENSSSCG00000003212
5: ENSSSCG00000003225
6: ENSSSCG00000003482
7: ENSSSCG00000004119
8: ENSSSCG00000004760
9: ENSSSCG00000006399
10: ENSSSCG00000006579
11: ENSSSCG00000007508
12: ENSSSCG00000007968
13: ENSSSCG00000008574
14: ENSSSCG00000009209
15: ENSSSCG00000009558
16: ENSSSCG00000009616
17: ENSSSCG00000012909
18: ENSSSCG00000012993
19: ENSSSCG00000013604
20: ENSSSCG00000014110
21: ENSSSCG00000014818
22: ENSSSCG00000015049
23: ENSSSCG00000015091
24: ENSSSCG00000015126
25: ENSSSCG00000016504
26: ENSSSCG00000016728
27: ENSSSCG00000017433
28: ENSSSCG00000017458
29: ENSSSCG00000017490
30: ENSSSCG00000023018
31: ENSSSCG00000024158
32: ENSSSCG00000024892
33: ENSSSCG00000024917
34: ENSSSCG00000025128
35: ENSSSCG00000026345
36: ENSSSCG00000026427
37: ENSSSCG00000026680
38: ENSSSCG00000029503
39: ENSSSCG00000029752
40: ENSSSCG00000029763
41: ENSSSCG00000029960
42: ENSSSCG00000030561
43: ENSSSCG00000030903
44: ENSSSCG00000031345
45: ENSSSCG00000032407
46: ENSSSCG00000034211
47: ENSSSCG00000035445
48: ENSSSCG00000035594
49: ENSSSCG00000035785
50: ENSSSCG00000036403
51: ENSSSCG00000036844
52: ENSSSCG00000037562
53: ENSSSCG00000038067
54: ENSSSCG00000038768
55: ENSSSCG00000038991
56: ENSSSCG00000040035
57: ENSSSCG00000040663
58: ENSSSCG00000045348
59: ENSSSCG00000050178
60: ENSSSCG00000050498
61: ENSSSCG00000051187

Fig 4. Communities detected for Liver 30dpf with the `cluster_fluid_communities()` function from the igraph R package. Each community is assigned a different colour. The legend on the right provides the encoding of the Ensembl gene IDs in the network

Following community detection, we focused on analyzing network statistics. In this section, we discuss human orthologs of genes shared between the identified communities, indicating their multi-functionality. We also provide results from the gene set enrichment analyses with `Enrichr` using DESCARTES. Additionally, we present results of the TF over-representation analysis with `ChEA3`. We present our findings per tissue at developmental stage.

Endoderm - liver

Starting with Liver 30dpf, two genes were shared between the communities, namely Cytochrome P450 family 1 subfamily A polypeptide 1 (CYP1A1) and Small Integral Membrane Protein 24 (SMIM24). While the CYP1A1 enzyme is significant throughout the entire duration of pregnancy playing a critical role in metabolizing xenobiotics within the placenta, the SMIM24 progenitor gene has been found to be expressed in the yolk sack pre-macrophages [70, 71]. The betweennesss centrality of the genes extracted for downstream analyses was between 3.45 and 33.50. Two cell types showed significant enrichment after adjusting the p-value using the Benjamini-Hochberg method, namely *Ductal Cells in Pancreas* and *Lymphoid Cells in Liver*, as seen in Table 5.

Table 5. Enrichment analysis results for the Liver 30dpf network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Ductal cells in Pancreas	0.0031	0.0347	28.06	162.11
2	Lymphoid cells in Liver	0.0033	0.0347	27.14	155.01
3	Corneal and conjunctival epithelial cells in Eye	0.0211	0.1200	10.15	39.17
4	Endocardial cells in Heart	0.0229	0.1200	47.97	181.23
5	Vascular endothelial cells in Lung	0.0372	0.1451	28.93	95.26

The enrichment for *Ductal Cells in Pancreas* is not surprising, as the endocrine, exocrine and ductal components of the pancreas originate from endoderm [72]. Furthermore, the fetal liver has been recognized as the major site of the development of the immune system, supporting the enrichment for *Lymphoid Cells in Liver* in a list of high betweennesss centrality genes from the Liver 30dpf network [73]. The innate immune lymphoid cells can also travel to other sites of organogenesis in the developing fetus through the blood [74]. Moving to the TFs, we identified KCNIP3, CREB3L3, MRFL, BARX2 and GRHL3 to be the most substantially over-represented for Liver 30dpf (see Table 6).

Table 6. TF target over-representation results for the Liver 30dpf network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	KCNIP3	16.50	2	LMTK3, LRRC4B
2	CREB3L3	33.67	5	F7, SMIM24, RORC, CYP1A1, CPN2
3	MRFL	45.00	2	SMIM24, MUC2
4	BARX2	50.00	3	TRIM29, PRSS22, MPZL2
5	GRHL3	52.33	3	TRIM29, PRSS22, MPZL2

cAMP responsive element-binding protein 3 like 3 (CREB3L3) is known to be primarily 501
expressed in the liver and small intestine, cooperating to maintain lipid metabolism 502
[75]. Moreover, BarH-like homeobox 2 (BARX2) is crucial during embryonic 503
development, being involved in the epithelial-mesenchymal interactions of organogenesis, 504
and cytoskeletal organization [76,77]. The expression of BARX2 is down-regulated by 505
CpG Island hypermethylation [76]. 506

Continuing with the next developmental stage, three genes were shared between the 507
two fluid communities with Napsin A Aspartic Peptidase (NAPSA) among them. In the 508
developing embryo NAPSA is expressed in mature B-cells in peripheral blood and 509
hematopoietic stem cells in the hematopoietic bone marrow [78]. The betweenness 510
centrality of Liver 70dpf genes extracted for downstream analyses was between 24.35 511
and 28.71. As presented in Table 7, the following cell types showed significant 512
enrichment after FDR correction: *Lymphoid cells in Heart*, *Lymphoid cells in Lung*, 513
Lymphoid cells in Pancreas, *Lymphoid cells in Intestine* and *Lymphoid cells in Adrenal*. 514

Table 7. Enrichment analysis results for the Liver 70dpf network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Lymphoid cells in Heart	0.0039	0.04758	24.54	136.34
2	Lymphoid cells in Lung	0.0071	0.0476	17.87	88.52
3	Lymphoid cells in Pancreas	0.0072	0.0476	17.63	86.88
4	Lymphoid cells in Intestine	0.0076	0.0476	17.17	83.74
5	Lymphoid cells in Adrenal	0.0097	0.0485	15.09	69.95

Although the adrenal medulla originates from mesoderm, while the adrenal cortex from 516
ectoderm, we already established that innate immune lymphoid cells can travel between 517
organogenesis sites [73,79,80]. Moreover, the fetal liver has been shown to 518
accommodate the progenitors of various types of lymphoid cells [73]. Next, we 519
identified IRPF9, ZNF831, BATF2, SCML4 and TBX21 to be the most substantially 520
over-represented TFs for Liver 70dpf (see Table 8). Interferon regulatory factor 9 521

(IRF9) belongs to the IRF TF family involved in the transcriptional regulation of the 522

immune system and cell growth, working at the intersection of innate and adaptive 523

immunity [81]. Moreover, Basic leucine zipper ATF-like transcription factor 2 (BATF2) 524

has been found to be a key factor in myeloid differentiation, therefore regulating 525

immune response [82]. 526

Table 8. TF target over-representation results for the Liver 70dpf network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	IRF9	14.75	7	NAPSA, IFI35, PTPRCAP, GPR37L1, MYO1F, HERPUD1
2	ZNF831	16.33	3	GPR171, PTPRCAP, MYO1F
3	BATF2	27.33	3	NAPSA, IFI35, MYO1F
4	SCML4	28.33	3	GPR171, PTPRCAP, MYO1F
5	TBX21	29.25	5	NAPSA, GPR171, PTPRCAP, MYO1F, HERPUD1

Endoderm - lung 527

Moving on to the lung networks, we identified five genes to be shared between the fluid 528

communities. Two of them are particularly interesting, namely Coagulation Factor X 529

(F10) and Thromboxane A Synthase 1 (TBXAS1). Both are involved in hemostasis, 530

with F10 encoding the vitamin K-dependent coagulation factor X [83,84]. TBXAS1 is 531

a member of the cytochrome P450 superfamily of enzymes, similar to CYP1A1 532

introduced in the Liver 30dpf section [84]. Since both lung and liver originate from the 533

endoderm, this is in line with the hematopoietic function of the liver during fetal 534

development. The betweenness centrality of the top genes from the Lung 70dpf network 535

was between 3.26 and 3.51. We identified no significantly enriched cell types after 536

adjusting the p-value using the Benjamini-Hochberg method. However, without the 537

correction (see Table 9), similar to Liver 30dpf, in Lung 70dpf network *Ductal cells in* 538

Pancreas was the most enriched term. 539

Table 9. Enrichment analysis results for the Lung 70dpf network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score	
1	Ductal cells in Pancreas	0.0057	0.1428	19.81	102.29	
2	PAEP MECOM positive cells in Placenta	0.0123	0.1541	13.16	57.87	540
3	Vascular endothelial cells in Lung	0.0501	0.2805	20.89	62.54	
4	Squamous epithelial cells in Stomach	0.0663	0.2805	15.58	42.28	
5	Lymphoid cells in Heart	0.0995	0.2805	10.13	23.37	

The second most significant term was *PAEP MECOM positive cells in Placenta* (p-value   541
= 0.01232). In male human fetus both *PAEP*-positive and *MECOM*-positive cells in the   542
placenta is associated with the expression of the *Xist* or *Tsix* gene [66]. *Xist* encodes a   543
long non-coding RNA (lncRNA) that coats one of the X chromosomes in females,   544
leading X chromosome inactivation to achieve dosage compensation between genders   545
[85]. *Tsix* is a lncRNA antisense to *Xist* [86]. The presence of these markers is of   546
maternal origin, and while *Xist* corresponds to maternal decidualized stromal cells, *Tsix*   547
corresponds to maternal endometrial epithelial cells [66]. Next, we identified FOXN1,   548
RBPJL, NR1D1, TBPL2 and CENPB to be the most substantially ovverepresented TFs   549
for Lung 70dpf (see Table 10). Forkhead box protein N1 (FOXN1) is responsible for   550
thymus organogenesis, by regulating thymic epithelial cells (TECs), therefore   551
contributing to in utero T-cell development, and aiding the immune response [87–89].   552

Table 10. TF target over-representation results for the Lung 70dpf network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	FOXN1	31.67	4	KRTDAP, CDSN, S100A3, VSIG8
2	RBPJL	33.33	4	RORC, SLC25A45, PRSS22, HERPUD1
3	NR1D1	49.33	5	KRTDAP, NAPSA, CDSN, PRSS22, HERPUD1
4	TBPL2	51.50	4	KRTDAP, NAPSA, CDSN, PRSS22, HERPA45
5	CENB	70.67	3	NAPSA, GPR37L1, SLC2A4RG

Next, the Lung NB network community detection resulted in the most overlapping   553
nodes (26 out of 61 present) from all networks. Therefore, we decided to conduct gene   554
ontology enrichment analysis of the human orthologs of these 26 genes using   555
`g:Profiler` [62]. However, we did not obtain any significant results after FDR   556
correction. Nevertheless, one should keep in mind that this is a relatively small gene set.   557
Continuing with nodes with the highest betweenness centrality, the same 26 genes had   558
betweenness centrality equal to 1.67, the lowest from all the tissues at developmental   559
stages. After FDR correction, no terms showed significant enrichment. The cell types   560
with the highest p value were *Ductal cells in Pancreas* (0.01056), similar to Liver 30dpf   561
and Lung 70dpf, and *IGFBP1 DKK1 positive cells in Placenta* (0.01107), as shown in   562
Table 11.   563

Table 11. Enrichment analysis results for the Lung NB network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Ductal cells in Pancreas	0.0106	0.1273	14.02	63.81
2	IGFBP1 DKK1 positive cells in Placenta	0.0111	0.1273	13.67	61.57
3	AFP ALB positive cells in Spleen	0.0418	0.2450	6.60	20.95
4	PDE1C ACSM3 positive cells in Stomach	0.0470	0.2450	22.15	67.72
5	Vascular endothelial cells in Lung	0.0679	0.2450	15.03	40.43

Insulin-like growth factor-binding protein 1 (IGFBP1), and Dickkopf-related protein 1   564
(DKK1) are marker genes associated with decidual cells, which the decidua (pregnant   565 566
endometrium) consists of, while the immune cells are one of the main cell types making   567
up the decidual cells [90, 91]. We subsequently identified OSR1, NR1I3, BATF2, IRF9,   568
and IRF7 to be the most overrepresented TFs for Lung NB (see Table 12).   569

Table 12. TF target over-representation results for the Lung NB network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	OSR1	26.00	4	RHBDF1, ANO1, FAM180A
2	NR1I3	29.33	4	DMGDH, IGFBP1, F7, CYP1A1, SLC25A45
3	BATF2	34.00	5	ZBP1, NAPSA, IFI35
4	IRF9	50.00	4	ZBP1, NAPSA, IFI35, GPR37L1, SLC25A45
5	IRF7	54.33	3	ZBP1, IFI35

Odd skipped-related 1 (OSR1) is associated with connective tissue formation and   570
myogenesis in limbs, pointing towards mesoderm development, whereas *constitutive*   571
androstane receptor (NR1I3) expression is enriched in adult mammalian liver and its   572
expression in human embryonic stem cells has been linked to differentiation and   573
maturation of hepatic-like cells [92, 93].   574

Mesoderm - kidney   575

Continuing with mesoderm networks, we identified six genes shared between the fluid   576
communities for Kidney 30dpf. On the basis of our previously presented results, we   577
decided to focus on two of them, namely Cytoglobin (CYGB) and the previously   578
introduced CYP1A1. CYGB is unique to vertebrates, encoding a protein called   579
hexacoordinate hemoglobin, distinguishing it from pentacoordinate hemoglobins   580
primarily engaged in oxygen transportation [94]. With its binding mechanism for   581
ligands varying significantly from pentacoordinate hemoglobins, the CYGB protein is   582

believed to safeguard cells during periods of oxidative stress [94]. CYGB has also been [583] revealed to activate hepatic stellate cells (HSC) during disturbances in oxygen [584] homeostasis, aiding in liver fibrogenesis [95]. Furthermore, CYGB has been shown to [585] protect kidney fibroblasts against oxidative stress caused by a lack of blood supply; and [586] therefore, a lack of oxygen [96]. As for CYP1A1, its expression in the kidney increases [587] in response to oxidative stress, whereas in the liver it is involved in the oxidation of [588] fatty acids and toxins [97, 98]. Next, we looked at 11 genes with the highest [589] betweenness centrality (between 27.82 and 32.32) in the Kidney 30dpf network. We did [590] not find any statistically significant terms after FDR correction. The top enriched cell [591] types after correction were *Smooth Muscle Cells in Heart*, *Oligodendrocytes in* [592] *Cerebellum*, *Vascular endothelial cells in Placenta*, *Lymphoid cells in Lung*, and [593] *Lymphoid cells in Placenta* (Table 13). [594]

Table 13. Enrichment analysis results for the Kidney 30dpf network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Smooth muscle cells in Heart	0.0198	0.0887	56.84	222.86
2	Oligodendrocytes in Cerebellum	0.0484	0.0887	22.55	68.29
3	Vascular endothelial cells in Placenta	0.0513	0.0887	21.25	63.11
4	Lymphoid cells in Lung	0.0725	0.0887	14.80	38.82
5	Lymphoid cells in Placenta	0.0739	0.0887	14.50	37.77

The enrichment for *Smooth Muscle Cells in Heart* is not surprising, since smooth muscle [596] cells (SMCs) play a significant role in the structural and functional makeup of numerous [597] organs throughout embryonic development, while the heart is of a mesodermal origin [598] [99, 100]. Moreover, oglidendrocytes wrap myelin, allowing fast and energy efficient [599] neuronal signaling, and nerve-muscle communication governs voluntary movements and [600] breathing [101, 102]. Like the kidney and heart, muscles originate from the mesoderm, [601] while cerebellum and the nervous system from the ectoderm [99, 103]. The development [602] of the nervous system from the ectoderm is initiated by a signal from mesoderm [103]. [603] Additionally, vascular endothelial cells (EC) form a single layer lining the interior of [604] arteries, veins, and capillaries, facilitating development and maintenance of the [605] circulatory system. The endothelium functions as an endocrine, i.e., hormonal organ as [606] well [104, 105]. Coupling this information with the observed enrichment for [607] lymphoid-associated cell types supports our results pointing towards the importance of [608] the immune system and interactions between various tissues (and germ layers) during [609]

fetal development. Eventually, we investigated the top enriched TFs for genes with high   610
betweenness centrality in the Kidney 30dpf network. In Table 14, we present the   611
results from this enrichment analysis.   612

Table 14. TF target over-representation results for the Kidney 30dpf network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	SP140L	21.50	2	MAP4K1, ZBP1
2	MEOX1	43.00	2	CYP1A1, CYGB
3	ZNF831	51.67	3	MAP4K1, ZBP1, MSLNL
4	CC2D1A	55.50	1	HR
5	TBX21	61.00	2	MAP4K1, ZBP1

The top enriched TFs for Kidney 30dpf were SP140L, MEOX1, ZNF831, CC2D1A, and   613
TBX21. The speckled protein (SP) family members are expressed primarily in   614
leukocytes, and the SP140 locus has been associated with multiple autoimmune diseases,   615
such as multiple sclerosis (MS) damaging the central nervous system, which is   616
considered to be a T-cell-mediated disease [106]. Furthermore, mesenchyme homeobox   617
1 (MEOX1) has been identified as a positive regulator of smooth muscle cell   618
differentiation during embryogenesis in mice [107].   619

With respect to the Kidney 70dpf genes, we did not identify any genes that were   620
shared between the two fluid communities. The betweenness centrality of 10 genes   621
extracted for downstream analyses was between 26.60 and 38.48. After applying FDR   622
correction, two cell types showed a significant enrichment, namely *Cardiomyocytes in*   623
Heart and *Megakaryocytes in Muscle* (see Table 15).   624

Table 15. Enrichment analysis results for the Kidney 70dpf network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Cardiomyocytes in Heart	0.007973	0.03446	147.96	714.91
2	Megakaryocytes in Muscle	0.008616	0.03446	17.22	81.88
3	Retinal pigment cells in Eye	0.04603	0.1003	23.77	73.18
4	Chromaffin cells in Adrenal	0.05129	0.1003	21.25	63.11
5	SATB2 LRRC7 positive cells in Heart	0.06267	0.1003	17.24	47.76

Cardiomyocytes are cardiac muscle cells responsible for involuntary contractions of the   626
heart, supplying the organism with blood [108]. Megakaryotes are rare cells found in   627
the bone marrow, responsible for the production of platelets, which aid the formation of   628
blood clots and wound healing [109]. In addition to assisting thrombosis, platelets are   629
know to influence the immune response as well by, e.g., directing T-cell activation and   630

differentiation [110, 111]. The circulatory, renal and musculoskeletal systems originate   631
from the mesoderm, and our results point towards the importance of interactions   632
between different tissues during fetal development [99]. Focusing on TF   633
over-representation analysis, the top five overrepresented TFs for Kidney 70dpf were   634
BORCS8-MEF2B, SNAI3, NKX2-5, HLF and THRA (see Table 16).   635

Table 16. TF target over-representation results for the Kidney 70dpf network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	BORCS8-MEF2B	51.00	2	RGS14, UNC13D
2	SNAI3	53.67	3	RGS14, TRIB3, UNC13I
3	NKX2-5	58.00	1	RNF207
4	HLF	66.67	1	TEF
5	THRA	74.75	2	TEF, LRRTM2

Snail family transcriptional repressor 3 (SNAI3) is a member of the SNAIL gene family,   636
known to be involved in orchestrating mesoderm formation during embryogenesis [112],   637
whereas NK2 homeobox 5 (NKX2-5) encodes a transcription factor involved in heart   638
formation [113].   639

Ectoderm - skin   640

Continuing with the Skin 70dpf networks, we identified two genes: solute carrier family   641
6 member 7 (SLC6A7) and lymphocyte specific protein 1 (LSP1) to be shared between   642
the computed fluid communities. SLC6A7 belongs to the gamma-aminobutyric acid   643
neurotransmitter gene family [114]. It is also responsible for the production of   644
L-proline transporter protein in the mammalian brain [114]. The presence of this gene   645
in the skin network is not surprising, since both the skin and the brain originate from   646
the ectoderm [115]. Additionally, LSP1 encodes a protein expressed in macrophages,   647
lymphocytes, and the endothelium; therefore, it is involved in the immune response   648
[116]. Moving to the 16 genes with the highest betweenness centrality in the Skin 70dpf   649
network (values between 6.40 and 6.52), we did not obtain any significant cell type   650
enrichment results (see Table 17). However, one must keep in mind that we   651
continuously worked with small gene sets in this study, therefore it is not surprising.   652

Table 17. Enrichment analysis results for the Skin 70dpf network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Squamous epithelial cells in Stomach	0.0560	0.2812	18.70	53.86
2	Megakaryocytes in Adrenal	0.0830	0.2812	12.38	30.82
3	Megakaryocytes in Heart	0.0889	0.2812	11.52	27.88
4	Ductal cells in Pancreas	0.0918	0.2812	11.13	26.57
5	Smooth muscle cells in Eye	0.1027	0.2812	9.88	22.47

The top five overrepresented TFs we identified for Skin 70dpf genes were KCNIP3, EBF4, BATF2, SOX9, and ARID5A (Table 18).

Table 18. TF target over-representation results for the Skin 70dpf network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	KCNIP3	45.50	1	NRGN
2	EBF4	58.00	3	NTNG2, ADGRG1, TNRC18
3	BATF2	77.00	2	NTNG2, IFI35
4	SOX9	77.20	3	ADGRG1, TNRC18, NRGN
5	ARID5A	80.50	3	SH3TC1, NR1H2, IFI35

We have already introduced BATF2 in the Liver 70dpf section as a regulator of the immune response. Furthermore, we found that Early B cell factor 4 (EBF4) is expressed in human immune cells, having a cytotoxic lymphocyte function, therefore aiding the immune response [117]. Additionally, SRY-box transcription factor 9 (SOX9) TF is known to be involved in a variety of developmental processes. Not only does SOX9 expression determine cell fate, driving tissue diversification in all germ layers, but it is also a crucial regulator of sex determination during mammalian embryonic development [118, 119].

Finally, we focused on analyzing the Skin NB network. Seven genes were found at the intersection of the two extracted communities. Here, we introduce fucose mutarotase (FUOM): an inhibitor of neuron differentiation [120] that is up-regulated between 70dpf and NB in our differential network visualization (see Supplementary File 1 Fig. S6, here as ENSSSCG00000010791). Moreover, we looked at 9 genes with the highest betweenness centrality in the Skin NB network. Their values were high compared to other tissues at developmental stages (between 62 and 74.2), however, we identified only one significantly enriched cell type after FDR correction, namely *Megakaryocytes in Muscle* (Table 19). We discussed megakaryotes in the Kidney 70dpf section.

Table 19. Enrichment analysis results for the Skin NB network

Index	Name	P-value	Adjusted p-value	Odds Ratio	Combined score
1	Megakaryocytes in Muscle	0.0069	0.0348	19.69	97.79
2	Skeletal muscle cells in Eye	0.0919	0.1714	11.66	27.84
3	AFP ALB positive cells in Spleen	0.1075	0.1714	9.87	22.02
4	Corneal and conjunctival epithelial cells in Eye	0.1371	0.1714	7.59	15.08
5	Hepatoblasts in Liver	0.1923	0.1923	5.21	8.60

Thereafter, we looked at overrepresented TFs in Skin NB networks, and identified ZNF605, ZNF154, ZNF221, ZNF780B, and ZNF626 (Table 20).

Table 20. TF target over-representation results for the Skin NB network

Rank	TF	Mean Rank	Overlapping Genes	Gene IDs
1	ZNF605	19.33	3	ZNF37A, PEG10, PCBD2
2	ZNF154	66.67	3	ZNF37A, GPX7, PCBD2
3	ZNF221	72.00	1	ZNF37A
4	ZNF780B	93.00	1	ZNF37A
5	ZNF626	98.67	3	ZNF37A, PEG10, PCBD2

Interestingly, all the top five TFs belong to the zinc finger proteins (ZNF) family, which is the largest TF family in the human genome [121]. In humans ZNF605, ZNF154, ZNF780B, and ZNF626 exhibit the highest expression in the thyroid (with the thyroid hormones modulating a variety of immune cells), whereas ZNF221 in the brain (originating from ectoderm, similar to skin) [115, 122, 123].

Discussion

As the omics revolution picks up pace, the need for novel approaches to analyzing high-throughput biological data ever-increasing in complexity and size, which can conquer the curse of dimensionality, becomes evident. At the same time, model interpretability and visually appealing analytical approaches are desirable. Therefore, we aimed to determine whether the joint fused ridge network extraction informed by the results from hierarchical clustering of RNA-Seq data can ensure the extraction of the relevant data signal in network modeling of DNA methylation data coming from multiple swine tissues and developmental stages.

The findings concerning hierarchical clustering of the RNA-Seq data revealed that liver, kidney and hindbrain samples clustered by tissue, whereas lung, ileum, muscle and skin clustered by developmental stage. This aligns with the results obtained by the

GENE-SWitCH, even though we worked with a subset of gene expression data, that is, CpG methylated at promoter regions DEGs, excluding genes located on the X chromosome [24]. Additionally, at 30dpf we observed the lung and ileum, which originate from endoderm, clustering together, suggesting that during early organogenesis, they have not yet fully differentiated.

Subsequently, we informed the penalty structures with results from hierarchical clustering to estimate the optimal penalty matrices for each germ layer separately. Ideally, we would have specified one common penalty structure for all tissues and developmental stages together; however, only one gene was shared among all data classes. In that sense, our analysis might be considered reductionist since we have ignored interplay between tissues originating from different germ layers. We observed favoring of class-specific regularization (ridge) over retainment of entry-wise similarities between data classes (fusion) in all germ layer-specific optimal penalty matrices. This suggests that all data classes (tissues at developmental stages) are quite distinct concerning methylation patterns, further emphasizing the dynamic nature of the (epi)genome. The highest class-specific ridge penalty was evaluated by LOOCV for Liver NB, indicating that the precision matrix approached the target matrix, suggesting negligible signal in the data. We also observed high ridge penalties for Lung 30dpf, Intestine 70dpf, Muscle 70dpf, Hindbrain 70dpf, Hindbrain NB, and Skin 30dpf, explaining empty networks obtained for the above-mentioned tissues at developmental stages. However, Kidney 70dpf also had a relatively high ridge penalty compared to other data classes ($\lambda_{gg} = 2.17 \times 10^6$), but we were able to extract the network. Notably, all partial correlation matrices estimated under the optimal penalty parameters determined for germ layer-specific network extraction were well-conditioned, as suggested by condition number plots. This illustrates that the l_2 regularization procedure implemented with `rags2ridges` led to well-conditioned estimates of the partial correlation matrices for network visualization and downstream analyses.

Continuing with cell type enrichment and TF over-representation analysis, using gene sets with the highest betweenness centrality from each data class specific network, we observed biologically feasible results for each tissue. Concerning the recurring trend of immune-response associated terms for all tissues, a fetus is considered vulnerable to infections in utero [124]. However, the maternal-fetal immunological landscape

involving, e.g., the placenta, lymphoid cells, and megakaryotes, supports the developing   725
fetus in all stages of pregnancy [125]. The fact that immune cells in mammals are   726
vertically transmitted from the mother to the fetus in utero, would also explain the   727
observed enrichment for terms associated with the placenta [126]. This transmission   728
promotes neonatal immunity, aiding the newborn in resistance against infections [126].   729
Additionally, a recurring trend of enrichment for terms associated with the circulatory   730
system, further supports our observations, since various types of immune cells are found   731
in blood, circulating through the developing organism [127]. This would also explain   732
why we obtain enrichment for a different tissue that we are analyzing at times, as every   733
tissue belongs to an organ system dependent on nutrient and oxygen supply by the   734
blood for healthy fetal development [128]. The lung and liver are the exceptions, not   735
being sustained by the fetal blood workflow; however, their waste products and carbon   736
dioxide will enter the fetal circulatory system [128]. The results obtained suggests that   737
our analytical framework allows for extraction of biologically feasible data signal from   738
high-dimensional methylation data spanning seven swine tissues at three developmental   739
stages.   740

However, our results should be approached with caution due to the limitations of   741
contemporary omics research and the methodological constraints of the project itself.   742
The sample size is the major issue to consider when aiming to accurately estimate   743
network parameters and, therefore, the network topology. As introduced before, omics   744
data analyses often suffer from the *curse of dimensionality*. Consequently, we   745
implemented joint fused ridge estimation. Nevertheless, conquering singularity does not   746
solve the issues of difficulty in collecting pig tissue samples, especially at early   747
developmental stages. The considerations for ethical animal experimentation call for the   748
adoption of the 4Rs principles (Reduction, Refinement, Replacement and   749
Responsibility) [129], which in essence, require the least feasible number of biological   750
replicates to be collected. Moreover, there is often not enough genetic material in a   751
single sample (which resulted in pooling of samples at 30dpf), and the desired sample   752
itself is hard to extract. To that end, a developing field of organoids, which are   753
lab-grown cell-based in vitro models mimicking the functioning of in-vivo tissues, arises   754
as a valuable alternative to traditional animal experiments [130]. Furthermore, genes   755
involved in early development are often less well annotated than genes expressed in   756

mature tissues. Additionally, literature on swine development at genomics resolution is ₇₅₇ scarce; therefore we conducted cell type enrichment and TF over-representation analyses ₇₅₈ on the human orthologs. Coupled with the small size of the gene sets analyzed, the ₇₅₉ insights obtained from our networks might not represent early developmental stages in ₇₆₀ swine accurately, since we reviewed literature focusing on human and mice development. ₇₆₁ These limitations highlight the importance and novelty of the efforts of projects such as ₇₆₂ GENE-SWitCH. We believe that our analyses would benefit from utilizing ₇₆₃ (epi)genomics data measured at single-cell resolution, therefore accounting for tissue ₇₆₄ heterogenity, allowing us to extract networks that more accurately represent ₇₆₅ methylation dynamics during pig fetal development. ₇₆₆

Nevertheless, one should realize that in this study, we did not test any biological ₇₆₇ hypothesis, but rather, we propose a novel exploratory analysis approach to extract a ₇₆₈ mechanistic view of the genes and their interactions based on methylation values. For ₇₆₉ that reason, we developed an interpretable and visually appealing workflow for the ₇₇₀ analysis of omics datasets spanning various tissues and developmental stages. And ₇₇₁ although there were limitations associated with gene annotations at early developmental ₇₇₂ stages, literature availability and the dataset itself, we were able to extract biologically ₇₇₃ feasible results with our approach. ₇₇₄

Future research would likely benefit from utilizing single-cell omics data to account ₇₇₅ for tissue heterogeneity. In that setting, we would recommend fusion between cell ₇₇₆ populations within tissues, not the tissues themselves. Additionally, to minimize ₇₇₇ biological noise in the collected samples, organoids could be considered. However, the ₇₇₈ question of how closely can organoids mimic the functioning of in-vivo tissues at a ₇₇₉ molecular level, remains unanswered. Moreover, single-cell sequencing and organoids, ₇₈₀ however promising, are an expensive alternative to more traditional technologies. With ₇₈₁ respect to the joint fused ridge, we identified a lack of fusion penalty parameter ₇₈₂ assessment methods available, which could open avenues for future research. ₇₈₃ Nevertheless, our proposed methodology could be implemented in other omics fields. ₇₈₄ The fused ridge approach has been used successfully in multiple omics domains, such as ₇₈₅ metabolomics, to study blood-based metabolomic signatures in Alzheimer's disease, and ₇₈₆ proteomics to identify sepsis biomarkers from patients' serum [131, 132]. We also see ₇₈₇ potential in exploring differences between tissues (differential networks between tissues), ₇₈₈

instead of focusing on differential molecular dynamics between developmental stages.   789

Conclusion   790

To conclude, we demonstrated that: 791

I. Utilizing RNA-Seq hierarchical clustering results to control the individual values   792
of the ridge and fusion penalties is a feasible approach to incorporate biologically   793
meaningful information in the penalty structures of time- and tissue-dependent   794
swine DNA methylation networks;   795

II. Joint fused ridge regularization procedure with the use of `rags2ridges` R package   796
allows for the extraction of the relevant data signal in an ill-posed problem, namely   797
network modeling of DNA methylation data subjected to class membership.   798

Furthermore, the results from our network analysis point toward the importance of the   799
immune and circulatory system in the developing fetus, shedding light on the dynamic   800
network of interactions between tissues, organ systems and germ layers.   801

List of Abbreviations   802

DNMT	DNA methyltransferase	803
CpG	5'—C—phosphate—G—3'	804
TSS	Transcription Start Site	805
FAANG	Functional Annotation of Animal Genomes consortium	806
RNA-Seq	RNA Sequencing	807
RRBS	Reduced Representation Bisulphite Sequencing	808
PCR	Polymerase Chain Reaction	809
30dpf	30 days post fertilization	810
70dpf	70 days post fertilization	811
NB	Newborn	812

ChIP-Seq	Chromatin Immunoprecipitation assays with Sequncing	813
H3K4me3	Histone 3 Lysine 4 trimethylation	814
bp	Base pair	815
DEG	Differentially Expressed Gene	816
GGM	Gaussian Graphical Model	817
NPN	Nonparanormal	818
CDF	Culmulative Distribution Function	819
ECDF	Empirical Culmulative Distribution Function	820
TPM	Transcripts Per Million	821
AWS	Average Silhouette Width	822
MLE	Maximum Likelihood Estimation	823
LOOCV	Leave One Out Cross Validation	824
lFDR	Local False Discovery Rate	825
FR algorithm	Fruchterman-Reingold algorithm	826
DESCARTES	Developmental Single Cell Atlas of Gene RegulaTion and Expression	827
TF	Transcription Factor	828
ChEA3	ChIP-X Enrichment Analysis Version 3	829
CYP1A1	Cytochrome P450 family 1 subfamily A polypeptide 1	830
SMIM24	and Small Integral Membrane Protein 24	831
CREB3L3	cAMP responsive element-binding protein 3 like 3	832
BARX2	BarH-like homeobox 2	833
NAPSA	Napsin A Aspartic Peptidase	834
IRF9	Interferon regulatory factor 9	835

BATF2	Basic leucine zipper ATF-like transcription factor 2	836
F10	Coagulation Factor XThromboxane A Synthase 1	837
TBXAS1	Thromboxane A synthase 1	838
lncRNA	Long non-coding RNA	839
FOXN1	Forkhead box protein N1	840
IGFBP1	Insulin-like growth factor-binding protein 1	841
DKK1	Dickkopf-related protein 1	842
OSR1	Odd skipped-related 1	843
NR1I3	Constitutive androstane receptor	844
CYGB	Cytoglobin	845
SMC	Smooth muscle cell	846
EC	Endothelial cells	847
SP	Speckled protein	848
MEOX1	Mesenchyme homeobox 1	849
SNAI3	Snail family transcriptional repressor 3	850
NKX2-5	NK2 homeobox 5	851
SLC6A7	Solute carrier family 6 member 7	852
LSP1	Lymphocyte specific protein 1	853
EBF4	Early B cell factor 4	854
SOX9	SRY-box transcription factor 9	855
FUOM	Fucose mutarotase	856
ZNF	Zinc finder protein	857
4Rs	Reduction, Refinement, Replacement, Responsibility	858

Supporting information

S1 File. Supplementary File 1 contains additional graphs and tables as indicated in manuscript text.

Statements and Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The data used in this study were collected by the GENE-SWitCH (`https://www.gene-switch.eu/`) and are available through the FAANG data portal at `https://data.faang.org/projects/GENE-SWitCH`. The accession code is: (i) PRJEB41822 for RRBS; (ii) PRJEB41970 for RNA-Seq; and (iii) PREJEB70458 for ChIP-Seq. The results supporting the conclusions of this article are included in Supplementary File 1. The code and generated, pre-processed data are available through a GitHub repository at `https://github.com/wachalak/methylation-networks/`.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

KMW contributed to the study's design, processed and analyzed the data, interpreted the results, and drafted the manuscript. JdV supervised KMW, contributed to data generation, and initial data analysis. OM conceived the GENE-SWitCH methylation study and supervised JdV. MFLD supervised KMW, assisted with data analysis and manuscript editing. CFWP supervised KMW, contributed to the study's conceptualization, supported results interpretation and manuscript editing. All authors reviewed and approved the final manuscript.

Acknowledgments   885

Not applicable   886

Funding   887

Jani de Vos and Ole Madsen were funded through the GENE-SWitCH project, which   888
was financed by the European Union's Horizon 2020 Research and Innovation Program   889
under the grant agreement no 817998.   890

References

1. Swindle MM, Makin A, Herron AJ, Clubb FJ, Frazier KS. Swine as Models in
   Biomedical Research and Toxicology Testing. Vet Pathol. 2012;49(2):344–356.
   doi:10.1177/0300985811402846.

2. Helke KL, Ezell PC, Duran-Struuck R, Swindle MM. Biology and Diseases of
   Swine. In: Laboratory Animal Medicine. 3rd ed. Academic Press; 2015:695–769.
   doi:10.1016/B978-0-12-409527-4.00016-X.

3. Gutierrez K, Dicks N, Glanzer WG, Agellon LB, Bordigon V. Efficacy of the
   porcine species in biomedical research. Front Genet. 2015;6:293.
   doi:10.3389/fgene.2015.00293.

4. Prather RS. Pig genomics for biomedicine. Nat Biotechnol. 2013;31:122–123.
   doi:10.1038/nbt.2490.

5. Warr A, Affara N, Aken B, et al. An improved pig reference genome sequence to
   enable pig genetics and genomics research. GigaScience. 2020;9(6):1-14.
   doi:10.1093/gigascience/giaa051.

6. Lenoir G, Flatres-Grall L, Friggens NC, David I. Robustness scores in fattening
   pigs based on routinely collected phenotypes: determination and genetic
   parameters. J Anim Sci. 2022;100(5). doi:10.1093/jas/skac157.

7. McGlone JJ. The Future of Pork Production in the World: Towards Sustainable,
   Welfare-Positive Systems. Animals. 2013;3(2):401–415. doi:10.3390/ani3020401.

8. van der Werf J. Genomic Selection in Animal Breeding Programs. Methods Mol Biol. 2013;1019:543–561. doi:10.1007/978-1-62703-447-0_26.

9. Johnsson M. Genomics in animal breeding from the perspectives of matrices and molecules. Hereditas. 2023;160(20). doi:10.1186/s41065-023-00285-w.

10. Triantaphyllopoulos KA, Ikonomopoulos I, Bannister AJ. Epigenetics and inheritance of phenotype variation in livestock. Epigenetics Chromatin. 2016;9(31). doi:10.1186/s13072-016-0081-5.

11. Goldberg AD, Allis CD, Bernstein E. Epigenetics: A Landscape Takes Shape. Cell. 2007;128:635–638. doi:10.1016/j.cell.2007.02.006.

12. Champroux A, Cocquet J, Henry-Berger J, Drevet JR, Kocer A. A Decade of Exploring the Mammalian Sperm Epigenome: Paternal Epigenetic and Transgenerational Inheritance. Front Cell Dev Biol. 2018;6(50). doi:10.3389/fcell.2018.00050.

13. Villicaña S, Bell JT. Genetic impacts on DNA methylation: research findings and future perspectives. Genome Biol. 2021;22(127). doi:10.1186/s13059-021-02347-6.

14. Wang S, Wu W. DNA Methylation Alterations in Human Cancers. In: Tollefsbol TO, ed. Epigenetics in Human Disease. 2nd ed. Academic Press; 2018:109-139. doi:10.1016/B978-0-12-812215-0.00005-4.

15. Jin B, Li Y, Robertson KD. DNA Methylation: Superior or Subordinate in the Epigenetic Hierarchy? Genes Cancer. 2011;2(6):607–617. doi:10.1177/1947601910393957.

16. Ito T, Kubiura-Ichimaru M, Murakami Y, et al. DNMT1 regulates the timing of DNA methylation by DNMT3 in an enzymatic activity-dependent manner in mouse embryonic stem cells. PLoS One. 2022;17(1). doi:10.1371/journal.pone.0262277.

17. Li E, Zhang Y. DNA methylation in mammals. Cold Spring Harb Perspect Biol. 2014;6(5). doi:10.1101/cshperspect.a019133.

18. Deaton AM, Bird A. CpG islands and the regulation of transcription. Genes Dev. 2011;25(10):1010–1022. doi:10.1101/gad.2037511.

19. Lim WJ, Kim KH, Kim JY, Jeong S, Kim N. Identification of DNA-Methylated CpG Islands Associated With Gene Silencing in the Adult Body Tissues of the Ogye Chicken Using RNA-Seq and Reduced Representation Bisulfite Sequencing. Front Genet. 2019;10(346). doi:10.3389/fgene.2019.00346.

20. Le NQK, Yapp EKY, Nagasundaram N, Yeh HY. Classifying Promoters by Interpreting the Hidden Information of DNA Sequences via Deep Learning and Combination of Continuous FastText N-Grams. Front Bioeng Biotechnol. 2019;7(305). doi:10.3389/fbioe.2019.00305.

21. de Mendoza A, Nguyen TV, Ford E, et al. Large-scale manipulation of promoter DNA methylation reveals context-specific transcriptional responses and stability. Genome Biol. 2022;23(163). doi:10.1186/s13059-022-02728-5.

22. Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. Nat Rev Genet. 2012;13(7):484–492. doi:10.1038/nrg3230.

23. Giuffra E, Tuggle CK, FAANG Consortium. Functional Annotation of Animal Genomes (FAANG): Current Achievements and Roadmap. Annu Rev Anim Biosci. 2019;7:65–88. doi:10.1146/annurev-animal-020518-114913.

24. Acloque H, Harrison PW, Lakhal W, et al. Extensive functional genomics information from early developmental time points for pig. In: Proceedings of 12th World Congress on Genetics Applied to Livestock Production (WCGALP). 2022. doi:10.3920/978-90-8686-940-4_550.

25. Mirza B, Wang W, Wang J, et al. Machine Learning and Integrative Analysis of Biomedical Big Data. Genes. 2019;10(2):87. doi:10.3390/genes10020087.

26. Donoho DL. High-Dimensional Data Analysis: The Curses and Blessings of Dimensionality. AMS Math Challenges Lecture. 2000;1-32.

27. Phan JH, Quo CF, Cheng C, Wang MD. Multiscale Integration of -Omic, Imaging, and Clinical Data in Biomedical Informatics. IEEE Rev Biomed Eng. 2012;5:74–87. doi:10.1109/RBME.2012.2212427.

28. Ogutu JO, Schulz-Streeck T, Piepho HP. Genomic selection using regularized linear regression models: ridge regression, lasso, elastic net and their extensions. BMC Proc. 2012;6. doi:10.1186/1753-6561-6-S2-S10.

29. van Wieringen WN. Lecture notes on ridge regression. arXiv preprint. 2023. doi:10.48550/arXiv.1509.09169.

30. Bodein A, Scott-Boyer MP, Perin O, Lê Cao KA, Droit A. Interpretation of network-based integration from multi-omics longitudinal data. Nucleic Acids Res. 2022;50(5). doi:10.1093/nar/gkab1200.

31. Price BS, Geyer CJ, Rothman AJ. Ridge Fusion in Statistical Learning. arXiv preprint. 2014. doi:10.48550/arXiv.1310.3892.

32. Kalavacharla VK, Subramani M, Ayyappan V, Dworkin MC, Hayford RK. Plant Epigenomics. In: Tollefsbol TO, ed. Handbook of Epigenetics. 2nd ed. Academic Press; 2017:245-258. doi:10.1016/B978-0-12-805388-1.00016-X.

33. Miyata K, Naito M, Miyata T, Mokuda S, Asahara H. Bisulfite Sequencing for DNA Methylation Analysis of Primary Muscle Stem Cells. Methods Mol Biol. 2017;1668:3–13. doi:10.1007/978-1-4939-7283-8_1.

34. Li Y, Tollefsbol TO. DNA methylation detection: bisulfite genomic sequencing analysis. Methods Mol Biol. 2011;11–21. doi:10.1007/978-1-61779-316-5_2.

35. Smith ZD, Gu H, Bock C, Gnirke A, Meissner A. High-throughput bisulfite sequencing in mammalian genomes. Methods. 2009;48(3):226–232. doi:10.1016/j.ymeth.2009.05.003.

36. Nakato R, Sakata T. Methods for ChIP-seq analysis: A practical workflow and advanced applications. Methods. 2021;187:44–53. doi:10.1016/j.ymeth.2020.03.005.

37. Xin D, Hu L, Kong X. Alternative Promoters Influence Alternative Splicing at the Genomic Level. PLoS One. 2008;3(6). doi:10.1371/journal.pone.0002377.

38. Iñiguez G, Battiston F, Karsai M. Bridging the gap between graphs and networks. Commun Phys. 2020;3(8). doi:10.1038/s42005-020-0359-6.

39. Hastie T, Tibshirani R, Friedman J. Undirected Graphical Models. In: The Elements of Statistical Learning. Springer New York, NY; 2009:625–648. doi:10.1007/978-0-387-84858-7_17.

40. Altenbuchinger M, Weihs A, Quackenbush J, Grabe HJ, Zacharias HU. Gaussian and Mixed Graphical Models as (multi-)omics data analysis tools. Biochim Biophys Acta Gene Regul Mech. 2020;1863(6). doi:10.1016/j.bbagrm.2019.194418.

41. Peeters CFW, Bilgrau AE, van Wieringen WN. rags2ridges: A One-Stop-l2-Shop for Graphical Modeling of High-Dimensional Precision Matrices. J Stat Softw. 2022;102(4):1–32. doi:10.18637/jss.v102.i04.

42. Cai Z, Li R, Zhang Y. A Distribution Free Conditional Independence Test with Applications to Causal Discovery. J Mach Learn Res. 2022;23(85):1-41.

43. Baba K, Shibata R, Sibuya M. Partial correlation and conditional correlation as measures of conditional independence. Aust N Z J Stat. 2004;46(4):657–664. doi:10.1111/j.1467-842X.2004.00360.x.

44. Li T, Qian C, Levina E, Zhu J. High-dimensional Gaussian graphical models on network-linked data. J Mach Learn Res. 2020;21:1-45.

45. Li X, Fu Y, Wang X, Qiu W. Robust joint score tests in the application of DNA methylation data analysis. BMC Bioinformatics. 2018;19(174). doi:10.1186/s12859-018-2185-3.

46. Weinhold L, Wahl S, Pechlivanis S, Hoffmann P, Schmid M. A statistical model for the analysis of beta values in DNA methylation studies. BMC Bioinformatics. 2016;17(480). doi:10.1186/s12859-016-1347-4.

47. Epskamp S, Fried EI. A Tutorial on Regularized Partial Correlation Networks. Psychol Methods. 2018;23(4):617–634. doi:10.1037/met0000167.

48. Czado C, Nagler T. Vine Copula Based Modeling. Annu Rev Stat Its Appl. 2022;9:453–477. doi:10.1146/annurev-statistics-040220-101153.

49. Liu H, Lafferty J, Wasserman L. The Nonparanormal: Semiparametric Estimation of High Dimensional Undirected Graphs. J Mach Learn Res. 2009;10:2295–2328.

50. Jiang H, Fei X, Liu H, et al. huge: High-Dimensional Undirected Graph Estimation. R package. 2021. URL: https://cran.r-project.org/web/packages/huge.

51. Zhao Y, Li MC, Konaté MM, et al. TPM, FPKM, or Normalized Counts? A Comparative Study of Quantification Measures for the Analysis of RNA-seq Data from the NCI Patient-Derived Models Repository. J Transl Med. 2021;19(269). doi:10.1186/s12967-021-02936-w.

52. Batool F, Henning C. Clustering with the Average Silhouette Width. Comput Stat Data Anal. 2021;158. doi:10.1016/j.csda.2021.107190.

53. Rousseeuw PJ. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. J Comput Appl Math. 1987;20:53–65. doi:10.1016/0377-0427(87)90125-7.

54. Bilgrau AE, Peeters CFW, Eriksen PS, Bøgsted M, van Wieringen WN. Targeted Fused Ridge Estimation of Inverse Covariance Matrices from Multiple High-Dimensional Data Classes. J Mach Learn Res. 2020;1–52. URL: http://jmlr.org/papers/v21/15-509.html.

55. Tsai K, Koyejo O, Kolar M. Joint Gaussian graphical model estimation: A survey. WIREs Comput Stat. 2022;14(6). doi:10.1002/wics.1582.

56. Nelder JA, Mead R. A Simplex Method for Function Minimization. Comput J. 1965;7(4):308–313. doi:10.1093/comjnl/7.4.308.

57. Fruchterman TMJ, Reingold EM. Graph drawing by force-directed placement. J Softw Pract Exp. 1991;21(11):1129–1164. doi:10.1002/spe.4380211102.

58. Csárdi G, Nepusz T. The igraph software package for complex network research. InterJournal Complex Syst. 2006;1695. URL: https://igraph.org.

59. Csárdi G, Nepusz T, Traag V, Horvát S, Zanini F, Noom D, Müller K. igraph: Network Analysis and Visualization in R. R package version 2.0.2. 2024. doi:10.5281/zenodo.7682609. URL: https://CRAN.R-project.org/package=igraph.

60. Radicchi F, Castellano C, Cecconi F, Loreto V, Parisi D. Defining and identifying communities in networks. Proc Natl Acad Sci U S A. 2004;101(9):2658–2663. doi:10.1073/pnas.0400054101.

61. Parés F, Gasulla DG, Vilalta A, et al. Fluid Communities: A Competitive, Scalable and Diverse Community Detection Algorithm. In: Cherifi C, Cherifi H, Karsai M, Musolesi M, eds. Complex Networks & Their Applications VI. Springer International Publishing; 2018:229-240. ISBN: 978-3-319-72150-7.

62. Kolberg L, Raudvere U, Kuzmin I, et al. g:Profiler—interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). Nucleic Acids Res. 2023;51(W1):W207–W212. doi:10.1093/nar/gkad347.

63. Chen EY, Tan CM, Kou Y, et al. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics. 2013;14(128). doi:10.1186/1471-2105-14-128.

64. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res. 2016;44(W1):W90-W97. doi:10.1093/nar/gkw377.

65. Xie Z, Bailey A, Kuleshov MV, et al. Gene set knowledge discovery with Enrichr. Curr Protoc. 2021;1(3). doi:10.1002/cpz1.90.

66. Cao J, O'Day DR, Pliner HA, et al. A human cell atlas of fetal gene expression. Science. 2020;370(6518). doi:10.1126/science.aba7721.

67. Keenan AB, Torre D, Lachmann A, et al. ChEA3: transcription factor enrichment analysis by orthogonal omics integration. Nucleic Acids Res. 2019;47(W1):W212–W224. doi:10.1093/nar/gkz446.

68. Wang G, Wang F, Huang Q, et al. Understanding Transcription Factor Regulation by Integrating Gene Expression and DNase I Hypersensitive Sites. Biomed Res Int. 2015. doi:10.1155/2015/757530.

69. Héberlé É, Bardet AF. Sensitivity of transcription factors to DNA methylation. Essays Biochem. 2019;63(6):727–741. doi:10.1042/EBC20190033.

70. Stejskalova L, Pavek P. The function of cytochrome P450 1A1 enzyme (CYP1A1) and aryl hydrocarbon receptor (AhR) in the placenta. Curr Pharm Biotechnol. 2011;12(5):715–730. doi:10.2174/138920111795470994.

71. Goh I, Botting RA, Rose A, Webb S, Engelbert J, Gitton Y, Stephenson E, Quiroga Londoño M, Mather M, Mende N, Imaz-Rosshandler I, Yang L, Horsfall D, Basurto-Lozada D, Chipampe NJ, Rook V, Lee JTH, Ton ML, Keitley D, Mazin P, Vijayabaskar MS, Hannah R, Gambardella L, Green K, Ballereau S, Inoue M, Tuck E, Lorenzi V, Kwakwa K, Alsinet C, Olabi B, Miah M, Admane C, Popescu DM, Acres M, Dixon D, Ness T, Coulthard R, Lisgo S, Henderson DJ, Dann E, Suo C, Kinston SJ, Park JE, Polanski K, Marioni J, van Dongen S, Meyer KB, de Bruijn M, Palis J, Behjati S, Laurenti E, Wilson NK, Vento-Tormo R, Chédotal A, Bayraktar O, Roberts I, Jardine L, Göttgens B, Teichmann SA, Haniffa M. Yolk sac cell atlas reveals multiorgan functions during human early development. Science. 2023;381(6659). doi:10.1126/science.add7564.

72. McCracken KW, Wells JM. Molecular pathways controlling pancreas induction. Semin Cell Dev Biol. 2012;23(6):656–662. doi:10.1016/j.semcdb.2012.06.009.

73. Gale RP. Development of the immune system in human fetal liver. Thymus. 1987;10(1-2):45–56. doi:10.1007/978-94-009-3365-1_6.

74. Miller D, Motomura K, Garcia-Flores V, Romero R, Gomez-Lopez N. Innate lymphoid cells in the maternal and fetal compartments. Front Immunol. 2018;9. doi:10.3389/fimmu.2018.02396.

75. Nakagawa Y, Wang Y, Han SI, Okuda K, Oishi A, Yagishita Y, Kumagai K, Ohno H, Osaki Y, Mizunoe Y, Araki M, Murayama Y, Iwasaki H, Konishi M, Itoh N, Matsuzaka T, Sone H, Yamada N, Shimano H. Enterohepatic transcription factor CREB3L3 protects atherosclerosis via SREBP competitive inhibition. Cell Mol Gastroenterol Hepatol. 2021;11(4):949–971. doi:10.1016/j.jcmgh.2020.11.004.

76. Ma J, Xia LL, Yao XQ, Zheng SM, Li S, Xu LS, Sha WH, Li ZS. BARX2 expression is downregulated by CpG island hypermethylation and is associated with suppressed cell proliferation and invasion of gastric cancer cells. Oncol Rep. 2020;43(6):1805–1818. doi:10.3892/or.2020.7558.

77. Naka T, Yokose S. Immunohistochemical localization of BARX2 in the developing fetal mouse submandibular glands. Acta Histochem Cytochem. 2009;42(2):47–53. doi:10.1267/ahc.08027.

78. Edgar R, Mazor Y, Rinon A, Blumenthal J, Golan Y, Buzhor E, Livnat I, Ben-Ari S, Lieder I, Shitrit A, Gilboa Y, Ben-Yehudah A, Edri O, Shraga N, Bogoch Y, Leshansky L, Aharoni S, West MD, Warshawsky D, Shtrichman R. LifeMap Discovery™: The Embryonic Development, Stem Cells, and Regenerative Medicine Research Portal. PLoS One. 2013;8(7):1-13. doi:10.1371/journal.pone.0066629.

79. Nicolaides NC, Willenberg HS, Bornstein SR. Adrenal Cortex: Embryonic Development, Anatomy, Histology and Physiology. In: Feingold KR, Anawalt B, Blackman MR, eds. Endotext [Internet]. South Dartmouth (MA): MDText.com, Inc; 2000. Available from: https://www.ncbi.nlm.nih.gov/books/NBK278945/.

80. Farraj KL, Zeltser R. Embryology, Heart Tube. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/books/NBK499934/.

81. Paul A, Tang TH, Ng SK. Interferon Regulatory Factor 9 Structure and Regulation. Front Immunol. 2018;9. doi:10.3389/fimmu.2018.01831.

82. Le DT, Florez MA, Kus P, Tran BT, Kain B, Zhu Y, Christensen K, Jain A, Malovannaya A, King KY. BATF2 promotes HSC myeloid differentiation by amplifying IFN response mediators during chronic infection. iScience. 2023;26(2). doi:10.1016/j.isci.2023.106059.

83. Bethesda (MD): National Library of Medicine (US). F10 coagulation factor X. 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/gene/2159.

84. Bethesda (MD): National Library of Medicine (US). TBXAS1 thromboxane A synthase 1. 2024. Accessed: 2024-03-09. Available from: https://www.ncbi.nlm.nih.gov/gene/6916.

85. Cerase A, Pintacuda G, Tattermusch A, Avner P. Xist localization and function: new insights from multiple levels. Genome Biol. 2015;16(166). doi:10.1186/s13059-015-0733-y.

86. Lee J, Davidow LS, Warshawsky D. Tsix, a gene antisense to Xist at the X-inactivation centre. Nat Genet. 1999;21:400–404. doi:10.1038/7734.

87. Vigliano I, Gorrese M, Fusco A, Vitiello L, Amorosi S, Panico L, Ursini MV, Calcagno G, Racioppi L, Del Vecchio L, Pignata C. FOXN1 mutation abrogates prenatal T-cell development in humans. J Med Genet. 2011;48(6):413–416. doi:10.1136/jmg.2011.089532.

88. Reis MD, Csomos K, Dias LP, Prodan Z, Szerafin T, Savino W, Takacs L. Decline of FOXN1 gene expression in human thymus correlates with age: possible epigenetic regulation. Immun Ageing. 2015;12:18. doi:10.1186/s12979-015-0045-9.

89. Moses A, Bhalla P, Thompson A, Lai L, Coskun FS, Seroogy CM, de la Morena MT, Wysocki CA, van Oers NSC. Comprehensive phenotypic analysis of diverse FOXN1 variants. J Allergy Clin Immunol. 2023;152(5):1273–1291. doi:10.1016/j.jaci.2023.06.019.

90. Wang Q, Li J, Wang S, Deng Q, An Y, Xing Y, Dai X, Li Z, Ma Q, Wang K, Liu C, Yuan Y, Dong G, Zhang T, Yang H, Du Y, Hou Y, Ke W, Shang Z. Single-cell transcriptional profiling reveals cellular and molecular divergence in human maternal-fetal interface. Sci Rep. 2022;12(10892). doi:10.1038/s41598-022-14516-z.

91. Farine T, Parsons M, Lye S, Shynlova O. Isolation of Primary Human Decidual Cells from the Fetal Membranes of Term Placentae. J Vis Exp. 2018. doi:10.3791/57443.

92. Vallecillo-García P, Orgeur M, Vom Hofe-Schneider S, Stumm J, Kappert V, Ibrahim DM, Börno ST, Hayashi S, Relaix F, Hildebrandt K, Sengle G, Koch M, Timmermann B, Marazzi G, Sassoon DA, Duprez D, Stricker S. Odd skipped-related 1 identifies a population of embryonic fibro-adipogenic progenitors regulating myogenesis during limb development. Nat Commun. 2017;8. doi:10.1038/s41467-017-01120-3.

93. Chen F, Zamule SM, Coslo DM, Chen T, Omiecinski CJ. The human constitutive androstane receptor promotes the differentiation and maturation of hepatic-like cells. Dev Biol. 2013;384(2):155–165. doi:10.1016/j.ydbio.2013.10.012.

94. Bethesda (MD): National Library of Medicine (US). Cytoglobin. 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/gene/114757.

95. Sufleţel RT, Melincovici CS, Gheban BA, Toader Z, Mihu CM. Hepatic stellate cells - from past till present: morphology, human markers, human cell lines, behavior in normal and liver pathology. Rom J Morphol Embryol. 2020;61(3):615–642. doi:10.47162/RJME.61.3.01.

96. Nishi H, Inagi R, Kawada N, Yoshizato K, Mimura I, Fujita T, Nangaku M. Cytoglobin, a novel member of the globin family, protects kidney fibroblasts against oxidative stress under ischemic conditions. Am J Pathol. 2011;178(1):128–139. doi:10.1016/j.ajpath.2010.11.011.

97. Siddarth M, Datta SK, Ahmed RS, Banerjee BD, Kalra OP, Tripathi AK. Association of CYP1A1 gene polymorphism with chronic kidney disease: a case control study. Environ Toxicol Pharmacol. 2013;36(1):164–170. doi:10.1016/j.etap.2013.03.008.

98. Huang B, Bao J, Cao YR, Gao HF, Jin Y. Cytochrome P450 1A1 (CYP1A1) Catalyzes Lipid Peroxidation of Oleic Acid-Induced HepG2 Cells. Biochemistry (Mosc). 2018;83(5):595–602. doi:10.1134/S0006297918050127.

99. Ferretti E, Hadjantonakis AK. Mesoderm specification and diversification: from single cells to emergent tissues. Curr Opin Cell Biol. 2019;61:110–116. doi:10.1016/j.ceb.2019.07.012.

100. Donadon M, Santoro MM. The origin and mechanisms of smooth muscle cell development in vertebrates. Development. 2021;148(7). doi:10.1242/dev.197384.

101. Nave KA, Asadollahi E, Sasmita A. Expanding the function of oligodendrocytes to brain energy metabolism. Curr Opin Neurobiol. 2023;83. doi:10.1016/j.conb.2023.102782.

102. Lepore E, Casola I, Dobrowolny G, Musarò A. Neuromuscular Junction as an Entity of Nerve-Muscle Communication. Cells. 2019;8(8). doi:10.3390/cells8080906.

103. de Vellis J, Carpenter E. General Development of the Nervous System. In: Basic Neurochemistry: Molecular, Cellular and Medical Aspects. Lippincott-Raven; 1999. Available from: https://www.ncbi.nlm.nih.gov/books/NBK28253/.

104. Krüger-Genge A, Blocki A, Franke RP, Jung F. Vascular Endothelial Cell Biology: An Update. Int J Mol Sci. 2019;20(18). doi:10.3390/ijms20184411.

105. Trimm E, Red-Horse K. Vascular endothelial cell development and diversity. Nat Rev Cardiol. 2023;20:197-210. doi:10.1038/s41569-022-00770-1.

106. Karaky M, Fedetz M, Potenciano V, Andrés-León E, Codina AE, Barrionuevo C, Alcina A, Matesanz F. SP140 regulates the expression of immune-related genes associated with multiple sclerosis and other autoimmune diseases by NF-κB inhibition. Hum Mol Genet. 2018;27:4012–4023. doi:10.1093/hmg/ddy284.

107. Dong K, Guo X, Chen W, Hsu AC, Shao Q, Chen JF, Chen SY. Mesenchyme homeobox 1 mediates transforming growth factor-β (TGF-β)-induced smooth muscle cell differentiation from mouse mesenchymal progenitors. J Biol Chem. 2018;293(22):8712–8719. doi:10.1074/jbc.RA118.002350.

108. Rip R, George T, Shumway KR, Sattar Y. Physiology, Cardiac Muscle. In: StatPearls [Internet]. 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/books/NBK572070/.

109. Davenport P, Liu ZJ, Sola-Visner M. Fetal vs adult megakaryopoiesis. Blood. 2022;139(22):3233–3244. doi:10.1182/blood.2020009301.

110. Ali RA, Wuescher LM, Worth RG. Platelets: essential components of the immune system. Curr Trends Immunol. 2015;16:65–78.

111. Koupenova M, Livada AC, Morrell CN. Platelet and Megakaryocyte Roles in Innate and Adaptive Immunity. Circ Res. 2022;130(2):288–308. doi:10.1161/CIRCRESAHA.121.319821.

112. Bethesda (MD): National Library of Medicine (US). Snail family transcriptional repressor 3. 2024. Accessed: 2024-03-09. Available from: https://www.ncbi.nlm.nih.gov/gene/333929.

113. Bethesda (MD): National Library of Medicine (US). NK2 homeobox 5. 2024. Accessed: 2024-03-09. Available from: https://www.ncbi.nlm.nih.gov/gene/1482.

114. Bethesda (MD): National Library of Medicine (US). Solute carrier family 6 member 7. 2024. Accessed: 2024-03-09. Available from: https://www.ncbi.nlm.nih.gov/gene/6534.

115. Elshazzly M, Lopez MJ, Reddy V, Caban O. Embryology, Central Nervous System. In: StatPearls [Internet]. 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/books/NBK526024/.

116. Bethesda (MD): National Library of Medicine (US). Lymphocyte specific protein 1. 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/gene/4046.

117. Kubo S, Kataria R, Yao Y, et al. Early B cell factor 4 modulates FAS-mediated apoptosis and promotes cytotoxic function in human immune cells. Proc Natl Acad Sci U S A. 2022;119(23). doi:10.1073/pnas.2208522119.

118. Jo A, Denduluri S, Zhang B, et al. The versatile functions of Sox9 in development, stem cells, and human diseases. Genes Dis. 2014;1(2):149-161. doi:10.1016/j.gendis.2014.09.004.

119. Jakob S, Lovell-Badge R. Sex determination and the control of Sox9 expression in mammals. FEBS J. 2011;278(7):1002–1009. doi:10.1111/j.1742-4658.2011.08029.x.

120. Bethesda (MD): National Library of Medicine (US). Fucose mutarotase. 2024. Accessed: 2024-03-08. Available from: https://www.ncbi.nlm.nih.gov/gene/282969.

121. Jen J, Wang YC. Zinc finger proteins in cancer progression. J Biomed Sci. 2016;23(53). doi:10.1186/s12929-016-0269-9.

122. Sayers EW, Bolton EE, Brister JR, et al. Database resources of the national center for biotechnology information. Nucleic Acids Res. 2022;50(D1):D20-D26. doi:10.1093/nar/gkab1112.

123. Jara EL, Muñoz-Durango N, Llanos C, et al. Modulating the function of the immune system by thyroid hormones and thyrotropin. Immunol Lett. 2017;184:76–83. doi:10.1016/j.imlet.2017.02.010.

124. Lu-Culligan A, Iwasaki A. The Role of Immune Factors in Shaping Fetal Neurodevelopment. Annu Rev Cell Dev Biol. 2020;36:441–468. doi:10.1146/annurev-cellbio-021120-033518.

125. Hussain T, Murtaza G, Kalhoro DH, et al. Understanding the Immune System in Fetal Protection and Maternal Infections during Pregnancy. J Immunol Res. 2022. doi:10.1155/2022/7567708.

126. Stelzer IA, Urbschat C, Schepanski S, et al. Vertically transferred maternal immune cells promote neonatal immunity against early life infections. Nat Commun. 2021;12. doi:10.1038/s41467-021-24719-z.

127. Janeway CA Jr, Travers P, Walport M, Shlomchik MJ. Immunobiology: The Immune System in Health and Disease. 5th ed. Garland Science; 2001. URL: https://www.ncbi.nlm.nih.gov/books/NBK27092/.

128. Remien K, Majmundar SH. Physiology, Fetal Circulation. StatPearls [Internet]. 2024. URL: https://www.ncbi.nlm.nih.gov/books/NBK539710/.

129. Kiani AK, Pheby D, Henehan G, et al. Ethical considerations regarding animal experimentation. J Prev Med Hyg. 2022;63(2 Suppl 3):E255-E266. doi:10.15167/2421-4248/jpmh2022.63.2S3.2768.

130. Zhao Z, Chen X, Dowbaj AM, et al. Organoids. Nat Rev Methods Primers. 2022;2(94). doi:10.1038/s43586-022-00174-y.

131. de Leeuw FA, Peeters CFW, Kester MI, et al. Blood-based metabolic signatures in Alzheimer's disease. Alzheimers Dement. 2017;8:196–207. doi:10.1016/j.dadm.2017.07.006.

132. Hayashi N, Yamaguchi S, Rodenburg F, et al. Multiple biomarkers of sepsis identified by novel time-lapse proteomics of patient serum. PLoS One. 2019;14(9). doi:10.1371/journal.pone.0222403.

133. Peeters CFW, van de Wiel MA, van Wieringen WN. The spectral condition number plot for regularization parameter evaluation. Comput Stat. 2020;35:629–646. doi:10.1007/s00180-019-00912-z.