Eurasian Spoonbill chicks receive parental care up to several months after fledging, but not into migration

Tamar Lok[1,2,3*], Petra de Goeij[4], Eldar Rakhimberdiev[5], Theunis Piersma[1,3,4] & Wouter Vansteelant[1,4,5]

[1] Rudi Drent Chair in Global Flyway Ecology, Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Nijenborgh 9, 9747 AG Groningen, Netherlands

[2] EcoMoves, Wilhelminalaan 78, 1791 AP Den Burg, The Netherlands

[3] Department of Coastal Systems, NIOZ Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB, Den Burg, Texel, The Netherlands

[4] BirdEyes, Centre for Global Ecological Change at the Faculties of Science & Engineering and Campus Fryslân, University of Groningen, Wirdumerdijk 34, 8911 CE, Leeuwarden, The Netherlands

[5] Department of Theoretical and Computational Ecology, Institute for Biodiversity and Ecosystems Dynamics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands

* Corresponding author, tamarlok@gmail.com

20 **Abstract**

Despite its potential role in affecting survival, habitat use and migration strategies of juvenile

birds, post-fledging parental care is poorly studied, as it requires that families can be followed

over large distances. Here we combine visual observations of colour-ringed chicks being fed by

a parent after fledging with GPS-tracking and accelerometer-based behavioural classification

of fledged chicks and their parent(s) to quantify post-fledging parental care in Eurasian

spoonbills and to test whether and how this is affected by age of the chick and sex of the chick

or parent. We show that the number of observed feedings and the amount of overall contact

between chicks and parents strongly decreased with chick age. Chicks were observed being fed

until 125 d old, always within 40 km of the natal colony. The amount of contact strongly varied

among the 16 GPS-tagged chick-parent pairs, on average decreasing from 11% of contact time

at 40 d old to <3% at 90 d. The last contact occurred at chick ages of 44-136 d (median: 88 d).

All contact occurred within 18 km of the natal colony except for the first outbound migratory

flight of one chick-parent pair. Both mothers and fathers engaged in post-fledging care, with

some evidence that mothers had slightly more contact with chicks than fathers, overall as well

as while the chick was begging or foraging. In 10 out of 11 cases that both chick and parent

embarked on autumn migration, they did not migrate together: post-fledging care ended on

average four weeks before the chick's departure on autumn migration.

38 **Keywords**

39 Animal tracking, biologging, development, Eurasian spoonbill, migration, parental care,

40 *Platalea leucorodia*, post-fledging, ontogeny

41 **Running title**

42 Post-fledging parental care in spoonbills

Introduction

Studies on parental care in birds have strongly focussed on the period when birds are bound to the nest (nest building, egg laying, incubation and feeding of chicks), likely due to the difficulty of following families after fledging, when they often travel large distances. After fledging, however, chicks must acquire foraging skills (Mellink, Castillo-Guerrero, & Peñaloza-Padilla, 2014), learn where to find suitable sites and habitats for foraging and resting, how to interact with conspecifics and other species, how to avoid predators and, in case of migratory species, acquire suitable migration routines. Not surprisingly, perhaps, young birds often experience considerable mortality shortly after fledging (Naef-Daenzer & Grüebler, 2016). Extended parental care during this period has been shown for a variety of species and has likely evolved to reduce this post-fledging mortality by (1) continued feeding of the chicks while they improve their own foraging skills (Mellink et al., 2014), (2) guiding them to suitable sites and habitats for foraging and resting, (3) protecting them from conspecifics, other species and predators and/or (4) guiding them on their first migration, i.e., teaching them suitable migration routines (Chetverikova, Babushkina, Galkina, Shokhrin, & Bojarinova, 2017). An experimental study on barn swallows (*Hirundo rustica*) confirmed that extended post-fledging parental care indeed increased post-fledging survival (Grüebler & Naef-Daenzer, 2010). On the other hand, providing longer post-fledging parental care may go at the expense of future fecundity of the parents (López-Idiáquez, Vergara, Fargallo, & Martínez-Padilla, 2018), thereby creating a parent-offspring conflict (Rehling et al., 2012; Riou, Chastel, & Hamer, 2012).

While in some species, post-fledging parental care lasts for only a few days, other species show much longer periods of post-fledging parental care, and family associations may even be maintained during and after the first outbound migration (Cramp, 1994) . While migrating in family groups was long thought to be restricted to Anseriformes (geese, swans; e.g. (Black & Owen, 1989; Earnst & Bart, 1991; Scott, 1980)), Gruiformes (cranes; e.g. (Alonso, Bautista, & Alonso, 2004)) and some species of terns (Ashmole & Tovar, 1968; Byholm, Beal, Isaksson, Lötberg, & Åkesson, 2022), recent studies revealed that it also occurs in a variety of other species, including great bustard (*Otis tarda*) (Palacín, Alonso, Alonso, Magaña, & Martín, 2011), and long-tailed tits (*Aegithalos caudatus*) (Chetverikova et al., 2017). In the case of terns, parents also feed their chicks well into the migration, and even as far as the non-breeding grounds (Ashmole & Tovar, 1968; Fijn & Kemper, 2023). At some point, parents may no longer have a caring role, but family bonds may nonetheless be maintained when the young may help with defending a (breeding or food) territory, warning for potential danger (Black & Owen, 1989), or raising genetically related individuals (i.e., cooperative breeding, Koenig & Dickinson, 2004).

The duration and type of post-fledging parental care (i.e., feeding at nesting site or guiding to

suitable sites for foraging and resting) will determine the extent to which juveniles learn suitable

habitats and/or migration routes from their own parent(s) or discover these through

independent exploration or social interactions with other conspecifics. This will influence the

social and genetic structure in a population as well as the capacity of populations to anticipate

to changing conditions. However, despite its presumed importance, functional role and

population-level consequences, only few studies have quantified the duration of post-fledging

parental care, as it is difficult to follow families, particularly when they move out of the territory

and disperse over large distances (Meyburg, Meyburg, Mizera, Maciorowski, & Kowalski, 2005).

Here, we investigate post-fledging parental care in Eurasian spoonbills (*Platalea*

leucorodia leucorodia). Eurasian spoonbills are colony-breeding birds that show obligate

biparental care during egg incubation and chick-rearing prior to fledging (Lok, van der Geest, de

Goeij, Rakhimberdiev, & Piersma, 2024). A recent study in a freshwater population in Hungary

showed that after fledging, at an age of *c.* 40 d old, juveniles were still fed by a (presumed)

parent at distances up until c. 150 km from the natal colony and at an age of 113 days after

hatching, with the last chick feedings observed in early October (Pigniczki, 2022). This is

considerably longer than what has so far been reported for closely related species (Cramp,

1994), which Pigniczki (2022) suggests to be caused by the long time it takes for young

spoonbills to master the technique of tactile foraging on often mobile prey. In contrast, the

most extreme examples of extended care involve species with rather basic foraging techniques

(e.g. grazing geese). As such, the observations from Hungary raise questions about how

common it is for spoonbills to extend care for several months after fledging, possibly extending

into (part of) the migratory period. If common, such care would have important implications for

the development of migratory routines of juvenile birds (Whiten, 2019). Spoonbills are known to

form post-breeding congregations to prepare for migration (de Boer, Vansteelant, & Piersma,

2024; Henriques, Piersma, Vansteelant, de Goeij, & Lok, in press) and to migrate in often mixed-

age flocks (Lagarde & Piersma, 2021). However, it remains to be confirmed whether (and how

often) parents and young migrate together in these flocks, and hence, whether juveniles might

learn routes, stop-overs and migratory destinations from their parents.

We combine GPS-tracking and acceleration-based behavioural classification of parents

and their young with visual observations of feedings of colour-ringed chicks (of known age and

origin) to quantify until what age and what distance from the colony spoonbill chicks receive

post-fledging parental care (in the form of being fed or being guided to post-breeding sites or on

migration), and whether this depends on the sex of the chick or the parent. As males are the

larger sex in spoonbills, male chicks require more food during growth (Krijgsveld, Dijkstra,

Visser, & Daan, 1998) and may therefore be more dependent on still receiving food from their

parents after fledging compared to female chicks. In addition, fathers and mothers may differ in

the amount of post-fledging parental care they provide (Byholm et al., 2022; Ersoy, Maag,

Boehly, Boucherie, & Bugnyar, 2021; Palacín et al., 2011), which could be related to the amount

of energy already spent on parental care before fledging, or sex-specific parental care roles that

may have evolved.

Methods

Study population

For this study, we use observations of colour-ringed Eurasian Spoonbill chicks from colonies

throughout The Netherlands between 1999 and 2023 (Photo 1) and GPS and acceleration data

of chicks and their parent(s) from the colony on Schiermonnikoog, The Netherlands (53°29'N,

6°15'E) between 2016 and 2018.

Feeding observations

125   To retrieve visual observations of feedings of colour-ringed chicks, we selected

resightings from the colour-ringing database of the Werkgroep Lepelaar that had feeding-

related English or Dutch words in the remarks (i.e., "feed", "fed", "voer") or where feeding was

entered as a behaviour. We then further checked the remarks on whether any information was

provided about the ring code of the parent (or its chick). This resulted in a dataset of 184 feeding

observations of 127 colour-ringed chicks born in The Netherlands between 1999 and 2024,

involving 77 colour-ringed parents and 33 feedings by unringed parents.

132   The sex of a bird was either determined molecularly (see below) or visually. For the birds

that were not molecularly sexed, we used the sex identification from visual observations to

assess a bird's (likely) sex. When a bird was visually sexed, it was often noted whether the bird

(1) was seen copulating, (2) was seen with partner with clear difference, (3) was seen with

partner with unclear difference, (4) was seen alone, (5) has a partner of which sex was

molecularly determined, (6) stopped or started incubating at the expected time for a certain sex

(starting in the morning or stopping in the evening = male), (7) incubated at night (based on

camera trap data) = female. For situations 1, 2, 5 or 7, the identified sex was considered to be

certain, all other visual sex identifications were considered uncertain. For each bird, we then

calculated the number of certain and uncertain female and male identifications. If a bird had

only certain identifications of a single sex, this was assumed to be the bird's sex. In all other

cases, when the total number of female identifications (certain and uncertain identifications

combined) differed by more than two from the total number of male identifications, the most

often identified sex was assumed to be the bird's sex. For 48 of the 54 birds (89%) for which sex

was determined both visually and molecularly, the sex was correctly identified based on visual

observations using the above-described procedure. The age of the chick was calculated from

the length of the 8[th] primary (P8) at colour-ringing, with the Gompertz-curve formula estimated

by Lok *et al.* (2014): age (days) = -ln(–ln($P8$/247))/0.095 + 19.3.

GPS tracking

Between 2012 and 2018, 31 adult spoonbills and between 2016 and 2018, 15 juvenile

spoonbills were equipped with a UvA-BiTS GPS/ACC tracker (Bouten, Baaij, Shamoun-Baranes,

& Camphuysen, 2013). The adult spoonbills were caught on the nest during the late egg

incubation stage (for more details, see Lok et al., 2024), while the juveniles were caught shortly

before they were able to fly. Upon capture, the birds were ringed with a unique combination of

colour-rings and a flag and measured. For molecular sex determination (Fridolfsson & Ellegren,

1999), a blood sample of 10-80 µl was collected from the brachial vein and stored in 96%

alcohol . The GPS-tracker was mounted on the back of the bird using a non-flexible Teflon

harness. The tracker including the neoprene pad and Teflon harness (5 g) weighed 31 ± 12 g

(mean ± SD) for the adults and 22 ± 0.9 g for the juveniles. With average body mass being 1646 ±

146 g and 1968 ± 162 g for the tagged adult female and male spoonbills, and 1371 ± 82 g and

1595 ± 133 g for the juvenile female and male spoonbills, the trackers contributed 0.9-2.8% to

body mass, thus not exceeding the 3% guideline (Phillips, Xavier, & Croxall, 2003). Data of

tagged birds used in this paper are shown in Table S1.

We selected juveniles for GPS-tagging that had at least one GPS-tagged parent. To this

aim, the chicks received a uniquely numbered temporary band of flexible plastic around each

leg when they were between 5 and 12 days old and still attached to their own nest. During this

visit, head-bill length was measured to estimate age (Lok et al., 2014). We then recaptured the

chicks when they were 30-32 days old and almost able to fly (at c. 35 days old) and equipped

them with a GPS tracker. As the chicks still grow somewhat bigger after fledging (they weighed

on average 83% of the adult sex-specific body mass), we left a bit more space between the

body and the harness than we did for the adults (i.e. *c.* 2 cm instead of 1 cm space between the

tracker and the backbone of the bird). Of the 15 juvenile spoonbills that were GPS-tagged, two

died prior to departure on migration and another 5 died later during their first year (Table S1).

This resulted in a first-year survival rate of 0.53 ± 0.13 (mean ± SE), which is not significantly

different from the first-year survival rate estimated from mark-recapture data and averaged over

the period 1988-2008 (0.56 ± 0.04; Lok, Overdijk, Tinbergen, & Piersma, 2013; Z=-0.20, p=0.84).

178   For this study, the trackers were set to collect a GPS location every 10 minutes along with

a 1.6 s sample of 20 Hz acceleration data. When the birds were within the UvA-BiTS antenna

network, all data were downloaded. Between 2012 and 2015, birds were equipped with GPS-

trackers whose data could only be downloaded via this antenna network. However, from 2016

onward, we used GPS-GSM trackers that were able to download a small part of the GPS

samples via the GSM-network when birds were out of range of the antenna network. This

allowed us to also track the birds during their southward migration, albeit at low resolution, as

trackers were set to transmit three GPS locations collected during the three hours prior to a

GSM session that occurred at 6-hour intervals, provided that the tracker was able to connect to

a GSM-network.

188   We used the habitat map of Lok et al. 2024 to determine whether birds were in foraging

habitat (i.e. freshwater, brackish or marine habitats) or on land. Moreover, we used the

behavioural classification model developed in Lok et al. 2023 to estimate a bird's behaviour on

the basis of 0.4 s samples of accelerometer data. As juveniles spend a considerable part of

their time walking, which is difficult to distinguish from foraging (Lok et al., 2023), for this study,

we 'only' distinguish whether birds are 'active on the ground' (i.e., either foraging, walking or

drinking, and for juveniles likely also including begging), 'flying' or 'resting'.

195   The day of departure on autumn migration was defined as the first day that the individual

travelled more than 100 km in southward direction between the first and last GPS fix of that day

and ended up more than 50 km south of the colony. The precise moment of departure was

defined as the date and time associated with the GPS fix preceding the first migratory flight,

which was defined as a flight (i.e., a bout of consecutive GPS intervals during which the average

speed was at least 5 km/h), with a net travel distance of at least 30 km that started on the day of

departure or the preceding day. Including the preceding day was necessary to include the few

cases where a bird departed on the evening or night before the departure day as defined above.

203   To identify contact moments between chick-parent pairs, and the behaviour of the chick

during these contact moments, we linked the GPS data of the chick and its parent(s) based on

the closest GPS timestamps between the two. We defined the chick to be in contact with its

parent when the distance between the two was less than 10 meters. We realize that this is a

rather arbitrary boundary distance that likely results in an underestimate of the actual amount

of contact as the GPS timestamps of the chick and its parent are never really at the same time.

Particularly when birds are walking, foraging or flying, the distance between the locations of the

chick and parent quickly becomes larger than 10 m. Therefore, we also did the analysis with a

contact distance of 50 m.

Behavioural classification

Lok *et al.* (2023) developed a random-forest model for the behavioural classification of

spoonbill behaviours from accelerometer data. This model accurately classified resting (sitting

and standing), flying and searching (F>0.90) and reasonably accurately identified the ingestion

of prey (F=0.73) and walking (F=0.65). Drinking and handling prey were poorly classified

(F<0.10). However, this model was trained on acceleration data associated with video-

annotated behaviour of adult spoonbills. As such, this model did not include the chick-specific,

and for this paper highly relevant, behaviour 'begging' for food at a parent, where the chick

moves its head up and down (also referred to as 'pumping') while standing next to the parent or

walking after the parent. Eventually, this begging may be followed by the parent feeding the

chick, but as this is such a rarely occurring (only a few times per 24 hours) and very short-lasting

(of a few seconds only) event compared to the amount of begging (usually lasting several

minutes) preceding this feeding event, we did not attempt to include 'feeding the chick' or

'being fed' into our behavioural assay. To be able to distinguish 'begging' from the

accelerometer data, we collected video-footage of GPS-tagged chicks and managed to get 110

s of video-material of begging behaviour, though from one chick only (6381). In addition to this

video-footage we visually annotated another 337 s of begging data that directly preceded our

video-footages, based on the very distinct acceleration signal of begging. We added these

video- and visually annotated 'begging' data to the annotated data used in Lok et al. (2023) and

trained a random-forest model to distinguish the behaviours 'search', 'handle', 'ingest', 'sit',

'stand' (also including cases where the bird was moving its body, e.g. when preening, drinking

or shaking feathers), 'fly-active', 'fly-passive', 'walk', as covered by Lok et al. (2023)), as well as

a new behavioural class 'beg'. As predictor variables in the models, we used the summary

statistics described in Lok et al. (2023), calculated over fixed-length segments (of 0.4 s, 0.8 s or

1.6 s) of the 20 Hz acceleration data measured over three axes, along with GPS-speed. As for

the current study, we are primarily interested in distinguishing begging and foraging from other

behaviours, we calculated the classification performance (using the F-measure as a balanced

measure of sensitivity and precision, see Lok *et al.* 2023) of the pooled behaviours 'forage'

(including 'search', 'handle' and 'ingest'), 'rest' (including 'sit' and 'stand') and 'fly' (including

'fly-active' and 'fly-passive'), and the unpooled behaviours 'walk' and 'beg'. We subsequently

applied the best-performing model on the chick data. As walking is often confused with foraging

(Lok et al., 2023), we assumed a chick was foraging when it was classified to be foraging on the

basis of its accelerometer data and GPS-speed and it was located in foraging habitat. All other

classifications of foraging were considered to be walking.

Data selection and analysis

To select the period after fledging, we only used data of chicks that were at least 35 days old.

Using the visual feeding observation dataset, we tested whether the number of observed

feedings decreased with chick age (randomly selecting one observation per known-age chick,

N=127) or depended on the chick's sex (randomly selecting one observation per sexed chick,

N=86) or parent's sex (randomly selecting one observation per sexed parent, N=57) using

generalized linear models with a Poisson error distribution. To have enough counts per age

class to make robust statistical inferences, we created chick age classes of 10 days, where 40 =

35-44 days old, 50 = 45-54 days old etc. Using linear models, we tested whether the distance

from the colony where a feeding took place increased with (unpooled) chick age.

Of the 15 GPS-tagged juvenile spoonbills, 12 had one GPS-tagged parent while 3 chicks

had two GPS-tagged parents, resulting in 18 chick-parent pairs. For the analysis of GPS-based

contact moments, we only used data where the absolute difference between the nearest

timestamp of the chick and that of its parent was less than 10 minutes, or where the parent had

already departed while the chick had not. In the latter case, despite a potentially larger time

difference than 10 minutes, it was certain that chick and parent were not in contact.

Furthermore, to derive reliable estimates of age-specific proportions of contact between a

chick and its parent, we only used days with at least 130 linked GPS locations per chick-parent

pair (there would be 144 if the GPS locations occurred at exactly 10-minute intervals and no

GPS fixes were missed). For modelling behaviour-specific contact, we only used days with at

least 100 linked GPS locations with an associated complete (1.6 s of 20 Hz) acceleration

sample for the chick. This dataset is smaller as acceleration measurements were sometimes

incomplete or missing, which occurred seemingly randomly, likely due to interference with the

process of downloading the data. Finally, we only included chick-parent pairs with at least 10

days of such nearly complete joint data and, due to the much lower data resolution for chicks

after their departure on autumn migration, only data prior to departure of the chick. As a result,

16 chick-parent pairs with on average 71 days of nearly complete overall contact and 68 days of

behaviour-specific contact data were retained for analysis (Table S1). The oldest chick in this

dataset that still had contact with a parent was 136 d old. Using data of chicks between 35 and

maximally 136 d old, we tested whether the proportion of time that chicks have contact, overall

or while the chick was begging or foraging, with their parent(s), and the distance to the nest

during contact, was explained by chick age, chick sex or parent sex. For this, we used

generalized linear mixed models with a binomial (for contact probabilities) and Gamma (for

distance to the nest, using a log-link) error distribution and correlated random variation in the

intercept and chick age slope among chicks. To avoid overfitting, and as we didn't have clear

expectations about this, we did not consider the two- and three-way interactions between chick

age, chick sex and parent sex. While we had not anticipated this beforehand, exploratory

analysis revealed strong annual variation in the amount of contact between chick and parent,

potentially caused by environmental variation (e.g., food availability). Therefore, we account for

this year effect by including it as a fixed effect (with three levels: 2016-2018) in all models. To

facilitate convergence, chick age was standardized in all models.

After departure on autumn migration, for most chicks and some parents, only GSM-data

were available for which the GPS-intervals varied between 1-6 hours. Moreover, because of the

high speed during migratory flights, even if the birds were in the same flock, a small time-

difference between the GPS-timestamp of the chick and the parent would already result in

more than 10 meters distance between the birds. Therefore, we assessed whether chicks and

parents may have been in contact during or after autumn migration by graphically exploring the

cases where the distance between the nearest GPS locations was less than 25 km.

All statistical analyses were performed in program R (v. 4.4.1, R Core Team, 2024) using

package lme4 (Bates, Maechler, Bolker, & Walker, 2015) to fit generalized linear mixed models

and package DHARMa (Hartig 2022) to confirm homogeneity of residuals and lack of

overdispersion. Model support was assessed on the basis of the Akaike information criterion

adjusted for small sample sizes (Burnham & Anderson, 2002). We selected the most

parsimonious model as the model within 2 ΔAIC_c of the best-supported model that had the

least parameters. Where applicable, post-hoc pairwise comparisons were performed using R-

package emmeans (Lenth, 2022).

Ethical note

This study was conducted under license numbers D6548 and AVD105002016446 following the

Dutch Animal Welfare Act and the ethical standards of the Dutch Centre for Avian Migration &

Demography. Animals were solely handled by persons with the required permits. To minimize

potentially negative effects on other breeding birds, birds were only captured during good

weather conditions (i.e. no rain, no strong winds (<6 Bft), and at mild temperatures (<25 °C)).

The maximum duration of catching birds in the colony per day was one hour for adults and <10

minutes for juveniles. The processing of the caught birds took place at c. 100 m from the

colony, during which the other birds returned to the colony. To minimize handling stress, the

bird's head was covered with a cotton cloth as much as possible during handling. Moreover,

banding, measuring, collecting blood samples and GPS-tracker mounting were performed as

quickly as possible (typically under 20 min handling time between capture and release). After

handling, the birds were released immediately <100 m from the colony (for adults) or in the

colony (for juveniles).

Results

Visual feeding observations

The number of observed feedings of colour-ringed chicks after fledging strongly decreased with

chick age (N=127, z=-8.87, p<0.001, Figure 1a), with the oldest chick observed to be fed being

125 days old. While accounting for this age effect, the number of observed feedings did not

significantly differ between female (N=40) and male chicks (N=46; z=0.56, p=0.58) nor between

female (N=23) and male parents (N=34; z=1.28, p=0.20; Figure S1). Of the 127 observed

feedings, 70% occurred within 10 km from the natal colony, and the remaining 30% between 10

and 40 km distance from the colony (Figure 1b, c). The distance from the colony at which chicks

are fed increased with chick age (R^2=0.33, $F_{1,125}$=62.06, p<0.001, Figure 1b).

Classification of begging behaviour

327   Begging behaviour was fairly accurately classified, with the highest classification performance

328   achieved when using the longest segments of 1.6 s (F=0.89; Table S2). The other pooled

329   behaviours were classified similarly well at 0.8 and 1.6 s (Table S2). As for this paper, the

330   primary interest is in distinguishing begging and foraging, we trained and applied the random-

331   forest model on 1.6 s acceleration data segments.

GPS-based contact moments

The overall probability of contact (i.e. a distance of less than 10 m) between a GPS-tagged chick

and its GPS-tagged parent was best explained by the model that included an effect of chick age

and parent sex (Table 1). The probability of contact generally decreased with increasing chick

age and mothers had somewhat more contact with their chick than fathers (Figure 2, Table S3).

Similar results were found for the probability of contact while the chick was classified to be

begging or foraging (Table 1), although the slope of the age effect was opposite for foraging

contact: older chicks were more likely to be foraging together with a parent than younger chicks

(Tables S4, S5). However, the parameter estimates of the fixed effects were usually smaller

than, or at most similar to, the estimated standard deviations (σ) for the random intercepts and

slopes (Tables S3-S5). This indicates that the variation among chicks in the probability to have

contact with their parent(s), and how this changed with age, was larger than what could be

344 explained by the fixed effects (Figures S2-S4). Indeed, the age at which a chick last had contact

345 with its parent strongly varied, between 44 and 136 d (median: 88 d).

346   Variation in distance to the nest during contact was also most parsimoniously explained

347 by chick age and sex of the parent (Table 1). The distance to the nest during contact increased

348 with chick age and contact with the father on average occurred further away from the nest than

349 contact with the mother (Figure 2; Table S6). When using a contact distance of 50 m instead of

350 10 m, the overall percentage of contact time (averaged over all chicks, irrespective of chick or

351 parent sex) increased from 11.4% to 14.0% for 40 d old chicks and from 2.5% to 3.1% for 90 d

352 old chicks. Model selection results were similar (Table 1 vs S7).

353 *Departure on autumn migration*

354 Thirteen of the 15 GPS-tagged chicks departed on autumn migration, at ages varying between

355 108 and 139 d (median: 116 d). Except for one chick-parent pair, the last contact occurred prior

356 to migration departure of either chick or parent: contact stopped 0 – 72 d (median: 28 d) prior to

357 the chick's departure and 0 – 70 d (median: 8 d) prior to the parent's departure.

358   Of the 11 GPS-tagged chick-parent pairs where both chick and parent(s) departed on

359 migration, only one chick departed on the same day as its parent (Figure 3). Visual inspection of

360 the timing and routes during their departing flights revealed that this chick (6295; a female) and

361 its parent (6288; a male) were in the same flock (Figure 4). However, while her father landed in

362 the Kwade Hoek in the Dutch Delta, 6295 continued flying another 75 km to the Baai van Heist

363 in northern Belgium. A few days later, they briefly met again at a roost along the Slikken van

364 Flakkee, after which 6295 headed further south on 30 September, while her father stayed for

365 another few days and departed on 2 October. So, even while they departed together from the

366 north of The Netherlands, their connection was already lost in the south of The Netherlands.

367   The distance between the nearest (in time) GPS locations of the other 10 chick-parent

368 pairs after their departure on migration, and of 6295-6288 after 30 September, was always >40

369 km, implying that chicks and parents did not meet again at later stages of the migration nor at

370 the wintering sites.

Discussion

372 Despite its potential role in affecting survival, habitat use and migration strategies of juvenile

373 birds, post-fledging parental care is an understudied aspect of avian life histories due to the

374 difficulty of following families over large distances. In this study, we investigated post-fledging

375 parental care in Eurasian spoonbills in The Netherlands through a combination of visual feeding

observations and GPS-tracking along with accelerometer-based classification of behaviours of

parent(s) and their chick. We found that the number of feedings and the amount of contact that

chicks still have with their parents strongly decreased with chick age and (usually) ends before

the onset of autumn migration. The amount of post-fledging parental care received (and

provided) strongly varied between chicks (and parents), with some evidence that mothers

provided more care than fathers.

382   In our study, all visually observed feedings of juvenile spoonbills occurred within 40 km

from the breeding colonies. This is similar to the Hungarian population, where the majority

(86%) of fledged colour-ringed chicks were fed within 30 km of the breeding colonies (Pigniczki,

2022). However, in this population, there were also two (out of 14) chicks observed being fed at

respectively 111 km north and 140 km south of their colony. As observations along the Central-

European flyway are scarce, this begged the question whether post-fledging parental care

might sometimes extend into the migratory period. Yet, despite considerable observation

efforts along the East-Atlantic flyway, with >10 000 observations of c. 4000 juvenile birds from

The Netherlands during their first autumn migration (i.e., observed south of The Netherlands

during the months July-December) between 1994 and 2024, there was not a single observation

of a juvenile being fed by a parent (Figure 1). Also for the GPS-tagged chicks and their parent(s),

all contact moments were within 18 km from the colony, except for the contact between chick

6295 and father 6288 during their joint outbound migratory flight (Figure S4). That some

juveniles in Hungary were fed at considerably larger distances from the colony might be

explained by potentially more variable food availability in this population, where spoonbills rely

entirely on freshwater habitats that might dry out or become densely vegetated in the course of

the summer. Contrastingly, in our dataset, the majority of observed feedings came from chicks

from the Wadden Sea area or the Dutch Delta, where many marine and brackish foraging

grounds remain available throughout the breeding season.

401   While the GPS-data revealed that female parents had more contact with their chick after

fledging than male parents, female parents were not more often observed feeding a chick

(N=23) than male parents (N=34). In fact, although the difference was not statistically

significant, the visual feeding observations even pointed in the opposite direction. In absence of

a strong bias in the percentage of females among the visually or molecularly sexed adults in the

entire resighting dataset (48%), there remain several explanations for this apparently

contrasting result. Firstly, contact does not necessarily mean that a chick is fed. However, the

GPS-tagged chicks were also more likely to be begging at their mother than at their father,

which may be assumed to reflect the probability of being fed. Alternatively, the estimated

parental sex effect in the GPS-data may not be very representative as it strongly driven by the

data of the two chicks of which both parents were GPS-tagged. In both cases, the mother had

more contact with the chick, overall and while the chick was begging or foraging, than the father

(Figs. S2-S4). But with N=2 only, this may not necessarily be a general pattern. Among chicks,

there was considerable variation in the amount of contact they still had with their parent,

whether it was with their father or their mother (Figs. S2-S4). Consequently, if we would treat

the two chicks of which both parents were GPS-tagged as four unique chick-parent pairs (i.e.,

by using chick-parent pair instead of chick as random effect), the effect of sex of the parent was

no longer supported (Table S8).

419   We used contact between GPS-tagged chicks and parent(s) as an indication of post-

fledging parental care. However, it is uncertain, at least when it comes to contact while the

chick was resting or foraging (the chick was on average only begging during c. 3% of its contact

time), whether this contact was 'deliberate' (i.e. that the parent was actively protecting or

guiding the chick, or the chick actively sought out or followed its parent) or 'coincidental'. The

latter could be caused by the sociality of spoonbills, irrespective of their relatedness. Shortly

after fledging, social structuring occurs between roosts near the colony and adjacent foraging

grounds in the Wadden Sea. Later on, spoonbills from multiple colonies congregate in larger

groups, where they moult and prepare for (e.g., fuel and seek companions to depart with on)

migration (de Boer et al., 2024; Henriques et al., in press). That at least part of the contact,

particularly at older ages, may be coincidental is suggested by the fact that six chick-parent

pairs showed a long period without contact after which there was contact again (Figures 3, S2).

In four of these cases (6295-6288, 6298.1-6118, 6304-763 and 6358.1-6066), this was caused

by the fact that the parent (in all four cases the father) no longer returned to the colony and at a

certain point, the chick dispersed away from the colony and met the parent again (Figure 5). In

the other two cases (6315-6289 and 6385-6291), the chick was the first to move away from the

colony, whereas the parent (in both cases the mother) followed later (Figure 5).

436   Our results indicate that the amount of post-fledging parental care received (and

provided) is highly variable among chicks (and parents). First of all, it should be realized that in

most cases, we only have information on the contact between the chick and one of its parents.

Therefore, it might be possible that the untagged parent compensates for the lack of care

provided by some of the GPS-tagged parents. Yet, in the two cases where both parents were

GPS-tagged, both provided post-fledging parental care, making this a rather unlikely scenario.

The variable care provided was also not explained by the fact that some parents had to divide

their care over multiple chicks: among the four GPS-tagged chicks that had a younger or older

sibling (6298.1, 6299, 6292 and 6381), three of them received hardly any post-fledging parental

care, whereas 6299 was among the chicks that received the most care. Similarly, among the ten

GPS-tagged chicks that were the only chick in the nest that survived until fledging, the amount

of care received also varied considerably.

It is also possible that parents compensate for differences in the amount of care they

already provided prior to fledging. During incubation and nest-bound chick-rearing, females and

males equally shared nest attendance duties, but females spent more time foraging (Lok et al.,

2024). As a result, female parents may be expected to contribute less to post-fledging parental

care than males, but this was not the case. In addition to potential sex-specific energetic

considerations, parents are known to vary in terms of migration distance (Lok, Veldhoen,

Overdijk, Tinbergen, & Piersma, 2017), with long-distance migrants potentially being more

constrained in terms of the amount of care they can afford to provide while they also invest in

moult and pre-migratory fuelling.

Rather than (or in addition to) being caused by variation in the availability or energetic

state of the parents, variation in post-fledging parental care could be caused by variation in the

energetic requirements of the chicks. When food availability is low, chicks may fledge in poorer

condition and have more difficulties in finding food for themselves, therefore begging their

parents to feed them for a longer period of time in order to survive. On the other hand, under

poor food conditions, parents may decide to secure their own survival and future reproductive

prospects at the cost of the survival of their current young, as is expected and commonly

observed in long-lived species (Stearns, 1992). That the latter might play a role in explaining the

observed variation is supported by the fact that in 2017, the year in which hardly any post-

fledging parental care was provided, breeding success (unpublished data) as well as estimated

prey ingestion rates during late summer in the Wadden Sea (Lok et al., 2023) were relatively low.

Although estimated prey ingestion rates in the Wadden Sea were similarly low in 2018, this

year's summer was extremely dry, causing the freshwater lake Westerplas on Schiermonnikoog

to dry out nearly completely with the fish being trapped in a very small layer of water: a foraging

feast for spoonbills (pers. obs.).

Rather than by energetic aspects, the observed inter-family variation in post-fledging care

could also be caused by different processes. It is now well established that birds often exhibit

consistent between-individual differences in boldness and explorativeness across different life

stages and environmental contexts (Réale, Reader, Sol, McDougall, & Dingemanse, 2007).

Some chicks may simply be more demanding than others, persistently following their parents

for weeks. Similarly, some parents may be more caring than others, regardless of their

energetic state.

Despite the considerable variation in the extent of contact prior to departure, in 10 of the

11 GPS-tagged families where both chick and parent departed on autumn migration, the chick

departed on a different day than (i.e. independently from) its parent. In the first half of the

migratory season (until 2 October), juveniles usually departed later than their GPS-tagged

parent, whereas later on, the juveniles (N=3) departed earlier. There was only one case where

the chick departed in the same flock as its parent, but they did not stop at the same place,

hence contact was already lost prior to leaving The Netherlands (Figure 3). Moreover, prior to

the two days preceding their joint southward departure, the chick and its parent had not been in

contact for 36 days (Figures 3, S2). This suggests that it was not the bond between chick and

parent, but rather coincidence due to spoonbills socially migrating in (large) flocks, that caused

them to be in the same flock during their departing flight. Therefore, our results suggest that,

unlike geese, swans, cranes and several other species (Alonso et al., 2004; Black & Owen,

1989; Byholm et al., 2022; Chetverikova et al., 2017; Earnst & Bart, 1991; Palacín et al., 2011;

Scott, 1980), spoonbill chicks and parents do not deliberately migrate together. Apparently, the

costs of prolonged care by the parents during migration are higher than the benefits gained from

it by either the juveniles (e.g., by receiving food and/or being guided during migration) or the

parents (e.g. through increased dominance status). However, as spoonbills usually migrate in

large flocks, consisting of both adult and juvenile birds (Piersma, de Goeij, Bouten, & Zuhorn,

2022), juveniles may use the information from other experienced adults to learn suitable

migration routes, to find staging and wintering sites, and to exploit local resources across a

variety of locations and environments – often very different from the environment in which they

were born and cared for.

To conclude, both parents contribute to post-fledging parental care, with considerable

variation in the amount of care provided. It remains unclear whether this variation is caused by

differences in the demands of chicks to still receive post-fledging parental care, the demands

of the parents, or by different processes entirely. The amount of parental care strongly

decreases with chick age and (usually) ends several weeks before departing on autumn

migration with other conspecifics.

Data availability

Tracking data generated and analyzed in this study are stored at Movebank

(www.movebank.org) under the studies "SPOONBILL_METAWAD -Eurasian Spoonbills (Platalea

leucorodia, Threskiornithidae) breeding on Schiermonnikoog, The Netherlands" (Movebank

study ID 2596955604) and "SPOONBILL_SPINOZA – Eurasian Spoonbills (Platalea leucorodia,

Threskiornithidae) born on Schiermonnikoog, The Netherlands" (Movebank study ID

2818737964) and will be made available upon reasonable request.

Analyses reported in this article can be reproduced using the code and additional data

published at GitHub (https://github.com/tamarlok/postfledging_care).

Acknowledgments

We thank Natuurmonumenten for permission to work in national park Schiermonnikoog,

Yvonne Verkuil for molecularly sexing, Willem Bouten for technical assistance on UvA-BiTS, Iris

Kromhout Van der Meer, Marnix Bosma, Remco Rood and many other volunteers for their

assistance in the field and Werkgroep Lepelaar and voluntary ring-readers for ringing and

resighting (and observing the behaviour and sex of) colour-ringed spoonbills. This research was

financially supported by the Metawad project awarded to TP by Waddenfonds (WF209925), the

Spinoza Premium 2014 awarded to TP, and a Veni grant awarded to TL (016.Veni.192.245) by

The Netherlands Organisation for Scientific Research. UvA-BiTS studies are facilitated by

infrastructures for e-Ecology, developed with the support of NLeSC and LifeWatch, and carried

out on the Dutch national e-infrastructure with the support of SURF Cooperative.

References

Alonso, J. C., Bautista, L. M., & Alonso, J. A. (2004). Family-based territoriality vs flocking in

wintering common cranes *Grus grus*. *Journal of Avian Biology, 35*, 434-444. doi:

10.1111/j.0908-8857.2004.03290.x

Ashmole, N. P., & Tovar, S. H. (1968). Prolonged parental care in royal terns and other birds.

Auk, 85, 90-&.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software, 67*, 1-48. doi: 10.18637/jss.v067.i01

Black, J. M., & Owen, M. (1989). Parent-offspring relationships in wintering barnacle geese.

Animal Behaviour, 37, 187-198. doi: 10.1016/0003-3472(89)90109-7

Bouten, W., Baaij, E. W., Shamoun-Baranes, J., & Camphuysen, K. C. J. (2013). A flexible GPS

tracking system for studying bird behaviour at multiple scales. *Journal of Ornithology,*

154, 571-580. doi: 10.1007/s10336-012-0908-1

Burnham, K., & Anderson, D. (2002). *Model selection and multi-model inference: a practical*

information-theoretic approach (2nd ed.). New York: Springer-Verlag.

Byholm, P., Beal, M., Isaksson, N., Lötberg, U., & Åkesson, S. (2022). Paternal transmission of
migration knowledge in a long-distance bird migrant. *Nature Communications, 13*. doi:
10.1038/s41467-022-29300-w
Chetverikova, R., Babushkina, O., Galkina, S., Shokhrin, V., & Bojarinova, J. (2017). Special case
among passerine birds: long-tailed tits keep family bonds during migration. *Behavioral
Ecology and Sociobiology, 71*. doi: 10.1007/s00265-017-2268-6
Cramp, S. (1994). *The birds of the Western Palearctic*. Oxford: Oxford University Press.
de Boer, A. P., Vansteelant, W. M. G., & Piersma, T. (2024). Primary moult of Eurasian spoonbills
Platalea l. leucorodia in the Wadden Sea in relation to age, breeding and migration.
Ardea, 112, 217-227, 211.
Earnst, S. L., & Bart, J. (1991). Costs and benefits of extended parental care in Tundra Swans
Cygnus columbianus columbianus. *Wildfowl Special Issue, 1*, 260-267.
Ersoy, S., Maag, N., Boehly, T., Boucherie, P. H., & Bugnyar, T. (2021). Sex-specific parental care
during postfledging in common ravens. *Animal Behaviour, 181*, 95-103. doi:
10.1016/j.anbehav.2021.09.004
Fijn, R. C., & Kemper, J. (2023). Sandwich Tern Feeds Juvenile on Wintering Grounds in Southern
Namibia. *Ardea, 111*, 549-553, 545.
Fridolfsson, A. K., & Ellegren, H. (1999). A simple and universal method for molecular sexing of
non-ratite birds. *Journal of Avian Biology, 30*, 116-121.
Grüebler, M. U., & Naef-Daenzer, B. (2010). Survival benefits of post-fledging care: experimental
approach to a critical part of avian reproductive strategies. *Journal of Animal Ecology, 79*,
334-341. doi: 10.1111/j.1365-2656.2009.01650.x
Henriques, M., Piersma, T., Vansteelant, W. M. G., de Goeij, P., & Lok, T. (in press). A place and
a time for everything: Eurasian spoonbills divide spring and summer activities across
different areas in the eastern Wadden Sea. *Ardea*.
Koenig, W. D., & Dickinson, J. L. (2004). *Ecology and evolution of cooperative breeding in birds*.
Cambridge: Cambridge University Press.
Krijgsveld, K. L., Dijkstra, C., Visser, G. H., & Daan, S. (1998). Energy requirements for growth in
relation to sexual size dimorphism in marsh harrier Circus aeruginosus nestlings.
Physiological Zoology, 71, 693-702.
Lagarde, F., & Piersma, T. (2021). Vocal signalling by Eurasian Spoonbills *Platalea leucorodia* in
flocks before migratory departure. *Ardea, 109*, 243-250, 248.
Lenth, R. V. (2022). emmeans: Estimated Marginal Means, aka Least-Squares Means. R
package version 1.7.2.

Lok, T., Overdijk, O., & Piersma, T. (2014). Interpreting variation in growth of Eurasian Spoonbill
chicks: disentangling the effects of age, sex and environment. *Ardea, 102*, 181-194. doi:
10.5253/arde.v102i2.a8

Lok, T., Overdijk, O., Tinbergen, J. M., & Piersma, T. (2013). Seasonal variation in density
dependence in age-specific survival of a long-distance migrant. *Ecology, 94*, 2358-2369.
doi: 10.1890/12-1914.1

Lok, T., van der Geest, M., Bom, R. A., de Goeij, P., Piersma, T., & Bouten, W. (2023). Prey
ingestion rates revealed by back-mounted accelerometers in Eurasian spoonbills. *Animal*
Biotelemetry, 11, 5. doi: 10.1186/s40317-022-00315-w

Lok, T., van der Geest, M., de Goeij, P., Rakhimberdiev, E., & Piersma, T. (2024). Sex-specific
nest attendance rhythm and foraging habitat use in a colony-breeding waterbird.
Behavioral Ecology, 35, arae020. doi: 10.1093/beheco/arae020

Lok, T., Veldhoen, L., Overdijk, O., Tinbergen, J. M., & Piersma, T. (2017). An age-dependent
fitness cost of migration? Old trans-Saharan migrating spoonbills breed later than those
staying in Europe, and late breeders have lower recruitment. *Journal of Animal Ecology,*
86, 998-1009.

López-Idiáquez, D., Vergara, P., Fargallo, J. A., & Martínez-Padilla, J. (2018). Providing longer
post-fledging periods increases offspring survival at the expense of future fecundity.
PLOS ONE, 13, e0203152. doi: 10.1371/journal.pone.0203152

Mellink, E., Castillo-Guerrero, J. A., & Peñaloza-Padilla, E. (2014). Development of diving
abilities by fledgling Brown Boobies (*Sula leucogaster*) in the Central Gulf of California,
México. *Waterbirds, 37*, 451-456.

Meyburg, B.-U., Meyburg, C., Mizera, T., Maciorowski, G., & Kowalski, J. (2005). Family break up,
departure, and autumn migration in Europe of a family of greater spotted eagles (*Aquila*
clanga) as reported by satellite telemetry. *Journal of Raptor Research, 39*, 462-466.

Naef-Daenzer, B., & Grüebler, M. U. (2016). Post-fledging survival of altricial birds: ecological
determinants and adaptation. *Journal of Field Ornithology, 87*, 227-250. doi:
10.1111/jofo.12157

Palacín, C., Alonso, J. C., Alonso, J. A., Magaña, M., & Martín, C. A. (2011). Cultural transmission
and flexibility of partial migration patterns in a long-lived bird, the great bustard Otis
tarda. *Journal of Avian Biology, 42*, 301-308. doi: 10.1111/j.1600-048x.2011.05395.x

Phillips, R. A., Xavier, J. C., & Croxall, J. P. (2003). Effects of satellite transmitters on albatrosses
and petrels. *The Auk, 120*, 1082. doi: 10.1093/auk/120.4.1082

Piersma, T., de Goeij, P., Bouten, W., & Zuhorn, C. (2022). *Sinagote. The biography of a*
spoonbill. Barcelona: Lynx Edicions.

Pigniczki, C. (2022). Observations on parental care of the Eurasian Spoonbill during the post-
fledging dispersal. *Ornis Hungarica, 30*, 146-157. doi: doi:10.2478/orhu-2022-0011

R Core Team. (2024). R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing.

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating
animal temperament within ecology and evolution. *Biol Rev Camb Philos Soc, 82*, 291-
318. doi: 10.1111/j.1469-185X.2007.00010.x

Rehling, A., Spiller, I., Krause, E. T., Nager, R. G., Monaghan, P., & Trillmich, F. (2012). Flexibility
in the duration of parental care: zebra finch parents respond to offspring needs. *Animal*
Behaviour, 83, 35-39. doi: 10.1016/j.anbehav.2011.10.003

Riou, S., Chastel, O., & Hamer, K. C. (2012). Parent–offspring conflict during the transition to
independence in a pelagic seabird. *Behavioral Ecology, 23*, 1102-1107. doi:
10.1093/beheco/ars079

Scott, D. K. (1980). Functional aspects of prolonged parental care in Bewicks swans. *Animal*
Behaviour, 28, 938-952. doi: 10.1016/s0003-3472(80)80156-4

Stearns, S. C. (1992). *The evolution of life histories*. Oxford: Oxford University Press.

Whiten, A. (2019). Cultural Evolution in Animals. In D. J. Futuyma (Ed.), *Annual Review of*
Ecology, Evolution, and Systematics, Vol 50 (Vol. 50, pp. 27-48).

Tables & Figures

Table 1.

Model selection results. The most parsimonious models are indicated in bold. "age" refers to age of the chick and "c" to the intercept-only "constant" model. For modelling contact probabilities, year was always included as a categorical fixed effect.

Model	K	Δ-2logL	ΔAIC$_c$	Akaike weight
(a) Total contact				
sex$_{parent}$ + age	**8**	**0.81**	**0.00**	**0.64**
sex$_{chick}$ + sex$_{parent}$ + age	9	0.00	1.19	0.35
sex$_{parent}$	7	11.50	8.69	0.01
sex$_{chick}$ + sex$_{parent}$	8	10.70	9.89	0.00
Age	7	600.49	597.68	0.00
sex$_{chick}$ + age	8	598.98	598.17	0.00
C	6	611.06	606.25	0.00
sex$_{chick}$	7	609.51	606.70	0.00
(b) Contact while begging				
sex$_{chick}$ + sex$_{parent}$ + age	9	0.00	0.00	0.51
sex$_{parent}$ + age	**8**	**2.51**	**0.51**	**0.39**
sex$_{chick}$ + age	8	5.61	3.61	0.08
Age	7	11.38	7.38	0.01
sex$_{chick}$ + sex$_{parent}$	8	11.98	9.98	0.00
sex$_{parent}$	7	14.32	10.32	0.00
sex$_{chick}$	7	17.69	13.69	0.00
C	6	23.22	17.22	0.00
(c) Contact while foraging				
sex$_{parent}$ + age	**8**	**0.75**	**0.00**	**0.57**
sex$_{chick}$ + sex$_{parent}$ + age	9	0.00	1.25	0.31
sex$_{parent}$	7	6.81	4.06	0.08
sex$_{chick}$ + sex$_{parent}$	8	5.86	5.11	0.04
age	7	41.00	38.25	0.00
sex$_{chick}$ + age	8	40.08	39.33	0.00
c	6	48.01	43.26	0.00
sex$_{chick}$	7	46.66	43.92	0.00
(d) Distance from nest during contact				
sex$_{parent}$ + age	**7**	**0.13**	**0.00**	**0.72**
sex$_{chick}$ + sex$_{parent}$ + age	8	0.00	1.88	0.28
sex$_{parent}$	6	25.69	23.56	0.00
sex$_{chick}$ + sex$_{parent}$	7	25.53	25.40	0.00
age	6	40.31	38.18	0.00
sex$_{chick}$ + age	7	40.20	40.07	0.00
c	5	65.66	61.53	0.00
sex$_{chick}$	6	65.53	63.40	0.00

Figure 1

(a) The number of observations of colour-ringed chicks that were fed by an adult and (b) the distance from the colony at which these feedings took place in relation to chick age, along with a (c) map of the actual feeding locations connected by black lines to the colony locations where these chicks were born. Plotted data and model estimates are based on one randomly selected observation per known-age chick (N=127).

Figure 2.

The proportion of time that a chick was in contact with its mother (a) or father (b) and the

distance to the nest during contact with the mother (c) or father (d). Error bars reflect 95% C.I.,

using the average proportion of time and distance to the nest during contact per day per chick

as raw data. For the proportion data, bootstrapping was used to determine 95% CI's. Sample

sizes (number of chick-parent pairs) for the calculation of proportion of contact (total bars) and

the distance to the nest during contact (darker colors) are shown above the panels. Only

reasonably reliable estimates are plotted, i.e., estimates that were based on at least 6 different

chick-parent pairs for the proportion of contact and at least 3 different chick-parent pairs for

the distance to the nest during contact.

Figure 3.

Autumn migration departure dates of spoonbill chicks and their parent(s) from The

Netherlands. The dashed line (y = x) is added to visualize whether parents departed earlier

(points above the line) or later (points below line) than their chicks.

Figure 4.

Detailed information about the timing and routes of the female chick 6295 (in orange) and her father 6288 (in blue) during the start of autumn migration from The Netherlands in 2016.

Figure 5.

Distance to the nest of chick (in black) and parent (fathers in blue, mothers in orange), with the

open red circles indicating the contact moments. Here the raw data are plotted, i.e. also the

days with less than 130 registered GPS locations for both chick and parent which was the

criterium to be included in the analyses. Some distances to the nest are not plotted as they

exceeded 50 km, including the last vertical lines representing the outbound migratory flights. All

contact moments are plotted except for the single contact moment between 6295 and 6288 in

the Dutch Delta. Although 6295 and 6288 were also in the same flock during their outbound

migratory flight, this was not estimated as contact as the distance between their locations was

larger than 10 m due to the time difference (varying between 1 and 4 minutes) between the

timestamps of chick and parent combined with the high speed during migration (see Figure S4).

Furthermore, although it may appear that 6354 attempted to depart with his mother 6298, this

was not the case: the departure location, time and routes of their flights were different.

Photo 1.

Adult spoonbill followed by two begging chicks during the post-fledging period at Braakman,

Hoek (51°196'N 3°445'E), 16 August 2015. The chick was colour-ringed on 6 July 2015 in the

colony at Van Cittershaven, Sloegebied (51°263'N 3°433'E), 13 km from the location of this

photo. Photographer: Gab de Croock.