Early development of direction selectivity in higher visual cortex

Authors: Dallas C. Khamiss[1,2,3], Augusto A. Lempel[1,2,3,4], Brandon R. Nanfito[1,2], Kristina J. Nielsen*[1,2]

Affiliations: [1] Solomon H. Snyder Department of Neuroscience, Johns Hopkins University School of

Medicine, Baltimore, MD 21205, USA

[2] Zanvyl Krieger Mind/Brain Institute, Johns Hopkins University, Baltimore, MD 21218, USA

[3]: shared first author

[4] current address: Max Planck Florida Institute for Neuroscience, Jupiter, FL 33458, USA

Corresponding author: Kristina J. Nielsen, knielse4@jhu.edu

Competing interests: The authors declare no competing financial and non-financial interests.

Acknowledgments: This work was funded by NIH EY027853 (KJN) and NIH EY035807 (KJN). We are

thankful for O. Garalde, R. Vistein and other members of the Nielsen lab for experimental support,

and thank J. Killebrew, W. Nash and W. Quinlan for their technical support.

Abstract

A fundamental aspect of visual motion processing is the computation of motion direction. In

ferrets, as in primates, selectivity for motion direction is found both in early cortical stages like the

primary visual cortex (V1) and in higher visual areas like primate MT and ferret PMLS. Little is known

about how this critical tuning function develops in higher visual cortex. Here, by studying the

development of the ferret's motion pathway, we first reveal the surprising finding that direction

selectivity develops earlier in PMLS than in V1, contrary to the areas' hierarchical positions. Our

data furthermore show that while direction selectivity is sensitive to visual experience in both

areas, the sensitivity profile differs between them: Moving stimuli, containing both spatial and

temporal cues, are required to promote direction selectivity development in V1, but flashing stimuli

that remain stationary spatially are sufficient for direction selectivity development in PMLS.

Collectively, our findings reveal that the development of the motion pathway is highly complex

when considered at the network level, and does not adhere to a simple feedforward model in which

the development of lower visual areas sets the pace for higher ones. Not only will developmental

models need to be updated to include this complexity, characterizing the link between the

development of different visual areas will likely also be crucial for understanding the consequences

of developmental disorders.

Introduction

Encoding of motion information is a core function of the visual system with high behavioral

relevance. Not surprisingly, circuits computing motion have been found in a wide range of species,

from flies [1], fish [2] to mammals, including rodents [3–5] and primates (e.g., [6–8]). A particular challenge

for motion computations is that they require not just precise processing of spatial cues but also of

temporal information, which might be one reason why deficits in motion processing occur in a

number of developmental disorders [9]. Because of their behavioral relevance and apparent

vulnerability, the development of circuits supporting motion computations is an important topic for

investigation.

One of the most fundamental forms of motion processing is the computation of motion

direction, which in primates and carnivores predominately occurs in cortex. It is found not just in

early stages like the primary visual cortex (V1) [10,11], but also in higher visual areas crucial for motion

vision like primate MT [12–14] and ferret PMLS (or PSS) [15,16]. Research on the development of direction

selectivity (DS) at the neural level has so far almost exclusively focused on V1 (e.g. [17–20]). As a

consequence, very little is currently known about its emergence in higher visual areas like MT or

PMLS, leaving a significant gap in our understanding of the development of a core motion

computation. Here, we address this issue by systematically following the maturation of DS in ferret

V1 and PMLS. Focusing on the same function in both areas also provides the unique opportunity to

directly compare the developmental mechanisms at play across the visual hierarchy.

In addition to the time course over which certain functions develop, a second important

aspect of visual development is the degree to which it is sensitive to visual information – both its

general presence, as well as its specific content. A number of studies have shown that DS

development in ferret V1 can indeed be influenced by providing or withholding certain types of

visual information [17,21–24]. DS in cat PMLS and monkey MT is also sensitive to altered visual

experience during development [25–27]. We have previously shown that the duration of visual

experience correlates with the developmental status of motion integration in ferret PMLS [28]. Visual

experience therefore also appears to play a role in circuit development in higher motion cortex.

However, all of these studies investigated a single area at a time, and a systematic comparison of

the effects of different types of visual experience across the visual hierarchy is lacking. Here, we

directly compare the impact of different visual cues on V1 and PMLS development, and use it as a

sensitive probe for the developmental mechanisms shaping DS in V1 and PMLS.

In general, our results show a surprising degree of divergence between the two areas, with

DS development occurring earlier, and under the influence of different types of visual information,

in PMLS than V1. DS development therefore does not simply follow a 'feedfoward' sweep in which

higher areas develop after lower ones. This observation provides insights into possible causes for

the vulnerability of motion functions to developmental disorders. It also highlights that the visual

hierarchy may undergo significant reconfiguration during development, important to consider for

models of visual development [29].

Results

<u>After eye opening, DS develops faster in PMLS than V1</u>

DS develops in ferret V1 in the week after eye opening, and matures around postnatal day (P) 37

[17,18]. While we have previously shown that PMLS DS is mature at the same age [28], the time course of

maturation has not yet been determined. Here, we systematically tracked PMLS DS development

and compared it directly to V1. To this end, we recorded responses to drifting gratings in both areas

from P29 (around eye opening) to P52 (young adult). Recordings were performed in anesthetized

animals using multi-channel silicon probes, in many cases collecting data from V1 and PMLS

simultaneously by recording from 2 probes. All data were collected with the eyes open, irrespective

of the natural eye opening state of an animal. Fig. 1B & C show example tuning functions for well

isolated single units (SU) in both areas from P30 to P37. As these plots show, we were able to find

neurons with reliable visual responses at all ages, which allowed us to quantify tuning properties.

The remainder of the analyses in this section will be limited to these visually responsive single

neurons (determined using an ANOVA across mean baseline-corrected firing rates for all conditions

including the blank with a criterion of $p<.05$; see Methods). The resulting data set sizes are listed in

Table 1.

Especially at the youngest ages, tuning functions were often poorly fit with standard

direction (or orientation) tuning functions. We therefore quantified DS by computing two different

metrics. The first was L_{dir} or 1-DirCircVar, a vector-based metric[30] that takes the entire tuning

function into account (see Methods for definition). Importantly, $L_{dir} = 1$ corresponds to high

selectivity (responses to a single direction only), while L_{dir} = 0 indicates no selectivity (equal

responses to all conditions). This metric is comparable to circular variance (CircVar), commonly

used to quantify orientation tuning strength [31]. The second metric used to quantify DS was the

direction selectivity index (DSI), which compares only the responses between the preferred

direction and the one opposing it (the null direction; see Methods). Results were generally identical

between the two metrics, so only the results based on L_{dir} (which is more robust because it is based

on the entire tuning curve) are discussed further (see Fig. S1 for the key DSI results). Both L_{dir} and

DSI values are indicated for each example neuron in Fig. 1B & C, and suggest a general increase in

DS with age in both areas.

106   To more accurately capture the time course of DS maturation and to compare it between

areas, we next analyzed DS as a function of age. As a first step, we computed the mean L_{dir} for each

individual animal by averaging across all visually responsive PMLS or V1 neurons collected in that

animal (results based on the median are comparable; data not shown). This makes it possible to

visualize developmental trends without having to group animals by age; it also more clearly shows

variability in developmental stage across animals of the same age. Fig. 1D plots these per-animal

means as a function of age, with symbol size indicating the number of neurons that passed the

selection criterion for each animal. The two horizontal bands correspond to the mean +/- 1

standard error (SEM) for animals P40 and older. Consistent with previous data[17,18] from ferret V1,

our data show V1 DS to be immature around P30. PMLS DS levels were similarly low at this age. L_{dir}

values eventually improved to mature levels in both areas, with some of the improvements

occurring very rapidly, as evident by significant changes in DS between animals with a single day

age difference. Unexpectedly, this sudden developmental improvement occurred earlier in PMLS

than V1 – around P34 for PMLS, P36 for V1. The faster PMLS development continued, in that adult

values were reached faster after the initial improvements than in V1. Judging by the overlap

between the per-animal mean and the adult range, PMLS matured around P35, but it took V1 until

at least P40 to reach this point.

123   This difference between PMLS and V1 development was also apparent when dividing single

unit L_{dir} values for each area into 3 age groups used previously for ferret V1 development (collapsing

data across animals). The youngest of the 3 groups corresponded to eye opening (P29-32), the

second to the phase of rapid DS development (P33-36), and the last sto animals with mature DS

levels (P37 and older). Fig. 1E & F plot the L_{dir} distributions for these groups for PMLS and V1. Age

had a general effect on the L_{dir} distribution in both areas (Kruskal-Wallis test across all 3 groups:

p<.001 for both areas; full statistics in Table 2). For PMLS, post-hoc rank-sum tests revealed that

the P29-32 group deviated from both other age groups (p<.001), which were not significantly

different from each other. In V1, on the other hand, the P29-32 & P33-35 groups were not

significantly different, but both differed from the oldest group (p<.001).

Since development is obviously dynamic and averaging across age groups cannot capture

all trends, Fig. 1G uses an alternate analysis approach to show the relative DS development in V1

and PMLS in more detail. Here, L_{dir} values were binned into age windows of 2 days, combining data

across animals. Since in adult animals DS is stronger in PMLS than V1, the comparison of the

developmental trajectories between the areas is complicated by the fact that they reach different

endpoints. To account for these endpoint differences, we chose to normalize L_{dir} values so that the

average L_{dir} in a 2-day window around eye opening (P29 to 30) was set to 0, and the average L_{dir}

values in young adults (here, P44-45) was set to 1 (see Methods). V1 and PMLS data were

normalized independently, and the normalized L_{dir} values were averaged in each age bin. These

normalized time courses again show that PMLS DS development occurred earlier than in V1: PMLS

L_{dir} values rapidly increased starting with the P33-34 window, while V1 L_{dir} values remained at the

eye-opening level until P35-36. A two-way ANOVA with factors area and age confirmed these

observations. For the ANOVA, the age windows used for normalization were excluded (since they

are forced to be 0 and 1, respectively), reducing the data to P31 – 39. For these data, the two-factor

ANOVA then resulted in a significant effect for age (p<.001), consistent with the general age-

dependent change in DS in both areas. Importantly, a significant effect of area (p<.001) and a

significant interaction between age and area (p=.003) support differences in DS levels between

areas during this time period, which because of the normalization cannot be explained by overall

higher DS levels in PMLS, but reflect developmental differences. Further post-hoc rank-sum tests

indeed revealed significantly higher normalized L_{dir} values in PMLS than in V1 at P33-34 (p<.001), as

well as a trend at P34-35 (p=.02), confirming the earlier development of DS in PMLS (see Table 3 for

full test statistics, including the p-values for all non-significant tests). It is worth noting that the time

courses in Fig. 1G also suggest that V1 DS development does not just start later, it also takes longer

to complete (note the shallower slope of the V1 time course).

At the most fundamental level, high DS requires strong responses to the preferred direction,

and weak responses to the opposite (null) direction. To probe some of the mechanisms underlying

DS development, we compared the baseline-corrected firing rates evoked by preferred and null

direction at different ages. To capture the overall trends, data were divided into the same 3 age

groups used for Fig. 1E & F. In both areas, the responses to the preferred stimulus became more

robust with age (Fig. 1H & I, see Table 2 for statistics). Kruskall-Wallis tests comparing the 3 age

groups were significant for both areas (both p<.001), with significant differences between the P29-

32 and the oldest group in both areas (rank-sum tests, both p<.001), as well as differences between

the P29-32 and P33-36 groups for V1 (p<.001). The increase in the V1 response to the preferred

direction is consistent with previous reports [18]; our data now show that a similar increase occurs in

PMLS, but at a later age than in V1.

Responses to the null stimulus changed in an area-specific manner as well (Fig. 1J & K). In

V1, null direction responses temporarily increased in the P33-36 group, a change that reached

significance in comparison to the oldest group, and trended toward significance relative to the P29-

32 group (Kruskall-Wallis test: p=.008; rank-sum tests P33-36 versus P37+: p=.006, P33-36 versus

P29-32: p=.01). These findings parallel a previous observation of lower V1 null responses after P35

than in younger animals[18]. In PMLS, on the other hand, null direction responses gradually

decreased with age (Kruskall-Wallis and post-hoc test between the P29-32 and the oldest age

group, p<.001; Table 2). This weaker response to the null direction, especially because it was paired

with an increase in the preferred direction response, would support better discriminability of the

two directions, and points to a role of null direction inhibition especially in the development of

PMLS DS (see also Discussion). In summary, our data show earlier and faster DS development in

PMLS than V1, in a process that involves area-specific decreases in null direction responses as well

as increases in preferred responses in both areas.

<u>Early PMLS development appears specific to DS</u>

Since both V1 and PMLS are tuned for orientation in addition to direction, the development of

orientation selectivity (OS) provide the opportunity to assess the areas' relative development along

a second tuning dimension, a crucial test for determining whether PMLS develops early in general,

or whether the observations made so far are specific for DS development. OS was quantified using

two metrics similar to those for DS, a vector-based metric[30,31] called L_{ori} or 1-CircVar, and an

orientation selectivity index (OSI). Again, analyses using either metric showed the same results and

only L_{ori} data will be shown here (for OSI figures, see Fig. S1).

Intriguingly, the relative development of OS in PMLS versus V1 followed a different pattern

than that of DS. In V1, consistent with previous reports[32,33], L_{ori} values were low up to P30. Starting

with P31, much higher L_{ori} values could be observed, even if there was significant variability across

animals (Fig. 2A). Consequently, dividing the data into the same 3 age groups used for L_{dir} before

(P29-32, 33-36, 37+; Fig. 2C revealed significant differences between the P29-32 group and both

other groups (Kruskall-Wallis test and both post-hoc rank-sum tests: p<.01; Table 4), but not

between the two older groups.

In PMLS, OS levels were as low as those found in V1 around P30 (Fig. 2A), with the exception

of 2 outliers (one animal contributing 2 neurons, the other 4). Since in all other respects (e.g., their

L_{dir} values) these animals were not outliers, their data was included in the OS data set as well. Other

than the 2 outliers, OS development in PMLS was delayed relative to V1, with consistent

improvements in PMLS L_{ori} values starting only at P34, later than the increase in V1 L_{ori} values. At

P34, however, PMLS L_{ori} values immediately assumed their mature levels in a very rapid

development, so that OS maturation was complete at the same time point in both areas. Indeed,

dividing the PMLS data set into the same 3 age groups as above (Fig. 2B) revealed differences only

between the P29-32 age groups and both other age groups, comparable to the results for V1

(Kruskall-Wallis test and both post-hoc rank-sum tests: p<.001; see also Table 4).

Given the rapid OS development especially in PMLS, we again turned to binning L_{ori} data into

2-day age windows, normalizing the data to assume 0 and 1 for eye opening and mature levels,

respectively, as before (Fig. 2D). Note that the 2 animals with high L_{ori} values around P30 caused the

normalized L_{ori} time course for PMLS to initially decrease at P31-32 before rapidly increasing in the

next age group. A two-factor ANOVA with age and area (excluding the endpoints used for) was again

used to compare development across the areas. It resulted in a significant main effect of age

(p<.001; see Table 5), consistent with the increase in OS with age. At same time, there was a

significant interaction between area and age (p=.003), due to significantly higher normalized L_{ori}

values in V1 in the P31-32 group (post-hoc rank-sum test, p<.001). To rule out possible effects of

including the two PMLS outliers in the normalization, we confirmed that a direct comparison of the

raw (non-normalized) L_{ori} values for animals aged P31-32 similarly resulted in a significant

difference between V1 and PMLS (rank-sum test, p=.003). Thus, the onset of OS maturation in PMLS

lags behind V1, which in turn highlights that the earlier development of PMLS DS is not due to

earlier development of the area in general, but that it reflects processes specific to processing of

motion direction.

The relative time courses of DS and OS maturation reveal another very intriguing difference

between V1 and PMLS development. Comparing these time courses within each area rather than

across areas (see Fig. S2 for a comparison of the normalized L_{dir} and L_{ori} time courses per area), first

replicated previous findings of OS development before DS development in V1 [17]. In PMLS, however,

our data showed no (or only minimal) delay between DS and OS development (e.g., both

normalized time courses show immature levels at P31-32 and significant improvement by P33-34).

The normalized L_{dir} and L_{ori} values can again be used to compare the relative development of OS

and DS without contamination by overall differences in tuning levels. For V1, a two-way ANOVA with

factors age and tuning property on these normalized data resulted in a significant main effect for

tuning property ($p<.001$) and age ($p=.006$), but no significant interaction between the factors ($p=.2$),

consistent with differences between the two tuning properties that are maintained throughout

development (again, the endpoints used for normalization were excluded from the ANOVA). In

contrast, in PMLS only the effect of age reached significance, reflecting that DS and OS developed

in parallel (age: $p<.001$; tuning: $p=.06$; interaction: $p=.3$). These differences in the sequence of DS

and OS development between V1 and PMLS suggest that there might be fundamental differences in

the organization of circuits computing direction information in V1 and PMLS (see Discussion).

<u>Developmental changes in response latency</u>

In the marmoset, MT has been shown to develop as early as V1, at least anatomically[34]. This early

development involves a transient, direct pathway from the thalamic pulvinar to MT[35,36]. This raises

the interesting possibility that early DS development in PMLS is supported by input sources

different from those found in adults. Resolving this question will ultimately require anatomical

studies. However, response latencies should at least provide some initial insights into network

configurations at different ages. As a next step, we therefore decided to analyze the evolution of

response latencies in V1 and PMLS with age. Latencies can be quite variable from neuron to

neuron. To provide the most robust estimates even at the youngest ages, we increased our sample

size by analyzing multi-unit (MU) instead of SU data. Here, MU activity is computed as the

thresholded but not spike-sorted activity for each recording channel, ensuring that each spike was

only counted on a single channel (see Methods). To be included in the latency data set, MU sites

had to meet the same visual responsiveness criterion used for SU above (see Table 1 for the

resulting data set size). Additionally, we limited the data set to recording sessions in which data was

collected simultaneously for V1 and PMLS so that within-animal comparisons were feasible in all

cases. This also minimizes confounds from changes in response dynamics that might be caused by

differences in anesthesia state.

For each included MU site, we then determined the onset latency for the site's best

stimulus (as the condition evoking the most robust responses). Latencies were computed by

binning spikes into 10 ms bins and computing the first bin in which responses exceeded the mean

of pre-stimulus baseline activity by two standard deviations, followed by at least two successive

bins exceeding baseline activity [37]. Fig. 3A shows raster plots for 3 example V1 MU sites at different

ages, as well as 3 example PMLS sites recorded in the same animals. These examples capture two

important trends present in the overall data set shown in Fig. 3B: First, latencies in both areas

became significantly shorter with age, until very short latencies were reached in adults (median V1

latency across all MU sites: 60ms; median PMLS latency: 70ms). Second, there was a large delay in

PMLS responses relative to V1 in young animals. As latencies in both areas gradually improved over

the next days, the gap in latencies between the areas ultimately narrowed and became quite small

in adults.

Fig. 3C & D plot how these findings play out at the per-neuron rather than per-animal level

(see Tables 6 & 7 for full statistics). First, Fig. 3C plots the distribution of latencies for the standard 3

age groups, demonstrating the gradual improvement in latencies with age in both areas (Kruskall-

Wallis test across groups per area and all post-hoc rank-sum tests : $p<.001$). Second, Fig. 3D shows

how average latencies, computed in 2-day windows (this time without normalization), evolve with

age, highlighting the gradual improvement in latencies between P30 and 35 in both areas, as well as

the larger delay in PMLS responses in this earlier phase. Indeed, significant differences between V1

and PMLS emerged only for the three age windows falling between P29 and P34 (2-way ANOVA with

factors age and area: all main effects and interaction $p<.001$; post-hoc rank-sum tests between

areas for P29-30, P31-32, P33-34: $p<.001$), but note that there is still a trend towards larger PMLS

latencies for older animals ($p<=.05$ for all subsequent age windows). Finally, we made use of the

fact that the data set was limited to cases with simultaneous recordings in V1 and PMLS to test for

each animal individually whether V1 and PMLS latencies differed systematically. For this analysis,

the PMLS and V1 latency distributions recorded in the same animal were compared against each

other using a rank-sum test, and the resulting p-values were Bonferroni corrected by the total

number of animals to account for multiple comparisons. Fig. 3E plots the corrected p-values as a

function of each animal's age, with the red line indicating $p=.01$. With the exception of one older

animal, significant differences between V1 and PMLS latencies using this criterion occurred only

before P35.

Taken together, our data show that in addition to age-dependent decreases in response

latencies in both areas, extra delays occur in PMLS in young animals, possibly reflecting changing

input sources and computational demands with age (see Discussion). Interestingly, for PMLS the

maturation of latency coincides with the maturation of DS and OS, while for V1 OS develops before

latency matures, and DS thereafter.

Visual experience accelerates DS development in both areas

An important aspect of the development of any visual function is the degree to which it reflects

endogenous mechanisms versus the impact of visual experience. In this regard, previous

experiments in the ferret have established a highly useful 'training' paradigm that allows controlled

tests of the impact of different types of visual experience [21]. In this paradigm, visually naïve animals

are exposed to a stimulus of interest for a prolonged period of time (while continuously under

anesthesia). Baseline tuning levels like DS are determined before and after stimulus exposure to

assess whether the chosen stimulus can exert a measurable influence over neural response

properties. For ferret V1, previous studies reported DS increases after exposure to a drifting grating

only – exposure to a gray screen was found to be ineffective, as was exposure to a stationary,

flashing grating (a grating with temporal modulation of its contrast) [21]. In identifying motion signals

as an important cue for V1 DS development, these results demonstrate that the paradigm is a

sensitive probe for which types of visual information might influence an area's development. For

this reason, we used it here to gain further insights into PMLS DS development.

Experiments using the training paradigm followed the procedures just described. More

precisely, we first determined pre-training DS levels in V1 and PMLS in visually naïve animals. We

then exposed the animals to a particular stimulus for 8 h. To minimize adaptation, the paradigm

was divided into 20 min long stimulation blocks alternating with 10 min of a blank screen (Fig. 4A).

Additionally, the stimulation block also consisted of alternating stimulus presentations (5 s) and

blank screen periods (10 s). At the conclusion of this training phase, DS levels were again measured

in V1 and PMLS and compared to the pre-training levels. Because of the duration of the

experiments, we did not attempt to track SU across pre- and post-training measurements. Instead,

we assessed DS at the MU level in each case as the more stable measure, reasoning that as

electrodes were not moved throughout the experiment, MU activity reflects activity in the same

population of neurons before and after exposure (see Methods for inclusion criteria for the training

data sets).

As a first test, we investigated the effect of motion cues on DS levels in both areas by

showing gratings drifting in one of 2 directions. Directions were chosen to be 180 deg apart from

each other so that the same orientation was shown throughout. Using this stimulus set, our data

replicated the previous findings of increased V1 DS levels after exposure. Fig. 4B plots the

cumulative L_{dir} distribution across MU sites before and after training, demonstrating a significant

increase in V1 DS levels after training (rank-sum test, p<.001). While this increase is consistent with

previous studies, the effect size appears to be significantly smaller in our animals than in those

earlier studies [21–24]. A possible explanation for this discrepancy is a difference in the amount of

natural visual experience per animal across studies. None of the animals studied here had naturally

opened their eyes at the start of the training experiments. In contrast, previous studies included

animals of the same age range, but with 1-3 days of visual experience ([22]; van Hooser, personal

communication). It is plausible that even brief amounts of natural visual experience can predispose

V1 circuits to be able to change more rapidly in response to the training stimuli, therefore resulting

in larger effects.

At the same time, since the effect size in our experiments is smaller, there might be a

concern that the population data mostly reflects changes in a single animal. In addition to

analyzing data across animals, we therefore also used a linear mixed-effect (LME) model to directly

control for grouping of data by animal. More precisely, we fit the before- and after-training L_{dir} data

with an LME model with a fixed effect for training phase (before/after training), allowing random

intercepts and slopes per animal (see Methods and Fig. S3A). On the full data set, the effect of

training phase indeed was not significant (p=.2). However, 4 out of the 5 animals showed an

increase in L_{dir} values after training (Table 8). If the 5[th] animal (which showed a decrease) is

excluded, the effect of training phase becomes significant (p=.009).

Importantly, our data show for the first time that exposure to drifting gratings also has a

measurable impact on PMLS DS levels. Combined across all animals, L_{dir} levels significantly

increased in PMLS with training (rank-sum test, p<.001). The LME model similarly returned a

significant effect of training phase (p<.001), and increases in L_{dir} values for all 5 animals (Table 8).

Thus, it appears that drifting gratings contain the necessary information to drive DS development

not just in V1, but also in PMLS.

To support this conclusion, it is important to compare the results to the effects of exposing

animals to a blank screen for 8 h, to determine whether the observed changes indeed were due to

information contained in the stimulus versus a more general effect of unspecific visual stimulation

(since the eyes are opened at the start of the experiment) or simply the passage of time. Fig. 4C and

S3B plot the L_{dir} changes observed before and after exposure to a homogenously gray screen for 8h

for both V1 and PMLS. Based on analyzing the population data irrespective of any per-animal

effects, V1 L_{dir} levels actually decreased during this time period (rank-sum test, p=.003). In PMLS, a

trend towards higher L_{dir} levels did not reach significance (rank-sum test, p=.01). The LME models,

which take per-animal effects into account, showed no significant change in the L_{dir} values for

either area (V1: p=.2, PMLS: p=.7), and positive and negative L_{dir} changes occurred with the same

frequencies across animals for both areas (Table 8). Thus, for PMLS as for V1, the changes induced

by the drifting grating indeed reflected the impact of the particular visual information presented

during the training, not unspecific effects.

PMLS DS development is sensitive to more visual cues than V1

The training experiments so far demonstrate that pattered visual experience can impact not just

V1 but also PMLS DS development. The next question then is whether certain types of visual

information are more important in this process than others. As mentioned above, exposure to a

training stimulus lacking motion signals proved ineffective in changing V1 DS levels. To probe

whether motion signals are similarly crucial for PMLS DS development, we performed a third set of

training experiments, in which animals were exposed to flashing gratings. These gratings were

generated by temporally modulating grating contrast between 0% and 100%. Importantly, no

contrast reversals were included. This ensured that while the stimulus contrast varied dynamically,

the spatial position of the stimulus remained fixed, and no motion cues were provided.

372   The results of these experiments are plotted in Fig. 4D and Fig. S3C. In agreement with the

earlier study by Li et al. [21], V1 L_{dir} levels remained unchanged after 8 h of exposure to the flashing

grating (rank-sum test, p=.9; LME model: p=.9, 3/6 animals with L_{dir} increases; Table 8). In contrast,

the same stimulus was highly effective in PMLS, with significant improvements in L_{dir} at the end of

the training experiment (rank-sum test, p<.001, LME model: p<.001, 7/7 animals with L_{dir} increases;

Table 8). Our data therefore strongly suggest that in contrast to V1, explicit motion cues were not

required for promoting DS development in PMLS. While the flashing gratings are stationary spatially,

they nonetheless dynamically changed over time. That means that of the spatiotemporal cues that

make up motion information, they at least contained the temporal component. Based on our data,

PMLS is sensitive to these temporal cues, while V1 is not (or to a much lesser degree). It remains to

be determined whether temporal cues alone are strong enough that a spatially uniform stimulus

that changes luminance in time, but lacks spatial information, would nonetheless be sufficient in

causing PMLS DS increases.

In addition to highlighting the role of different types of visual information, the differential

impact of flashing gratings on V1 and PMLS provides a striking example of dissociation in DS

development between the two areas reminiscent of the observations during normal development.

The training data generally provide an opportunity to further test this idea, because they can be

used to ask whether L_{dir} changes induced by the training protocol in V1 are correlated with L_{dir}

changes in PMLS. Fig. 4E plots the L_{dir} changes in V1 against the changes in PMLS for each animal

(with training effect sizes determined from the per-animal LME models; see Methods). Note that for

this plot, we only used animals in which data from both areas was available. The lack of correlation

of effect sizes is evident in this figure (r=-0.09, p=0.8), and supports the notion that DS development

in PMLS does not track V1 DS development.

As intriguing as the different effects of the flashing gratings on V1 and PMLS are, it needs to

be ruled out that the flashing gratings simply fail to activate V1, therefore rendering the stimulus

ineffective in driving L_{dir} change. To address this question, we turned to MU responses recorded

during the stimulus presentation periods in the training blocks. Responses to flashing gratings were

compared against those to drifting gratings, as a stimulus that could generate L_{dir} changes in V1.

Fig. 4F plots the average response time courses evoked by both types of stimuli for the two areas

(averaging across all blocks and all MU sites in all animals for the 2 training cohorts). These time

courses clearly show that flashing gratings did not fail to drive V1, but rather that both types of

stimuli evoked reliable activity in both areas. In fact, while peak rates differed between the two

areas (Fig. S4), they did not differ between stimulus types (two-way ANOVA with factors area and

stimulus type; area: p<.001, type: p=0.1, interaction: p=0.3; Table 10). The major difference

between the stimulus types appeared to be the duration of the evoked response (quantified as the

time with firing rates above baseline; see Methods), and as a consequence the average rates (Fig.

S4, Table 9). More precisely, response durations were significantly shorter for flashing than drifting

gratings in both areas (ANOVA with factors area and stimulus type, both main effects and

interaction p<.001; individual rank-sum tests per area: p<.001; Table 9). Interestingly, the difference

in response duration was more pronounced in V1 than PMLS. However, if lower V1 drive due to

shorter response durations indeed was the reason for a lack of DS change after training with

flashing gratings, then response duration should be correlated with the training-induced change in

L_{dir}. As is obvious from Fig. 4G, in V1 this is not the case for both stimuli combined or individually

(both conditions combined: r=-0.07, p=0.8; drifting: r=-0.7, p = 0.2; flashing: r=-0.3, p = 0.5). The

same is the case for PMLS (both conditions: r=-0.6, p=0.04; drifting: r=0.001, p=1; flashing: r=-0.5,

p=0.3). Thus, the different effect of exposure to flashing gratings on DS levels in V1 and PMLS

cannot simply be attributed to a failure to drive V1 response, but indeed appears to reflect

sensitivity to different stimulus attributes across areas.

Discussion

A key feature of the experiments presented here is the direct comparison of multiple stages of the

ferret's motion pathway. This network-level perspective reveals a complex picture of development,

one that does not agree with a simple feedforward sweep in which early stages develop before later

ones. Neither are our data consistent with an equally simple reversal of the hierarchy early in life, in

which PMLS becomes the primary driver. DS develops earlier in PMLS than in V1, in violation of a

strictly feedforward developmental sequence. Yet, PMLS OS does not develop early, and neither

does response latency. Paired with different sensitivity to visual cues, it is clear that the

developmental status of one area cannot be predicted by that of the other. An important issue to

resolve in the future is whether (and which) aspects of V1 and PMLS development indeed proceed

independently, and whether any independent development is supported through specific

(potentially age-dependent) input channels.

433   The latter is of particular interest because of the anatomical results for primate MT

development discussed above. Specifically, based on anatomical data marmoset MT develops

structurally as early as V1[34], consistent with our observation of the early functional development of

PMLS. This early MT development is supported by a transient pathway providing retinal input to MT

through the thalamic pulvinar, different from the standard flow of information through the lateral

geniculate nucleus (LGN) and V1 to MT found in adults [35,36]. Whether the input circuits to PMLS

undergo similar changes remains to be investigated. The latency data are interesting in this respect:

Response latencies in both areas shorten with age, something that has also been observed for

monkey V1 and V2 and might suggest general circuit maturation [38]. At the same time, while there is

considerable overlap in V1 and PMLS latencies in adults (consistent with data for primate V1 and

MT[37,39,40]), up to P35 PMLS latencies are significantly slower than in V1. An intriguing hypothesis then

is that the extra delay reflects the cost incurred by driving PMLS activity through different input

sources compared to adults (interesting candidates include LGN, pulvinar or superior colliculus).

Since DS is sparse at least in ferret LGN from eye opening on [41,42], the extra delay may also reflect

that different input sources require PMLS to compute direction signals *de novo*, rather than relying

on 'precomputed' direction signals provided by V1. Finally, it is worth mentioning that around eye

opening, PMLS provides the largest fraction of feedback to V1, before closer areas like area 18

become dominant later in life [43]. While further experiments are needed to answer these questions,

our data, in conjunction with developmental data from primates, nonetheless strongly argue that

development of the visual hierarchy is best described as a dynamic sequence of unique networks

that can be configured quite differently from the adult, rather than a gradual, layer-by-layer

emergence of the adult hierarchy [29].

One of the unexpected results of our study are the differences in sensitivity to visual cues

between V1 and PMLS. Motion computations require the joint processing of spatial and temporal

signals. Clearly, around eye opening V1 requires stimuli that provide full spatiotemporal cues to

develop DS. It is possible that this changes in slightly older animals, as the role of visual experience

switches from instructive (building circuits that reflect the characteristics of the visual input) to

permissive (enabling circuit development) around eye opening[24]. Here, animals had not opened

their eyes at the time of recording. However, previous studies similarly showing a lack of DS

development after exposure to flashing gratings were done in animals with 1-3 days of visual

experience[21], therefore suggesting that temporal information in isolation remains ineffective even

after visual experience switches to a permissive role. In PMLS, though, the same purely temporal

inputs are sufficient in driving DS development, and at least for the specific stimulus properties and

animals studied here, might actually be more effective than stimuli with joint spatial and temporal

information. On the one hand, this difference in sensitivity could mean that altered visual

experience during development could have very different consequences for PMLS and V1 function.

On the other hand, it is interesting to speculate about what this sensitivity to temporal cues means

for the impact of normal visual experience on PMLS before eye opening. Previous studies have

found that reliable visual responses can be recorded in ferret LGN and V1 starting around P17[44,45].

At these ages, ferret kits have been found to make scanning head movements from side to side[45],

and at least based on neural responses, the corresponding visual signals, even through the closed

eye lids, contain enough structure to result in measurable OS and DS[44]. Nonetheless, the filter

applied by the eye lids will likely remove spatial structure to a much larger degree than temporal

transients. Thus, natural input before eye opening might contain more of the relevant cues for PMLS

than V1 DS development.

In addition to these insights into visual hierarchy development, our data may also provide

valuable clues regarding motion processing in adult cortex. One important observation in this

respect is the relative developmental time course of OS and DS. In V1, we and others observe OS

development before DS[17,18]. At least in the ferret, this sequence is reflected in the functional

organization of the area: Orientation maps develop first, and show minimal discontinuities (singular

points or pinwheels) [17,33,46]. The later developing direction maps are then constrained by the existing

orientation maps. As a consequence, they are nestled inside the orientation maps, with each

orientation column split into subregions selective for opposite directions of motion (that share the

same orientation). As a result, the discontinuities for the direction map take the form of linear

fractures at which directions shift by 180 deg [46]. In PMLS, DS and OS mature at the same time. It is

therefore possible that the functional architecture is optimized for direction rather than orientation,

a question to be resolved in future imaging studies. In monkey MT, extracellular recordings show

that neighboring neurons either have very similar direction preferences or differ by nearly 180 deg [47],

which would be most consistent with a direction map with linear fractures. Indeed, optical imaging

from owl monkey MT shows orientation and direction maps very similar to those found in ferret V1

[48]. Yet, recent two-photon data from the marmoset reported predominantly smooth changes in

direction preference, rather than sudden 180 deg reversals, at least at the scale of several hundred

micrometers [49]. Whether PMLS/MT organization is optimized for direction or orientation therefore

remains an open question.

Another observation of interest for adult PMLS function is the fact that concurrent with the

increase in DS, the response to the null direction decreases. Null direction inhibition is thought to

play a role in the generation of DS in primate MT [50–52], and we have suggested the same might be the

case in ferret PMLS [16]. While the anatomical and functional development of inhibitory neurons in

ferret V1 has received some attention (e.g., [20,53–55]), there are currently no comparable data for

PMLS. Further studies are therefore necessary to resolve whether the decreased null direction

response is indeed the consequence of the development of selective inhibition (rather than e.g. a

change in feedforward drive).

Finally, the complex network-level development we observed will be important to consider

in the context of both developmental disorders and plasticity mechanisms in the adult brain. Early

development, paired with sensitivity to a broad range of visual cues, likely means increased

vulnerability to disruptions of normal development early in life – consistent with the observation

that motion vision is impaired in a range of developmental disorders (see e.g. [9,29] for reviews). On

the other hand, the circuit mechanisms that support at least the early DS development may still be

available for reactivation later in life and provide mechanisms for phenomena like blindsight [56,57] or

the ability to regain some visual function after V1 stroke through training on a motion task [58].

References

1.   Borst, A. & Groschner, L. N. How Flies See Motion. *Annu. Rev. Neurosci.* **46**, 17–37 (2023).

2.   Bollmann, J. H. The Zebrafish Visual System: From Circuits to Behavior. *Annu. Rev. Vis. Sci.* **5**,

269–293 (2019).

3.   Wei, W. & Feller, M. B. Organization and development of direction-selective circuits in the

retina. *Trends Neurosci.* **34**, 638–645 (2011).

4.   Vaney, D. I., Sivyer, B. & Taylor, W. R. Direction selectivity in the retina: symmetry and

asymmetry in structure and function. *Nat. Rev. Neurosci.* **13**, 194–208 (2012).

5.   Niell, C. M. & Scanziani, M. How Cortical Circuits Implement Cortical Computations: Mouse

Visual Cortex as a Model. *Annu. Rev. Neurosci.* **44**, 517–546 (2021).

6.   Born, R. T. & Bradley, D. Structure and Function of Visual Area MT. *Annu. Rev. Neurosci.* **28**, 157–

189 (2005).

7.   Orban, G. Higher Order Visual Processing in Macaque Extrastriate Cortex. *Physiol. Rev.* **88**, 59–

89 (2008).

8.   Chaplin, T. A., Rosa, M. G. P. & Lui, L. L. Auditory and Visual Motion Processing and Integration

in the Primate Cerebral Cortex. *Front. Neural Circuits* **12**, (2018).

9.   Atkinson, J. The Davida Teller Award Lecture, 2016: Visual Brain Development: A review of

"Dorsal Stream Vulnerability"—motion, mathematics, amblyopia, actions, and attention. *J. Vis.*

17, 26 (2017).

10. Hubel, D. H. & Wiesel, T. N. Receptive fields and functional architecture of monkey striate

cortex. *J. Physiol.* **195**, 215–43 (1968).

11. De Valois, R. L., Yund, E. W. & Hepler, N. The orientation and direction selectivity of cells in

macaque visual cortex. *Vision Res.* **22**, 531–544 (1982).

12. Dubner, R. & Zeki, S. M. Response properties and receptive fields of cells in an anatomically

defined region of the superior temporal sulcus in the monkey. *Brain Res.* **35**, 528–532 (1971).

13. Maunsell, J. & Van Essen, D. C. Functional properties of neurons in middle temporal visual area

of the macaque monkey. I. Selectivity for stimulus direction, speed, and orientation. *J.*

Neurophysiol. **49**, 1127–1147 (1983).

14. Albright, T. Direction and orientation selectivity of neurons in visual area MT of the macaque. *J.*

Neurophysiol. **52**, 1106–30 (1984).

15. Philipp, R., Distler, C. & Hoffmann, K.-P. A motion-sensitive area in ferret extrastriate visual

cortex: an analysis in pigmented and albino animals. *Cereb. Cortex* **16**, 779–90 (2006).

16. Lempel, A. A. & Nielsen, K. J. Ferrets as a Model for Higher-Level Visual Motion Processing.

Curr. Biol. **29**, 179–191 (2019).

17. Li, Y., Fitzpatrick, D. & White, L. E. The development of direction selectivity in ferret visual cortex

requires early visual experience. *Nat. Neurosci*. **9**, 676–681 (2006).

18. Clemens, J. M., Ritter, N. J., Roy, A., Miller, J. M. & Van Hooser, S. D. The Laminar Development

of Direction Selectivity in Ferret Visual Cortex. *J. Neurosci*. **32**, 18177–18185 (2012).

19. Smith, G. B. *et al*. The development of cortical circuits for motion discrimination. *Nat. Neurosci*.

18, 252–261 (2015).

20. Chang, J. T. & Fitzpatrick, D. Development of visual response selectivity in cortical GABAergic

interneurons. *Nat. Commun*. **13**, 3791 (2022).

21. Li, Y., Van Hooser, S. D., Mazurek, M., White, L. E. & Fitzpatrick, D. Experience with moving

visual stimuli drives the early development of cortical direction selectivity. *Nature* **456**, 952–6

(2008).

22. Van Hooser, S. D. *et al*. Initial neighborhood biases and the quality of motion stimulation jointly

influence the rapid emergence of direction preference in visual cortex. *J. Neurosci*. **32**, 7258–66

(2012).

23. Ritter, N. J., Anderson, N. M. & Van Hooser, S. D. Visual Stimulus Speed Does Not Influence the

Rapid Emergence of Direction Selectivity in Ferret Visual Cortex. *J. Neurosci*. **37**, 1557–1567

(2017).

24. Roy, A., Wang, S., Meschede-Krasa, B., Breffle, J. & Van Hooser, S. D. An early phase of

instructive plasticity before the typical onset of sensory experience. *Nat. Commun*. **11**, 11

(2020).

25. Spear, P. D., Tong, L., McCall, M. A. & Pasternak, T. Developmentally induced loss of direction-

selective neurons in the cat's lateral suprasylvian visual cortex. *Dev. Brain Res*. **20**, 281–285

(1985).

26. El-Shamayleh, Y., Kiorpes, L., Kohn, A. & Movshon, J. A. Visual Motion Processing by Neurons in

Area MT of Macaque Monkeys with Experimental Amblyopia. *J. Neurosci.* **30**, 12198–12209

(2010).

27. Grootel, T. J. V., Raghavan, R. T., Kelly, J. G., Movshon, J. A. & Kiorpes, L. Responses to visual

motion of neurons in the extrastriate visual cortex of macaque monkeys with experimental

amblyopia. 2024.07.01.601564 Preprint at https://doi.org/10.1101/2024.07.01.601564 (2024).

28. Lempel, A. A. & Nielsen, K. J. Development of visual motion integration involves coordination of

multiple cortical stages. *eLife* **10**, e59798 (2021).

29. Bourne, J. A. *et al.* Development of Higher-Level Vision: A Network Perspective. *J. Neurosci.* **44**,

e1291242024 (2024).

30. Mazurek, M., Kager, M. & Van Hooser, S. D. Robust quantification of orientation selectivity and

direction selectivity. *Front. Neural Circuits* **8**, (2014).

31. Batschelet, E. *Circular Statistics in Biology*. (Academic Press, New York, NY, US, 1981).

32. Chapman, B. & Stryker, M. P. Development of orientation selectivity in ferret visual cortex and

effects of deprivation. *J. Neurosci.* **13**, 5251 (1993).

33. Chapman, B., Stryker, M. P. & Bonhoeffer, T. Development of orientation preference maps in

ferret primary visual cortex. *J. Neurosci.* **16**, 6443–53 (1996).

34. Bourne, J. A. & Rosa, M. G. P. Hierarchical development of the primate visual cortex, as revealed

by neurofilament immunoreactivity: early maturation of the middle temporal area (MT). *Cereb.*

Cortex **16**, 405–14 (2006).

35. Warner, C. E., Kwan, W. C. & Bourne, J. A. The Early Maturation of Visual Cortical Area MT is

Dependent on Input from the Retinorecipient Medial Portion of the Inferior Pulvinar. *J. Neurosci.*

32, 17073–17085 (2012).

36. Mundinano, I.-C. *et al.* Transient visual pathway critical for normal development of primate

grasping behavior. *Proc. Natl. Acad. Sci.* 201717016 (2018) doi:10.1073/pnas.1717016115.

37. Raiguel, S. E., Xiao, D.-K., Marcar, V. L. & Orban, G. A. Response Latency of Macaque Area

MT/V5 Neurons and Its Relationship to Stimulus Parameters. *J. Neurophysiol.* **82**, 1944–1956

(1999).

38. Zhang, B., Smith, E. L. & Chino, Y. M. Postnatal development of onset transient responses in

macaque V1 and V2 neurons. *J. Neurophysiol.* **100**, 1476–87 (2008).

39. Raiguel, S. E., Lagae, L., Gulyàs, B. & Orban, G. A. Response latencies of visual cells in

macaque areas V1, V2 and V5. *Brain Res.* **493**, 155–159 (1989).

40. Schmolesky, M. T. *et al.* Signal Timing Across the Macaque Visual System. *J. Neurophysiol.* **79**,
3272–3278 (1998).

41. Murphy, A. J., Hasse, J. M. & Briggs, F. Physiological characterization of a rare subpopulation of
doublet-spiking neurons in the ferret lateral geniculate nucleus. *J. Neurophysiol.* **124**, 432–442
(2020).

42. Stacy, A. K., Schneider, N. A., Gilman, N. K. & Van Hooser, S. D. Impact of Acute Visual
Experience on Development of LGN Receptive Fields in the Ferret. *J. Neurosci.* **43**, 3495–3508
(2023).

43. Khalil, R. & Levitt, J. B. Developmental remodeling of corticocortical feedback circuits in ferret
visual cortex. *J. Comp. Neurol.* **522**, 3208–3228 (2014).

44. Krug, K., Akerman, C. J. & Thompson, I. D. Responses of Neurons in Neonatal Cortex and
Thalamus to Patterned Visual Stimulation Through the Naturally Closed Lids. *J. Neurophysiol.*
85, 1436–1443 (2001).

45. Akerman, C. J., Smyth, D. & Thompson, I. D. Visual Experience before Eye-Opening and the
Development of the Retinogeniculate Pathway. *Neuron* **36**, 869–879 (2002).

46. Weliky, M., Bosking, W. & Fitzpatrick, D. A systematic map of direction preference in primary
visual cortex. *Nature* **379**, 725–728 (1996).

47. Albright, T., Desimone, R. & Gross, C. G. Columnar organization of directionally selective cells
in visual area MT of the macaque. *J. Neurophysiol.* **51**, 16–31 (1984).

48. Malonek, D., Tootell, R. B. H. & Grinvald, A. Optical Imaging Reveals the Functional Architecture
of Neurons Processing Shape and Motion in Owl Monkey Area MT. *Proc. R. Soc. Lond. B Biol.*
Sci. **258**, 109–119 (1994).

49. Pattadkal, J. J., Zemelman, B. V., Fiete, I. & Priebe, N. J. Primate neocortex performs balanced
sensory amplification. *Neuron* **112**, 661-675.e7 (2024).

50. Livingstone, M. S., Pack, C. C. & Born, R. T. Two-Dimensional Substructure of MT Receptive
Fields. *Neuron* **30**, 781–793 (2001).

51. Thiele, A., Distler, C., Korbmacher, H. & Hoffmann, K.-P. Contribution of inhibitory mechanisms
to direction selectivity and response normalization in macaque middle temporal area. *Proc.*
Natl. Acad. Sci. **101**, 9810–9815 (2004).

52. Thiele, A., Herrero, J. L., Distler, C. & Hoffmann, K.-P. Contribution of Cholinergic and GABAergic
Mechanisms to Direction Tuning, Discriminability, Response Reliability, and Neuronal Rate
Correlations in Macaque Middle Temporal Area. *J. Neurosci.* **32**, 16602–16615 (2012).

53. Gao, W. J., Newman, D. E., Wormington, A. B. & Pallas, S. L. Development of inhibitory circuitry
in visual and auditory cortex of postnatal ferrets: immunocytochemical localization of
GABAergic neurons. *J. Comp. Neurol.* **409**, 261–73 (1999).
54. Gao, W. J., Wormington, A. B., Newman, D. E. & Pallas, S. L. Development of inhibitory circuitry
in visual and auditory cortex of postnatal ferrets: Immunocytochemical localization of
calbindin- and parvalbumin-containing neurons. *J. Comp. Neurol.* **422**, 140–157 (2000).
55. Mulholland, H. N., Hein, B., Kaschube, M. & Smith, G. B. Tightly coupled inhibitory and
excitatory functional networks in the developing primary visual cortex. *eLife* **10**, e72456 (2021).
56. Weiskrantz, L., Barbur, J. L. & Sahraie, A. Parameters affecting conscious versus unconscious
visual discrimination with damage to the visual cortex (V1). *Proc. Natl. Acad. Sci.* **92**, 6122–
6126 (1995).
57. Weiskrantz, L. Blindsight revisited. *Curr. Opin. Neurobiol.* **6**, 215–220 (1996).
58. Huxlin, K. R. *et al.* Perceptual Relearning of Complex Visual Motion after V1 Damage in
Humans. *J. Neurosci.* **29**, 3981–3991 (2009).
59. Du, J., Blanche, T. J., Harrison, R. R., Lester, H. A. & Masmanidis, S. C. Multiplexed, High Density
Electrophysiology with Nanofabricated Neural Probes. *PLOS ONE* **6**, e26204 (2011).
60. nielsenlabmbi. Spike. (2024).
61. Brainard, D. H. The Psychophysics Toolbox. *Spat. Vis.* **10**, 433–436 (1997).
62. Pelli, D. G. The VideoToolbox software for visual psychophysics: transforming numbers into
movies. *Spat. Vis.* **10**, 437–442 (1997).
63. Ellaway, P. H. Cumulative sum technique and its application to the analysis of peristimulus
time histograms. *Electroencephalogr. Clin. Neurophysiol.* **45**, 302–304 (1978).

658
659

Methods
Animal preparation and surgery
All procedures adhered to the guidelines of the national Institute of Health and were approved by
the Animal Care and Use Committee at Johns Hopkins University. Experiments were performed in
male and female ferrets (*Mustela putoris furo*) aged 29 – 52 days. Animals used in the training
experiments were 28- - 33 days old; none of these animals had naturally opened their eyes at the
start of the experiment. Recordings were performed in anesthetized ferrets following established
procedures described previously [16]. Briefly, animals were anesthetized using isoflurane (during

surgery: 1.5-3%, during recordings: 0.5-2%), and paralyzed using pancuronium bromide (0.15

mg/kg/hr) to prevent eye movements. A range of vital parameters (heart rate, SpO2, EtCO2, EEG)

were monitored continuously to ensure adequate anesthetic depth during the experiments.

Neosynephrine and atropine were applied to the eyes to retract the nictitating membranes and

dilate the pupils. Eyes were then covered with contact lenses. Craniotomies were made over V1

and PMLS, targeting related visual field positions.

Extracellular recordings and extraction of neural responses

Neural responses were recorded using multichannel silicon probes obtained from the Masmanidis

lab at UCLA [59], plated with gold to reach impedances around 150-300 kΩ. Signals from one or two

probes were recorded with a RHD2000 amplifier (intan Technologies) using sampling rates of 20-30

kHz. Data was filtered offline between 250 Hz and 5kHz. Based on these filtered signals, two types

of neural responses were extracted. Single units were isolated using a custom-written processing

pipeline in Matlab (MathWorks) for manual spike sorting (available on github [60]). In this pipeline,

detection thresholds were first set manually for each recording channel based on noise levels.

Spikes detected on channels with distances up to 50um that occurred within +/- 0.2 ms of each

other were assigned to the channel with the largest energy and eliminated from other channels.

Spikes were then manually clustered using multiple waveform characteristics like through

amplitude, energy, or trough-peak amplitude, as well as recording position on the probe

(determined as center-of-mass of the spike energy or trough amplitude computed across

neighboring probe channels). Quality of isolation was confirmed by analyzing inter-spike intervals

and eliminating any units with intervals below 1.2 ms. Multi-unit signals were extracted by first

applying an automatically computed detection threshold to every channel. The threshold was set to

4 times the standard deviation of each channel's bandpass-filtered signal, using the median

absolute deviation scaled for a normal distribution to estimate the standard deviation. Again,

events detected within +/-0.2 ms on channels within 50 um were assigned only to the channel with

the largest energy. No further spike sorting was performed on this multi-unit signal.

Visual stimuli and experiment design

Visual stimulus generation: Stimuli were generated using the Psychophysics Toolbox extension to

Matlab [61,62] and displayed on a 23 inch LCD monitor with a refresh rate of 120 Hz, placed 20 – 35 cm

in front of the ferret. The monitor was gamma corrected using a SpectraScan 655 (PhotoResearch).

Stimuli for selectivity measurements: Except for the stimulation blocks of the exposure paradigm,

all data was collected by presenting gratings drifting in 12 or 16 directions. Gratings were shown

full-field (about 65 x 50 deg), with the exception of 3 experiments that also presented an additional

smaller size. For those experiments, only the full-field data was analyzed. With the exception of 2

experiments (ages P45 & P48) that used a sine-wave profile, gratings were square-wave gratings

shown at 100% contrast. Spatial and temporal frequency were optimized per recording, and ranged

between 0.01 – 0.1 cyc/deg and 1 - 6 cyc/s, respectively. In addition to the different directions, each

experiment also contained a blank condition. All conditions were shown in a pseudo-random

sequence with 5 repetitions per condition. Within each condition, each presentation of a stimulus

was preceded and followed by presentation of a gray screen of equal luminance. Stimuli were

shown for 1 s, with inter-stimulus intervals of 3 - 10 s (longer intervals were needed for the youngest

animals in which adaptation to the stimulus is more pronounced).

Stimuli during training paradigm: For the animals exposed to drifting or flashing stimuli, we

presented full-field, square-wave gratings during the stimulation blocks. Stimuli were shown for 5 s,

with 10 s inter-stimulus intervals. Spatial frequency was again adjusted based on responses in an

individual animal, and ranged between 0.05 – 0.08 cyc/deg for both groups. For the drifting gratings,

temporal frequency ranged between 3 – 6 cyc/deg. Drifting gratings were always shown at 100%

contrast. For the flashing gratings, we modulated contrast between 0 and 100% (no contrast

reversal) using a sinusoidal function with a temporal frequency of 3 Hz.

Data analysis

<u>Inclusion criteria and data set size</u>

For the analysis of SU responses, neural responses to every stimulus were calculated as the firing

rate during the stimulus presentation period (1s long) minus the firing rate during the last second of

the pre-stimulus period. SU needed to pass two criteria to be included in the analyses. First, we

computed an ANOVA across the responses in all stimulus conditions, including the blank. Only SU

with $p < .05$ for the ANOVA were included. Additionally, the average firing rate for the best condition

had to exceed 0 Hz.

For the analysis of response latency using MU data, we used the same approach to test for

visual responsiveness (ANOVA for the baseline-corrected firing rates with $p < .05$, firing rate for best

stimulus > 0Hz). Additionally, the data set as limited to simultaneous recordings from V1 and PMLS.

This MU data set includes the same data used for SU analysis of DS development, but contains

additional data in which no SU could be isolated or in which no SU passed the criteria.

MU data was also used to analyze the effects of the exposure paradigm. In this case, the

long duration of the experiments generally made responses more variable, which required an

adjustment in the inclusion criteria. For these data, we tested for visual responsiveness by

comparing the stimulus-evoked firing rates (not baseline corrected) to the spontaneous activity

during the 1s preceding each stimulus presentation (blanks were excluded from this analysis). MU

with p<.05 and with responses to the preferred stimulus exceeding 0Hz were included in

subsequent analyses. Further, data were restricted to experiments in which visually responses sites

could be identified before and after the exposure. This meant that for a few animals only PMLS data

(1 animal in the flashing and gray cohort each) or V1 data (1 animal in the gray cohort) was

available. These cases were eliminated when comparing exposure effect sizes between V1 and

PMLS (Fig, 4E, G). Finally, no responsiveness criterion was applied to the MU collected during the

exposure blocks. The resulting number of animals and neurons for normal development and

exposure experiments are listed in Table 1.

<u>Measurements of DS and OS</u>

DS was quantified using 2 metrics, L_{dir} and DSI. L_{dir} was computed as [30]

751
$$L_{dir} = \left| \frac{\sum_k R(\theta_k) \exp(i\theta_k)}{\sum_k R(\theta_k)} \right|,$$

752   where $R(\theta_k)$ represents the response to angle θ_k. DSI was computed as

753
$$DSI = 1 - \frac{R(N)}{R(P)},$$

754   where R(P) is the response to the preferred and R(N) the response to the null direction. Matching

755   metrics were computed for OS:

756
$$L_{ori} = \left| \frac{\sum_k R(\theta_k) \exp(2i\theta_k)}{\sum_k R(\theta_k)} \right| \text{ and } OSI = 1 - \frac{R(O)}{R(P)},$$

with R(O) the response to the orthogonal direction.

<u>Normalized DS and OS metrics</u>

To compare development of DS and OS across areas without the effects of different selectivity

levels in adults, normalized L_{dir} and L_{ori} values were computed. For this analysis, SU were first

binned into 6 age windows, P29-30, P31-32, P33-34, P35-36, P37-39, P44-45. The age window from

P37-39 was expanded to include a single V1 animal at P39. We then computed the average L_{dir} value

for the first and last bin. All L_{dir} values were then normalized by computing

765
$$norm\ L_{dir} = \frac{L_{dir} - \bar{L}_{dir,EO}}{\bar{L}_{dir,adult} - \bar{L}_{dir,EO}}$$

where $\bar{L}_{dir,EO}$ refers to the average L_{dir} value in the first age group around eye opening, and $\bar{L}_{dir,adult}$

to the average value in the last, adult group. The same analysis was repeated to compute

normalized L_{ori} values.

<u>Measurements of latency</u>

Latency was computed for each MU site based on the responses to its best condition. Latencies

were computed using cumulative sum analysis [63] using similar statistical criteria as used previously

for primate MT neurons [37]. First, responses were binned into 10 ms bins to first compute the PSTH.

Second, mean and standard deviation of the baseline rate were computed for the 1 s baseline

preceding stimulus onset, and the baseline mean was subtracted from the PSTH before computing

the cumulative sum of the PSTH. The latency was then taken as the first bin after stimulus onset in

which the cumulative PSTH was larger than two standard deviations (of the baseline rate), followed

by 2 increasing bins. For the analysis of latency development in shorter age windows, data from V1

and PMLS was binned into the following age windows: P29-30, P31-32, P33-34, P35-36, P37-39,

P42-52. These are the same age windows used for the normalized DS and OS metric, but the adult

window was expanded to include more animals in which paired V1 and PMLS data were available.

<u>Analysis of MU responses in stimulation blocks</u>

To analyze response patterns during the stimulation blocks, responses were computed during the 5

s of stimulus presentation. To compute average rate, the firing rate was computed for each 5 s

stimulus presentation period, baseline corrected by the firing rate in the preceding 2 s baseline

period, and averaged across all stimulus presentation periods. To compute peak rate and response

duration, we first computed the PSTH across all stimulus presentation periods using 10 ms bins.

The peak rate was set to the maximum of the PSTH, baseline corrected as before. Duration was

quantified as the number of bins during the stimulus presentation exceeding the baseline rate by 2

standard deviations, converted to seconds. Finally, all metrics were averaged across all blocks for

each channel. This assumes that there are no significant movements of the probe throughout the

experiment. Treating each block as an independent measurement (no averaging across blocks)

does not affect the results.

<u>Statistics</u>

For all normal development data, we first used an ANOVA or Kruskall-Wallis test to establish high-

level differences between groups or areas (see Tables 2-6). Post-hoc tests (where justified) were

then performed using Wilcoxon rank-sum tests. The outcome of all statistical tests is listed in

Tables 2-6. For the training data, we performed two different kinds of analyses. First, we compared

before and after training data, combined across all animals, for each area using a rank-sum test.

Second, we fit the L_{dir} MU data with a LME model (lmefit, Matlab), with training phase (before/after

exposure to the stimulus) as fixed effect and by-animal random intercepts and slopes. We

compared this model against one omitting the by-animal random slopes; in most instances, the

model including random slopes outperformed the one without (based on likelihood-ratio

comparisons), so for consistency, we used the model with random slopes in all cases. Models were

fit using maximum likelihood estimation, and the significance of the fixed effect was assessed

using an ANOVA. Effect size per animal was computed as the sum of the fixed effect slope and the

random, per-animal slope.

Figure 1

Figure 2

Figure 3

Figure 4

Tables

Figure	Exp. Group	N Animals	N SU/MU
1D & 2A, S1A & B	PMLS	24	128 (SU)
1D & 2A, 3, S1A & B	V1	25	140 (SU)
1E, H & J, 2C, S1C & D	PMLS, P29-32	10	35 (SU)
1E, H & J, 2C, S1C & D	PMLS, P33-36	6	22 (SU)
1E, H & J, 2C, S1C & D	PMLS, P37-52	8	71 (SU)
1F, I & K, 2D, S1E & F	V1, P29-32	12	42 (SU)
1F, I & K, 2D, S1E & F	V1, P33-36	6	45 (SU)
1F, I & K, 2D, S1E & F	V1, P37-52	7	53 (SU)
1G & 2B, S2A	PMLS	2-6	8-28 per age group (SU)
1G & 2B, S2A	V1	2-7	15-23 per age group (SU)
3B, E	PMLS	30	1065 (MU)
3B, E	V1	30	1904 (MU)
3C	PMLS, P29-32	17	439 (MU)
3C	PMLS, P33-36	6	337 (MU)
3C	PMLS, P37-52	7	289 (MU)
3C	V1, P29-32	17	1006 (MU)
3C	V1, P33-36	6	464 (MU)
3C	V1, P37-52	7	436 (MU)
3D	PMLS	2-9	95-242 per age group (MU)
3D	V1	2-9	121-517 per age group (MU)
4B	PMLS, before	5	297 (MU)
4B	PMLS, after	5	297 (MU)
4B	V1, before	5	320 (MU)
4B	V1, after	5	312 (MU)
4C	PMLS, before	7	404 (MU)
4C	PMLS, after	7	380 (MU)
4C	V1, before	6	382 (MU)
4C	V1, after	6	383 (MU)
4D	PMLS, before	5	344 (MU)
4D	PMLS, after	5	463 (MU)
4D	V1, before	5	278 (MU)
4D	V1, after	5	467 (MU)
4E, G	PMLS	4-6	47-237 per animal (MU)
4E, G	V1	4-6	31-114 per animal (MU)
4F	V1, drifting	5	320 (MU)
4F	V1, flashing	6	384 (MU)
4F	PMLS, drifting	5	319 (MU)
4F	PMLS, flashing	6	384 (MU)
4G	V1, resp. dur.	5-6	64 per animal (MU)
4G	PMLS, resp. dur.	5-6	63-64 per animal (MU)

Table 1: Data set sizes for all figures and analyses

Metric	Group	Test	PMLS p value	PMLS other	V1 p value	V1 other
L_{dir}	All age groups	Kruskall-Wallis	<0.001	X(2)=56.5	<0.001	X(2)=30.2
L_{dir}	P29-32 vs P33-36	Rank sum	<0.001		0.3	
L_{dir}	P29-32 vs P37-52	Rank sum	<0.001		<0.001	
L_{dir}	P33-36 vs P37-52	Rank sum	0.2		<0.001	
Pref.	All age groups	Kruskall-Wallis	<0.001	X(2)=14.5	<0.001	X(2)=34.3
Pref.	P29-32 vs P33-36	Rank sum	0.02		<0.001	
Pref.	P29-32 vs P37-52	Rank sum	<0.001		<0.001	
Pref.	P33-36 vs P37-52	Rank sum	0.5		0.03	
Null	All age groups	Kruskall-Wallis	<0.001	X(2)=21	0.008	X(2)=9.6
Null	P29-32 vs P33-36	Rank sum	0.02		0.01	
Null	P29-32 vs P37-52	Rank sum	<0.001		0.6	
Null	P33-36 vs P37-52	Rank sum	0.5		0.006	

Table 2: Statistics for Figs. 1E, F, H-K

Metric	Group	Test	p value	Other
Norm. L_{dir}	V1 vs PMLS	ANOVA: area, age	Age: <0.001 Area: <0.001 IA: 0.003	Age: F(3)=19.6 Area: F(1)=18.1 IA: F(3)=4.9
Norm. L_{dir}	V1 vs PMLS, P31-32	Rank sum	0.52	
Norm. L_{dir}	V1 vs PMLS, P33-34	Rank sum	<0.001	
Norm. L_{dir}	V1 vs PMLS, P35-36	Rank sum	0.02	
Norm. L_{dir}	V1 vs PMLS, P37-39	Rank sum	0.14	

Table 3: Statistics for Figs. 1G (IA: Interaction)

Metric	Group	Test	PMLS p value	PMLS other	V1 p value	V1 other
L_{ori}	All age groups	Kruskall-Wallis	<0.001	X(2)=24.9	0.004	X(2)=10.9
L_{ori}	P29-32 vs P33-36	Rank sum	<0.001		0.009	
L_{ori}	P29-32 vs P37-52	Rank sum	<0.001		0.002	
L_{ori}	P33-36 vs P37-52	Rank sum	0.9		0.6	

Table 4: Statistics for Figs. 2B & C

Metric	Group	Test	p value	Other
Norm. L_{ori}	V1 vs PMLS	ANOVA: area, age	Age: <0.001 Area: 0.07 IA: 0.003	Age: F(3)=8.9 Area: F(1)=3.4 IA: F(3)=4.8
Norm. L_{ori}	V1 vs PMLS, P31-32	Rank sum	<0.001	
Norm. L_{ori}	V1 vs PMLS, P33-34	Rank sum	0.6	
Norm. L_{ori}	V1 vs PMLS, P35-36	Rank sum	0.95	
Norm. L_{ori}	V1 vs PMLS, P37-39	Rank sum	0.9	

Table 5: Statistics for Figs. 2D (IA: Interaction)

Metric	Group	Test	PMLS p value	PMLS other	V1 p value	V1 other
Latency	All age groups	Kruskall-Wallis	<0.001	X(2)=468.3	<0.001	X(2)=816.4
Latency	P29-32 vs P33-36	Rank sum	<0.001		<0.001	
Latency	P29-32 vs P37-52	Rank sum	<0.001		<0.001	
Latency	P33-36 vs P37-52	Rank sum	<0.001		<0.001	

Table 6: Statistics for Fig. 3C

Group	Test	p value	Other
All age groups	ANOVA: area, age	Area: <0.001 Age: <0.001 IA: <0.001	Area: F(1)=228.4 Age: F(5)=354.2 IA: F(5)=35.8
V1 vs PMLS, P29-30	Rank sum	<0.001	
V1 vs PMLS, P31-32	Rank sum	<0.001	
V1 vs PMLS, P33-34	Rank sum	<0.001	
V1 vs PMLS, P35-36	Rank sum	0.05	
V1 vs PMLS, P37-39	Rank sum	0.03	
V1 vs PMLS, P40-52	Rank sum	0.02	

Table 7: Statistics for Fig. 3D (IA: interaction)

Training stim.	Parameter	V1	PMLS
Drifting	Phase: Estimate	Full data: 0.02 Reduced data: 0.04	0.03
Drifting	Phase: p value	Full data: 0.2 Reduced data: 0.009	<0.001
Drifting	Slope per animal	0.02, 0.09, -0.03, 0.02, 0.03	0.04, 0.03, 0.05, 0.03, 0.01
Gray	Phase: Estimate	-0.03	0.01
Gray	Phase: p value	0.2	0.7
Gray	Slope per animal	-0.1, 0.01, -0.02, 0.03, -0.08	0.09, -0.07, -0.002, -0.03, 0.07
Flashing	Phase: Estimate	0.0008	0.09
Flashing	Phase: p value	0.9	<0.001
Flashing	Slope per animal	-0.02, 0.02, 0.02, -0.01, -0.01, 0.006	0.09, 0.11, 0.10, 0.08, 0.12, 0.06, 0.09

Table 8: LME results for Fig. 4B-D. Phase: Slope for the fixed effect training phase (before/after); slope: fixed effect slope + random effect slope per animal.

Metric	Group	Test	PMLS p value	PMLS other	V1 p value	V1 other
DSI	All age groups	Kruskall-Wallis	<0.001	X=38.7	<0.001	X=27.3
DSI	P29-32 vs P33-36	Rank sum	<0.001		0.6	
DSI	P29-32 vs P37-52	Rank sum	<0.001		<0.001	
DSI	P33-36 vs P37-52	Rank sum	0.2		<0.001	
OSI	All age groups	Kruskall-Wallis	<0.001	X=19.7	0.03	X=6.9
OSI	P29-32 vs P33-36	Rank sum	0.001		0.01	
OSI	P29-32 vs P37-52	Rank sum	<0.001		0.05	
OSI	P33-36 vs P37-52	Rank sum	0.8		0.6	

Table 9: Statistics for Figs. S1C-F

Area	Metric	Test	p value	Other
V1 & PSS	Peak rate	ANOVA: area, stimulus type	Area: <0.001 Stim type: 0.12 IA: 0.3	Area: F(1)=282.4 Stim type: F(1)=2.4 IA: F(1)=1.2
V1 & PMLS	Response duration	ANOVA: area, stimulus type	Area: <0.001 Stim type: <0.001 IA: <0.001	Area: F(1)=9733.9 Stim type: F(1)=400.4 IA: F(1)=93.7
V1 & PMLS	Average rate	ANOVA: area, stimulus type	Area: <0.001 Stim type: <0.001 IA: <0.001	Area: F(1)=312.0 Stim type: F(1)=63.7 IA: F(1)=18.5
V1	Response duration	Rank sum	<0.001	
V1	Average rate	Rank sum	<0.001	
PMLS	Response duration	Rank sum	<0.001	
PMLS	Average rate	Rank sum	<0.001	

Table 10: Statistics for Fig. S4 (IA: interaction). All rank-sum tests compare drifting versus flashing gratings.