30th April 2025

30th April 2025 — corrected below

Insights from aquaporin structures into drug-resistant sleeping sickness

Modestas Matusevicius[1#], Robin A. Corey[6,14#], Marcos Gragera[7#], Keitaro Yamashita[2#],

Teresa Sprenger[4,13#], Marzuq A. Ungogo[3,16,17], James N. Blaza[10], Pablo Castro-Hartmann[11],

Dimitri Y. Chirgadze[4], Sundeep Chaitanya Vedithi[1], Pavel Afanasyev[12], Roberto Melero[7],

Rangana Warshamanage[2,15], Anastasiia Gusach[2], Jose Maria Carazo[7], Mark Carrington[4], Tom

L. Blundell[1], Garib Murshudov[2], Phillip Stansfeld[9], Mark Sansom[6], Harry P. de Koning[3],

Christopher G. Tate[2] & Simone N. Weyand[1,4,5,8]*

[1]Department of Medicine, University of Cambridge, Victor Phillip Dahdaleh Building, Heart

& Lung Research Institute, Papworth Road, Cambridge Biomedical Campus,

Cambridge, CB2 0BB, UK

[2]MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge CB2 0QH, UK

[3]School of Infection and Immunity, University of Glasgow, Glasgow, UK

[4]Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge, UK

[5]Cambridge Institute for Medical Research, Keith Peters Building, Biomedical Campus, Hills

Rd, Cambridge CB2 0XY, UK

[6]Department of Biochemistry, University of Oxford, Oxford OX1 3QU, UK

[7]Instruct Image Processing Centre (I2PC), Unidad de Biocomputación, Centro Nacional de

Biotecnología, c/ Darwin, 3 (Universidad Autónoma de Madrid) 28049 Cantoblanco,

Madrid, Spain

[8]EMBL's European Bioinformatics Institute (EMBL-EBI), Wellcome Genome Campus,

Hinxton, Cambridgeshire, CB10 1SD, UK

[9]School of Life Sciences, Gibbet Hill Campus, The University of Warwick, Coventry, CV4

7AL, UK

[10]Structural Biology Laboratory and York Biomedical Research Institute, Department of

Chemistry, The University of York, Heslington, York, YO10 5DD, UK

[11]Materials and Structure Analysis, Thermofisher Scientific, Eindhoven, 5651 GG,

Netherlands

[12]ETH Zuerich, ScopeM, Cryo-EM Knowledge Hub, HPM C 51, Otto-Stern-Weg 3, 8093

Zurich, Switzerland

30th April 2025

[13]Present address: Eurofins BioPharma Product Testing Germany, Behringstraße 6/8, 82152
Planegg/ Munich, Germany
[14]School of Physiology, Pharmacology and Neuroscience, Biomedical Sciences Building,
Bristol BS8 1TD
[15]Present address: Science & Technology Facilities Council, Research Complex at Harwell,
Didcot OX11 0FA, England
[16]Department of Veterinary Pharmacology and Toxicology, Ahmadu Bello
University, Zaria, Nigeria
[17]Present address: Division of Immunology, The Roslin Institute, University of Edinburgh,
Edinburgh, UK
43
44 [#] Equal contribution
45 * Corresponding author
46
47 Correspondence for the manuscript:
48 Dr. S. N. Weyand
49
50 Email   sw644@cam.ac.uk
51 Telephone   +44-(0)7717217072
52
53

30th April 2025

Abstract

Trypanosoma brucei is the causal agent of African trypanosomiasis in humans and animals, the latter resulting in significant negative economic impacts in afflicted areas of the world. Resistance has arisen to the trypanocidal drugs pentamidine and melarsoprol through mutations in the aquaglyceroporin TbAQP2 that prevent their uptake. Here we use cryogenic electron microscopy to determine the structure of TbAQP2 from *Trypanosoma brucei*, bound to either the substrate glycerol or to the sleeping sickness drugs, pentamidine or melarsoprol. The drugs bind within the AQP2 channel at a site completely overlapping that of glycerol. Mutations leading to a drug-resistant phenotype were found in the channel lining. Molecular dynamics simulations showed the channel can be traversed by pentamidine, with a low energy binding site at the centre of the channel, flanked by regions of high energy association at the extracellular and intracellular ends. Drug-resistant TbAQP2 mutants still bind pentamidine, but the much weaker binding in the centre of the channel is insufficient to compensate for the high energy processes of ingress and egress, hence impairing transport at pharmacologically relevant concentrations. These structures of an aquaporin bound to a drug and represent a novel paradigm for drug-transporter interactions and could provide new mechanisms for targeting drugs into other pathogens or human cells.

Introduction

Human African trypanosomiasis (HAT) remains a neglected tropical disease[1]. It is transmitted by tsetse flies and, once symptoms are apparent, treatment is necessary to avoid fatality. The infection starts with a haemolymphatic stage that progresses to a late stage infection of the central nervous system, at which point severe neurological symptoms occur with rapid progression to coma and death[2]. Early-stage infections are commonly treated with pentamidine and the late stage with melarsoprol, although for most cases this is now being replaced with fexinidazole[3]. Animal African trypanosomiasis (AAT) causes billions of dollars in damage to agriculture throughout the tropics[4,5]. For both HAT and AAT, chemotherapy remains the only control option and most of the medication is old, inadequate and threatened by resistance[3,4,6,7].

For most trypanocidal drugs, resistance is associated with mutations in the membrane transporters through which the drugs enter the trypanosome[8]. The uptake of pentamidine and melarsoprol was initially attributed to a High Affinity Pentamidine Transporter (HAPT)[9,10], which was subsequently identified as the aquaglyceroporin TbAQP2[11,12]. Deletions and

chimeric rearrangements between TbAQP2 and the adjacent gene TbAQP3 were found to be

responsible for melarsoprol-pentamidine cross-resistance (MPXR; Fig. S1) [6,12-15]. To

understand the differences between TbAQP2 (water/glycerol/drug transporter) and TbAQP3

(classical aquaglyceroporin transporting water/glycerol[16] but not pentamidine or

melarsoprol[11,12,17]), mutants were made exchanging amino acids between the proteins[17].

Mutation of TbAQP2 to add any of the TbAQP3 selectivity filter residues Tyr250, Trp102 or

Arg256 resulted in inhibition of melarsoprol and pentamidine uptake, whereas TbAQP3

containing all three of their TbAQP2 counterparts (Leu250, Ile102, Leu256) permitted the

uptake of pentamidine. However, the molecular details of the interactions between TbAQP2

and drugs as well as the resistance mechanism remained unclear, with alternative models

proposing that TbAQP2 acts as a channel[8,17,18] or receptor[19] for the drugs. To resolve this

quandary and to resolve the molecular interactions underlying drug uptake by aquaporins, we

determined the structures of TbAQP2 bound to either melarsoprol, pentamidine or the native

substrate, glycerol.

Cryo-EM structures of TbAQP2 and drug binding

TbAQP2 was expressed using the baculovirus expression system in insect cells and

purified in the presence of either a drug (melarsoprol or pentamidine) or the substrate

glycerol (see Methods, Fig. S2a,b). Structures of TbAQP2 bound to either glycerol,

melarsoprol, or pentamidine were determined by electron cryogenic microscopy (cryo-EM)

to overall resolutions of 3.2 Å, 3.2 Å and 3.7 Å, respectively (Figs. S2c, S3 and S4a-c). The

resolution was sufficient to see the majority of the side chains and the bound drug or glycerol.

All of the cryo-EM structures were virtually identical (rmsd 0.2 Å) and represented the

aquaporin tetramer (Fig. 1a,b) that is typical for this family of channels[20]. The tetramer

contained four channels, each consisting of a single TbAQP2 polypeptide folded into a

bundle of eight α-helices (Fig 1b), two of which (helices H3 and H7) penetrated the plasma

membrane only partially, as is typical for aquaporins[20], the others crossing the membrane

fully.

Melarsoprol and pentamidine are bound in a channel formed by the TbAQP2 protomer,

completely overlapping the binding site for the substrate glycerol (Fig. 1c,d). The densities of

the substrates were distinct from one another and are consistent with the structures of the

substrates (Figs. S5 and S6). The position of drug binding is consistent with a hotspot

analysis performed on the structures (see Methods and Fig. S2d,e). This analysis defines

likely regions where molecules may bind based on *in silico* docking scores of small molecule

fragments docking to a model of TbAQP2[21] (see Methods). Binding of the two glycerol

molecules is mediated by a total of six hydrogen bonds, four to backbone carbonyl groups

(residues Gly126, Gly127, His128, Leu129) and two to the side chain of His128 (Fig. 2a).

These are on a polar strip of residues running down the channel, that also includes Val133,

which makes van der Waals contacts with the glycerol molecule closest to the cytoplasm. The

remainder of the contacts are van der Waals interactions (Val222, Phe226, Ile241, Val245) on

the opposite side of the channel. It is striking that all eleven amino acid residues within 3.9 Å

of the two glycerol molecules also make contacts to both pentamidine and melarsoprol (Fig.

2a). Whereas the TbAQP2-glycerol contacts are equally polar and non-polar, van der Waals

interactions dominate contacts between the drugs and TbAQP2. The exceptions are a

hydrogen bond between the backbone carbonyl of Ala259 and both drugs, and a backbone

carbonyl hydrogen bond between Gly126 and melarsoprol that is also seen between glycerol

and TbAQP2. It is also remarkable that an overlay between the three different structures

shows a very close alignment between glycerol, pentamidine and melarsoprol (Fig. 2b) which

is consistent with the channel structure being virtually identical regardless of whether a drug

or glycerol is bound. The size of the channel with pentamidine bound is very similar to the

glycerol-bound *Plasmodium falciparum* AQP structure (Fig. S2f). The close fit of the drugs

within the channel is consistent with the extremely limited modification of the drugs that is

possible whilst retaining the ability to pass through the pore, including modifications that

reduce flexibility[17]. Diamidines with fixed-angle structures such as diminazene and

furamidine are not substrates for TbAQP2 but are exclusively accumulated through

aminopurine transporter TbAT1[22,23]. A number of amino acid residues interact only with

pentamidine and/or melarsoprol and not with glycerol (Leu84, Ile110, Val114, Leu122,

Leu218, Ala259, Asn261, Met260, Leu264), which is a result of the drugs being longer and

extending further along the channel towards the extracellular surface when compared with

glycerol (Figs. 1d and 2b). Towards the intracellular surface, only Leu122 makes contacts to

the drugs and not to glycerol. In all, 19 residues in TbAQP2 interact with pentamidine,

compared with 18 residues interacting with melarsoprol and 11 to the two molecules of

glycerol.

The structure of AQP2 from *Plasmodium falciparum* (PfAQP) [24] with glycerol bound is

the closest structural homologue to TbAQP2 available (rmsd 0.8 Å, PDB ID 3CO2, 33%

amino acid sequence identity). The overall arrangement of α-helices in TbAQP2 is identical

to that in PfAQP (Fig. 3a-d) and other aquaglyceroporins whose structures have been

determined (human AQP7[25], human AQP10[26] and *Escherichia coli* GlpF[27]). Despite the low

identity in amino acid sequence, there are striking similarities between the structures of

TbAQP2 and PfAQP in the conserved NPA/NPA motif and the positions of bound glycerol

molecules in the channel. The NPA/NPA motif is found at the ends of the half-helices H3 and

H7, with TbAQP2 and PfAQP similarly differing from the canonical sequences (NPS/NSA in

TbAQP2 and NPS/NLA in PfAQP; Fig 3a,b). Superposition of TbAQP2 and PfAQP shows

the position and rotamer of Asn-Pro-Ser in the NPS motif and both Asn and Ala in the

NSA/NLA motifs are identical (Fig. 3c). Some of the network of hydrogen bonds that

maintain the structure of this region are also conserved, such as the hydrogen bonds in

TbAQP2 between Asn130 and the backbone carbonyl of 260 and the backbone amine of

Ala132, the hydrogen bond between Ser263 and Gln155 and the hydrogen bond between

Asn261 and the backbone amide of Ser263. However, both Asn residues in the NPS/NLA

motif of PfAQP form hydrogen bonds with one of the glycerol molecules in the channel,

whereas the Asn residues in the NPS/NSA motif in TbAQP2 form weak van der Waals

interactions with the glycerol molecules, and Asn261 also makes a hydrogen bond to the

backbone carbonyl of Leu129. The position of the intracellular loop between H2 and H3 is

also conserved between TbAQP2 and PfAQP (Fig. 3c) with the backbone carbonyls of

residues 126, 127 and 128 aligning and all forming hydrogen bonds in both channels to

glycerol molecules. The conservation in structure is reflected in the conserved position of the

glycerol molecules in the intracellular half of the channel in TbAQP2 and PfAQP, although

the orientation of the hydroxyl groups of the glycerol molecules differs. A divergence in the

position of the extracellular loop between H6 and H7 in TbAQP2 and PfAQP (Fig. 3c) leads

to a wider channel in TbAQP2 and no ordered glycerol molecules are observed in the channel

at this point, unlike in other aquaglyceroporins (Fig. 3d). Another difference between the

transport of glycerol by TbAQP2 and PfAQP is that there is a water molecule coordinated

between each glycerol molecule in PfAQP, whereas in TbAQP2 the glycerol molecules

interact directly. Any role of water molecules in the transport of glycerol by TbAQP2

remains to be elucidated as the resolution of the current cryo-EM structure is insufficient for

resolving them.

Molecular dynamics simulations

To assess the dynamic properties of the structurally-resolved pentamidine binding pose,

all-atom molecular dynamics (MD) simulations were performed on TbAQP2 in a tetrameric

state (see Methods). In these simulations, both the protein and pentamidine are very stable

(protein RMSD = 0.17 ± 0.01 nm and pentamidine RMSD = 0.16 ± 0.03 nm, see Fig. S7a,e)

and pentamidine binding prevents water flux through the channel (Fig. 4a,b). The stability of

the pentamidine is enabled by a number of extremely high occupancy interactions with

specific residues lining the TbAQP2 pore (Fig. S7b). Hydrogen bond analysis reveals around

23 hydrogen bonds may be made between pentamidine and the channel (Fig. S8a,b). In

contrast, water shows a different pattern of high occupancy interactions in the pore, with MD

snapshots highlighting several key positions of where waters interact with the channel as they

flow through (Fig. S8c). The presence of pentamidine both severely reduces the number of

waters in the channel and slightly impacts their orientation (Fig. S8d,e). It was not possible to

perform MD simulations of TbAQP2 in the presence of melarsoprol, because the presence of

an arsenic atom causes force field parameterization issues.

We then explored the translocation of pentamidine through the TbAQP2 channel. The

positively charged state of pentamidine suggests that it might be translocated by the $\Delta\Psi$

component of the membrane potential. To test this, we applied a z-dimension electric field to

simulations containing monomeric TbAQP2 in a membrane, using either a positive field (the

opposite direction to the physiological membrane) or a physiologically-correct negative field.

In all simulations, the resulting potential causes the pentamidine to translocate through

TbAQP2, and in some simulations (1 out of 3 repeats for each condition) the pentamidine

fully leaves the channel (Fig. 4c). For the physiologically correct negative field, the

pentamidine moves downwards (Fig. 4c), suggesting that the membrane potential is able to

import pentamidine into the cell, consistent with experimental evidence that TbAQP2-

mediated pentamidine uptake is sensitive to ionophores that depolarise the plasma

membrane[10,17].

Drug resistant mutations in TbAQP2

T. brucei contains two related aquaglyceroporins, TbAQP2 and TbAQP3, with only

TbAQP2 being able to transport pentamidine and melarsoprol into the trypanosome[11,12]. Drug

resistance has arisen through recombination between the two genes and the expression of

TbAQP2-AQP3 chimeras[13-15]. There are 68 amino acid residues that differ between TbAQP2

and TbAQP3 and the drug-resistant chimeras can contain 20-43 differences to TbAQP2 (Fig.

S1). A number of these residues have been mutated individually in attempts to identify key

residues involved in drug translocation and thus potentially resistance. At the intramembrane

ends of the two transmembrane half-helices H3 and H7 are the aforementioned NSA/NPS

motifs, Asn130-Ser131-Ala132 and Asn261-Pro262-Ser263, respectively (Fig. 3a). In

aquaporins such as TbAQP3 these conserved residues are the canonical NPA/NPA motifs,

and mutations were made (S131P+S263A) to convert the NSA/NPS motif in TbAQP2 to

NPA/NPA. This resulted in a 95% reduction in pentamidine transport by the mutant, although

this was still measurably higher than that observed in a TbAQP2 knockout cell line[17] and a

similar reduction in the transport of the melarsoprol analogue cymelarsan (Fig. S9). The cryo-

EM structures show that both Ser131 and Ser263 are critical in stabilising this region of the

transporter through specific hydrogen bonds with Gln155 and the backbone amine group of

Ala132 (Fig. 3a) and do not make any direct interactions either to the drugs or the glycerol

molecules. Thus, the decrease in pentamidine uptake observed in the S131P+S263A TbAQP2

mutant is probably more complex than previously thought, as it seems unlikely to have arisen

through simple changes in transporter-drug interactions or a narrowing of the channel.

Based on a previous modelling study, several other mutations were predicted to

potentially affect drug uptake in the TbAQP2-TbAQP3 chimeras[17]. Based on the cryo-EM

structures, the mutated residues fall into three distinct groups: those not in the channel

(I190T, W192G), those lining the channel and within van der Waals distance of pentamidine

(I110W, L258Y, L264R) and those lining the channel, >3.9 Å from pentamidine and

observed to make contacts to pentamidine during MD simulations (L84W/M, L118W/M,

L218W/M). In all instances, there was a >90% decrease in pentamidine uptake) and in many

cases no residual pentamidine transport was detected (e.g. I110W, L264R, L84W) [17]. To

understand the molecular basis for the dramatic decreases observed in pentamidine transport,

we performed molecular dynamics simulations on a subset of the mutants.

Three mutations were chosen for molecular dynamics simulations (I110W, L258Y,

L264R), all of which were defective in uptake of cymelarsan (Fig. S9) as well as

pentamidine[17]. These three residues all make high occupancy contacts to pentamidine in the

MD simulations (Fig. S7b) and residues Ile110 and Leu264 make van der Waals contacts

with pentamidine in the cryo-EM structure (Leu258 is 5.5 Å away from pentamidine). The

double mutant L258Y+L264R was used in the MD simulations as this is found in a number

of drug-resistant *T. brucei* strains such as P1000 and Lister 427MR[12]. The mutant I110W is

found in the *T. brucei* drug resistant strain R25[28], but in the absence of the L258Y+L264R

double mutation (Fig 1). Therefore, we performed atomistic MD simulations of monomeric

TbAQP2 with either the I110W (hereafter IW) or L258Y+L264R (hereafter LY/LR)

mutations present, as well as with wild-type monomeric TbAQP2. Each protein was

simulated both with pentamidine bound and removed (apo). The mutations do not affect

TbAQP2 stability (Fig. S7e), but do destabilise pentamidine binding, causing it to shift away

from the initial binding pose along the z-axis (Fig. S7c), suggesting a lowered affinity for the

structural pose. The pore radii are unchanged by the mutations (Fig. S7d), meaning that this

effect is likely to be a loss of specific interactions rather than a more general effect on protein

structure.

We next computed energy landscapes for pentamidine moving along the z-axis through

the channel using potential of mean force (PMF) calculations with umbrella sampling. The

data for WT TbAQP2 reveals a clear energy well for the structurally resolved pentamidine at

-0.2 nm (Fig. 4d and Fig. S10c). The depth of this well (-40 kJ/mol) suggests a very high

affinity binding site, consistent with previous kinetic analyses[9,10,29], and is much higher than

previously investigated docked poses[17] of pentamidine in models of TbAQP2. Notably, this

energy well is much shallower for both the IW (-20 kJ/mol) and LY/LR mutants (-25 kJ/mol),

representing a relative reduction in binding likelihood of about 2,800-fold and 400-fold,

respectively. In biochemical terms, these changes would result in a K_m shift from ~50 nM to

50 or 300 μM, respectively, all but abolishing uptake at pharmacologically relevant

concentrations; the experimental K_m for pentamidine on wild type TbAQP2 is 36 nM[10]. The

reduction in pentamidine binding likelihood is consistent with the increased z-axis dynamics

seen in Fig. S7c. Flanking the central energy well are large energy barriers at about -0.7 nm

and 1 nm (Fig. 4d), similar to those seen for urea translocation in HsAQP1[30]. These would

presumably slow pentamidine entry into and through the channel, and previous studies did

reveal a slow transport rate for pentamidine[10,29]. The energy barriers are highest in the IW

mutant, and interestingly the inner (-0.7 nm) barrier is lower in the LY/LR mutant (Fig.

S10c). In all cases, especially WT, small energy minima are observed at about -1.35 nm and

+2.7 nm; these represent the drug binding at the entrance and exit of the channel (Fig. S10d).

The differences in landscape between the WT and mutants helps explain how these mutations

confer a resistance to pentamidine, mostly through reduced binding affinity.

Discussion

285   It has only recently become accepted that the aquaglyceroporin AQP2 in *T. brucei* is

286   acting as a *bona fide* transporter of pentamidine and melarsoprol that is essential for

287   translocation of the drugs into the cell[17]. The cryo-EM structures of TbAQP2 identify the

288   binding site for melarsoprol and pentamidine in a wide channel that has, at least for part of

the trajectory, similar dimensions to channels in other aquaglyceroporins (Fig. 3d). The overall similarity between TbAQP2 and the structures of other aquaglyceroporins suggests a similarity in the mechanism of glycerol transport, despite differences in the amino acid residues lining the channel. It also suggests that there is scope for pharmacological targeting of other aquaglyceroporins, either for uptake of cytotoxic drugs or inhibition of glycerol uptake. Leishmaniasis, for instance, is another neglected tropical disease, caused by parasitic protozoa of the genus *Leishmania*; it is commonly treated by pentavalent antimonials meglumine antimoniate and sodium stibogluconate. The antimony that is released in the acidic parasitophorous vacuole enters the parasite through its AQP1 aquaporin[31]. Widespread antimonial drug resistance in *Leishmania donovani* in India has been linked to a frameshift mutation in LdAQP1[32]. In another example, *Plasmodium spp.* contain a single aquaglyceroporin and studies in mouse models have shown that a knockout of the aquaglyceroporin leads to a significant loss of virulence and reduction in erythrocyte infectivity by merozoites[33], suggesting that a specific AQP inhibitor could be part of antimalarial treatments. Interestingly, pentamidine is a known antimalarial[34] and although a substrate of the *P. falciparum* choline transporter, does not act through inhibition of that carrier[35]; inhibition/permeation of PfAQP by pentamidine has not been investigated to date. The structure determination of TbAQP2 highlights new opportunities for drug development targeting the aquaporin family to treat a number of parasitic diseases whose current treatments often suffer from severe side effects and growing problems in drug resistance. A thorough understanding of the interaction of extended, flexible drugs like pentamidine and melarsoprol with AQP channel linings could contribute to the design of new substrates able to penetrate not just protozoan but also human AQPs.

After this work was completed, a paper was published describing the structures of TbAQP2 bound to pentamidine and melarsoprol[36]. The structures of TbAQP2 are very similar to the ones described here (RMSD 0.6 Å), despite the environments of the purified channels being significantly different. In the work described here, TbAQP2 was purified in detergent and then incorporated into saposin A (SapA) [37] lipid nanoparticles (see Methods), whereas Chen *et al.* [36] incorporated TbAQP2 into lipid nanodiscs formed from a helical membrane scaffold protein (MSP) [38,39]. Lipid nanodiscs and nanoparticles are considered better mimetics of a lipid bilayer than detergent micelles, although biophysical studies indicate that there are significant differences between these lipidic structures and a biological membrane[40,41]. Comparison of the pentamidine in the different structures indicates good agreement in its position within the channel, although the pose of the drug does differ slightly

(Fig. 5a). In contrast, there is a significant difference in the position and pose of melarsoprol

between the different structures (Fig. 5b). Melarsoprol in the TbAQP2 structure from Chen *et*

al. [36] is shifted further towards the extracellular surface of the channel and is rotated

approximately 180° about an axis in the membrane plane and centred on the arsenic atom

(Fig 5b). This unexpected difference in orientation of melarsoprol in the two structures may

perhaps not be so surprising as could be initially thought. The two structures represent the

thermodynamically most stable states under the buffer conditions and detergent/lipid

nanoparticles used for protein purification, which were different. In addition, the site of

localisation of the drugs in the TbAQP2 channel is not a physiologically relevant 'binding

site' as used to describe, for example, the orthosteric binding site in G protein coupled

receptors where a hormone or neurotransmitter binds. Thus, evolution has not evolved a high-

affinity binding site in the TbAQP2 channel and neither were the drugs designed to bind to

TbAQP2. The channel merely facilitates the entry of the drugs into the cell through passive

diffusion and it is not necessary for its mode of action to bind to TbAQP2 in a specific pose.

Indeed, *a priori*, there is no reason to think that melarsoprol has to traverse the channel in any

particular orientation, although under cellular conditions where a membrane potential is

present it might be anticipated that the melamine moiety may enter the channel first given its

net positive charge. Regrettably, in the absence of molecular dynamics simulations (due to

the issues with parameterization of the arsenic atom) these ideas have to remain untested.

Acknowledgements

S.N.W.'s lab was funded by a Sir Henry Dale fellowship from the Wellcome Trust and the

Royal Society London (Grant Number 101234/Z/13/Z) and by the Isaac Newton Trust

Cambridge (Grant Number 15.40(a)). The work in C.G.T.'s laboratory was supported by core

funding from the Medical Research Council (MRC U105197215). T.S. was supported by an

MRC-DTP PhD fellowship, a Cambridge Trust Vice-Chancellor's Award, the Sackler fund

and a King's College Cambridge PhD overrun fund. R.A.C, M.S.P.S, and P.J.S are supported

by Wellcome (208361/Z/17/Z). Aquaporin-related work in H.P.d.K. lab was supported by the

Medical Research Council (Grant Number G0701258) and a BBSRC Impact Accelerator

award (BB/S506734/1). R.W. was supported by the Wellcome Trust (208398/Z/17/Z). MC is

a Wellcome Investigator (217138/Z/19/Z). This project made use of time on ARCHER2

granted via the UK High-End Computing Consortium for Biomolecular Simulation,

HECBioSim (http://hecbiosim.ac.uk), supported by EPSRC (EP/R029407/1). In addition,

simulations were performed using the computational facilities of the Advanced Computing

Research Centre, University of Bristol (http://www.bristol.ac.uk/acrc/). This work benefited

from access to the Instruct Image Processing Centre (I2PC), an Instruct-ERIC centre;

financial support was provided by PID12540.

Author contributions

M.C. and S.N.W conceived the project. T.S. and S.N.W. cloned, overexpressed, purified, and

analysed protein samples. J.N.B., P.C.-H., D.Y.C. and S.N.W. prepared grids, screened and

collected data. M.G. and R.M. processed the raw data. M.M. and P.A. built the models and

performed refinements. A.G performed further structural analyses. K.Y. and R.W. identified

and fitted the substrates and performed refinements. J.M.C. advised on the processing and

G.M. advised on the refinement process. V.S.C. performed hot spot analysis and T.L.B.

advised. H.P.d.K. and M.C. advised the mutational studies. R.A.C. performed the molecular

dynamics simulations, and R.A.C., P.S. and M.S. did the analysis of the MD simulation

trajectories. M.A.U. performed cymelarsan uptake experiments. C.G.T. carried out structure

analysis and S.N.W. managed the overall project. The manuscript was written by S.N.W.,

H.P.d.K., R.A.C. and C.G.T., and included contributions from all the authors.

Data availability

Structures have been deposited in the Protein Data Bank (PDB; https://www.rcsb.org/), and

the associated cryo-EM data have been deposited in the Electron Microscopy Data Bank

(EMDB; https://www.ebi.ac.uk/pdbe/emdb/) and the Electron Microscopy Public Image

Archive (EMPIAR; https://www.ebi.ac.uk/empiar/): SapA-TbAQP2-NP bound to glycerol

(8OFZ, EMD-16864 and EMPIAR-11410); SapA-TbAQP2-NP bound to pentamidine

(8OFY, EMD-16863 and EMPIAR-11412); SapA-TbAQP2-NP bound to melarsoprol

(8OFX, EMD-16862 and EMPIAR-11411). There are no restrictions on data availability.

METHODS

Data reporting

Sample size was not predetermined by statistical methods, no randomization was
applied to experiments and investigators were not blinded to experimental allocation or
outcome assessment.

Cloning and expression of wild-type AQP2

A construct encoding wild-type *T. brucei* AQP2 (residues 1–312) that included a
tobacco etch virus (TEV) cleavage site, eGFP, and a decahistidine tag (TbAQP2-TEV-eGFP-
His10) was generated and cloned into a pFastBac vector (Thermo Fisher) using standard
ligation procedures. Recombinant pFastBac-TbAQP2-TEV-eGFP-His10 vector was
transformed into Stellar Competent Cells (Takara) following the Bac-to-Bac Baculovirus
Expression System manual (Thermo Fisher). Expression of AQP2 in *Spodoptera frugiperda* 9
(Sf9) cells (Thermo Fisher Scientific) was performed following the Bac-to-Bac Baculovirus
Expression System manual (Thermo Fisher). The cells were then collected by centrifugation
(10 min, 4°C, 3000 $\times g$), flash-frozen in liquid nitrogen, and stored at -80 °C until further use.

Purification of AQP2 bound to either glycerol, pentamidine or melarsoprol and
reconstitution into saposin-A nanoparticles (SapA-NP)

All steps described below were carried out at 4°C. *Sf*9 cells were defrosted on ice,
resuspended in cold homogenization buffer (100 mM NaCl, 20 mM HEPES pH 7.5, 5% (v/v)
glycerol, 2 mM PMSF, 1× Complete Protease Inhibitor Cocktail (Sigma)) and homogenised
by applying 20 strokes in a Potter-Elvehjem glass homogenizer on ice. Membranes were
solubilised using 1% (w/v) final concentration n-dodecyl-β-D-maltopyranoside (DDM,
Anatrace) and a final concentration of 0.03% (w/v) cholesteryl hemisuccinate (CHS,
Anatrace) for 4 hours at 4°C on a roller platform. Insoluble material was removed by
ultracentrifugation (100,000 $\times g$, 60 min, 4 °C). The solubilisate was supplemented with 20
mM imidazole (final concentration) and incubated (16 h, 4°C) with 2 ml of Ni^{2+}-NTA
agarose resin (Qiagen) per litre of *Sf*9 cell culture. The resin was then applied to a gravity-
flow column and washed three times with 20 column volumes washing buffer (100 mM
NaCl, 20 mM HEPES pH 7.5, 2% (v/v) glycerol, 2 mM PMSF, 1× Complete Protease

Inhibitor Cocktail (Sigma), 0.05% (w/v) DDM, 0.03% (w/v) CHS, 100 mM imidazole) and

was eluted twice with one column volume elution buffer (100 mM NaCl, 20 mM HEPES pH

7.5, 2% (v/v) glycerol, 1× Complete Protease Inhibitor Cocktail (Sigma), 2 mM PMSF, 500

mM imidazole) each. Both eluates were combined and concentrated in an Amicon Ultra 4 mL

centrifugal filter (100,000 MWCO, Merck) and aggregates were removed using a syringe-

driven filter units (Millex-GV, 0.22 μm, PVDF, 13 mm, Merck). The filtrate was loaded at

0.4 ml/min onto a Superdex Increase S200 10/300 (GE Healthcare) size exclusion

chromatography (SEC) column, pre-equilibrated with 0.05% (w/v) DDM, 100 mM NaCl, 20

mM HEPES pH 7.5. The peak fractions containing AQP2 were pooled and concentrated as

above. Protein concentration was estimated by the Pierce BCS assay according to the

manufacturer's instructions.

SapA-NPs were prepared and stored in SapA-NP buffer consisting of 150 mM NaCl,

50 mM sodium acetate pH 4.8) [42]. Purified TbAQP2 tetramer was combined at a

TbAQP2:SapA:lipid molar ratio of 0.1:1:10 and dialysed (20 h, 4°C) against SapA-

reconstitution buffer (100 mM NaCl, 20 mM HEPES pH 7.5). SapA-NPs, unincorporated

TbAQP2 and SapA were removed by SEC and SapA-TbAQP2-NP tetramer complex was

concentrated using Amicon Ultra centrifugal filters (100,000 MWCO; Merck). Melarsoprol

or pentamidine was added to the purified SapA-TbAQP2-NP tetramer complex (final

concentration 1 mM, 2 h, 4°C) applied to a Superdex Increase S200 10/300 SEC column (GE

Healthcare) and fractions containing the complex were concentrated using Amicon Ultra

centrifugal filters (100,000 MWCO; Merck) to final concentrations between 0.8 mg/ml and

1.6 mg/ml. Final samples were centrifuged (1 h, 18,000 ×*g*, 4°C) immediately prior to cryo-

EM grid preparation.

Vitrified sample preparation and data collection

Cryo-EM grids were prepared by applying 2 μl of purified SapA-TbAQP2-NP with

either glycerol, pentamidine or melarsorol bound (0.8-1.6 mg/ml) onto either glow-

discharged PEGylated UltrAuFoil R1.2/1.3 300 Mesh Gold grids (Agar scientific, No.

AGS187-3) or glow-discharged Quantifoil R1.2/1.3 300 Mesh Gold grids (Agar scientific,

No. AGS143-8). Grids were blotted with filter paper for 4 s before plunge-freezing in liquid

ethane (at −181 °C) using an Thermo Fisher Scientific Vitrobot Mark IV (95% relative

humidity, 4°C). All cryo-EM datasets were collected on Titan Krios microscopes (Thermo

Fisher Scientific) operating at 300 kV. SapA-TbAQP2-NP with glycerol was collected at the

Nanoscience Centre (University of Cambridge) with a Falcon 3 detector to obtain dose-

fractioned movies. 2881 movies were recorded with 75,000× magnification at 1.07 Å per

pixel using an exposure time of 60 s resulting in a total exposure of (30 e/pixel) and a target

defocus range of -1.4 and -3.2 μm. Datasets of SapA-TbAQP2-NP with bound pentamidine

or melarsoprol were collected at the cryo-EM facility at the Department of Biochemistry,

University of Cambridge; images were recorded using a K2 detector in counting mode,

operating with GIF Quantum LS imaging filter (Gatan) with a slit width of 20 eV to obtain

dose-fractioned movies with a 100 μm objective aperture. 2400 movies were recorded for

SapA-TbAQP2-NP with pentamidine bound and 3000 movies for melarsoprol bound

complex. Micrographs for both datasets were recorded at a magnification of 130,000× (1.07

Å per pixel) as dose-fractioned movie frames with an exposure time of 15 s and 12 s,

respectively, resulting in a total exposure of 54 e/Å^2 and 57.5 e/Å^2 with target defocus ranges

of -1.5 and -3.3 μm, and -1.2 and -3 μm, respectively.

Cryo-EM data processing and 3D reconstruction

Cryo-EM data processing was conducted inside the Scipion platform[43]. Image stacks

(2,881 glycerol-bound, 4,086 melarsoprol-bound and 3,262 pentamidine-bound) were

subjected to beam-induced motion correction using Relion's implementation of

MotionCor2[44] by dividing each frame into 5 × 5 patches. CTF parameters were estimated

from dose-weighted micrographs as follows: estimations made with GCTF[45] with equiphase

averaging and with CTFFIND-4.1[46,47] were compared using «xmipp-ctf consensus». Outputs

from these consensus (micrographs for which the two algorithms agreed in their estimations)

undergone an additional round of ctf estimation with «xmipp-ctf estimation»[48], followed by a

second round of xmipp-ctf consensus for micrograph curation based on defocus, astigmatism,

resolution and ice thickness. As a result, the number of micrographs for downstream analysis

was 2102, 3586 and 2983 for SapA-TbAQP2-NP bound to either glycerol, melarsoprol or

pentamidine, respectively.

Autopicking was performed using crYOLO[49], that yielded 849,607 particles, 1,927,102

particles and 1,263,534 particles for SapA-TbAQP2-NP bound to either glycerol, melarsoprol

or pentamidine, respectively. Particles were extracted in a box-size of 250 px at the original

sampling rate of the micrographs (1.07 Å/px) and subjected to several rounds of 2D

classifications with cryoSPARC[50]. To further curate the particle set (ligand-free: 100,075;

melarsoprol: 202,752; pentamidine: 114,206). Particles were re-extracted with a box of 320

px before 3D classifications.

For the glycerol-bound dataset, an *ab initio* 3D model was generated using stochastic

gradient descent algorithm implemented in cryoSPARC, followed by non-uniform

refinement[51] with C4 symmetry and a static mask encompassing the transmembrane domains

and nanodiscs belt, while excluding the mobile GFP tags. With these settings, a 3.2 Å-

resolution map was obtained. For the melarsoprol and pentamidine datasets, six *ab initio*

models were generated. Particles from the best classes (melarsoprol: 126,551; pentamidine:

83845) were used for non-uniform-refinement with the same strategy as for the glycerol-

bound particles, yielding a 3.2 and 3.7 Å-resolution map, respectively.

The global resolution was calculated following the FSC gold standard threshold of

0.143. The refined maps were sharpened using deepEMhancer[52] to facilitate model building.

Finally, voxel size for melarsoprol and pentamidine was re-adjusted to 1.042 Å/px using

Relion post-process[53]. Local resolution was estimated with MonoRes[54].

490 Structure determination and model refinement

The monomer structure of TbAQP2 was predicted using the AlphaFold2 program as a

starting point for further refinement[55]. The final cryo-EM map was converted to an mtz file

with a range of sharpening and blurring options using CCPEM[56,57]. EMDA was used for

fitting maps[58]. Real-space refinement of the atomic coordinates into the cryo-EM map was

performed using Phenix[59] and Coot software[60]. Finally, models in the asymmetric unit were

refined against unsharpened and unweighted half maps using Servalcat Refmac5 pipeline[61]

with *C*4 symmetry constraints. The reference structure restraints were prepared with

ProSmart[62] using AlphaFold2 predicted models from the AlphaFold DB[63]. The final models

were evaluated for geometry, close contacts, and bond parameters using MolProbity[64].

Graphic representations of the fitted coordinates into electron density and the final cryo-EM

map were generated using PyMOL[65] and UCSF ChimeraX[66,67] packages.

502 Molecular dynamics simulations

The model for MD simulations was based on the structure of TbAQP2 with

pentamidine bound and was built into simulation boxes using CHARMM-GUI[68,69] (see Table

S2 for a list of systems built). Protein atoms were described with the CHARMM36m force

field[70,71] TbAQP2 was built either as a tetramer or a monomer. Side chain pK_as were assessed

using propKa3.1 and side chain side charge states were set to their default[72]. The pentamidine

molecule used existing parameters available in the CHARMM36 database under the name

PNTM with a charge state of +2. For the ligand-free simulations, the pentamidine molecule

was removed prior to simulation.

The proteins were built into membranes comprising 20% cholesterol and 80% POPC,

solvated with explicit water molecules using the TIP3P model, and neutralised with K^+ and

Cl^- to 150 mM. Box sizes were initially set to $10 \times 10 \times 10$ nm for the tetrameric systems and

$6.5 \times 6.5 \times 10$ nm for the monomeric systems. The use of an all-atom fixed charge force field

with explicit solvent and membrane is optimal to understanding the protein-drug interactions

central to this study. Each system was minimized and equilibrated according to the standard

CHARMM-GUI protocol[68,69]. Production simulations using random seeds were run in the

NPT ensemble with 2 fs time steps, temperatures held at 303.5 K using a velocity-rescale

thermostat and a coupling constant of 1 ps, and pressure maintained at 1 bar using a semi-

isotropic Parrinello-Rahman pressure coupling with a coupling constant of 5 ps[73,74]. Short

range van der Waals and electrostatics were cut off at 1.2 nm, and the particle mesh Ewald

(PME) method was used for long-range Lennard-Jones interactions[75]. Hydrogen bonds were

constrained using the LINCS algorithm.

Membrane potential simulations were run using the computational electrophysiology

protocol. An electric field of 0.1 V/nm was applied in the z-axis dimension only, to create a

membrane potential of about 1 V (see Fig. S10a). Note that this is higher than the

physiological value, and was chosen to improve the sampling efficiency of the simulations.

The protein and lipid molecules were visually confirmed to be unaffected by this voltage.

Potential of mean force (PMF) calculations were built from initial steered MD pulls

using an umbrella potential along a coordinate of the COM of the pentamidine molecule and

the COM of select residues in TbAQP2 which form the pentamidine binding site (Leu84,

Ile110, Val114, Leu122, Gly127, Leu129, Leu218, Val222, Phe226, Leu244, Val245,

Ala259, Met260, and Leu264). Pulls were run in the z-axis in either a positive or negative

direction, at a rate of 1 nm per ns and with a force constant of 1000 kJ/mol/nm^2. Snapshots

were taken along the coordinates every 0.05 nm, using a script from Dr. Owen Vickery

(https://doi.org/10.5281/zenodo.3592318), with a total of 167 snapshots to construct the

reaction coordinate. End points for the coordinate were determined based on the pentamidine

no longer making any contacts (<0.4 nm) with any atoms from the protein or membrane.

Simulations were run for 40 ns per window with a 5,000 kJ/mol/nm^2 umbrella restraint

imposed to keep the system in place with respect to the reaction coordinate. An atom in

Leu218 was chose as a reference for the treatment of periodic boundary conditions. Forces

were written every 1 ps. PMF profiles were constructed using the weighted histogram

analysis method (WHAM) [76] implemented in *gmx wham*[77] between -3.5 nm to 4.5 nm of the

coordinate and with cyclisation applied. The first 50 ns of each window was discarded as

equilibration time and 200 Bayesian bootstraps were used to estimate the PMF error.

Adequate coverage of the landscape was assessed based on histogram overlap (Fig. S10b),

and convergence was determined based on re-running the WHAM calculations on longer

windows until the landscapes converged (Fig. S10c).

All simulations were run in Gromacs 2020.3

(https://doi.org/10.5281/zenodo.7323409)[78]. Data were analysed using Gromacs tools and

VMD[79]. Pentamidine interactions with TbAQP2 were assessed using PyLipID[80]. Pore radii

were assessed using the CHAP package[81]. Plots were made using Matplotlib[82] or Prism 9

(GraphPad). MD input and output coordinate files are available at https://osf.io/zc235/.

Docking and hotspots experiments

Fragment Hotspot Maps (FHMs) is a powerful tool for predicting the binding hotspots

of small molecules on protein surfaces[21]. FHMs are based on the concept that certain

chemical fragments are more likely to interact with specific regions of a protein's surface, or

'hotspots', than others. By systematically mapping the interactions between a library of

fragment molecules and a target protein, FHMs can identify key hotspots and help guide the

design of new small molecules with improved binding affinity and specificity. The

development of FHMs has been driven by advances in computational methods for predicting

protein-ligand interactions, as well as by improvements in experimental techniques for

mapping the 3D structure of protein-ligand complexes. FHMs have been used to guide drug

discovery efforts in a variety of therapeutic areas, including oncology, infectious diseases,

and neurology.

One key advantage of FHMs is that they can help to identify "druggable" regions of a

protein's surface that may be otherwise difficult to target with small molecules. For example,

many protein-protein interaction interfaces are large and flat, making it challenging to design

small molecules that can disrupt these interactions. FHMs can help to identify the specific

regions of the interface that are most important for binding and guide the design of molecules

that can effectively target these regions. Another advantage of FHMs is that they can be used

to predict the binding modes of small molecules that have not yet been synthesized or tested

experimentally. This can be particularly useful in early-stage drug discovery, where

computational methods are often used to screen large databases of compounds for potential hits. Overall, FHMs represent a powerful approach for predicting and optimizing small molecule binding to protein surfaces.

We used FHMs program to generate hotspots for AQP2 monomer and probe protein's potential ligand binding sites. The superposition of the final structures and the predicted hotspot maps show a clear overlap between the hotspots and the two drugs (Fig. S2).

Lysis assay with arsenical drugs

The effects of cymelarsan and phenylarsine oxide (PAO) on *T. brucei* Lister 427 bloodstream forms was measured as the decrease in cell absorbance at λ 750 nm over time, based on the reduction of cell motility and increased cell lysis leading to reduced light scatter and absorbance in the cuvette[83], exactly as described[84]. Briefly, 2×10^6 cells in 200 μL buffer was placed in the wells of a 96-well plate and absorbance recorded in a PHERAstar plate reader (BMG Labtech, Durham, NC, USA) for 10 min before the addition of 20 μL of 100 μM of either cymelarsan or PAO in aqueous buffer, or buffer alone.

References

Organisation, W. H. Neglected tropical diseases. https://www.who.int/news-room/questions-and-answers/item/neglected-tropical-diseases, (2023).

Kennedy, P. G. Clinical features, diagnosis, and treatment of human African trypanosomiasis (sleeping sickness). Lancet Neurol **12**, 186-194, (2013).

De Koning, H. P. The Drugs of Sleeping Sickness: Their Mechanisms of Action and Resistance, and a Brief History. Trop Med Infect Dis **5**, (2020).

Giordani, F., Morrison, L. J., Rowan, T. G., De Koning, H. P. & Barrett, M. P. The animal trypanosomiases and their chemotherapy: a review. Parasitology **143**, 1862-1889, (2016).

Desquesnes, M. *et al.* Trypanosoma evansi and surra: a review and perspectives on transmission, epidemiology and control, impact, and zoonotic aspects. Biomed Res Int **2013**, 321237, (2013).

Baker, N., de Koning, H. P., Maser, P. & Horn, D. Drug resistance in African trypanosomiasis: the melarsoprol and pentamidine story. Trends Parasitol **29**, 110-118, (2013).

Carruthers, L. V. *et al.* Diminazene resistance in Trypanosoma congolense is not caused by reduced transport capacity but associated with reduced mitochondrial membrane potential. Mol Microbiol **116**, 564-588, (2021).

Munday, J. C. *et al.* Functional analysis of drug resistance-associated mutations in the Trypanosoma brucei adenosine transporter 1 (TbAT1) and the proposal of a structural model for the protein. Mol Microbiol **96**, 887-900, (2015).

Bridges, D. J. *et al.* Loss of the high-affinity pentamidine transporter is responsible for high levels of cross-resistance between arsenical and diamidine drugs in African trypanosomes. Mol Pharmacol **71**, 1098-1108, (2007).

10 De Koning, H. P. Uptake of pentamidine in Trypanosoma brucei brucei is mediated
by three distinct transporters: implications for cross-resistance with arsenicals. Mol
Pharmacol **59**, 586-592, (2001).
11 Baker, N. *et al.* Aquaglyceroporin 2 controls susceptibility to melarsoprol and
pentamidine in African trypanosomes. Proc Natl Acad Sci U S A **109**, 10996-11001,
(2012).
12 Munday, J. C. *et al.* Trypanosoma brucei aquaglyceroporin 2 is a high-affinity
transporter for pentamidine and melaminophenyl arsenic drugs and the main genetic
determinant of resistance to these drugs. J Antimicrob Chemother **69**, 651-663,
(2014).
13 Graf, F. E. *et al.* Chimerization at the AQP2-AQP3 locus is the genetic basis of
melarsoprol-pentamidine cross-resistance in clinical Trypanosoma brucei gambiense
isolates. Int J Parasitol Drugs Drug Resist **5**, 65-68, (2015).
14 Graf, F. E. *et al.* Aquaporin 2 mutations in Trypanosoma brucei gambiense field
isolates correlate with decreased susceptibility to pentamidine and melarsoprol. PLoS
Negl Trop Dis **7**, e2475, (2013).
15 Pyana Pati, P. *et al.* Melarsoprol sensitivity profile of Trypanosoma brucei gambiense
isolates from cured and relapsed sleeping sickness patients from the Democratic
Republic of the Congo. PLoS Negl Trop Dis **8**, e3212, (2014).
16 Uzcategui, N. L. *et al.* Cloning, heterologous expression, and characterization of three
aquaglyceroporins from Trypanosoma brucei. J Biol Chem **279**, 42669-42676, (2004).
17 Alghamdi, A. H. *et al.* Positively selected modifications in the pore of TbAQP2 allow
pentamidine to enter Trypanosoma brucei. Elife **9**, (2020).
18 Munday, J. C., Settimo, L. & de Koning, H. P. Transport proteins determine drug
sensitivity and resistance in a protozoan parasite, Trypanosoma brucei. Front
Pharmacol **6**, 32, (2015).
19 Song, J. *et al.* Pentamidine Is Not a Permeant but a Nanomolar Inhibitor of the
Trypanosoma brucei Aquaglyceroporin-2. PLoS Pathog **12**, e1005436, (2016).
20 Gonen, T. & Walz, T. The structure of aquaporins. Q Rev Biophys **39**, 361-396,
(2006).
21 Radoux, C. J., Olsson, T. S. G., Pitt, W. R., Groom, C. R. & Blundell, T. L.
Identifying Interactions that Determine Fragment Binding at Protein Hotspots. J Med
Chem **59**, 4314-4325, (2016).
22 de Koning, H. P. *et al.* The trypanocide diminazene aceturate is accumulated
predominantly through the TbAT1 purine transporter: additional insights on
diamidine resistance in african trypanosomes. Antimicrob Agents Chemother **48**,
1515-1519, (2004).
23 Ward, C. P., Wong, P. E., Burchmore, R. J., de Koning, H. P. & Barrett, M. P.
Trypanocidal furamidine analogues: influence of pyridine nitrogens on trypanocidal
activity, transport kinetics, and resistance patterns. Antimicrob Agents Chemother **55**,
2352-2361, (2011).
24 Newby, Z. E. *et al.* Crystal structure of the aquaglyceroporin PfAQP from the
malarial parasite Plasmodium falciparum. Nat Struct Mol Biol **15**, 619-625, (2008).
25 de Mare, S. W., Venskutonyte, R., Eltschkner, S., de Groot, B. L. & Lindkvist-
Petersson, K. Structural Basis for Glycerol Efflux and Selectivity of Human
Aquaporin 7. Structure **28**, 215-222 e213, (2020).
26 Gotfryd, K. *et al.* Human adipose glycerol flux is regulated by a pH gate in AQP10.
Nat Commun **9**, 4749, (2018).
27 Fu, D. *et al.* Structure of a glycerol-conducting channel and the basis for its
selectivity. Science **290**, 481-486, (2000).

28 Unciti-Broceta, J. D. *et al.* Specific Cell Targeting Therapy Bypasses Drug Resistance
Mechanisms in African Trypanosomiasis. PLoS Pathog **11**, e1004942, (2015).

29 Teka, I. A. *et al.* The diamidine diminazene aceturate is a substrate for the high-
affinity pentamidine transporter: implications for the development of high resistance
levels in trypanosomes. Mol Pharmacol **80**, 110-116, (2011).

30 Hub, J. S. & de Groot, B. L. Mechanism of selectivity in aquaporins and
aquaglyceroporins. Proc Natl Acad Sci U S A **105**, 1198-1203, (2008).

31 Plourde, M. *et al.* Generation of an aquaglyceroporin AQP1 null mutant in
Leishmania major. Mol Biochem Parasitol **201**, 108-111, (2015).

32 Imamura, H. *et al.* Evolutionary genomics of epidemic visceral leishmaniasis in the
Indian subcontinent. Elife **5**, (2016).

33 Promeneur, D., Mlambo, G., Agre, P. & Coppens, I. Aquaglyceroporin PbAQP is
required for efficient progression through the liver stage of Plasmodium infection. Sci
Rep **8**, 655, (2018).

34 Bray, P. G., Barrett, M. P., Ward, S. A. & de Koning, H. P. Pentamidine uptake and
resistance in pathogenic protozoa: past, present and future. Trends Parasitol **19**, 232-
239, (2003).

35 Biagini, G. A. *et al.* Characterization of the choline carrier of Plasmodium falciparum:
a route for the selective delivery of novel antimalarial drugs. Blood **104**, 3372-3377,
(2004).

36 Chen, W. *et al.* Structural insights into drug transport by an aquaglyceroporin. Nat
Commun **15**, 3985, (2024).

37 Frauenfeld, J. *et al.* A saposin-lipoprotein nanoparticle system for membrane proteins.
Nat Methods **13**, 345-351, (2016).

38 Bayburt, T. H., Grinkova, Y. V. & Sligar, S. G. Self-assembly of discoidal
phospholipid bilayer nanoparticles with membrane scaffold proteins. Nano Lett **2**,
853-856, (2002).

39 Denisov, I. G., Grinkova, Y. V., Lazarides, A. A. & Sligar, S. G. Directed self-
assembly of monodisperse phospholipid bilayer Nanodiscs with controlled size. J Am
Chem Soc **126**, 3477-3487, (2004).

40 Stepien, P. *et al.* Complexity of seemingly simple lipid nanodiscs. Biochim Biophys
Acta Biomembr **1862**, 183420, (2020).

41 Denisov, I. G., McLean, M. A., Shaw, A. W., Grinkova, Y. V. & Sligar, S. G.
Thermotropic phase transition in soluble nanoscale lipid bilayers. J Phys Chem B **109**,
15580-15588, (2005).

42 Chien, C. H. *et al.* An Adaptable Phospholipid Membrane Mimetic System for
Solution NMR Studies of Membrane Proteins. J Am Chem Soc **139**, 14829-14832,
(2017).

43 de la Rosa-Trevin, J. M. *et al.* Scipion: A software framework toward integration,
reproducibility and validation in 3D electron microscopy. J Struct Biol **195**, 93-99,
(2016).

44 Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. Nat Methods **14**, 331-332, (2017).

45 Zhang, K. Gctf: Real-time CTF determination and correction. J Struct Biol **193**, 1-12,
(2016).

46 Sorzano, C. O. S. *et al.* Image Processing in Cryo-Electron Microscopy of Single
Particles: The Power of Combining Methods. Methods Mol Biol **2305**, 257-289,
(2021).

47 Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. J Struct Biol **192**, 216-221, (2015).

48 Sorzano, C. O., Jonic, S., Nunez-Ramirez, R., Boisset, N. & Carazo, J. M. Fast,
robust, and accurate determination of transmission electron microscopy contrast
transfer function. J Struct Biol **160**, 249-262, (2007).
49 Wagner, T. *et al.* SPHIRE-crYOLO is a fast and accurate fully automated particle
picker for cryo-EM. Commun Biol **2**, 218, (2019).
50 Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms
for rapid unsupervised cryo-EM structure determination. Nat Methods **14**, 290-296,
(2017).
51 Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization
improves single-particle cryo-EM reconstruction. Nat Methods **17**, 1214-1221,
(2020).
52 Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM
volume post-processing. Commun Biol **4**, 874, (2021).
53 Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
determination in RELION-3. Elife **7**, (2018).
54 Vilas, J. L. *et al.* MonoRes: Automatic and Accurate Estimation of Local Resolution
for Electron Microscopy Maps. Structure **26**, 337-344 e334, (2018).
55 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. Nature
596, 583-+, (2021).
56 Wood, C. *et al.* Collaborative Computational Project for Electron cryo-Microscopy.
Acta Crystallogr D **71**, 123-126, (2015).
57 Burnley, T., Palmer, C. M. & Winn, M. Recent developments in the CCP-EM
software suite. Acta Crystallogr D **73**, 469-477, (2017).
58 Warshamanage, R., Yamashita, K. & Murshudov, G. N. EMDA: A Python package
for Electron Microscopy Data Analysis. J Struct Biol **214**, 107826, (2022).
59 Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons
and electrons: recent developments in Phenix. Acta Crystallogr D **75**, 861-877,
(2019).
60 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of
Coot. Acta Crystallographica Section D-Biological Crystallography **66**, 486-501,
(2010).
61 Yamashita, K., Palmer, C. M., Burnley, T. & Murshudov, G. N. Cryo-EM single-
particle structure refinement and map calculation using Servalcat. Acta Crystallogr D
77, 1282-1291, (2021).
62 Nicholls, R. A., Fischer, M., McNicholas, S. & Murshudov, G. N. Conformation-
independent structural comparison of macromolecules with ProSMART. Acta
Crystallogr D **70**, 2487-2499, (2014).
63 Varadi, M. & Velankar, S. The impact of AlphaFold Protein Structure Database on
the fields of life sciences. Proteomics, (2022).
64 Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular
crystallography. Acta Crystallogr D **66**, 12-21, (2010).
65 DeLano, W. PyMol: An Open-Source Molecular Graphics Tool DeLano Scientific,
San Carlos, California USA.
66 Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. Protein
Sci **32**, e4792, (2023).
67 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. Protein Sci **30**, 70-82, (2021).
68 Jo, S., Kim, T. & Im, W. Automated builder and database of protein/membrane
complexes for molecular dynamics simulations. PLoS One **2**, e880, (2007).

69   Lee, J. *et al.* CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER,
767   OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive
768   Force Field. J Chem Theory Comput **12**, 405-413, (2016).

769   70   Best, R. B. *et al.* Optimization of the additive CHARMM all-atom protein force field
770   targeting improved sampling of the backbone phi, psi and side-chain chi(1) and chi(2)
771   dihedral angles. J Chem Theory Comput **8**, 3257-3273, (2012).

772   71   Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically
773   disordered proteins. Nat Methods **14**, 71-73, (2017).

774   72   Sondergaard, C. R., Olsson, M. H., Rostkowski, M. & Jensen, J. H. Improved
775   Treatment of Ligands and Coupling Effects in Empirical Calculation and
776   Rationalization of pKa Values. J Chem Theory Comput **7**, 2284-2295, (2011).

777   73   Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity
778   rescaling. J Chem Phys **126**, 014101, (2007).

779   74   Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new
780   molecular dynamics method. . J. Appl. Phys. **52**, 7182-7190, (1981).

781   75   Darden, T., York, D. & Pedersen, L. Particle Mesh Ewald - an N.Log(N) Method for
782   Ewald Sums in Large Systems. Journal of Chemical Physics **98**, 10089-10092,
783   (1993).

784   76   Kumar, S., Bouzida, D., Swendsen, R. H., Kollman, P. A. & Rosenberg, J. M. The
785   Weighted Histogram Analysis Method for Free-Energy Calculations on Biomolecules
786   .1. The Method. J Comput Chem **13**, 1011-1021, (1992).

787   77   Hub, J. S., de Groot, B. L. & van der Spoel, D. g_wham-A Free Weighted Histogram
788   Analysis Implementation Including Robust Error and Autocorrelation Estimates.
789   Journal of Chemical Theory and Computation **6**, 3713-3720, (2010).

790   78   Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through
791   multi-level parallelism from laptops to supercomputers. SoftwareX, 1–2 19–25
792   (2015).

793   79   Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. J Mol
794   Graph Model **14**, 33-38, (1996).

795   80   Song, W. L. *et al.* PyLipID: A Python Package for Analysis of Protein-Lipid
796   Interactions from Molecular Dynamics Simulations. Journal of Chemical Theory and
797   Computation **18**, 1188-1201, (2022).

798   81   Klesse, G., Rao, S. L., Sansom, M. S. P. & Tucker, S. J. CHAP: A Versatile Tool for
799   the Structural and Functional Annotation of Ion Channel Pores. J Mol Biol **431**, 3353-
800   3365, (2019).

801   82   Hunter, J. D. Matplotlib: A 2D graphics environment. Comput Sci Eng **9**, 90-95,
802   (2007).

803   83   Fairlamb, A. H., Carter, N. S., Cunningham, M. & Smith, K. Characterisation of
804   melarsen-resistant Trypanosoma brucei brucei with respect to cross-resistance to other
805   drugs and trypanothione metabolism. Mol Biochem Parasitol **53**, 213-222, (1992).

806   84   Ungogo, M. A., Campagnaro, G. D., Alghamdi, A. H., Natto, M. J. & de Koning, H.
807   P. Differences in Transporters Rather than Drug Targets Are the Principal
808   Determinants of the Different Innate Sensitivities of Trypanosoma congolense and
809   Trypanozoon Subgenus Trypanosomes to Diamidines and Melaminophenyl
810   Arsenicals. Int J Mol Sci **23**, (2022).

811   85   Smart, O. S., Goodfellow, J. M. & Wallace, B. A. The pore dimensions of gramicidin
812   A. Biophys J **65**, 2455-2460, (1993).

813   86   Carter, N. S. & Fairlamb, A. H. Arsenical-resistant trypanosomes lack an unusual
814   adenosine transporter. Nature **361**, 173-176, (1993).

815   87   de Koning, H. P., MacLeod, A., Barrett, M. P., Cover, B. & Jarvis, S. M. Further
816   evidence for a link between melarsoprol resistance and P2 transporter function in
817   African trypanosomes. Mol Biochem Parasitol **106**, 181-185, (2000).
818
819

30th April 2025

820 **Main Figures**

Fig. 1 | Cryo-EM structures of TbAQP2 bound to either glycerol, melarsoprol or
pentamidine. a, Overall structure of the TbAQP2 tetramer viewed either from the
extracellular surface or within the membrane plane. **b**, Structure of protomer A of TbAQP2
viewed as a cartoon (rainbow colouration) with glycerol (sticks) bound. The cryo-EM density
for glycerol is shown as a grey surface. **c**, Cut-away views of the channel in each of the
TbAQP2 structures showing bound substrates and drugs (spheres) with atoms coloured
according to type: red, oxygen; yellow, sulphur; blue, nitrogen; purple, arsenic; carbon, grey
(glycerol), orange (melarsoprol) or cyan (pentamidine). **d**, Cryo-EM densities (grey surface)
for glycerol, melarsoprol and pentamidine in their respective structures. See Figures S5 and
S6 for different views of the substrates and comparisons between densities.

Fig. 2 | Interactions between TbAQP2 and bound substrates and drugs. a, Amino acid
residues containing atoms ≤ 3.9 Å from substrate or drug model in the cryo-EM structures are
depicted and coloured according to the chemistry of the side chain: hydrophobic, green,
positively charged, blue; polar, orange. Interactions with a backbone carbonyl group is shown
as CO (red) and potential hydrogen bonds are shown as red dashed lines. Residues
highlighted in an additional colour (grey, pale blue, pink or dark orange) make contacts in
two or less of the structures (as indicated on the figure), whilst those without additional
highlighting make contacts in all three structures. **b**, Protomer A from each structure was
aligned and the positions of the two glycerol molecules (grey), melarsoprol (orange) and
pentamidine (cyan) are depicted.

Fig. 3 | Comparison between conserved glycerol-binding regions of TbAQP2 and PfAQP. a, The structure of TbAQP2 is depicted (rainbow colouration) with residues shown that make hydrogen bonds (yellow dashed lines) to either glycerol (grey) or in the NPS/NSA motifs. CO, backbone carbonyl groups involved in hydrogen bond formation. **b,** The structure of PfAQP2 is depicted (rainbow colouration) with residues shown that make hydrogen bonds (yellow dashed lines) to either glycerol (yellow sticks) or water (red spheres), or in the NPS/NLA motifs. CO, backbone carbonyl groups involved in hydrogen bond formation. **c,** Structures of TbAQP2 (rainbow colouration) and PfAQP are superimposed to highlight structural conservation of the NPS and NSA/NLA motifs. **d,** Outline of channel cross-sections from aquaglyceroporins containing bound glycerol molecules (sticks): TbAQP2 (grey); PfAQP (yellow; PDB ID 3CO2); human AQP7 (magenta, PDB ID 6QZI); human AQP10 (green, PDB ID 6F7H); *Escherichia coli* GlpF (purple, PDB ID 1FX8).

Fig. 4 | Molecular dynamics simulations of pentamidine-bound TbAQP2. a, Density of
water molecules (blue) through the TbAQP2 central cavity from molecular dynamics. Bound
pentamidine (sticks, green) abolishes the ability of TbAQP2 to transport water molecules. **b**,
View of pentamidine (cyan, carbon; red, oxygen; blue, nitrogen; white, hydrogen) bound to
TbAQP2. Cyan mesh shows the density of the molecule across the MD simulation. **c**, Upon
application of a membrane potential, the pentamidine centre of mass (COM) moves along the
z-dimension in relation to the COM of the channel, with three independent repeats shown in
different shades of blue. The bottom graph is for the potential in the physiological direction
(negative intracellular). **d**, Energy landscapes for pentamidine through the TbAQP2 central
cavity as calculated using umbrella sampling. Separate calculations were made for
monomeric WT TbAQP2 and TbAQP2 with I110W (IW) or L258Y/L264R (LY/LR)
mutations. Each trace is built from 167 x 40 ns windows, with the histogram overlap and
convergence plotted in Fig. S10a,b. The position of the membrane phosphates is shown as
grey bars, and the structural binding pose is shown as a dotted line.

30 April 2025

Fig. 5 | Comparison between drug-bound AQP2 structures. Superposition of an AQP2 protomer from this work (cyan) with a protomer determined by Chen *et al.*[36] (grey), with the positions of drugs shown in stick representation: **a**, pentamidine; **b**, melarsoprol.

30th April 2025

968   **Supplementary Figures and Tables**

Fig. S1 | Alignment of aquaporin amino acid sequences. Amino acid sequences of AQP2 and AQP3 from *Trypanosoma brucei brucei* (Tbb) are aligned with chimeric aquaporins from drug resistant strains (P1000, 427MR, R25) and a chimera from drug-resistant *T. b. gambiense* (TbgAQP2/3). Three identical sequences are shown of TbbAQP2 with residues making contact to either glycerol, pentamidine or melarsoprol highlighted (cyan, green or yellow, respectively). Red bars represent transmembrane regions. Residues highlighted in grey are conserved in all sequences. The conserved NSA/NPS motif in TbAQP2 and the analogous NPA/NPA motif in TbAQP3 are in purple font and the residues mutated for the MD simulation studies (I110W, L258Y and L264R) are in red. Sequence information was obtained from Munday et al. [12] except R25[28] and TbgAQP2/3 (678-800) [15].

Fig. S2 | Purification of TbAQP2 and identification of the drug binding site by hotspot analysis. a, SEC trace of affinity purified TbAQP2. **b**, Coomassie-stained SDS-PAGE gel of fractions during detergent solubilisation (Sol), Ni^{2+}-affinity chromatography and from the SEC column. **c**, Cryo-EM electron micrograph showing the particle distribution of TbAQP2. **d**, A region of TbAQP2 was identified as being most likely to bind a ligand is shown (yellow surface) is mapped on to the structure of TbAQP2 determined here (rainbow colouration. **e**, as in **d**, but also containing pentamidine (cyan sticks). **f**, HOLE[85] trace for our resolved TbAQP2 structure, compared to a previously released TbAQP2 structure ('8yj7')[36] and *Plasmodium falciparum* AQP[24].

1093 **Fig. S3 | Flow chart of the cryo-EM image processing and structure determination.**

1118 **Fig. S4 | Local resolution maps of TbAQP2 structures. a**-**c**, 2D class averages for each of
1119 the structures, FSC curves of the reconstructions with estimates for resolution determined
1120 using an FSC of 0.143 and local resolution estimations as calculated by Relion.
1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145 **Fig. S5 | Depictions of the side chain and substrate densities.**

1146

1147
1148
1149

a Melarsoprol compared to glycerol

1161

b Pentamidine compared to glycerol

Fig. S6 | Comparisons between substrate densities. a, Comparison between melarsoprol and glycerol. **b**, comparison between pentamidine and glycerol.

Fig. S7 | Molecular dynamics simulations. a, RMSDs for simulations of tetrameric WT TbAQP2. Trajectories were fitted on the Cα atoms and RMSDs were calculated for either Cα atoms for apo TbAQP2 (top), Cα atoms for pentamidine (pent) bound TbAQP2 (middle), or for the pent molecule itself (bottom). Five independent repeats are shown as blue traces. **b**, Percentage of time (occupancy) a given residue is within 3.9 Å of either pentamidine, water or glycerol during the MD simulations. **c**, Plotting the z-axis position of pentamidine when bound to TbAQP2 in a WT, IW, or LY/LR background. **d**, Pore radius profiles from simulations of apo monomeric WT, IW, or LY/LR TbAQP2. Analyses were run using the CHAP package[81] with traces showing the mean (black line), standard deviation (dark grey), and range (light grey) for all frames over 5 x 800 ns of simulation. **e**, As panel **a**, but for Cα

1214 atoms from monomeric TbAQP2 in a WT background, I110W (IW) or L258Y/L264R
1215 (LY/LR) mutations.

Fig. S8 | Water in molecular dynamics simulations a, hydrogen-bond analysis from *gmx*
hbond between the pentamidine and channel. Across the 5 independent MD runs of the
monomeric TbAQP2 MD simulations, 23 hydrogen-bonds are formed, but only 6 appear to
be high occupancy. These are highlighted on the image along with their index number, and
their quantification is shown below. **b**, hydrogen-bond distances from the bonds in panel **a**
plotted as a frequency plot. The average bond length was 0.29±0.02 nm, suggesting moderate
strength. **c**, snapshots from MD showing water progression through the TbAQP2 channel in
the absence of pentamidine. Three snapshots are shown. Residues that form especially high
contact with water molecules (values given in brackets) are highlighted in each snapshot.
Water-water hydrogen bonds, as computed using VMD, are shown in yellow dashed lines. **d**,
number of water molecules along the TbAQP2 channel ±pentamidine, as computed using
gmx h2order on the monomeric TbAQP2 MD simulations. **e**, orientation of the waters in each
of the slices along with channel, as computed using *gmx h2order*.

Fig. S9 | Molecular dynamics simulations. a, Plot of the membrane potential (V) as applied
via an electric field, and calculated using the Gromacs tool *gmx potential*. Shown are
potentials produced by an electric field applied in the forward (-ve; solid line) or backward
(+ve; dashed line) direction. For context, the densities of lipid phosphate atoms and pent
atoms are shown as orange and green traces, as measured using *gmx density*. **b**, Histograms
for the umbrella sampling simulations used to generate the landscapes in Fig. 4d. The data
show considerable overlap between windows. **c**, Conversion plots for the PMF landscapes.
Shown are landscapes calculated for increasing lengths of simulations (length in ns). After
about 30 ns, the landscapes converged. **d**, Snapshots taken from post umbrella sampling
windows for WT TbAQP2 at -1.35 nm and +2.7 nm.

1281

Fig. S10 | Lysis assay of *T. brucei* cells. a-d, Lysis assay with *T. b. brucei* wild-type and various AQP2 mutant cell lines treated with arsenical compounds: **a** and **c**, cymelarsan; **b** and **d**, phenylarsine oxide, a different arsenic-containing trypanocide that enters the parasite independently of TbAQP2[8,10,12] and is thus used as a control for transporter-related arsenic resistance versus resistance to arsenic *per se*. The cells were placed in a cuvette and treated with either compound at t = 10 min. All points shown are the average of triplicate determinations and SD. When error bars are not visible, they fall within the symbol. The slow decline with cymelarsan over time in AQP2-KO and the mutant cell lines is attributable to residual uptake of the compound through the TbAT1/P2 transporter[86,87].

1319

1320 **Table S1 | Cryo-EM data collection, refinement and validation statistics**

1321

1322

	TbAQP2 + glycerol (EMDB: EMD-16864) (PDB: 8OFZ)	TbAQP2 + melarsoprol (EMDB: EMD-16862) (PDB: 8OFX)	TbAQP2 + pentamidine (EMDB: EMD-16863) (PDB: 8OFY)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron exposure ($e^-/Å^2$)	25,2	57,6	57,6
Defocus range (μm)	[-0.6 , -4]	[-0.6 , -3.6]	[-0.8 , -4]
Pixel size (Å)	1.07	1.042*	1.042*
Symmetry imposed	C4	C4	C4
Initial particle images (no.)	849,607	1,927,102	1,263,534
Final particle images (no.)	101,075	126,551	83,845
Map resolution (Å)	3.2	3.2	3.7
FSC threshold	0.143	0.143	0.143
Map resolution range (Å)†	3.1 – 4.4	3.0 – 8.3	3.5 – 9.8
Refinement			
Initial model used	AlphaFold	AlphaFold	AlphaFold
Model composition in the ASU			
Non-hydrogen atoms	1842	1852	1855
Protein residues	244	244	244
Substrates	2	1	1
B factors ($Å^2$)			
Protein	159.7	140.2	170.4
Ligand	180.8	175.0	209.6
R.m.s. deviations			
Bond lengths (Å)	0.0074	0.0099	0.0060
Bond angles (°)	1.730	1.903	1.525
Validation			
MolProbity score	1.99	1.79	1.79
Clashscore	4.34	6.80	4.75
Poor rotamers (%)	5.52	1.66	2.21
Ramachandran plot			
Favored (%)	96.69	96.28	95.87
Allowed (%)	3.31	3.72	3.72
Disallowed (%)	0	0	0.41

1323

1324 Footnotes:

1325 *Original pixel size in movies: 1.07 Å/px. Calibrated for modelling

1326 †Local resolution range

1327

1328

1329

30th April 2025

1330

1331 **Table S2 | Details of molecular dynamics simulations on TbAQP2.**

1332

1333

System	Box size	No. atoms	Setup	Simulation length
AQP2 WT tetramer	10x10x10 nm	ca. 95,000	Pentamidine	5 x ca. 300 ns
AQP2 WT tetramer	10x10x10 nm	ca. 95,000	Apo	5 x ca. 300 ns
AQP2 WT monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine	5 x 800 ns
AQP2 WT monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine and +ve electric field	3 x ca. 600 ns
AQP2 WT monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine and -ve electric field	3 x ca. 1400 ns
AQP2 WT monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine steered MD	1 x 5 ns pull
AQP2 WT monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine PMF	167 x 40 ns windows
AQP2 WT monomer	6.5x6.5x10 nm	ca. 45,000	Apo	5 x 800 ns
AQP2$_{L258Y\_L264R}$ monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine	5 x 800 ns
AQP2$_{L258Y\_L264R}$ monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine steered MD	1 x 5 ns pull
AQP2$_{L258Y\_L264R}$ monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine PMF	167 x 40 ns windows
AQP2$_{L258Y\_L264R}$ monomer	6.5x6.5x10 nm	ca. 45,000	Apo	5 x 800 ns
AQP2$_{I110W}$ monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine	5 x 800 ns
AQP2$_{I110W}$ monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine steered MD	1 x 5 ns pull
AQP2$_{I110W}$ monomer	6.5x6.5x10 nm	ca. 45,000	Pentamidine PMF	167 x 40 ns windows
AQP2$_{I110W}$ monomer	6.5x6.5x10 nm	ca. 45,000	Apo	5 x 800 ns

1334

1335

1336