Brachyury expression levels predict lineage potential and axis-forming ability

of *in vitro* derived neuromesodermal progenitors

running title: Brachyury dictates neuromesodermal lineage specification

Anahí Binagui-Casas*[1], Anna Granés*[1], Alberto Ceccarelli[2], Eleni Karagianni[3], Daniel Lopez Ramajo[1],

Rosa Portero[1], Matthew French[4], Jen Annoh[1], Frederick C.K. Wong[5], Sally Lowell[1], Osvaldo Chara[2,6],

Valerie Wilson[1]

1. Institute for Regeneration and Repair, Institute for Stem Cell Research, School of Biological

Sciences, University of Edinburgh, 5 Little France Drive, Edinburgh EH16 4UU, Scotland

2. School of Biosciences, University of Nottingham, Sutton Bonington Campus, Nottingham LE12, UK

3. GSK, Hills Rd, Cambridge CB2 0QQ, UK

4. Department of Genetics, University of Cambridge, Cambridge, CB2 3EH UK,

5. The Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge CB10 1SA, UK

6. Instituto de Tecnología, Universidad Argentina de la Empresa, Buenos Aires, Argentina.

* joint first authors

co-corresponding authors: a.binagui-casas@ed.ac.uk, v.wilson@ed.ac.uk

Keywords: Neuromesodermal progenitors, Axis elongation, Pluripotent stem cells, Mesoderm,

Neurectoderm, In vitro differentiation, Single cell

Abstract

Neuromesodermal progenitors (NMPs) produce the spinal cord and musculoskeleton in the elongating
anterior-posterior axis. *In vivo*, NMPs possess dual potency, coinciding with regions coexpressing SOX2
and TBXT. *In vitro*, SOX2[+]TBXT[+] cells can be produced from pluripotent cells and, like their *in vivo*
counterparts, colonise neural and mesodermal embryonic regions. However, the potency of individual
in vitro SOX2[+]TBXT[+] cells is unclear, confounding comparisons with *in vivo* data. To address this, we
developed a dual *Sox2/Tbxt* reporter mouse ESC line and demonstrate that single *in vitro* derived
Sox2/Tbxt-expressing cells are bipotent NMPs. *In vitro* derived NMPs also exhibit lateral plate
mesoderm potency when cultured in inductive conditions. Surprisingly, we identified two successively
higher *Tbxt* threshold expression levels that correspond to mesoderm bias and specification,
respectively. Moreover, clonally plated Sox2[+]/Tbxt[+] cells self-propagate. Media and substrate
composition alter the specification thresholds and consequent differentiation outcomes of *in vitro*
derived mouse NMPs. Using this live reporter line, we uncover rapid inhibition of axis elongation
following TBXT downregulation in response to WNT or NOTCH inhibition. In summary, this tractable
cell line provides evidence supporting a key role for TBXT in tuning cell fate to drive axis elongation.

Introduction

In vertebrates, the head-to-tail (or anteroposterior, AP) body axis forms during early embryonic development. In mice, where this process is well characterised (Wymeersch et al., 2021), a transient population of progenitors, collectively called neuromesodermal competent cells (NMCs), produces both paraxial mesoderm and neurectoderm as the AP axis elongates in post-gastrulation stages. NMC populations across vertebrates harbour cells with functional heterogeneity, including bona fide neuromesodermal progenitors (NMPs) (Binagui-Casas et al., 2021). Fate mapping has shown that neuromesodermal potency is found at the posterior end of various vertebrate embryo species and this region coincides with the co-expression of the transcription factors SOX2 and Brachyury (TBXT) (Wymeersch et al., 2021). Importantly, the hallmark of NMPs is their developmental potential to generate both neuroectodermal and mesodermal lineages, two separate germ layers, at post-gastrulation stages (Guillot et al., 2021; Tzouanacou et al., 2009). Current literature suggests that NMPs in amniotes (chick, mouse) and anamniotes (fish, frog) are distinct. In amniotes, individual bi-potent NMPs are dual neural/mesodermal fated. In contrast, in anamniotes neuromesodermal competency is maintained in axial progenitors in the tailbud, but individual cells adopt either neural or mesodermal fates. (Attardi et al., 2018; Davis & Kirschner, 2000; Guillot et al., 2021; Martin & Kimelman, 2012; Tzouanacou et al., 2009; Wymeersch et al., 2016).

FGF and WNT treatment of pluripotent stem cells *in vitro* generates a population of cells that resemble *in vivo* NMCs both phenotypically and functionally: a high percentage of cells co-express SOX2/TBXT transcription factors, and these populations can differentiate into neural and mesodermal lineages (for a comprehensive list of protocols see Wymeersch et al., 2021).

Recent single cell RNA-seq studies have found that *in vitro* cultures harbouring NMC populations are transcriptionally heterogeneous. One interpretation of this data is that *in vitro* NMC populations contain contaminating -non-NMP cells (Edri, Hayward, Baillie-Johnson, et al., 2019; Edri, Hayward, Jawaid, et al., 2019). Alternatively, *Sox2* and *Tbxt* expressing cells themselves are transcriptionally heterogeneous, perhaps reflecting functional heterogeneity. A proportion of in vitro-derived NMP-like cells expressing *Tbxt* (monitored with a fluorescent reporter) have dual potential for neural and mesodermal differentiation (Tsakiridis & Wilson, 2015). However, the bi-potency of individual NMP-like cells co-expressing *Sox2* and *Tbxt* remains unproven.

Addressing these questions requires the use of cells that report on both SOX2 and TBXT expression. Previous reporters have employed either knock-in constructs that disrupt the function of these genes (Avilion et al., 2003; Fehling et al., 2003), or are additive transgenics where there is a risk that expression does not accurately mirror endogenous expression (Koch et al., 2017).

Here we use a newly generated minimally disruptive mouse embryonic stem cell (mESC) dual

Sox2/Tbxt gene expression reporter cell line to examine live cell behaviours, population heterogeneity

and functional potency of *in vitro* NMP-like cells. Additionally, we show that rapid deceleration of axis

elongation in gastruloids on WNT or NOTCH inhibition corresponds in time with *Tbxt* inhibition, while

Sox2 level dynamics are slower. By clonal plating (using Fluorescence-Activated Cell Sorting -FACS) in

vitro-derived Sox2/Tbxt- expressing putative NMC populations, we demonstrate that individual

Sox2/Tbxt dual positive cells are bona fide bipotent progenitors. We uncover previously elusive lateral

plate mesoderm potency within descendants of NMPs, an attribute previously shown only at the

population level (Row et al., 2018). Interestingly, the *Sox2/Tbxt* co-expressing population is not

functionally homogeneous: the levels of *Tbxt*-GFP predict clonal lineage bias towards mesoderm in

individual progenitors. This bias is not fixed and can be tuned by the culture conditions and cell

substrates, mirroring the *in vivo* plasticity of NMC populations (Wymeersch et al., 2016). Together, we

demonstrate previously unsuspected influences on the fate choices of individual NMPs, and provide

a robust basis for further investigating these factors.

Results

1. Generation and Validation of STR-KI mESC line: a novel *Sox2/Tbxt* gene expression

reporter

To characterise the dynamics of neuromesodermal competent population emergence from mESC, we

generated a novel dual reporter: Sox2::mCherry; Tbxt::GFP, hereafter named Sox-Tbxt Reporter

Knock-In **(STR-KI)**. We employed CRISPR knock-in transgenesis to replace the stop codon of the

endogenous gene loci with T2A self-cleavable peptide fluorescent protein fusions, retaining the *Tbxt*

and *Sox2* open reading frames in their entirety. This minimises the chance of adverse phenotypic

effects of the transgenes and maximises the faithfulness of the reporters (Fig. 1A) (Dewari et al., 2018).

High contribution of STR-KI mESCs in embryo chimeras, whose phenotype was normal, demonstrates

that the gene targeting was indeed minimally disruptive to SOX2 and TBXT protein function.

Furthermore, the fluorescent reporters in pre-streak and somite stage embryos recapitulated the

endogenous in vivo expression pattern of *Tbxt* and *Sox2* in mesodermal and neural lineages,

respectively (Fig. 1B, Suppl. Fig. 1A,B) (Wymeersch et al., 2016). In pre-streak high contributing

chimeras, *Sox2*-mCHERRY was expressed in the entire epiblast whereas *Tbxt*-GFP expression was

undetectable (Suppl Fig. 1A). By early somite stages onwards, *Sox2*-mCherry was predominantly

restricted to neural tissues, and primordial germ cells at the base of the allantois, whereas *Tbxt*-GFP

was expressed in the node, notochord and nascent mesoderm at the posterior end (Fig.1 C and Suppl.

1B). Culturing the STR-KI line following established(Gouti et al., 2014)(Gouti et al., 2014)   produced

mCherry and GFP double positive cells (putative NMP-like cells). Immunostaining of these in vitro-
differentiated NMP-like cultures showed that the reporters faithfully co-localised with endogenous
SOX2 and TBXT protein in colonies (Fig. 1C, D).

We next tracked the reporters' dynamics over a time course of NMP differentiation (Fig. 1 E). Briefly,
this entails differentiation of ESC to an epiblast-like state followed by treatment with the WNT agonist
CHIR from day (d) 2 onwards (Gouti et al., 2014). Cells apparently moved along a differentiation
trajectory from pluripotent ESC (Sox2^{mCherry} high) to a Sox2^{mCherry} low (putative epiblast) state, followed
by appearance of Sox2^{mCherry}/Tbxt^{GFP} double positive cells within 3 days, resembling the SOX2 and TBXT
level dynamics seen in epiblast NMC-containing regions *in vivo,* where there is heterogeneity at the
transcriptional and protein levels for Sox2 and Tbxt, likely denoting their functional potential (French
et al., 2024; Wymeersch et al., 2016). At d3, cells exhibited a spectrum of Tbxt^{GFP} levels before exiting
the double-positive state towards Sox2^{mCherry} negative Tbxt^{GFP} high mesoderm from d4. Following the
established Gouti et al. 2014 protocol with the STR-KI mESC line reproducibly resulted in the
production of ~75% of the population co-expressing mCherry and GFP putative NMCs by day 3,
consistent with previous reports (Fig. 1F).

Figure 1

Figure 1. STR-KI, a novel Sox2 and Tbxt mESC fluorescent reporter cell line that allows the study of

NMP-like cell differentiation trajectories. A) *Schematic illustration of the targeting construct for the*

insertion of GFP and mCherry in the Tbxt and Sox2 loci, respectively. **B)** *Embryonic day 9 STR-KI*

chimaera showing high contribution of STR-KI to the entire embryo and showing appropriate location

of the fluorescent reporters. Note high mCherry in neural tissues like brain, neural tube and primordial

germ cells (PGCs), whereas GFP labels the notochord and Tailbud (TB). **C)** *Protein correlation between*

endogenous SOX2 and TBXT transcription factors, and Sox2-mCherry and Tbxt-GFP fluorescent

proteins, respectively. Cells for analyses taken from in vitro differentiation assays. **D)** *Example of*

colonies derived from following Gouti et al. 2014 NMP differentiation protocol. Stained for TBXT and

*GFP, and SOX2 and mCherry to show co-localisation in immunostaining. **E)** Flow cytometry time-course*

to monitor reporter expression along the Gouti et al. 2014 differentiation protocol from mESC (Day0)

into Day 5. A representative example (from n=4) is shown together with a trajectory summary that

*highlights the population dynamics. **F)** Efficiency of STR-KI mESC differentiation into NMP-like (Sox2-*

mCherry and Tbxt-GFP double positive) at Day 3 following Gouti et al. 2014 protocol. n=26

2. Gene expression and lineage potential analysis of Sox2^{mCherry}/Tbxt^{GFP} fluorescent subpopulations

during differentiation

To characterise the cell populations that emerge in the differentiation trajectory, we performed FACS

during the differentiation protocol and analysed the gene expression profiles of various populations

across the trajectory we identified in Fig. 1E via qPCR of diagnostic marker genes at d2-d4 (Fig. 2A,B;

Suppl. 2 A,B). The marker genes analysed were: pluripotency-associated *Nanog*, *Nodal* and *Oct4*; axial

progenitor markers *Sox2*, *Tbxt*, *Fgf8*, *Evx1*, *Cdx2*, *Wnt3a* and *Nkx1.2*; paraxial mesoderm marker *Tbx6*

and lateral mesoderm marker *Flk-1*; and neural-associated *Otx2, Sox1* and *Pax6*. In all sorted

populations, the endogenous gene expression activity of *Sox2* and *Tbxt* was accurately recapitulated

by mCherry and GFP transcripts. mCherry transcription was higher in populations that also exhibited

high Sox2, remained Tbxt negative and expressed pluripotency or neural markers across all days tested

(Sox2^{mCherry}+;Tbxt^{GFP} negative). The putative NMC population (Sox2^{mCherry}+;Tbxt^{GFP}+) displayed

relatively constant Sox2-mCherry fluorescence but a spectrum of Tbxt-GFP levels by flow cytometry.

To further characterise the expression profile of cells that gradually increase *Tbxt*-GFP fluorescence

along the differentiation trajectory, we subdivided the NMC-like population into low (Sox2^{mcherry} +

Tbxt^{GFP}-low) and high (Sox2^{mcherry} + Tbxt^{GFP}-high) expressing subpopulations. Since the levels of *Tbxt*-

GFP in the double-positive population increased before the appearance of *Tbxt*-GFP single positive

mesoderm cells, we hypothesised that low and high NMC-like population represented maturation

states within the same population. Specifically, we proposed that high *Tbxt*-GFP positive cells were

further along a mesoderm differentiation trajectory than Tbxt-GFP low cells, as suggested *in vivo*

(Romanos et al., 2021; Wymeersch et al., 2016) . Bulk gene expression analyses by qPCR

supported this hypothesis, since although both Sox2^{mcherry}+Tbxt^{GFP} low and high populations expressed

similar levels of NMC markers, Tbxt^{GFP} cells were enriched for mesodermal markers such as *Wnt3a*

and *Tbx6*, while expression of neural genes like *Sox1* were low. Finally, the absence of Sox2, coupled

with the high expression of gene markers like *Tbx6* and *Flk-1* in Sox2^{mCherry} negative Tbxt^{GFP}+ and

Sox2^{mCherry} negative Tbxt^{GFP} negative (double negative) populations, together with the gradual

downregulation of NMC/caudal markers like *Fgf8* and *Wnt3a* from day 3 to day 4, suggested that these

two populations represented nascent and differentiated mesoderm respectively. The expression of

pluripotency- and early mesoderm/node marker *Nodal*, together with endoderm markers *Sox17* and

Foxa2 was low, suggesting low-to-negligible fractions of contaminant non-NMC cells. Thus, gene

expression in populations undergoing differentiation confirms that our double reporter cell line

faithfully recapitulates endogenous gene expression and can be used to purify expected neural,

mesodermal and NMC-like cell populations based on *Sox2-mCherry* and *Tbxt-GFP* expression levels

(Fig. 2B, Suppl. 2A,B).

To test whether the apparently dynamic changes in Sox2^{mcherry} and Tbxt^{GFP} cell populations during NMP

differentiation (Fig.1 E) represent one or more lineage trajectories, the lineage potentials of sorted

subpopulations at day 3 of NMP differentiation were assayed over 3 further days in continuous neural

and mesodermal (FGF and CHIR) inducing conditions   (Gouti et al., 2014; Tsakiridis & Wilson, 2015)

(Fig 2C, Suppl. Fig. 2C). These subpopulations were: 1) Sox2^{mcherry}+ Tbxt^{GFP} negative; 2) Sox2^{mCherry}+

Tbxt^{GFP}-low; 3) Sox2^{mCherry}+ Tbxt^{GFP}-high; and 4) Sox2^{mcherry} negative Tbxt^{GFP}+. Sorted populations

showed differential responses to these stimuli. Sox2^{mcherry} single positive cells mainly remained single

positive. Together with the enrichment of Sox1 in the starting population, this suggests a neural

identity for these cells. However, a proportion of the progeny lost expression of both reporters (double

negative cells), suggesting that CHIR/FGF induced further differentiation to a Sox2-negative state.

Sox2^{mCherry} + Tbxt^{GFP}-low cells also produced Sox2^{mcherry} single positive cells by day 3, but also a

recapitulation of the pattern shown in Fig. 1E: emergence of Sox2^{mCherry} + Tbxt^{GFP} -high cells, followed

by a high proportion of Sox2^{mCherry} negative Tbxt^{GFP}+ and later double negative cells. This suggests that

Sox2^{mCherry} + Tbxt^{GFP}-low NMCs produce both neural and mesodermal cells. Interestingly, Sox2^{mCherry} +

Tbxt^{GFP}-high cells produced few Sox2^{mcherry} single positive cells but followed a predominantly

mesoderm trajectory, producing a Sox2^{mCherry} negative Tbxt^{GFP}+ and then double negative cells. Finally,

Sox2^{mcherry} negative Tbxt^{GFP}+ only produced double negative cells. Together with its enrichment in

expression of mesoderm markers, it suggests this is an already mesoderm-committed population.

Taken together, using our dual reporter, we have captured the *in vitro* NMC-lineage trajectory of the

progenitors and their progeny (Fig 2D, see summary). Interestingly, populations expressing low and

high levels of Tbxt^{GFP} at d3 show distinct potency: Sox2^{mCherry} Tbxt^{GFP}-low cells produced a higher

proportion of Sox2^{mch} positive neural descendants compared to Sox2^{mCherry} Tbxt^{GFP}-high cells. This

supports our hypothesis that the *Tbxt*-high NMC population is predisposed towards mesoderm

differentiation. Thus, within the Sox2^{mch}Tbxt^{gfp} double positive NMC population, there are cells with

differing competence to differentiate to neural or mesodermal outcomes. It had been suggested that

different in vitro NMC-like populations are biased in lineage potential depending on their starting

pluripotent state (Edri, Hayward, Baillie-Johnson, et al., 2019). Our data suggests an alternative

possibility: that the bias within the NMC-like population depends on the transcriptional state of cells

within the same population. Thus, heterogeneous competence to respond to stimuli can affect the

outcome of assays testing the functional properties of NMP populations.

Treatment of pluripotent cells with FGF and CHIR leads to the expression of both paraxial and lateral

mesoderm markers (Edri, Hayward, Baillie-Johnson, et al., 2019; Gouti et al., 2017). Since NMPs

predominantly produce paraxial mesoderm, the expression of lateral mesoderm markers has been

attributed to other in-vitro derived progenitor populations that emerge alongside NMP-like cells in

these protocols. In vivo, while homotopic grafts of NMCs overwhelmingly give rise to paraxial

mesoderm (Cambray & Wilson, 2007; Wymeersch et al., 2016), heterotopic grafts to the posterior

primitive streak lead to lateral mesoderm differentiation (Row et al., 2018). Theoretically, this result

can arise from population heterogeneity or from lateral mesoderm potency of NMPs. To distinguish

these possibilities, we subjected FACS-enriched $Sox2^{mCherry}$+ $Tbxt^{GFP}$-high cells, containing putative

NMP-like cells to examine their potential to produce lateral plate mesoderm lineages. We followed a

previously reported protocol, which includes bulk *in vitro* NMP differentiation followed by BMP

treatment (Row et al., 2018). Seeding 25 to 1000 $Sox2^{mCherry}$+ $Tbxt^{GFP}$-high cells, we observed not only

Flk-1 protein labelling in colonies but also the formation of primitive tubes similar to those seen in

classical endothelial assays (Taoudi et al., 2005) (Suppl. Fig. 2E). These findings indicate that Sox2/Tbxt-

positive NMP-like cells can produce lateral mesoderm, favouring the hypothesis that in vitro NMP-like

cells themselves (and not accompanying differentiated cells from other origin) are the source of lateral

mesoderm. Thus, we demonstrate a previously unrecognised differentiation potential within enriched

Sox2+Tbxt+ populations *in vitro,* mirroring the lineage potential of unsorted Sox2+Tbxt+ regions *in*

vivo (Row et al., 2018; Wymeersch et al., 2016) .

Since our data suggests that we can live-track the lineage potential of *in vitro* derived NMC

populations, we hypothesised that we can now test factors that might have an effect in the NMC

population differentiation outcomes. Using our STR-KI line we dynamically tracked Sox2 and Tbxt

during in mESC-derived gastruloid models of axis elongation. In the standard gastruloid protocols

(Baillie-Johnson et al., 2015; van den Brink et al., 2014), aggregates of mESCs grown in N2B27

remain Sox2-positive and Tbxt negative until 48 hours post aggregation (hpa), then exit

pluripotency. At 48 hours, a 24-hour pulse of CHIR is used to activate WNT globally and ensure

robust symmetry breaking. WNT activation nduces Tbxt, transiently in all cells, which then

resolves into polarized expression at the posterior pole by 72 hpa and is followed by

elongation. We first investigated the dynamics of mesoderm differentiation in gastruloids in the

presence/absence of global WNT activation via CHIR (Suppl. Fig. 2F). After treatment with CHIR, most

cells became Tbxt^{GFP} + by 72hpa, and down-regulated Sox2^{mCherry}. In the absence of CHIR, STR-KI

cells remained Sox2^{mCherry} high, but a small portion of cells began to express Tbxt^{GFP}. These trends

were maintained at the population level by 96hpa (Suppl. Fig. 2F). We also observed general events

such as downregulation of Sox2^{mCherry}, symmetry breaking and polarisation of Tbxt^{GFP} to the posterior

growing axis only upon the CHIR pulse. Thus, our cell line is suitable for the generation of gastruloids.

We next used STR-KI gastruloids to track the dynamics of reporter expression in gastruloids upon

manipulation of signalling pathways with known roles in mesoderm differentiation from NMPs: WNT

and Notch(Cooper et al., 2024).We next used STR-KI gastruloids to track the dynamics of reporter

expression in gastruloids upon manipulation of signalling pathways with known roles in mesoderm

differentiation from NMPs: WNT (Wymeersch et al., 2016) and Notch (Cooper et al., 2024). We treated

gastruloids that had broken symmetry and started elongating with either WNT or Notch inhibitors

(IWP2 and LY, respectively) and live imaged them every hour for 24 hours to monitor major axis

length and changes in reporter expression (Fig. 2E,F). Both inhibition of WNT and NOTCH

decelerated major axis growth (elongation) starting as early as between 2 and 3 h post treatment

(Fig. 2G).

Segmentation of Sox2^{mCherry} and Tbxt^{GFP} fluorescence in individual cells to determine axial

progenitor regions, allowed monitoring of double reporter dynamics during gastruloid axis

growth. Treatment with IWP2 or LY rapidly decreased Tbxt^{GFP} signal (within ~2-3h), while

Sox2^{mCherry} signal increased after ~10-15 hours, compared to the DMSO control (Fig2. H). This

coincides in time with growth deceleration (Fig2. H, I). Thus, the rapid Tbxt^{GFP} signal

downregulation coupled with slowdown of axis elongation in response to WNT or NOTCH

inhibition suggests AP axis elongation is driven predominantly by *Tbxt*-expressing cells.

Figure 2

Figure 2. In vitro derived Sox2-mCherry and Tbxt-GFP co-expressing cells are NMCs, and their lineage trajectory can be captured for potency studies. A) *Differentiation protocol scheme and representative gating strategy for FACS and further gene expression analyses performed at day 3 of the differentiation*

*protocol. **B)** qPCR gene expression analysis of the populations in A. Gene markers of Pluripotency,*

Neuromesodermal competent cells (NMC) and differentiated cells (neural, mesodermal, endodermal)

*were chosen for this characterisation. n=3 per each population **C)** Differentiation protocol scheme and*

*representative gating strategy for FACS and further time-course lineage potency studies. **D)** Flow*

cytometry data on one time-course experiment (see Suppl. Fig 2C for 2 more experiments), following

the Sox2-mCherry and Tbxt-GFP expression dynamics. A summary of the lineage outcomes across the

time-course and 3 experiments is provided. Thickness of arrows represents contribution to the

trajectory of the arrow, dashed arrow indicates a trajectory that is likely to happen given the

*proportions of cells at the end of the experiment, but we have not formally proven here. **E)***

Experimental set up for the generation of Gastruloids and embedding in 10% matrigel at 96 hours

with DMSO, Wnt signaling inhibitor IWP2 (5uM) or Notch signaling inhibitor LY (150nm).

*Gastruloids were imaged on a confocal microscope every hour for 24 hours after embedding. **F)***

Quantification pipeline of 3D gastruloid confocal image stacks. Whole tissue segmentation of

phase contrast confocal images using the Segment anything model (arXiv:2304.02643) and

quantification of tissue properties. Then the extending Tbxt-GFP positive pole was automatically

detected, shown over a median z-projection of Tbxt-GFP Sox2-mCherry signal. Along this pole

orientation, a 50um wide sample trace of the gastruloid was isolated, and the distance from the

*gastruloid edge was calculated. All scale bars indicate 100um. **G)** Treatment of gastruloids with*

IWP2 or LY decreases major axis length of the gastruloid with IWP2 showing a greater effect on

*gastruloid eccentricity or 'roundness'. **H)** Heatmaps show the average Tbxt-GFP or Sox2-*

mCherry signal per binned distance from the end pole along time post Matrigel embedding.

Treatment with IWP2 or LY shows a decrease in Tbxt-GFP signal and increase in Sox2-mCherry

*signal compared to the DMSO control. **I)** Average intensity of the isolated gastruloid tip reveals a*

delayed onset of Sox2-mCherry increase in the IWP2 or LY treated conditions in contrast to a near

immediate decrease in Tbxt-GFP signal.

3. Clonal tracing of in vitro derived NMCs reveals balance/bias of NMPs dependent on matrix
and T levels

Our findings at the bulk level above suggest there are lineage biases within the NMC (*Sox2/Tbxt* double

positive) population and that nuances in *Tbxt* levels influence neural or mesodermal differentiation

outcomes. Therefore, we asked whether lineage bias is reflected across the whole *in vitro* NMC-like

population or limited to a subset of already lineage-specified cells expressing either the very highest

or lowest levels of the reporters. We therefore asked whether the levels of Tbxt (or Sox2 influence the
degree of neural versus mesoderm differentiation.

To track the progeny of individual *Sox2* and *Tbxt* co-expressing cells we used FACS coupled to single-
cell deposition into 96-well plates (Fig. 3A). We then correlated this information with the lineage
output observed by immunofluorescence and high throughput microscopy in each well after 5 days in
culture conditions that support both neural and mesodermal differentiation (Fig. 3B). This
experimental set-up allowed for a definitive examination of potency via immunostaining for SOX2 or
TBXT at the clonal level in *in vitro* derived NMCs. A variety of substrates are used in different NMC-
generating protocols, and we therefore compared the clonal lineage output of Sox2^{mCherry} Tbxt^{GFP} co-
expressing cells on fibronectin, gelatin, geltrex and matrigel. Sox2^{mCherry} + Tbxt^{GFP} low and Sox2^{mCherry} +
Tbxt^{GFP} high input cells showed distinct lineage output profiles. Sox2^{mCherry} + Tbxt^{GFP} high cells gave rise
to a greater proportion of Tbxt^{GFP} only colonies, confirming our previous data showing that the
increased mesoderm differentiation in Sox2^{mCherry} + Tbxt^{GFP} high cells at the population level is a
function of high TBXT expression in individual cells (Fig3. C). Interestingly, Sox2^{mCherry} + Tbxt^{GFP} high
cells also showed higher plating efficiency than Sox2^{mCherry} + Tbxt^{GFP} low cells, suggesting different
adhesive properties to the cell substrate (Fig3. C, Suppl. Fig 3A, B). Substrate composition differentially
affected cell phenotype: specifically, matrigel led to a large number of colonies where
Sox2^{mCherry}Tbxt^{GFP} double positive cells persisted over the culture period. Intriguingly, although the
plating efficiency of Sox2^{mCherry} + Tbxt^{GFP} low or high was particularly poor on gelatin, this substrate
appeared to enrich for colonies containing double positive cells (Fig3. C, Suppl. Fig. 3A, C).
Furthermore, the number of cells per colony was high using gelatin as substrate (Suppl. Fig. 3A), and
cells grew as compact colonies rather than dispersed cells, as on other substrates, suggesting that
once established from Sox2^{mCherry} + Tbxt^{GFP} low cells, double positive cells are favoured by self-
adherent growth. Sox2^{mCherry} + Tbxt^{GFP} high colony survival on geltrex and matrigel was comparable,
although matrigel led to more than two-fold higher clone survival of Sox2^{mCherry} + Tbxt^{GFP}-low cells.
Cells in the low-to-medium range of Tbxt^{GFP} expression values were abundant enough to permit index
sorting, recording the fluorescence intensity of each reporter for individual sorted cells. Thus, we could
correlate fates to specific input fluorescence values (Fig 3D). Overall, the trends in differentiation
outcome were evident at individual colony level. For example, TbxtGFP high cells either gave rise to
TBXT+ or double negative cells, consistent with the population analysis that showed cells express
Tbxt^{GFP} transiently before differentiating towards presumptive mesoderm. At lower input values,
SOX2+ cells and double positive cells were evident. This confirmed that Sox2^{mCherry} + Tbxt^{GFP} + cells, at
least at lower values of TBXT ^{GFP}, are dual potent, and that these ox2^{mCherry} + Tbxt^{GFP} low cells can self

propagate but further suggested that the outcomes of differentiation appear to be sensitive to
differing levels of TBXT^{GFP}.

We next used the low-to-medium level Tbxt^{GFP} index sorted cells to mathematically determine
whether the clonal lineage of individual NMP progeny is controlled by the initial levels of Tbxt or *Sox2*
expression and the cell substrate. Tbxt^{GFP} and Sox2^{mCherry} expression levels recorded at the time of
plating (time = 0), will be designated *T* and *S*, respectively. After 5 days, each of these seeded cells
gave rise to varying numbers of cells (n) with four phenotypes based on their expression of TBXT and
SOX2: n_{T+S-}, n_{S+T-}, n_{T+S+} and n_{T-S-} .
To determine whether the clonal lineage at day 5 post index sort (time = 5 days) was predominantly
mesodermal or neural, we compared the difference between numbers of Tbxt^{GFP} and Sox2^{mCherry} (nT+S-
versus nT-S+) single positive cells per colony (n_{T+S-} - n_{S+T-}). The histogram of n_{T+S-} - n_{S+T-} determined at
time = 5 days, resembled a normal distribution almost centred at n_{T+S-} - n_{S+T-}=zero (Suppl. Fig. 3 E).

Of note, many NMPs had small n_{T+S-} - n_{S+T-} values (close to zero) and it would be rather arbitrary to
identify these NMPs as mesodermal or neural biased. Instead, the tails of the distribution should
correspond with high confidence to mesodermal (n_{T+S-} >> n_{S+T-}) or neural (n_{T+S-} << n_{S+T-}) lineages.
Therefore, we used the standard deviation of the distribution as a cut-off to assign NMP lineages, even
though this excludes colonies from our analyses. By fitting a Gaussian distribution to the n_{T+S-} - n_{S+T-}
data, we estimated a standard deviation of 47 cells (Suppl. Fig. 3 E). Therefore, we classified the lineage
of the progeny as mesodermal if n_{T+S} > n_{S+T} + 47 cells. Conversely, we consider the lineage to be neural
when n_{S+T-} > n_{S+T} + 47 cells. To investigate whether the lineage of the NMP progeny at time = 5 days
depends on the Tbxt^{GFP} or Sox2^{mCherry} level of the initial index sorted NMP cell at time = 0, we plotted
the lineage classification of each seeded NMP as a function of both Sox2^{mCherry} and Tbxt^{GFP}. To
implement this, we discretized the scales of Sox2^{mCherry} and Tbxt^{GFP} by selecting bins of equal numbers
of colonies.

The mesoderm or neural bias of the NMP progeny did not appear to be affected by the level of
Sox2^{mCherry} in the Sox2^{mCherry}Tbxt^{GFP} double positive starting population, in line with our experiments
at the bulk level and as already suggested *in vivo (French et al., 2024; Kinney et al., 2020; Romanos et*
al., 2021) (Suppl. Fig. 3 F). In contrast, the starting level of Tbxt^{GFP} strongly influenced
neural/mesodermal bias in the clones (Fig. 3D, E). At low values of Tbxt^{GFP}, most NMPs gave rise to
neural-biased clones (Fig. 3E). As Tbxt^{GFP} approached a value of 1760 arbitrary units in the index sort,
the proportion of NMPs contributing to the mesodermal lineages rapidly increased at the expense of
the neural fate (Fig. 3D, E). These results suggest the existence of a Tbxt^{GFP} expression threshold value

above which NMP cells are more likely to produce mesodermal cells, in addition to the higher levels

of Tbxt^{GFP} (Fig. 3D) at which cells are specified towards mesoderm differentiation.

Our results also showed that neural-biased colonies were more abundant on fibronectin and matrigel,

whereas we did not observe a particular bias towards either lineage when cells were plated on gelatin

or geltrex, suggesting that cell substrate can modulate the outcome of lineage-biased

Sox2^{mCherry}Tbxt^{GFP} double positive cells (Suppl. Fig. 3G)

To gain mechanistic insight into fate decisions in *in vitro* derived NMPs, we took a modelling approach.

We developed a minimal mathematical model of the proliferation and differentiation dynamics of

individual NMPs (see Materials and Methods for details). We assumed that each NMP cell can divide

symmetrically to: i) self-renew, giving rise to 2 NMPs (SOX2+ TBXT+); ii) differentiate into 2 mesoderm

cells (TBXT single positive); iii) differentiate into neuroectoderm, (SOX2 single positive); and iv)

differentiate into double negative cells (SOX2 negative TBXT negative). Thus, our model has 4

parameters, which are the rates by which NMP cells give rise to each outcome. To check whether the

model is sufficient to reproduce the dependence of the neural/mesodermal decision on the initial

Tbxt^{GFP} levels in NMPs, we fitted our model to the experimental data.

When we grouped NMPs according to cell substrate, we observed that gelatin was associated with

the highest rates of self-propagation of NMPs and their differentiation towards mesoderm (SOX2

negative TBXT+), as well as descendants of nascent mesoderm, which are double negative (Suppl.Fig.

3H). Matrigel, on the other hand, produced the highest rate of differentiation towards neuroectoderm

(SOX2+ and TBXT negative) (Suppl.Fig. 3H). When we classified NMPs according to whether they had

Tbxt^{GFP} levels below or above the Tbxt^{GFP} threshold reported above (regardless of substrate), we

observed that *Tbxt-GFP* values above the threshold lead to a higher differentiation rate towards

mesoderm, whereas *Tbxt-GFP* values below the threshold lead to neuroectoderm (Fig. 3.F), consistent

with our results shown in Fig. 3E. Interestingly, for both populations, cells produced more NMPs at the

highest rates, and cells above the threshold produced NMPs at a higher rate than those below the

threshold. This suggests that the threshold in differentiation bias does not affect the ability of NMPs

to self-propagate, and distinguishes this lower threshold from the high-Tbxt^{GFP} threshold where cells

predominantly exit the NMP state towards mesoderm differentiation. Despite the ability of

fibronectin and matrigel to bias cells towards neural differentiation, the mid-level threshold still holds

when the experimental data are disaggregated according to the coating used (Supp. Fig. 3I).

In summary, clonal analysis and mathematical modelling demonstrates that individual

Sox2^{mCherry}Tbxt^{GFP} double positive cells in vitro are NMPs, and that the levels of *Tbxt*, but not *Sox2*,

determine whether they are neural-biased, mesoderm-biased or mesoderm-specified. At low to mid-

levels of *Tbxt*, they can self-propagate, but at high levels, they preferentially form mesoderm. The cell

substrate can modulate the proportions of neural versus mesoderm cells formed, while gelatin, a

relatively challenging substrate for NMP adherence, favours self-adherence and more efficient NMP

self-propagation.

Figure 3

394   **Figure 3. In vitro derived Sox2/Tbxt co-expressing cells are bipotent at clonal level and their lineage**

395   **output is determined by Tbxt expression. A)** Differentiation protocol scheme and representative

396   gating strategy for FACS and index sorting according to $Sox2^{mCherry}$ and $Tbxt^{GFP}$ expression. Cells were

individually seeded into different substrates and cultured further for 5 days. Colony lineage output was
assessed by immunofluorescence of endogenous SOX2 and TBXT proteins. After imaging, the data was
analysed by segmenting individual cells in colonies and scoring by their lineage output. as SOX2+,
SOX2+TBXT+ (Double positive-DP), TBXT+, and no fluorescent (no signal detected). **B)** *Example of a*
colony at the end of the experiment and SOX2 /TBXT stained for imaging. **C)** *Lineage output of colonies*
derived from single cell deposition of sorted Sox2^{mCherry}+Tbxt^{GFP} high and Sox2^{mCherry}+Tbxt^{GFP}. Scoring
was done as SOX2+, SOX2+TBXT+ (Double positive-DP), TBXT+, and no fluorescent (no signal detected).
Wells with no colony growth were also scored as 'no growth'. For Sox2^{mCherry}+Tbxt^{GFP} high colonies, n
scored = 120. For Sox2^{mCherry}+Tbxt^{GFP} low colonies, gelatin and fibronectin n=90, matrigel and geltrex
n=118. **D)** *Total individual colonies scored with their lineage output, classified by coating and ordered*
by Tbxt-GFP levels at FACS. Total number of cells per colony is also indicated. Note that the plateau in
GFP levels corresponds to progenitors that were FACS for Sox2^{mCherry}Tbxt^{GFP}high but not index sorted.
Given the colony lineage output, this corresponds to a high threshold that demarks mesoderm
commitment. **E)** *NMP lineage output as a function of initial Tbxt^{GFP} levels. Each dot represents one*
colony/experiment. The coating used is shown in colours dark green for matrigel, light green for
geltrex, pink for felatin and fuchsia for fibronectin. The Tbxt^{GFP} level reported is the middle value in the
bin. **F)** *Rates estimated with a mathematical model indicating whether the initial individual NMP was*
above (in red) or below (in blue) a Tbxt level threshold of 1760 arbitrary units (au) as function of their
colony output at the end of the experiment. The dots and the error bars correspond to the mean and
the standard error of the mean, respectively. stars indicate significant differences (p < 0.05) obtained
after applying a Mann-Whitney statistical test.

Discussion

We report here a new minimally disruptive cell line, STR-KI, that faithfully reports on *Tbxt* and *Sox2*
expression. This cell line mirrors endogenous protein expression and minimises phenotypic effects,
lending confidence to the studies of lineage dynamics performed here. Reduction in SOX2 function
negatively affects ESC maintenance and differentiation (Corsinotti et al., 2017), while reduction of *Tbxt*
levels disrupts tail length (Gruneberg, 1958) and cell movement of *Tbxt* +/- cells in chimeras is
defective (Wilson et al., 1995). Therefore, it is likely that normal protein function is preserved in this
cell line since no abnormal phenotype or distribution of cells was detected in vivo or in vitro.

We show here that NMCs generated *in vitro* from mESC (1) are bi-potent and (2) can self-renew in
clonal analyses. These are functional attributes used to define *in vivo* NMPs. Our study of the
differentiating population over time indicates that ESCs during NMP-like differentiation, while

exposed to FGF, first reduce *Sox2-mCherry* expression, as expected in the epiblast *in vivo* (Corsinotti

et al., 2017). Then, after CHIR addition, the double positive *Sox2-mCherry Tbxt-GFP* population

emerges from these *Sox2-mCherry* low cells. Over the next 48 hours, *Tbxt-GFP* expression increases,

until *Sox2-mCherry* expression disappears from *Tbxt*-GFP high cells. *Tbxt-GFP* is later extinguished,

resulting in a double-negative population. This cellular trajectory fits well with the known fate of *in*

vivo epiblast cells that will eventually become NMPs: distal epiblast cells express relatively low *Sox2*

levels and exit pluripotency to enter a bipotent NMP state in the caudal lateral epiblast or node-streak

border. If they are incorporated in the primitive streak at the midline they will upregulate *Tbxt*,

undergo epithelial to mesenchymal transition and exit the streak (and its WNT-high environment) as

mesoderm, thereafter downregulating *Tbxt* (French et al., 2024; Kispert et al., 1995; Wymeersch et

al., 2016).

In vivo, axial elongation is sensitive to perturbation of both WNT and NOTCH signalling, at least partly

because NMPs require these pathways for maintenance (WNT) and continued mesoderm production

(WNT, NOTCH) (Cooper et al., 2024; Martins-Costa et al., 2024; Wymeersch et al., 2016). TBXT is

directly activated by WNT signalling (Yamaguchi et al., 1999) (Yamaguchi et al. 1999), and optimal

levels require NOTCH signalling (Cooper et al., 2024), and therefore an effect of inhibiting WNT or

NOTCH signalling on TBXT is expected. However, the rapid (2-3h) suppression of Tbxt^{GFP} by both

inhibitors, while Sox2^{mCherry} levels were unchanged for 10-15h, indicates gastruloids require both

pathways to be intact to maintain high levels of TBXT, and furthermore, the immediate slowing of

gastruloid axial elongation suggests that mesoderm production by TBXT is a key driver of continued

gastruloid elongation.

We show that single *Tbxt/Sox2* expressing cells are capable of differentiation towards SOX2-positive

neurectoderm and TBXT-positive (and subsequently double-negative) mesoderm. This formally

demonstrates that double-positive cells generated using standard protocols from mESC are bipotent

NMPs. Interestingly, however, we have also shown, through indexing double positive cells during

sorting, and mathematical modelling of the outcomes, that higher *Tbxt^{GFP}* levels predispose cells

towards mesoderm differentiation. This seems to occur at two distinct levels of Tbxt^{GFP}. Firstly, double-

positive cells with the highest Tbxt^{GFP} fluorescence appear specified for mesoderm differentiation,

reminiscent of a minority of cells at the primitive streak which express high levels of TBXT yet still

express SOX2, but are not bipotent on transplantation to NM-fated regions (Cambray & Wilson, 2007;

French et al., 2024). Secondly, a threshold, mid-level of GFP appears to bias, but not commit, cells to

a mesoderm fate. This is unexpected because in vivo NMCs, even from predominantly mesoderm-

fated regions (which express higher TBXT than those in neural-fated regions), are still potent to
produce neurectoderm on transplantation to more neural-fated regions (French et al., 2024;
Wymeersch et al., 2016). Our study suggests that a previously hidden nuance exists within the *in vitro*
derived NMC population, whereby SOX2/TBXT-positive NMCs exhibit (at least) three functional
phenotypes: neural-biased, mesoderm-biased and mesoderm-specified. The primacy of TBXT dosage
in influencing the outcome of differentiation recalls previous experiments in vivo where levels of TBXT
dictated cell behaviours: cells stay in the primitive streak/tail bud at low TBXT dosage, and exit as
mesoderm at higher doses (Wilson & Beddington, 1997; Wilson et al., 1995). Our study indicates that
this works at individual cell level, whereby cells sense cues in their environment (e.g. WNT, NOTCH),
respond by tuning TBXT up or down, which in turn influences the differentiation outcome at single cell
level.

In conclusion, this dual reporter ESC line provides new opportunities to develop strategies for the
study of the molecular mechanisms underlying self-renewal and differentiation of NMPs.

Materials and methods

<u>Reporter mESC Generation</u>

To generate the double reporter, we performed CRISPR transgenesis in two targeting rounds (one for
each gene locus) in E14tg2a ESC following a stablished protocol (Dewari et al., 2018). The donor DNA
and sgRNA are listed in Table S1. Briefly, we replaced the stop codon at the C-terminus of the targeted
genes by homology recombination with a reporter gene containing T2A-mCherry (in the case of Sox2)
or T2A-H2B-GFP (in the case of *Tbxt*). The *Tbxt* targeting vector contained a puromycin resistant
cassette which was used for clone selection. *Sox2+* Selection was performed by FACS of individual
mCherry+ cells into 96-well plates. Single clones were picked, expanded, and checked for correct
transgene integration by PCR and Sanger sequencing.

<u>Chimaera generation</u>

C57BL/6 female mice (Charles River) were superovulated (100 IU/ml PMSG and 100 IU/ml hCG
intraperitoneal injections 48 h apart) and crossed with wild-type stud male mice. Pregnant mice were
culled at embryonic day 2.5 (E2.5) by cervical dislocation, ovaries with oviducts were dissected and
collected in pre-warmed M2 medium. Oviducts were flushed using PBS and a 20-gauge needle
attached to a 1 ml syringe and filled with PB1 (Whittingham, 1974). E2.5 embryos were collected and

washed in PB1, their zona pellucida removed using acidic Tyrode's solution and transferred to a plate

with incisions where two clumps of approximately eight sender and eight receiver cells were added to

each embryo. Embryos were then incubated at 37°C in 5% CO2 for 48 h prior to fixation, or for 24h

prior to transfer to pseudopregnant females for the generation of post-implantation chimaeras.

Females carrying the transferred embryos were culled following schedule 1 procedures and chimaeric

embryos taken out for analyses. The sex of embryos used in this study was not determined. All

reagents are listed in Table S1

<u>Cell culture, differentiation and flow cytometry</u>

Double reporter mESCs were maintained on 0.1% gelatin-coated plates in Glasgow Minimum Essential

Medium (GMEM, Sigma-Aldrich #G5154) supplemented with 10% foetal calf serum (Gibco #10270-

106), 100 U/mL LIF (made in-house), 100µM 2-mercaptoethanol (Gibco #31350-010), 1X non-essential

amino acids (Gibco #11140-035), 2mM L-Glutamine (Invitrogen #25030-024) and 1mM Sodium

Pyruvate (Invitrogen #11360-039). mESC were passaged every other day with 1X trypsin (Sigma

#59429C) after washing with PBS.

For NMP differentiation, following Gouti et al. 2014, reporter mESCs were plated at a density of 5.000

cells/cm2 on 0.1% gelatin-coated plates in N2B28 medium (1:1 mixture of Neurobasal (Thermofisher

#21103049) and Advanced Dulbecco's Modified Eagle Mediu F12 (Thermofisher #12634028)

supplemented with 2 mM L-Glutamine, 50 µM 2-mercaptoethanol, 0.5X B27 (Thermofisher

#17504001) and 0.5X N2 (Thermofisher #17502048). Cells were supplemented with 10 ng/mL FGF2

(R&D Systems #3718-FB) for 48 hours, and for 24 additional hours with both 10 ng/mL FGF2 and 5µM

CHIRON99021 (Axon #1386).

For single cell plating experiments, differentiated cells were detached using TrypLE Express Enzyme

1X (Gibco 12604013) and sorted on CellCarrier 96-well plates coated with 0.1% gelatin, 16.6 µL/mL

fibronectin (Sigma #F1141) in PBS, 200 µg/mL Matrigel (Corning #354277) in PBS or 120 µg/mL geltrex

(Thermofisher #A1413201) in advanced DMEM/F12, for 1 hour at 37ºC. Cells were cultured in N2B27

supplemented with 100 U/mL penicillin/streptomycin (Invitrogen, #15140-122)), 10 µM ROCK

inhibitor Y-27632 (Tocris #1254/10) and either 10 ng/mL FGF2 or both 10 ng/mL FGF2 and 5µM

CHIRON99021. After 24 hours, media was changed without ROCK inhibitor. The ROCK inhibitor Y-

27632 is used to enhance survival of single cells (Watanabe et al., 2007) and we use it in FACS

experiments as a means of counteracting dissociation and cell stress induced apoptosis.

All cells were maintained under standard culture condition (37ºC and 5% CO2).

For flow cytometry and FACS analysis, cells were detached using trypsin on mESC or TrypLE on
differentiated cells, quenched and pelleted by centrifugating at 300 g for 3 minutes. For flow
cytometry analysis, cells were resuspended in 2% FCS in PBS with 0.1 µg/Ml Draq7 and kept in ice. For
FACS sorting single cells, cells were resuspended in N2B27 with 0.1 µg/mL Draq7.

Cells were analysed with 4 laser LSR Fortessa (BD) flow cytometer using B 530/30-A, Y/G 610/20-A,
and R780/60-A laser/filter combinations or sorted with Cell Sorter BD FACS Aria II using B 525/50-A,
Y/G 610/20-A, and R780/60-A laser/filter combinations.

<u>qPCR</u>

200 cells were sorted into PCR tubes containing 10 µL of 2X Reaction Mix (CellsDirect One-Step qRT-
PCR Kit; Invitrogen #11753100) with 0.4U/µl of RNAse inhibitor (Invitrogen #AM2694) and flash-frozen
on dry ice. Reverse transcription reactions were performed using the Superscript III RT/Platinum Taq
mix (CellsDirect One-Step qRT-PCR Kit; Invitrogen #11753100). qPCR was performed using the
LightCycler 480 SYBR Green I Master (Roche, #4887352001) in LightCycler 480 Multiwell Plate 384
(Roche #04729749001) using the Roche LightCycler 480 Real-time PCR System. TATA-box binding
protein (TBP) gene was used as an endogenous control and relative gene expression was calculated
using the 2-ΔΔCt method.

<u>Immunocytochemistry and microscopy</u>

Cells were fixed with 4% paraformaldehyde for 10 minutes, washed with PBS two times for a total of
10 minutes and permeabilised with 0.5% Triton X-100 in PBS for 10 minutes. After washing two times
with PBS, cells were blocked for 30 minutes with 5% donkey serum and 0.1% Triton X-100 in PBS.
Primary antibodies in blocking solution were added overnight at 5ºC. After washing two times with
PBS, secondary antibodies diluted in blocking solution were added for 2 hours at room temperature.
Cells were washed 2 times in PBS and DAPI diluted in PBS was added for 5 minutes at room
temperature. Finally, cells were washed two times with PBS. Imaging of the 96-well plates was
performed using the High-Content Imaging Opera Phenix Plus from the IRR Imaging Core Facility.
554

<u>Gastruloid culture and live imaging</u>

Gastruloid culture was carried out as previously described (van den Brink et al., 2020 Vianello et al
2020 [dx.doi.org/10.17504/protocols.io.9j5h4q6]). 2i/LIF/FCS CELL LINE cells are washed in PBS before

adding 0.05% Trypsin EDTA solution. After ES colony detachment, 5-10 volumes of fresh 2i/LIF/FCS

were added to quench the Trypsin EDTA solution. The cells and media were transferred to a universal

tube and pelleted by centrifugation at 300g for 3 minutes. The media was aspirated, and the pellet

was resuspended in cold PBS to create a single cell suspension. This PBS wash was repeated once

more, then the cell pellet was resuspended in prewarmed N2B27 to create a single cell solution. The

cells were then counted and then diluted to a solution of 8,250 cells/mL in N2B27 medium. 40μL of

this medium, containing ~330 cells, was plated into untreated u-bottom 96 wells and incubated for 48

hours at 36 °C and 5% CO2. At 48 hours, 150μL of N2B27 and 3uM CHIR99021 (Axon) were added into

the well, the last half of the media was expelled forcefully to dislodge the aggregate but without

spilling the medium. The aggregates were cultured for a further 24 hours to Day 3, after which 150μL

of media was removed and replaced with 150μL fresh N2B27. At Day 4, gastruloids were washed in

N2B27 and then transferred to flat-bottom and black-walled 96-well plates (PerkinElmer) containing

ice-cold N2B27 + 10% Matrigel supplemented with LY 411575 (150nM), IWP2 (2.5μM), or an equal

amount of DMSO. Gastruloids were left for 5 minutes to drop to the bottom of the well and then

placed into an incubation chamber at 36 °C and 5% CO2 to set the Matrigel. After this period,

gastruloids were imaged on a confocal microscope (Opera Phenix Plus) every hour for 24 hours.

Transmitted light and fluorescence signal was collected using epifluorescence and a 20x objective.

Excitations of 488nm and 561nm, emissions of 522nm and 599nm, and exposure times of 0.1s and

0.2s were used for GFP and mCherry respectively. Voxel dimensions were 0.89x0.89x5 microns.

<u>Gastruloid image analysis</u>

Image analysis was performed in python using custom scripts and open source libraries. Whole tissue

segmentation masks were generated using with the Segment Anything Model (SAM). Extending poles

of gastruloids were inferred by median projecting *Tbxt*-GFP signal z-stacks to 2D, smoothing with a

Gaussian filter, and isolating pixels over the 90th percentile. Traces were collected in a 50μm wide

linear line with an orientation defined by the gastruloid mask centroid to the extending-pole centroid.

Extending pole "ends" were isolated as a bin between 80um and 160um from the end of the pole.

<u>Mathematical modelling</u>

In this study, we propose a minimal mathematical model of the dynamics of NMP cells and their

progeny to investigate the cell fate decision process. The main assumption is that each NMP cell

divides symmetrically, resulting in i) 2 NMPs, or ii) 2 cells that are simultaneously T+ and Sox2-, or iii)

2 cells that are simultaneously T- and Sox2+, or iv) 2 cells that are simultaneously T- and Sox2-. We

also assume that, with the exception of NMPs, the other cells do not divide within the time window

of our study. The model is encoded in the following linear system of ordinary differential equations

describing the time course of NMPs and their progeny:

$\frac{dNMP}{dt} = \alpha * NMP - \beta * NMP - \gamma * NMP - \delta * NMP$ (Eq. 1)

$\frac{dn_{S+T-}}{dt} = 2 * \beta * NMP$ (Eq. 2)

$\frac{dn_{S-T+}}{dt} = 2 * \gamma * NMP$ (Eq. 3)

$\frac{dn_{S-T-}}{dt} = 2 * \delta * NMP$ (Eq. 4)

Where t is time, NMP, n_{S+T-}, n_{S-T+} and n_{S-T-} are the number of NMPs, the number of cells that are

simultaneously Sox2+ and T-, the number of cells that are simultaneously Sox2- and T+ and the number

of cells that are simultaneously Sox2- and T-, respectively. The model has 4 parameters: α, β, γ and δ,

which correspond to the rates at which NMP cells give rise to more NMP cells, cells positive for Sox2

but negative for T (*i.e.*, neural lineage), cells negative for Sox2 but positive for T (*i.e.*, mesodermal

lineage) and cells negative for both Sox2 and T (double negative).

To test whether the model was sufficient to reproduce the dependence of the neural/mesodermal

decision on the *T* level of the NMP, we fitted the model to the experimental data.

This system has an analytical solution given by

$NMP(t) = NMP_0 * e^{(\alpha-\beta-\gamma-\delta)t}$ (Eq. 5)

$n_{S+T-}(t) = \frac{2*\beta*NMP_0*e^{(\alpha-\beta-\gamma-\delta)t}}{(\alpha-\beta-\gamma-\delta)} + n_{S+T-0}$ (Eq. 6)

$n_{S-T+}(t) = \frac{2*\gamma*NMP_0*e^{(\alpha-\beta-\gamma-\delta)t}}{(\alpha-\beta-\gamma-\delta)} + n_{S-T+0}$ (Eq. 7)

$n_{S-T-}(t) = \frac{2*\delta*NMP_0*e^{(\alpha-\beta-\gamma-\delta)t}}{(\alpha-\beta-\gamma-\delta)} + n_{S-T-0}$ (Eq. 8)

Where NMP_0 is the initial number of NMP cells, n_{S+T-0} is the initial number of Sox2 positive but T

negative cells, n_{S-T+0} is the initial number of Sox2 negative but T positive cells and n_{S-T-0} is the initial

number of double negative cells.

Finally, we assume the following initial condition, reflecting the experiments described in section 3.

$NMP_0 = 1$ (Eq. 9)

$n_{S+T-_0} = 0$ (Eq. 10)

$n_{S-T+_0} = 0$ (Eq. 11)

$n_{S-T-_0} = 0$ (Eq. 12)

To fit the model to the experimental data, we minimized the distance function δ between the
experimental number of cells and the model predicted number of all cell types for each experiment.
The distance d was defined as:

$$\delta = \sqrt{(NMP_e - NMP_t)^2 + \left(n_{S+T-_e} - n_{S+T-_t}\right)^2 + \left(n_{S-T+_e} - n_{S-T+_t}\right)^2 + \left(n_{S-T-_e} - n_{S-T-_t}\right)^2}$$ (Eq.
13)

Where NMP_e and NMP_t are the experimental and model predicted number of NMP cells, n_{S+T-_e} and
n_{S+T-_t} are the experimental and model predicted number of cells positive for SOX2 but negative for
TBXT, n_{S-T+_e} and n_{S-T+_t} are the experimental and model predicted number of cells positive for T but
negative for SOX2, and finally n_{S-T-_e} and n_{S-T-_t} are the experimental and model predicted number
of cells negative for both SOX2 and TBXT. From all best-fitting parameter values that minimized d for
each experiment, we calculated the mean and standard error of the mean.

A Jupyter Notebook (http://jupyter.org/) containing the source code used to generate the
mathematical model plots in Fig. 3 and Supp. Fig. 3 can be found at Zenodo (Ceccarelli & Chara, 2024).

Table S1

Reagent	Source	Identifier or catalog number
DNeasy Blood & Tissue Kit	QIAGEN	69504
Q5® Hot Start High-Fidelity 2X Master Mix	BioLabs	M0494S
Agarose	Invitrogen	16500-500
SYBR™ Safe DNA Gel Stain	Invitrogen	S33102
Zymoclean Gel DNA Recovery Kit Protocol	Zymo Research	D4001
Qubit 1X dsDNA HS Assay Kit	Invitrogen	Q33230
Glasgow Minimum Essential Medium	Sigma-Aldrich	G5154
Foetal calf serum	Gibco	10270-106
LIF	Nade in-house	N/A

2-mercaptoethanol	Gibco	31350-010
Non-essential amino acids	Gibco	11140-035
L-Glutamine	Invitrogen	25030-024
Sodium Pyruvate	Invitrogen	11360-039
Trypsin	Sigma	59429C
Neurobasal	Thermofisher	21103049
Advanced Dulbecco's Modified Eagle Mediu F12	Thermofisher	12634028
B27	Thermofisher	17504001
N2	Thermofisher	17502048
FGF2	R&D Systems	3718-FB
CHIRON99021	Axon	1386
TrypLE Express Enzyme 1X	Gibco	12604013
Fibronectin	Sigma	F1141
Matrigel	Corning	354277
Geltrex	Thermofisher	A1413201
Penicillin/streptomycin	Invitrogen	15140-122
ROCK inhibitor Y-27632	Tocris	1254/10
2X Reaction Mix (CellsDirect One-Step qRT-PCR Kit)	Invitrogen	11753100
RNAse inhibitor	Invitrogen	AM2694
Superscript III RT/Platinum Taq mix (CellsDirect One-Step qRT-PCR Kit)	Invitrogen	11753100
LightCycler 480 SYBR Green I Master	Roche	4887352001
LightCycler 480 Multiwell Plate 384	Roche	04729749001

Acknowledgements

The authors thank the IRR core facilities for their important contributions to this work, particularly

The IRR Flow cytometry, Tissue Culture and High content Facilities.

Competing interests

No competing interests declared.

Funding

This work was supported by the Medical Research Council (MR/S008799/1 to A.B-C, A.G., V.W,

MR/K011200E.K, F.C.K.W), Carnegie Trust for the Universities of Scotland (RIG012666 to A.B-C), the

Agencia Nacional de Promoción Científica y Tecnológica of Argentina (PICT-2019-03828 to A.S.C. and

O.C), Biotechnology and Biological Sciences Research Council (BB/X014908/1 to O.C.), Wellcome Trust

Senior Fellowship (220298 to M.F., J.A. and S.L), Erasmus Exchange fund (to A.G. and D.L.R), Gurdon

summer studentship (to A.G).

Data availability

Data availability: All relevant data can be found within the article and its supplementary information.

Supplementary Figures

Suppl. Figure 1

A **Pre-streak stage STR-KI chimaera**

B **Somite stage STR-KI chimaera**

654 **Suppl. Figure 1. STR-KI cell line contributes highly to chimaeras. A)** mouse **p**re-streak and **B) E8.5 (2-**

655 **6)** somite stage chimeras appear morphologically normal, show high contribution of STR-KI cells, and

656 correct location of the reporter fluorescent proteins. Asterisk in B shows a non-contributing littermate

657 as negative control.

Suppl. Figure 2

Suppl. Figure 2. *In vitro* derived *Sox2*-mCherry and *Tbxt*-GFP co-expressing cells are NMC progenitors, and their lineage trajectory can be captured for potency studies **A)** qPCR gene expression analysis of populations on day 2 and day 4 using the same gating strategy and differentiation protocol in Fig2A. Classical gene markers of Pluripotency, Neuromesodermal competent cells (NMC) and differentiated cells (neural, mesodermal, endodermal) were chosen for this characterisation. n=3 per each population **C)** Differentiation protocol scheme and representative gating strategy for FACS and further time-course lineage potency studies. **D)** Repeats of the time-course experiment described in Fig. 2C-D, note the variability between 3 experiments, which is inherent to the differentiation protocol. However, cell populations did follow the same trajectory trends (summarised in Fig. 2D). A representative purity check of the initial FACS populations is shown. **D)** Scheme of the experimental set up for FACS $Sox2^{mCherry}Tbxt^{GFP}$ double positive progenitors and testing of lateral plate mesoderm (LPM) potency. **E)** Example of colonies stained for the LPM marker Flk-1. In some cases, cells within colonies were Flk-1+ without changes in morphology (top) whereas in other cases elongated Flk-1+ cells were observes, which could indicate endothelial cell differentiation. **F)** Experimental setup for the generation of gastruloids to test symmetry breaking and $Tbxt^{GFP}$ activation upon CHIR activation. Flow cytometry on an average of 60 gastruloids were pooled and bulk analysed for 72hpa and 30 gastruloids for 96hpa. n=1 **G)** Immunostaining of Wild type (WT) or STR-KI derived gastruloids for $Sox2^{mCherry}$ and $Tbxt^{GFP}$ at 72h and 96h. Note the absence of Tbxt^{GFP} in both WT and no CHIR treated gastruloids. Axis elongation is visible between 96-120, depending on the cell line use. STR-KI does show elongation at 96h upon CHIR treatment.

Suppl. Figure 3

Suppl. Figure 3. Quantitative analysis and mathematical modelling of clone data.

A) Number of cells per colony segregated by coating. B) Number of cells per colony segregated by colony phenotype. C) number of SOX2+TBXT+ cells per colony by coating. D) number of SOX2+TBXT+ cells per colony by colony phenotype. **E)** Histogram for n_{S-T+} - n_{S+T-}. The blue line is the gaussian function fitted to the data. **F)** Clonally plated Sox2 ^{mCherry} Tbxt^{GFP} (NMP) lineage output as a function of Sox2^{mCherry} levels in individually plated NMPs. Each dot represents one colony/experiment. The Sox2^{mCherry} level reported is the middle value in the bin. **G)** Clonally plated Sox2 ^{mCherry} Tbxt^{GFP} (NMP) lineage output as a function of the coting used. **H)** Rates estimated with a mathematical model, coloured by coating used: dark green for matrigel, light green for geltrex, pink for gelatin and fuchsia for fibronectin. Dots are the mean value obtained, and the error bars show the standard error of the mean. **I)** Rates estimated with a mathematical model disaggregated by coating used: fibronectin, gelatin, geltrex and matrigel. The initial NMP cells were above (in red) or below (in blue) a $Tbxt^{GFP}$ level threshold of 1760 arbitrary units (au). The dots are the mean value obtained, and the error bars show the standard error of the mean. Stars indicate significant differences ($p < 0.05$) obtained after applying a Mann-Whitney statistical test.

References

Attardi, A., Fulton, T., Florescu, M., Shah, G., Muresan, L., Lenz, M. O., Lancaster, C., Huisken, J., van Oudenaarden, A., & Steventon, B. (2018). Neuromesodermal progenitors are a conserved source of spinal cord with divergent growth dynamics. *Development*, *145*(21). https://doi.org/10.1242/dev.166728

Baillie-Johnson, P., van den Brink, S. C., Balayo, T., Turner, D. A., & Martinez Arias, A. (2015). Generation of Aggregates of Mouse Embryonic Stem Cells that Show Symmetry Breaking, Polarization and Emergent Collective Behaviour In Vitro. *J Vis Exp*(105). https://doi.org/10.3791/53252

Binagui-Casas, A., Dias, A., Guillot, C., Metzis, V., & Saunders, D. (2021). Building consensus in neuromesodermal research: Current advances and future biomedical perspectives. *Current Opinion in Cell Biology*, *73*, 133-140. https://doi.org/10.1016/j.ceb.2021.08.003

Cambray, N., & Wilson, V. (2007). Two distinct sources for a population of maturing axial progenitors. *Development*, *134*(15), 2829-2840. https://doi.org/10.1242/dev.02877

Cooper, F., Souilhol, C., Haston, S., Gray, S., Boswell, K., Gogolou, A., Frith, T. J. R., Stavish, D., James, B. M., Bose, D., Kim Dale, J., & Tsakiridis, A. (2024). Notch signalling influences cell fate decisions and HOX gene induction in axial progenitors. *Development*, *151*(3). https://doi.org/10.1242/dev.202098

Corsinotti, A., Wong, F. C., Tatar, T., Szczerbinska, I., Halbritter, F., Colby, D., Gogolok, S., Pantier, R., Liggat, K., Mirfazeli, E. S., Hall-Ponsele, E., Mullin, N. P., Wilson, V., & Chambers, I. (2017). Distinct SoxB1 networks are required for naive and primed pluripotency. *Elife*, *6*. https://doi.org/10.7554/eLife.27746

Davis, R. L., & Kirschner, M. W. (2000). The fate of cells in the tailbud of Xenopus laevis. *Development*, *127*(2), 255-267. https://doi.org/10.1242/dev.127.2.255

Dewari, P. S., Southgate, B., McCarten, K., Monogarov, G., O'Duibhir, E., Quinn, N., Tyrer, A., Leitner, M. C., Plumb, C., Kalantzaki, M., Blin, C., Finch, R., Bressan, R. B., Morrison, G., Jacobi, A. M., Behlke, M. A., von Kriegsheim, A., Tomlinson, S., Krijgsveld, J., & Pollard, S. M. (2018). An efficient and scalable pipeline for epitope tagging in mammalian stem cells using Cas9 ribonucleoprotein. *Elife*, *7*. https://doi.org/10.7554/eLife.35069

Edri, S., Hayward, P., Baillie-Johnson, P., Steventon, B. J., & Martinez Arias, A. (2019). An epiblast stem cell-derived multipotent progenitor population for axial extension. *Development*, *146*(10). https://doi.org/10.1242/dev.168187

Edri, S., Hayward, P., Jawaid, W., & Arias, A. M. (2019). Neuro-mesodermal progenitors (NMPs): a comparative study between pluripotent stem cells and embryo-derived populations. *Development*, *146*(12). https://doi.org/ARTN dev180190

10.1242/dev.180190

French, M., Portero, R., Dale, J. K., Blin, G., Wilson, V., & Lowell, S. (2024). A toolkit for mapping cell identities in relation to neighbours reveals Notch-dependent heterogeneity within neuromesodermal progenitor populations. *bioRxiv*, 2024.2009.2003.610492. https://doi.org/10.1101/2024.09.03.610492

Gouti, M., Delile, J., Stamataki, D., Wymeersch, F. J., Huang, Y. L., Kleinjung, J., Wilson, V., & Briscoe, J. (2017). A Gene Regulatory Network Balances Neural and Mesoderm Specification during Vertebrate Trunk Development. *Developmental Cell*, *41*(3), 243-+. https://doi.org/10.1016/j.devcel.2017.04.002

Gouti, M., Tsakiridis, A., Wymeersch, F. J., Huang, Y. L., Kleinjung, J., Wilson, V., & Briscoe, J. (2014). In Vitro Generation of Neuromesodermal Progenitors Reveals Distinct Roles for Wnt Signalling in the Specification of Spinal Cord and Paraxial Mesoderm Identity. *Plos Biology*, *12*(8). https://doi.org/ARTN e1001937

10.1371/journal.pbio.1001937

Gruneberg, H. (1958). Genetical studies on the skeleton of the mouse. XXIII. The development of brachyury and anury. *J Embryol Exp Morphol*, *6*(3), 424-443. https://www.ncbi.nlm.nih.gov/pubmed/13575656

Guillot, C., Djeffal, Y., Michaut, A., Rabe, B., & Pourquie, O. (2021). Dynamics of primitive streak regression controls the fate of neuromesodermal progenitors in the chicken embryo. *Elife*, *10*. https://doi.org/10.7554/eLife.64819

Kinney, B. A., Al Anber, A., Row, R. H., Tseng, Y. J., Weidmann, M. D., Knaut, H., & Martin, B. L. (2020). Sox2 and Canonical Wnt Signaling Interact to Activate a Developmental Checkpoint Coordinating Morphogenesis with Mesoderm Fate Acquisition. *Cell Rep*, *33*(4), 108311. https://doi.org/10.1016/j.celrep.2020.108311

Kispert, A., Ortner, H., Cooke, J., & Herrmann, B. G. (1995). The chick Brachyury gene: developmental expression pattern and response to axial induction by localized activin. *Dev Biol*, *168*(2), 406-415. https://doi.org/10.1006/dbio.1995.1090

Martin, B. L., & Kimelman, D. (2012). Canonical Wnt signaling dynamically controls multiple stem cell fate decisions during vertebrate body formation. *Dev Cell*, *22*(1), 223-232. https://doi.org/10.1016/j.devcel.2011.11.001

Martins-Costa, C., Wilson, V., & Binagui-Casas, A. (2024). Neuromesodermal specification during head-to-tail body axis formation. *Curr Top Dev Biol*, *159*, 232-271. https://doi.org/10.1016/bs.ctdb.2024.02.012

Romanos, M., Allio, G., Roussigne, M., Combres, L., Escalas, N., Soula, C., Medevielle, F., Steventon, B., Trescases, A., & Benazeraf, B. (2021). Cell-to-cell heterogeneity in Sox2 and Bra expression guides progenitor motility and destiny. *Elife*, *10*. https://doi.org/10.7554/eLife.66588

Row, R. H., Pegg, A., Kinney, B. A., Farr, G. H., Maves, L., Lowell, S., Wilson, V., & Martin, B. L. (2018).
BMP and FGF signaling interact to pattern mesoderm by controlling basic helix-loop-helix
transcription factor activity. *Elife*, *7*. https://doi.org/ARTN e31018

10.7554/eLife.31018
Taoudi, S., Morrison, A. M., Inoue, H., Gribi, R., Ure, J., & Medvinsky, A. (2005). Progressive
divergence of definitive haematopoietic stem cells from the endothelial compartment does
not depend on contact with the foetal liver. *Development*, *132*(18), 4179-4191.
https://doi.org/10.1242/dev.01974
Tsakiridis, A., & Wilson, V. (2015). Assessing the bipotency of in vitro-derived neuromesodermal
progenitors. *F1000Res*, *4*, 100. https://doi.org/10.12688/f1000research.6345.2
Tzouanacou, E., Wegener, A., Wymeersch, F. J., Wilson, V., & Nicolas, J. F. (2009). Redefining the
Progression of Lineage Segregations during Mammalian Embryogenesis by Clonal Analysis.
Developmental Cell, *17*(3), 365-376. https://doi.org/10.1016/j.devcel.2009.08.002
van den Brink, S. C., Baillie-Johnson, P., Balayo, T., Hadjantonakis, A. K., Nowotschin, S., Turner, D. A.,
& Martinez Arias, A. (2014). Symmetry breaking, germ layer specification and axial
organisation in aggregates of mouse embryonic stem cells. *Development*, *141*(22), 4231-
4242. https://doi.org/10.1242/dev.113001
Whittingham, D. G. (1974). Embryo banks in the future of developmental genetics. *Genetics*, *78*(1),
395-402. https://doi.org/10.1093/genetics/78.1.395
Wilson, V., & Beddington, R. (1997). Expression of T protein in the primitive streak is necessary and
sufficient for posterior mesoderm movement and somite differentiation. *Developmental
Biology*, *192*(1), 45-58. https://doi.org/DOI 10.1006/dbio.1997.8701
Wilson, V., Manson, L., Skarnes, W. C., & Beddington, R. S. P. (1995). The T-Gene Is Necessary for
Normal Mesodermal Morphogenetic Cell Movements during Gastrulation. *Development*,
121(3), 877-886. <Go to ISI>://WOS:A1995QN06700023
Wymeersch, F. J., Huang, Y. L., Blin, G., Cambray, N., Wilkie, R., Wong, F. C. K., & Wilson, V. (2016).
Position-dependent plasticity of distinct progenitor types in the primitive streak. *Elife*, *5*.
https://doi.org/ARTN e10042

10.7554/eLife.10042
Wymeersch, F. J., Wilson, V., & Tsakiridis, A. (2021). Understanding axial progenitor biology in vivo
and in vitro. *Development*, *148*(4). https://doi.org/ARTN dev180612

10.1242/dev.180612
Yamaguchi, T. P., Takada, S., Yoshikawa, Y., Wu, N., & McMahon, A. P. (1999). T (Brachyury) is a
direct target of Wnt3a during paraxial mesoderm specification. *Genes Dev*, *13*(24), 3185-
3190. https://doi.org/10.1101/gad.13.24.3185

805