Telomerase reverse transcriptase is required for resistance to mycobacterial infection

Darryl JY Han [1], David M Costa [1], Geyang Luo [2], Kathryn Wright [3], Manitosh Pandey [1], Denise Wee [1], Yao Chen [1], Semih Can Akincilar [4], Vinay Tergaonkar [4], Rajkumar Dorajoo [5,6], Guillaume Carissimo [1], Amit Singhal [1], Philip M Elks [7], Pamela S Ellis [8], Catarina M Henriques [8], Bo Yan [2], Serge Mostowy [3], Stefan H Oehlers [1]*

[1] A*STAR Infectious Diseases Labs (A*STAR ID Labs), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore.
[2] Center for Tuberculosis Research, Shanghai Public Health Clinical Center, Fudan University, Shanghai, People's Republic of China.
[3] Department of Infection Biology, London School of Hygiene & Tropical Medicine, London, United Kingdom.
[4] Laboratory of NF-κB Signalling, Institute of Molecular and Cell Biology (IMCB), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore.
[5] Genome Institute of Singapore (GIS), Agency for Science, Technology and Research (A*STAR), Singapore.
[6] Department of Paediatrics, Yong Loo Lin School of Medicine, National University of Singapore, Singapore.
[7] The Bateson Centre, School of Medicine and Population Health, Faculty of Health, University of Sheffield, Sheffield, United Kingdom.
[8] The Bateson Centre, MRC-Arthritis Research UK Centre for Integrated Research Into Musculoskeletal Ageing (CIMA) and Healthy Lifespan Institute (HELSI), School of Medicine and Population Health, University of Sheffield, Sheffield, UK.
* Corresponding author contact: stefan_oehlers@idlabs.a-star.edu.sg

Abstract

Age is an important risk factor for infections such as tuberculosis (TB). Telomerase is expressed in immune cells yet leukocyte telomere length declines during ageing suggesting an age-dependent loss of telomerase activity in the immune system. Leukocyte telomere length has been correlated with worse outcomes in TB patients, however the mechanisms linking telomere biology to TB susceptibility and response to therapy are unexplored. Here we use the zebrafish-*Mycobacterium marinum* model to investigate the role of telomerase in TB resistance. We find depletion and inhibition of Tert, the catalytic subunit of telomerase, increases bacterial burden in zebrafish embryos while small molecule activation of Tert decreases bacterial burden. Depletion of shelterin components did not recapitulate the infection susceptibility phenotype, and infection susceptibility could not be rescued by p53 or STING depletion. Consistent with a previously described role for Tert in developmental hematopoiesis, we find Tert is necessary for demand-driven emergency myelopoiesis to support containment of extended mycobacterial infection. Our findings establish a previously undescribed role for host telomerase in supporting infection demand-driven hematopoiesis to control infection.

Introduction

Ageing is one of the most important risk factors for tuberculosis (TB) susceptibility and severity (*1*). Animal models and clinical measurements have identified inflammageing, a persistent low grade interferonopathy triggered by accumulated DNA damage, and immune anergy, caused qualitatively by declining lymphocyte function and quantitatively by the biased production of myeloid cells, as potential mechanisms underlying age-dependent TB susceptibility (*2, 3*). Recent work across guinea pig and human systems have demonstrated that TB accelerates ageing suggesting that *Mycobacterium*

53 *tuberculosis* may actively benefit from subverting host ageing pathways and that
54 interception of this process could yield novel targeted adjunctive therapies for an important
55 at-risk group (*4*).
56
Leukocyte telomere length (LTL) decreases during ageing and influences immune
function, specifically affecting susceptibility to infectious diseases driven by host
immunopathology including TB (*5-12*). Some aspects of the relationship between LTL and
TB biology may be ethnicity specific as while South African patients with longer LTL are
more likely to have a successful response to TB treatment, South East Asian and East
Asian patients with longer LTL experienced more treatment side effects (*13-15*). As
telomerase activity is required for maintenance of LTL, we hypothesized decreased
leukocyte expression of the telomerase reverse transcriptase (Tert) may drive the
increased susceptibility of the elderly and LTL individuals to mycobacterial infection.
66
Expression of *tert* by hematopoietic stem cells is necessary for telomere elongation and
likely protects chromosomes during hematopoiesis. There is an increasing body of
evidence that Tert and the RNA template *TERC* execute important non-canonical functions
in immunity as an inflammation-responsive gene aiding the expression of key
hematopoietic differentiation genes (*16-18*). Telomerase-expressing macrophages (T-
MACS) have been previously described and loss of telomerase expression has been
associated with reduced immune function (*19*).
74
75 Here we utilize the zebrafish-*M. marinum* model of human TB to study the role of Tert in
76 mycobacterial infection. We find that Tert is expressed at the site of infection and loss of
77 Tert is causes reduced control of infection. Tert-dependent infection demand-driven
78 hematopoiesis can be supplemented by *tert* mRNA providing evidence for a role for Tert in
79 the resilience of the hematopoietic system during chronic infections.
80
Results
Telomerase reverse transcriptase is expressed at the host-mycobacterial interface
To determine if T-MACs are present in mycobacterial granulomas we analyzed a
previously published scRNAseq dataset of dissected zebrafish-*M. marinum* granulomas
(*20*). We found *tert* is most expressed in proliferative cells, as expected from canonical
function in telomere elongation of proliferating cells and matching the expression of the
DNA polymerase subunit *pola2* (Figure 1A). Expression of *tert* was detected in myeloid-
derived macrophage subsets, many of which lacked *pola2* expression suggesting a non-
proliferative role of *tert* expression. There appeared to be comparable levels of *tert*
expression across epithelialized macrophages (Epmacs, Cluster 3, and cdh1+ve) which
make up the ring-like structure of mature granulomas compared to the "non-structural"
myeloid cells (monocytes, inflammatory macrophages, DCs). The overall level of *tert*
expression in the myeloid cell clusters was less than half of the level of *tert* expression
proliferating cell clusters.
95
We analyzed the expression of *tert* in adult zebrafish-*M. marinum* granulomas by *in situ*
hybridization and found expression in both the flat inner layer of primarily macrophage-
derived epithelialized cells and in round cells around the periphery of the granuloma,
presumably the proliferating cells identified in the scRNAseq dataset or monocyte-derived
cells (Figure 1B). As expected from the heterogenous nature of previously described T-
MACs, there were fewer *tert* positive cells detected within granulomas than were seen in
previously published images of macrophages visualized with the pan monocyte marker
mpeg1 illustrating expression only in a subset of macrophages (*21*).
104

Figure 1: A subset of zebrafish macrophages express *tert* in mycobacterial granulomas.
(A) Mean expression value of *tert* and *pola2* across clusters of granuloma cells identified
by scRNAseq of adult zebrafish-*M. marinum* granulomas from (*20*).
(B) Detection of *tert* expression (blue staining) in (i) necrotic and (ii) non-necrotic adult
zebrafish-*M. marinum* granulomas by *in situ* hybridization. Yellow dashed line indicates
edge of necrotic core and red arrows indicate epithelioid macrophage layer.

Telomerase deficiency increases the severity of pathogenic mycobacterial infection
We next examined the role of *tert* in the immune response to mycobacterial infection by
knocking down expression with a pooled gRNA-CRISPR approach (Supplementary Figure
1A). Knockdown of *tert* increased *M. marinum* burden in zebrafish larvae at 5 days post
infection (dpi) (Figure 2A).

In contrast, infection with two relatively non-pathogenic mycobacteria, *Mycobacterium
smegmatis* and smooth colony variant *Mycobacterium abscessus*, was comparable
between scramble control and *tert* crispant embryos (Figure 2B and 2C). Infection with the
more virulent rough colony variant *M. abscessus* resulted in higher bacterial burden in *tert*
crispant embryos (Figure 2D). These data suggest the mycobacterial infection
susceptibility is specific to infections capable of forming persistent inflammatory lesions.

We next tested the ability of *tert* crispants to control Chikungunya virus (CHIKV) infection,
a well characterized wide host range RNA virus capable of replicating in zebrafish embryos
(*22*). Quantification of infected cell area by fluorometry revealed comparable CHIKV

129 infection levels between scramble control and *tert* crispant embryos, demonstrating a
130 specific immune defect to infection by pathogenic bacteria (Figure 2E).
131

132
133
Figure 2: Knockdown of *tert* increases zebrafish susceptibility to pathogenic mycobacterial
infection
(A) i. Quantification of *M. marinum* burden in 5 dpi *tert* crispant embryos. ii. Representative
images of fluorescent *M. marinum* signal in 5 dpi *tert* crispant embryos. Dashed lines
indicate outline of embryos.
(B) Quantification of *M. smegmatis* burden in 5 dpi *tert* crispant embryos.
(C) Quantification of smooth *M. abscessus* burden in 5 dpi *tert* crispant embryos.
(D) Quantification of rough *M. abscessus* burden in 5 dpi *tert* crispant embryos
(E) Quantification of CHIKV burden in 1-3 dpi *tert* crispant embryos (n= 29 scramble, n= 35
tert crispants).
144
Inhibition of telomerase reverse transcriptase phenocopies the knockdown
mycobacterial infection susceptibility phenotype and activation reverses
susceptibility
We complemented our CRISPR-Cas9 knockdown approach with small molecule TERT
inhibitors BIBR1532 and MST-312 (*23, 24*). Treatment of infected embryos increased *M.*
marinum burden consistent with our genetic approach data (Figure 3A and 3B).
151
152 We next targeted the RNA biogenesis factor *naf1* by CRISPR-Cas9 mutagenesis as it is
153 necessary for TERT function in other models (*25*). Knockdown of *naf1* by CRISPR-Cas9
154 mutagenesis also increased the susceptibility of zebrafish embryos to *M. marinum*
155 infection (Figure 3C, Supplementary Figure 1B).
156
Conversely, to determine if telomerase activation could protect against mycobacterial
infection, we treated infected zebrafish embryos with cycloastragenol, a small molecule
with potential telomerase activating properties in zebrafish (*26*). Cycloastragenol treatment
of infected embryos reduced *M. marinum* burden (Figure 3D).

Genetic supplementation of endogenous *tert* by microinjection with mRNA encoding
zebrafish *tert* also reduced *M. marinum* burden (Figure 3E and 3F). Interestingly,
supplementation of catalytically dead "VAA" *tert* mRNA reduced *M. marinum* burden,
suggesting a non-canonical function of Tert could be protective against infection by
pathogenic mycobacteria.

168
Figure 3: Small molecule and genetic inhibition of Tert increases susceptibility to
mycobacterial infection while activation increases resistance to mycobacterial infection.
(A) Quantification of *M. marinum* burden in 5 dpi embryos treated with BIBR1532.

(B) Quantification of *M. marinum* burden in 5 dpi embryos treated with MST-312.
(C) Quantification of *M. marinum* burden in 5 dpi *naf1* crispant embryos.
(D) i. Quantification of *M. marinum* burden in 5 dpi embryos treated with cycloastragenol
(CAG). ii. Representative images of fluorescent *M. marinum* signal in 5 dpi embryos
treated with the indicated concentrations of cycloastragenol.
(E) i. Quantification of *M. marinum* burden in 5 dpi embryos injected with control *gfp*, *tert*,
or VAA *tert* mRNA. ii. Quantification of *tert* transcript abundance in 1 dpf embryos injected
with control *gfp*, *tert*, or VAA *tert* mRNA.
180
Telomere protection is not associated with resistance against mycobacterial
infection
Chronic infections reduce LTL and DNA damage-induced interferonopathy plays a critical
role in TB pathogenesis (*27, 28*), however previous reports have demonstrated Tert-
dependent telomere shortening is only observed in adult zebrafish and not at the embryo
stage (*16, 29, 30*). We next investigated the hypothesis that canonical telomere-protective
functions of Tert were necessary for resistance to *M. marinum* infection in the context of an
increased genotoxic microenvironment caused by infection. Consistent with the presence
of an inflammatory DNA damage response we observed SPiDER-βGal staining around *M.*
marinum granulomas demonstrating the presence of the senescent cell marker beta
galactosidase activity (Figure 4A).
192
Expression of the DNA damage response gene *tp53* was increased in infected *tert* crispant
embryos compared to uninfected *tert* crispants indicating a small global increase in DNA
damage in the infection-susceptible *tert* crispants (Figure 4B). However, functional
examination of the DNA damage response in *tp53* mutants and *sting* crispants failed to
rescue the *tert* knockdown-induced susceptibility to *M. marinum* infection (Figure 4C and
4D, Supplementary Figure 1C).
199
Knockout of zebrafish shelterin genes induces premature ageing in adult animals and
provides an alternative experimental approach to examine the canonical role of Tert in
telomere protection (*31*). Knockdown of the shelterin genes *acd*, *pot1*, and *trf1* did not
affect control of *M. marinum* infection in zebrafish embryos (Figure 4E and Supplementary
Figure 1D). Together these data suggest the increased susceptibility to infection seen in
Tert-deficient embryos could be due to non-canonical Tert functions.
206

207
208
Figure 4: Increased susceptibility to infection in *tert* crispants is not driven by DNA damage
response pathways.
(A) Fluorescent SPiDER-βGal staining (green) around *M. marinum* granuloma (magenta)
in a 5 dpi embryo.
(B) Quantification of *tp53* transcript abundance in 5 dpi *tert* crispant embryos infected with
M. marinum.
(C) Quantification of *M. marinum* burden in 5 dpi *tp53* mutant embryos.
(D) Quantification of *M. marinum* burden in 5 dpi *sting* crispant embryos.
(E) Quantification of *M. marinum* burden in 5 dpi i. *acd*, ii. *pot1*, and iii. *trf1* crispant
embryos.
219

Tert supports infection-induced myelopoiesis

Zebrafish *tert* and *terc* are known to be necessary for developmental myelopoiesis (*16, 17, 32, 33*), and *M. marinum* infection control requires replenishment of the myeloid cell compartment (*34*). We next analyzed infection-induced myelopoiesis as a potential mechanism of susceptibility in *tert* crispants using *Tg(lyzc:GFP)^{nz117}* and *Tg(lyzc:dsred)^{nz50}* lines, where neutrophils are fluorescently labelled (*35*), and the *TgBAC(mpeg1.1:EGFP)^{vcc7}* line where macrophages are fluorescently labelled (*36*). Caudal hematopoietic tissue-resident neutrophil and macrophages were reduced in *tert* crispants late in infection (Figure 5A and 5B).

To establish if the loss of *lyzc* and *mpeg1.1* positive cells was due to reduced haematopoietic stem cell (HSC) numbers or a lack of differentiation of myeloid cells, we enumerated *Tg(Mmu.Runx1:NLS-mCherry)^{cz2010}* positive HSCs (Figure 5C) (*37*). Tert crispants exhibited a decline in HSC counts at 5 dpi, corresponding with the observed loss of myeloid cells.

Genetic supplementation of endogenous *tert* by microinjection with mRNA encoding full length and catalytically dead zebrafish *tert* increased neutrophil numbers late in infection (Figure 5D). This increase in leukocyte number corresponded with protection from infection (Figure 3E) placing Tert in the control pathogenic mycobacterial infection via supporting infection-induced myelopoiesis.

242
243
Figure 5: Expression of *tert* correlates with infection demand-driven leukocyte abundance.
(A) i. Quantification of CHT-resident neutrophils in 5 dpi *Tg(lyzC:dsred) tert* crispant
embryos infected with *M. marinum*. ii. Representative images of fluorescent neutrophils.
Bracket demarcates CHT region used for counting.

(B) i. Quantification of CHT-resident macrophages in 5 dpi *TgBAC(mpeg1.1:gfp) tert*
crispant embryos infected with *M. marinum*. ii. Representative images of fluorescent
macrophages in the CHT region.
(C) i. Quantification of CHT-resident HSCs in 5 dpi *Tg(Mmu.Runx1:NLS-mCherry)*
embryos infected with *M. marinum*. ii. Representative images of fluorescent HSCs in the
CHT region.
(D) i. Quantification of CHT-resident neutrophils in 5 dpi *Tg(lyzC:dsred)* embryos injected
with control eGFP, *tert*, or VAA *tert* mRNA and infected with *M. marinum*. ii. Representative
images of fluorescent neutrophils.
257
Discussion
Our study used depletion of zebrafish telomerase reverse transcriptase to find a novel role
for telomerase in the control of mycobacterial infection via supporting infection demand-
driven myelopoiesis. These data provide a mechanism that may contribute to the
increasing susceptibility to TB during ageing and explain the more general association of
short LTL with infection susceptibility. The data further suggest supplementation of
telomerase or myelopoietic factors could be used to bolster anti-TB immunity in the elderly
(*34*).
266
While disruption of telomerase activity by small molecule inhibition and *naf1* depletion
suggest a canonical telomere-shortening mechanism of infection susceptibility, our use of
the VAA active site mutant *tert* to curtail *M. marinum* infection and the lack of effect of
shelterin mutation on *M. marinum* infection outcome point towards a non-canonical
function of Tert in infection demand-driven hematopoiesis. We have previously reported a
non-canonical role of zebrafish Tert in supporting gut macrophage proliferation in adult
animals (*19*), an organ with constant turnover and immune stimulation, and consistent with
the defect in infection demand-driven hematopoiesis found in this study. Non-canonical
telomerase, non-coding RNA component (TERC), and shelterin functions span a wide
range of cellular activities, while there are similarities in the DNA-based themes of these
functions there is also clear mechanistic independence of their non-canonical activities
(*38, 39*).
279
Disseminated *M. tuberculosis* drives epigenetic reprogramming of HSCs to impair the
myelopoietic response of infection (*40*). Our zebrafish-*M. marinum* infection model data
adds to this concept by demonstrating loss of HSCs sensitized by the loss of host
telomerase, suggesting TB infection may accelerate the onset of clonal hematopoiesis in
the elderly. Our finding that the premature aging model of telomerase deficiency is a
susceptibility factor for pathogenic mycobacterial infection ties together with prior human
TB and guinea pig-*M. tuberculosis* infection model data finding *M. tuberculosis* infection
prematurely ages the host (*4*). Together, these data suggest TB actively ages the host as a
virulence mechanism to trigger premature decline of hematopoietic function.
289
A commonality of hematopoietic diseases is an association with DNA-associated genes
demonstrating the importance of safeguarding HSC DNA for maintaining the cellular
components of the immune system (*41*). The loss of HSCs in our chronic mycobacterial
infection model suggests TERT may safeguard HSCs from infection-induced damage in
addition to performing previously described roles as an inflammatory transcription factor
(*42*). Additionally, we must include the possibility that loss of canonical telomerase activity
renders HSCs susceptible to premature replicative senescence in context of chronic
infection, which provides an alternative explanation for the infection susceptibility
phenotype seen in small molecule telomerase inhibitor-treated and *naf1* depleted embryos
(Supplementary Figure 2). Further phenotyping of the HSC compartment in the context of

infection is required to determine the range of roles played by telomerase in protecting
HSCs.
302
Multiple studies have linked short telomere length as a proxy for reduced telomerase
expression to viral infection susceptibility in species as diverse as humans and rock
oysters (*5, 10-12, 43*), however little is known about the mechanistic basis for this
association. The only prior mechanistic data connecting telomerase to infection
susceptibility is the cell-intrinsic hijacking of host telomerase to support viral replication
during lytic human cytomegalovirus infection (*44*). Our bacterial infection experiments add
a new dimension to the literature documenting interactions between telomerase and
infection across host and pathogen species. We hypothesize the relative contributions of
HSC and T-MAC telomerase expression will manifest in a disease-specific manner and
encourage more investigation of telomerase function in diverse disease settings.
313
Methods
<u>Animal husbandry</u>
Zebrafish experiments were approved under the A*STAR IACUC protocols 211667 and
221694; Shanghai Public Health and Clinical Center, Fudan University
GB/T 35823-2018 and 2022-A002-01; UK Home Office Project License PPL PP5900632.
Breeding adults were housed under 14 hour light / 10 hour dark cycles in a 28°C
recirculating aquarium. Eggs were produced by natural spawning and embryos were
raised in E3 media at 28°C and supplemented with PTU from 24 hours post fertilization.
322
<u>Bacterial culture</u>
M. abscessus, M. marinum, and *M. smegmatis* were cultured in 7H9 or on 7H10
supplemented with OADC and hygromycin to select for pTEC fluorescent protein plasmids
(L. Ramakrishnan, plasmids are available through Addgene
https://www.addgene.org/Lalita_Ramakrishnan/). Single cell suspensions of mycobacteria
were prepared and frozen at -70°C in aliquots for individual experiments as previously
described (*45*).
330
<u>Viral culture</u>
Chikungunya virus (CHIKV) virus was produced from an infectious clone plasmid
containing a duplicated subgenomic promoter driving ZsGreen (*Anthozoa* reef coral
fluorescent protein) expression between the non-structural and structural genes. The viral
genome was flanked by a CMV and a HDV ribozyme to launch transcription. BHK cells
(RRID: CVCL_1914) were transfected with 500 ng of plasmid per well in a 6 well plate
using lipofectamine 2000 (Thermofisher). At 3 days post transfection when CPE reaches
approximately 40%, supernatant was cleared of cell debris and used to expand CHIKV
culture on VeroE6 cells (RRID: CVCL_0574) for 44 hours. Cleared supernatant was
tittered by standard plaque assay on VeroE6 cells. VeroE6 and BHK were grown in
DMEM high glutamine (Gibco) with 10% FBS (Hyclone).
342
<u>Infection</u>
Zebrafish embryos were infected by microinjection with approximately 200 CFU *M.*
marinum, 1000 CFU DESX1 *M. marinum* or *M. abscessus.* Primary infection was
performed by injection into the caudal vein or neural tube above the yolk sac extension at
2 dpf. Infection was assayed by stereomicroscopy of immobilized zebrafish embryos on a
Nikon SMZ25 stereoscope or an inverted Nikon Eclipse Ti2 at 5 dpi unless otherwise
mentioned. Image analysis was performed in ImageJ/FIJI with bacterial fluorescent pixel
count (FPC) carried out as previously described (*46*).
351

352 HSC counts were carried out on a Zeiss Celldiscoverer 7 (CD7) widefield microscope
353 equipped with a 20x/0.95 plan-apochromat objective and 0.5x tube lens to acquire a 3 Z-
354 stacks every 5 µm.
355
In situ hybridization
DIG-labeled RNA antisense probes were synthesized by *in vitro* transcription with DIG-
labeled mix (Roche 11277073910) and the MEGA shortscript T7 Transcription kit (Thermo
Fisher AM1354) from *tert* cDNA cloned into a pBluescript SK vector (primers referred to as
ISH-tert in Table 1). Probe preparation and *in situ* hybridization of paraffin sections were
carried out as described (*21*).
362
CRISPR-Cas9 mutagenesis
Gene-specific oligos for *in vitro* transcription of gRNAs were designed with IDT or selected
from Wu *et al.* (*47*). HiScribe T7 High Yield RNA Synthesis Kit (NEB E2040) was used for
in vitro transcription and Alt-R CRISPR-Cas9 crRNA was purchased commercially (IDT).
Alt-R S.p. Cas9 Nuclease V3 (IDT 1081059) was combined with gRNA/crRNA and injected
into the yolk of 1-2 cell stage zebrafish embryos.
369
Small molecule treatment
BIBR1532 (Selleckchem S1186), MST-312 (Sigma M3949), and cycloastragenol (Sigma
SML1448) were dissolved in DMSO. Small molecules were added to the media of infected
embryos at final concentrations of 600 nM BIBR1532, 2.5 µM MST-312 and 25 or 50 µM
cycolastragenol. Small molecules were refreshed in complete media changes every
second day.
376
In vitro mRNA synthesis
A full length *tert* cDNA (referred to as FL-tert in Table 1) was cloned into pGEMTeasy using
Q5 polymerase (New England Biolabs). This was subcloned into a pSP64 Poly(A) plasmid
(Promega P1241) using NEBuilder HiFi DNA Assembly (NEB E2621). Site-directed
mutagenesis was carried out with the Q5 Site-Directed Mutagenesis Kit (NEB E0554) to
mutate the ATGAT bases to CCGCC resulting in a predicted change in aspartate to alanine
recreating the VAA catalytically dead TERT mutation (*48*). *In vitro* transcription of capped
mRNA was performed with mMESSAGE mMACHINE Sp6 Transcription Kit (Invitrogen
AM1340). 300 pg of mRNA was injected into the yolk of 1-2 cell stage zebrafish embryos.
386
DNA damage assays
SPiDER-βGal staining was carried out with the Cellular Senescence Detection Kit -
SPiDER-βGal (Dojindo SG03-10) as per manufacturer's instructions on fixed zebrafish
embryos equilibrated to McIlvain buffer pH6 and stained for 30 minutes at 37°C.
TUNEL staining was carried out with the In Situ Cell Death Detection Kit, TMR red (Roche
12156792910) as per manufacturer's instruction on fixed zebrafish embryos with
incubation for 1 hour at 37°C.
394
395 Gene expression analysis
396 Quantitative PCR was performed with reverse transcription using the High Capacity cDNA
397 Kit (Applied Biosystems 4368813) and PCR amplification with PowerUp SYBR Green
398 Master Mix (Thermo Fisher A25742).
399
400 Statistics
401 Analyses of quantitative data were carried out with Graphpad Prism using T-tests for
402 pairwise comparisons or ANOVA for multiple comparisons. All data are representative of at

403 least 3 biological replicates unless otherwise stated in the captions. Error bars on graphs
404 represent standard deviation.
405
406

Acknowledgements
This study was funded by the Singapore Ministry of Health's National Medical Research
Council under its individual research grant scheme (OFIRG22jul-0081) to S.H.O.,
European Research Council Consolidator Grant (772853 - ENTRAPMENT) and Wellcome
Discovery Award (226644/Z/22/Z) to S.M., the National Natural Science Foundation of
China (82372263 to B.Y; 82402628 to G.L).
The authors thank A*STAR IMCB Zebrafish Core Facility discussions and the LSHTM
Biological Services Facility for expert animal husbandry; Dr Sudipto Roy for providing the
tp53 mutant zebrafish line; Drs Miguel Ferreira, Oliver Dreesen, Dennis Kappei, and Maya
Jeitany for helpful telomerase discussions; Mostowy lab members for helpful discussions;
Ms Terumi Nakashima, and Drs Caroline Wee, Shifeng Xue, and Zhipeng Tai for technical
assistance.

Author's Contributions
Darryl JY Han: experimentation, designed experiments, wrote draft
David M Costa: experimentation, designed experiments, approved draft
Geyang Luo: experimentation, approved draft
Kathryn Wright: experimentation, approved draft
Manitosh Pandey: experimentation, approved draft
Denise Wee: experimentation, approved draft
Yao Chen: experimentation, approved draft
Semih Can Akincilar: provided tools, designed experiments, approved draft
Vinay Tergaonkar: provided tools, approved draft
Rajkumar Dorajoo: experimentation, approved draft
Guillaume Carissimo: provided tools, approved draft
Amit Singhal: experimentation, approved draft
Philip M Elks: experimentation, approved draft
Pamela S Ellis: provided tools, approved draft
Catarina M Henriques: provided tools, designed experiments, approved draft
Bo Yan: experimentation, approved draft
Serge Mostowy: experimentation, approved draft
Stefan H Oehlers: conceived study, experimentation, wrote draft

References
1.   A. M. Olmo-Fontanez, J. Turner, Tuberculosis in an Aging World. *Pathogens* **11**,
(2022).
2.   I. M. Orme, Aging and immunity to tuberculosis: increased susceptibility of old mice
reflects a decreased capacity to generate mediator T lymphocytes. *J Immunol* **138**,
4414-4418 (1987).
3.   J. B. Torrelles *et al.*, The Impact of Aging on the Lung Alveolar Environment,
Predetermining Susceptibility to Respiratory Infections. *Front Aging* **3**, 818700
(2022).
4.   C. A. Bobak *et al.*, Increased DNA methylation, cellular senescence and premature
epigenetic aging in guinea pigs and humans with tuberculosis. *Aging (Albany NY)*
14, 2174-2193 (2022).
5.   S. Cohen *et al.*, Association between telomere length and experimentally induced
upper respiratory viral infection in healthy adults. *JAMA* **309**, 699-705 (2013).

6.   J. Helby, B. G. Nordestgaard, T. Benfield, S. E. Bojesen, Shorter leukocyte telomere
length is associated with higher risk of infections: a prospective study of 75,309
individuals from the general population. *Haematologica* **102**, 1457-1465 (2017).
7.   G. A. Dos Santos *et al.*, Shorter leukocyte telomere length is associated with
severity of COVID-19 infection. *Biochem Biophys Rep* **27**, 101056 (2021).
8.   S. Liu *et al.*, Peripheral blood leukocyte telomere length is associated with survival
of sepsis patients. *Eur Respir J* **55**, (2020).
9.   L. Freimane *et al.*, Telomere length and mitochondrial DNA copy number in
multidrug-resistant tuberculosis. *Tuberculosis (Edinb)* **131**, 102144 (2021).
10.   G. A. Noppert *et al.*, Pathogen burden and leukocyte telomere length in the United
States. *Immun Ageing* **17**, 36 (2020).
11.   A. E. Aiello *et al.*, The impact of pathogen burden on leukocyte telomere length in
the Multi-Ethnic Study of Atherosclerosis. *Epidemiol Infect* **145**, 3076-3084 (2017).
12.   J. B. Dowd *et al.*, Persistent Herpesvirus Infections and Telomere Attrition Over 3
Years in the Whitehall II Cohort. *J Infect Dis* **216**, 565-572 (2017).
13.   P. Katoto *et al.*, Telomere length and outcome of treatment for pulmonary
tuberculosis in a gold mining community. *Scientific reports* **11**, 4031 (2021).
14.   W. Udomsinprasert *et al.*, Leukocyte telomere length as a diagnostic biomarker for
anti-tuberculosis drug-induced liver injury. *Scientific reports* **10**, 5628 (2020).
15.   X. Chen *et al.*, Relationship between common telomere length-related genetic
variations, telomere length, and risk of antituberculosis drug-induced hepatotoxicity
in Chinese Han population: As assessed for causality using the updated Roussel
Uclaf Causality Assessment Method. *Fundam Clin Pharmacol* **37**, 858-867 (2023).
16.   S. Imamura *et al.*, A non-canonical function of zebrafish telomerase reverse
transcriptase is required for developmental hematopoiesis. *PLoS ONE* **3**, e3364
(2008).
17.   E. Martinez-Balsalobre *et al.*, Telomerase RNA-based aptamers restore defective
myelopoiesis in congenital neutropenic syndromes. *Nat Commun* **14**, 5912 (2023).
18.   M. Anchelin, L. Murcia, F. Alcaraz-Perez, E. M. Garcia-Navarro, M. L. Cayuela,
Behaviour of telomere and telomerase during aging and regeneration in zebrafish.
PLoS ONE **6**, e16955 (2011).
19.   P. S. Ellis *et al.*, A subset of gut leukocytes has telomerase-dependent "hyper-long"
telomeres and require telomerase for function in zebrafish. *Immun Ageing* **19**, 31
(2022).
20.   M. R. Cronan *et al.*, A non-canonical type 2 immune response coordinates
tuberculous granuloma formation and epithelialization. *Cell* **184**, 1757-1774 e1714
(2021).
21.   G. Luo *et al.*, Temporal and cellular analysis of granuloma development in
mycobacterial infected adult zebrafish. *J Leukoc Biol* **115**, 525-535 (2024).
22.   N. Palha *et al.*, Real-time whole-body visualization of Chikungunya Virus infection
and host interferon response in zebrafish. *PLoS Pathog* **9**, e1003619 (2013).
23.   A. Amin *et al.*, Short-Term TERT Inhibition Impairs Cellular Proliferation via a
Telomere Length-Independent Mechanism and Can Be Exploited as a Potential
Anticancer Approach. *Cancers (Basel)* **15**, (2023).
24.   S. Giunco *et al.*, Anti-Proliferative and Pro-Apoptotic Effects of Short-Term Inhibition
of Telomerase In Vivo and in Human Malignant B Cells Xenografted in Zebrafish.
Cancers (Basel) **12**, (2020).
25.   S. E. Stanley *et al.*, Loss-of-function mutations in the RNA biogenesis factor NAF1
predispose to pulmonary fibrosis-emphysema. *Science translational medicine* **8**,
351ra107 (2016).

26. J. Wu *et al.*, Cycloastragenol protects against glucocorticoid-induced osteogenic
differentiation inhibition by activating telomerase. *Phytother Res* **35**, 2034-2044
(2021).
27. P. Ilmonen, A. Kotrschal, D. J. Penn, Telomere attrition due to infection. *PLoS ONE*
3, e2143 (2008).
28. S. Liu *et al.*, Mycobacterium tuberculosis suppresses host DNA repair to boost its
intracellular survival. *Cell Host Microbe* **31**, 1820-1836 e1810 (2023).
29. M. Anchelin *et al.*, Premature aging in telomerase-deficient zebrafish. *Dis Model*
Mech **6**, 1101-1112 (2013).
30. C. M. Henriques, M. C. Carneiro, I. M. Tenente, A. Jacinto, M. G. Ferreira,
Telomerase is required for zebrafish lifespan. *PLoS Genet* **9**, e1003214 (2013).
31. J. Ma, D. Tang, P. Gao, S. Liang, R. Zhang, Knockout of Shelterin subunit genes in
zebrafish results in distinct outcomes. *Biochem Biophys Res Commun* **617**, 22-29
(2022).
32. F. Alcaraz-Perez *et al.*, A non-canonical function of telomerase RNA in the
regulation of developmental myelopoiesis in zebrafish. *Nat Commun* **5**, 3228
(2014).
33. J. Garcia-Castillo *et al.*, Telomerase RNA recruits RNA polymerase II to target gene
promoters to enhance myelopoiesis. *Proc Natl Acad Sci U S A* **118**, (2021).
34. A. J. Pagan *et al.*, Myeloid Growth Factors Promote Resistance to Mycobacterial
Infection by Curtailing Granuloma Necrosis through Macrophage Replenishment.
Cell Host Microbe **18**, 15-26 (2015).
35. C. Hall, M. V. Flores, T. Storm, K. Crosier, P. Crosier, The zebrafish lysozyme C
promoter drives myeloid-specific expression in transgenic fish. *BMC Dev Biol* **7**, 42
(2007).
36. K. Sugimoto, S. P. Hui, D. Z. Sheng, M. Nakayama, K. Kikuchi, Zebrafish FOXP3 is
required for the maintenance of immune tolerance. *Dev Comp Immunol* **73**, 156-162
(2017).
37. O. J. Tamplin *et al.*, Hematopoietic stem cell arrival triggers dynamic remodeling of
the perivascular niche. *Cell* **160**, 241-252 (2015).
38. S. C. Akincilar, C. H. T. Chan, Q. F. Ng, K. Fidan, V. Tergaonkar, Non-canonical
roles of canonical telomere binding proteins in cancers. *Cell Mol Life Sci* **78**, 4235-
4257 (2021).
39. C. Cao *et al.*, Canonical and non-canonical functions of the non-coding RNA
component (TERC) of telomerase complex. *Cell Biosci* **15**, 30 (2025).
40. N. Khan *et al.*, M. tuberculosis Reprograms Hematopoietic Stem Cells to Limit
Myelopoiesis and Impair Trained Immunity. *Cell* **183**, 752-770 e722 (2020).
41. S. Jaiswal, B. L. Ebert, Clonal hematopoiesis in human aging and disease. *Science*
366, (2019).
42. A. Ghosh *et al.*, Telomerase directly regulates NF-kappaB-dependent transcription.
Nat Cell Biol **14**, 1270-1281 (2012).
43. A. Dupoue *et al.*, Reproductive aging weakens offspring survival and constrains the
telomerase response to herpesvirus in Pacific oysters. *Sci Adv* **10**, eadq2311
(2024).
44. C. M. Cavanaugh *et al.*, Effect of host telomerase inhibition on human
cytomegalovirus. *J Virol* **99**, e0157824 (2025).
45. K. Takaki, J. M. Davis, K. Winglee, L. Ramakrishnan, Evaluation of the
pathogenesis and treatment of Mycobacterium marinum infection in zebrafish. *Nat*
Protoc **8**, 1114-1124 (2013).
46. M. A. Matty, S. H. Oehlers, D. M. Tobin, Live Imaging of Host-Pathogen Interactions
in Zebrafish Larvae. *Methods Mol Biol* **1451**, 207-223 (2016).

47.   R. S. Wu *et al.*, A Rapid Method for Directed Gene Knockout for Screening in G0
556   Zebrafish. *Dev Cell* **46**, 112-125 e114 (2018).
557   48.   W. C. Drosopoulos, V. R. Prasad, The active site residue Valine 867 in human
558   telomerase reverse transcriptase influences nucleotide incorporation and fidelity.
559   *Nucleic Acids Res* **35**, 1155-1168 (2007).
560   49.   A. T. McCurley, G. V. Callard, Characterization of housekeeping genes in zebrafish:
561   male-female differences and effects of tissue type, developmental stage and
562   chemical treatment. *BMC Mol Biol* **9**, 102 (2008).
563   50.   B. Novoa *et al.*, Rag1 immunodeficiency-induced early aging and senescence in
564   zebrafish are dependent on chronic inflammation and oxidative stress. *Aging Cell*
565   **18**, e13020 (2019).
566
567

568
569
Supplementary Figure 1: Editing efficiencies of pooled gRNA-Cas9 gene editing.
(A) High resolution melt analysis of *tert* gRNA2 target site, n=35, 100% efficiency, control
clusters: Green & Blue.
(B) Quantification of *naf1* transcripts in 5 dpi embryos infected with *M. marinum*.
(C) High resolution melt analysis of *sting* gRNA1 target site, n=18, 77.77% efficiency,
control clusters: Green & Blue.
(D) i. High resolution melt analysis of *acd* gRNA2 target site, n=32, 84.3% efficiency,
control clusters: Green & Blue.
ii. High resolution melt analysis of *pot1* gRNA2 target site, n=24 100% efficiency,
control cluster: Green.
iii. High resolution melt analysis of *trf1* gRNA2 target site, n= 25 100% efficiency,
control clusters: Green & Pink.

582
583
584
585

586
Supplementary Figure 2: Hypothesis for modulation of infection susceptibility by *tert* loss-
and gain-of-function
(A) In the control wildtype setting, infection stress drives HSC proliferation to produce
myeloid cells that control infection.
(B) In the Tert loss-of-function setting, loss of telomere elongation and/or protection from
infection-associated genotoxic stress, and Tert-dependent hematopoietic transcription
precipitates hematopoietic collapse and subsequent loss of infection control.
(C) In the Tert gain-of-function setting, increased telomere elongation and/or protection
from infection-associated genotoxic stress, and Tert-dependent hematopoietic transcription
increases myelopoiesis which protects against infection.
597

 Table 1: Oligonucleotides

Use	Name	Sequence 5'-3'	Reference/notes
Cloning eGFP	eGFP Fw	acacatataccttcatacttaattgtcaagTTATTTGTATAGTTCATCCATGCCATGTG	This study, upper case letters indicate coding sequence
Cloning eGFP	eGFP Fw	ctctgcaaagcaagagattaataacccatcATGAGTAAAGGAGAAGAACTTTTCACTG	This study, upper case letters indicate coding sequence
Cloning full length zebrafish *tert* transcript for IVT	FL-Tert Fw	TAAGCATGTCTGGACAGTACTCGACAGATG	This study
Cloning full length zebrafish *tert* transcript for IVT	FL-Tert Rv	CAGGTTTTTTTTACACCCGC	This study
Cloning full length zebrafish *tert* transcript for ISH	ISH-Tert Fw	caagagattaataacccatcATGTCTGGACAGTACTCGACAGAT	This study, upper case letters indicate coding sequence
Cloning full length	ISH-Tert Rv	cttcatacttaattgtcaagTTACACCCGCATGCTGAAATCCAG	This study, upper

zebrafish *tert* transcript for ISH			case letters indicate coding sequence
Gene expression *actb*	Bactin 1 qFW	CGAGCAGGAGATGGGAACC	McCurley and Callard (*49*)
Gene expression *actb*	Bactin 1 qRv	CAACGGAAACGCTCATTGC	McCurley and Callard (*49*)
Gene expression *tert*	tert qFw	GCGCTCCTTATATCCAGTCG	This study
Gene expression *tert*	tert qRv	GGCGCATACAATAAGTCCACT	This study
Gene expression *tp53*	tp53 qFw	GATGGTGAAGGACGAAGGAA	Novoa *et al.* (*50*)
Gene expression *tp53*	tp53 qRv	AAATGACCCCTGTGACAAGC	Novoa *et al.* (*50*)
Gene expression *naf1*	naf1 qFw	GAAAGAGCAAAAGAGGCGCC	This study
Gene expression *naf1*	naf1 qRv	GGAAGTTGCTGTGGAAACCG	This study
gRNA *acd*	acd_Target_1	TAATACGACTCACTATAGGTGAAACGTCGCCGAATGGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *acd*	acd_Target_2	TAATACGACTCACTATAGGTTTCTGAACCCCCCCCGCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *acd*	acd_Target_3	TAATACGACTCACTATAGGTCTAGTGCAAACACGACCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *acd*	acd_Target_4	TAATACGACTCACTATAGGGCTAAACTGGGTCGGCTTGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *naf1*	naf1_Target_1	TAATACGACTCACTATAGGTTGGTCTCATCTTGATCCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *naf1*	naf1_Target_2	TAATACGACTCACTATAGGAGTCTGGCACTGGATCGTGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *naf1*	naf1_Target_3	TAATACGACTCACTATAGGTGCTTGATATTGTGCCGAGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)

gRNA *naf1*	naf1_Target_4	TAATACGACTCACTATAGGACCAGGAACCTCCCCCTGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA scaffold	sgRNA Scaffold Primer	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC	Wu *et al.* (*47*)
gRNA Scrambled	scramble_target_1	TAATACGACTCACTATAGGCAGGCAAAGAATCCCTGCCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA Scrambled	scramble_target_2	TAATACGACTCACTATAGGTACAGTGGACCTCGGTGTCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA Scrambled	scramble_target_3	TAATACGACTCACTATAGGCTTCATACAATAGACGATGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA Scrambled	scramble_target_4	TAATACGACTCACTATAGGTCGTTTTGCAGTAGGATCGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *sting*	sting_Target_1	TAATACGACTCACTATAGGGACTGGGTTTTCTTACTCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *sting*	sting_Target_2	TAATACGACTCACTATAGGCATGTTGCTCAATAGCCTGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *sting*	sting_Target_3	TAATACGACTCACTATAGGCAGAAGGATATGTAGACGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *sting*	sting_Target_4	TAATACGACTCACTATAGGAGTTTCTCACCGTTGAGCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *tert*	tert_Target_1	TAATACGACTCACTATAGGAACCGGCTTTCGGCCGTCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *tert*	tert_Target_2	TAATACGACTCACTATAGGACATTGCGAGGGTCGAGGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *tert*	tert_Target_3	TAATACGACTCACTATAGGGCAGCCCCAGAGCTCGTGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *tert*	tert_Target_4	TAATACGACTCACTATAGGTTCAGGCCTGCCATTGGCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *trf1*	trf1_Target_1	TAATACGACTCACTATAGGTTAGAGAAAGGTCTTCGGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *trf1*	trf1_Target_2	TAATACGACTCACTATAGGAACAGACGGTTCTCTGAGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *trf1*	trf1_Target_3	TAATACGACTCACTATAGGAAGAAACCTTGCAGTGGCGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
gRNA *trf1*	trf1_Target_4	TAATACGACTCACTATAGGTTTCTTACTCGAAGTATGGTTTTAGAGCTAGAAATAGC	Wu *et al.* (*47*)
HRM *acd*	HRM_Acd_Fw	TGCGTTTAACGGGTAGTGTTAAT	This study
HRM *acd*	HRM_Acd_Rv	CGTTTACAGTTCAGTGCAAGTGTA	This study
HRM *sting*	HRM_Sting_Fw	CTGTGATGGGAGAAGACGCT	This study
HRM *sting*	HRM_Sting_Rv	TCACTGAACTTGTCTGAGTCCA	This study
HRM *tert*	HRM_Tert_Fw	GTTGGAAACCAGCGGATCAG	This study

HRM *tert*	HRM_Tert_Rv	AGGAAGTTCTTCATGCCCCG	This study
HRM *trf1*	HRM_Trf1_Fw T2	TGTGTTTAGGTTTAGTGGATGAGC	This study
HRM *trf1*	HRM_Trf1_Rv T2	CGTCCATCACTCTGGCTAGG	This study
HRM *pot1*	HRM_Pot1_Fw T2/2a	AAGCCGATAAGCGCAGAGTT	This study
HRM *pot1*	HRM_Pot1_Rv T2/2a	CTATGCATGACAGCCTTGGC	This study
ISH probe *tert*	ISH_Tert_Fw	TGGCGGATTCAGGCCGGTTTTG	This study
ISH probe *tert*	ISH_Tert_Rv	TATTGAACATTGCGAGGGTCGAG	This study
Site directed mutagenesis of zebrafish *tert* transcript	SDM Tert 'AA' Fw	GAGGCTGATTG[CCGCC]TTTTTGCTCATTACTCC	This study, bracketed sequence indicates mutant sequence
Site directed mutagenesis of zebrafish *tert* transcript	SDM Tert 'AA' Rv	ATCAGACACCCTCCTTTAG	This study

599
600