Fine-tuning quantitative agronomic traits by manipulating gene copy number in rice

Chihiro Nomura[1], Hiroyuki Kanzaki[1], Eiko Kanzaki[1], Motoki Shimizu[1], Kaori Oikawa[1], Hiroe Utsushi[1], Kazue Ito[1], Yusaku Sugimura[1], Ryohei Terauchi[1,2], Akira Abe[1*]

[1] Iwate Biotechnology Research Center, Kitakami, Iwate, Japan

[2] Crop Evolution Laboratory, Kyoto University, Muko, Kyoto, Japan

* Corresponding author

E-mail: a-abe@ibrc.or.jp

Abstract

Although recent plant pan-genomes have revealed extensive copy number variations, studies addressing their phenotypic consequences are limited. Here, we manipulated gene copy number for *OsMADS18* through gene editing and evaluated the effects. The presence of one to three *OsMADS18* tandem copies was reflected in stepwise higher transcript levels and concomitant trait values for leaf blade and culm length, demonstrating that manipulating gene copy number can fine-tune important agronomic traits.

Main

Since the advent of high-throughput sequencing, numerous single-nucleotide polymorphisms (SNPs) and short insertion/deletion polymorphisms (indels) have been identified that affect phenotypes in many organisms, from humans[1] to crop plants[2]. Recent

developments in long-read sequencing technologies such as the PacBio and Oxford

Nanopore Technologies platforms have helped reveal copy number variations (CNVs)

and large structural variations in genomes, which were previously difficult to capture

using short-read sequencing methods, such as those involving the Illumina platform[3–5].

However, current *in silico* analyses using genome sequences obtained by long-read

sequences across plants to understand the relationships between CNV and phenotype are

based on materials with diverse genetic backgrounds, making it difficult to demonstrate

a direct dosage effect of CNV[6]. There is therefore a pressing need to study how variation

in gene dosage caused by CNVs results in different transcript levels and ultimately

influences plant phenotypes[7].

Genome editing is a powerful tool to introduce mutations at target genomic

locations[8]. Among gene editing methods, the clustered regularly interspaced short

palindromic repeats (CRISPR)/CRISPR-associated nuclease 9 (Cas9) system is the most

widely used for genetic research[9–11]. CRISPR/Cas9 typically cuts the genome near a target

site bound by a single guide RNA (sgRNA), after which the repair machinery introduces

mutations such as indels. CRISPR/Cas9-mediated gene editing can knock out genes to

explore gene function[12,13]. However, a complete loss-of-function mutant often displays

drastic changes in phenotypes compared to the wild type[14–17] and thus is not suitable for

studying the quantitative phenotypic effects associated with subtle changes in gene

expression. In addition to knocking out gene function, CRISPR/Cas9-mediated genome

editing can also delete larger genomic fragments by using two sgRNAs targeting two sites

flanking the genomic sequence to be deleted[11]. We hypothesized that this approach could

be used to increase or decrease the copy number of target genes, thereby revealing the

quantitative effects of gene CNV on expression levels and phenotypes.

MADS-box family proteins are transcription factors widely distributed across

eukaryotes. Among them, OsMADS18 from rice (*Oryza sativa*) is involved in the

transition from the vegetative to the reproductive stages[18,19] and exerts pleiotropic effects

on germination and tillering[20]. A previous study of a pan-genome from 33 genetically

diverse rice accessions reported CNVs for *OsMADS18* via tandem duplication[5]. Here, we

performed *de novo* genome assemblies for the elite *japonica* rice cultivar 'Hitomebore'

and the *japonica* cultivar 'Sasanishiki' using long reads from Oxford Nanopore

Technologies and compared the results to the rice reference genome IRGSP-1.0 from the

cultivar 'Nipponbare'[21] (Extended Data Fig. 1a,b). We also assessed sequencing coverage

with Illumina-based short reads between Hitomebore and Sasanishiki (Extended Data Fig.

1c and Supplementary Figs. 1 and 2). From these analyses, we confirmed the presence of

a 20-kb tandem duplication at 24.776–24.796 Mb on chromosome 7 in the Hitomebore

genome and its absence from the Sasanishiki and IRGSP-1.0 reference genomes.

According to the rice annotation project database (RAP-DB)[21,22], the only full-length

protein-coding gene contained in the 20-kb fragment was *OsMADS18*, which is

duplicated specifically in Hitomebore (Extended Data Fig. 1a–c and Supplementary Figs.

1–3). We detected only one SNP and three indels across the two tandem repeats of the 20-

kb fragment, with no variation in the two copies of the *OsMADS18* coding region

(Extended Data Fig. 1d). Given that *OsMADS18* is the only full-length protein-coding

gene within the duplicated region and that sequence variation between the two tandem

copies is minimal, we speculated that CNV of *OsMADS18* might offer an attractive model

for the genetic evaluation of gene dosage effects. We thus attempted to locally manipulate

gene copy number for *OsMADS18* using CRISPR/Cas9 and study how changing gene

copy number would affect plant phenotypes.

To this end, we designed an sgRNA targeting a site 12 kb upstream of the ATG of

OsMADS18, within the region tandemly duplicated in Hitomebore, for CRISPR/Cas9-

mediated editing in this cultivar. Since the two copies of *OsMADS18* have the same

sequence, this sgRNA has two target sites in this cultivar (Fig. 1a). From T_0 individuals

regenerated from Hitomebore calli, we checked for individuals with altered *OsMADS18*

copy number by quantitative PCR (qPCR), using two primer sets (Extended Data Fig.

2a): one set amplifying an amplicon outside of the duplicated region and thus having a

constant copy number and another with primers designed to anneal on either side of the

junction of the duplicated region to specifically amplify the junction if a tandem

duplication exists. We calculated the relative amplification ratio of these two primer sets

using an efficiency-corrected Pfaffl method[23]. When the relative amplification ratio of

Hitomebore (with two homozygous copies of the duplication) was set to 1, deviations of

the relative ratio from 1 in 0.5-unit increments or decrements reflected a change in

OsMADS18 copy number (Extended Data Fig. 2a). The qPCR analysis identified nine T_0

plants with a relative ratio of ~0.5, out of 107 regenerated T_0 plants obtained from

different calli or from the same callus with different culture periods, as well as one T_0

plant with a relative ratio of ~1.5, suggesting a loss or gain of one *OsMADS18* copy,

respectively (Extended Data Fig. 2b,c). We allowed these plants to self-pollinate and

repeated the qPCR analysis on their progeny. From the parents with a relative ratio of

~0.5, we obtained progeny with relative amplification ratios of 0 (reflecting absence of

the junction sequence of the duplicated region), ~0.5 or ~1 (Extended Data Fig. 2d); from

the parent with a relative ratio of ~1.5, we obtained progeny with relative amplification

ratios of ~1, ~1.5 or ~2 (Extended Data Fig. 2e). Furthermore, plant lines with relative

amplification ratios of ~0, ~1 or ~2 produced progeny with the same relative amplification

ratios, suggesting that these plants are homozygous for one, two or three copies of

OsMADS18, respectively (Fig. 1b). These results demonstrate that our qPCR-based

approach is a practical tool for efficiently detecting of genomic CNVs.

To confirm these copy numbers, we checked the sequencing coverage depth over the

OsMADS18 genomic region using Illumina short reads from homozygous plants with an

estimated one, two or three tandem copies. The relative sequencing depth within the

OsMADS18 region decreased in plants with one copy and increased in plants with three

copies, relative to those with two copies (Fig. 1c). A *de novo* genome assembly using

Nanopore long reads for the genomes from homozygous individuals with an estimated

one or three *OsMADS18* copies in the Hitomebore background showed that the

OsMADS18 copy number was indeed one and three, respectively, as evidenced by a dot

matrix analysis (Extended Data Fig. 3).

To study the effect of CNV at *OsMADS18* on its transcript levels, we performed

reverse transcription qPCR (RT-qPCR) on individuals with one to three copies of the gene,

normalizing to *UBQ5*[24,25], with the mean relative *OsMADS18* transcript level for one-

copy plants set to 1. The relative *OsMADS18* transcript levels gradually rose from 1, on

average, to 2 or 3 in plants homozygous for one, two or three copies of *OsMADS18*,

respectively, thus neatly correlating with gene copy number (Fig. 1d). These findings

suggest that simple genome editing introduced CNV with stepwise changes in

OsMADS18 expression levels.

We assessed the agronomic traits of lines with CNV (from one to three copies) at

OsMADS18. We observed no major difference in overall plant architecture among the rice

lines (Extended Data Fig. 4a). However, the flag leaf blade was significantly shorter when

the *OsMADS18* copy number was higher (Fig. 1e,f), as was the length of the leaf blade

just below the flag leaf (Extended Data Fig. 4b). By contrast, culm length was

significantly greater with more *OsMADS18* copies (Fig. 1g). To our knowledge, an effect

of *OsMADS18* on leaf blade size and culm length has not been previously reported.

Previous studies have reported that *OsMADS18* expression is upregulated during the

transition from vegetative to reproductive growth and is involved in the number of days

to heading[18,19]. However, in this study, we measured no significant difference in the

number of days to heading among the three sets of lines with CNV at *OsMADS18*

(Extended Data Fig. 4c).

Most agronomic traits are quantitative in nature, and their variations are controlled

by subtle differences in the expression levels of multiple genes. Thus, manipulating gene

expression is important to adjust traits. Recently, promoter editing has been developed to

generate a large allelic series of plants with different expression levels of a target gene by

inserting random mutations in the *cis*-regulatory elements of promoters[26]. However,

because this approach targets *cis*-regulatory elements, it is difficult to predict the direction

and degree of the changes in expression levels in the edited plants; moreover, the

spatiotemporal gene expression pattern may differ from that driven by the intact promoter.

Here, we propose the manipulation of gene copy number as a powerful alternative method

to create lines with a range of copy numbers leading to stepwise changes in the expression

levels of a target gene without randomly perturbing its spatiotemporal expression patterns.

This method should allow the fine-tuning of important traits by optimizing gene copy

number, thus accelerating plant breeding.

Methods

Plant materials and growth conditions

The genomic structure over the target *OsMADS18* region was compared between the rice

(*Oryza sativa*) ssp. *japonica* cultivars Hitomebore and Sasanishiki. Genome editing was

performed using Hitomebore, which carries two tandem copies of *OsMADS18*. Plants

with *OsMADS18* CNVs were selected from 107 regenerated gene-edited independent

lines. The selected individuals were allowed to self-pollinate to establish homozygous

lines with different *OsMADS18* copy numbers. These lines were grown in a plant

incubator when seedlings. After transplanting, they were grown in a temperature-

controlled greenhouse under short-day conditions, with day/night temperatures of

~27°C/~23°C, and *OsMADS18* transcript levels and various phenotypes were evaluated.

Plasmid construction and plant transformation

To generate lines with CNV at *OsMADS18*, one sgRNA targeting the duplicated genomic

region was designed using CRISPRdirect (https://crispr.dbcls.jp/)[27]. To increase the

probability of mutations, a binary vector containing two sgRNA expression cassettes was

constructed based on a previous study[28]. The *OsU6a* and *OsU6b* promoters, amplified

using pYLsgRNA-OsU6a and pYLsgRNA-OsU6b as templates, were each fused to the

target sequence and sgRNA scaffold. The ccdB cassette of the pYLCRISPR/Cas9P$_{ubi}$-H

vector was replaced with the two sgRNA cassettes via the BsaI site using Golden Gate

assembly. This vector was introduced into *Agrobacterium tumefaciens* strain EHA105,

and rice transformation into Hitomebore was performed according to a previous study[29].

The pYLCRISPR/Cas9P$_{ubi}$-H (RRID:Addgene_66187), pYLsgRNA-OsU6a

RRID:Addgene_66194) and pYLsgRNA-OsU6b (RRID:Addgene_66196) vectors were

kindly provided by Yao-Guang Liu. The primers used in this study are listed in

Supplementary Tables 1.

De novo assembly

To reconstruct the Hitomebore and Sasanishiki genome, *de novo* assembly of each

genome was performed using Nanopore long reads and Illumina short reads following a

previously published method[30]. To extract high-molecular-weight genomic DNA from

rice leaf tissue for Nanopore sequencing, leaf blades were carefully ground using a mortar

and pestle while being cooled with liquid nitrogen, and genomic DNA was extracted using

a NucleoBond high-molecular-weight DNA kit (Macherey-Nagel, Düren, Nordrhein-

Westfalen, Germany). Following DNA extraction, low-molecular-weight DNA was

eliminated using a Short Read Eliminator Kit XL (Circulomics, Baltimore, Maryland,

USA). For Hitomebore, genomic library preparation was performed using a ligation

sequencing kit (SQK-LSK-114; Oxford Nanopore Technologies [ONT], Oxford,

Oxfordshire, UK) according to the manufacturer's instructions, and sequencing was

performed using PromethION 2 Solo (ONT). Additionally, long reads for Hitomebore

produced on a PromethION (ONT) platform, as reported by Sugihara et al.[30], were also

used. For Sasanishiki, genomic library preparation was performed using a ligation

sequencing kit (SQK-LSK-109; ONT), and sequencing was performed on a MinION

instrument.

Base-calling of the Nanopore long reads was conducted using Dorado v0.9.0

(https://github.com/nanoporetech/dorado). The first 50-bp of each read and reads with an

average read quality score below 10 were removed using Chopper v0.9.0[31]. The resulting

clean Nanopore long reads were assembled using NECAT v0.0.1[32], with the genome size

set to 380 Mb and using reads longer than 3,000 bp. To improve the accuracy of the

assembly, Racon v1.4.20[33] was employed for error correction using Nanopore reads,

followed by Medaka v1.4.1 (https://github.com/nanoporetech/medaka) for misassembly

correction. Two rounds of consensus correction were then performed using BWA

v0.7.17[34,35] and HyPo v1.0.3[36] with the Illumina short reads. Redundant contigs were

removed using purge-haplotigs v1.1.1[37], resulting in a *de novo* assembly of 376.2 Mb

comprising 59 contigs for Hitomebore and 378.7 Mb consisting of 49 contigs for

Sasanishiki.

De novo assembly was also performed for Hitomebore gene-edited lines with one or

three copies of the *OsMADS18* region. In both cases, high-molecular-weight genomic

DNA was extracted as described above. The library for the one-copy line was prepared

using an SQK-LSK110 kit and sequenced on MinION, while the library for the three-

copy line was prepared using an SQK-LSK114 kit and sequenced on a PromethION 2

Solo. Base-calling was performed using the same method as described above, and *de novo*

assembly was conducted using Flye v2.9.5[38].

Illumina-based DNA sequencing

Total genomic DNA was extracted from rice leaf blades using the CTAB-based method

with a NucleoSpin Plant II kit (Macherey-Nagel). Sequencing libraries were prepared

using a Collibri ES DNA Library Prep Kit for Illumina Systems (Thermo Fisher Scientific,

Waltham, Massachusetts, USA). Libraries were sequenced on a MiSeq (Illumina, San

Diego, California, USA) instrument in 251-bp paired-end mode for Hitomebore and in

301-bp paired-end mode for Sasanishiki and on a NovaSeq X Plus (Illumina) instrument

in 151-bp paired-end mode for lines with CNV (from one to three copies) at *OsMADS18*.

The resulting raw reads from the MiSeq instrument were filtered using Trimmomatic

v0.36[39] with the paired-end option. Adapter sequences were removed with the

ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10 option. Bases with quality scores of <20 at

the beginning and end of reads were removed (LEADING:20, TRAILING:20). A sliding

window of four bases was used to remove reads with average quality scores of <20

(SLIDINGWINDOW:4:20). Reads were removed if the read length after trimming was <

36 (MINLEN:36) and cut to 200 bases if the read length was > 200 bases (CROP:200).

The resulting raw reads from the NovaSeq X Plus instrument were filtered using fastp

v0.22.0[40] with default settings supplemented with the --detect_adapter_for_pe option to

automatically detect and trim adapter sequences and --trim_poly_g option to trim polyG

tails. The processed reads were aligned to the rice reference genome IRGSP-1.0 using

BWA v0.7.18. The obtained SAM files were converted to BAM format, sorted, and

indexed using SAMtools v1.9[41].

Sequencing depth analysis

236 The sequencing depth was calculated using mosdepth v0.3.3[42] with the BAM files

237 generated from high-throughput sequencing and subsequent processing. The sliding

238 window used in the analysis was created with BEDTools v2.31.1[43] using the command

239 'makewindows'. Relative sequencing depth between target and control was calculated

240 using the following equation:

$$D_{\text{t}} = \left(\frac{d_{\text{t}}}{\overline{D_{\text{t}}}}\right) \times \left(\frac{\overline{D_{\text{c}}}}{d_c}\right) \tag{1}$$

242 where D_{t} is the relative sequencing depth in the region of the target genome, d_{t} is the

243 sequencing depth in the region of the target genome, $\overline{D_{\text{t}}}$ is the average sequencing depth

244 of the whole target genome, d_{c} is the sequencing depth in the target region of the control

245 genome and $\overline{D_{\text{c}}}$ is the average sequencing depth of the whole control genome.

Comparison of genome structures

The genomic structure over the target *OsMADS18* region was compared using MUMmer4

v4.0.1[44]. First, the command 'blastn' from BLAST+ v2.13.0[45] was used to identify the

contig containing *OsMADS18* from each of the four *de novo* assemblies. The identified

contigs were then extracted individually using SeqKit v2.4.0[46]. Each contig was

compared to chromosome 7 of the IRGSP-1.0 reference genome using the command

'nucmer' of MUMmer4. The results were filtered with the command 'delta-filter', setting

the minimum alignment identity to 95% and the minimum alignment length to 5,000 bp.

Finally, the filtered results were visualized using the command 'mummerplot'.

To identify SNPs and indels within the duplicated regions of the Hitomebore

genome, the duplicated sequences were separated into two segments corresponding to the

upstream and downstream copies. These sequences and the corresponding region from

the IRGSP-1.0 reference genome were aligned using MAFFT v7[47]. Sequence differences

were then identified based on the alignment results. Visualization of the alignment of the

tandem duplication was performed using the Integrative Genomics Viewer (IGV)

v2.16.2[48], with the IRGSP-1.0 reference genome.

qPCR-based analysis of target genomic copy number

qPCR was conducted using a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific) and Luna Universal qPCR Master Mix (New England Biolabs, Ipswich, Massachusetts, USA). To estimate CNV of *OsMADS18*, the relative amplification ratio was calculated using qPCR data from primer sets designed to span the duplicated junction and a non-duplicated control region. The analysis was based on the Pfaffl approach[23], which incorporates primer-specific amplification efficiencies. The mean C_t value of wild-type Hitomebore plants (homozygous for two copies of *OsMADS18*, $n = 3$) was used as the calibrator. The C_t difference (ΔC_t) for each primer set was calculated using Equations 2 and 3:

$$\Delta C_{t,junction} = C_{t,junction,calibrator} - C_{t,junction,sample} \tag{2}$$

$$\Delta C_{t,control} = C_{t,control,calibrator} - C_{t,control,sample} \tag{3}$$

The relative amplification ratio (R) was then calculated using Equation 4:

$$R = \frac{\left(E_{junction}\right)^{\Delta C_{t,junction}}}{\left(E_{control}\right)^{\Delta C_{t,control}}} \tag{4}$$

where $E_{junction}$ and $E_{control}$ represent the amplification efficiencies of the junction and control primer sets, respectively.

Further details are provided in Extended Data Fig. 2a, and the primer sequences used in the experiments are listed in Supplementary Table 1.

RT-qPCR analysis

Leaf blade samples were bulk collected from the uppermost fully expanded leaf of the main stem, with three individuals per sample, and immediately frozen in liquid nitrogen. After each sample was ground to powder, total RNA was extracted using a NucleoSpin

RNA Plant kit (Macherey-Nagel). Total RNA was then reverse transcribed into first-

strand cDNA using PrimeScript RT Master Mix (Takara Bio, Kusatsu, Shiga, Japan).

qPCR was performed using a QuantStudio 3 Real-Time PCR System with Luna Universal

qPCR Master Mix. The ΔΔCt method was used to quantify the transcript levels, and the

relative transcript levels of OsMADS18 were calculated using $UBQ5$[24,25] as the internal

control. The mean C_t value of biological replicates from plants with one homozygous

copy of the $OsMADS18$ was used as the calibrator. The primers used in the experiment

are listed in Supplementary Table 1.

Phenotyping

After heading, fully expanded leaf blades were collected, scanned and analysed using a

custom-developed automated measurement program in Python to calculate their

morphology, including length. Leaf blade length was compared among individuals with

the same total number of leaves on the main stem. After heading was fully completed, the

length from the ground level to the panicle neck node was measured and defined as culm

length. The number of days from sowing to heading was recorded as the heading date.

All traits were measured on the main stem.

Statistical analysis

Trait comparisons were made between lines with different $OsMADS18$ copy numbers,

and statistical significance was determined using the Tukey-Kramer test in R v.4.4.2

(https://www.r-project.org/). Details of the test are provided in the figure legends.

Data availability

The raw sequencing reads generated in this study have been deposited in the DNA Data

Bank of Japan (DDBJ) under BioProject ID PRJDB 13864 and PRJDB20676. A detailed

list of accession numbers for all samples is provided in Supplementary Table 2. Nanopore-

based *de novo* genome assemblies for four samples are available at Zenodo (DOI:

https://doi.org/10.5281/zenodo.15335260). The IRGSP-1.0 reference genome used in this

study was obtained from RAP-DB (https://rapdb.dna.affrc.go.jp/download/irgsp1.html).

Source data of this study are provided with this paper.

References

1. Koboldt, D. C., Steinberg, K. M., Larson, D. E., Wilson, R. K. & Mardis, E. R. The

    next-generation sequencing revolution and its impact on genomics. *Cell* **155**, 27–38

    (2013).

2. Singh, R., Kumar, K., Bharadwaj, C. & Verma, P. K. Broadening the horizon of crop

    research: a decade of advancements in plant molecular genetics to divulge phenotype

    governing genes. *Planta* **255**, 46 (2022).

3. Alonge, M. *et al.* Major impacts of widespread structural variation on gene

    expression and crop improvement in tomato. *Cell* **182**, 145–161.e23 (2020).

4. Liu, Y. *et al.* Pan-genome of wild and cultivated soybeans. *Cell* **182**, 162–176.e13

    (2020).

5. Qin, P. *et al.* Pan-genome analysis of 33 genetically diverse rice accessions reveals

hidden genomic variations. *Cell* **184**, 3542–3558.e16 (2021).

6. Yuan, Y., Bayer, P. E., Batley, J. & Edwards, D. Current status of structural variation

studies in plants. *Plant Biotechnol. J.* **19**, 2153–2163 (2021).

7. Maino, E. *et al.* Targeted genome editing *in vivo* corrects a Dmd duplication restoring

wild-type dystrophin expression. *EMBO Mol. Med.* **13**, e13228 (2021).

8. Gaj, T., Gersbach, C. A. & Barbas, C. F., 3rd. ZFN, TALEN, and CRISPR/Cas-based

methods for genome engineering. *Trends Biotechnol.* **31**, 397–405 (2013).

9. Jinek, M. *et al.* A programmable dual-RNA–guided DNA endonuclease in adaptive

bacterial immunity. *Science* **337**, 816–821 (2012).

10. Doudna, J. A. & Charpentier, E. The new frontier of genome engineering with

CRISPR-Cas9. *Science* **346**, 1258096 (2014).

11. Belhaj, K., Chaparro-Garcia, A., Kamoun, S., Patron, N. J. & Nekrasov, V. Editing

plant genomes with CRISPR/Cas9. *Curr. Opin. Biotechnol.* **32**, 76–84 (2015).

12. Shalem, O., Sanjana, N. E. & Zhang, F. High-throughput functional genomics using

CRISPR-Cas9. *Nat. Rev. Genet.* **16**, 299–311 (2015).

13. Bao, A. *et al.* The CRISPR/Cas9 system and its applications in crop genome editing.

Crit. Rev. Biotechnol. **39**, 321–336 (2019).

14. Pnueli, L. *et al.* The *SELF-PRUNING* gene of tomato regulates vegetative to

reproductive switching of sympodial meristems and is the ortholog of *CEN* and

TFL1. *Development* **125**, 1979–1989 (1998).

15. Jeon, J. S. *et al. leafy hull sterile1* is a homeotic mutation in a rice MADS box gene

affecting rice flower development. *Plant Cell* **12**, 871–884 (2000).

16. Komatsu, M., Chujo, A., Nagato, Y., Shimamoto, K. & Kyozuka, J. *FRIZZY*

PANICLE is required to prevent the formation of axillary meristems and to establish

floral meristem identity in rice spikelets. *Development* **130**, 3841–3850 (2003).

17. Qi, J. *et al.* Mutation of the rice *Narrow leaf1* gene, which encodes a novel protein,

affects vein patterning and polar auxin transport. *Plant Physiol.* **147**, 1947–1959

(2008).

18. Fornara, F. *et al.* Functional characterization of *OsMADS18*, a member of the

AP1/SQUA subfamily of MADS box genes. *Plant Physiol.* **135**, 2207–2219 (2004).

19. Kobayashi, K. *et al.* Inflorescence meristem identity in rice is specified by

overlapping functions of three *AP1/FUL*-like MADS box genes and *PAP2*, a

SEPALLATA MADS box gene. *Plant Cell* **24**, 1848–1859 (2012).

20. Yin, X. *et al.* OsMADS18, a membrane-bound MADS-box transcription factor,

modulates plant architecture and the abscisic acid response in rice. *J. Exp. Bot.* **70**,

3895–3909 (2019).

21.   Kawahara, Y. *et al.* Improvement of the *Oryza sativa* Nipponbare reference genome

using next generation sequence and optical map data. *Rice* **6**, 4 (2013).

22.   Sakai, H. *et al.* Rice Annotation Project Database (RAP-DB): an integrative and

interactive database for rice genomics. *Plant Cell Physiol.* **54**, e6 (2013).

23.   Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT-

PCR. *Nucleic Acids Res.* **29**, e45 (2001).

24.   Jain, M., Nijhawan, A., Tyagi, A. K. & Khurana, J. P. Validation of housekeeping

genes as internal control for studying gene expression in rice by quantitative real-

time PCR. *Biochem. Biophys. Res. Commun.* **345**, 646–651 (2006).

25.   Jain, N., Vergish, S. & Khurana, J. Validation of house-keeping genes for

normalization of gene expression data during diurnal/circadian studies in rice by RT-

qPCR. *Sci. Rep.* **8**, 3203 (2018).

26.   Rodríguez-Leal, D., Lemmon, Z. H., Man, J., Bartlett, M. E. & Lippman, Z. B.

Engineering quantitative trait variation for crop improvement by genome editing.

Cell **171**, 470–480.e8 (2017).

27.   Naito, Y., Hino, K., Bono, H. & Ui-Tei, K. CRISPRdirect: software for designing

CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics* **31**, 1120–

1123 (2015).

28. Ma, X. *et al.* A robust CRISPR/Cas9 system for convenient, high-efficiency

multiplex genome editing in monocot and dicot plants. *Mol. Plant* **8**, 1274–1284

(2015).

29. Toki, S. *et al.* Early infection of scutellum tissue with *Agrobacterium* allows high-

speed transformation of rice. *Plant J.* **47**, 969–976 (2006).

30. Sugihara, Y. *et al.* Disentangling the complex gene interaction networks between rice

and the blast fungus identifies a new pathogen effector. *PLoS Biol.* **21**, e3001945

(2023).

31. De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C.

NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* **34**,

2666–2669 (2018).

32. Chen, Y. *et al.* Efficient assembly of nanopore reads via highly accurate and intact

error correction. *Nat. Commun.* **12**, 60 (2021).

33. Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome

assembly from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

34. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics* **26**, 589–595 (2010).

35. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. Preprint at https://doi.org/10.48550/arXiv.1303.3997 (2013).

36.   Kundu, R., Casey, J. & Sung, W.-K. HyPo: Super fast & accurate polisher for long

read genome assemblies. Preprint at https://doi.org/10.1101/2019.12.19.882506

(2019).

37.   Roach, M. J., Schmidt, S. A. & Borneman, A. R. Purge Haplotigs: allelic contig

reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 460

(2018).

38.   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone

reads using repeat graphs. *Nat. Biotechnol.* **37**, 540–546 (2019).

39.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

40.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

41.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008

(2021).

42.   Pedersen, B. S. & Quinlan, A. Mosdepth: quick coverage calculation for genomes

and exomes. *Bioinformatics* **34**, 867–868 (2017).

43.   Quinlan, A. R. BEDTools: the Swiss-army tool for genome feature analysis. *Curr.*

Protoc. Bioinformatics **47**, 11.12.1–11.12.34 (2014).

44.   Marçais, G. *et al.* MUMmer4: A fast and versatile genome alignment system. *PLoS*

Comput. Biol. **14**, e1005944 (2018).

45.   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**,

421 (2009).

46.   Shen, W., Sipos, B. & Zhao, L. SeqKit2: A Swiss army knife for sequence and

alignment processing. *Imeta* **3**, e191 (2024).

47.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

48.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

Acknowledgements

We appreciate Yao-Guang Liu for providing the pYLCRISPR/Cas9P$_{ubi}$-H, pYLsgRNA-OsU6a and pYLsgRNA-OsU6b plasmids, and Yumiko Ogasawara and Miyuki Saito for technical support. Part of the analysis was performed using the National Institute of Genetics (NIG) supercomputer at the ROIS National Institute of Genetics. This research was partially supported by the following grants: the research program on development of innovative technology (JPJ007097) from the Project of the Bio-oriented Technology Research Advancement Institution (BRAIN); and JSPS KAKENHI (JP22K20584 to C.N. and JP23K26882 to A.A.).

Author information

Authors and Affiliations

Iwate Biotechnology Research Center, Kitakami, Iwate, Japan

Chihiro Nomura, Hiroyuki Kanzaki, Eiko Kanzaki, Motoki Shimizu, Kaori Oikawa, Hiroe Utsushi, Kazue Ito, Yusaku Sugimura, Ryohei Terauchi & Akira Abe

Crop Evolution Laboratory, Kyoto University, Muko, Kyoto, Japan

Ryohei Terauchi

Contributions

C.N. and A.A. conceived and designed the research. C.N., H.K., E.K., M.S., K.O., H.U., K.I., Y.S., and A.A. performed the experiments. C.N. and A.A. conducted bioinformatics analyses. C.N., R.T., and A.A. wrote and revised the manuscript. All authors read and approved the final version of the manuscript.

Corresponding author

Correspondence to Akira Abe.

Ethics declarations

Competing interests

The authors declare no competing interests.

Figure legends

Fig. 1 | **Phenotypic variation associated with manipulation of *OsMADS18* copy**

number.

a, Diagram of *OsMADS18* CNV. The sgRNA targets two identical sequences within the

tandem-duplicated *OsMADS18* genomic region. CRISPR/Cas9-mediated editing can

result in plants with either one or three copies of *OsMADS18*. Target sites of the sgRNA

are indicated with scissor icons. **b**, Relative amplification ratio of the tandem-duplicated

OsMADS18 region, as determined by qPCR (n = 18 for each CNV). **c**, Relative

sequencing depth of Illumina short reads around *OsMADS18*. A sliding window with a

window size of 1,000 bp and a step size of 500 bp was used. Thin lines represent

individual plants, and the thick line indicates the mean depth for plants with each

OsMADS18 CNV (n = 5). The mean value of plants with the one homozygous copy of

the *OsMADS18* CNV is used as the control. The orange shaded area marks the position

of *OsMADS18*. **d**, Relative *OsMADS18* transcript levels in leaf blades, as determined by

RT-qPCR, normalized to *UBQ5*, with the mean transcript level of one-copy plants set to

1. Leaf blades were bulk collected from three individuals at 58 days after sowing (n = 10;

total of 30 plants per *OsMAD18* CNV). **e**, Representative photograph of flag leaves. **f,g**,

Lengths of the flag leaf (n = 29, 34, and 29 for 1, 2, and 3 CNV, respectively) (**f**) and culm

(n = 30, 35, and 36 for 1, 2, and 3 CNV, respectively) (**g**). "1 CNV," "2 CNV" and "3

CNV" refer to plants with one, two and three homozygous copies of the *OsMADS18* CNV,

respectively, for (**b**)–(**g**). Different lowercase letters indicate significant differences ($P <$

0.05) by Tukey-Kramer test for (**d**), (**f**) and (**g**). All violin plots were generated using

kernel density estimation with a smoothing adjustment factor of 1.5.

Extended data

Extended Data Fig. 1 | **Comparison of the genomic regions flanking the**

OsMADS18 **locus on chromosome 7 in the Hitomebore and Sasanishiki cultivars.**

a,b, Dot matrix plots of the *OsMADS18* genomic region between Hitomebore and IRGSP-

1.0 (**a**) and between Sasanishiki and IRGSP-1.0 (**b**), using one contig from each cultivar.

Grid lines are spaced every 0.03 Mb. **c**, Relative sequencing depth of Hitomebore as target

to Sasanishiki as control using Illumina short reads around *OsMADS18*. A sliding window

with a window size of 5,000 bp and a step size of 500 bp was used. The orange shaded

area marks the position of *OsMADS18* for (**a**)–(**c**). **d**, One SNP and three indels were

detected between the *OsMADS18* tandem duplications in Hitomebore, representing

sequence differences observed in duplicated region 2 when aligned with duplicated region

1.

Extended Data Fig. 2 | **Determination of *OsMADS18* copy number by qPCR.**

a, Diagram of the method used to determine *OsMADS18* copy number by qPCR. When

the relative amplification ratio of Hitomebore (homozygous for two *OsMADS18* copies)

is set to 1, the values for different genotypes are as follows: homozygous for one copy, 0;

heterozygous for one and two copies, 0.5; heterozygous for two and three copies, 1.5;

homozygous for three copies, 2. **b**, Frequency distribution of the relative amplification

ratio for copy number determination in transgenic plants (T_0), as determined by qPCR. **c**,

Summary table showing the numbers and percentages of T_0 plants with estimated copy

number alterations. T_0 plants derived from different calli or from the same callus with

different culture periods were counted as independent lines. The relative amplification

ratio was used to classify heterozygous plants as follows: heterozygous for one copy/two

copies, $0.3 < x < 0.7$; heterozygous for two copies/three copies, $1.3 < x < 1.7$; x represents

the relative amplification ratio. **d**, Frequency distribution of progeny (n = 400) derived

from estimated heterozygous for one copy/two copies (relative amplification ratio is ~0.5).

e, Frequency distribution of progeny (n = 12) derived from estimated heterozygous for

two copies/three copies (relative amplification ratio is ~1.5).

Extended Data Fig. 3 │ Genomic structure of the *OsMADS18* locus in IRGSP-1.0

and CNV-edited lines.

a,b, Dot matrix plots comparing the *OsMADS18* genomic region in the Nipponbare

reference genome (IRGSP-1.0) to that in plants with one homozygous copy (1 CNV) **(a)**

or three homozygous copies (3 CNV) **(b)** of *OsMADS18*, using a contig containing the

OsMADS18 from each edited line. Grid lines are spaced at 0.03-Mb intervals. The orange

shaded region indicates the position of *OsMADS18*.

Extended Data Fig. 4 │ Effects of *OsMADS18* CNV on other agronomic traits.

a, Representative photograph of overall plant architecture with different *OsMADS18* copy

number in the Hitomebore background. **b**, Leaf blade length of the leaf just below the

flag leaf (n = 29, 35, and 29 for 1, 2, and 3 CNV, respectively). **c**, Days to heading (n =

30, 35, and 36 for 1, 2, and 3 CNV, respectively). Different lowercase letters indicate

significant differences ($P < 0.05$) based on Tukey-Kramer test. "1 CNV," "2 CNV" and

"3 CNV" refer to plants with one, two and three homozygous copies of the *OsMADS18*

CNV, respectively. All violin plots were generated using kernel density estimation with a

smoothing adjustment factor of 1.5.

Supplementary information

Supplementary Information

Supplementary Figs. 1–3, and Figs. 1 and 2.

Fig.1

Extended Data Fig.1

Extended Data Fig. 2

a

b

Extended Data Fig. 3

Extended Data Fig. 4