A Neurocognitive Shift in Midlife: Linking Cognitive Flexibility and Functional-Metabolic Adaptation with the SENECA model

Clément Guichet[1],

Sophie Achard[2], Martial Mermillod[1] & Monica Baciu[1,3*]

[1]Univ. Grenoble Alpes, CNRS LPNC UMR 5105, 38000 Grenoble, France

[2]Univ. Grenoble Alpes, CNRS, INRIA, Grenoble INP, LJK, 38000 Grenoble, France

[3]Neurology Department, CMRR, Grenoble Hospital, 38000 Grenoble, France

*Corresponding Author, CA

Monica Baciu MD PhD

LPNC UMR CNRS 5105

Université Grenoble Alpes

1251 Avenue Centrale, 38400 Grenoble Cedex 09 France

Tel: +33 476 748 145

e-mail: monica.baciu@univ-grenoble-alpes.fr

Graphical abstract

Abstract (184 words)

Cognitive flexibility in the human brain engages dynamic interactions between the Default Mode Network (DMN) and the Fronto-Parietal Network (FPN), a functional architecture that is metabolically demanding and thus potentially susceptible to age-related decline. How the aging brain reorganizes its functional architecture to sustain cognitive flexibility under metabolic constraints remains an open question. In this study, we modeled resting-state functional flexibility across the adult lifespan (ages 18–88) using structural balance theory. Our findings align with the predictions of the SENECA model (*Synergistic, Economical, Nonlinear, Emergent, Cognitive Aging*), revealing a midlife neurocognitive transition: (i) from a metabolically costly, flexible DMN-FPN architecture, toward (ii) a more redundant configuration dominated by low-cost, sensory-driven interactions. The medio-parietal DMN and the Cingulo-Opercular Network (CON) are crucial to this transition, contributing to maintain global brain activity near a critical dynamic regime in older adulthood that optimizes for cognitive flexibility in face of declining metabolic resources. These findings advance a theoretical and methodological framework for understanding neurocognitive flexibility in aging and underscore the importance of multimodal fMRI-PET studies in midlife. They also open promising avenues for translational applications in neuropathology.

Keywords: cognitive flexibility, cognitive aging, resting-state fMRI, metabolism, SENECA model

1. Introduction

A core idea in network neuroscience is that resting-state brain activity is organized to support rich functional dynamics at a minimal wiring cost – a phenomenon known as the *wiring economy* (Achard & Bullmore, 2007; Bullmore & Sporns, 2012; Shine & Poldrack, 2018; Takagi, 2021). Such economical dynamics position the brain in a near-critical regime – notably in the Default Mode Network (DMN) (Deco et al., 2011; Deco & Jirsa, 2012) – that enables rapid exploration and switching between different perceptual, motor, and cognitive cross-system configurations (Cocchi et al., 2017; Hancock et al., 2022; O'Byrne & Jerbi, 2022), and therefore facilitates a flexible functional and cognitive repertoire – i.e., neurocognitive flexibility (Q. Chen et al., 2025; Deco et al., 2013; Ionescu, 2017; R. Wang et al., 2021).

An emerging body of research suggests that such neurocognitive flexibility is regulated by high-level cognitive systems that involve self- and goal-oriented processes (Kupis et al., 2021; Nieberlein et al., 2024; Xia et al., 2022, 2024). Self-oriented processes typically engage the Default Mode Network (DMN) (Menon, 2023; Raichle, 2015); whereas goal-oriented processes primarily engage the Fronto-Parietal Network (FPN), known for its role in cognitive control (Camilleri et al., 2018). Therefore, the interplay between the DMN and FPN lends the control resources necessary for switching between externally-directed – when the DMN is deactivated – and internally-directed streams of cognition – when the DMN is activated. To be clear, external cognition is supported by *anticorrelated* DMN-FPN interactions – a competitive dynamic that suppresses goal-irrelevant thoughts (via DMN deactivation) to facilitate goal-directed attentional processes (via FPN activation) (Buckner & DiNicola, 2019; Finc et al., 2020; Raichle, 2015). In comparison, internal cognition, used for semantic memory, creative thinking (Adnan et al., 2019; Chen et al., 2025), or mental time travel (Chen et al., 2022), typically involves *correlated* activity DMN-FPN interactions – a cooperative dynamic that facilitates spontaneous, yet controlled cognition (Spreng et al., 2014).

This flexible switch between cooperative and competitive DMN-FPN dynamics gives rise to complex cognitive functions that draw on substantial neural flexibility (Cocchi et al., 2013; Luo et al., 2024; Paquola et al., 2025; Smallwood et al., 2021). One example of such cognitive function is semantic control – the ability to selectively access and manipulate learned semantic knowledge (Reilly et al., 2023). Many studies suggest that semantic control is particularly vulnerable to disruptions in neurocognitive flexibility as it engages regions adjacent to both the DMN and FPN (Branzi & Lambon Ralph, 2023; Chiou et al., 2018, 2023; Wang et al., 2024). More broadly, this suggests that complex cognitive functions that heavily rely on semantic control processes, such as langage, offer a valuable framework linking cognitive flexibility to real-word behavior (Benítez-Burraco & Ivanova, 2024), recapitulating the information flow between DMN-semantic and FPN-control systems beyond the traditional left-lateralized language network (Aliko et al., 2023; Dudschig, 2022; Forkel & Hagoort, 2024; Hasson et al., 2018; Hsieh et al., 2024; Krieger-Redwood et al., 2025; Roger, Rodrigues De Almeida, et al., 2022; X. Wang et al., 2020).

As the global population ages (United Nations, 2023), understanding the cognitive relevance of disruption in this DMN-FPN architecture becomes increasingly important. This could guide the development of interventions to sustain cognitive flexibility (Politakis et al., 2022; Uddin, 2021), and *in extenso*, prevent word retrieval and generation challenges in the older population (Alhanai et al., 2017; Baciu & Roger, 2024; Martínez-Ferreiro, 2024).

At a cognitive level, the accumulation of life experiences and knowledge about the world expands the semantic store (Cutler et al., 2025; Gollan & Goldrick, 2019). This is often reflected by preserved or even enhanced vocabulary and langage comprehension (Diaz et al., 2016; Harada et al., 2013; Hartshorne & Germine, 2015; Park & Reuter-Lorenz, 2009). In parallel, older adults show difficulties in inhibitory control processes, therefore reducing their ability to inhibit learned semantic associations that not relevant to the goal (Badre & Wagner, 2007;

Benítez-Burraco & Ivanova, 2023; Hoffman & MacPherson, 2022; Loaiza, 2024; S. Martin et al., 2022; Salthouse, 2019; Shafto & Tyler, 2014; Verhaegen & Poncelet, 2013).

At a brain level, influential models such as the DECHA (*Default-Executive Coupling Hypothesis in Aging*; Spreng & Turner, 2019) suggests that this reduced inhibition reduces the ability to inhibit the DMN at rest, thus disrupting competitive DMN-FPN interactions that uphold externally-directed processes crucial to semantic control (Guichet, Banjac, et al., 2024).

In this study, we argue that further insights on neurocognitive aging can be gained by considering metabolic constraints on this architecture (Camandola & Mattson, 2017; Palombit et al., 2022; Pellerin, 2025; Raz & Daugherty, 2018; Volpi et al., 2024; Wan et al., 2024). Metabolism is particularly relevant in the context of neurocognitive aging as the DMN accounts for up to 80% of the brain's energy consumption at rest (Luo et al., 2024; Shulman et al., 2004; Thomas Yeo et al., 2011; Zang et al., 2004) and shows the strongest coupling between neural activity and energy use of any large-scale network (Shokri-Kojori et al., 2019). Task-related DMN deactivation has also been linked to an increased metabolic response, suggesting that competitive DMN-FPN interactions represents a metabolically expensive mechanism for cognitive flexibility (Leonards et al., 2023; Stiernman et al., 2021). Previous fMRI-PET studies also suggest that GABAergic inhibition of the DMN is a high-cost mechanism, particularly regarding the posterior medial (pm) DMN, which is sensitive to the FPN control demands (Godbersen et al., 2023; Hahn et al., 2024). In sum, these studies suggests that competitive DMN-FPN dynamics are vulnerable to age-related decline in metabolism (Deery et al., 2024; Diniz & Crestani, 2023; Stranahan & Mattson, 2012). Therefore, we argue that comprehensive neurocognitive models should be explicit about how the brain reorganizes this DMN-FPN architecture when metabolic resources decline in older adults, and further explain its impact on cognitive flexibility.

In this regard, the SENECA model *(Synergistic, Economical, Nonlinear, Emergent, Cognitive Aging)* was recently introduced based on resting-state fMRI evidence (Guichet, Banjac, et al., 2024). While initially developed to explain age-related differences in lexical production, the model has the potential to encompass the broader domain of cognitive flexibility (Ionescu, 2017; Xia et al., 2024).

The present study

In the context of cognitive flexibility, the SENECA model makes three predictions that we propose to evaluate in this study.

- (**Prediction 1**) In face of declining metabolic resources, older adults maintain a *global wiring economy* to support flexible dynamics across brain systems, reflecting a form of whole-brain functional homeostasis. Thus, we expected no changes in flexibility at the whole-brain level.

- (**Prediction 2**) To maintain such homeostasis, the brain architecture may undergo spatially distributed changes that downregulates metabolically expensive mechanisms. Specifically, we anticipate a prevalence of flexible DMN-FPN dynamics in younger adulthood, particularly in the pmDMN, before the adoption of a brain architecture with lower metabolic demands around midlife. This new architecture may translate to increase flexibility in lower-level networks in older adulthood (Guichet, Banjac, et al., 2024), emphasizing the SMN for sensorimotor integration and the Cingulo-Opercular Network (CON) for cognitive control instead of the FPN (Sestieri et al., 2014; Wallis et al., 2015).

- (**Prediction 3**) The posterior cingulate cortex (PCC) is a hub linking the DMN and FPN (Fan et al., 2019; Goelman et al., 2024), crucial for semantic control (Krieger-Redwood et al., 2016) and highly-dependent on metabolic activity (Basten et al., 2015; Utevsky

et al., 2014). Because its dorsal (dPCC) and ventral (vPCC) components are differently integrated within the DMN (Godbersen et al., 2023), we expect that the dorsal PCC (dPCC), being more connected to FPN control, will become more rigid with age, while the ventral PCC (vPCC), tied to self-oriented representations, may increase its flexibility within the DMN (Leech et al., 2011; Leech & Sharp, 2014).

To operationalize the concept of neural flexibility, we analyzed resting-state functional connectomes from healthy adults aged 18-88 (Cam-CAN et al., 2014) using *structural balance theory* – a promising graph-theoretical approach (Saberi et al., 2024). This approach proposes to examine flexibility as the interaction between cooperative/correlated and competitive/anti-correlated dynamics (Cocchi et al., 2013; Fox et al., 2005; Luppi, Sanz Perl, et al., 2024; Sporns & Betzel, 2016). Their interaction can be classified into two motifs (Figure 1):

- **Rigid motifs**: Rigid configurations with minimal tension between regions (even number of anti-correlated connections).

- **Flexible motifs**: Flexible configurations that support dynamic reconfigurations between regions (odd number of anti-correlated connections).

Figure 1. Neural flexibility framework. Based on the framework described in Saberi et al. (2024). Orange and Blue represents correlated and anti-correlated patterns of functional connectivity respectively.

2. Results

2.1. Prediction 1: Whole-brain level

A Bayesian practical equivalence test indicated that the ROPE (Region of Practical Equivalence) fully covered the 89% HDI of posterior samples [-1e-4; 1e-4], suggesting no change in whole-brain flexibility at rest throughout the adult lifespan (see Method). In comparison, the ratio of flexibility between versus within networks declined beyond midlife ($F = 76.42$, $p < .001$, $edf = 1.93$, signed partial $R^2 = - .19$). Collectively, this suggests that distributed changes in flexibility within and between networks contribute to maintain homeostasis at the whole-brain level (see Figure 2).

Figure 2. Whole-brain analysis

Bayesian equivalence test found no strong evidence of whole-brain changes in flexibility across the lifespan. The blue trajectory plots the mean expectation of posterior predictive distribution.

In comparison, Between vs. Within-network flexibility begins to decline around age 45 and accelerates in midlife. The orange trajectory plots the fitted values of a Generalized Additive Model (GAM). The bar on top represents the window of significant age-related changes and was determined by computing the 1st derivative of the fitted GAM smooth function. The inflection point in red is the age with the highest significant 2nd derivative.

2.2. Prediction 2: Network level

We decomposed whole-brain flexibility (i) within networks, and (ii) between networks (Figure 3). Model statistics reported in Table 1 in Supplementary Materials. Overall, results suggest

that most age-related changes in flexibility occur around midlife, with an inflection point around age 52.

Between networks. Between networks, only DMN-FPN had a substantial decline in flexibility beyond midlife (see upper right panel on Figure 3); whereas flexibility across lower-level networks exhibited important increases, especially considering Auditory-SMN and SMN-Visual couplings (see bottom panels on Figure 3).

Within networks. Age-related changes in within-network flexibility also occurred within lower-level networks with a similar inflection point in midlife, especially within the SMN, Auditory, Language and Visual 2 networks (see upper left panel on Figure 3).

Figure 3. Age-related within and between-network flexibility

The trajectories plots the predicted values of Generalized Additive Models (GAM). Bars on top represent the window of significant age-related changes and were determined by computing the 1st derivative of the fitted GAM smooth function. We only illustrate the top significant trajectories.

2.3. Prediction 2 & 3: Regional flexibility in DMN-FPN & within DMN

DMN-FPN. Figure 4A shows that the decline in DMN-FPN flexibility is primarily driven by posterior cingulate regions, especially bilateral areas 7m and the bilateral parieto-occipital sulci (POS2). Overall, regions with stronger contributions were more posterior ($r_{(76)} = .52$ [.34;.67] 95% CI, $p < .001$), superior ($r_{(76)} = .32$ [.11;.51] 95% CI , $p < .001$), and medial ($r_{(76)} = .32$ [.1;.5] 95% CI , $p < .001$).

Within the DMN. Figure 4B shows heterogeneity of the PCC activity within the DMN, but only in the left hemisphere. The left dPCC showed substantial decline until midlife, indicating more rigidity, at which point the left vPCC showed great flexibility.

Figure 4. Regions contributing to changes in (A) DMN-FPN and (B) within DMN flexibility. Brain illustrations made with ggseg R-package (Mowinckel & Vidal-Piñeiro, 2020).

2.4. Complementary analysis: Regional flexibility at the whole-brain level

We evaluated the contribution of each region to flexible versus rigid motifs using a novel metric, ***Region Flexibility***. In line with our previous results, Figure 5 illustrates that most changes have an inflection point in midlife, with more flexibility in sensorimotor, auditory and visual cortices; and more rigidity in mostly posterior DMN and FPN areas. A thousand spin

tests preserving spatial autocorrelation revealed that these changes were strongly and significantly correlated to changes along the sensory-transmodal (G1) ($r_s = .49$, $p_{spin} < .001$).

Figure 5. Age-related changes in Region Flexibility. (A & B) Results from GAM models. Red indicates an increase in flexibility, and Blue in rigidity. Brain illustrations made with ggseg R-package (Mowinckel & Vidal-Piñeiro, 2020). **(C)** Resting-state network description of panel A & B. The age-related trajectory for each region is available at: https://github.com/LPNC-LANG/SENECA_network_flexibility.

2.4. Robustness analysis

These findings remain valid when thresholding connectivity matrices using null models that preserve spatial and temporal autocorrelation (Shinn et al., 2023). When applying the more stringent correlation screening approach (Lbath et al., 2024), whole-brain and network-level results result also held. However, robustness could not be assessed at the region level using this approach as there was not enough significant connections to conduct reliable analyses (see Supplementary materials).

3. Discussion

To adapt to varying cognitive demands, resting-state dynamics are thought to reside in a critical regime that supports flexible reconfigurations of cognitive, perceptual, and motor systems (Thiebaut de Schotten & Forkel, 2022; Wang et al., 2021). In this study, we investigated how metabolic resources constrain the resting-state dynamics to optimize for neurocognitive flexibility throughout life. Specifically, we tested the predictions of the SENECA model (Guichet, Banjac, et al., 2024) using cross-sectional resting-state fMRI data from 628 healthy adults aged 18-88 and applied a novel graph-theoretical framework (Saberi et al., 2024).

Our findings support the model's predictions: (1) Changes in flexibility occur locally but not globally, suggesting that spatially distributed changes may contribute to preserve whole-brain homeostasis across the lifespan. (2) These local changes delineates two neurocognitive mode: a high-level DMN-FPN architecture that is more prevalent in younger adulthood, and a sensory-driven architecture that is more prevalent in older adulthood. (3) A similar shift was observed within the posterior medial (pm) DMN, reaffirming the unique role of the DMN in regulating cross-systems interactions.

Taken together, our results suggest that healthy neurocognitive aging is a dynamic phenomenon that implies substantial modifications to optimize for cognitive flexibility given available metabolic resources – marking midlife as a cognitive, functional, and "metabolic cliff" (Huo et al., 2020; Mattson, 2025; Shichkova et al., 2025). This challenges the narrative of a spatially-uniform, and linear neurocognitive decline across the lifespan. Below, we discuss our main findings before integrating our resulting within the SENECA model and highlighting perspectives for future works.

Maintenance of a whole-brain equilibrium across the lifespan

Our main finding is that the resting-state remains in a near-critical regime throughout life that sits in-between rigidity and flexibility (see Figure 1) (Cocchi et al., 2013; Luppi, Sanz Perl, et al., 2024). This regime appears to be sustained differently with age, through spatially distributed adjustments around midlife. Specifically, we noted (i) reduced flexibility between higher-level networks, and (ii) heightened flexibility within and between lower-level networks.

Such finding aligns with the idea of "local cost-efficiency trade-offs" for maintaining global functional homeostasis (Ramchandran et al., 2019), likely reflecting how the aging brain adapts its wiring economy mechanisms to meet for the decline in metabolic resources (Deery et al., 2024). Computational modeling further supports this interpretation, indicating that changes in brain energy induces local connectivity changes based on ongoing cognitive processing, while brain functional activity remains stable globally (Ashourvan et al., 2017; Beim Graben et al., 2019; Volpi et al., 2024). In sum, our study adds to compelling evidence that the brain energy supplying functional activity is heterogeneously distributed at rest (Aiello et al., 2015; Kleinridders et al., 2018; Riedl et al., 2014), in line with functional networks engaging distinct metabolic pathways (Shokri-Kojori et al., 2019).

A synergistic, DMN-FPN architecture in younger adults

With age, the DMN and FPN showed a reduced ability to flexible modulate their between-network connectivity, confirming reports of more rigid DMN-FPN connectivity in older adulthood (e.g., Spreng & Turner, 2019). This gain in rigidity could be further interpreted as a loss of synergistic communication between transmodal areas (Luppi et al., 2022; Margulies et al., 2016), as previously noted in older adulthood (Gatica et al., 2021, 2022), and could highlight difficulties sustaining the high metabolic demands of long-range competitive dynamics (Li et al., 2023; Luppi, Sanz Perl, et al., 2024).

Previous studies suggest that transmodal cortices draw from substantial glucose metabolism compared to lower-level networks (see the section *Additional metabolic considerations*; Luppi, Rosas, et al., 2024). Their important metabolic activity can be traced back to their relative independence from underlying structural constraints (for a review, see Fotiadis et al., 2024), facilitating "top-down projections that tend to jump synaptic levels" (Paquola et al., 2019). While this promotes denser interconnectivity across spatially distributed areas, and thus great flexibility, this comes at the cost of greater energy constraints that makes these networks vulnerable to aging (Diniz & Crestani, 2023; Takagi, 2021; Tomasi et al., 2013).

At the region level, complementary results clarified that age-related decline in DMN-FPN flexibility is driven by posterior medial regions in the PCC. This is consistent with our third hypothesis, highlighting the PCC as a key hub for cognitive flexibility in healthy younger adults, perhaps owing to its unique cortical morphology within the brain (Willbrand et al., 2022). Relatedly, age-related decline could reflect difficulties stimulating GABAergic inhibitory processes linked to the DMN glucose metabolism (Bi et al., 2020; Fortel et al., 2023; Xie et al., 2025), in turn affecting competitive DMN-FPN dynamics.

A redundant, sensory-driven architecture in older adults

Compared to the DMN and FPN, lower-level networks (SMN, auditory, visual, language) showed both increased within- and between-network flexibility beyond midlife. Particularly, observed increases in auditory-SMN, auditory-language, language-SMN, and SMN-visual flexibility could reflect the key role of the SMN in modulating the integration of auditory and visual stimuli in older adulthood.

This could represent the establishment of a sensory-driven processing strategy in older adulthood that demands less metabolic resources. Specifically, we propose to interpret our results as evidence for the processing of similar information within distinct sensory systems,

facilitating a multi-sensory representation. In this context, this strategy may stand as a more redundant approach for supporting cognitive flexibility, favoring robustness over precise neural code (Johnston et al., 2020; Johnston & Freedman, 2023).

This interpretation further agrees with a large body of work showing that the use of multiple sensory channels represents a compensatory strategy to overcome uni-sensory deficits during aging (De Dieuleveult et al., 2017; Hirst et al., 2019; Laurienti et al., 2006). Moreover, previous resting-state MEG teamwork (Guichet, Harquel, et al., 2024) also identified a similar sensory-driven mechanism thought to depend on alpha-band oscillatory activity, potentially compensating for inhibitory control deficits (Jensen & Mazaheri, 2010; Klimesch, 2012; Klimesch et al., 2007; Zioga et al., 2024). These considerations are in line with growing evidence supporting the facilitating role of multi-sensory integration, and particularly the importance of the sensorimotor system in neurocognition (Ferré et al., 2020; Goelman et al., 2024; Roger, Banjac, et al., 2022; Zapparoli et al., 2022).

The DMN and CON drive the midlife shift in dynamics

So far, our discussion gathered evidence showing that younger adults rely on a metabolically expensive, DMN-FPN synergistic mode, while older adults rely on a metabolically inexpensive, sensory-driven architecture with more redundancies for supporting cognitive flexibility. Additional results reveal that (i) posterior cingulate cortex (PCC) dynamics within the DMN, and (ii) between-network dynamics between lower-level networks and the cingulo-opercular network (CON) play a key role in this transition.

Regarding the DMN, dPCC connectivity showed more rigidity beyond midlife while the ventral portion was more engaged (see Figure 4). This contributes to challenge the traditional view of a cohesive task-negative DMN network (Chiou et al., 2020; Kernbach et al., 2018; Williams et al., 2022), and instead aligns with studies highlighting the crucial role of the DMN in cognitive

transitions between internal and external contexts (Crittenden et al., 2015; Kurtin et al., 2023; Paquola et al., 2025; Smith et al., 2018; Zhou et al., 2025). Specifically, our results suggest that medio-parietal DMN activity could support a young-like and older-like mode of cognitive flexibility, in line with prior conceptualizations of a "learning-to-prediction" (Brown et al., 2022), "exploration-to-exploitation" shift in midlife (Hills et al., 2015; Spreng & Turner, 2021; Wyatt et al., 2024).

In younger adults, greater flexibility of the dorsal medio-parietal circuitry may modulate the PMN subnetwork of the DMN (Parietal Memory Network), providing support for updating semantic representations with recent experiences (Zheng et al., 2021) – that is geared towards a *learning or explorative mode*. In comparison, in older adults, enhanced flexibility of ventral medio-parietal circuitry may facilitate the integration of self-related processes by modulating the CAN subnetwork of the DMN (Contextual Association Network) (Braga et al., 2019; Braga & Buckner, 2017), likely playing a key role for generating predictions based on a lifetime of learned associations between objects and associated contexts – that is geared towards a *prediction or exploitative mode.* This could help instantiate "situation models" in older adulthood – a form of embodied memory representation less prone to working memory deficits and generally well-preserved with age (Baltes & Lindenberger, 1997; Kintsch & Van Dijk, 1978; Lindenberger & Baltes, 1994; Lindenberger & Mayr, 2014; A. Martin, 2016; Wright, 2016).

Regarding the CON, increased between-network flexibility with the Auditory and SMN networks confirms its crucial role in age-related controlled cognition (Cao & Cannon, 2021; Crittenden et al., 2016; Hausman et al., 2022; Saberi et al., 2024). From a metabolic perspective, the CON may operate as a lower-cost, bottom-up system for attentional regulation, lending processing resources to goal-relevant networks in a more efficient manner than global FPN coordination (Cocuzza et al., 2020). From a cognitive perspective, one potential interpretation

is that it facilitates somatosensory feedback for preparing and evaluating action outcomes a preferential recruitment through its action-oriented subnetwork in older adulthood (D'Andrea et al., 2023; Dosenbach et al., 2024) and establishes a more embodied form of cognition via tight connectivity to the somato-cognitive action network (SCAN) (Gordon et al., 2023).

SENECA: A Neuroenergetic model of healthy cognitive aging

Overall, our study establishes novel links between cognitive flexibility, resting-state functional brain dynamics, and metabolic resource constraints. The proposed SENECA model posits that age-related declines in metabolic resources may drive shifts in neurocognitive strategies that balances the cost of cross-systems interactions between control-, self-, and action-oriented aspects of cognition (Buckner & DiNicola, 2019; Dosenbach et al., 2025; Gordon et al., 2020; Williams et al., 2022). Below, we propose a hypothetical scenario (see also the Graphical Abstract).

When metabolic resources are high in younger adulthood, the DMN can flexibly engage or disengage the dPCC to regulate cross-region coordination with the FPN. Cooperative DMN-FPN dynamics may provide the attentional resources for integrating novel internal experiences while competitive DMN-FPN dynamics may inhibit task-irrelevant interferences when more top-down control is required to discern between complex semantic relationships (Krieger-Redwood et al., 2019) and provide feedback control (Luppi, Sanz Perl, et al., 2024). This may correspond to a **synergistic mode** as a dynamic interaction between the DMN and FPN is needed to support cognitive flexibility.

Beyond midlife, as metabolic resources decline, inhibiting DMN activity becomes more difficult, thus leading to less competitive DMN-FPN dynamics. To support cognitive flexibility, the older adult brain may engage a different strategy that runs with less metabolic demands. This strategy may correspond to a **redundant mode** where similar information is processed in

distinct lower-level networks to facilitate a more robust multi-sensory mechanism. At the cognitive level, this could provide sufficient flexibility for comparing multi-sensory inputs against stored multimodal representations in the vPCC, exploiting memory-guided processes to generate enacted predictions to boost "working memory capacity limits" (Bar, 2007; Bubic, 2010; Chan et al., 2021; Yang et al., 2024).

Additional metabolic considerations

Most age-related changes occurred between lower-level, unimodal or higher-level, transmodal cortices. Such dissociation could be attributed to the higher costs for sustaining dynamic neural activity between sensory and transmodal areas, further highlighting this gradient as a cost-effective principle for cognitive flexibility (Ceballos et al., 2025; Margulies et al., 2016; Sydnor et al., 2021). From a metabolic perspective, this gradient aligns with the brain-wide distribution of oxygen metabolism (Camandola & Mattson, 2017; Vaishnavi et al., 2010), preferably promoting synaptic formation and growth in transmodal cortices (Goyal et al., 2014). Further studies showed that this oxygen mechanism persists across the lifespan (Goyal et al., 2017; Goyal & Raichle, 2013; Marder & Goaillard, 2006) but at significantly lower rates beyond early adulthood (Goyal et al., 2017; Takada et al., 1992). Thus, oxygen metabolism does not appear to explain the observed in neurocognitive mode as it remains the primary source of energy for sustaining brain activity (Dienel, 2019).

Recently, Luppi, Rosas, Noonan, et al. (2024) suggested that, as the metabolic demands in transmodal cortices exceeds the capacity of oxygen delivery by the vascular system, a mechanism known as nonoxidative glycolysis (i.e., that do not require oxygen) supplies the additional energy needed for synergistic cognitive processes. In comparison, lower-level networks show better management of energy supplied by oxygen (Fletcher et al., 2021). Thus, the shift in neurocognitive mode could reflect a lower reliance of such nonoxidative metabolic

activity. Further studies support this interpretation. For example, BOLD-fMRI responses – largely driven by regional cerebral blood flow (rCBF) – shows tight covariation with glucose metabolism during functional activity in comparison to oxygen metabolism (Dienel, 2012; Magistretti & Allaman, 2015; Paulson et al., 2010). Research also revealed that lower rCBF and glucose hypometabolism is a key element of older adults' decline in cognitive flexibility (Deery et al., 2024; Juttukonda et al., 2021; Tarumi & Zhang, 2018), especially in midlife (Huo et al., 2020; Thambisetty et al., 2013). However, further multimodal imaging approaches are needed to validate these claims (Sala et al., 2023; Voigt et al., 2023).

Limitations & Perspectives

The proposed model assumes that age-related declines in metabolic resources induces a different cognitive strategy. However, since metabolic activity cannot be directly manipulated in a provoked independent variable, the causal direction remains uncertain. It is equally plausible that shifts in cognitive activities, for instance based on heuristics – such as an increased reliance on a fast, intuitive (System 1) compared to slow, deliberative (System 2) processing in older adulthood (Kahneman, 2012), or a tendency toward cognitive closure over openness to experience – may themselves shape the neural architecture over time and, as a consequence, influence metabolic flexibility (Mattson, 2025).

Relatedly, numerous studies demonstrate that lifestyle factors – such as education, socioeconomic status, and physical activity – mediate the relationship between openness and enhanced cognitive functioning in older adulthood (Ihle et al., 2016, 2019; Montoliu et al., 2023; Stolz et al., 2023), with downstream effects on cognitive flexibility (Chung et al., 2012) and language performances (Iso-Markku et al., 2024; Khalili et al., 2024; Perez et al., 2025; Rahman et al., 2023). Another study also suggests that exercise and dietary factors could regulate the brain's metabolic efficiency through neurotrophic factor signaling, giving rise to

the concept of a "metabolic reserve" that complements cognitive reserve aspects (Stranahan & Mattson, 2012). In sum, future research is needed to disentangle how metabolic constraints shape cognitive preferences, or in turn, how enduring cognitive lifestyles actively sculpt brain metabolism and functional organization across the lifespan.

Additionally, we caution that the BOLD-fMRI response does not exclusively reflect glucose metabolism (Volpi et al., 2024). Regional CBF and lactate production also correlate with functional activity (Boumezbeur et al., 2010; Magistretti & Allaman, 2015) and independently vary across the lifespan (Ross et al., 2010; Shichkova et al., 2024). Thus, additional metabolic and vascular mechanisms, unaccounted for in our model, could also play a role in our results (Shichkova et al., 2025). Multimodal datasets could offer a more comprehensive assessment of metabolic alterations related to neurocognitive aging while mitigating vascular confounds, for example targeting age-related changes in glutamate and GABA concentrations using functional MR spectroscopy (Stanley & Raz, 2018).

Conclusion

The interplay between cognitive flexibility, functional brain dynamics, and metabolic constraints across the lifespan remains poorly understood. In this study, we investigated the flexibility of resting-state fMRI activity in a population-based sample of healthy adults aged 18-88. Our results reaffirm previous reports of a midlife shift in neurocognitive processing: (i) younger adults tend to rely on a metabolically costly, DMN-FPN strategy to flexibly adapt to internal and external task demands, whereas (ii) older adults increasingly engage a more energy-efficient, sensory-driven circuitry that supports multi-sensory integration strategy and exploits a lifetime of learned semantic associations. This shift may contribute to preserve whole-brain resting-state activity near a critical regime throughout life that optimizes for neurocognitive flexibility despite declining metabolic resources. Notably, the medio-parietal DMN and the CON may play key roles in orchestrating this shift.

Taken together, our study offers a novel theoretical and empirical perspective on healthy neurocognitive aging, opening avenues for extending the proposed SENECA model to neuropathology.

4. Material and Methods

4.1. Participants

Our dataset included 628 healthy adults (age range: 18-88; 310 males, 318 females) from the Cambridge Center for Ageing and Neuroscience Project (http://www.mrc-cbu.cam.ac.uk/: Cam-CAN et al., 2014). Details on participant recruitment are provided in Taylor et al. (2017).

4.2. MR acquisition and preprocessing

MR acquisition. As the data was sourced from Cam-CAN et al. (2014), we refer the reader to the latter reference for MR acquisition details. Data were preprocessed using the *fMRIPrep* software (https://fmriprep.org/en/stable/: Esteban et al., 2019). T1-weighted images underwent skull stripping, tissue segmentation, and spatial normalization. Resting-state fMRI data underwent motion correction, slice timing correction, susceptibility distortion correction, co-registration, and spatial normalization.

Functional connectomes. Denoised timeseries were aggregated using the 360-region HCP-MMP 1.0 atlas (Glasser et al., 2016).We used custom scripts to co-register the atlas to subject-space with ANTs (https://github.com/ANTsX/ANTsPy) and applied a confound removal strategy via Nilearn (https://nilearn.github.io/). Confounds included high-pass filtering, the full 24 motion parameters, and the full 8 white matter and CSF parameters as recommended in Wang et al. (2024). No global signal regression was performed as it introduces artefactual negative connections (Chai et al., 2012; Murphy & Fox, 2017; Saberi et al., 2022). We performed full scrubbing with a framewise displacement threshold set to 0.5 mm and the standardized DVARs to 1.5. In the event that all segments were removed with full scrubbing, the scrubbing parameter was decremented. Mean framewise displacement was also extracted and used as a covariate in subsequent statistical models. Spatial smoothing and standardization was applied to the signal using a 6mm full-width-at-half-maximum (FWHM) kernel. Finally,

functional connectomes were obtained by computing the Pearson correlation between time series.

4.3. Network flexibility modeling

Computations were performed in Python (version 3.9.12).

4.3.1.   Whole-brain level

In line with Saberi et al. (2024), we define the flexibility of signed and fully-weighted and signed graph in Equation 1:

$$(1) \qquad Flexibility\ (F)\ =\ -\frac{1}{\binom{R}{3}}\sum_{i<j<k}\varphi(F_{ijk})$$

$$with\ \varphi(F_{ijk})\ =\ sign(F_{ijk})\cdot\left|F_{ijk}\right|^{1/3}$$

With $F_{ijk}\ =\ W_{ij}W_{ik}W_{jk}$ representing the product of the pairwise Pearson correlations (*W*) in the motif defined by regions *i, j,* and *k*. Each *W* can be positive or negative, thereby representing correlated or anti-correlated function activity. The value of each motif defined by $\varphi(F_{ijk})$ can be positive, indicating a prevalence of rigid motifs, or negative, indicating a prevalence of flexible motifs. The final flexibility metric (F) is obtained by taking the average of all motifs and applying the negative sign in front of the equation. This negative sign enables a more intuitive interpretation whereby higher values index greater flexibility in the system. Thus, the higher the F value, the greater the flexibility. Conversely, the lower the F value, the greater the rigidity.

4.3.2.   Resting-state network level

We further examined the flexibility *within* networks and *between* networks. Within networks, we examined all motifs made up of regions from a single network N (see Equation 2a).

Let N_α represent a single network, and let I_α be the set of indices for regions in network N_α.

$$(2a) \qquad F_\alpha = -\frac{1}{\binom{|I_\alpha|}{3}} \left(\sum_{\substack{i,j,k \in I_\alpha \\ i<j<k}} \varphi(F_{ijk}) \right)$$

Between networks, we examined motifs made up of at least one region from each network N (see Equation 2b). This allowed us to examine the motifs that resume the joint connectivity between pairs of networks.

Let N_α and N_β represent two networks, and let I_α and I_β be the sets of indices for regions in networks N_α and N_β respectively.

$$(2b) \qquad F_{\alpha\beta} = -\frac{1}{T} \left(\sum_{\substack{i \in I_\alpha, j,k \in I_\beta \\ i<j<k}} \varphi(F_{ijk}) + \sum_{\substack{i \in I_\beta, j,k \in I_\alpha \\ i<j<k}} \varphi(F_{ijk}) \right)$$

With $T = \binom{|I_\alpha|}{1} \cdot \binom{|I_\beta|}{2} + \binom{|I_\beta|}{1} \cdot \binom{|I_\alpha|}{2}$ representing the two ways to select motifs spanning both networks: (i) a motif made of one region from N_α and two regions from N_β; and (b) a motif made of one region from N_β and two regions from N_α.

Between-Within network. We quantified the tendency of a network to exhibit more flexibility between versus within networks. We begin by making all values positive to avoid inconsistencies (Equation 3a). Then, we compute the average flexibility of all between and within components (Equation 3b & 3c) and their normalized ratio (Equation 3d).

Let F_{within} be the set of all within-network flexibility values, $F_{between}$ that of all between-network values, and F_{all} their union.

$$(3a) \qquad \widetilde{F_{all}} = F_{all} + |min(F_{all})| + \varepsilon \quad if \quad min(F_{all}) < 0$$

$$(3b) \qquad F_{average\_between} = \frac{1}{|F_{between}|} \sum_{\alpha < \beta} \widetilde{F_{between,N_{\alpha\beta}}}$$

$$(3c) \qquad F_{average\_within} = \frac{1}{|F_{within}|} \sum_{\alpha} \widetilde{F_{within,N_\alpha}}$$

(3d) **Between vs. Within Energy** $= \frac{1}{|N|} \cdot \frac{F_{average\_between}}{F_{average\_within}}$

with |N| the number of unique networks

4.3.3. Novel graph energy metric

We propose one novel metric at the region level called the **Region Flexibility**. It quantifies the contribution of a region to flexible versus rigid motifs in the network (see Equation 4c). We first define the signed contribution of a region *r* as follows:

$$\text{(4a)} \qquad \frac{1}{T_r \cdot I} \cdot \sum_{\substack{r \in \{i,j,k\} \\ i < j < k}} \varphi\big(F_{ijk}\big) \cdot I$$

With $T_r = \binom{R-1}{2}$ representing the number of motifs to which a region *r* contributes. I is an indicator function observing whether the motif is strictly negative or positive. Then, we defined the **Region Flexibility** as:

$$\text{(4b)} \qquad \frac{\frac{1}{T_r^{(-)}}|Flexible_r|}{\frac{1}{T_r^{(+)}}Rigid_r} \quad \text{with} \quad \begin{array}{l} Flexible_{r,}\ T_r^{(-)}\ when\ F_{ijk} < 0 \\ Rigid_r, T_r^{(+)}\ when\ F_{ijk} > 0 \end{array}$$

This simplifies to:

$$\text{(4c)} \qquad \textbf{\textit{Region Energy}} = \frac{|Flexible_r|}{Rigid_r} \cdot \frac{T_r^{(+)}}{T_r^{(-)}}$$

4.4. Statistical analysis

Statistical analyses were carried out in R (version 4.3.2). All models included sex, handedness, total intracranial volume, and mean framewise displacement as covariates.

Hypothesis 1. We computed the whole-brain network flexibility (see Equation 1) for each individual and built a Bayesian linear regression model with age as a predictor with the *brms* package (version 2.22.0; Bürkner, 2017). Using the *bayestestR* package (version 0.13.2; Makowski et al., 2019), we conducted a practical equivalence test (Kruschke, 2018) with a 89%

HDI (Highest Density Interval) to test whether energy consumption remains stable across the lifespan. The ROPE (Region of Practical Equivalence) limits were defined automatically using the *rope_range()* function. We also computing the global between versus within-network flexibility (see Equation 3) and built a generalized additive model (GAM) with age as a smooth predictor (parameters: *k = 3, method = REML*).

Hypothesis 2. We computed the flexibility within and between networks (see Equation 2) for 9 of the 12 networks mapped by Ji et al. (2019). We did not consider the lesser known networks (i.e., ventro-multimodal, posterior-multimodal, and orbito-affective). Then, we followed the same GAM procedure to examine the age effect on each network or pair of network. Significance thresholds were adjusted at the False Discovery Rate (FDR) across models that examined similar neural elements (that is either within-network or between-network).

Region-level hypothesis. We computed the proposed **Region Flexibility** (see Equation 4) for DMN and FPN regions with respect to DMN-FPN motifs. Similarly, we computed the flexibility of posterior-medial (pm) DMN regions considering DMN motifs. pmDMN regions were taken from the Posterior Cingulate division of the HCP-MMP1.0 atlas (Glasser et al., 2016). Then, we followed the same GAM procedure to examine the age effect as described previously.

4.5. Null models

We sought to replicate our analyses using thresholded matrices to minimize the number of functional connections caused by noise. We determined the statistical significance of each correlation coefficient following two approaches. In the first approach, we generated surrogates by randomizing timeseries, thereby preserving spatial and temporal autocorrelation as these measures are very sensitive to age in our dataset (Shinn et al., 2023). In the second approach,

we determined an absolute threshold (± 0.27) through the correlation screening method, with the goal of limiting false-positives due to intra-regional correlation (Lbath et al., 2024).

We added the subject-specific density as a covariate to preclude the effect of different network sizes between individuals (Rubinov & Sporns, 2010; Stumme et al., 2022; van den Heuvel et al., 2017). Indeed, the elimination of non-significant edges may distort graph theoretical measures due to differences in the number of remaining edges, and generally lead to more random network characterization in network in older adults due to overall low FC.

Data and code availability

Code is made publicly available at:
https://github.com/LPNC-LANG/SENECA_network_flexibility

Supplementary information

Supplementary materials

CRediT author statement

Conceptualization: CG; Methodology, Formal Analysis & Data Curation: CG; Writing – Original Draft: CG, MB; Writing – Review & Editing: CG, MB, MM, SA; Funding Acquisition: MB, MM, SA; Supervision: MB, MM.

Declaration of interests

The authors declare no competing interests.

Acknowledgments

This work was supported by the ANR project ANR-15-IDEX-02 and MIAI @Grenoble Alpes (ANR-19-P3IA-0003). This project has received financial support from the CNRS through the

MITI interdisciplinary programs. The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) research was supported by the Biotechnology and Biological Sciences Research Council (grant number BB/H008217/1).

References

Achard, S., & Bullmore, E. (2007). Efficiency and Cost of Economical Brain Functional Networks. *PLOS Computational Biology*, *3*(2), e17. https://doi.org/10.1371/journal.pcbi.0030017

Adnan, A., Beaty, R., Silvia, P., Spreng, R. N., & Turner, G. R. (2019). Creative aging : Functional brain networks associated with divergent thinking in older and younger adults. *Neurobiology of Aging*, *75*, 150-158. https://doi.org/10.1016/j.neurobiolaging.2018.11.004

Aiello, M., Salvatore, E., Cachia, A., Pappatà, S., Cavaliere, C., Prinster, A., Nicolai, E., Salvatore, M., Baron, J.-C., & Quarantelli, M. (2015). Relationship between simultaneously acquired resting-state regional cerebral glucose metabolism and functional MRI : A PET/MR hybrid scanner study. *NeuroImage*, *113*, 111-121. https://doi.org/10.1016/j.neuroimage.2015.03.017

Alhanai, T., Au, R., & Glass, J. (2017). Spoken language biomarkers for detecting cognitive impairment. *2017 IEEE Automatic Speech Recognition and Understanding Workshop (ASRU)*, 409-416. https://doi.org/10.1109/ASRU.2017.8268965

Aliko, S., Wang, B., Small, S. L., & Skipper, J. I. (2023). The entire brain, more or less, is at work *: 'Language regions' are artefacts of averaging* [Preprint]. Neuroscience. https://doi.org/10.1101/2023.09.01.555886

Ashourvan, A., Gu, S., Mattar, M. G., Vettel, J. M., & Bassett, D. S. (2017). The energy landscape underpinning module dynamics in the human brain connectome. *NeuroImage*, *157*, 364-380. https://doi.org/10.1016/j.neuroimage.2017.05.067

Baciu, M., & Roger, E. (2024). Finding the Words : How Does the Aging Brain Process Language? A Focused Review of Brain Connectivity and Compensatory Pathways. *Topics in Cognitive Science*, tops.12736. https://doi.org/10.1111/tops.12736

Badre, D., & Wagner, A. D. (2007). Left ventrolateral prefrontal cortex and the cognitive control of memory. *Neuropsychologia*, *45*(13), 2883-2901. https://doi.org/10.1016/j.neuropsychologia.2007.06.015

Baltes, P. B., & Lindenberger, U. (1997). Emergence of a powerful connection between sensory and cognitive functions across the adult life span : A new window to the study of cognitive aging? *Psychology and Aging*, *12*(1), 12-21. https://doi.org/10.1037/0882-7974.12.1.12

Bar, M. (2007). The proactive brain : Using analogies and associations to generate predictions. *Trends in Cognitive Sciences*, *11*(7), 280-289. https://doi.org/10.1016/j.tics.2007.05.005

Basten, U., Hilger, K., & Fiebach, C. J. (2015). Where smart brains are different : A quantitative meta-analysis of functional and structural brain imaging studies on intelligence. *Intelligence*, *51*, 10-27. https://doi.org/10.1016/j.intell.2015.04.009

Beim Graben, P., Jimenez-Marin, A., Diez, I., Cortes, J. M., Desroches, M., & Rodrigues, S. (2019). Metastable Resting State Brain Dynamics. *Frontiers in Computational Neuroscience*, *13*, 62. https://doi.org/10.3389/fncom.2019.00062

Benítez-Burraco, A., & Ivanova, O. (2023). Revisiting the hypothesis of language retrogenesis from an evolutionary perspective. *Neuropsychology*, *37*(5), 501-518. https://doi.org/10.1037/neu0000886

Benítez-Burraco, A., & Ivanova, O. (2024). Language in healthy and pathological ageing : Methodological milestones and challenges. *International Journal of Language & Communication Disorders*, *59*(1), 4-12. https://doi.org/10.1111/1460-6984.13003

Bi, D., Wen, L., Wu, Z., & Shen, Y. (2020). GABAergic dysfunction in excitatory and inhibitory (E/I) imbalance drives the pathogenesis of Alzheimer's disease. *Alzheimer's & Dementia*, *16*(9), 1312-1329. https://doi.org/10.1002/alz.12088

Boumezbeur, F., Petersen, K. F., Cline, G. W., Mason, G. F., Behar, K. L., Shulman, G. I., & Rothman, D. L. (2010). The Contribution of Blood Lactate to Brain Energy Metabolism in Humans Measured by Dynamic[13] C Nuclear Magnetic Resonance Spectroscopy. *The Journal of Neuroscience*, *30*(42), 13983-13991. https://doi.org/10.1523/JNEUROSCI.2040-10.2010

Braga, R. M., & Buckner, R. L. (2017). Parallel Interdigitated Distributed Networks within the Individual Estimated by Intrinsic Functional Connectivity. *Neuron*, *95*(2), 457-471.e5. https://doi.org/10.1016/j.neuron.2017.06.038

Braga, R. M., Van Dijk, K. R. A., Polimeni, J. R., Eldaief, M. C., & Buckner, R. L. (2019). Parallel distributed networks resolved at high resolution reveal close juxtaposition of distinct regions. *Journal of Neurophysiology*, *121*(4), 1513-1534. https://doi.org/10.1152/jn.00808.2018

Branzi, F. M., & Lambon Ralph, M. A. (2023). Semantic-specific and domain-general mechanisms for integration and update of contextual information. *Human Brain Mapping*, *44*(17), 5547-5566. https://doi.org/10.1002/hbm.26454

Brown, R. M., Gruijters, S. L. K., & Kotz, S. A. (2022). Prediction in the Aging Brain : Merging Cognitive, Neurological, and Evolutionary Perspectives. *The Journals of Gerontology: Series B*, *77*(9), 1580-1591. https://doi.org/10.1093/geronb/gbac062

Bubic. (2010). Prediction, cognition and the brain. *Frontiers in Human Neuroscience*. https://doi.org/10.3389/fnhum.2010.00025

Buckner, R. L., & DiNicola, L. M. (2019). The brain's default network : Updated anatomy, physiology and evolving insights. *Nature Reviews Neuroscience*, *20*(10), 593-608. https://doi.org/10.1038/s41583-019-0212-7

Bullmore, E., & Sporns, O. (2012). The economy of brain network organization. *Nature Reviews Neuroscience*. https://doi.org/10.1038/nrn3214

Bürkner, P.-C. (2017). **brms** : An *R* Package for Bayesian Multilevel Models Using *Stan*. *Journal of Statistical Software*, *80*(1). https://doi.org/10.18637/jss.v080.i01

Camandola, S., & Mattson, M. P. (2017). Brain metabolism in health, aging, and neurodegeneration. *The EMBO Journal*, *36*(11), 1474-1492. https://doi.org/10.15252/embj.201695810

Cam-CAN, Shafto, M. A., Tyler, L. K., Dixon, M., Taylor, J. R., Rowe, J. B., Cusack, R., Calder, A. J., Marslen-Wilson, W. D., Duncan, J., Dalgleish, T., Henson, R. N., Brayne, C., & Matthews, F. E. (2014). The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) study protocol : A cross-sectional, lifespan, multidisciplinary examination of healthy cognitive ageing. *BMC Neurology*, *14*(1), 204. https://doi.org/10.1186/s12883-014-0204-1

Camilleri, J. A., Müller, V. I., Fox, P., Laird, A. R., Hoffstaedter, F., Kalenscher, T., & Eickhoff, S. B. (2018). Definition and characterization of an extended multiple-demand network. *NeuroImage*, *165*, 138-147. https://doi.org/10.1016/j.neuroimage.2017.10.020

Cao, H., & Cannon, T. D. (2021). Distinct and temporally associated neural mechanisms underlying concurrent, postsuccess, and posterror cognitive controls : Evidence from a stop-signal task. *Human Brain Mapping*, *42*(9), 2677-2690. https://doi.org/10.1002/hbm.25347

Ceballos, E. G., Luppi, A. I., Castrillon, G., Saggar, M., Misic, B., & Riedl, V. (2025). The control costs of human brain dynamics. *Network Neuroscience*, 1-23. https://doi.org/10.1162/netn_a_00425

Chai, X. J., Castañón, A. N., Öngür, D., & Whitfield-Gabrieli, S. (2012). Anticorrelations in resting state networks without global signal regression. *NeuroImage*, *59*(2), 1420-1428. https://doi.org/10.1016/j.neuroimage.2011.08.048

Chan, J. S., Wibral, M., Stawowsky, C., Brandl, M., Helbling, S., Naumer, M. J., Kaiser, J., & Wollstadt, P. (2021). Predictive Coding Over the Lifespan : Increased Reliance on Perceptual Priors in Older Adults—A Magnetoencephalography and Dynamic Causal Modeling Study. *Frontiers in Aging Neuroscience*, *13*, 631599. https://doi.org/10.3389/fnagi.2021.631599

Chen, Q., Kenett, Y. N., Cui, Z., Takeuchi, H., Fink, A., Benedek, M., Zeitlen, D. C., Zhuang, K., Lloyd-Cox, J., Kawashima, R., Qiu, J., & Beaty, R. E. (2025). Dynamic switching between brain networks predicts creative ability. *Communications Biology*, *8*(1), 54. https://doi.org/10.1038/s42003-025-07470-9

Chen, R., Singh, M., Braver, T. S., & Ching, S. (2025). Dynamical models reveal anatomically reliable attractor landscapes embedded in resting-state brain networks. *Imaging Neuroscience*, *3*, imag_a_00442. https://doi.org/10.1162/imag_a_00442

Chen, T., Huang, J., Cui, J., Li, Z., Wang, Y., Irish, M., & Chan, R. C. K. (2022). Functional Coupling between the Fronto-Parietal Network and Default Mode Network Is Associated with Balanced Time Perspective. *Brain Sciences*, *12*(9), 1201. https://doi.org/10.3390/brainsci12091201

Chiou, R., Humphreys, G. F., Jung, J., & Lambon Ralph, M. A. (2018). Controlled semantic cognition relies upon dynamic and flexible interactions between the executive 'semantic control' and hub-and-spoke 'semantic representation' systems. *Cortex*, *103*, 100-116. https://doi.org/10.1016/j.cortex.2018.02.018

Chiou, R., Humphreys, G. F., & Lambon Ralph, M. A. (2020). Bipartite Functional Fractionation within the Default Network Supports Disparate Forms of Internally Oriented Cognition. *Cerebral Cortex*, *30*(10), 5484-5501. https://doi.org/10.1093/cercor/bhaa130

Chiou, R., Jefferies, E., Duncan, J., Humphreys, G. F., & Lambon Ralph, M. A. (2023). A middle ground where executive control meets semantics : The neural substrates of semantic control are topographically sandwiched between the multiple-demand and default-mode systems. *Cerebral Cortex*, *33*(8), 4512-4526. https://doi.org/10.1093/cercor/bhac358

Chung, S.-H., Su, Y.-F., & Su, S.-W. (2012). The Impact of Cognitive Flexibility on Resistance to Organizational Change. *Social Behavior and Personality: An International Journal*, *40*(5), 735-745. https://doi.org/10.2224/sbp.2012.40.5.735

Cocchi, L., Gollo, L. L., Zalesky, A., & Breakspear, M. (2017). Criticality in the brain : A synthesis of neurobiology, models and cognition. *Progress in Neurobiology*, *158*, 132-152. https://doi.org/10.1016/j.pneurobio.2017.07.002

Cocchi, L., Zalesky, A., Fornito, A., & Mattingley, J. B. (2013). Dynamic cooperation and competition between brain systems during cognitive control. *Trends in Cognitive Sciences*, *17*(10), 493-501. https://doi.org/10.1016/j.tics.2013.08.006

Cocuzza, C. V., Ito, T., Schultz, D., Bassett, D. S., & Cole, M. W. (2020). Flexible Coordinator and Switcher Hubs for Adaptive Task Control. *The Journal of Neuroscience*, *40*(36), 6949-6968. https://doi.org/10.1523/JNEUROSCI.2559-19.2020

Crittenden, B. M., Mitchell, D. J., & Duncan, J. (2015). Recruitment of the default mode network during a demanding act of executive control. *eLife*, *4*, e06481. https://doi.org/10.7554/eLife.06481

Crittenden, B. M., Mitchell, D. J., & Duncan, J. (2016). Task Encoding across the Multiple Demand Cortex Is Consistent with a Frontoparietal and Cingulo-Opercular Dual Networks Distinction. *The Journal of Neuroscience*, *36*(23), 6147-6155. https://doi.org/10.1523/JNEUROSCI.4590-15.2016

Cutler, R. A., Mirjalili, S., Pham, P., Devulapalli, H., Zafar, S., & Duarte, A. (2025). Semantic Memory Space Becomes Denser with Age. *Neuropsychologia*, 109083. https://doi.org/10.1016/j.neuropsychologia.2025.109083

D'Andrea, C. B., Laumann, T. O., Newbold, D. J., Nelson, S. M., Nielsen, A. N., Chauvin, R., Marek, S., Greene, D. J., Dosenbach, N. U. F., & Gordon, E. M. (2023). *Substructure of the brain's Cingulo-Opercular network*. https://doi.org/10.1101/2023.10.10.561772

De Dieuleveult, A. L., Siemonsma, P. C., Van Erp, J. B. F., & Brouwer, A.-M. (2017). Effects of Aging in Multisensory Integration : A Systematic Review. *Frontiers in Aging Neuroscience*, *9*. https://doi.org/10.3389/fnagi.2017.00080

Deco, G., & Jirsa, V. K. (2012). Ongoing Cortical Activity at Rest : Criticality, Multistability, and Ghost Attractors. *The Journal of Neuroscience*, *32*(10), 3366-3375. https://doi.org/10.1523/JNEUROSCI.2523-11.2012

Deco, G., Jirsa, V. K., & McIntosh, A. R. (2011). Emerging concepts for the dynamical organization of resting-state activity in the brain. *Nature Reviews Neuroscience*, *12*(1), 43-56. https://doi.org/10.1038/nrn2961

Deco, G., Jirsa, V. K., & McIntosh, A. R. (2013). Resting brains never rest : Computational insights into potential cognitive architectures. *Trends in Neurosciences*, *36*(5), 268-274. https://doi.org/10.1016/j.tins.2013.03.001

Deery, H. A., Liang, E., Di Paolo, R., Voigt, K., Murray, G., Siddiqui, M. N., Egan, G. F., Moran, C., & Jamadar, S. D. (2024). The association of regional cerebral blood flow and glucose metabolism in normative ageing and insulin resistance. *Scientific Reports*, *14*(1), 14574. https://doi.org/10.1038/s41598-024-65396-4

Deery, H., Liang, E., Siddiqui, N. M., Murray, G., Voigt, K., Di Paolo, R., Moran, C., Egan, G. F., & Jamadar, S. D. (2024). *Metabolic connectivity in ageing*. https://doi.org/10.1101/2024.06.16.599247

Diaz, M. T., Rizio, A. A., & Zhuang, J. (2016). The Neural Language Systems That Support Healthy Aging : Integrating Function, Structure, and Behavior: Neural Language Systems in Aging. *Language and Linguistics Compass*, *10*(7), 314-334. https://doi.org/10.1111/lnc3.12199

Dienel, G. A. (2012). Fueling and Imaging Brain Activation. *ASN Neuro*, *4*(5), AN20120021. https://doi.org/10.1042/AN20120021

Dienel, G. A. (2019). Brain Glucose Metabolism : Integration of Energetics with Function. *Physiological Reviews*, *99*(1), 949-1045. https://doi.org/10.1152/physrev.00062.2017

Diniz, C. R. A. F., & Crestani, A. P. (2023). The times they are a-changin' : A proposal on how brain flexibility goes beyond the obvious to include the concepts of "upward" and "downward" to neuroplasticity. *Molecular Psychiatry*, *28*(3), 977-992. https://doi.org/10.1038/s41380-022-01931-x

Dosenbach, N. U. F., Raichle, M. E., & Gordon, E. M. (2025). The brain's action-mode network. *Nature Reviews Neuroscience*. https://doi.org/10.1038/s41583-024-00895-x

Dosenbach, N. U. F., Raichle, M., & Gordon, E. M. (2024). *The brain's cingulo-opercular action-mode network*.

Dudschig, C. (2022). Language and non-linguistic cognition : Shared mechanisms and principles reflected in the N400. *Biological Psychology*, *169*, 108282. https://doi.org/10.1016/j.biopsycho.2022.108282

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep : A robust preprocessing pipeline for functional MRI. *Nature Methods*, *16*(1), 111-116. https://doi.org/10.1038/s41592-018-0235-4

Fan, Y., Borchardt, V., Von Düring, F., Leutritz, A. L., Dietz, M., Herrera-Meléndez, A. L., Bajbouj, M., Li, M., Grimm, S., & Walter, M. (2019). Dorsal and Ventral Posterior Cingulate Cortex Switch Network Assignment via Changes in Relative Functional Connectivity Strength to Noncanonical Networks. *Brain Connectivity*, *9*(1), 77-94. https://doi.org/10.1089/brain.2018.0602

Ferré, P., Jarret, J., Brambati, S. M., Bellec, P., & Joanette, Y. (2020). Task-Induced Functional Connectivity of Picture Naming in Healthy Aging : The Impacts of Age and Task Complexity. *Neurobiology of Language*, *1*(2), 161-184. https://doi.org/10.1162/nol_a_00007

Finc, K., Bonna, K., He, X., Lydon-Staley, D. M., Kühn, S., Duch, W., & Bassett, D. S. (2020). Dynamic reconfiguration of functional brain networks during working memory training. *Nature Communications*, *11*(1), 2435. https://doi.org/10.1038/s41467-020-15631-z

Fletcher, J. L., Makowiecki, K., Cullen, C. L., & Young, K. M. (2021). Oligodendrogenesis and myelination regulate cortical development, plasticity and circuit function. *Seminars in Cell & Developmental Biology*, *118*, 14-23. https://doi.org/10.1016/j.semcdb.2021.03.017

Forkel, S. J., & Hagoort, P. (2024). Redefining language networks : Connectivity beyond localised regions. *Brain Structure and Function*, s00429-024-02859-4. https://doi.org/10.1007/s00429-024-02859-4

Fortel, I., Zhan, L., Ajilore, O., Wu, Y., Mackin, S., & Leow, A. (2023). Disrupted Excitation-Inhibition Balance in Cognitively Normal Individuals at Risk of Alzheimer's Disease. *Journal of Alzheimer's Disease*, *95*(4), 1449-1467. https://doi.org/10.3233/JAD-230035

Fotiadis, P., Parkes, L., Davis, K. A., Satterthwaite, T. D., Shinohara, R. T., & Bassett, D. S. (2024). Structure–function coupling in macroscale human brain networks. *Nature Reviews Neuroscience*, *25*(10), 688-704. https://doi.org/10.1038/s41583-024-00846-6

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences*, *102*(27), 9673-9678. https://doi.org/10.1073/pnas.0504136102

Gatica, M., Cofré, R., Mediano, P. A. M., Rosas, F. E., Orio, P., Diez, I., Swinnen, S. P., & Cortes, J. M. (2021). High-Order Interdependencies in the Aging Brain. *Brain Connectivity*, *11*(9), 734-744. https://doi.org/10.1089/brain.2020.0982

Gatica, M., E. Rosas, F., A. M. Mediano, P., Diez, I., P. Swinnen, S., Orio, P., Cofré, R., & M. Cortes, J. (2022). High-order functional redundancy in ageing explained via alterations in the connectome in a whole-brain model. *PLOS Computational Biology*, *18*(9), e1010431. https://doi.org/10.1371/journal.pcbi.1010431

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171-178. https://doi.org/10.1038/nature18933

Godbersen, G. M., Klug, S., Wadsak, W., Pichler, V., Raitanen, J., Rieckmann, A., Stiernman, L., Cocchi, L., Breakspear, M., Hacker, M., Lanzenberger, R., & Hahn, A. (2023). Task-evoked metabolic demands of the posteromedial default mode network are shaped by dorsal attention and frontoparietal control networks. *eLife*, *12*, e84683. https://doi.org/10.7554/eLife.84683

Goelman, G., Dan, R., Bezdicek, O., Jech, R., & Ekstein, D. (2024). Directed functional connectivity of the default-mode-network of young and older healthy subjects. *Scientific Reports*, *14*(1), 4304. https://doi.org/10.1038/s41598-024-54802-6

Gollan, T. H., & Goldrick, M. (2019). Aging deficits in naturalistic speech production and monitoring revealed through reading aloud. *Psychology and Aging*, *34*(1), 25-42. https://doi.org/10.1037/pag0000296

Gordon, E. M., Chauvin, R. J., Van, A. N., Rajesh, A., Nielsen, A., Newbold, D. J., Lynch, C. J., Seider, N. A., Krimmel, S. R., Scheidter, K. M., Monk, J., Miller, R. L., Metoki, A., Montez, D. F., Zheng, A., Elbau, I., Madison, T., Nishino, T., Myers, M. J., … Dosenbach, N. U. F. (2023). A somato-cognitive action network alternates with effector regions in motor cortex. *Nature*, *617*(7960), 351-359. https://doi.org/10.1038/s41586-023-05964-2

Gordon, E. M., Laumann, T. O., Marek, S., Raut, R. V., Gratton, C., Newbold, D. J., Greene, D. J., Coalson, R. S., Snyder, A. Z., Schlaggar, B. L., Petersen, S. E., Dosenbach, N. U. F., & Nelson, S. M. (2020). Default-mode network streams for coupling to language and control systems. *Proceedings of the National Academy of Sciences*, *117*(29), 17308-17319. https://doi.org/10.1073/pnas.2005238117

Goyal, M. S., Hawrylycz, M., Miller, J. A., Snyder, A. Z., & Raichle, M. E. (2014). Aerobic Glycolysis in the Human Brain Is Associated with Development and Neotenous Gene Expression. *Cell Metabolism*, *19*(1), 49-57. https://doi.org/10.1016/j.cmet.2013.11.020

Goyal, M. S., & Raichle, M. E. (2013). Gene expression-based modeling of human cortical synaptic density. *Proceedings of the National Academy of Sciences*, *110*(16), 6571-6576. https://doi.org/10.1073/pnas.1303453110

Goyal, M. S., Vlassenko, A. G., Blazey, T. M., Su, Y., Couture, L. E., Durbin, T. J., Bateman, R. J., Benzinger, T. L.-S., Morris, J. C., & Raichle, M. E. (2017). Loss of Brain Aerobic Glycolysis in Normal Human Aging. *Cell Metabolism*, *26*(2), 353-360.e3. https://doi.org/10.1016/j.cmet.2017.07.010

Guichet, C., Banjac, S., Achard, S., Mermillod, M., & Baciu, M. (2024). Modeling the neurocognitive dynamics of language across the lifespan. *Human Brain Mapping*, *45*(5), e26650. https://doi.org/10.1002/hbm.26650

Guichet, C., Harquel, S., Achard, S., Mermillod, M., & Baciu, M. (2024). *Lifespan Oscillatory Dynamics in Lexical Production : A Population-based MEG Resting-State Analysis*. https://doi.org/10.1101/2024.07.28.605484

Hahn, A., Reed, M. B., Vraka, C., Godbersen, G. M., Klug, S., Komorowski, A., Falb, P., Nics, L., Traub-Weidinger, T., Hacker, M., & Lanzenberger, R. (2024). High-temporal resolution functional PET/MRI reveals coupling between human metabolic and hemodynamic brain response. *European Journal of*

Nuclear Medicine and Molecular Imaging, *51*(5), 1310-1322. https://doi.org/10.1007/s00259-023-06542-4

Hancock, F., Rosas, F. E., Mediano, P. A. M., Luppi, A. I., Cabral, J., Dipasquale, O., & Turkheimer, F. E. (2022). May the 4C's be with you : An overview of complexity-inspired frameworks for analysing resting-state neuroimaging data. *Journal of The Royal Society Interface*, *19*(191), 20220214. https://doi.org/10.1098/rsif.2022.0214

Harada, C. N., Natelson Love, M. C., & Triebel, K. L. (2013). Normal Cognitive Aging. *Clinics in Geriatric Medicine*, *29*(4), 737-752. https://doi.org/10.1016/j.cger.2013.07.002

Hartshorne, J. K., & Germine, L. T. (2015). When Does Cognitive Functioning Peak? The Asynchronous Rise and Fall of Different Cognitive Abilities Across the Life Span. *Psychological Science*, *26*(4), 433-443. https://doi.org/10.1177/0956797614567339

Hasson, U., Egidi, G., Marelli, M., & Willems, R. M. (2018). Grounding the neurobiology of language in first principles : The necessity of non-language-centric explanations for language comprehension. *Cognition*, *180*, 135-157. https://doi.org/10.1016/j.cognition.2018.06.018

Hausman, H. K., Hardcastle, C., Albizu, A., Kraft, J. N., Evangelista, N. D., Boutzoukas, E. M., Langer, K., O'Shea, A., Van Etten, E. J., Bharadwaj, P. K., Song, H., Smith, S. G., Porges, E., DeKosky, S. T., Hishaw, G. A., Wu, S., Marsiske, M., Cohen, R., Alexander, G. E., & Woods, A. J. (2022). Cingulo-opercular and frontoparietal control network connectivity and executive functioning in older adults. *GeroScience*, *44*(2), 847-866. https://doi.org/10.1007/s11357-021-00503-1

Hills, T. T., Todd, P. M., Lazer, D., Redish, A. D., & Couzin, I. D. (2015). Exploration versus exploitation in space, mind, and society. *Trends in Cognitive Sciences*, *19*(1), 46-54. https://doi.org/10.1016/j.tics.2014.10.004

Hirst, R. J., Setti, A., Kenny, R. A., & Newell, F. N. (2019). Age-related sensory decline mediates the Sound-Induced Flash Illusion : Evidence for reliability weighting models of multisensory perception. *Scientific Reports*, *9*(1), 19347. https://doi.org/10.1038/s41598-019-55901-5

Hoffman, P., & MacPherson, S. E. (2022). What determines cognitive estimation ability? Changing contributions of semantic and executive domains as a function of age. *Journal of Neuropsychology*, *16*(3), 481-497. https://doi.org/10.1111/jnp.12279

Hsieh, J. K., Prakash, P. R., Flint, R. D., Fitzgerald, Z., Mugler, E., Wang, Y., Crone, N. E., Templer, J. W., Rosenow, J. M., Tate, M. C., Betzel, R., & Slutzky, M. W. (2024). Cortical sites critical to language function act as connectors between language subnetworks. *Nature Communications*, *15*(1), 7897. https://doi.org/10.1038/s41467-024-51839-z

Hsu, C.-W., Huang, C.-C., Hsu, C.-C. H., Bi, Y., Tzeng, O. J.-L., & Lin, C.-P. (2025). Revisiting human language and speech production network : A meta-analytic connectivity modeling study. *NeuroImage*, *306*, 121008. https://doi.org/10.1016/j.neuroimage.2025.121008

Huo, Z., Rana, B. K., Elman, J. A., Dong, R., Engelman, C. D., Johnson, S. C., Lyons, M. J., Franz, C. E., Kremen, W. S., & Zhao, J. (2020). Metabolic Profiling of Cognitive Aging in Midlife. *Frontiers in Aging Neuroscience*, *12*, 555850. https://doi.org/10.3389/fnagi.2020.555850

Ihle, A., Oris, M., Fagot, D., Maggiori, C., & Kliegel, M. (2016). The association of educational attainment, cognitive level of job, and leisure activities during the course of adulthood with cognitive performance in old age : The role of openness to experience. *International Psychogeriatrics*, *28*(5), 733-740. https://doi.org/10.1017/S1041610215001933

Ihle, A., Zuber, S., Gouveia, É. R., Gouveia, B. R., Mella, N., Desrichard, O., Cullati, S., Oris, M., Maurer, J., & Kliegel, M. (2019). Cognitive Reserve Mediates the Relation between Openness to Experience and Smaller Decline in Executive Functioning. *Dementia and Geriatric Cognitive Disorders*, *48*(1-2), 39-44. https://doi.org/10.1159/000501822

Ionescu, T. (2017). The Variability-Stability-Flexibility Pattern : A Possible Key to Understanding the Flexibility of the Human Mind. *Review of General Psychology*, *21*(2), 123-131. https://doi.org/10.1037/gpr0000110

Iso-Markku, P., Aaltonen, S., Kujala, U. M., Halme, H.-L., Phipps, D., Knittle, K., Vuoksimaa, E., & Waller, K. (2024). Physical Activity and Cognitive Decline Among Older Adults : A Systematic Review and Meta-Analysis. *JAMA Network Open*, *7*(2), e2354285. https://doi.org/10.1001/jamanetworkopen.2023.54285

Jensen, O., & Mazaheri, A. (2010). Shaping Functional Architecture by Oscillatory Alpha Activity : Gating by Inhibition. *Frontiers in Human Neuroscience*, *4*. https://doi.org/10.3389/fnhum.2010.00186

Ji, J. L., Spronk, M., Kulkarni, K., Repovš, G., Anticevic, A., & Cole, M. W. (2019). Mapping the human brain's cortical-subcortical functional network organization. *NeuroImage*, *185*, 35-57. https://doi.org/10.1016/j.neuroimage.2018.10.006

Johnston, W. J., & Freedman, D. J. (2023). Redundant representations are required to disambiguate simultaneously presented complex stimuli. *PLOS Computational Biology*, *19*(8), e1011327. https://doi.org/10.1371/journal.pcbi.1011327

Johnston, W. J., Palmer, S. E., & Freedman, D. J. (2020). Nonlinear mixed selectivity supports reliable neural computation. *PLOS Computational Biology*, *16*(2), e1007544. https://doi.org/10.1371/journal.pcbi.1007544

Juttukonda, M. R., Li, B., Almaktoum, R., Stephens, K. A., Yochim, K. M., Yacoub, E., Buckner, R. L., & Salat, D. H. (2021). Characterizing cerebral hemodynamics across the adult lifespan with arterial spin labeling MRI data from the Human Connectome Project-Aging. *NeuroImage*, *230*, 117807. https://doi.org/10.1016/j.neuroimage.2021.117807

Kahneman, D. (2012). *Thinking, fast and slow*. Penguin Books.

Kernbach, J. M., Yeo, B. T. T., Smallwood, J., Margulies, D. S., Thiebaut De Schotten, M., Walter, H., Sabuncu, M. R., Holmes, A. J., Gramfort, A., Varoquaux, G., Thirion, B., & Bzdok, D. (2018). Subspecialization within default mode nodes characterized in 10,000 UK Biobank participants. *Proceedings of the National Academy of Sciences*, *115*(48), 12295-12300. https://doi.org/10.1073/pnas.1804876115

Khalili, R., Kehayia, E., & Roig, M. (2024). Exercise and Language Performance in Healthy Aging, Stroke and Neurodegenerative Conditions : A Scoping Review. *Canadian Geriatrics Journal*, *27*(3), 324-344. https://doi.org/10.5770/cgj.27.707

Kintsch, W., & Van Dijk, T. A. (1978). Toward a model of text comprehension and production. *Psychological Review*, *85*(5), 363-394. https://doi.org/10.1037/0033-295X.85.5.363

Kleinridders, A., Ferris, H. A., Reyzer, M. L., Rath, M., Soto, M., Manier, M. L., Spraggins, J., Yang, Z., Stanton, R. C., Caprioli, R. M., & Kahn, C. R. (2018). Regional differences in brain glucose metabolism determined by imaging mass spectrometry. *Molecular Metabolism*, *12*, 113-121. https://doi.org/10.1016/j.molmet.2018.03.013

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored information. *Trends in Cognitive Sciences*, *16*(12), 606-617. https://doi.org/10.1016/j.tics.2012.10.007

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations : The inhibition–timing hypothesis. *Brain Research Reviews*, *53*(1), 63-88. https://doi.org/10.1016/j.brainresrev.2006.06.003

Krieger-Redwood, K., Jefferies, E., Karapanagiotidis, T., Seymour, R., Nunes, A., Ang, J. W. A., Majernikova, V., Mollo, G., & Smallwood, J. (2016). Down but not out in posterior cingulate cortex : Deactivation yet functional coupling with prefrontal cortex during demanding semantic cognition. *NeuroImage*, *141*, 366-377. https://doi.org/10.1016/j.neuroimage.2016.07.060

Krieger-Redwood, K., Lanzoni, L., Gonzalez Alam, T. R. J., Jackson, R. L., Smallwood, J., & Jefferies, E. (2025). Divergent and convergent creativity relate to different aspects of semantic control. *Imaging Neuroscience*, *3*, imag_a_00502. https://doi.org/10.1162/imag_a_00502

Krieger-Redwood, K., Wang, H.-T., Poerio, G., Martinon, L. M., Riby, L. M., Smallwood, J., & Jefferies, E. (2019). Reduced semantic control in older adults is linked to intrinsic DMN connectivity. *Neuropsychologia*, *132*, 107133. https://doi.org/10.1016/j.neuropsychologia.2019.107133

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation. *Advances in Methods and Practices in Psychological Science*, *1*(2), 270-280. https://doi.org/10.1177/2515245918771304

Kupis, L., Goodman, Z. T., Kornfeld, S., Hoang, S., Romero, C., Dirks, B., Dehoney, J., Chang, C., Spreng, R. N., Nomi, J. S., & Uddin, L. Q. (2021). Brain Dynamics Underlying Cognitive Flexibility Across the Lifespan. *Cerebral Cortex*, *31*(11), 5263-5274. https://doi.org/10.1093/cercor/bhab156

Kurtin, D. L., Araña-Oiarbide, G., Lorenz, R., Violante, I. R., & Hampshire, A. (2023). Planning ahead : Predictable switching recruits task-active and resting-state networks. *Human Brain Mapping*, *44*(15), 5030-5046. https://doi.org/10.1002/hbm.26430

Laurienti, P. J., Burdette, J. H., Maldjian, J. A., & Wallace, M. T. (2006). Enhanced multisensory integration in older adults. *Neurobiology of Aging*, *27*(8), 1155-1163. https://doi.org/10.1016/j.neurobiolaging.2005.05.024

Lbath, H., Petersen, A., & Achard, S. (2024). Large-scale correlation screening under dependence for brain functional connectivity network inference. *Statistics and Computing*, *34*(2), 90. https://doi.org/10.1007/s11222-024-10411-x

Leech, R., Kamourieh, S., Beckmann, C. F., & Sharp, D. J. (2011). Fractionating the Default Mode Network : Distinct Contributions of the Ventral and Dorsal Posterior Cingulate Cortex to Cognitive Control. *The Journal of Neuroscience*, *31*(9), 3217-3224. https://doi.org/10.1523/JNEUROSCI.5626-10.2011

Leech, R., & Sharp, D. J. (2014). The role of the posterior cingulate cortex in cognition and disease. *Brain*, *137*(1), 12-32. https://doi.org/10.1093/brain/awt162

Leonards, C. A., Harrison, B. J., Jamieson, A. J., Steward, T., Lux, S., Philipsen, A., & Davey, C. G. (2023). A distinct intra-individual suppression subnetwork in the brain's default mode network across cognitive tasks. *Cerebral Cortex*, *33*(8), 4553-4561. https://doi.org/10.1093/cercor/bhac361

Li, X., Salami, A., & Persson, J. (2023). Hub architecture of the human structural connectome : Links to aging and processing speed. *NeuroImage*, *278*, 120270. https://doi.org/10.1016/j.neuroimage.2023.120270

Lindenberger, U., & Baltes, P. B. (1994). Sensory functioning and intelligence in old age : A strong connection. *Psychology and Aging*, *9*(3), 339-355. https://doi.org/10.1037/0882-7974.9.3.339

Lindenberger, U., & Mayr, U. (2014). Cognitive aging : Is there a dark side to environmental support? *Trends in Cognitive Sciences*, *18*(1), 7-15. https://doi.org/10.1016/j.tics.2013.10.006

Loaiza, V. M. (2024). An overview of the hallmarks of cognitive aging. *Current Opinion in Psychology*, *56*, 101784. https://doi.org/10.1016/j.copsyc.2023.101784

Luo, W., Liu, B., Tang, Y., Huang, J., & Wu, J. (2024). Rest to Promote Learning : A Brain Default Mode Network Perspective. *Behavioral Sciences*, *14*(4), 349. https://doi.org/10.3390/bs14040349

Luppi, A. I., Mediano, P. A. M., Rosas, F. E., Holland, N., Fryer, T. D., O'Brien, J. T., Rowe, J. B., Menon, D. K., Bor, D., & Stamatakis, E. A. (2022). A synergistic core for human brain evolution and cognition. *Nature Neuroscience*, *25*(6), 771-782. https://doi.org/10.1038/s41593-022-01070-0

Luppi, A. I., Rosas, F. E., Noonan, M. P., Mediano, P. A. M., Kringelbach, M. L., Carhart-Harris, R. L., Stamatakis, E. A., Vernon, A. C., & Turkheimer, F. E. (2024). Oxygen and the Spark of Human Brain Evolution : Complex Interactions of Metabolism and Cortical Expansion across Development and Evolution. *The Neuroscientist*, *30*(2), 173-198. https://doi.org/10.1177/10738584221138032

Luppi, A. I., Sanz Perl, Y., Vohryzek, J., Mediano, P. A. M., Rosas, F. E., Milisav, F., Suarez, L. E., Gini, S., Gutierrez-Barragan, D., Gozzi, A., Misic, B., Deco, G., & Kringelbach, M. L. (2024). *Competitive interactions shape brain dynamics and computation across species*. https://doi.org/10.1101/2024.10.19.619194

Magistretti, P. J., & Allaman, I. (2015). A Cellular Perspective on Brain Energy Metabolism and Functional Imaging. *Neuron*, *86*(4), 883-901. https://doi.org/10.1016/j.neuron.2015.03.035

Makowski, D., Ben-Shachar, M., & Lüdecke, D. (2019). bayestestR : Describing Effects and their Uncertainty, Existence and Significance within the Bayesian Framework. *Journal of Open Source Software*, *4*(40), 1541. https://doi.org/10.21105/joss.01541

Marder, E., & Goaillard, J.-M. (2006). Variability, compensation and homeostasis in neuron and network function. *Nature Reviews Neuroscience*, *7*(7), 563-574. https://doi.org/10.1038/nrn1949

Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., Bezgin, G., Eickhoff, S. B., Castellanos, F. X., Petrides, M., Jefferies, E., & Smallwood, J. (2016). Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences*, *113*(44), 12574-12579. https://doi.org/10.1073/pnas.1608282113

Martin, A. (2016). GRAPES—Grounding representations in action, perception, and emotion systems : How object properties and categories are represented in the human brain. *Psychonomic Bulletin & Review*, *23*(4), 979-990. https://doi.org/10.3758/s13423-015-0842-3

Martin, S., Saur, D., & Hartwigsen, G. (2022). Age-Dependent Contribution of Domain-General Networks to Semantic Cognition. *Cerebral Cortex*, *32*(4), 870-890. https://doi.org/10.1093/cercor/bhab252

Martínez-Ferreiro, S. (2024). Naming as a window to word retrieval changes in healthy and pathological ageing : Methodological considerations. *International Journal of Language & Communication Disorders*, *59*(1), 68-83. https://doi.org/10.1111/1460-6984.12827

Mattson, M. P. (2025). Pathways toward and away from a metabolic cliff during brain aging. *Frontiers in Science*, *3*, 1571009. https://doi.org/10.3389/fsci.2025.1571009

Menon, V. (2023). 20 years of the default mode network : A review and synthesis. *Neuron*, *111*(16), 2469-2487. https://doi.org/10.1016/j.neuron.2023.04.023

Montoliu, T., Zapater-Fajarí, M., Hidalgo, V., & Salvador, A. (2023). Openness to experience and cognitive functioning and decline in older adults : The mediating role of cognitive reserve. *Neuropsychologia*, *188*, 108655. https://doi.org/10.1016/j.neuropsychologia.2023.108655

Mowinckel, A. M., & Vidal-Piñeiro, D. (2020). Visualization of Brain Statistics With R Packages *ggseg* and *ggseg3d*. *Advances in Methods and Practices in Psychological Science*, *3*(4), 466-483. https://doi.org/10.1177/2515245920928009

Murphy, K., & Fox, M. D. (2017). Towards a consensus regarding global signal regression for resting state functional connectivity MRI. *NeuroImage*, *154*, 169-173. https://doi.org/10.1016/j.neuroimage.2016.11.052

Nieberlein, L., Martin, S., Williams, K. A., Gussew, A., Cyriaks, S. D., Scheer, M., Rampp, S., Prell, J., & Hartwigsen, G. (2024). *Semantic integration demands modulate large-scale network interactions in the brain*. https://doi.org/10.1101/2024.11.20.624508

O'Byrne, J., & Jerbi, K. (2022). How critical is brain criticality? *Trends in Neurosciences*, *45*(11), 820-837. https://doi.org/10.1016/j.tins.2022.08.007

Palombit, A., Silvestri, E., Volpi, T., Aiello, M., Cecchin, D., Bertoldo, A., & Corbetta, M. (2022). Variability of regional glucose metabolism and the topology of functional networks in the human brain. *NeuroImage*, *257*, 119280. https://doi.org/10.1016/j.neuroimage.2022.119280

Paquola, C., Garber, M., Frässle, S., Royer, J., Zhou, Y., Tavakol, S., Rodriguez-Cruces, R., Cabalo, D. G., Valk, S., Eickhoff, S., Margulies, D. S., Evans, A., Amunts, K., Jefferies, E., Smallwood, J., & Bernhardt, B. C. (2025). The architecture of the human default mode network explored through cytoarchitecture, wiring and signal flow. *Nature Neuroscience*. https://doi.org/10.1038/s41593-024-01868-0

Paquola, C., Vos De Wael, R., Wagstyl, K., Bethlehem, R. A. I., Hong, S.-J., Seidlitz, J., Bullmore, E. T., Evans, A. C., Misic, B., Margulies, D. S., Smallwood, J., & Bernhardt, B. C. (2019). Microstructural and functional gradients are increasingly dissociated in transmodal cortices. *PLOS Biology*, *17*(5), e3000284. https://doi.org/10.1371/journal.pbio.3000284

Park, D. C., & Reuter-Lorenz, P. (2009). The Adaptive Brain : Aging and Neurocognitive Scaffolding. *Annual Review of Psychology*, *60*(1), 173-196. https://doi.org/10.1146/annurev.psych.59.103006.093656

Paulson, O. B., Hasselbalch, S. G., Rostrup, E., Knudsen, G. M., & Pelligrino, D. (2010). Cerebral Blood Flow Response to Functional Activation. *Journal of Cerebral Blood Flow & Metabolism*, *30*(1), 2-14. https://doi.org/10.1038/jcbfm.2009.188

Pellerin, L. (2025). The future of healthy brain aging will be metabolic : A predictive lesson from data-driven modeling. *Frontiers in Science*, *3*, 1561735. https://doi.org/10.3389/fsci.2025.1561735

Perez, D. C., Hernandez, J. J., Wulfekuhle, G., & Gratton, C. (2025). Variation in brain aging : A review and perspective on the utility of individualized approaches to the study of functional networks in aging. *Neurobiology of Aging*, *147*, 68-87. https://doi.org/10.1016/j.neurobiolaging.2024.11.010

Politakis, V. A., Slana Ozimič, A., & Repovš, G. (2022). Cognitive Control Challenge Task Across the Lifespan. *Frontiers in Psychology*, *12*, 789816. https://doi.org/10.3389/fpsyg.2021.789816

Rahman, F., Tsvetanov, K. A., Feron, J., Mullinger, K., Joyce, K., Gilani, A., Fernandes, E. G., Wetterlin, A., Wheeldon, L., Lucas, S. J. E., & Segaert, K. (2023). *Lifestyle and brain health determinants of word-finding failures in healthy ageing*. https://doi.org/10.1101/2023.12.08.570799

Raichle, M. E. (2015). The Brain's Default Mode Network. *Annual Review of Neuroscience*, *38*(1), 433-447. https://doi.org/10.1146/annurev-neuro-071013-014030

Ramchandran, K., Zeien, E., & Andreasen, N. C. (2019). Distributed neural efficiency : Intelligence and age modulate adaptive allocation of resources in the brain. *Trends in Neuroscience and Education*, *15*, 48-61. https://doi.org/10.1016/j.tine.2019.02.006

Raz, N., & Daugherty, A. M. (2018). Pathways to Brain Aging and Their Modifiers : Free-Radical-Induced Energetic and Neural Decline in Senescence (FRIENDS) Model - A Mini-Review. *Gerontology*, *64*(1), 49-57. https://doi.org/10.1159/000479508

Reilly, J., Shain, C., Borghesani, V., Kuhnke, P., Vigliocco, G., Peelle, J. E., Mahon, B., Buxbaum, L., Majid, A., Brysbaert, M., Borghi, A. M., De Deyne, S., Dove, G., Papeo, L., Pexman, P. M., Poeppel, D., Lupyan, G., Boggio, P., Hickock, G., … Vinson, D. (2023). *What we mean when we say semantic : A Consensus statement on the nomenclature of semantic memory* [Preprint]. PsyArXiv. https://doi.org/10.31234/osf.io/xrnb2

Riedl, V., Bienkowska, K., Strobel, C., Tahmasian, M., Grimmer, T., Förster, S., Friston, K. J., Sorg, C., & Drzezga, A. (2014). Local Activity Determines Functional Connectivity in the Resting Human Brain : A Simultaneous FDG-PET/fMRI Study. *The Journal of Neuroscience*, *34*(18), 6260-6266. https://doi.org/10.1523/JNEUROSCI.0492-14.2014

Roger, E., Banjac, S., Thiebaut de Schotten, M., & Baciu, M. (2022). Missing links : The functional unification of language and memory (LUM). *Neuroscience & Biobehavioral Reviews*, *133*, 104489. https://doi.org/10.1016/j.neubiorev.2021.12.012

Roger, E., Rodrigues De Almeida, L., Loevenbruck, H., Perrone-Bertolotti, M., Cousin, E., Schwartz, J. L., Perrier, P., Dohen, M., Vilain, A., Baraduc, P., Achard, S., & Baciu, M. (2022). Unraveling the functional attributes of the language connectome : Crucial subnetworks, flexibility and variability. *NeuroImage*, *263*, 119672. https://doi.org/10.1016/j.neuroimage.2022.119672

Ross, J. M., Öberg, J., Brené, S., Coppotelli, G., Terzioglu, M., Pernold, K., Goiny, M., Sitnikov, R., Kehr, J., Trifunovic, A., Larsson, N.-G., Hoffer, B. J., & Olson, L. (2010). High brain lactate is a hallmark of aging and caused by a shift in the lactate dehydrogenase A/B ratio. *Proceedings of the National Academy of Sciences*, *107*(46), 20087-20092. https://doi.org/10.1073/pnas.1008189107

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity : Uses and interpretations. *NeuroImage*, *52*(3), 1059-1069. https://doi.org/10.1016/j.neuroimage.2009.10.003

Saberi, M., Khosrowabadi, R., Khatibi, A., Misic, B., & Jafari, G. (2022). Pattern of frustration formation in the functional brain network. *Network Neuroscience*, *6*(4), 1334-1356. https://doi.org/10.1162/netn_a_00268

Saberi, M., Rieck, J. R., Golafshan, S., Grady, C. L., Misic, B., Dunkley, B. T., & Khatibi, A. (2024). The brain selectively allocates energy to functional brain networks under cognitive control. *Scientific Reports*, *14*(1), 32032. https://doi.org/10.1038/s41598-024-83696-7

Sala, A., Lizarraga, A., Caminiti, S. P., Calhoun, V. D., Eickhoff, S. B., Habeck, C., Jamadar, S. D., Perani, D., Pereira, J. B., Veronese, M., & Yakushev, I. (2023). Brain connectomics : Time for a molecular imaging perspective? *Trends in Cognitive Sciences*, *27*(4), 353-366. https://doi.org/10.1016/j.tics.2022.11.015

Salthouse, T. A. (2019). Trajectories of normal cognitive aging. *Psychology and Aging*, *34*(1), 17-24. https://doi.org/10.1037/pag0000288

Sestieri, C., Corbetta, M., Spadone, S., Romani, G. L., & Shulman, G. L. (2014). Domain-general Signals in the Cingulo-opercular Network for Visuospatial Attention and Episodic Memory. *Journal of Cognitive Neuroscience*, *26*(3), 551-568. https://doi.org/10.1162/jocn_a_00504

Shafto, M. A., & Tyler, L. K. (2014). Language in the aging brain : The network dynamics of cognitive decline and preservation. *Science*, *346*(6209), 583-587. https://doi.org/10.1126/science.1254404

Shichkova, P., Coggan, J. S., Kanari, L., Boci, E., Favreau, C., Antonel, S. M., Keller, D., & Markram, H. (2025). Breakdown and repair of metabolism in the aging brain. *Frontiers in Science*, *3*, 1441297. https://doi.org/10.3389/fsci.2025.1441297

Shichkova, P., Coggan, J. S., Markram, H., & Keller, D. (2024). Brain Metabolism in Health and Neurodegeneration : The Interplay Among Neurons and Astrocytes. *Cells*, *13*(20), 1714. https://doi.org/10.3390/cells13201714

Shine, J. M., & Poldrack, R. A. (2018). Principles of dynamic network reconfiguration across diverse brain states. *NeuroImage*, *180*, 396-405. https://doi.org/10.1016/j.neuroimage.2017.08.010

Shinn, M., Hu, A., Turner, L., Noble, S., Preller, K. H., Ji, J. L., Moujaes, F., Achard, S., Scheinost, D., Constable, R. T., Krystal, J. H., Vollenweider, F. X., Lee, D., Anticevic, A., Bullmore, E. T., & Murray, J. D. (2023). Functional brain networks reflect spatial and temporal autocorrelation. *Nature Neuroscience*, *26*(5), 867-878. https://doi.org/10.1038/s41593-023-01299-3

Shokri-Kojori, E., Tomasi, D., Alipanahi, B., Wiers, C. E., Wang, G.-J., & Volkow, N. D. (2019). Correspondence between cerebral glucose metabolism and BOLD reveals relative power and cost in human brain. *Nature Communications*, *10*(1), 690. https://doi.org/10.1038/s41467-019-08546-x

Shulman, R. G., Rothman, D. L., Behar, K. L., & Hyder, F. (2004). Energetic basis of brain activity : Implications for neuroimaging. *Trends in Neurosciences*, *27*(8), 489-495. https://doi.org/10.1016/j.tins.2004.06.005

Smallwood, J., Bernhardt, B. C., Leech, R., Bzdok, D., Jefferies, E., & Margulies, D. S. (2021). The default mode network in cognition : A topographical perspective. *Nature Reviews Neuroscience*, *22*(8), 503-513. https://doi.org/10.1038/s41583-021-00474-4

Smith, V., Mitchell, D. J., & Duncan, J. (2018). Role of the Default Mode Network in Cognitive Transitions. *Cerebral Cortex*, *28*(10), 3685-3696. https://doi.org/10.1093/cercor/bhy167

Sporns, O., & Betzel, R. F. (2016). Modular Brain Networks. *Annual Review of Psychology*, *67*(1), 613-640. https://doi.org/10.1146/annurev-psych-122414-033634

Spreng, R. N., DuPre, E., Selarka, D., Garcia, J., Gojkovic, S., Mildner, J., Luh, W.-M., & Turner, G. R. (2014). Goal-Congruent Default Network Activity Facilitates Cognitive Control. *The Journal of Neuroscience*, *34*(42), 14108-14114. https://doi.org/10.1523/jneurosci.2815-14.2014

Spreng, R. N., & Turner, G. R. (2019). The Shifting Architecture of Cognition and Brain Function in Older Adulthood. *Perspectives on Psychological Science*, *14*(4), 523-542. https://doi.org/10.1177/1745691619827511

Spreng, R. N., & Turner, G. R. (2021). From exploration to exploitation : A shifting mental mode in late life development. *Trends in Cognitive Sciences*, *25*(12), 1058-1071. https://doi.org/10.1016/j.tics.2021.09.001

Stanley, J. A., & Raz, N. (2018). Functional Magnetic Resonance Spectroscopy : The "New" MRS for Cognitive Neuroscience and Psychiatry Research. *Frontiers in Psychiatry*, *9*, 76. https://doi.org/10.3389/fpsyt.2018.00076

Stiernman, L. J., Grill, F., Hahn, A., Rischka, L., Lanzenberger, R., Panes Lundmark, V., Riklund, K., Axelsson, J., & Rieckmann, A. (2021). Dissociations between glucose metabolism and blood oxygenation in the human default mode network revealed by simultaneous PET-fMRI. *Proceedings of the National Academy of Sciences*, *118*(27), e2021913118. https://doi.org/10.1073/pnas.2021913118

Stolz, C., Bulla, A., Soch, J., Schott, B. H., & Richter, A. (2023). Openness to Experience is associated with neural and performance measures of memory in older adults. *Social Cognitive and Affective Neuroscience*, *18*(1), nsad041. https://doi.org/10.1093/scan/nsad041

Stranahan, A. M., & Mattson, M. P. (2012). Metabolic Reserve as a Determinant of Cognitive Aging. *Journal of Alzheimer's Disease*, *30*(s2), S5-S13. https://doi.org/10.3233/JAD-2011-110899

Stumme, J., Krämer, C., Miller, T., Schreiber, J., Caspers, S., & Jockwitz, C. (2022). Interrelating differences in structural and functional connectivity in the older adult's brain. *Human Brain Mapping*, *43*(18), 5543-5561. https://doi.org/10.1002/hbm.26030

Sydnor, V. J., Larsen, B., Bassett, D. S., Alexander-Bloch, A., Fair, D. A., Liston, C., Mackey, A. P., Milham, M. P., Pines, A., Roalf, D. R., Seidlitz, J., Xu, T., Raznahan, A., & Satterthwaite, T. D. (2021).

Neurodevelopment of the association cortices : Patterns, mechanisms, and implications for psychopathology. *Neuron*, *109*(18), 2820-2846. https://doi.org/10.1016/j.neuron.2021.06.016

Takada, H., Nagata, K., Hirata, Y., Satoh, Y., Watahiki, Y., Sugawara, J., Yokoyama, E., Kondoh, Y., Shishido, F., Inugami, A., Fujita, H., Ogawa, T., Murakami, M., Lida, H., & Kanno, L. (1992). Age-related decline of cerebral oxygen metabolism in normal population detected with positron emission tomography. *Neurological Research*, *14*(2), 128-131. https://doi.org/10.1080/01616412.1992.11740031

Takagi, K. (2021). Energy constraints on brain network formation. *Scientific Reports*, *11*(1), 11745. https://doi.org/10.1038/s41598-021-91250-y

Tarumi, T., & Zhang, R. (2018). Cerebral blood flow in normal aging adults : Cardiovascular determinants, clinical implications, and aerobic fitness. *Journal of Neurochemistry*, *144*(5), 595-608. https://doi.org/10.1111/jnc.14234

Taylor, J. R., Williams, N., Cusack, R., Auer, T., Shafto, M. A., Dixon, M., Tyler, L. K., Cam-CAN, & Henson, R. N. (2017). The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data repository : Structural and functional MRI, MEG, and cognitive data from a cross-sectional adult lifespan sample. *NeuroImage*, *144*, 262-269. https://doi.org/10.1016/j.neuroimage.2015.09.018

Thambisetty, M., Beason-Held, L. L., An, Y., Kraut, M., Metter, J., Egan, J., Ferrucci, L., O'Brien, R., & Resnick, S. M. (2013). Impaired glucose tolerance in midlife and longitudinal changes in brain function during aging. *Neurobiology of Aging*, *34*(10), 2271-2276. https://doi.org/10.1016/j.neurobiolaging.2013.03.025

Thiebaut de Schotten, M., & Forkel, S. J. (2022). The emergent properties of the connected brain. *Science*, *378*(6619), 505-510. https://doi.org/10.1126/science.abq2591

Thomas Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125-1165. https://doi.org/10.1152/jn.00338.2011

Tomasi, D., Wang, G.-J., & Volkow, N. D. (2013). Energetic cost of brain functional connectivity. *Proceedings of the National Academy of Sciences*, *110*(33), 13642-13647. https://doi.org/10.1073/pnas.1303346110

Uddin, L. Q. (2021). Cognitive and behavioural flexibility : Neural mechanisms and clinical considerations. *Nature Reviews Neuroscience*, *22*(3), 167-179. https://doi.org/10.1038/s41583-021-00428-w

United Nations (Éd.). (2023). *Leaving no one behind in an ageing world*. United Nations.

Utevsky, A. V., Smith, D. V., & Huettel, S. A. (2014). Precuneus Is a Functional Core of the Default-Mode Network. *The Journal of Neuroscience*, *34*(3), 932-940. https://doi.org/10.1523/JNEUROSCI.4227-13.2014

Vaishnavi, S. N., Vlassenko, A. G., Rundle, M. M., Snyder, A. Z., Mintun, M. A., & Raichle, M. E. (2010). Regional aerobic glycolysis in the human brain. *Proceedings of the National Academy of Sciences*, *107*(41), 17757-17762. https://doi.org/10.1073/pnas.1010459107

van den Heuvel, M. P., de Lange, S. C., Zalesky, A., Seguin, C., Yeo, B. T. T., & Schmidt, R. (2017). Proportional thresholding in resting-state fMRI functional connectivity networks and consequences for patient-control connectome studies : Issues and recommendations. *NeuroImage*, *152*, 437-449. https://doi.org/10.1016/j.neuroimage.2017.02.005

Verhaegen, C., & Poncelet, M. (2013). Changes in Naming and Semantic Abilities With Aging From 50 to 90 years. *Journal of the International Neuropsychological Society*, *19*(2), 119-126. https://doi.org/10.1017/S1355617712001178

Voigt, K., Liang, E. X., Misic, B., Ward, P. G. D., Egan, G. F., & Jamadar, S. D. (2023). Metabolic and functional connectivity provide unique and complementary insights into cognition-connectome relationships. *Cerebral Cortex*, *33*(4), 1476-1488. https://doi.org/10.1093/cercor/bhac150

Volpi, T., Silvestri, E., Aiello, M., Lee, J. J., Vlassenko, A. G., Goyal, M. S., Corbetta, M., & Bertoldo, A. (2024). The brain's "dark energy" puzzle : How strongly is glucose metabolism linked to resting-state brain activity? *Journal of Cerebral Blood Flow & Metabolism*, *44*(8), 1433-1449. https://doi.org/10.1177/0271678X241237974

Wallis, G., Stokes, M., Cousijn, H., Woolrich, M., & Nobre, A. C. (2015). Frontoparietal and Cingulo-opercular Networks Play Dissociable Roles in Control of Working Memory. *Journal of Cognitive Neuroscience*, *27*(10), 2019-2034. https://doi.org/10.1162/jocn_a_00838

Wan, B., Riedl, V., Castrillon, G., Kirsch, M., & Valk, S. (2024). *Bridging Glucose Metabolism and Intrinsic Functional Organization of the Human Cortex*. https://doi.org/10.1101/2024.09.26.615152

Wang, H.-T., Meisler, S. L., Sharmarke, H., Clarke, N., Gensollen, N., Markiewicz, C. J., Paugam, F., Thirion, B., & Bellec, P. (2024). Continuous evaluation of denoising strategies in resting-state fMRI connectivity using fMRIPrep and Nilearn. *PLOS Computational Biology*, *20*(3), e1011942. https://doi.org/10.1371/journal.pcbi.1011942

Wang, R., Liu, M., Cheng, X., Wu, Y., Hildebrandt, A., & Zhou, C. (2021). Segregation, integration, and balance of large-scale resting brain networks configure different cognitive abilities. *Proceedings of the National Academy of Sciences*, *118*(23), e2022288118. https://doi.org/10.1073/pnas.2022288118

Wang, X., Krieger-Redwood, K., Cui, Y., Smallwood, J., Du, Y., & Jefferies, E. (2024). Macroscale brain states support the control of semantic cognition. *Communications Biology*, *7*(1), 926. https://doi.org/10.1038/s42003-024-06630-7

Wang, X., Margulies, D. S., Smallwood, J., & Jefferies, E. (2020). A gradient from long-term memory to novel cognition : Transitions through default mode and executive cortex. *NeuroImage*, *220*, 117074. https://doi.org/10.1016/j.neuroimage.2020.117074

Willbrand, E. H., Parker, B. J., Voorhies, W. I., Miller, J. A., Lyu, I., Hallock, T., Aponik-Gremillion, L., Koslov, S. R., Alzheimer's Disease Neuroimaging Initiative, Bunge, S. A., Foster, B. L., & Weiner, K. S. (2022). Uncovering a tripartite landmark in posterior cingulate cortex. *Science Advances*, *8*(36), eabn9516. https://doi.org/10.1126/sciadv.abn9516

Williams, K. A., Numssen, O., & Hartwigsen, G. (2022). Task-Specific Network Interactions across Key Cognitive Domains. *Cerebral Cortex*, bhab531. https://doi.org/10.1093/cercor/bhab531

Wright, H. H. (Éd.). (2016). *Cognition, language and aging*. John Benjamins Publishing Company.

Wyatt, L. E., Hewan, P. A., Hogeveen, J., Spreng, R. N., & Turner, G. R. (2024). Exploration versus exploitation decisions in the human brain : A systematic review of functional neuroimaging and neuropsychological studies. *Neuropsychologia*, *192*, 108740. https://doi.org/10.1016/j.neuropsychologia.2023.108740

Xia, H., He, Q., & Chen, A. (2022). Understanding cognitive control in aging : A brain network perspective. *Frontiers in Aging Neuroscience*, *14*, 1038756. https://doi.org/10.3389/fnagi.2022.1038756

Xia, H., Hou, Y., Li, Q., & Chen, A. (2024). A meta-analysis of cognitive flexibility in aging : Perspective from functional network and lateralization. *Human Brain Mapping*, *45*(14), e70031. https://doi.org/10.1002/hbm.70031

Xie, K., Royer, J., Rodriguez-Cruces, R., Horwood, L., Ngo, A., Arafat, T., Auer, H., Sahlas, E., Chen, J., Zhou, Y., Valk, S. L., Hong, S., Frauscher, B., Pana, R., Bernasconi, A., Bernasconi, N., Concha, L., & Bernhardt, B. C. (2025). Temporal Lobe Epilepsy Perturbs the Brain-Wide Excitation-Inhibition Balance : Associations with Microcircuit Organization, Clinical Parameters, and Cognitive Dysfunction. *Advanced Science*, 2406835. https://doi.org/10.1002/advs.202406835

Yang, T., Allen, R. J., Waterman, A. H., Graham, A. J., Su, X., & Gao, Y. (2024). Exploring techniques for encoding spoken instructions in working memory : A comparison of verbal rehearsal, motor imagery, self-enactment and action observation. *Memory*, *32*(1), 41-54. https://doi.org/10.1080/09658211.2023.2273763

Zang, Y., Jiang, T., Lu, Y., He, Y., & Tian, L. (2004). Regional homogeneity approach to fMRI data analysis. *NeuroImage*, *22*(1), 394-400. https://doi.org/10.1016/j.neuroimage.2003.12.030

Zapparoli, L., Mariano, M., & Paulesu, E. (2022). How the motor system copes with aging : A quantitative meta-analysis of the effect of aging on motor function control. *Communications Biology*, *5*(1), 79. https://doi.org/10.1038/s42003-022-03027-2

Zheng, A., Montez, D. F., Marek, S., Gilmore, A. W., Newbold, D. J., Laumann, T. O., Kay, B. P., Seider, N. A., Van, A. N., Hampton, J. M., Alexopoulos, D., Schlaggar, B. L., Sylvester, C. M., Greene, D. J., Shimony, J. S., Nelson, S. M., Wig, G. S., Gratton, C., McDermott, K. B., … Dosenbach, N. U. F. (2021). Parallel hippocampal-parietal circuits for self- and goal-oriented processing. *Proceedings of the National Academy of Sciences*, *118*(34), e2101743118. https://doi.org/10.1073/pnas.2101743118

Zhou, A. X., Duncan, J., & Mitchell, D. J. (2025). Default mode network activation at task switches reflects mental task-set structure. *Imaging Neuroscience*, *3*, imag_a_00515. https://doi.org/10.1162/imag_a_00515

Zioga, I., Kenett, Y. N., Giannopoulos, A., & Luft, C. D. B. (2024). The role of alpha oscillations in free- and goal-directed semantic associations. *Human Brain Mapping*, *45*(10), e26770. https://doi.org/10.1002/hbm.26770