ASPPs multimerize protein phosphatase 1

Derek T. Wei[1,2], Kayleigh N. Morrison[1,4], Gwendolyn M. Beacham[1,2,#a], Erika

Beyrent[1,2,#b], Ying Zhang[3], Laurence Florens[3], Gunther Hollopeter[1,2,4]*

[1] Department of Molecular Medicine, Cornell University, Ithaca, New York, United States

of America; [2] Field of Biochemistry, Molecular, and Cell Biology, Cornell University,

Ithaca, New York, United States of America; [3] Stowers Institute for Medical Research,

Kansas City, Missouri, United States of America; [4] Field of Biomedical and Biological

Science, Cornell University, Ithaca, New York, United States of America.

[#a] Current address: Department of Medicine, Section of Hematology and Medical

Oncology, Boston University Chobanian and Avedisian School of Medicine and Boston

Medical Center, Boston, Massachusetts, United States of America.

[#b] Current address: Penn State Neuroscience Institute – University Park, The

Pennsylvania State University and Department of Engineering Science and Mechanics,

The Pennsylvania State University, University Park, Pennsylvania, United States of

America.

*Corresponding author.

Email: gh383@cornell.edu (GH)

Abstract

Protein Phosphatase 1 (PP1) activity is thought to be spatiotemporally defined by hundreds of different regulatory subunits, but their mechanisms of action are largely unknown. The Ankyrin repeat, SH3-domain, and Proline-rich region containing Proteins (ASPPs) bind and localize PP1 to cell-cell junctions. Here, we show ASPPs bind superstoichiometric amounts of PP1. Missense mutations in the ankyrin repeats of ASPPs, that were previously isolated from a forward genetic screen in *Caenorhabditis elegans*, reduce the stoichiometry of PP1 binding. Forcing PP1 oligomerization restores mutant ASPP function *in vivo*. We propose that ASPPs multimerize PP1 to establish a concentrated hub of phosphatase activity at cell-cell junctions.

Author Summary

We have elucidated a new mechanism governing protein phosphatase 1 (PP1) activity. A family of proteins called the ASPPs function to spatially regulate PP1 by recruiting active phosphatase to specific subcellular locations. Critically, we observed that ASPPs promote the formation of higher-order PP1 assemblies – a previously unrecognized regulatory mechanism. We identified specific ASPP mutants in our nematode model organism that disrupt PP1 oligomerization, leading to altered development. However, inducing PP1 clustering was sufficient to rescue these ASPP mutants, underscoring the functional significance of ASPP-mediated PP1 oligomerization. These results provide new insights into the intricate control of cellular signaling pathways mediated by PP1

45 and may have implications for understanding diseases associated with dysregulated

46 phosphatase activity.

Introduction

There are far fewer protein phosphatases than protein kinases in the human genome

[1,2], so how dephosphorylation reactions are sculpted in order to match kinase

diversity is an open question. The predominant serine/threonine protein phosphatase

catalytic subunit, Protein Phosphatase 1 (PP1), is thought to achieve spatiotemporal

precision by binding hundreds of different regulatory subunits [3,4]. These regulatory

subunits have so far been shown to specify subcellular localization and substrate

selection of the catalytic subunit. For example, a glycogen-binding regulatory subunit is

thought to bring the PP1 catalytic subunit to glycogen in order to focus phosphatase

activity on nearby metabolic enzymes [5]. Another example is spinophilin, which directs

PP1 to act on glutamate receptor 1 while preventing reactions with other substrates,

such as phosphorylase a [6]. However, mechanisms of action for most regulatory

subunits are unknown [7].

The Ankyrin repeat, SH3-domain, and Proline-rich region containing Proteins (ASPPs)

are a family of medically important PP1 regulatory subunits with an unclear function.

There are three vertebrate ASPP homologs – ASPP1, ASPP2, and iASPP. Forward-

genetic studies across cows [8–10], mice [11,12], and humans [13,14] connect

mutations in iASPP, the most conserved homolog [15], to a lethal cardiocutaneous

disease. While studies originally reported that the ASPPs interact with p53 [16–18], it

has been difficult to attribute p53 dysregulation to phenotypes associated with loss of

iASPP [19]. The ASPP C-terminal ankyrin repeats and SH3 domain bind PP1 with an

affinity that is ~100-fold higher than p53 [20–25]. Additionally, mutating ASPP residues

predicted to interact with PP1 produces a phenotype similar to the ASPP null mutant in

Drosophila [24]. These data suggest that ASPPs may regulate PP1, but the mechanism

is unclear.

Several groups have proposed that ASPPs function at cellular junctions. iASPP

colocalizes with desmosomes in cardiomyocytes and keratinocytes [19,26]. iASPP

knockout and knockdown also results in reduced expression of numerous junctional

components that are found in desmosomes, tight junctions, GAP junctions, and

adherens junctions [19,27,28]. ASPP2 also localizes to apical cell-cell junctions in

epithelial cells and associates with the PAR polarity complex [29–31]. Deletion of

ASPP2 in mice reduces PAR3 localization to apical cell junctions and slows the

formation of tight junctions [30,32]. ASPP2 null mice are similar to iASPP mutants in

that they are postnatally lethal and have skin and heart defects [29,33–35]. In

Drosophila melanogaster, the sole ASPP homolog is also reported to colocalize with

adherens junction components [36] and its knockout leads to disorganized epithelial cell

junctions [37]. Therefore, ASPPs appear to have a conserved role in regulating cellular

junctions.

One holistic model proposed for the ASPPs is that they bind and localize PP1 to cell

junctions in order to dephosphorylate specific junctional components [24,32,38,39].

Consistent with this model, we found that phosphatase localization to epithelial cell-cell

junctions is dependent on the sole ASPP homolog in *C. elegans* [25]. However, we

demonstrated that mere localization of phosphatase to epithelial junctions could not

recapitulate ASPP function [25]. Instead, we proposed that the highly conserved ASPP

C-terminal ankyrin repeats and SH3 domain (hereafter called the ASPP 'C-terminus') is

required to modulate PP1 activity through an additional unknown mechanism.

In this study, we find that ASPPs perform a new, unanticipated function – clustering of

PP1. In addition to localizing and promoting PP1 activity at epithelial junctions, we

discover that ASPPs bind superstoichiometric amounts of PP1. We show that semi-

dominant ASPP mutations, identified in a *C. elegans* suppression screen, reduce the

ratio of PP1 to ASPP. Forcing oligomerization of the phosphatase catalytic subunit

restores ASPP function in these mutants. We therefore propose that ASPPs multimerize

PP1 to form dephosphorylation centers at epithelial cell junctions.

Results

APE-1 function requires an N-terminal helix that localizes to

epithelial junctions

Previously, we found that the N-terminal 519 residues of the sole *C. elegans* ASPP

homolog (APE-1) are required for both function and localization [25]. However, the

protein sequence mediating localization was unclear as this region is poorly conserved.

Using AlphaFold structural prediction, we noted that a region predicted to form a helix of

~100 residues is embedded within the first half of APE-1 (Fig 1A, S1A). We truncated

the endogenous APE-1 using CRISPR to see if this region might encode a localization

signal (Fig 1A). We found that the presumptive helix does localize a GFP tag to

epithelial junctions, marked by MLT-4::RFP (Fig 1B).

We then assessed which pieces of APE-1 are needed for function. Loss of APE-1

suppresses jowls – overt cyst-like bulges near the head of the animal caused by

hyperactivation of the Inversin homolog, MLT-4 [40]. Our hyperactive alleles of MLT-4

include a C-terminal red fluorescent protein (RFP) tag or a missense mutation (E470K).

The MLT-4::RFP allele also results in reduced body length. Therefore, we can assay for

APE-1 function by quantifying the percentage of animals with jowls or measuring their

body lengths. We found that neither APE-1's N-terminal helix nor the C-terminus is

functional on its own, according to our jowls and body length assays (Fig 1C, S1 Fig).

However, fusing the N-terminal helix to APE-1's C-terminal domains creates a mini-

construct that not only localizes to epithelial junctions (Fig 1B), but is also functional (Fig

1C, S1 Fig). Therefore, APE-1 function likely requires its N-terminal helix to localize the

APE-1 C-terminus, with bound phosphatase, to epithelial junctions. These results

suggest that APE-1 confers subcellular localization to PP1, one of the canonical roles

proposed for phosphatase regulatory subunits.

133   ASPPs promote PP1 activity

PP1 regulatory subunits are also thought to control the activity state of the PP1 catalytic

subunits. We and others have shown that ASPPs bind PP1 through the highly

conserved ASPP C-terminus [20,24,41]. While we proposed that the ASPP C-terminus

is required for modulating PP1, it was unclear if this modulation promoted or inhibited

PP1 activity [25].

We hypothesized that if ASPPs stimulate PP1 activity, then deletion of the *C. elegans*

PP1 homolog (GSP-2) would suppress jowls. However, since null alleles of GSP-2 are

lethal, we degraded GSP-2 specifically in the *C. elegans* hyp7 epithelium using the

auxin-inducible degron system (Fig 2A). We found that hyp7-specific degradation of

GSP-2 suppressed jowls in our two mutants (Fig 2A). Therefore, GSP-2 is required in

the epithelium for APE-1 function. These data are consistent with ASPPs promoting

PP1 activity at epithelial junctions.

PP1 is thought to be inactivated by a phosphorylation event at T320 [42,43]. To test if

our model that ASPPs promote PP1 activity is correct, we queried whether ASPPs bind

PP1 that is dephosphorylated (active). We transfected HEK293T cells with a plasmid of

the C-teriminus of iASPP – the most conserved mammalian ASPP homolog [15] – fused

to a HaloTag. We isolated the iASPP C-terminus and any bound PP1 via HaloTag

purification. We found that only dephosphorylated (active) PP1 co-purified with the

iASPP C-terminus (Fig 2B). This result is consistent with ASPPs recruiting active PP1.

Thus, ASPPs specify both the localization and activity state of PP1, consistent with the

two modes of action posited for regulatory subunits [4].

ASPPs multimerize PP1

We previously showed that localizing PP1 to epithelial junctions is not sufficient to

recapitulate ASPP function in *C. elegans* [25]. Instead, the ASPP C-terminus appears to

be required to exert some additional effect on PP1. We had identified missense

mutations in two highly conserved residues (N583 and H591) of the ASPP ankyrin

repeats that each independently render APE-1 inactive [25]. Yet, each mutant is still

able to bind and localize PP1, implying that these residues somehow inactivate a key

aspect of PP1 regulation. One potential model is that these missense mutations disrupt

ASPP selectivity, allowing for the recruitment of phosphorylated (inactive) PP1. To test

this model, we included each homologous missense mutation in the iASPP C-terminus

(N657K and H665Y) and performed HaloTag purifications from HEK293T cells as

described above. We found no change in the phosphorylation state of copurified PP1

with each mutation (Fig 2B). These data suggest the missense mutations alter some

other regulatory function of the ASPPs.

Interestingly, while identifying ASPP interactors using semi-quantitative MudPIT

proteomics, we noted that both APE-1 and the three vertebrate homologs all co-purified

superstoichiometric amounts of PP1 from *C. elegans* (Fig 3A) and HEK293T cell

lysates, respectively [25]. Furthermore, introducing either missense mutation identified

in our genetic screen reduced the amount of GSP-2 copurified with APE-1 (Fig 3A).

Since the residues in the ankyrin repeats that seem to control phosphatase

oligomerization are highly conserved, we asked whether the analogous mutations would

also reduce the stoichiometry of PP1 bound to iASPP, the most conserved vertebrate

homolog. We cotransfected HEK293T cells with plasmids to express PP1 fused to an

HA tag and the iASPP C-terminus fused to a HaloTag and HA tag. We performed

HaloTag purifications and quantitatively blotted for the HA tags on both iASPP and PP1.

We found that the iASPP C-terminus was sufficient to co-purify superstoichiometric

amounts of PP1 (Fig 3B). Introducing either of the homologous mutations identified in

our genetic screen reduced the ratio of PP1 to iASPP (Fig 3B). Therefore, both *Homo*

sapien iASPP and *C. elegans* APE-1 appear to form a complex with superstoichiometric

amounts of PP1 that depends on two key residues in the ASPP ankyrin repeats.

To approximate the size of the ASPP:PP1 complex, we solubilized HEK293T cells co-

transfected as described above and covalently labeled the HaloTagged iASPP C-

terminal constructs with the fluorescent dye JFX549. We performed fluorescent size

exclusion chromatography (fSEC) on these labeled samples by monitoring for JFX549

fluorescence in the elutions. We compared the elution times of fluorescent ASPP:PP1

complexes with protein standards of known molecular weights. The wild type complex

eluted earlier than the 160 kDa standard, (Fig 3C) – consistent with the iASPP C-

terminus complexing with superstoichiometric amounts of PP1. Introducing one of the

homologous missense mutations, N657K, into the iASPP C-terminus shifted the

complex elution time to between the 75 kDa and 160 kDa standards (Fig 3C) –

consistent with a reduction in the amount of PP1 bound to iASPP. A 1:1 ratio would be

~100 kDa. Interestingly, introducing the second homologous missense mutation,

H665Y, into the iASPP C-terminus did not overtly shift the complex's size (Fig 3C).

These data suggest the H665Y mutation is weaker at disrupting the ASPP:PP1 complex

than the N657K mutation. Indeed, the H665Y missense mutation reduces the amount of

PP1 in the ASPP:PP1 complex to a lesser extent than the N657K missense mutation in

both the semi-quantitative proteomics (Fig 3A) and quantitative blotting (Fig 3B).

Together, these data suggest ASPPs multimerize PP1 and that the ASPP ankyrin

repeats assist in phosphatase oligomerization.

210   Single-molecule analysis shows superstoichiometric

211   ASPP:PP1 complexes

We next evaluated ASPP:PP1 complexes at the single molecule level using Total

Internal Reflection Fluorescence (TIRF) microscopy (Fig 4A). We isolated endogenous

APE-1, tagged with 3xflag and GFP, from *C. elegans* whole-worm lysates using

coverslips functionalized with anti-flag antibody. These animals also had endogenously

HaloTagged GSP-2 labeled with the far-red dye JF646. We identified spots where APE-

1 (green) colocalized with GSP-2 (far-red). We reasoned that if ASPP can bind more

than one PP1, then, under continuous excitation, green spots exhibiting one

photobleaching event should colocalize with far-red spots exhibiting multiple

photobleaching events. We trained two Convolutional Neural Networks (CNNs) to

classify stepwise photobleaching events in both green and far-red fluorescence intensity

traces. Our "CNN-Green" and "CNN-FarRed" showed mean validation accuracies of

85.6 +/- 1.2% and 73.2 +/- 1.0%, respectively, during 5-fold cross validations with our

pre-labeled training data (S4 Fig, S1 and S2 Files). Using our trained CNNs, we found

that 39% of ASPP:PP1 complexes, which contained a single APE-1, colocalized with

multiple GSP-2 molecules (Fig 4B). Introducing the N583K missense mutation into APE-

1 resulted in a reduction in the amount of complexes containing oligomeric GSP-2 (Fig

4B, p = 0.008). The H591Y missense mutation in APE-1 did not significantly alter the

stoichiometry of the complex, which is similar to our other *in vitro* assays where this

mutation had less of an effect on the ASPP:PP1 complex. In conclusion, our single

molecule analysis of the ASPP:PP1 complex supports the model that ASPPs can bind

multiple PP1 catalytic subunits and that the ASPP ankyrin repeats promote this

oligomerization.

Phosphatase oligomers restore function to ASPP missense

mutants *in vivo*

If a key function of the ASPP C-terminus is to oligomerize PP1, we hypothesized that

forcing GSP-2 oligomerization should restore function to the APE-1 proteins containing

the ankyrin repeat missense mutations (Fig 5A). To test this hypothesis, we fused extra-

oligomerization domains (EODs) to the N-terminus of endogenous GSP-2 in *C. elegans*

using CRISPR. EODs are domains from a variety of proteins that have been shown to

favor homo-oligomers of various sizes *in vivo* [44]. We selected EODs that would favor

a dimeric, trimeric, tetrameric, or hexameric GSP-2. Fusing the dimeric EOD to GSP-2

was not sufficient to restore jowls, but did reduce body length suggesting APE-1

function was not fully recapitulated (Fig 5B, S5 Fig). Instead, fusing a trimeric or higher-

order EOD to GSP-2 was sufficient to reconstitute APE-1 function in the presence of

either missense mutation, as indicated by both jowls and body length assays (Fig 5B,

S5 Fig). As expected, GSP-2 oligomers failed to bypass the complete deletion of APE-1

(Fig 5C), since the regulatory subunit is also required to recruit and activate the GSP-2

complex at epithelial junctions (Figs 1 and 2). These *in vivo* data complement the *in*

vitro data to suggest that, in addition to localizing active PP1, the core function of

ASPPs is to multimerize PP1.

Discussion

Our study refines and expands the model for ASPP regulation of PP1. We found that

the *C. elegans* ASPP contains a presumptive N-terminal helix that specifies localization

to epithelial junctions (Fig 1) and that ASPPs likely promote PP1 activity (Fig 2).

Interestingly, we also discovered that ASPPs oligomerize PP1 via their ankyrin repeats

(Figs 3 and 4) and that we can bypass inactivating mutations in this conserved region

by engineering PP1 oligomers in *C. elegans* (Fig 5). Altogether, our data are consistent

with a new model whereby ASPPs cluster PP1 activity at epithelial junctions.

AlphaFold predicts that the vertebrate ASPPs also contain N-terminal helices (S1 Fig),

yet whether the vertebrate helices specify subcellular localization or are required for

function is unclear. Some studies have shown that regions of the ASPP N-termini bind

proteins such as desmoplakin [26] and PAR-3 [29,30,38]. These proteins may act as

molecular anchors allowing the vertebrate ASPP:PP1 complex to localize to epithelial

junctions. We were unable to localize GFP to epithelial cell junctions in *C. elegans* using

the predicted vertebrate helices. One possibility is that the binding partners required to

localize the vertebrate helices are not present in the worm hypodermis. Alternatively,

the binding partners are present, but the interacting residues are not conserved – the N-

terminal regions of the ASPPs are poorly conserved. Future studies will be required to

further clarify the molecular mechanisms of ASPP localization.

Our data are consistent with ASPPs promoting PP1 activity – epithelial GSP-2 is

required for APE-1 function (Fig 2A). However, it is formally possible that the

phosphatase is required for a non-enzymatic function. Perhaps oligomerized

phosphatase provides a structural feature required for junctional organization that is

independent of enzymatic activity. We also showed the iASPP C-terminus specifically

co-purifies PP1 that is dephosphorylated at T320 (Fig 2B), which has been considered

to be an active form of PP1 [42,43]. However, to validate the model that ASPPs

promote PP1 activity, a substrate at epithelial junctions needs to be identified. If ASPPs

promote PP1 activity, then we predict that loss of ASPP will result in the accumulation of

one or more phosphorylated targets in affected tissues. Identification of phosphorylated

proteins enriched in ASPP mutants may reveal candidate substrates for the ASPP:PP1

complex and lead to potential therapeutic interventions for the cardiocutaneous disease

using specific kinase inhibitors.

Our *in vitro* (Figs 3 and 4) and *in vivo* (Fig 5) data suggest that ASPPs oligomerize PP1,

yet precisely how ASPPs induce PP1 oligomerization and the molecular arrangement of

the complex remains unclear. Previously, crystal structures of ASPPs bound to PP1

showed a 1:1 complex [24,45]. These structures could represent an initial seeding

intermediate or regulatory state distinct from what we observe. The superstoichiometric

ASPP:PP1 complex might consist of an ordered array of PP1. Indeed, a *Drosophila*

protein phosphatase named Herzog homo-oligomerizes into fibrillar structures via its

prion-like domain [46]. The ASPP:PP1 complex relies on a slightly different mechanism

whereby the independent regulatory subunit, ASPP, somehow clusters many PP1

molecules into a superstoichiometric complex. Alternative structural approaches such

as cryo-electron microscopy might reveal these higher order states whereas

crystallization conditions may favor the heterodimeric state. The ASPPs could also

induce phase separation of PP1, in which case a structure may be difficult to resolve.

Our *in vitro* assays report different stoichiometries for the ASPP:PP1 complex. Perhaps

the ASPP:PP1 stoichiometry is highly sensitive to the method of purification, which

varies between our assays. It is also possible that there is inherent variability in

ASPP:PP1 stoichiometry. Perhaps subunit stoichiometry changes depending on the

tissue in which the complex is expressed or the developmental state of the organism.

We used whole-worm lysates of unsynchronized populations in our MudPIT proteomics

and single molecule analysis of the worm ASPP:PP1 complex. Therefore, these

analyses lack tissue specificity and resolution in developmental timing. Similarly, our

biochemical assays using HEK293T cells capture a snapshot of ASPP:PP1 behavior in

a single cell type. Nonetheless, all *in vitro* assays point to a shared conclusion – ASPPs

can oligomerize PP1. Our assays also consistently show that including either of two

semi-dominant suppressor mutations in the ASPP ankyrin repeats reduces PP1

oligomerization, with one mutation (*C. elegans* N583K/*H. sapien* N657K) having a

greater effect than the other (*C. elegans* H591Y/*H. sapien* H665Y). These data suggest

that even a slight decrease in PP1 oligomerization reduces ASPP function. Our *in vivo*

data complement our *in vitro* assays in that fusing a trimeric or higher-order EOD to

GSP-2 restores APE-1 function in these mutants, while fusing a dimeric EOD to GSP-2

is less effective (Fig 5B, S5 Fig). Our data are consistent with the model that APE-1

induces GSP-2 oligomerization beyond some threshold to elicit a functional output. It

seems likely there are active forms of the ASPP:PP1 complex that remain to be

determined.

The purpose of ASPP-induced PP1 oligomerization is unclear. In the case of Herzog,

oligomerization is thought to enhance the phosphatase's enzymatic activity [46]. It is

possible that ASPP-induced PP1 oligomerization also enhances PP1 activity. Perhaps a

target substrate is highly concentrated at epithelial junctions and requires substantial

dephosphorylation to reach a desired phospho-state. However, an alternative model

that remains consistent with our data is that oligomerized PP1 may function in a non-

enzymatic, structural capacity. While the functional significance of the ASPP:PP1

complex remains an open question, our discovery of PP1 oligomerization by ASPPs

may represent a general mechanism of PP1 regulation.

Materials and methods

Worm strains, maintenance, and CRISPR-Cas9 transgenics

Strains were maintained at room temperature on 60 mm petri dishes containing

nematode growth media (NGM) seeded with a bacterial food source (OP50). In Fig 2A,

strains were also maintained on petri dishes as described above with 1 mM auxin.

CRISPR-Cas9 edits were generated as described in [47] with modifications described in

[25]. Alleles were verified by Sanger sequencing of a PCR amplicon of the modified

locus. A complete list of components of the RNP complexes, repair strategies, and

resulting strains and alleles can be found in S3 File.

Structural prediction and molecular visualization

346 Structural predictions of *C. elegans* APE-1 (AF-Q9XVN3-F1-v4) and the *Homo sapien*

347 ASPP1 (AF-Q96KQ4-F1-v4), ASPP2 (AF-Q13625-F1-v4), and iASPP (AF-Q8WUF5-F1-

348 v4) were acquired from the AlphaFold repository [48,49]. Molecules were visualized in

349 ChimeraX [50,51].

DIC and live fluorescent imaging

For DIC imaging, adult worms were mounted on 2% agarose pads in M9 buffer

containing 20 mM sodium azide on glass slides. Worms were imaged at room

temperature on a Nikon Eclipse Ti-S microscope equipped with a Nikon DS-Qi2 camera

using a 40x DIC objective (Nikon, NA = 0.75) within 20 minutes of mounting and overlay

of No. 1.5 glass coverslip. Images were taken using the NIS-Elements D 5.41.00

software and processed in Fiji [52].

For live fluorescent imaging, adult animals were mounted on 8% agarose pads in 3 µL

1x PBS containing 1.3% (w/v) 0.10 µm Polybead® Microspheres (Polysciences Inc.,

00876) on glass slides. We imaged red and green channels using a custom-built RM21

TIRF microscope (MadCity Labs) with a 60x oil immersion objective (Nikon, NA = 1.49),

488 and 552 nm lasers (Coherent OBIS), and an Orca Fusion BT sCMOS camera

(Hamamatsu). Images were collected at room temperature using Micro-Manager 2.0.

MLT-4::tagRFP::HA was used to identify the focal plane for the hyp7 and seam cell

junction. Images were then collected in the red and green channels. For image analysis,

the MLT-4 signal was used as a marker for apical junctions. A segmented line (5 pixels

wide) was traced along the MLT-4 signal as a region of interest (ROI) in Fiji. The ROI

was duplicated and moved into the cytosolic region of the hyp7 cell syncytium. Both

ROIs were overlaid onto the images collected in the green channel. The mean intensity

along each ROI was quantified and the ratio of junctional to cytosolic mean intensities

calculated.

Body Length Assays

375   Body length assays were conducted as in [25]. Briefly, adult animals were mounted on

376   2% agarose pads in M9 buffer containing 20% (v/v) sodium azide on glass slides.

377   Samples were imaged at room temperature using the BZ-X810 Keyence microscope

378   system with a 10x objective (Nikon, NA = 0.45). Body length was measured by drawing

379   a line through the middle of the worm from anterior to posterior in Fiji.

Jowls assays

In Fig 1, two plates of synchronized worms for each strain were scored for the presence

of jowls using a dissection microscope (Nikon SMZ800N). In Fig 2A, 30 adult worms

were scored for each condition except GUN1663 with no auxin, which had 20 worms

scored. In Fig 5, 40 L4 larvae were scored for each strain. Tagging the endogenous

GSP-2 with trimeric, tetrameric, or hexameric EODs made adult animals unhealthy and

more difficult to score phenotypically. Therefore, phenotypic scoring was conducted on

L4 larvae when jowls were easily observable.

Whole Worm Immunoprecipitations

Worm samples were prepared as described in [25]. Briefly, worms were grown at room

temperature in an OP50 *E. coli* culture mixed with chicken eggs. Worms were collected

and washed in ice-cold H150 buffer (50 mM HEPES pH 7.6 and 150 mM KCl). Worms

were resuspended in 25 mM HEPES pH 7.6, 75 mM KCl, and 5% glycerol with protease

inhibitors (Roche, A32965, 1 tablet per 10 ml lysis buffer) and frozen in liquid nitrogen as

small pellets. Frozen worm samples were lysed using mortar and pestle pre-chilled with

liquid nitrogen until no intact worms were visible under a dissection microscope. Lysates

were diluted 5-fold with 50 mM HEPES pH 7.6, 150 mM KCl, 10% glycerol, 0.005%

IGEPAL (CA-630, Sigma) and clarified via centrifugation at 19,650 × g for 20 min at 4°C.

Clarified lysates were incubated with equilibrated M2 magnetic FLAG beads (Sigma-

Aldrich, M8823) (100 µl of 50% slurry for every ~2.5 g starting worm pellet) overnight at

4°C. Beads were washed twice with 50 mM HEPES pH 7.6, 150 mM KCl, 10% glycerol

and once with TBS (Tris pH 7.6 and 150 mM NaCl). Protein samples were eluted with

150 ng/µl 3xFLAG peptide (Sigma-Aldrich, F4799) in TBS for 30 min at 4°C with

rotation.

MudPIT analysis

Samples were prepared for MudPIT proteomics as described in [25]. Briefly, the Tris

concentration of samples was adjusted to 100 mM, tris(2-carboxyethyl)phosphine

(TCEP-HCl) (Pierce) was added to 10 mM, and samples were incubated at 55°C for 1

hr. 2-chloroacetamide (Sigma-Aldrich) was added to 25 mM and samples incubated for

30 min at room temperature, protected from light. Six volumes of pre-chilled acetone

were added and precipitation proceeded overnight at -20 °C. Samples were then

centrifuged at 8000 x g for 10 min at 4°C. The acetone was aspirated and protein pellets

dried. Subsequent protein preparation and downstream analysis were performed

exactly as in [25]. Raw data and search results will be deposited to the Proteome

Xchange via the MassIVE repository at the time of publication. Mass Spectrometry data

will also be accessible at the Stowers Original Data Repository at the time of

publication.

Tissue Culture Maintenance, Plasmid Generation, and

Pulldowns

HEK293T cells were cultured at 37 °C with 5% CO_2 in 150 mm petri dishes with

mammalian growth media (DMEM, 10% Fetal Bovine Serum, 100 U/mL Penicillin, 100

µg/mL Streptomycin, 10 mM HEPES).

Mammalian vectors expressing HaloTag and HA-tagged proteins were generated via

Gibson cloning [53]. The control plasmid (pDTW117) contains a linker sequence

followed by an HA tag, second linker, TEV cleavage site, and a HaloTag. cDNAs of the

H. sapien wild type iASPP (accession number: NP_001135974.1) C-terminus

(pDTW123) and mutant iASPP C-termini (pDTW124, and pDTW125) were then cloned

into the control plasmid upstream of the initial linker sequence as described in [25]. The

HA-tagged PP1 plasmid (pDTW131) was constructed via a one-piece Gibson assembly

using an amplicon that was obtained from a plasmid containing *M. musculus* PPP1CA

(accession number: NP_114074.1) with oDTW457 and oDTW458. HA-tagged iASPP C-

terminus (pDTW231) was constructed via a one-piece Gibson assembly using an

amplicon that was obtained from the HaloTagged iASPP C-terminus (pDTW81) with

oDTW1113 and oDTW1114. The HA-tagged iASPP C-terminus containing the N657K

missense mutation (pDTW232) was made via a one-piece Gibson assembly using an

amplicon obtained from the N657K mutant iASPP C-terminus plasmid (pDTW94) with

oDTW1113 and oDTW1114. The HA-tagged iASPP C-terminus containing the H665Y

missense mutation (pDTW233) was constructed via a one-piece Gibson assembly using

an amplicon that was obtained from the H665Y mutant iASPP C-terminus plasmid

(pDTW87) with oDTW1113 and oDTW1114.

HEK293T cell transfections and pulldowns were conducted as described in [25]. Briefly,

150 mm dishes of HEK293T cells (~65% confluent) were transfected with a total of

10 µg of plasmid using 60 µg of linear Polyethylenimine 25 kDa (23966, Polysciences

Inc.) mixed in 2 ml Opti-MEM (31985070, ThermoFisher). For cotransfection

experiments, 5 µg of each plasmid was cotransfected for a total of 10 µg. Approximately

24 hr post-transfection, cells were pelleted and frozen at −80°C. Cell pellets were lysed,

clarified, and bound to HaloLink Magnetic Beads (Promega) according to the

manufacturer's instructions. Magnetic beads were washed in TBS with 0.09% IGEPAL

(CA-630, Sigma). Samples were cleaved (2 h at 21°C with 1,100 RPM shaking) with

AcTEV protease (20 units in 100 µl, Invitrogen) to release baits and any bound proteins.

Western Blotting

All SDS-PAGE experiments were run on precast polyacrylamide gels (4-12% Bis-Tris,

Invitrogen). Samples were denatured at 95°C for 1 min in 1x Bolt™ LDS Sample Buffer

(ThermoFisher Scientific; B0007) containing 50 mM dithiothreitol. Protein was

transferred to PVDF membranes (MilliporeSigma; IPFL85R) using the Pierce Power

Blot Cassette system (ThermoFisher Scientific). Blocking and antibody incubation were

conducted in Intercept® (TBS) Blocking Buffer (Fig 2B, LICOR; 927-60001) or EveryBlot

Blocking Buffer (Fig 3B, BioRad, 12010020). Blots were washed using TBS containing

0.1% Tween-20 (v/v) and imaged using the Bio-Rad ChemiDoc™ MP Imaging System.

Band intensities were quantified using the accompanying Image Lab software.

Primary antibodies and dilutions included anti-PP1 (1:1,000; Santa Cruz; Clone E-9; sc-

7482), anti-PPP1CA (phospho T320) (1:1,000; Abcam; Clone EP1512Y; ab62334), and

anti-HA::Peroxidase (1:500; Roche; clone 3F10; 1201381900). Secondary antibodies

and dilutions included the goat anti-mouse IRDye® 800CW (1:20,000; LICOR; 926-

32210) and the

goat anti-mouse IgG (H + L)-HRP Conjugate (1:20,000; Bio-Rad;1706516).

Chemiluminescent images were taken following development using the SuperSignal®

West Dura Extended Duration Substrate (Thermo Scientific; 34075).

Fluorescent size exclusion chromatography

HEK293T cells were transfected and collected as described in [25]. Cell pellets were

thawed in 300 µL Mammalian Lysis Buffer (PAG9381, Promega) with 6 µL 50x Protease

Inhibitor Cocktail (G6521, Promega) and sheared using a 25 gauge needle. JFX549

(gifted from Luke Lavis, Janelia Research Campus) was resuspended to 200 µM in

DMSO and then diluted 1:40 in TBS. Lysates were labeled for 1 hr on ice in the dark

with 1 µL of diluted JFX549. 1 mg/mL ovalbumin, 1 mg/mL aldolase, and 1 mg/mL

conalbumin were prepared from Cytiva's Gel Filtration High Molecular Weight kit

(28403842, Cytiva). All samples were clarified by centrifugation at 17,000 x g for 5 min

at 4°C. Clarified samples were further spun in a Beckman Optima TLX Ultracentrifuge at

200,312 x g for 23 min at 4°C. 100 µL of the resulting supernatants was injected on a

Superdex™ 200 Increase 10/300 GL column using the Shimadzu SIL-20AC HT

Prominence Autosampler. Samples were eluted in filter-sterilized (0.22 µm) and

degassed TBS for 75 min at 4°C at a rate of 0.5 mL/min using the Shimadzu LC-20AD

Prominence Liquid Chromatograph. Fluorescence in elutions was monitored using the

Shimadzu RF-20A XS Prominence Fluorescence Detector. General protein was

detected using Ex = 288 nm, Em = 350 nm. JFX549 labeled protein was detected using

Ex = 549 nm, Em = 571 nm.

Single Molecule TIRF Microscopy

Two NGM plates per worm strain were maintained daily at 15°C. The night before an

imaging experiment, 200 µL of M9 buffer containing 1.5 µM JF649 (GA1121, Promega)

was added dropwise to each plate. Plates of stained worms were wrapped in aluminum

foil and stored upright at 15°C overnight. Worms were washed off of each plate using 1

mL 50 mM HEPES pH 7.6, 150 mM KCl and spun at 100 x g for 1 min at 4°C. Worm

pellets from two plates were transferred to a new tube and spun at 100 x g for 1 min at

4°C. Excess buffer was removed and worm pellets were resuspended in 30 µL of 50

mM HEPES pH 7.6, 150 mM KCl, 10% (v/v) glycerol, Pierce Protease Inhibitor Tablet (1

tablet for every 12.5 mL buffer, ThermoFisher, A32965) and frozen at -80°C. Worms

were thawed on ice and sonicated with a Branson Digital Sonifier using a cup-horn

adaptor with 70% power for ~10 min with 3 sec on/off at 4°C. Sonication was repeated

until intact worms were no longer visible under a dissection microscope. 30 µL of wash

buffer (50 mM HEPES pH 7.6, 150 mM KCl, 10% (v/v) glycerol) was added to each

sample before centrifugation at 17,000 x g for 5 min at 4°C. The supernatant was

collected and loaded into single molecule multi-chamber devices for imaging on a TIRF

microscope (described below).

24 mm x 50 mm No. 1.5 glass coverslips (Corning) were marked on one corner with a

diamond stylist and placed in a glass holder. Coverslips were treated for 20 min in a

Harrick Plasma Plasma Cleaner (PDC-001) on the "high" setting with a pressure

between 0.4-0.8 Torr, established with a IDP-3 Dry Scroll Pump and Harrick Plasma

Vacuum Gauge (PDC-VCG). Cleaned coverslips were then PEGylated by applying 45

µL of a 1:100 mixture of 1% biotin-PEG-silane in ethanol (Laysan Bio Biotin-PEG-SIL-

2K-1g) and PEG-silane (85%, VWR 77035-498) between two coverslips such that their

marked sides are in contact with the PEGylation solution. Coverslips incubated at room

temperature in the dark for 1 hr. Coverslips were washed thoroughly in RO water and

dried with nitrogen gas before storage at room temperature in the dark in a glass

Tupperware container with Drierite desiccant (VWR) until use.

Multi-chamber devices were prepared as in [40]. Briefly, multi-chamber devices of ~30

µL volumes were fashioned by adhering PEGylated coverslips to glass slides using ~4

mm strips of double-sided adhesive tape orthogonal to the long axis. Chambers were

equilibrated with two volumes of wash buffer by capillary action. Chambers were

washed twice after each of the following steps: 10 min incubation with 0.2 mg/mL

neutravidin, 10 min incubation with biotinylated flag antibody (F9291, Sigma Aldrich)

diluted 1:100 in wash buffer, and 10 min incubation with blocking buffer (50 mM HEPES

pH 7.6, 150 mM KCl, 10% (v/v) glycerol, 20 mg/mL BSA). 30 µL of clarified whole worm

lysates (as described above) were loaded into chambers and incubated at room

temperature for 60-80 min in a dark humidified container prior to imaging. Immediately

before imaging, the sample chamber was washed once with 30 µL of wash buffer.

Samples were imaged on a custom-built TIRF microscope (as described above) with

the addition of a Zero-order quarter-wave plate (WPQ10M-633, ThorLabs) to circularly

polarize the 647 nm laser (Coherent OBIS). Samples were imaged continuously at 20

frames/sec for ~50 sec while simultaneously exciting with 488 nm (~4 mW) and 647 nm

(~21 mW) lasers for a total of 1000 frames/channel. Laser powers were tuned such that

the bleach half-time of our JF646 labeled HaloTag::GSP-2 was ~1/5 of our total imaging

time and our APE-1::GFP::3xflag was ~1/10 of our total imaging time. Bleach half-times

were calculated using a half-time calculation software at https://github.com/dickinson-

lab/SiMPull-Analysis-

Software/blob/master/Static_Analysis/Visualization/bleachHalfTime.m [54]. Each single

molecule imaging session began by acquiring a registration image, in which 0.1 µm

TetraSpeck fluorescent microsphere beads (T7279, ThermoFisher) were imaged in

green and far-red channels, to allow for proper colocalization analysis in subsequent

steps.

Training data for our CNNs was collected across 50 technical replicates for each of two

biological replicates of the strain expressing APE-1::GFP::3xflag; HaloTag::GSP. We

used a modified version of the analyze_batch.m function at

https://github.com/dickinson-lab/SiMPull-Analysis-Software/blob/master/Static_Analysis

[54], in which the automated step-counting feature is disabled and the "maxSpots"

variable is set equal to "imgArea / 3e5." We also modified the coloc_spot.m utility

function by setting the "colocDistance" variable equal to 2 pixels. These modified

functions were used to identify and catalog colocalized green and far-red spots along

with their corresponding fluorescence intensity traces.

We then employed custom software to randomly display either green or far-red Z-score

normalized fluorescence intensity traces from colocalized spots. Traces were manually

labeled based on the number of discrete photobleaching events – defined as stepwise,

sustained (>25 counts) decreases in fluorescence. Traces were labeled as having "at

least x" steps, where x is the value of photobleaching events across the entire trace or

until early stopping. Counting stopped early when the intensity trace increased by a

step.

Far-red traces were binned into four classes: rejected, 1-step, 2-step, or 3-and-higher-

step. Far-red traces were labeled as rejected if no steps were counted or if the signal

variability prevented visualization of discrete drops in fluorescence. We combined 3-

step with higher steps traces into one class to simplify CNN-training. Green traces were

binned into two classes: rejected or 1-step. Green traces were labeled as rejected if no

steps were counted, if the signal variability prevented visualization of discrete drops in

fluorescence, or if there was more than 1 photobleaching event in the trace.

In total, 30,135 far-red traces from both biological replicates were labeled: 12,563

rejected, 10,926 1-step, 5,053 2-step, and 1,593 3-and-higher-step. 13,773 green traces

from one biological replicate were labeled: 4,329 rejected and 9,444 1-step. To account

for class imbalances, we applied Inverse Class Frequency Weighting, which assigns

greater weight to underrepresented classes when calculating loss values during

training.

CNN training for each fluorescent channel began with an exploratory phase in which we

tested 39 distinct model architectures. Each model architecture tested different kernel

sizes, number of 1D filters, or the configuration of fully connected (dense) layers. For

each model architecture, we tested various learning rates and batch sizes to optimize

the model training protocol. Each configuration used a stratified, randomized 80/20

train-validation split of the respective training dataset. We selected the top five

configurations based on validation accuracy. Final configuration selection for each

channel was made using 5-fold cross validation, evaluating average accuracy and F1

scores across folds. Then, the final CNN-FarRed and CNN-Green models were trained

on 100% of their respective labeled datasets using the selected architecture and

hyperparameter configuration.

9-10 biological replicates, each consisting of 7 technical replicates, were collected for

each experimental sample. Colocalized far-red and green spots were identified using

the modified analyze_batch.m function described above. We used custom Python

scripts to perform the following actions: 1) extract and Z-score normalize far-red traces

from colocalized spots; 2) use the CNN-FarRed to assign far-red traces into appropriate

classes; 3) extract and Z-score normalize green traces from colocalized spots; 4) use

the CNN-Green to assign green traces into appropriate classes; 5) filter data such that

only cases where a non-rejected green trace accompanied by its respective non-

rejected far-red trace remain for final analysis. The total number of traces for each class

of far-red traces was summed. The percentage of traces in each class for each replicate

was calculated.

Statistical Analysis

Localization data of APE-1 to epithelial junctions was analyzed in GraphPad Prism

using one-way ANOVA with Tukey's adjustment for multiple comparison. Jowls assays

were analyzed in RStudio as previously described in [25]. Briefly, a generalized linear

model was fit to a binomial distribution via RStudio for jowls assays. Body length assays

were analyzed using one-way ANOVAs with Tukey's adjustment for multiple

comparisons. Each CNN was evaluated using a custom python script to perform 5-fold

cross validation. Single molecule photobleaching data were analyzed in GraphPad

Prism using a nested one-way ANOVA with Tukey's adjustment for multiple

comparisons.

Acknowledgements

We thank the labs of Toshi Kawate, Maurine Linder, Carrie Adler, Carolyn Sevier, Richa

Sardana, Josh Chappie, and Natasza Kurpios for sharing space and reagents. We

thank Scott D. Emr and Chris Fromme for valuable advice throughout the project. For

technical advice and assistance, we thank Wendy Greentree (tissue culture) and

Jacqueline Ehrlich (fSEC). We thank Dan J. Dickinson, Eric Drier, and Jeff Lange for

guidance and advice in our single molecule TIRF microscopy. We also thank Dan J.

Dickinson for sharing plasmids containing the EODs. We thank Luke Lavis for sharing

various fluorescent dyes. We thank Matt Thomas at the Cornell Statistical Consulting

Unit for advice on statistical analyses. ChimeraX was developed by the Resource for

Biocomputing, Visualization, and Informatics at the University of California, San

Francisco, with support from National Institutes of Health R01-GM129325 and the Office

of Cyber Infrastructure and Computational Biology, National Institute of Allergy and

Infectious Diseases.

References

1.   Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. The protein kinase

complement of the human genome. Science. 2002;298: 1912–1934.

doi:10.1126/science.1075762

2.   Chen MJ, Dixon JE, Manning G. Genomics and evolution of protein phosphatases.

Sci Signal. 2017;10. doi:10.1126/scisignal.aag1796

3.   Hendrickx A, Beullens M, Ceulemans H, Den Abt T, Van Eynde A, Nicolaescu E, et

al. Docking motif-guided mapping of the interactome of protein phosphatase-1.

Chem Biol. 2009;16: 365–371. doi:10.1016/j.chembiol.2009.02.012

4.   Verbinnen I, Ferreira M, Bollen M. Biogenesis and activity regulation of protein

phosphatase 1. Biochem Soc Trans. 2017;45: 89–99. doi:10.1042/BST20160154

5.   Hubbard MJ, Cohen P. Regulation of protein phosphatase-1G from rabbit skeletal

muscle. 2. Catalytic subunit translocation is a mechanism for reversible inhibition of

activity toward glycogen-bound substrates. Eur J Biochem. 1989;186: 711–716.

doi:10.1111/j.1432-1033.1989.tb15264.x

6.   Ragusa MJ, Dancheck B, Critton DA, Nairn AC, Page R, Peti W. Spinophilin directs

protein phosphatase 1 specificity by blocking substrate binding sites. Nat Struct Mol

Biol. 2010;17: 459–464. doi:10.1038/nsmb.1786

7.   Bertolotti A. The split protein phosphatase system. Biochem J. 2018;475: 3707–

3723. doi:10.1042/BCJ20170726

8.   Morrow CJ, McOrist S. Cardiomyopathy associated with a curly hair coat in Poll

Hereford calves in Australia. Vet Rec. 1985;117: 312–313.

doi:10.1136/vr.117.12.312

9.   Whittington RJ, Cook RW. Cardiomyopathy and woolly haircoat syndrome of Poll

Hereford cattle: electrocardiographic findings in affected and unaffected calves.

Aust Vet J. 1988;65: 341–344. doi:10.1111/j.1751-0813.1988.tb14260.x

10. Simpson MA, Cook RW, Solanki P, Patton MA, Dennis JA, Crosby AH. A mutation

in NFkappaB interacting protein 1 causes cardiomyopathy and woolly haircoat

syndrome of Poll Hereford cattle. Anim Genet. 2008;40: 42–46. doi:10.1111/j.1365-

2052.2008.01796.x

11. Herron BJ, Rao C, Liu S, Laprade L, Richardson JA, Olivieri E, et al. A mutation in

NFkB interacting protein 1 results in cardiomyopathy and abnormal skin

development in wa3 mice. Hum Mol Genet. 2005;14: 667–677.

doi:10.1093/hmg/ddi063

12. Toonen J, Liang L, Sidjanin DJ. Waved with open eyelids 2 (woe2) is a novel

spontaneous mouse mutation in the protein phosphatase 1, regulatory (inhibitor)

subunit 13 like (Ppp1r13l) gene. BMC Genet. 2012;13: 76. doi:10.1186/1471-2156-

13-76

13. Falik-Zaccai TC, Barsheshet Y, Mandel H, Segev M, Lorber A, Gelberg S, et al.

Sequence variation in PPP1R13L results in a novel form of cardio-cutaneous

syndrome. EMBO Mol Med. 2017;9: 319–336. doi:10.15252/emmm.201606523

14. Robinson HK, Zaklyazminskaya E, Povolotskaya I, Surikova Y, Mallin L, Armstrong

C, et al. Biallelic variants in PPP1R13L cause paediatric dilated cardiomyopathy.

Clin Genet. 2020;98: 331–340. doi:10.1111/cge.13812

15. Bergamaschi D, Samuels Y, O'Neil NJ, Trigiante G, Crook T, Hsieh J-K, et al.

iASPP oncoprotein is a key inhibitor of p53 conserved from worm to human. Nat

Genet. 2003;33: 162–167. doi:10.1038/ng1070

16. Iwabuchi K, Bartel PL, Li B, Marraccino R, Fields S. Two cellular proteins that bind

to wild-type but not mutant p53. Proc Natl Acad Sci U S A. 1994;91: 6098–6102.

doi:10.1073/pnas.91.13.6098

17. Gorina S, Pavletich NP. Structure of the p53 tumor suppressor bound to the ankyrin

and SH3 domains of 53BP2. Science. 1996;274: 1001–1005.

doi:10.1126/science.274.5289.1001

18. Samuels-Lev Y, O'Connor DJ, Bergamaschi D, Trigiante G, Hsieh JK, Zhong S, et

al. ASPP proteins specifically stimulate the apoptotic function of p53. Mol Cell.

2001;8: 781–794. doi:10.1016/s1097-2765(01)00367-7

19. Dedeić Z, Sutendra G, Hu Y, Chung K, Slee EA, White MJ, et al. Cell autonomous

role of iASPP deficiency in causing cardiocutaneous disorders. Cell Death Differ.

2018;25: 1289–1303. doi:10.1038/s41418-017-0039-6

20. Helps NR, Barker HM, Elledge SJ, Cohen PT. Protein phosphatase 1 interacts with

p53BP2, a protein which binds to the tumour suppressor p53. FEBS Lett. 1995;377:

295–300. doi:10.1016/0014-5793(95)01347-4

21. Tidow H, Veprintsev DB, Freund SMV, Fersht AR. Effects of Oncogenic Mutations

and DNA Response Elements on the Binding of p53 to p53-binding Protein 2

(53BP2) *. J Biol Chem. 2006;281: 32526–32533. doi:10.1074/jbc.M604725200

22. Patel S, George R, Autore F, Fraternali F, Ladbury JE, Nikolova PV. Molecular

interactions of ASPP1 and ASPP2 with the p53 protein family and the apoptotic

promoters PUMA and Bax. Nucleic Acids Res. 2008;36: 5139–5151.

doi:10.1093/nar/gkn490

23. Skene-Arnold TD, Luu HA, Uhrig RG, De Wever V, Nimick M, Maynes J, et al.

Molecular mechanisms underlying the interaction of protein phosphatase-1c with

ASPP proteins. Biochem J. 2013;449: 649–659. doi:10.1042/BJ20120506

24. Bertran MT, Mouilleron S, Zhou Y, Bajaj R, Uliana F, Kumar GS, et al. ASPP

proteins discriminate between PP1 catalytic subunits through their SH3 domain and

the PP1 C-tail. Nat Commun. 2019;10: 771. doi:10.1038/s41467-019-08686-0

25. Beacham GM, Wei DT, Beyrent E, Zhang Y, Zheng J, Camacho MMK, et al. The

Caenorhabditis elegans ASPP homolog APE-1 is a junctional protein phosphatase

1 modulator. Genetics. 2022;222. doi:10.1093/genetics/iyac102

26. Notari M, Hu Y, Sutendra G, Dedeić Z, Lu M, Dupays L, et al. iASPP, a previously

unidentified regulator of desmosomes, prevents arrhythmogenic right ventricular

cardiomyopathy (ARVC)-induced sudden death. Proc Natl Acad Sci U S A.

2015;112: E973–81. doi:10.1073/pnas.1408111112

27. Chikh A, Matin RNH, Senatore V, Hufbauer M, Lavery D, Raimondi C, et al.

iASPP/p63 autoregulatory feedback loop is required for the homeostasis of

stratified epithelia. EMBO J. 2011;30: 4261–4273. doi:10.1038/emboj.2011.302

28.   Notari M, Hu Y, Koch S, Lu M, Ratnayaka I, Zhong S, et al. Inhibitor of apoptosis-

stimulating protein of p53 (iASPP) prevents senescence and is required for

epithelial stratification. Proc Natl Acad Sci U S A. 2011;108: 16645–16650.

doi:10.1073/pnas.1102292108

29.   Sottocornola R, Royer C, Vives V, Tordella L, Zhong S, Wang Y, et al. ASPP2

binds Par-3 and controls the polarity and proliferation of neural progenitors during

CNS development. Dev Cell. 2010;19: 126–137. doi:10.1016/j.devcel.2010.06.003

30.   Cong W, Hirose T, Harita Y, Yamashita A, Mizuno K, Hirano H, et al. ASPP2

regulates epithelial cell polarity through the PAR complex. Curr Biol. 2010;20:

1408–1414. doi:10.1016/j.cub.2010.06.024

31.   Hauri S, Wepf A, van Drogen A, Varjosalo M, Tapon N, Aebersold R, et al.

Interaction proteome of human Hippo signaling: modular control of the co-activator

YAP1. Mol Syst Biol. 2013;9: 713. doi:10.1002/msb.201304750

32.   Royer C, Sandham E, Slee E, Schneider F, Lagerholm CB, Godwin J, et al. ASPP2

maintains the integrity of mechanically stressed pseudostratified epithelia during

morphogenesis. Nat Commun. 2022;13: 941. doi:10.1038/s41467-022-28590-4

33.   Vives V, Su J, Zhong S, Ratnayaka I, Slee E, Goldin R, et al. ASPP2 is a

haploinsufficient tumor suppressor that cooperates with p53 to suppress tumor

growth. Genes Dev. 2006;20: 1262–1267. doi:10.1101/gad.374006

34. Kampa KM, Acoba JD, Chen D, Gay J, Lee H, Beemer K, et al. Apoptosis-

stimulating protein of p53 (ASPP2) heterozygous mice are tumor-prone and have

attenuated cellular damage-response thresholds. Proc Natl Acad Sci U S A.

2009;106: 4390–4395. doi:10.1073/pnas.0809080106

35. Zak J, Vives V, Szumska D, Vernet A, Schneider JE, Miller P, et al. ASPP2

deficiency causes features of 1q41q42 microdeletion syndrome. Cell Death Differ.

2016;23: 1973–1984. doi:10.1038/cdd.2016.76

36. Langton PF, Colombani J, Chan EHY, Wepf A, Gstaiger M, Tapon N. The dASPP-

dRASSF8 complex regulates cell-cell adhesion during Drosophila retinal

morphogenesis. Curr Biol. 2009;19: 1969–1978. doi:10.1016/j.cub.2009.10.027

37. Langton PF, Colombani J, Aerne BL, Tapon N. Drosophila ASPP regulates C-

terminal Src kinase activity. Dev Cell. 2007;13: 773–782.

doi:10.1016/j.devcel.2007.11.005

38. Yamashita K, Mizuno K, Furukawa K, Hirose H, Sakurai N, Masuda-Hirata M, et al.

Phosphorylation and dephosphorylation of Ser852 and Ser889 control the

clustering, localization and function of PAR3. J Cell Sci. 2020;133.

doi:10.1242/jcs.244830

39. Royer C, Koch S, Qin X, Zak J, Buti L, Dudziec E, et al. ASPP2 links the apical

lateral polarity complex to the regulation of YAP activity in epithelial cells. PLoS

One. 2014;9: e111384. doi:10.1371/journal.pone.0111384

40. Beyrent E, Wei DT, Beacham GM, Park S, Zheng J, Paszek MJ, et al. Dimerization

activates the Inversin complex in C. elegans. Mol Biol Cell. 2024;35: ar127.

doi:10.1091/mbc.E24-05-0218

41. Llanos S, Royer C, Lu M, Bergamaschi D, Lee WH, Lu X. Inhibitory member of the

apoptosis-stimulating proteins of the p53 family (iASPP) interacts with protein

phosphatase 1 via a noncanonical binding motif. J Biol Chem. 2011;286: 43039–

43044. doi:10.1074/jbc.M111.270751

42. Dohadwala M, da Cruz e Silva EF, Hall FL, Williams RT, Carbonaro-Hall DA, Nairn

AC, et al. Phosphorylation and inactivation of protein phosphatase 1 by cyclin-

dependent kinases. Proc Natl Acad Sci U S A. 1994;91: 6408–6412.

doi:10.1073/pnas.91.14.6408

43. Yamano H, Ishii K, Yanagida M. Phosphorylation of dis2 protein phosphatase at the

C-terminal cdc2 consensus and its potential role in cell cycle regulation. EMBO J.

1994;13: 5310–5318. doi:10.1002/j.1460-2075.1994.tb06865.x

44. Chang Y, Dickinson DJ. A particle size threshold governs diffusion and segregation

of PAR-3 during cell polarization. Cell Rep. 2022;39: 110652.

doi:10.1016/j.celrep.2022.110652

45. Zhou Y, Millott R, Kim HJ, Peng S, Edwards RA, Skene-Arnold T, et al. Flexible

Tethering of ASPP Proteins Facilitates PP-1c Catalysis. Structure. 2019;27: 1485–

1496.e4. doi:10.1016/j.str.2019.07.012

46. Nil Z, Hervás R, Gerbich T, Leal P, Yu Z, Saraf A, et al. Amyloid-like Assembly

Activates a Phosphatase in the Developing Drosophila Embryo. Cell. 2019;178:

1403–1420.e21. doi:10.1016/j.cell.2019.08.019

47. Ghanta KS, Mello CC. Melting dsDNA donor molecules greatly improves precision

genome editing in Caenorhabditis elegans. Genetics. 2020;216: 643–650.

doi:10.1534/genetics.120.303564

48. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly

accurate protein structure prediction with AlphaFold. Nature. 2021;596: 583–589.

doi:10.1038/s41586-021-03819-2

49. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al.

AlphaFold Protein Structure Database: massively expanding the structural

coverage of protein-sequence space with high-accuracy models. Nucleic Acids

Res. 2022;50: D439–D444. doi:10.1093/nar/gkab1061

50. Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al.

UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein

Sci. 2018;27: 14–25. doi:10.1002/pro.3235

51. Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al. UCSF

ChimeraX: Structure visualization for researchers, educators, and developers.

Protein Sci. 2021;30: 70–82. doi:10.1002/pro.3943

52. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.

Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9:

676–682. doi:10.1038/nmeth.2019

53. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA 3rd, Smith HO.

Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat

Methods. 2009;6: 343–345. doi:10.1038/nmeth.1318

54. Stolpner N, Dickinson DJ. Single-cell single-molecule pull-down (sc-SiMPull) for

detection of protein complexes from embryonic lysates. Methods Mol Biol.

2022;2438: 59–81. doi:10.1007/978-1-0716-2035-9_4

Fig 1. APE-1 function requires its N-terminal helix.

(A) Experimental design. Endogenous APE-1 was truncated and GFP-tagged via

CRISPR. APE-1 constructs were then assessed for localization to epithelial junctions

between the seam and hyp-7 cell syncitia via TIRF microscopy. A region of interest

(ROI) was drawn over cellular junctions using MLT-4::RFP as a marker. This ROI was

duplicated and moved to a nearby hyp-7 cytosolic region. The mean GFP intensity was

then calculated for both ROIs and reported as a ratio of junctional intensity over

cytosolic intensity. APE-1 constructs were also assayed for function via a jowls assay.

(B) Localization assay. White arrowheads indicate seam and hyp-7 junctions. * indicates

autofluorescent alae. Data represent the mean and S.E.M. (black bars) of 10 biological

replicates. Scale bar = 10 μm. NS = not significant. (C) Jowls assay. Red arrow heads

indicate jowls. Data represent the percentage and 95% confidence interval (black and

gray bars) of jowls in populations of adult animals (n = 247, 76, 142, and 164,

respectively). Scale bar = 15 μm.

Fig 2. ASPPs promote PP1 activity.

(A) Skin-specific depletion of GSP-2. Endogenous GSP-2, tagged with an Auxin

Inducible Degron (AID), is ubiquitinated by skin-specific TIR-1 when animals are fed

auxin. Data represent the percentage and 95% confidence interval (black and gray bars,

respectively) of jowls in adult animals (n = 30 each). (B) HEK293T cells were

transfected with plasmids encoding HaloTagged iASPP C-terminal constructs. Input and

elutions were blotted for PP1 phosphorylated at T320, total PP1, and an HA tag on the

iASPP bait.

Fig 3. ASPPs multimerize protein phosphatase 1.

(A) Multi-dimensional Proten Identification Technology (MudPIT) analysis of 3xflag

immunoprecipitations from whole-worm lysates. Data represent the amount of bait and

GSP-2 as a percentage of total protein found in each sample. Ratios of GSP-2 to APE-1

are shown below the table. Note that the control and APE-1 MudPIT data are from [25].

(B) Quantitative HA blot of Halotag purifications from HEK293T cells. Ratios of PP1 to

iASPP C-terminus are shown below the blot. (C) Fluorescent size exclusion

chromatography of JFX549-labeled HEK293T cell lysates. Peaks for molecular weight

standards (described in methods) are represented as vertical black lines with the top

10% of signal shaded in light gray.

Fig 4. Single molecule analysis shows superstoichiometric ASPP:PP1 complexes.

(A) Experimental design for single molecule analysis of APE-1:GSP-2 complex

stoichiometry. Worms with wild type or mutant APE-1 were fed far-red fluorescent dye

(JF646) to label HaloTag::GSP-2. Whole worm lysates were loaded on coverslips

functionalized with anti-flag antibody. Coverslips were imaged on a TIRF microscope in

the green and far-red channels. Fluorescence intensity traces from colocalized green

and far-red spots (yellow circles) were isolated for analysis. A convolutional neural

network (CNN), trained to recognize single-step photobleaching events in green

intensity traces, selected protein complexes containing monomeric APE-1. A second

CNN, trained to recognize single and multi-step photobleaching events in far-red

intensity traces, quantified the number of monomeric APE-1 molecules colocalized with

mulitple GSP-2. Scale bar = 5 µm. (B) Quantification of APE-1:GSP-2 complexes

containing monomeric APE-1 and oligomeric GSP-2. Data represent the mean and

S.E.M. (black bars) of 9-10 biological replicates (outlined spots), each consisting of 7

technical replicates (small spots). (C-E) Example green (top panels) and far-red (bottom

panels) normalized fluorescence intensity (Z-score) traces from colocalized spots.

Fig 5. GSP-2 oligomers bypass APE-1 missense mutations, but not APE-1

deletion.

(A) Extra-oligomerization domains (EODs) favoring different homo-oligomers are

attached to the endogenous GSP-2 of animals where jowls had been suppressed by

mutations in APE-1. Animals are then assessed for the restoration of jowls. (B) Jowls

assay for APE-1 missense mutants. (C) Jowls assay for APE-1 deletion mutant. Data

represent percent jowls and 95% confidence interval (Black and gray bars) in L4

animals (n = 40 each).

Supporting information

S1 Fig. APE-1 function requires its N-terminal helix.

(A) Alphafold structural domain predictions mapped onto the primary sequences of

ASPP1, ASPP2, and iASPP from H. sapiens as well as APE-1 from C. elegans.

Predicted structures of the N-terminal alpha helices are shown in the insets above each

domain map with predicted residues numbers indicated. Represented in different colors

are the beta-grasp domain (teal), N-terminal alpha helix (tan), undefined alpha-helical

regions (brown), ankyrin repeats (purple), and SH3 domain (magenta). (B) Body length

assay. Data represent mean and S.E.M. (black bars) of 23-29 biological replicates.

S4 Fig. Convolutional Neural Network (CNN) confusion matrices.

(A) CNN-Green mean confusion matrix of raw counts averaged across 5-fold cross

validation during training with GFP fluorescence intensity traces. (B) Row-normalized

mean confusion matrix from panel A showing recall for each class during K-fold cross

validation of the CNN-Green. (C) CNN-FarRed mean confusion matrix of raw counts

averaged across 5-fold cross validation during training with far-red fluorescence

intensity traces. (D) Row-normalized mean confusion matrix from panel C showing

recall for each class during K-fold cross validation of the CNN-FarRed.

S5 Fig. GSP-2 oligomers bypass APE-1 missense mutations

(A) Body length assay.Data represent the mean and S.E.M. (black bars) of 27-36

biological replicates. **** indicates $p<0.0001$. * indicates $p<0.05$.

S1 File. Training metrics from CNN-Green K-fold cross validation.

S2 File. Training metrics from CNN-FarRed K-fold cross validation.

S3 File. Strains, alleles, and reagents.

Fig 1

Fig 2

Fig 3

A

Grow worms expressing APE-1::GFP::3xflag

Solubilize & 3xflag IP

	3xflag-tagged bait			
	Control (GFP)	APE-1	APE-1 N583K	APE-1 H591Y
Bait	39.38	0.54	0.22	0.12
GSP-2	0.01	10.79	0.87	0.87
GSP-2:APE-1		20:1	4:1	7:1

B

Co-transfect HA-tagged constructs

Solubilize & Halotag Pulldown

iASPP(C) bait

WT   N657K   H665Y

HA::PP1

HA::iASPP(C)

8:1   2:1   4:1   PP1:ASPP ratio

C

Co-transfect HA-tagged constructs

Solubilize & Label Halotag

PP1
iASPP(C)
HT

1:1	2:1	3:1
MW: 100	137	174

160 kDa   75 kDa   40 kDa

Fluorescence Intensity (a.u.)

Time (min)

— WT   — N657K   — H665Y   — Halotag

Fig 4

A Grow
APE-1::GFP::3xflag
Halotag::GSP-2

JF646 labeling

Bind lysate to functionalized coverslip

HT
GSP-2
APE-1
3xflag
GFP
α-flag

Single molecule photobleaching
of green and far-red channels

APE-1::GFP::3xflag

Halotag::GSP-2

Data analysis

Isolate intensity traces
from colocalized spots

Select monomeric
green spots
(APE-1)

Quantify far-red
photobleaching steps
(GSP-2)

B

APE-1:GSP-2 complexes with oligomeric GSP-2 (%)

p=0.531

p = 0.008

APE-1

APE-1
N583K

APE-1
H591Y

C

Green (APE-1) Normalized fluorescence intensity (Z-score)

1-step

1-step

Far-red (GSP-2)

Time (sec)

D

1-step

2-step

E

1-step

3-step

Fig 5

S4 Fig

S5 Fig