Patient-specific iPSC models of neural tube defects identify underlying deficiencies in neuroepithelial cell shape regulation and differentiation

Ioakeim Ampartzidis[1*], Elliott M Thompson[1], Yashica Gupta[1], Andrea Krstevski[1], Nicola Elvassore[2,3], Eirini Maniou[2,3], Paolo de Coppi[4-8], Gabriel L Galea[1,#]

1. Developmental Biology and Cancer, UCL GOS Institute of Child Health, London, UK

2. Department of Industrial Engineering, University of Padua, Padua, Italy

3. Veneto Institute of Molecular Medicine, Padua, Italy.

4. Stem Cells and Regenerative Medicine, UCL GOS Institute of Child Health, London, UK

5. Department of Development and Regeneration, Woman and Child and UZ Leuven Clinical Department of Obstetrics and Gynaecology, KU Leuven, Leuven, Belgium.

6. Specialist Neonatal and Paediatric Surgery, Great Ormond Street Hospital for Children NHS Foundation Trust, London, UK.

7. Medical and Surgical Department of the Fetus, Newborn and Infant, Ospedale Pediatrico Bambino Gesù, IRCCS, Rome, Italy.

8. NIHR Great Ormond Street Hospital Biomedical Research Centre, London, UK

Corresponding author: g.galea@ucl.ac.uk

* Current address = Department of Biochemistry, University of Cambridge, Hopkins Building, Tennis Court Road, Cambridge CB2 1QW, UK

Key words: Neural tube defects, apical constriction, iPSC, planar cell polarity

Abstract

Spina bifida and anencephaly are neural tube defects caused by failure of embryonic

neural tube closure. Successful closure requires elongation and apical constriction of

neuroepithelial cells. These critical behaviours have not been studied in human, patient-

specific models. We characterise a human iPSC-derived neuroepithelial morphogenesis

model which is highly reproducible between lines. Differentiated neuroepithelial cells

actively undergo ROCK-dependent apical constriction. ROCK acts downstream of planar

cell polarity/VANGL2 in other species. We find that VANGL2 is up-regulated and

phosphorylated during iPSCs-to-neuroepithelial differentiation. The patient-identified

VANGL2-R353C mutation does not alter its expression, localisation or phosphorylation,

but reduces myosin-II phosphorylation and apical constriction relative to congenic control

iPSCs. Non-congenic comparisons and forward genetic testing are also informative in this

reproducible model. We compare lines reprogrammed from amniocytes of two patients

with spina bifida, versus two controls. One patient-derived line forms a morphologically

normal neuroepithelium, but fails to differentiate into neurons. The second fails to undergo

apicobasal elongation, and harbours compound heterozygous mutations in the *MED24*

gene previously implicated in neuroepithelial elongation in mice. Thus, iPSC-derived

neuroepithelial modelling provides mechanistic insights into conserved cell behaviours,

links genetically-impaired apical constriction with a human disease, and establishes

patient-specific models which recapitulate failures of heterogenous neuroepithelial

functions.

Introduction

Neural tube defects (NTDs), such as spina bifida and anencephaly, affect approximately 1:1,000 births globally[1]. These severe neurodevelopmental malformations are caused by abnormal morphogenesis of the neural tube during embryonic development. The most severe NTDs result from failure of the physical process of neural tube closure, which in humans is completed by approximately 30 days gestation[2,3]. Completion of this morphogenetic process during largely inaccessible stages of pregnancy has limited our ability to study it in humans. This challenge is compounded by the multigenic causation of NTDs, as well as interactions of both maternal and embryonic genetics with environmental factors[4-6]. Consequently, few patients receive an interpretable genetic diagnosis. Typically, patient-identified mutations have been functionally validated in transgenic animal models [5,7-9].

Mice and other vertebrates have been invaluable models for understanding the process of neural tube closure. Both mice and humans close their open neural tube regions – called neuropores – through progressive elevation of neural folds which meet at the dorsal midline and zipper along the body axis[2,10]. The progression of closure requires the action of different cell types, particularly the surface ectoderm (future epidermis) and neuroepithelium (future brain and spinal cord)[3]. These cells coordinate multiple behaviours within and between each other: neuroepithelial cells thicken along their apicobasal axis as they differentiate into a pseudostratified epithelium which undergoes interkinetic nuclear migration synchronised with cell cycle progression, superimposed on active apical constriction[11-17].

Neuroepithelial apical constriction is an evolutionarily conserved force-generating mechanism which contributes to neural tube closure in chordates including amphibians, birds and mammals[3]. It requires targeted localisation and phosphorylation of non-muscle myosin II in the cell's apical domain. Myosin binds filamentous (F-)actin, cross-linking filaments and generating contractile mechanical force through power strokes[18]. Neuroepithelial apical enrichment of F-actin and phosphorylation of myosin both require Rho-associated kinase (ROCK)[19], which is in turn apically localised at least in part by signalling through the non-canonical wingless homology (Wnt)/planar cell polarity (PCP)

pathway[20,21]. PCP components promote neuroepithelial apical constriction in *Xenopus*

frogs[22,23], chickens[24] and mice[15,20], but this pathway has been difficult to study in human

epithelial models due to lack of quantifiable readouts relevant to this pathway's

functions *in vivo*.

PCP was originally described in *Drosophila* as a regulator of polarised cell features such

as hair cell bristle orientation[25-27]. Pathway components were later shown to be

conserved in vertebrates and contribute to neural tube closure[28-31]. The core PCP

pathway involves signalling through asymmetric receptor complexes on adjacent cells,

pairing van Gogh-like (VANGL) and Cadherin EGF LAG seven-pass G-type receptor

(CELSR) proteins on one cell, with frizzled (FZD)/CELSR complexes on the adjacent

cell[32]. Both VANGL and FZD branches of this pathway are involved in neural tube

closure[33,34], although the roles of VANGL2 are best understood in mice. VANGL2 can be

activated through phosphorylation following binding of PCP Wnt ligands[35,36].

PCP signalling drives two functionally distinct processes in neuroepithelial cells

essential for mammalian NT closure: early convergent extension and later apical

constriction. Global genetic deletion of *Vangl2* precludes medial convergence and

rostro-caudal elongation necessary to initiate neural tube closure in mouse embryos[37],

although this effect may be partly mediated by the adjacent mesoderm[38]. At later

developmental stages, however, VANGL2 promotes cytoskeletal reorganisation

involving F-actin planar polarisation and phosphorylation of non-muscle myosin IIB,

likely by ROCK[20,39-41]. We previously reported that VANGL2-dependent enhancement of

apical constriction increases apical tension required to elevate the neural folds in the

spinal region of mice[20], and others have implicated it in apical constriction of the anterior

neuroepithelium[22].

Mutations in PCP genes including *VANGL2*[42-47] have been identified in patients who

have NTDs and damaging variants in actomyosin-related cytoskeletal regulation are

enriched in individuals with spina bifida[48]. However, it has been challenging to study the

regulation of apical constriction in human neuroepithelial cell models. A previous study

described elegant modelling of neural tube closure by differentiation of induced

pluripotent stem cells (iPSCs) on micropatterned surfaces[49]. That model's formation of

neuroepithelium enveloped within a larger cyst limits access to its apical surface for
functional measurements, such as inference of mechanical tension through laser
ablation assays. Similarly, previous studies have demonstrated that disruption of
actomyosin regulation dilates apical surfaces on the inside of human iPSC-derived
neuroepithelial spheres[50,51]. Several models have also been described which enable
studies into later stages of neuroepithelial function as they differentiate into neural
progenitors and then neurons[52-55]. While these neurogenesis models provide insights
into later differentiation, they do not capture cell behaviours crucial for neural tube
closure.

We previously characterised a minimal model of neuroepithelial morphogenesis in
which a monolayer of cuboidal iPSCs apicobasally elongate into a complex,
pseudostratified neuroepithelium with a planar morphology[17]. Importantly, this model
presents readily accessible neuroepithelial apical surfaces which are directly
comparable in size with those of mouse embryos[17]. Moreover, the same protocol can be
extended to differentiate neuroepithelial cells into cortical neurons[56]. This enables
sequential assessment of neuroepithelial functions relevant to generating mechanical
deformations which close the neural tube and their subsequent differentiation which
normally occurs after completion of closure. Here, we show that neuroepithelial
development in this model is reproducible, even when comparing non-congenic iPSC
lines. We validate use of the model to study PCP components including ROCK and
VANGL2, finding that a *VANGL2* point mutation previously identified in a patient[46]
diminishes neuroepithelial apical constriction. Extension of the model to two iPSC lines
derived from individuals who have spina bifida identifies line-specific deficiencies
potentially relevant to their malformation. Strikingly, all three NTD-relevant lines –
VANGL2 point mutation and two patient-derived lines – present failures of distinct
neuroepithelial functions, recapitulating the heterogeneity of these conditions.

Results

Lineage-specific marker expression precedes morphological changes during iPSC to neuroepithelium differentiation

Intra-line variability often limits the use of iPSCs to model multigenic diseases, such as NTDs, in which congenic controls are often impractical. In contrast, neuroepithelial differentiation through dual SMAD inhibition (Figure 1a)[17,56] is highly reproducible between different iPSC lines. We illustrate this using three previously reported iPSC lines from healthy individuals, selected to represent different origins and reprogramming methods (Figure 1b). Line HO193b was reprogrammed from male (46XY) amniocytes through microfluidic delivery of modified mRNA[57,58]. SFC086-03-01 was reprogrammed from adult female (46XX) fibroblasts using non-integrating Sendai virus[17,59]. 802-30F was from endothelial cells of a Hispanic female (46XX) reprogrammed through StemRNA[60]. Despite their varied origins, all three undergo equivalent, temporally predictable changes in epithelial morphology during neuroepithelial differentiation (Figure 1a-h).

These cells begin their differentiation as an apicobasally polarised cuboidal monolayer approximately 25 μm thick with large apical areas approximately 60 μm^2 ('day 2', Figure 1c-g). During differentiation, they double their apicobasal length into a pseudostratified epithelium[17] approximately 55 μm thick, with small apical surfaces around 10 μm^2 ('day 8', Figure 1c-g). Apicobasal elongation increases total epithelial volume, accommodating ongoing cell division[17], yet the volume of individual cells decreases from 1,550 μm^3 at the start of differentiation to 500 μm^3 at the end (Figure 1h). This reduction in total cell volume is partly due to a halving of nuclear volume (Supplementary Figure 1).

Neuroepithelial apical size and apicobasal thickness – properties whose dynamic control enables neural tube closure in vivo[20,24,61,62] – are indistinguishable between non-congenic iPSC lines at each stage of differentiation analysed (Figure 1c-g). Equivalent neuroepithelial thickness and apical areas are also obtained in cultures differentiated by different investigators (Supplementary Figure 2). All lines show a non-linear morphological evolution during differentiation: apical area and apicobasal thickness

remain similar between days 2 and 4 of differentiation, but then change rapidly to a

plateau by day 6 of differentiation (Figure 1e-g). In contrast, their expression of

pluripotency markers such as NANOG and OCT4 are rapidly lost between days 2 and 4

of differentiation (Figure 1i,j)[17]. The cadherin CDH1 is similarly down-regulated early

during differentiation, switching to expression of the neuroepithelial CDH2 from day 4 of

differentiation (Figure 1i). The neuroepithelial marker PAX6 is detectably expressed

from day 4 and its expression increases between day 4 and 8 (Figure 1i,k).

This suggests that cells' expression of molecular neuroepithelial markers precedes their

morphological changes in apicobasal elongation and reduction in apical size.

Figure 1: Reproducible morphological and molecular evolution of iPSC differentiation into neuroepithelial cells. iPSC1 is HO193b, iPSC2 is SFC086-03-01 and iPSC3 is 802-30F.
a, Schematic representation of the neuroepithelial induction protocol. Sub-confluent human iPSCs (in grey) are passaged and seeded at full confluency from the start of differentiation and

cultured for eight days, until reaching neuroepithelial maturity. Further passage at this stage
*forms neural rosettes and allows further differentiation into post-mitotic neurons[56]. **b,** Summary*
*of control lines used to validate reproducibility of differentiation into neuroepithelial cells. **c,d,***
*Representative high-resolution confocal top-view (**c**) and cross-section (**d**) images of day 2-8*
cultures fluorescently labelled to show F-actin with inverted gray LUT. P values from one-way
*ANOVA with post-hoc Bonferroni. **e,** Dot plot representing cell's apical area across timepoints of*
neuroepithelial differentiation, in three iPSC lines. Black bars represent the median apical area
*across three culture plates of each line. **f,** Heatmaps of hiPSC2 representing the apical area*
*size changes during differentiation, in a contiguous area of 1,200 μm² per timepoint. **g,***
Quantification of epithelial thickness across neuroepithelial differentiation of control cell lines.
*Three culture replicates were measured for each time point and cell line. **h,** Dot plot*
representing the cell volume decrease across timepoints of differentiation, in control cell lines,
*with representative 3D volumetric images at day 2 and day 8 (**h'**). **i,** Representative western blot*
panel of NANOG, OCT4, CDH1, CDH2, PAX6, and LMNB protein levels across timepoints of
neuroepithelial differentiation, in the iPSC2 line. NANOG expression is normalised to day 2 per
*line, whereas PAX6 is normalised to day 8 as it is not detected at day 2. **k,l,** Dot plots*
*representing normalised band intensity of NANOG (**k**), and PAX6 (**l**), across differentiation for*
three control cell lines. Protein levels across all the cell lines were normalised to day 2. P values
*from one-way ANOVA (**e,g**) or Wilcoxon paired test (**j,k**) with Bonferroni correction (***: p ≤*
0.001, *: p ≤ 0.005). Scale bar = 10 μm.

iPSC-derived neuroepithelial cells actively generate mechanical tension through
apical constriction

While reproducible between cultures, apical areas are variable within individual cultures
(Figure 1e): a hallmark of asynchronous apical constriction *in vivo*[63]. Through live
imaging, we observe that iPSC-derived neuroepithelial cells spontaneously and
dynamically alter their apical area (Figure 2a). Pharmacological inhibition of the myosin-
activating kinase ROCK increases neuroepithelial apical area (Figure 2b-c), as
previously observed in mouse embryos with equivalent ROCK inhibition *in vivo*[11].

The functional outcome of apical constriction is generation of mechanical tension which
is transmitted between adjacent cells and pulls on the surrounding tissue[20,64].
Contractile neuroepithelial apical tension can be inferred through annular laser ablation
assays[11,20], in which release of a small cluster of cells from their neighbours causes
rapid retraction of the cell borders within the ablated ring (Figure 2d). Their tension is

significantly diminished following a short period of ROCK inhibition (Figure 2e),

confirming that it is actively established through apical constriction.

In vivo, the Rho/ROCK/myosin pathway can be activated by PCP/VANGL2

signalling[20,24], and is well-established to promote apical constriction[20,22,65]. We observe

that VANGL2 and its binding partner scribbled (SCRIB) are expressed at low levels from

early stages of neuroepithelial differentiation and are significantly up-regulated by the

end of the differentiation period (Figure 2f,g). Both are co-localised on the cell's apical

cortex (Supplementary Figure 3). Serine-79/82/84 tri-phosphorylation of VANGL2 also

increases significantly during differentiation, with a particularly large up-regulation

between days 4 and 6 (Figure 2h), coinciding with the reduction in apical area (Figure

1e). Thus, iPSC-derived neuroepithelial cells express the cellular machinery required for

physiological control of apical constriction and actively establish apical tension *in vitro*.

*Figure 2: iPSC-derived neuroepithelial cells undergo ROCK-dependent apical constrictions and up-regulate expression of the PCP co-receptor VANGL2. **a,** Live imaging of day 8 neuroepithelial cultures stained with CellMask Actin dye. **b,** Representatives confocal images of neuroepithelial cultures immunolabelled against ROCK1/2. Top view shows the distribution of protein of interest across the apical areas and cross-section annotates protein localisation along the apicobasal axis. Dashed line indicates the basal side. **c,** Representative F-actin-stained images of cultures treated vehicle or 10 μM ROCK inhibitor (Y27632) for 4 hr. **d,** Dot plot representing the median apical area of day 8 SFC086 neuroepithelial cultures. **e,** Live-imaged membrane-stained day 8 neuroepithelial cultures before and after a circular laser ablation, shown in red dash line. Magenta and cyan outlined areas indicate the cell cluster recoil before and after the ablation, respectively. **f,** Median apical cluster displacement of neuroepithelial cells, across three independent replicates of. Each point represents the median displacement quantified in each plate after at least three laser ablations. **g,h,** Representative confocal images of SCRIB and VANGL2 immunolabelled cultures at day 2 and day 8, in of top-view (**g**) and cross-section (**h**). Top view shows protein distribution across the apical areas and cross-section annotates protein's localisation along the apicobasal axis. Dashed lines indicate the basal side. **i,** Representative western blot panel of VANGL2, phosphorylated VANGL2 (pVANGL2), and GAPDH protein levels across timepoints of neuroepithelial differentiation, in cell line. All analyses shown are in iPSC2. P values from Wilcoxon paired test with Bonferroni correction (**: $p \leq 0.01$, *: $p \leq 0.005$). Scale bar = 10 μm.*

An NTD-associated *VANGL2* mutation impairs neuroepithelial apical constriction

To our knowledge, no genetic mutations identified in human patients have previously been functionally associated with failure of apical constriction. The *VANGL2-R353C* mutation (Figure 3a) was initially reported in a fetus which had both anencephaly and spina bifida[46], and impairs other functions of PCP signalling in fruit flies[66]. This point mutation causes a basic to polar amnio acid switch in a C-terminal region (Supplementary Figure 4) which is highly conserved across species and mediates interactions with pathway adaptor proteins[46,66]. We therefore considered it a high priority target for testing in a human system.

We find that *VANGL2-R353C* mutation does not alter overall VANGL2 protein levels, phosphorylation (Figure 3b,c), or apical localisation (Supplementary Figure 5) in two iPSC clonal knock-in lines homozygous for this mutation compared with a congenic control line. Knock-in and congenic control lines retain their ability to differentiate into different germ layers (Supplementary Figure 6), achieve equivalent changes in neuroepithelial apicobasal thickness (Supplementary Figure 7a-c) and nuclear pseudostratification (Supplementary Figure 7d,e) during differentiation. The *VANGL2-*

R353C mutation does not significantly alter cell volume, nor does it diminish extension

of sub-apical protrusions which are characteristic of neuroepithelial cells[67]

(Supplementary Figure 8).

However, *VANGL2-R353C* knock-in lines have significantly larger apical areas (Figure

3d,e) and less apical localisation of phosphorylated myosin regulatory light chain

(Figure 3f,g) than their congenic controls. Laser ablation assays confirm that apical

tension is significantly lower in *VANGL2-R353C* knock-in lines than their congenic

controls (Figure 3h,i). Thus, this *VANGL2* mutation associated with human NTDs

selectively impairs neuroepithelial apical constriction.

269
Figure 3: VANGL2-R353C mutation selectively impairs neuroepithelial apical constriction.
a, Schematic representation of PCP pathway, starting from non-canonical Wnt ligands (e.g.
Wnt5a/11) interacting with membrane co-receptors and activating cytoskeletal remodeling
responses. The transmembrane protein VANGL2 interacts with CELSR, which results to
downstream interaction with cytoplasmic proteins such as PRICKLE and SCRIB. The VANGL2-
R353C mutation is localised at the C-terminal region of the protein, in the prickle binding domain
(Pk-D). b, Representative western blot against VANGL2 and pVANGL2 at day 8 of
neuroepithelial differentiation in the isogenic control cell line (WT) and two knock in clones with
the R353C mutation (KI-1 and KI-2). c, Dot plots representing VANGL2 and pVANGL2 relative
band intensity levels after normalisation to GAPDH, respectively, across three replicates.
Intensity levels across the three lines were normalised to the WT levels of each replicate. d,
Representative high resolution confocal images of ZO-1 immunolabelled neuroepithelial cultures
at day 8, across the WT and KI cell lines. e, Dot plot representing the median apical area size of
day 8 neuroepithelial cultures, across the WT and KI cell lines. Lines link points from replicate
cultures, with the WT intensity set at 1. f, Representatives confocal images of neuroepithelial

cultures immunolabelled for phosphor-myosin light chain (pMLC)II, comparing the WT and KI-1
and KI-2 cell lines. Top view shows the distribution of protein of interest across the apical areas
and cross-section illustrates localisation along the apicobasal axis. Scale bar = 10 µm. **g,** *Dot*
plot showing normalised paired relative apical area pMLCII intensity, across the WT and KI-1
and KI-2 cell lines, on day 8 of differentiation. Relative apical intensity was normalised to WT
and set at 1, for each paired analysis, across four plates. **h,** *Live-imaged membrane-stained*
neuroepithelial sheet before and after a circular laser ablation, with overlayed image of post-
ablation timepoint and particle image velocimetry (PIV) analysis. The size and colour of arrows
in PIV analysis represent the direction and magnitude of recoil. Scale = 20 µm. **i,** *Median apical*
cluster displacement of neuroepithelial cells, across seven independent replicates. Each point
represents three individual quantifications in each plate and connected lines indicate paired
cultures. P values from Wilcoxon paired test with Bonferroni correction (***: p ≤ 0.001, *: p ≤
0.005, n.s.: not significant).

Establishment of patient-derived models of neuroepithelial dysfunction

As well as reverse genetic studies, an additional application of this highly reproducible

model is patient-specific testing of key neuroepithelial behaviours (Figure 4a).

Importantly, short-loop reprogramming of amniotic fluid stem cells with microfluidic

delivery of modified mRNA[57] enables rapid generation of patient-specific models in the

~4 months between spina bifida diagnosis and birth. Having patient-specific models for

functional genetic testing may be particularly useful to stratify post-natal management,

particularly of individuals with non-isolated NTDs[68]. As proof of principle, we used short-

loop reprogramming to compare two iPSC lines derived from amniotic fluid cells of

individuals who do not have spina bifida (here called GOC1-2 for simplicity) with two

lines of individuals treated at Great Ormond Street Hospital who underwent foetal

surgery for spina bifida (GOSB1-2). All four lines express comparably high levels of the

iPSC marker NANOG before differentiation, and differentiate to form endoderm,

ectoderm and mesoderm (Supplementary Figure 6).

Both control lines and the first spina bifida line - GOSB1 - achieve indistinguishable

apical areas and thicknesses (Figure 4b-e), further testament to the excellent

reproducibility of this modelling system when comparing non-congenic lines. In contrast,

the spina bifida line GOSB2 fails to apicobasally elongate as much as other lines and

retains larger apical areas (Figure 4b-e). We consider this is unlikely to represent an

apical constriction defect: laser ablation assays show comparable retraction in the

GOSB2 line compared with the other lines tested (Supplementary Figure 9). The

epithelia they form include mitotic cells which are not apically-located, in contrast to all
other cells studied which are pseudostratified and localise their mitoses apically
(Supplementary Figure 10). Unlike all other lines tested, differentiated GOSB2 cultures
also show abnormally heterogenous expression of CDH2 while partially retaining lamin
A/C (Figure 4f) at the end of the standard differentiation protocol. Attempts to adapt the
protocol to 'rescue' this phenotype are beyond the scope of this study given the
excellent reproducibility of differentiation in the seven other lines presented here.

We performed whole exome sequencing of all four amniocyte-derived lines to identify
coding variants of potential clinical relevance in GOSB1 and GOSB2. Variants were
compared with a list of genes previously associated with NTDs in mice and/or humans
(Figure 4g, Supplementary Table 1).

GOSB1 harbours 851 variant loci within 747 genes (Supplementary Table 2; Figure 4h).
Of all genes with variants, 27 are NTD-associated, such as *ZEB2* (variant 2-144389425-
T-C) and *ZIC5* (13-99970996-G-T). None of the variants in NTD-associated genes are
predicted to have a high transcript impact and all are heterozygous.

GOSB2, which has a more severe neuroepithelial phenotype, has 1122 variant loci in
939 unique genes of which 43 are in NTD-associated genes (Figure 4h). In this line, six
NTD-associated genes harbour variants predicted to have a high impact on transcript
sequence: *CSNK2A1*, *FAM20C*, *LFNG*, *MED24* and *PCSK5*.

Of these, *CSNK2A1* (20- 503643-G-GCATATTT), *MED24* (17-40019203-C-CACAT) and
PCSK5 (9-76175222-AAATGGAATGGAATGAAATGG-A) have gnomAD variant allele
frequencies <0.1. *MED24* (formerly called *TRAP100)* is of particular interest because its
deletion in mice results in a thin neuroepithelium which is not pseudostratified[69],
equivalent to our observations during GOSB2 differentiation. This gene is part of the
mediator complex involved in RNA polymerase II transcriptional co-activation. GOSB2
cells have compound heterozygous frameshift mutations in the same repetitive region of
the final exon of this gene. We additionally confirmed that the same *MED24* variants are
present in independent exome sequencing of amniocytes from which GOSB2 had been
derived (Supplementary Table 4). Both frameshifts are in the 3' untranslated region but
are predicted to alter RNA structure, causing loss of a multi-branch loop (Figure 4i), and

may therefore diminish binding of translation-regulating proteins. To test this, we

compared MED24 protein levels between GOSB2 iPSCs and three control lines, finding

reduced protein abundance compared to all three controls (Figure 4j,k). Although

genetic rescue experiments are beyond the scope of this study, particularly given the

likelihood of gene-gene interactions, equivalence of the cellular phenotype we observed

with a published mouse model and confirmation of reduced protein abundance suggest

the *MED24* variants in this line are pathogenic.

357
358 ***Figure 4: Spina bifida patient amniocyte-derived iPSC lines show can identify line-***
359 ***specific deficiencies in neuroepithelial cell function. a,*** *Experimental timeline of patient*

*amniocyte-derived iPSC lines, following microfluidic reprogramming. Whole exome sequencing was performed before and after reprogramming to identify line-specific deficiencies at genomic level and pair them to neuroepithelial and/or neural function level. **b,** Representative high-resolution top-view and cross-section images of day 8 neuroepithelial cultures visualizing F-actin. Two control cell lines (GOC1 and GOC2) were compared with two SB patient-derived hiPSCs cell lines (GOSB1 and GOSB2). **c,** Dot plot representing cell apical areas across the GOC1, GOC2, GOSB1, and GOSB2 cell lines at day 8 of neuroepithelial differentiation. Three independent plates were quantified for each line and the median apical area for each plate is shown with a black bar. **d,** Heatmaps representing the apical area size differences, across the four cell lines, in a contiguous area of 2,908 μm^2. **e,** Quantification of epithelial thickness on day 2 and day 8 samples, across four cell lines, GOC1, GOC2, GOSB1, and GOSB2. Points represent independent plate quantification for each line. Three paired replicates were measured for each time point and each cell line. Statistical test used for median values across quantifications was Wilcoxon paired test values with post-hoc Bonferroni **f,** Representative confocal images of LMN A/C, CDH2, SOX2, and LMNB immunolabelling in GOC2, and GOSB2, at day 8 neuroepithelial differentiation. **g,** Venn diagrams of GOSB1 and GOSB2 (**g'**) gene lists, after exome-sequencing, with at least one genetic variation observed in other datasets. Other datasets include previously associated mutations identified in humans under after search for the term "Neural Tube Defect" and "Spina Bifida" in ClinVar dataset, Lee and Geeson[5]. **h,** Filtering of variants in GOSB1 and GOSB2 to identify NTD-associated variants predicted to have a high impact on the transcript and which have a frequency <0.1 in gnomAD exome datasets. **i,** RNA structure of the MED24 major allele and rare variant (gnomAD reference 17-40019203-C-CACAT, allele exome frequency 0.018) predicted by RNAFold[70]. **j,** Bar plots representing normalised band intensity of MED24 across four cell lines. GOC2's intensity levels were set as 1 in each of three replicates. **k,** Western blot panel of iPSCs from four cell lines, GOC1, GOC2, GOSB2, and SFC086-06 showing MED24 protein and housekeeping GAPDH levels. Statistical test used for values was one-way ANOVA with post-hoc Bonferroni (***: p ≤ 0.001, **: p ≤ 0.001). Scale = 20 μm.*

As expected, given poor neuroepithelial morphogenesis of cells from the GOSB2 patient line, further differentiation does not produce polarised neural rosettes and results in only sparse differentiation of PAX6-positive neural progenitor cells with few neurites positive for the neuronal marker TUJ1 (Supplementary Figure 11). In contrast, GOSB1 formed prominent neuronal rosettes equivalently to the control lines (Figure 5a). However, by day 35 of neuronal differentiation, GOSB1 cultures show diminished TUJ1 positivity, without a change in the proportion of cells positive for the progenitor marker SOX2, compared with the two control lines differentiated in parallel (Figure 5b-d). Poor neuronal maturation in cells from the GOSB1 line is clearly evident by day 45 of differentiation, when control lines form dense aggregates interconnected by axons while GOSB1 cells form sparse and poorly-connected aggregates (Figure 5e,f). Thus, both the spina bifida patient-derived lines tested, in addition to *VANGL2-R353C* knock-in

model, demonstrate quantifiable yet distinct abnormalities in neuroepithelial functions

(Figure 5g).

Figure 5: Diminished neuronal maturation suggests post-neurulation deficit of an iPSC
lines derived from a patient with spina bifida. a, *Representative confocal images of GOC1,*
GOC2, and GOSB1 neuronal rosette cultures (day 20) immunolabelled against PAX6 and F-
actin. Scale = 100 µm. ***b,*** *Representative immunofluorescence panel of SOX2 and TUJ1*
immunofluorescence of day 30 cultures, showing GOC1, GOC2, and GOSB1 lines. Scale = 50
µm. ***c, d,*** *Dot plot representing the percentage of cells positive for TUJ1 (**c**) and SOX2 (**d**).*
Percentage positive nuclei were calculated in three independent replicates shown as an
individual dots and the average is shown with the bars. ***e,*** *Widefield images of day 45 GOC1*
and GOSB1 neuronal cultures immunolabelled for TUJ1. Scale = 200 µm. ***f,*** *Box plot*
representing the TUJ1 positive area quantified across three independent cultures of GOC1 and
GOSB1 cell lines, on day 45 of differentiation. P values from one-way ANOVA with post-hoc
Bonferroni (***: p ≤ 0.001, **: p ≤ 0.001, n.s.: not significant). ***g,*** *Summary of patient-associated*
hiPSCs lines used in this study along with the stage of failure (red rhombi) to form a functional
neuroepithelium or undergo mature neuronal formation in vitro.

Discussion

We present a multifaceted yet tractable and reproducible model of neuroepithelial

morphogenesis which is sufficiently sensitive to identify NTD-associated dysfunctions in

both forward and reverse genetic studies. All three of the NTD-associated models we

compare – knock-in of a patient-identified *VANGL2* mutation and two patient-derived

iPSC lines – identify abnormalities which are specific to that line, and which are not

present in a panel of control lines tested. The abnormalities observed relate to apical

constriction, apicobasal elongation, and neuronal maturation.

Neuroepithelial apical constriction is required for closure of both the cranial and spinal

neural tube[11,14,15,21]. Mouse and *Xenopus* models establish VANGL2 as a conserved

enhancer of apical constriction[20,22,65] but, to our knowledge, this had never been shown

in a human system and no human genetic mutations have been linked to apical

constriction as a likely pathogenic mechanism. The mechanisms by which VANGL2

promotes apical localisation and activation of myosin[20] remain incompletely understood.

They may include both PCP-specific functions of *VANGL2* equivalent to those originally

identified for its orthologue in *Drosophila*, and potentially vertebrate-specific roles[71,72].

The neuroepithelial model presented here does not produce planar features commonly

used as characteristic readouts of PCP signalling *in vivo*[26,40]. Nonetheless, our

observation of increasing VANGL2 poly-phosphorylation suggests ligand-induced

pathway activation during differentiation[35]. This contrasts with decreasing VANGL2

phosphorylation previously reported during *in vitro* differentiation of lactogenic cells[71].

The diminished apical constriction phenotype we document following *VANGL2-R535C*

mutation occurs despite achieving wildtype-equivalent VANGL2 localisation, apicobasal

elongation and cell volume. These cellular phenotypes are consistent with previously-

reported mutations in the protein's C-terminus domain in mice[20,39,40]. VANGL2's C-

terminus contains the prickle-binding domain (directly affected by the R353C mutation)

and PDZ domain, which interacts with other components of the PCP pathway including

SCRIB[73]. SCRIB also enhances apical constriction[15,74].

We observe that apical constriction correlates with apicobasal epithelial thickening

during iPSC-to-neuroepithelial differentiation. This inter-relation is well-established in

Drosophila ventral furrow formation[75,76], and can be replicated following optogenetic

induction of apical constriction in cultured mouse cells[77]. However, we provide two lines

of evidence that this inter-relation during neuroepithelial differentiation is not simply due

to volume conservation, as had been suggested[76]. Firstly, cell volume decreases during

neuroepithelial differentiation as thickness increases. Secondly, neither epithelial

thickness nor cell volume are significantly altered in *VANGL2-R535C* neuroepithelial

cells with diminished apical tension. Mechanistic separability of apical constriction and

apicobasal elongation is consistent with biomechanical modelling of *Xenopus* neural

tube closure showing that both are independently required for tissue bending[61]. The

molecular regulation of neuroepithelial apicobasal elongation, and simultaneous

pseudostratification, has not been extensively studied. In *Xenopus,* microtubule-

regulating proteins are required for elongation[78], and in mice global deletion of *Med24*

results in a non-pseudostratified and thinner neuroepithelium[69]. Finding that a patient-

derived iPSC line with diminished MED24 expression fails to apicobasally elongate and

pseudo-stratify by the end of the standard differentiation period suggests this cell shape

change is also important in human neural tube closure.

Future work will be needed to determine whether *MED24* mutation is the only genetic

change relevant to neuroepithelial morphogenesis in GOSB2. Another limitation of our

model, shared by many *in vitro* systems, is the poor ability to replicate maternal

environmental influences. For example, the validated cell culture media used for this

model contain 3.4 mg/L folic acid instead of folate. This may mask NTD predispositions

caused by maternal folate deficiency or genetically impaired placental transport, while

accentuating cellular deficiencies which diminish folic acid conversion into a one-carbon

donor[79]. We cannot exclude *in vitro* accentuation of the cellular phenotype of the

GOSB2 line, as it is difficult to imagine that such severe failure of neuroepithelial

morphogenesis would produce a sufficiently normal spinal cord to benefit from fetal

surgery performed on the patient these cells were derived from. Alternatively,

compensation may be possible *in vivo*: for example, *Med1* knockout mouse embryos

also have thinned neuroepithelium but still achieve neural tube closure[80]. MED family

proteins are increasingly implicated in human neural tube defects: *MED12* and *MED13L*

likely pathogenic variants have been reported in patients[9,81]. Their functions during

neural tube closure, including through known interactions with Wnt and PCP

pathways[82], remain incompletely understood.

MED protein functions in neurogenesis and neuronal maturation are more well-

established[80,83-85]. Others previously reported that iPSC lines derived from individuals

who have spina bifida are less able to differentiate into neurons than those from healthy

controls[86]. This is consistent with our findings in the GOSB1 patient-derived line.

Although gene rescue experiments are beyond the scope of this study, potentially

deleterious variants identified in the GOSB1 patient line, including a *ZIC5* variant, may

impair neuronal development. The reduction in TUJ1 positive cells observed in this line

compared with non-congenic controls is at a point in differentiation which is highly

comparable between individuals and produces deep-layer cortical neurons[87]. Individuals

with spina bifida commonly have thinning of the neocortex and other supratentorial brain

abnormalities[88]. What is less clear is whether diminished neurogenesis, which normally

occurs after closure of the neural tube, may relate to causation of NTDs. The converse

– that open spina bifida impairs neural development – is directly supported by

experimental evidence[89]. Nonetheless, impaired neuronal differentiation of patient-

derived iPSC lines raises the potential for shared genetic/epigenetic causation of the

spinal lesion and cortical brain abnormalities, potentially compounded by abnormalities

in cerebrospinal fluid drainage and amniotic fluid exposure.

In summary, our findings in a human neuroepithelial model corroborate animal studies

identifying diminished apical constriction as a cause of NTDs, implicate VANGL2 acting

upstream of myosin phosphorylation in this behaviour, identify diminished

neuroepithelial thickness in a patient-specific disease model and corroborate the

importance of MED family proteins known to enhance this process as an emerging

genetic contributor to human NTDs. The highly reproducible modelling platform we

evaluate – which is robust to differences in iPSC reprogramming method, sex and

ethnicity – represents a valuable tool for future mechanistic insights and personalised

disease modelling applications.

Materials and Methods

Human iPSC culture maintenance

All iPSCs lines were grown in E8 medium (ThermoFisher, A1517001), using 0.5%
Matrigel Growth Factor Reduced (Corning, 354277) coated plates, according to a
previously published protocol (Beers et al., 2012). Cells were routinely grown in 6 well
plates (SARSTEDT, 833920300), unless otherwise stated. E8 medium was fully
replaced daily. At ~80% confluency cells were washed with PBS without $CaCl_2$/$MgCl_2$
(ThermoFisher,14190094) and dissociated with 0.5 mM EDTA (ThermoFisher,
15575020) for 5 min. Cells were resuspended in E8 medium and re-plated at a 1:10
split ratio. All cell lines were tested monthly for mycoplasma contamination, using the
MycoAlert® kit (Lonza, LT07-318). A summary with the characteristics and
reprogramming technologies of the hiPSCs lines used on this study can be found in
Table 1.

Table 1: hiPSCs line information.

Cell line name	Designation[#]	Source	Karyotype	Reprogramming method
SFC086-03-01	iPSC2	Livesey Lab[17]	46, XX	Non-integrating Sendai virus
HO193b	iPSC1, GOC1	NIHR GOSH BRC*	46, XY	Microfluidic mmRNAs
HO471b	GOSB1	NIHR GOSH BRC*	46, XX	Microfluidic mmRNAs
HO220	GOC2	NIHR GOSH BRC*	46, XY	Microfluidic mmRNAs
HO541	GOSB2	NIHR GOSH BRC*	46, XY	Microfluidic mmRNAs
802-30F, parental	iPSC3, WT	SYNTHEGO	46, XX	StemRNA™ 3rd Gen
802-30F, VANGL2-R353C, clone E2	KI-1	SYNTHEGO	46, XX	StemRNA™ 3rd Gen
802-30F, VANGL2-R353C, clone F1	KI-2	SYNTHEGO	46, XX	StemRNA™ 3rd Gen

Shorthand nomenclature used in this paper for simplicity. *National Institute of Health
Research Great Ormond Street Hospital Biomedical Research Centre

VANGL2 knock-in cell line generation

The selection of the mutation was made according to a previously identified gene alteration in a patient with craniorachischisis (Lei et al., 2010). The specific point mutation was found in heterozygosity and was located on chromosome 1, at nucleotide 160,421,174 in the genomic sequence (Supplementary Figure 3). Although other *Vangl2* mutations have been found to exert dominant negative effects[90], we generated homozygous mutant lines to facilitate interpretation of results obtain. In collaboration with SYNTHEGO, an established hiPSC line 802-30F was genetically engineered to introduce a homozygous point mutation by replacing a cytosine (C) with a thymine (T) (CGC>TGC), at a specific genetic locus. After high knock-in efficiency was confirmed, specific sgRNA with Cas9 were electroporated to the cells. For the generation of the isogenic control line (WT), the cells were electroporated with SpCas9 only and confirmed to be unedited at target locus. Approximately two weeks later the cells colonies have expanded and genotyping of the pool of clones was undertaken. Individual clones were then isolated and clonally expanded to confirm the presence of the knock-in genetic alteration. Those clones positive for the genetic alteration were further Sanger sequenced to confirm stable *VANGL2* editing.

Adherent neural induction protocol

Direct differentiation of hiPSCs to neuroepithelial sheets was as we previously reported[17]. Briefly, hPSCs were lifted with EDTA and plated at 100% confluency, ~300,000 cells/cm^2, in E8 medium. It is critical to ensure that 100 % confluency is reached at seeding (considered 'day 1'). After 24 hours, cells were washed twice with PBS without $CaCl_2$/$MgCl_2$ and the medium was replaced with 2 ml of neural maintenance media (NMM, see Table 2), supplemented with 1 µM Dorsomorphin (Tocris, 3093) and 10 µM SB431542 (Tocris, 1614). NMM with these supplements is referred as neural induction medium (NIM) and was refreshed daily until day 8. At that timepoint the cultures were terminated or passaged for neuronal differentiation.

Neuronal differentiation was performed as previously described[56]. The neuroepithelial

sheets were passaged at day 8, using Dispase (ThermoFisher, 17105041) for 15-20

minutes at 37°C. After washing with PBS, neuroepithelial clumps were plated on 1%

Matrigel-coated 6-wells in NIM, ensuring sufficiently sparse seeding to give clumps

space to form rosettes. After 24 hours of passaging, the medium was changed to NMM

supplemented daily with 20 ng/ml of FGF2 (PeproTech, AF-100-18B) for the next four

days. After day 12, neural rosettes were passaged to 1% Matrigel coated plates, using

Dispase. NMM was then changed every 48 hours. This process was repeated at least

three-times, when confluency reaches 80 %, until day 25 when the neural progenitor

cells mature. At that stage, once the confluency reached 80 %, cells were passaged

with Accutase (Sigma, A6964) to dissociate the rosettes into single cells. At that stage

(~day 30) cells can be cryopreserved or plated for the last time at 25,000-75,000

cells/cm^2 on Laminin-521 (Gibco, A29249) coated plates.

Table Error! No text of specified style in document.**:** Composition of neural maintenance
medium.

Product	Volume (ml)	Company	Cat. Number	Storage (°C)
Solution A (500ml):				
2-Mercaptoethanol	1	Gibco	21985-023	4
DMEM/F12+Glutamax	481.25	Gibco	A15169-01	4
Insulin	0.25	Sigma	19278	4
N2 supplement (100x)	5	Gibco	17502-048	-20
Non-Essential amino Acids	5	Gibco	11140-050	4
Penicillin/Streptomycin	2.5	Gibco	15140-122	-20
Sodium Pyruvate	5	Sigma	S8636	4
Solution B (500ml):				
B27-supplement	10	Gibco	17504-044	-20
L-Glutamine	5	Gibco	25030-024	-20
Neurobasal	482.5	Gibco	12348-017	4
Penicillin/Streptomycin	2.5	Gibco	15140-122	-20

Immunofluorescent staining and confocal microscopy

Adherent cultures were stained for a maximum of 20 minutes in cold 4%

paraformaldehyde (PFA) and immunofluorescent staining as previously described

(Ampartzidis et al., 2023). Primary antibodies used were rabbit anti-ROCK (Abcam

antibody ab45171), rabbit anti-Ser19 pMLC (Cell Signalling Technology antibody

#3671), rabbit anti-SOX2 (Merck antibody ab5603), mouse anti-CDH2 (Cell Signaling

Technology antibody #14215), mouse anti-Lamin A/C (Santa Cruz antibody sc7292),

goat anti-Lamin B1 (Santa Cruz antibody sc6216), rabbit anti-ZO1 (Thermo Fisher

Scientific antibody #40–2200), goat anti-scribble (Santa Cruz antibody sc-11049), rat

anti-VANGL2 (Merck antibody, MABN750), rabbit anti-Pax6 (Biolegend antibody

901301), and mouse anti-TUJ1 (BioLegend MMS-435P), and detected with Alexa

Fluor™ conjugated secondary antibodies (Thermo Fisher Scientific). All primary and

secondary antibodies were used in 1:200 and 1:500 dilution, respectively. Nuclei were

labelled with DAPI (Severn Biotech Ltd 17057) and F-actin was labelled with Alexa

Fluor™-647 conjugated phalloidin (Thermo Fisher Scientific). Confocal salt and pepper

noise was removed using a 'remove outliers' filter and linear adjustments to brightness

and contrast were applied to parts of each figure panel equally.

Confocal images were captured on a Zeiss Examiner LSM880 confocal, using a

20x/NA1.0 Plan Apochromat dipping objective with AiryScan Fast. Widefield images of

the neural rosettes and neuronal cultures were captured on a Zeiss Observer inverted

using 10x/NA0.45 objective or a Nikon Ti2 Eclipse widefield microscope using

10x/NA0.30 or 20x/NA0.45 Plan Apochromat λD objectives.

Laser ablations and live imaging

Annual laser ablations were performed as previously reported[11,20] on day 8

neuroepithelial sheets. Briefly, cells were homogenously stained with CellMask

membrane dye (Invitrogen, C10046) and a single apical z-slice was selected for the

target region. The ablation circle was randomly placed in the stained neuroepithelial

sheet and three ablations were performed per dish then averaged (no predictable

change in tension was observed between repeated ablations in different fields of view).

The percentage reduction in cell area within the ablated annulus immediately after

ablation was quantified. The imaging order for each cell line was alternated between

each independent replicate to ensure no bias in the analysis and recoils were analysed

blinded to genotype/treatment.

For live-imaging sessions, day 8 neuroepithelial sheets were stained for 5 minutes with

CellMask Deep Red Actin Tracking stain (ThermoFisher, A57245) in NIM. Imaging was

performed with the Zeiss LSM880 confocal microscope using 20x/NA1.0 Plan

Apochromat dipping objective. The microscope chamber was pre-warmed at 37 $^{\circ}$C and

CO_2 was set at 5 %. The water-immersive objective was cleaned with 70 % ethanol

solution and let to air dry to remove any potential contaminants. A plexiglass chamber

was assembled around the culture plate to maintain stable culture condition. Damp

cotton wool was put in the plexiglass set up to create a humidified chamber and avoid

media evaporation. ZEN Black v2.3 was used to operate the microscope and capture

images. Apical region images were acquired every 5 minutes.

Particle intensity velocimetry (PIV) analysis

Particle Image Velocimetry (PIV) examined the flow of apical areas of day 8

neuroepithelial sheets after laser ablations. Briefly, we identify changes in pixel flow

before and after the ablation. For the PIV analysis used in this report a pre-installed plug

in FIJI was used (Tseng et al., 2012). The analysis considers the displacement of pixels

within a specific region, termed 'interrogation window'. This area in the present study

was specified between two frames, before and immediately after the circular laser

ablation. For the analysis a 'Pure Denoise' filter (Cycle-spins: 4, Multiframe: 2) was used

to ensure that non-specific random noise is not interpreted as movement.

Quantitative cell morphology analysis

Apical area analyses were performed in F-actin stained neuroepithelial sheets, using

the polygon selection tool in FIJI and outlining the apical cap of individual cells and

saving them as ROIs. At least 100 contiguous cells were analysed from each image.

Epithelial apicobasal thickness, confocal images were optically resliced and their cross-

section outline manually annotated. Thickness was calculated using the 'bounding

rectangle' (width) tool in ImageJ/Fiji. This was performed in three images across each
dish and the average height was calculated for each culture.

For fluorescent signal polarisation analysis, the plot profile command in ImageJ/FIJI was
applied to an optically-resliced section of equal thickness within each dataset.

TUJ1 and SOX2 neuronal quantification

For the quantification of TUJ1+ axons of neural progenitor, the total area of axons was
quantified relative to the area imaged. The TUJ1 signal was binarized using the
'Threshold' command, and the total area of the axons was summed. Finally, the TUJ1+
area was normalised to the total number of cells per field of view.

For the quantification of SOX2 nuclear marker in neuronal progenitor cultures, a multi-
step pipeline was used. Initially, all nuclei were identified in the DAPI-stained channel
and using the command 'Find Maxima' in FIJI. A binarized image was generated, with
individual points within each nucleus. Each point was enlarged and used to generate a
ROI within each nucleus The individual points were used as a reference to quantify the
SOX2 intensity within each cell. A threshold was set manually after identifying the
nucleus with the lowest positive signal. Finally, the percentage for positive nuclei was
calculated after subtracting the total cell numbers with the positive proportion and
dividing by the overall cell number.

Western blotting and quantification

Western blotting was performed as previously described[91]. Neuroepithelial cells were
lysed on ice for 30 min in 50 mM Tris Base pH 7.6, 150 mM NaCL, 1% Triton X-100,
0.02% Sodium Azide, 1x protease inhibitor cocktail (Merck Life Science UK Ltd), 1 mM
sodium orthovanadate, 25 mM sodium fluoride. The protein concentration was
determined using Pierce BCA Protein Assay kits (Thermo Fisher). Proteins were
resolved by SDS-PAGE using 10-12% polyacrylamide gels and transferred to PVDF
membranes. Membranes were blocked in TBS (15.4 mM Trizma-HCL, 4.62 mM Tris-

base, 150 mM NaCl, pH 7.6) containing 10% w/v milk powder (Merck Life Science UK

Ltd). Membranes were incubated with primary antibody overnight and then with

horseradish peroxidase (HRP)-conjugated secondary antibodies (Agilent Technologies,

Stockport, UK) for 1 hour. Primary antibodies were used rabbit anti-pVangl2 (Merck

antibody SAB5701946), mouse anti-NANOG (Abcam antibody ab173368), rabbit anti-

OCT4 (Abcam antibody ab19857), mouse anti-CDH1 (R&D FAB18381A), mouse anti-

GAPDH (Merck antibody MAB374), rabbit anti-MED24 (BETHYL antibody A301-472A).

HRP was detected using ECL Prime (Cytiva, Amersham, UK). To strip and re-probe,

membranes were incubated in 0.2 M NaOH for 20 min at 37 °C and then 20 min at room

temperature before re-blocking and re-use.

For the quantification of western blot films, the band area method was applied using

ImageJ/FIJI. A box was drawn around the bands and after ensuring all bands fit within it,

and the intensity profile was plotted. The total area between the line graph and x axis,

after removing background, was quantified for each sample. The protein level was

normalised to the housekeeping protein, LMNB or GAPDH. The normalised protein

levels were compared between samples, and a control value was normalised to 1.

Protein quantifications were performed in paired replicates or in three independent cell

lines.

Whole exome sequencing

Approximately 500 ng of DNA from each cell line, namely GOC1, GOC2, GOSB1, and

GOSB2, were purified. The subsequent steps of fragmentation, library preparation,

Illumina sequencing were performed using established pipelines by Azenta, UK. Upon

receipt of the samples, a quality control (QC) screening was executed, and all samples

successfully pass the QC threshold including concentration and maintaining high DNA

integrity number (DIN, >7.8). FASTQ data QC, genomic alignment, variant colling and

annotation was also performed by Azenta, UK. For all samples, >97.5% of target

regions achieved a read depth ≥20X. Variants were annotated using the Sequence

Ontology consensus terms. The following were considered to have a high impact:

transcript_ablation (SO:0001893), splice_acceptor_variant (SO:0001574),

splice_donor_variant (SO:0001575), stop_gained (SO:0001587), frameshift_variant

(SO:0001589), stop_lost (SO:0001578), start_lost (SO:0002012), and

transcript_amplification (SO:0001889).

NTD associated gene sorting workflow

Lists of genes with annotated variants in GOSB1 and GOSB2 were compared with

GOC1 and GOC2. Only variants not shared with GOC1/2 were considered in order to

eliminate batch effects and very common variants. Each of resulting gene lists was then

cross-referenced with the most current compilation of mouse models associated with

NTDs, as documented in Lee and Gleeson[5]. Two additional gene lists were generated

based on searches in ClinVar database for entries linked to "Neural Tube Defect" and

"Spina Bifida" (Supplementary Table 1). Collectively, these identified a subset of variant

genes associated with NTDs. These were subsequently filtered by sequency ontology

severity score, and the allele frequencies of high impact variants was determined using

gnomAD v4.1.0[92].

Statistical analysis and reproducibility

Comparisons between two groups were by Student's t-test accounting for homogeneity

of variance in SPSS (IBM Statistics 25). Comparison of multiple groups was first tested

for normality and homogeneity of variance in Origin 2020 (Origin Labs). According to the

result if samples were normally distributed an ANOVA test was performed, while for non-

parametric datasets a Kruskal-Wallis ANOVA was used in Origin 2020 (Origin Labs).

Post-hoc pairwise analysis was performed to determine the statistical significance. P

values <0.05 were considered statistically significant. All graphs were generated in

Origin 2020. Replicates were defined as technical replicates (same iPSC line cultured at

the same time) or biological replicates (different iPSC lines or same line cultured

independently). In the case of VANGL2 knock-in lines, two clones were compared with

their congenic control line. To avoid bias, blinding and randomisation was performed

were possible by asking a lab member to rename copied files, without disclosing cell

line information prior to analysis. However, changes of neuroepithelial thickness across

time points days was too dramatic to not know which day was analysed and

morphological differences of GOSB3 also precluded blinding.

Acknowledgements

The authors would like to thank Prof F Livesey and Dr F Paonessa for their assistance

in establishing the dual-SMAD inhibition protocol in our lab and provision of SFC086-03-

01 as a reference line. We thank Jessica Reeves for technical assistance with western

blots, Dr Teresa Pereira da Silva for scientific discussions and cell culture assistance, Dr

Timo N. Kohler and Clara Munger for their critical reading of the manuscript, and Dr

Dale Moulding for assistance with imaging. IA was funded by the NIHR GOSH

Biomedical Research Centre. Views expressed in this manuscript are solely those of the

authors and not necessarily those of the NHS, the NIHR or the Department of Health.

GLG acknowledges funding from the Royal Society (RG\R2\232082), Rosetrees

(2023\100266), NC3Rs (NC/Y50063X/1) and Wellcome Trust (211112/Z/18/Z).

Author contributions

IA, EMT, YG, AK and GLG performed and analysed experiments. GLG, PDC and NE

conceived the study and designed experiments with IA. GLG, PDC and IA secured

funding. GLG, PDC, NE and EM supervised staff. GLG and IA wrote the manuscript with

input from all co-authors.

References

1   Zaganjor, I. *et al. PLoS One* **11**, e0151586, doi:10.1371/journal.pone.0151586
(2016).
2   Santos, C. *et al.*   (eLife Sciences Publications, Ltd, 2024).
3   Nikolopoulou, E. *et al. Development* **144**, 552-566, doi:10.1242/dev.145904
(2017).
4   Chen, Z. *et al. Cell Res* **28**, 1039-1041, doi:10.1038/s41422-018-0061-3 (2018).
5   Lee, S. & Gleeson, J. G. *Trends Neurosci* **43**, 519-532,
doi:10.1016/j.tins.2020.04.009 (2020).
6   Finnell, R. H. *et al. Front Genet* **12**, 659612, doi:10.3389/fgene.2021.659612
(2021).
7   Cao, X. *et al. Hum Mutat* **41**, 786-799, doi:10.1002/humu.23969 (2020).
8   Crane-Smith, Z. *et al. Hum Mol Genet* **32**, 2681-2692, doi:10.1093/hmg/ddad094
(2023).
9   Tian, T. *et al. Front Cell Dev Biol* **9**, 641831, doi:10.3389/fcell.2021.641831
(2021).
10   Maniou, E. *et al. Development* **150**, doi:10.1242/dev.202139 (2023).
11   Butler, M. B. *et al. J Cell Sci* **132**, doi:10.1242/jcs.230300 (2019).
12   McShane, S. G. *et al. Dev Biol* **404**, 113-124, doi:10.1016/j.ydbio.2015.06.003
(2015).
13   Ferreira, M. A. *et al. PLoS Comput Biol* **15**, e1007171,
doi:10.1371/journal.pcbi.1007171 (2019).
14   Brooks, E. R. *et al. Elife* **9**, doi:10.7554/eLife.60234 (2020).
15   Lesko, A. C. *et al. Dev Biol* **478**, 59-75, doi:10.1016/j.ydbio.2021.05.013 (2021).
16   Smith, J. L. *et al. J Comp Neurol* **342**, 144-151, doi:10.1002/cne.903420113
(1994).
17   Ampartzidis, I. *et al. Dev Biol* **494**, 60-70, doi:10.1016/j.ydbio.2022.12.002
(2023).
18   Martin, A. C. & Goldstein, B. *Development* **141**, 1987-1998,
doi:10.1242/dev.102228 (2014).
19   Escuin, S. *et al. J Cell Sci* **128**, 2468-2481, doi:10.1242/jcs.164574 (2015).
20   Galea, G. L. *et al. Nat Commun* **12**, 1159, doi:10.1038/s41467-021-21372-4
(2021).
21   McGreevy, E. M. *et al. Biol Open* **4**, 186-196, doi:10.1242/bio.20149589 (2015).
22   Kowalczyk, I. *et al. Development* **148**, doi:10.1242/dev.195008 (2021).
23   Ossipova, O. *et al. Nat Commun* **5**, 3734, doi:10.1038/ncomms4734 (2014).
24   Nishimura, T. *et al. Cell* **149**, 1084-1097, doi:10.1016/j.cell.2012.04.021 (2012).
25   Boutros, M. *et al. Cell* **94**, 109-118, doi:10.1016/s0092-8674(00)81226-x (1998).
26   Adler, P. N. *Bioessays* **14**, 735-741, doi:10.1002/bies.950141103 (1992).
27   Gubb, D. & Garcia-Bellido, A. *J Embryol Exp Morphol* **68**, 37-57 (1982).
28   Murdoch, J. N. *et al. Hum Mol Genet* **10**, 2593-2601,
doi:10.1093/hmg/10.22.2593 (2001).
29   Kibar, Z. *et al. Nat Genet* **28**, 251-255, doi:10.1038/90081 (2001).
30   Wallingford, J. B. *et al. Nature* **405**, 81-85, doi:10.1038/35011077 (2000).
31   Rothbacher, U. *et al. Dev Biol* **170**, 717-721, doi:10.1006/dbio.1995.1249 (1995).

32 Aw, W. Y. & Devenport, D. *Curr Opin Cell Biol* **44**, 110-116,
doi:10.1016/j.ceb.2016.08.002 (2017).
33 Yu, H. *et al. Development* **137**, 3707-3717, doi:10.1242/dev.052001 (2010).
34 Murdoch, J. N. *et al. Dis Model Mech* **7**, 1153-1163, doi:10.1242/dmm.016758
(2014).
35 Yang, W. *et al. Cell Res* **27**, 1466-1484, doi:10.1038/cr.2017.127 (2017).
36 Humphries, A. C. *et al. PLoS Genet* **19**, e1010849,
doi:10.1371/journal.pgen.1010849 (2023).
37 Ybot-Gonzalez, P. *et al. Development* **134**, 789-799, doi:10.1242/dev.000380
(2007).
38 Nychyk, O. *et al. Dis Model Mech* **15**, doi:10.1242/dmm.049194 (2022).
39 Lopez-Escobar, B. *et al. Development* **145**, doi:10.1242/dev.157487 (2018).
40 Galea, G. L. *et al. Dis Model Mech* **11**, doi:10.1242/dmm.032219 (2018).
41 Baumholtz, A. I. *et al. Dev Biol* **428**, 25-38, doi:10.1016/j.ydbio.2017.05.013
(2017).
42 Doudney, K. *et al. Am J Med Genet A* **136**, 90-92, doi:10.1002/ajmg.a.30766
(2005).
43 De Marco, P. *et al. Birth Defects Res A Clin Mol Teratol* **100**, 633-641,
doi:10.1002/bdra.23255 (2014).
44 Iliescu, A. *et al. Biochemistry* **53**, 5356-5364, doi:10.1021/bi500400g (2014).
45 Tian, T. *et al. Neurogenetics* **21**, 217-225, doi:10.1007/s10048-020-00613-6
(2020).
46 Lei, Y. P. *et al. N Engl J Med* **362**, 2232-2235, doi:10.1056/NEJMc0910820
(2010).
47 Kibar, Z. *et al. Clin Genet* **80**, 76-82, doi:10.1111/j.1399-0004.2010.01515.x
(2011).
48 Aguiar-Pulido, V. *et al. Proc Natl Acad Sci U S A* **118**,
doi:10.1073/pnas.2106844118 (2021).
49 Karzbrun, E. *et al. Nature* **599**, 268-272, doi:10.1038/s41586-021-04026-9
(2021).
50 Takla, T. N. *et al. Cells* **12**, doi:10.3390/cells12131697 (2023).
51 Tidball, A. M. *et al. Stem Cell Reports* **18**, 2498-2514,
doi:10.1016/j.stemcr.2023.10.020 (2023).
52 Mirdass, C. *et al. Emerg Top Life Sci* **7**, 423-437, doi:10.1042/ETLS20230087
(2023).
53 Abdel Fattah, A. R. *et al. Cell Rep* **42**, 113334, doi:10.1016/j.celrep.2023.113334
(2023).
54 Krammer, T. *et al. Dev Cell* **59**, 1940-1953 e1910,
doi:10.1016/j.devcel.2024.04.021 (2024).
55 Rabeling, A. *et al. Stem Cell Rev Rep* **20**, 2045-2061, doi:10.1007/s12015-024-
10785-5 (2024).
56 Shi, Y. *et al. Nat Protoc* **7**, 1836-1846, doi:10.1038/nprot.2012.116 (2012).
57 Gagliano, O. *et al. Nat Protoc* **14**, 722-737, doi:10.1038/s41596-018-0108-4
(2019).
58 Maniou, E. *et al. Nat Mater* **23**, 1575-1581, doi:10.1038/s41563-024-01942-9
(2024).

59 Bluhmki, T. *et al. Methods Mol Biol* **1994**, 101-115, doi:10.1007/978-1-4939-9477-
9_9 (2019).
60 Zhang, P. P. *et al. Acta Pharmacol Sin* **45**, 282-297, doi:10.1038/s41401-023-
01172-w (2024).
61 Inoue, Y. *et al. Biomech Model Mechanobiol* **15**, 1733-1746, doi:10.1007/s10237-
016-0794-1 (2016).
62 Schoenwolf, G. C. & Franks, M. V. *Dev Biol* **105**, 257-272, doi:10.1016/0012-
1606(84)90284-7 (1984).
63 Matsuda, M. *et al. Nat Commun* **14**, 8475, doi:10.1038/s41467-023-43973-x
(2023).
64 Campas, O. *et al. Nat Rev Mol Cell Biol* **25**, 252-269, doi:10.1038/s41580-023-
00688-7 (2024).
65 Ossipova, O. *et al. Development* **142**, 99-107, doi:10.1242/dev.111161 (2015).
66 Humphries, A. C. *et al. Elife* **9**, doi:10.7554/eLife.53532 (2020).
67 Kasioulis, I. *et al. J Cell Sci* **135**, doi:10.1242/jcs.259897 (2022).
68 Che, X. *et al. Biology (Basel)* **11**, doi:10.3390/biology11091371 (2022).
69 Ito, M. *et al. EMBO J* **21**, 3464-3475, doi:10.1093/emboj/cdf348 (2002).
70 Hofacker, I. L. & Stadler, P. F. *Bioinformatics* **22**, 1172-1176,
doi:10.1093/bioinformatics/btl023 (2006).
71 Rubio, S. *et al. Cells* **13**, doi:10.3390/cells13030222 (2024).
72 Walton, A. *et al. J Biol Chem* **300**, 106792, doi:10.1016/j.jbc.2024.106792 (2024).
73 How, J. Y. *et al. Biochem J* **478**, 1321-1332, doi:10.1042/BCJ20200816 (2021).
74 Boeda, B. *et al. J Cell Biol* **222**, doi:10.1083/jcb.202211113 (2023).
75 He, B. *et al. Nature* **508**, 392-396, doi:10.1038/nature13070 (2014).
76 Gelbart, M. A. *et al. Proc Natl Acad Sci U S A* **109**, 19298-19303,
doi:10.1073/pnas.1205258109 (2012).
77 Martinez-Ara, G. *et al. Nat Commun* **13**, 5400, doi:10.1038/s41467-022-33115-0
(2022).
78 Suzuki, M. *et al. Development* **137**, 2329-2339, doi:10.1242/dev.048769 (2010).
79 Leung, K. Y. *et al. Brain* **136**, 2836-2841, doi:10.1093/brain/awt209 (2013).
80 Ito, M. *et al. Mol Cell* **5**, 683-693, doi:10.1016/s1097-2765(00)80247-6 (2000).
81 Ha, Y. J. *et al. Nature*, doi:10.1038/s41586-025-08676-x (2025).
82 Rocha, P. P. *et al. Development* **137**, 2723-2731, doi:10.1242/dev.053660 (2010).
83 Haque, F. *et al. Sci Adv* **8**, eadd1466, doi:10.1126/sciadv.add1466 (2022).
84 Hamada, N. *et al. Dev Neurosci* **43**, 43-52, doi:10.1159/000515188 (2021).
85 Zhu, W. *et al. Development* **142**, 465-476, doi:10.1242/dev.112946 (2015).
86 Sahakyan, V. *et al. Sci Rep* **8**, 2942, doi:10.1038/s41598-018-21103-8 (2018).
87 Strano, A. *et al. Cell Rep* **31**, 107732, doi:10.1016/j.celrep.2020.107732 (2020).
88 Fietz, S. A. *et al. J Neurosci* **40**, 1766-1777, doi:10.1523/JNEUROSCI.0192-
19.2019 (2020).
89 Clark, M. *et al. bioRxiv*, 2025.2001.2006.631442,
doi:10.1101/2025.01.06.631442 (2025).
90 Yin, H. *et al. PLoS One* **7**, e31988, doi:10.1371/journal.pone.0031988 (2012).
91 Thompson, E. M. *et al. FEBS Open Bio* **12**, 1388-1405, doi:10.1002/2211-
5463.13418 (2022).
92 Chen, S. *et al. Nature* **625**, 92-100, doi:10.1038/s41586-023-06045-0 (2024).

Supplementary Figures and Legends

Supplementary Figure 1: Nuclear volume decreases during iPSC differentiation. a.
Visualisation of nuclei on day 2 and 8 of neuroepithelial differentiation. Dashed white lines
indicate nuclear outlines. Representative of multiple iPSC lines including iPSC1-3. Scale = 25
*μm. **b**, Quantification of nuclear volume measured by manually drawing around each nucleus in*
AiryScan Z-stacks spread throughout the apicobasal thickness of the epithelium. Points
represent individual nuclei (30 nuclei per day, 10 nuclei from three differentiation experiments).

Supplementary Figure 2: Comparison of neuroepithelial morphology in cultures
*differentiated by three different investigators. a-b. Neuroepithelial thickness (**a**) and apical*
*area (**b**) in iPSC2 cells differentiated by ET, GG and IA. All were analysed by GG. Points*
*represent different cultures (**a**) and individual cells (**b**, 300 cells from 3 cultures per investigator,*
with bars showing medians). No groups are significantly different from others by T-test.

Supplementary Figure 5: VANGL2-R353C mutation does not alter apical localization. a.
VANGL2 immunofluorescence showing apical and lateral views of iPSC-derived
*neuroepithelium from the wildtype (WT) and two knock-in (KI) lines. Scale bar = 50 µm. **b.***
Profile intensity quantification of VANGL2 immunofluorescence in three cultures of each line.
Points represent the mean ± standard deviation. All plots are aligned so that their maximum
values are in the same position.

Supplementary Figure 6: Tri-lineage differentiation of hiPSC lines. *a. Schematic*
representation of tri-lineage differentiation protocol, using STEMdiff™ Trilineage Differentiation
patient-derived cell lines after tri-lineage differentiation, with cells stained for NANOG (grey), FOXA2
(orange), T (cyan), and PAX6 (magenta) markers. Scale = 100 µm.

Supplementary Figure 7: VANGL2-R353C mutation does not impair neuroepithelial apicobasal elongation. a,b, Optical cross-section of neuroepithelial sheets on day 2 (a) and day 8 (b), after staining with F-actin (grey), respectively. Scale = 20 µm. c, Quantification of epithelial thickness on day 2 and day 8 samples, across all cell lines. Points represent independent plate quantification for each line. Three paired replicates were measured for each time point and each cell line. Paired T-test was performed for mean epithelial thickness for each cell line across day 2 and day 8 samples, and one-way ANOVA with post-hoc Bonferroni for the day 8 comparison across the cell lines. d, e. Optical cross-section of neuroepithelial sheets on day 2 (D) and day 8 (E), after staining with DAPI (grey). Dots annotate individual nuclei across the epithelium. Scale = 20 µm. f, Quantification of nuclei dispersed in a row (nuclear index) of day 2 and day 8 samples. Two sample T-test performed for mean nuclear index for each cell line across day 2 and day 8 samples (***: p ≤ 0.0001, n.s.: not significant).

Supplementary Figure 8: VANGL2-R353C mutation does not alter neuroepithelial cell volume. a. Bar graph overlaid with dot plot represent the quantification of neuroepithelial cell volume across the WT, F1, and E2 cell lines, with 15 cells analysed in each cell line. Statistical test used for individual cell volume and sub-apical protrusion length was Mann Whitney U-test with post-hoc Bonferroni (n.s.: not significant). b. Representative image of subapical cross-section of neuroepithelial cells stained for Bio-Tracker (grey). Arrowheads indicate sub-apical protrusions. Scale = 20 µm.

Supplementary Figure 9: Neuroepithelial cells differentiated from the control and spina bifida patient-derived iPSC lines show equivalent apical retraction following laser ablation. a. Live-imaged CellMask-stained neuroepithelial sheet before and after a linear circular laser ablation (white dash line). Scale = 20 μm. **b.** Particle image velocimetry (PIV) analysis represent the direction recoil, in the last panel. **c.** Quantification of median apical cluster displacement of neuroepithelial cells, across three independent biological plates. Statistical test used for mean values across quantifications was one way ANOVA with post-hoc Bonferroni (n.s.: not significant).

Supplementary Figure 10: Sub-apical localisation of mitotic figure in GOSB2 spina bifida patient-derived iPSC line. a,b. *Representative high-resolution confocal cross-section (A) and top-view (B) images of day 8 GOSB2 cultures stained against F-actin and DAPI.* ***c, d.*** *Representative high-resolution confocal cross-section (**a**) and top-view (**b**) images of day 8 GOC2 cultures stained against F-actin and DAPI. Dashed lines indicate basal site of neuroepithelial culture. The arrow lines (white and green) in both merged images correspond to equivalent apical and basal slices shown in panel **b** and **d**, respectively for GOSB2 and GOC2. Asterisk annotates the mitotic figure during metaphase.*

974
975 ***Supplementary Figure 11: Neuronal rosette characterisation between GOC2 and GOSB2***
976 ***patient-derived cell lines. a.*** *Representative immunofluorescence panel of day 20, neuronal*
977 *rosettes stage cultures stained against PAX6 (magenta) and DAPI (grey). Scale = 50 µm.* ***b.***
978 *Representative immunofluorescence panel of day 20, neuronal rosettes stage cultures stained*
979 *against TUJ1 (cyan) and DAPI (grey). Arrows indicate TUJ1+ outgrowths. Scale = 50 µm.*