Parkin-dependent ubiquitination of TAX1BP1 affects the degradation pathway of

defective mitochondria

Anna Lechado-Terradas[1, *], Bianca Lemke[2], Katharina I. Zittlau[2, #], Boris Macek[2], Philipp J.

Kahle[1, 3, *]

[1]Laboratory of Functional Neurogenetics, Department of Neurodegeneration, Hertie

Institute for Clinical Brain Research and German Center for Neurodegenerative Diseases,

Faculty of Medicine, University of Tübingen, Tübingen, Germany.

[2]Quantitative Proteomics Group, Department of Biology, Interfaculty Institute of Cell

Biology, University of Tübingen, Tübingen, Germany.

[3]Department of Biochemistry, Faculty of Science, University of Tübingen, Tübingen,

Germany.

[#]present address: Laboratory for Structural Mass Spectrometry, Department of

Biomolecular Sciences, Weizmann Institute of Science, Rehovot, Israel.

*Corresponding author(s): philipp.kahle@uni-tuebingen.de, anna.lechado-terrads@uni-

tuebingen.de

Keywords: Autophagosome, endosome, mitophagy, parkin, TAX1BP1, ubiquitination

Abstract

In stressed cells, the recessive Parkinson disease (PD) associated gene products PINK1 and parkin mediate the autophagic removal of damaged mitochondria (mitophagy). Upon mitochondrial membrane potential disruption, PINK1 phosphorylation activates the ubiquitin ligase parkin which ubiquitinates various mitochondrial protein substrates. These feed-forward modifications on the mitochondria surface attract ubiquitin-binding autophagy receptors that target ubiquitinated mitochondria for degradation. Investigating post-translational protein modifications during this process, we detected transient ubiquitination of K549 within the third coiled-coil domain (CC3) of TAX1BP1 in HeLa cells expressing wild-type (WT) but not catalytically inactive parkin. Parkin-dependent ubiquitination did not target TAX1BP1 to proteasomal degradation but was rather indicative of a regulatory modification. In cells with the full complement of autophagy receptors, TAX1BP1 plays only a minor role in mitophagy. However, when expressed as a sole autophagy receptor, both WT and ubiquitination deficient TAX1BP1 were capable of promoting mitophagy, albeit mitochondria degradation was slightly delayed under mutant conditions. Use of the lysosomal inhibitor bafilomycin A indicated classical autophagolysosomal targeting of damaged mitochondria mediated by WT TAX1BP1. However, for the ubiquitination-deficient TAX1BP1, we observed an increased prevalence of enlarged endolysosomal vesicles carrying accumulated TAX1BP1-positive autophagosomes filled with mitochondrial material. Thus, while ubiquitination of the CC3 domain of TAX1BP1 is not essential for complete mitophagy, the lack of CC3 in TAX1BP1 reroutes the degradation flux to a less efficient endolysosmal degradative pathway. Interestingly, the PD gene product VPS35 becomes prominently engaged in this alternative mitophagy pathway.

Keywords

Autophagosome, endosome, mitophagy, parkin, TAX1BP1, ubiquitination.

Abbreviations

5KO: penta knock-out HeLa; ATG9A: autophagy-related protein 9A; BafA: Bafilomycin A;

CALCOCO2/NDP52: calcium binding and coiled-coil domain 2; CC3: third coiled-coil

domain of TAX1BP1; CCCP: carbonyl cyanide m-chlorophenylhydrazone; CD63: CD63

antigen; COX4: cytrochrome c oxidase subunit 4; CS: citrate synthase; HSP60: heat shock

protein 60; LC3B: microtubule-associated protein 1 light chain 3 beta; MFN1: mitofusin-1;

MMF: membrane fraction; MTF: mitochondria fraction; MYO6: unconventional myosin-VI;

NBR1: NBR1 autophagy cargo receptor; OMM: outer mitochondrial membrane; OPTN:

optineurin; SQSTM1/p62: sequestosome-1; PD: Parkinson disease; PINK1:

serine/threonine-protein kinase PINK1; PRKN: E3-ligase parkin; PTM: post-translational

modification; RAB7: ras-related protein rab-7a; SSBP1: single-stranded DNA-binding

protein; TAX1BP1: Tax1-binding protein 1 homolog; TBC1D15: TBC1 domain family

member 15; TOM20: translocase of outer membrane 20 kDa subunit; UBD: ubiquitin binding

domain; VDAC: voltage-dependent anion channel; VPS35: Vacuolar protein sorting-

associated protein 35; WDR45B: WD repeat domain phosphoinositide-interacting protein 3;

WT: wild-type.

Introduction

Mitochondrial quality control and turnover are essential for cellular homeostasis. Cellular stress associated with mitochondrial membrane depolarization induces a pathway leading to removal of damaged mitochondria activated by the two most common recessive Parkinson disease (PD) associated gene products, PINK1 (phosphatase and tensin homolog induced kinase 1) and PRKN (E3 ligase parkin) [1–4]. Under basal conditions, the nuclear encoded PINK1 protein is efficiently imported into mitochondria, where it is rapidly degraded. Upon dissipation of the mitochondrial membrane potential, PINK1 lacks the driving force for mitochondrial import and hence accumulates on the mitochondrial outer membrane (OMM). There, the PINK1 kinase domain faces the cytosol to phosphorylate ubiquitin and the ubiquitin-like domain of parkin, a process which activates the ubiquitin ligase parkin recruited to the OMM of depolarized mitochondria [5–8]. Various ubiquitin linkages target proteins from depolarized mitochondria to proteasomal and autophagic degradation [9]. Five major selective autophagy receptors have been implicated to connect ubiquitinated mitophagy cargo with the autophagy machinery, including SQSTM1/p62 (sequestosome-1), OPTN (optineurin), CALCOCO2/NDP52 (calcium binding and coiled-coil domain 2), NBR1 (NBR1 autophagy cargo receptor), and TAX1BP1 (Tax1 binding protein 1) [2,10–12]. These selective autophagy receptors do not statically bridge ubiquitinated mitophagy cargo with membranes containing the lipidated form of LC3B (microtubule-associated protein 1 light chain 3 beta), but can be dynamically regulated by post-translational modifications (PTMs). These PTMs are critical modulators of their function and contribute to the recruitment of autophagy-related proteins involved in autophagosome biogenesis (i.e: ATG9 vesicles or the unc-51 like autophagy activating kinase 1 complex) [13–16]. For example, phosphorylation

by TANK binding kinase 1 contributes to important regulatory dynamics of OPTN and

NDP52 for selective autophagic removal of damaged mitochondria [10,12,16].

Investigating the phospho-proteome and ubiquitylome throughout the time course of

mitophagy in catalytically active parkin expressing HeLa cells [17], we detected transient

ubiquitination of TAX1BP1 at K549 within the third coiled-coil domain (CC3) of this

autophagy receptor. Validation of TAX1BP1 ubiquitination in cells expressing wild-type

(WT) or a ligase-dead parkin mutant (C431A), confirmed parkin-dependent ubiquitination

of the autophagy receptor at early stages of mitophagy. Removal of the highly ubiquitinated

CC3 domain of TAX1BP1 reduced TAX1BP1 distribution on mitochondrial material and

induced the engulfment of accumulated autophagosomes into enlarged endolysosomal

vesicles (potential amphisomes), apparently rerouting the mitophagy pathway to the

endolysosomal system. Thus, the ubiquitinatable CC3 of TAX1BP1 directs depolarized

mitochondrial material to autophagosomal degradation, whereas CC3 perturbation mediates

the degradation of accumulated autophagosomes with damaged mitochondrial cargo through

a less efficient pathway involving endolysosomal trafficking into enlarged endolysosomes.

Interestingly, the PD gene product VPS35 (VPS35 retromer complex component) became

prominently engaged in the alternative mitophagy pathway described in this study, clearly

recognizing the surroundings of the identified enlarged degradative vesicles.

Results

Transient ubiquitination of TAX1BP1 in Parkin-dependent mitophagy

Examining the ubiquitylome of HeLa cells stably expressing parkin treated with the

mitochondrial uncoupler carbonyl cyanide m-chlorophenylhydrazone (CCCP) [17], we

detected transient ubiquitination of TAX1BP1 at K549 (**Figure 1A and B**). The

ubiquitination-indicating di-glycine motifs were detected by mass spectrometry

predominantly for TAX1BP1 peptides containing K549, with only minor signals for K571,

also located in the CC3 region, and K439, upstream of CC3 (**Figure 1A and B**). TAX1BP1

ubiquitination only occurred in cells expressing WT parkin, but not the catalytically inactive

form (C431A-parkin) (**Figure 1B**). Other studies previously reported K549-TAX1BP1 to be

increasingly ubiquitinated in a parkin-dependent manner and to follow a similar time course

as the one described here [18]. Validation of parkin-dependent ubiquitination of TAX1BP1

via western blotting confirmed the differences between WT- and C431A-parkin observed in

the mass-spectrometry screen (**Figure 1C**). In HeLa cells stably expressing active parkin, but

not ligase-inactive C431A-parkin, CCCP treatment caused a band shift compatible with

TAX1BP1 poly-ubiquitination peaking around 4 h after mitochondrial membrane

depolarization. TAX1BP1 ubiquitination vanished at later time points of mitophagy, when

parkin substrates like TOM20 (translocase of outer mitochondrial membrane 20) and VDAC

(voltage dependent anion channel) were degraded (**Figure 1C**).

To confirm that the observed TAX1BP1 band shifts reflect ubiquitination of the autophagy

adaptor, ubiquitin pulldown experiments were performed in the WT-Parkin HeLa cell line

(**Figure 1D**). After short CCCP depolarization, some parkin-dependent substrates were

pulled down such as ubiquitinated (Ub-) TOM20 as well as ubiquitin in its unmodified form

(Total Ub) or the Ser65-phosphorylated version (pS65-Ub) (**Figure 1D**). The appearance of

a smear-like band shift for TAX1BP1 was also detected under the same conditions, which

contrasted with the absence of a smear observed under basal or bead control pulldown

conditions (**Figure 1D**). This data confirms the ubiquitination nature of

endogenousTAX1BP1 occurring in a parkin-dependent mitophagy context.

Overall, TAX1BP1 levels decreased parallelly to VDAC and TOM20 degradation during

early depolarization (2-4 h CCCP) but increased after extended depolarization (8-18 h CCCP)

(**Figure 1C**), indicating that early steps of mitophagy might also involve some degradation

of the autophagy receptor along with mitochondrial cargo. As early stages of PINK1/parkin

mitophagy rely on proteasomal degradation of immediate-early parkin substrates, we

checked if the identified ubiquitination of TAX1BP1 could target the autophagy receptor for

early proteasomal degradation. In order to test this, HeLa cells stably expressing parkin were

transfected with TAX1BP1 and subjected to mitophagy in the presence or absence of the

proteasomal inhibitor MG132 (**Fig. S1**). Here, while endogenous TAX1BP1 was not reduced

after short depolarization (2 h CCCP), a slight reduction of overexpressed TAX1BP1 levels

was observed at this early time point (**Fig. S1**). Importantly, proteasome inhibition did not

stabilize poly-ubiquitinated TAX1BP1, neither in the endogenous or overexpressed form,

contrary to what was observed for the known proteasome target MFN1 (mitofusin 1) (**Fig.**

S1). Thus, the rapid TAX1BP1 ubiquitination observed in this study, seemed to not be part

of the initial proteasome-targeting ubiquitination program of OMM-associated proteins, but

rather appears to be a regulatory PTM of TAX1BP1.

Ubiquitinated TAX1BP1 is enriched in depolarized mitochondria

Mitochondria depolarization triggers the activation of parkin-dependent ubiquitination of mitochondrial proteins that will be recognized by the ubiquitin binding domains of autophagy receptors for further successful delivery of mitochondrial cargo to autophagosomes [19]. As expected, endogenous TAX1BP1 translocated to mitochondria after depolarization (**Figure 2A**). To investigate if the ubiquitinated form of TAX1BP1 was also present on mitochondria, sucrose fractionation experiments were performed. After sucrose fractionation, mitochondrial proteins (i.e: TOM20, VDAC and CS [citrate synthase]) were highly enriched in the mitochondria fraction (MTF) (**Figure 2B**). Under no stress conditions, TAX1BP1 was enriched in the membrane fraction (MMF), suggesting that TAX1BP1 could localize directly to membranous compartments under basal conditions. However, upon short mitochondrial depolarization, TAX1BP1 was enriched in the MTF, also in the ubiquitinated form (**Figure 2B**). These data indicated that ubiquitinated TAX1BP1 localized on depolarized mitochondria where it may contribute to parkin-dependent mitophagy.

Loss of the TAX1BP1 ubiquitinatable domain reduces clearance of ubiquitinated TOM20

TAX1BP1 is one of the autophagy receptors capable of recognizing ubiquitinated mitochondria and triggering its selective degradation during PINK1/parkin mitophagy [19]. However, among the identified five selective autophagy receptors involved in mitophagy, OPTN, NDP52 and TAX1BP1 are characterized as the only ones capable of inducing mitophagy on their own, with TAX1BP1 being the least efficient of the three [11]. Our results are in accordance with the hypothesis of TAX1BP1 having a minor role amongst other

autophagy receptors, as siRNA mediated knock-down of TAX1BP1 in WT-parkin HeLa cells

did not abolish the degradation of the outer mitochondrial proteins TOM20 and VDAC (**Fig.**

S2A). On the other hand, overexpression of TAX1BP1 had a positive effect on mitophagy,

as we observed an increased mitochondrial clustering upon short depolarization (**Fig. S2B**).

These results confirmed the involvement of TAX1BP1 in parkin-mediated mitophagy and

strengthened the notion of TAX1BP1 having a minor role when all other autophagy receptors

are present. Thus, in order to be able to characterize TAX1BP1 functionality isolated in a

null autophagy receptor background, HeLa cells lacking all five major autophagy receptors

were used (from now on, referred as 5KO HeLa) [11].

To understand the relevance of the identified TAX1BP1 ubiquitination events, lysine

substitution (TAX1BP1^{K549R}) and deletion (TAX1BP1ΔCC3) mutants were generated

(**Figure 3A**). Here, 5KO HeLa cells were transiently co-transfected with WT-parkin

combined with the indicated TAX1BP1 mutants and mitophagy was induced via CCCP-

mediated depolarization at the indicated time points. Upon co-expression of WT-parkin and

WT or TAX1BP1^{K549R} after early mitochondrial depolarization (4 h CCCP), TAX1BP1

ubiquitination was observed, even in the absence of all other four complementary autophagy

receptors (**Figure 3B**). In contrast, ubiquitination was reduced upon expression of

TAX1BP1ΔCC3 under the same conditions (**Figure 3B-C**). Likewise, TAX1BP1ΔCC3

ubiquitination was significantly reduced in 5KO HeLa stably expressing parkin (from now

on 5KO-PRKN HeLa) (**Figure 3D-E**). In ubiquitin pulldown experiments with 5KO-PRKN

cells, positive controls such as Ub-TOM20, pS65-Ub and total-Ub were captured upon short

CCCP-mediated depolarization but not under basal or control bead conditions (**Figure 3D**).

Importantly, when analyzing TAX1BP1 ubiquitination status in pulldown experiments,

additional higher molecular bands were observed for both WT and TAX1BP1[K549R], while

only a residual smear after long exposure was observed for TAX1BP1ΔCC3 (**Figure 3D**).

Further quantification of these identified higher molecular bands after long TAX1BP1

exposure revealed a significant ubiquitination reduction in TAX1BP1ΔCC3 only (**Figure**

3E). While we cannot discard that residual ubiquitination also occurs in TAX1BP1ΔCC3 -as

it is evidenced by the residual smear observed upon depolarization-, a reduced ubiquitination

signal is observed in TAX1BP1ΔCC3 when similar TAX1BP1 intensity bands are compared

(**Figure 3D**, red vs. blue arrowheads), and this effect is further confirmed upon quantification

(**Figure 3E**). The effects observed for TAX1BP1ΔCC3, contrast with the ubiquitination

status observed for TAX1BP1[K549R], which did not show any significant difference when

compared to WT (**Figure 3D-E**). Overall, these data indicate that most of the TAX1BP1

ubiquitination events observed upon mitochondria depolarization were localized at the CC3

domain of the autophagy receptor, but were not restricted to K549. It is important to note that

the CC3 domain of TAX1BP1 is particularly lysine-rich compared to CC1 or CC2 and that

the lysines encoded in the CC3 domain of TAX1BP1 are highly conserved across species

(**Fig. S3**). Thus, it is very possible that the sole substitution of a single lysine (i.e: K549R) in

this domain is not enough to have an impact on the total TAX1BP1 ubiquitination levels, as

many other lysines could be alternatively ubiquitinated in this condition.

Next, we focused on investigating if and how the ubiquitination status of TAX1BP1 could

affect the degradation of specific mitochondrial proteins. For this purpose, we quantified

protein levels of the outer and inner mitochondrial proteins: TOM20, VDAC, CS and COX4

(cytochrome c oxidase subunit 4I1) upon co-expression of WT-Parkin and the different

TAX1BP1 mutants after CCCP-mediated depolarization. Under control conditions, where

neither TAX1BP1 nor parkin were expressed, protein levels of all analyzed proteins (VDAC,

TOM20, CS and COX4) were not significantly reduced after 4 h of CCCP (**Figure 3B and**

3F-I). As expected, upon expression of parkin alone or together with WT TAX1BP, an

increase in OMM protein degradation was observed (i.e: TOM20 and VDAC) while inner

mitochondrial proteins (COX4 and CS) remained unaffected after 4 h of CCCP, when

compared to empty controls (**Figure 3F-I**). These results are in agreement with previous

studies indicating that the degradation of inner mitochondrial targets in a parkin-expressing

HeLa cell model, require longer depolarization exposure [9,17]. Expression of

TAX1BP1^{K549R} did not have any impact on VDAC or TOM20 degradation (**Figure 3F-G**).

VDAC degradation was increased upon addition of WT or any TAX1BP1 mutant, when

compared to parkin transfection alone. However, changes were found to not be significant,

suggesting that the autophagy receptor may not be required for the elimination of this

particular OMM protein (**Figure 3B and F**). Similarly, no significant changes in TOM20

degradation were observed upon TAX1BP1 over-expression, as TOM20 degradation was

already enhanced with parkin overexpression alone (**Figure 3B and G**), in agreement with

previous studies [11]. However, despite no observed changes in protein degradation under

these conditions, a significant increase in Ub-TOM20 was identified in TAX1BP1ΔCC3

(**Figure 3G and J**). Thus, these data suggest that the loss of the CC3 ubiquitinatable domain

on TAX1BP1 (TAX1BP1ΔCC3) has an impact on the targeting of ubiquitinated TOM20,

potentially delaying TOM20 degradation.

Modulation of the ubiquitination pattern of TAX1BP1 has mild effects on its translocation

behavior to mitochondria

The engulfment of mitochondrial material inside autophagosomes is a step that depends on

autophagy receptor functionality and that allows: (i) the recognition of ubiquitinated

mitochondrial proteins and (ii) the recruitment of autophagy effectors, which will promote

autophagosome biogenesis and eventually allow cargo engulfment [20]. Considering this, we

hypothesized that the slight defects observed in mitochondrial degradation could be due to a

delay in the TAX1BP1-dependent recognition of mitochondrial material. For this purpose,

TAX1BP1 translocation behavior and mitochondria clustering were analyzed.

First, basic immunofluorescence was used to visualize TAX1BP1 translocation on

mitochondrial material upon co-expression of WT-parkin (**Figure 4A and Fig. S4A**). Under

basal conditions, puncta-like structures were observed for all three TAX1BP1 constructs

(**Fig. S4A**). Short mitochondria depolarization induced the translocation of all three

TAX1BP1 constructs (**Figure 4A**), indicating that both TAX1BP1^{K549R} and TAX1BP1ΔCC3

were capable of recognizing depolarized mitochondria. However, TAX1BP1ΔCC3

translocation to mitochondria was rather localized in specific regions of the SSBP1 cluster

and, unlike parkin translocation, did not co-localize with most of SSBP1 material in some

cases (**Figure 4A, zoom panel iv**).

Next, mitochondrial morphology analysis was performed by quantifying mitochondria area

after depolarization. In this case, given that matrix protein degradation was not significantly

affected after short depolarization (**Figure 3B and H**), the matrix protein SSBP1 was used

for mitochondrial area quantification by using CellProfiler as previously performed [21].

Here, after short depolarization and co-expression of parkin with WT TAX1BP1, we

observed a strong reduction of SSBP1 area when compared to control conditions, indicative

of mitochondrial clustering and successful mitophagy (**Fig. S4B**). Importantly, while a

reduction of mitochondrial area was also observed with parkin expression only,

mitochondrial clustering was not as prominent as when WT TAX1BP1 was co-expressed

(**Fig. S4B**), indicative of delayed or impaired mitophagy in absence of the autophagy

receptor. When analyzing mitochondrial clustering upon co-expression of WT and mutant

TAX1BP1, an increase in SSBP1 area and perimeter was observed for TAX1BP1ΔCC3

while no changes were observed for TAX1BP1^{K549R} (**Figure 4B-D**). These data indicated

that TAX1BP1ΔCC3 had a delayed efficacy in promoting mitophagy.

Above results seem to indicate that the recognition of depolarized mitochondria is not

impaired by any of the mutants, as both TAX1BP1^{K549R} and TAX1BP1ΔCC3 were co-

localizing with mitochondrial material upon depolarization (**Figure 4A**). However, upon

expression of TAX1BP1ΔCC3 and short depolarization, we observed an increase in Ub-

TOM20 (**Figure 3B and J**) as well as a general delay in mitochondrial clustering, indicated

by an increase in SSBP1 foci area and perimeter (**Figure 4B-D**). Additionally, a partial

TAX1BP1 translocation to specific SSBP1 cluster regions was observed (**Figure 4A, zoom**

panel iv). Thus, we hypothesized that the observed TAX1BP1ΔCC3 delay in mitophagy

progression may be related to TAX1BP1 capability to recognize SSBP1 clusters. For this

purpose, the distribution of TAX1BP1 on depolarized mitochondria material was analyzed

with CellProfiler. In general, all TAX1BP1 constructs seemed to recognize SSBP1 material

in a similar manner, with distribution percentages ranging from 35-45% of SSBP1 area (**Fig.**

S4C). While no significant differences were observed between WT and TAX1BP1 mutants,

a slight reduction of the mitochondrial area recognized by TAX1BP1ΔCC3 was observed,

when compared to WT and TAX1BP1^{K549R} (**Fig. S4C**). To exclude that the observed

differences in TAX1BP1 mitochondrial distribution could be due to distinct mitophagy

progression stages, parkin distribution on mitochondrial material was additionally analyzed.

In this case, no overall differences were observed upon transfection of WT or TAX1BP1

mutants, as parkin-dependent recognition of SSBP1 area was found between 60-65% for all

cases (**Fig. S4D**). We additionally monitored the percentage of total TAX1BP1 involved in

the recognition of depolarized mitochondria. Here, no differences were observed for WT and

TAX1BP1 mutant, showing 45-50% of total TAX1BP1 to be engaged with depolarized

mitochondria (**Fig. S4E**). We further hypothesized that the differences of TAX1BP1

distribution on SSBP1 foci could be caused by differences in TAX1BP1 capability to

translocate to mitochondria. For this purpose, the recognized SSBP1 area (X) was plotted

against the percentage translocated TAX1BP1 on mitochondria (Y) and correlation analyses

were performed for each condition. Indeed, a significant correlation was observed for WT

TAX1BP1, with increasing TAX1BP1 being able to recognize 40-50% of SSBP1 area,

followed by a TAX1BP1 translocation decrease covering up to 60-80% of SSBP1 area

(**Figure 4E**). Correlation analysis was also significant for TAX1BP1^{K549R}, which behaved

similarly as WT but with an increased variability on the percentage of translocated

TAX1BP1^{K549R} (Y axis), after reaching a maximum of 70% translocated TAX1BP1 (**Figure**

4E). Surprisingly, unlike for WT and TAX1BP1^{K549R}, no significant correlation was

observed for TAX1BP1ΔCC3 (**Figure 4E**). These data suggested an inefficient capability of

TAX1BP1ΔCC3 to actively translocate to depolarized mitochondria and thus, distinct

binding behavior in comparison to WT TAX1BP1. Additionally, to discard that the

differences observed could be due to upstream differences in parkin translocation, a similar

correlation analysis was performed for parkin (**Fig. S4F**). Here, significant correlation was

found in all cases and no drastic differences were observed for parkin translocation upon

expression of all three TAX1BP1 constructs (**Fig. S4F**). In this case, approximately >80% of

SSBP1 was found recognized by parkin (**Fig. S4F**), indicating a comparable parkin

translocation and thus a similar mitophagy progression stage for all three TAX1BP1

expression cases.

Next, to validate the previously observed differences in TAX1BP1ΔCC3 binding behavior

in a context of stable parkin expression, TAX1BP1 mutants were transfected in 5KO-PRKN

HeLa cells and TAX1BP1 translocation was analyzed after 4 h of CCCP treatment. To

confirm mitophagy progression was at an advanced level, only cells with mitochondrial

COX4-positive clusters were taken into account. Here, cells were classified based on

TAX1BP1 translocation to mitochondria (complete vs. localized distribution) (**Figure 4F**).

Interestingly, a significant increase of cells with localized TAX1BP1 recognition was

observed under conditions of TAX1BP1^{K549R} and TAX1BP1ΔCC3 when compared to WT

(**Figure 4F**). These data indicated that the recognition of depolarized mitochondria was

slightly impaired in TAX1BP1^{K549R} and TAX1BP1ΔCC3 (**Figure 4E and F**).

Blocking autophagy flux in TAX1BP1ΔCC3 promotes the formation of enlarged

TAX1BP1 structures

Next, we sought to examine if the observed differences in mitophagy progression and binding

behavior of TAX1BP1^{K549R} and TAX1BP1ΔCC3 would translate in the capability of the

autophagy receptor to promote complete elimination of mitochondria. For this purpose,

mitochondrial degradation after extended depolarization was analyzed in 5KO-PRKN HeLa

cells transfected with WT and TAX1BP1 mutants. Interestingly, all TAX1BP1 constructs

were able to induce mitochondria degradation after extended CCCP treatment, as an

increased number of cells without mitochondria were observed in comparison to control

conditions (**Figure 5A and C**). However, when blocking the autophagy flux with

bafilomycin A (BafA), we could observe the presence of small and enlarged TAX1BP1$^+$

structures co-localizing with the matrix protein SSBP1 (**Figure 5B**). Importantly, the

percentage of cells harboring enlarged TAX1BP1$^+$ structures was significantly higher in

TAX1BP1ΔCC3 compared to WT or TAX1BP1^{K549R} (**Figure 5D**). In addition, TAX1BP1

ubiquitination status at that time point was validated by means of western blotting. Here, in

accordance to what was observed at an earlier stage of mitophagy (**Figure 3B**), ubiquitination

of TAX1BP1 was conserved upon autophagy flux blockage for WT and TAX1BP1^{K549R},

while a strong ubiquitination reduction was seen for TAX1BP1ΔCC3 under these conditions

(**Figure 5E**).

To confirm that the observed phenotype was not derived from CCCP side effects, additional

experiments were performed using a combination of Oligomycin and Antimycin A (OA) in

addition to the CCCP treatment (**Figure 5F-G and Figure S5).** Here, similarly as observed

in previous experiments, we could identify an increase in TAX1BP1ΔCC3 puncta size when

compared to WT or TAX1BP1^{K549R} under both treatment conditions (**Figure 5F**).

Importantly, the identified TAX1BP1ΔCC3 enlarged puncta were positive for HSP60

(**Figure 5G).** In addition, while TAX1BP1 puncta size significantly increased upon

depolarization for both WT and TAX1BP1 mutants, no size differences were observed under

BafA control conditions among WT TAX1BP1 and mutants (**Figure S5A-B**).

Next, CellProfiler was used to stratify TAX1BP1 puncta according to their size into enlarged,

intermediate or small puncta (**Figure S5C and D**). Such unbiased classification confirms the

previous eye-based quantification results (**Figure 5D**). Consistently, an increased

intracellular percentage of enlarged TAX1BP1 puncta was observed under TAX1BP1ΔCC3

expression (**Figure S5E**). Following TAX1BP1 puncta size-stratification, cells with more

than 25% of enlarged TAX1BP1 puncta were quantified upon CCCP and OA depolarization

(**Figure 5H**). Here, a significant increase of cells with enlarged TAX1BP1ΔCC3 puncta was

observed under CCCP treatment, while a less prominent increase was identified under OA

conditions (**Figure 5H**).

Overall, these data suggested that, while TAX1BP1ΔCC3 was able to promote mitochondria

elimination as TAX1BP1^{K549R} and WT TAX1BP1 after long depolarization, the degradative

mechanisms driven by TAX1BP1ΔCC3 could differ, as its expression induces a significantly

higher prevalence of enlarged TAX1BP1$^+$ structures carrying mitochondrial material.

Enlarged TAX1BP1ΔCC3 structures are not simple autophagosomes

Given the fact that TAX1BP1 is considered the least effective autophagy receptor to be able

to initiate mitophagy on its own in a 5KO-HeLa cell model [11], it was tempting to speculate

that the enlarged TAX1BP1$^+$ structures carrying mitochondrial material could be indicative

of mitochondrial degradation occurring via a distinct autophagy pathway. Thus, several

markers were used to investigate the nature of these enlarged TAX1BP1$^+$ structures.

Mitochondrial material was identified by using the matrix markers SSBP1 or HSP60 (60 kDa

heat shock protein, mitochondrial). As BafA is known to inhibit autophagic flux, the first

marker to test was the autophagosome marker LC3B. The enlarged TAX1BP1$^+$ HSP60$^+$

structures of WT, TAX1BP1^{K549R} and TAX1BP1ΔCC3 were LC3B$^+$ (**Figure 6A**). In all

cases, these structures seemed to be of a more complex nature, as they were composed of

multiple LC3B$^+$ entities structures across the Z plane (**Figure 6B, Fig. S6 and**

Supplementary videos1-4). These results suggest the enlarged TAX1BP1$^+$ structures were

formed by an accumulation of multiple autophagosomes (**Figure 6B, Fig. S6 and**

Supplementary videos1-4). On the other hand, all small TAX1BP1$^+$ HSP60$^+$ structures were

engaged in single cup-shaped LC3B$^+$ rings, suggesting they were indeed individual

autophagosomes (**Figure 7A**).

Later steps of autophagy-mediated degradation rely on the fusion of sealed autophagosomes

with lysosomes [19]. However, prior to autophagosome-lysosome fusion, autophagosome

maturation takes place and sealed autophagosomes may additionally fuse with endosomes,

generating a new degradative vesicles termed amphisome [22]. Thus, we decided to prove

for the late endosomal marker RAB7 (RAB7A, member RAS oncogene family), the member

of the retromer complex VPS35 (which is mutated in rare forms of PD) and the late

endosomal and marker CD63/LAMP-3 (CD63 antigen or Lysosomal-associated membrane

protein 3). Surprisingly, all enlarged TAX1BP1$^+$ structures carrying mitochondrial material

were surrounded by RAB7 and VPS35 signals, suggesting additional vesicles were engulfing

the identified accumulated LC3B$^+$ TAX1BP1$^+$ HSP60$^+$ structures (**Figure 6C and D**). These

patterns contrasted with the ones observed for CD63, which rather seemed to be part of the

cargo (**Figure 6E**). On the other hand, all regular sized TAX1BP1$^+$ structures were positive

for all tested endolysosomal markers (**Figure 7B-D),** with the only particularity that the

VPS35 and RAB7 rings engulfed single HSP60$^+$ and TAX1BP1$^+$ structures. These data

indicate that the enlarged TAX1BP1$^+$ structures are vesicles of endolysosomal nature filled

with multiple autophagosomes carrying mitochondrial cargo (amphisomes), while the

regular-sized TAX1BP1$^+$ structures correspond to individual vesicles or mature

autophagosomes.

Enlarged TAX1BP1ΔCC3$^+$ vesicles were also observed in parental HeLa cells with all

other complementary autophagy receptors

To confirm the relevance of the previously described phenotype in 5KO-PRKN HeLa cells,

similar experiments were performed in WT-Parkin HeLa cells endogenously expressing all

other complementary autophagy receptors (NDP52, OPTN, NBR1, SQSTM1/p62,

TAX1BP1) (**Fig. S7A**) [11]. Here, siRNA-based transient knockdown experiments were

followed by re-transfection of the desired TAX1BP1 constructs and subsequent CCCP-

mediated depolarization combined with BafA (**Figure 8A**). Confirmation of efficient

silencing was additionally performed via protein expression of total TAX1BP1 (**Figure 8A,**

lower panel). Overall, TAX1BP1 puncta in all conditions co-localized with the

mitochondrial marker HSP60 (**Figure 8A**). Surprisingly, the mean size of all observed

TAX1BP1$^+$ puncta per cell significantly increased upon siRNA knockdown and further

TAX1BP1 re-transfection in comparison to controls (**Figure 8B**). Indeed, when classifying

TAX1BP1 puncta size between small (<50 px^2), intermediate (≤150 px^2) or enlarged (> 150

px^2) (**Fig. S7B and C**), the majority of TAX1BP1$^+$ puncta were of small or intermediate

sizes, while larger puncta represented a much reduced percentage of the total TAX1BP1$^+$

puncta in all conditions (**Fig. S7C**). Of note, there was a 2-fold significant increase of the

intracellular percentage of larger TAX1BP1ΔCC3$^+$ puncta in comparison to WT TAX1BP1

conditions, reaching 5% of total TAX1BP1ΔCC3$^+$ puncta identified (**Figure 8C**). An

additional increase of the percentage of larger puncta profiles was observed upon WT

TAX1BP1 transfection when compared to transfection and silencing controls, suggesting the

overexpression of TAX1BP1 could increase its puncta size (**Figure 8C**). Furthermore, while

the number of cells with large TAX1BP1$^+$ puncta was slightly higher in TAX1BP1ΔCC3 in

comparison to re-transfected WT or TAX1BP1^{K549R}, the differences were only significant

when compared to siRNA controls (**Figure 8D**). Consistent with previous experiments in

5KO-PRKN Hela, TAX1BP1$^+$ enlarged puncta also co-localized with the mitochondrial

proteins HSP60 in WT-Parkin HeLa (**Figure 8A**). Next, to relate TAX1BP1 mitochondrial

removal efficiency upon expression of the different TAX1BP1 constructs in WT-Parkin

HeLa, co-localization analysis of HSP60 and TAX1BP1 was performed. First, the percentage

of HSP60 foci co-localizing with TAX1BP1$^+$ puncta of all sizes was analyzed (**Figure 8E**).

An increased percentage of HSP60 co-localizing with TAX1BP1 was observed upon

silencing and TAX1BP1 re-transfection in comparison to controls, with no significant

differences among all three TAX1BP1 conditions (**Figure 8E**). These data indicated that all

TAX1BP1 constructs enhanced the recognition of HSP60 material, in agreement with

previous experiments where overexpression of TAX1BP1 had a positive effect on

mitochondrial degradation (**Fig. S2B and C**). Next, HSP60 foci co-localizing with enlarged

TAX1BP1$^+$ puncta were analyzed (**Figure 8F**). Here, while there was a slight increase of the

percentage of HSP60 foci recognized by enlarged TAX1BP1ΔCC3$^+$ puncta, no significant

differences were found among WT and mutants under silencing conditions (**Figure 8F**).

However, it is worth noting that the mean area of the identified HSP60 foci co-localizing

with TAX1BP1ΔCC3 and TAX1BP1^{K549R} were larger than the ones observed in WT or

control conditions, while no differences were observed for WT compared to controls (**Figure**

8G).

In short, we observed an increased prevalence of enlarged TAX1BP1ΔCC3 structures

capable of recognizing increased amounts of HSP60 in comparison to WT. Similarly,

TAX1BP1^{K549R} enlarged structures are able to engage with increased amounts of HSP60,

however the prevalence of TAX1BP1^{K549R} enlarged structures is less abundant than

TAX1BP1ΔCC3, suggesting a milder phenotype for the site-substitution mutant. Thus, these

data indicated that the enlarged TAX1BP1ΔCC3^{+} structures observed in 5KO-PRKN HeLa

cells were of dominant nature, despite the partial rescue observed in WT-parkin HeLa, where

all complementary autophagy receptors are present.

Reduced autophagosome formation in TAX1BP1ΔCC3

Next, we hypothesized that the observed accumulation of enlarged amphisomes in

TAX1BP1ΔCC3 upon autophagy blockage could be due to defects in autophagy initiation

flux at earlier stages. In order to check this, short mitochondria depolarization was induced

by CCCP or OA treatment and co-localization of TAX1BP1 and LC3B on mitochondria was

further analyzed.

As expected, TAX1BP1-HSP60 co-localization upon CCCP or OA depolarization was

significantly increased when compared to control conditions (**Figure 9A and Fig. S8A-B**).

However, while all TAX1BP1 mutants were able to translocate to mitochondria upon

depolarization, a significant reduction on TAX1BP1-HSP60 overlap was observed for

TAX1BP1ΔCC3 while no changes in intensity correlation were detected (**Figure 9B-C**).

These data are in agreement with previous experiments where we observed a decrease in

TAX1BP1ΔCC3 overlap on COX4$^+$ or SSBP1$^+$ mitochondrial material (**Figure 4E-F**).

Along those lines, mitochondrial area was found increased in TAX1BP1ΔCC3 upon CCCP

treatment only (**Fig. S8C-D**), in agreement with previous experiments (**Figure 4B-D**).

Next, to further understand the potential defects in TAX1BP1ΔCC3-dependent autophagy

initiation, LC3B colocalization with HSP60 was analyzed (**Figure 9D-E** and **Fig. S8E**). As

expected, LC3B-HSP60 overlap upon CCCP or OA depolarization was significantly

increased when compared to control conditions (**Figure S8E**). On the other hand, no changes

in LC3B overlap with mitochondria were observed for any of the mutants in comparison to

WT TAX1BP1 (**Figure 9D**). However, there were significant changes in LC3B-HSP60

signal correlation for both TAX1BP1 mutants, as a significant increase in LC3B-HSP60

intensity correlation was observed for TAX1BP1ΔCC3 under both CCCP and OA

depolarization (**Figure 9E**). Changes in TAX1BP1^{K549R} were only significant under OA

treatment conditions (**Figure 9E**). Importantly, this was not a result of increased LC3B

intensity in TAX1BP1ΔCC3, as we observed a significant decrease in LC3B intensity for

both TAX1BP1 mutants in all conditions (**Figure 9F** and **Fig. S8F**). Similarly, there were

also important changes on the LC3B signal area (**Figure 9G**). As expected, TAX1BP1

expression alone induced a significant increase in LC3B area and this was accentuated upon

mitochondria depolarization (**Fig. S8G**). However, such increase was not as pronounced in

TAX1BP1ΔCC3 after depolarization, as we observed a significant reduction of the total

LC3B area in TAX1BP1ΔCC3 when compared to WT TAX1BP1 under both CCCP and OA

treatment conditions (**Figure 9G**).

These data strongly suggest that autophagosome formation induced by TAX1BP1ΔCC3 may

be compromised due to a reduced recognition of depolarized mitochondria under these

conditions, which results in a generally reduced autophagosome production.

Mass-spectrometry analysis of TAX1BP1ΔCC3 pulldown reveals defects in

intracellular trafficking processes

In order to get an idea of the global processes occurring upon TAX1BP1ΔCC3 expression

and try to pinpoint specific TAX1BP1 interactors that could be indicative of distinct non-

classical autophagic processes, mass-spectrometry analysis of TAX1BP1 precipitates was

performed. Here, HIS-tagged TAX1BP1 was co-precipitated under native conditions by

means of Ni-NTA pulldown and the pulled down fractions were further analyzed by mass-

spectrometry. Both conditions of depolarization only (16 h CCCP) or depolarization

combined with autophagic flux inhibition (16 h CCCP + BafA) were analyzed.

Identified protein groups and peptides showed a minor increase for cells expressing WT and

mutant TAX1BP1 when compared to the empty plasmid (**Fig. S9A-B**). Furthermore,

comparisons of the numbers of protein groups connected to mitochondria as well as the

autophagy and vesicular transport machineries revealed no sample specific differences (**Fig.**

S9A). The overall sample correlations were high and principal component analysis indicated

that the main differences between samples stemmed from variations between replicates (**Fig.**

S9C-D). As further quality control of the dataset, the number of identified peptides for

TAX1BP1 peptides and protein quantity were investigated. As expected, TAX1BP1 was

found to be absent in the control samples (**Fig. S9E-F**). Surprisingly, TAX1BP1^{K549R} quantity

values were lower in comparison to WT and TAX1BP1ΔCC3 (**Figure S9F**). Statistical

differences were analyzed between TAX1BP1ΔCC3 and WT upon CCCP treatment only

(**Fig. S10A and Table S2**) or in combination with BafA (**Figure 10A and Table S1**). Here,

when comparing TAX1BP1ΔCC3 with WT after extended depolarization only, an increased

co-precipitation of inner mitochondrial proteins was identified (**Fig. S10A**). These findings

were in line with the identified delayed mitochondrial degradation phenotype of

TAX1BP1ΔCC3 during earlier depolarization (**Figure 4B-D**). On the other hand, when

comparing TAX1BP1ΔCC3 and WT upon CCCP combined with BafA inhibition, multiple

proteins involved in vesicle-mediated transport, mitophagy or mitochondria localized

proteins were significantly regulated (**Figure 10A**).

Significantly up- and downregulated proteins were further analyzed for statistical

overrepresentation on GO biological processes for the condition of depolarization combined

with autophagy flux blockage only, using the panther database (www.pantherdb.org, version

19.0) [23]. Upon comparing the protein signatures interacting with TAX1BP1ΔCC3 against

WT under CCCP and BafA treatment combined, GO-term analysis indicated a

downregulation of several vesicle-related biological processes, some of them related to

retrograde transport or vesicle budding. Moreover, an upregulation of DNA replication,

repair and translation processes and an additional upregulation of mitophagy response

processes related to mitochondria depolarization were identified (**Figure 10B**).

When analyzing the protein signatures interacting with TAX1BP1ΔCC3 under these

conditions in more detail, a downregulated co-precipitation of TAX1BP1 with several

proteins involved in intracellular trafficking was identified (**Figure 10A**). Those included the

Rab-GTPases RAB8A, RAB14 and RAB1A, proteins involved in membrane organization

(TMEM33 [Transmembrane protein 33], VPS33A [vacuolar protein sorting-associated

protein 33A]) and other proteins involved in endosome budding from the Golgi apparatus

(ARF1 [ADP-ribosylation factor 1], ARF4 [ADP-ribosylation factor 4]). On the other hand,

an upregulated interaction with ubiquitin and inner mitochondrial proteins was found (**Figure**

10A). This later finding correlated with previous data showing an increased amount of

enlarged TAX1BP1ΔCC3 positive vesicles carrying increased amounts of mitochondrial

material (**Figure 5A and G and Fig. 8G**) as well as with the presence of Ub-TOM20 after

short depolarization (**Figure 3B and J**). Additionally, several vesicular-trafficking related

proteins were identified to increasingly co-precipitate with TAX1BP1ΔCC3. Those were:

three COPII vesicle-related proteins (SEC31A, SEC24D and SEC24A [protein transport

protein Sec24A/D/31A]) [24]; the endocytosis-related protein EPN1 (epsin-1) [25]; ERC1

(ELKS/Rab6-intercting/CAST family member 1) known to be involved in exocytosis

[26,27]; and the GTPase activating protein ACAP2 (Arf-GAP with coiled-coil, ANK repeat

and PH domain-containing protein 2), involved in endosomal trafficking [28] (**Figure 10A**).

Moreover, at a lower significance level (p-value<0.1), an upregulation of autophagy-related

protein WDR45B/WIPI3 (WD repeat domain phosphoinositide-interacting protein 3) as well

as the Rab7 GTPase activator TBC1D15 (TBC1 domain family member 15) were identified

(**Figure 10A**), both also implicated in mitophagy or general autophagy [29–32]. Importantly,

no significant interaction with ubiquitin or vesicle-related proteins was observed for

TAX1BP1[K549R] while an additional interaction with mitochondrial proteins was identified

(**Fig. S10B and Table S1**).

These data indicated that TAX1BP1ΔCC3 under these conditions was capable of

dysregulating vesicle-trafficking processes while being able to increasingly interact with

ubiquitinated mitochondrial targets. Overall, these findings reinforced the hypothesis that

removing the highly ubiquitinatable domain CC3 of TAX1BP1 impaired its mitochondrial

targeting efficacy which may additionally have an effect at later stages of autophagy, when

autophagosomes are formed and vesicular transport is essential for autophagosome

maturation.

Discussion

Complete mitochondria elimination during PINK1/Parkin mitophagy is subject to autophagy

receptor-dependent recognition of ubiquitinated mitochondria. In this context, five

autophagy receptors have been linked to PINK1/parkin mitophagy: NBR1, SQSTM1/p62,

OPTN, NDP52 and TAX1BP1 [33]. PTMs are known to play a regulatory role on autophagy

receptor functionality during selective autophagy [34]. Here we show that, in a context of

PINK1/parkin mitophagy, TAX1BP1 ubiquitination at its CC3 domain can regulate the

delivery of depolarized mitochondria to either a classical autophagy-dependent degradative

pathway or to an alternative pathway that is more dependent on the endolysosomal

machinery, without dramatically compromising the elimination of defective mitochondria.

TAX1BP1 ubiquitination at K549 or other lysines within its CC3 domain have been

described under similar mitochondria depolarizing conditions as the ones reported in this

study (**Figure 1A-B**) [18,35], strongly suggesting a regulatory role for this PTM. Indeed, we

could observe that TAX1BP1 ubiquitination did not direct the autophagy receptor for early

proteasomal degradation (**Fig. S1A**) but instead, ubiquitinated TAX1BP1 was found enriched

in mitochondria fractions upon short depolarization (**Figure 2B**). The limited effects of the

K549 point mutation suggests complexity and redundance in the TAX1BP1 ubiquitination

pattern. This study relies on the deletion of the whole CC3 domain to understand the role of

TAX1BP1 ubiquitination and the data presented here indicate that the CC3 deletion does not

impair the overall functionality of the autophagy adaptor. However, we cannot discard that

the entire CC3 domain deletion subtly perturbs TAX1BP1 structure, accounting for potential

effects beyond ubiquitination. Nevertheless, the CC3 domain is richer in lysins in comparison

to the other coiled-coiled regions of TAX1BP1, and the lysines within the CC3 domain are

highly conserved across mammalian species (**Fig. S3**). Indeed, when deleting the CC3

domain of TAX1BP1, a clear decrease in total TAX1BP1 ubiquitination was observed, both

after short (4 h CCCP) and extended depolarization combined with autophagy flux inhibition

(16 h CCCP + BafA) (**Figure 3B-E, Figure 5E and Fig. S2C**). Interestingly, when checking

TAX1BP1 predicted structure in AlphaFold [36], the structure of the regions connecting the

different CC domains and the ubiquitin binding domain (UBD) with the CC3 are of lowest

confidence (**Figure 11A**). This could be indicative of a highly dynamic behavior of those

regions, which could in turn influence the orientation of the distinct CC and UBD regions.

Thus, while there is no evidence for a potential structural interaction between the CC3 and

UBD regions based on the AlphaFold model, it is tempting to speculate that the ubiquitination

of multiple lysines at the CC3 could influence the orientation or availability of the UBD

region (**Figure 11B**).

Abolishing TAX1BP1 ubiquitination at CC3 domain by deleting its domain did not impair

the degradation of OMM proteins but had a negative impact in the targeting of Ub-TOM20

(**Figure 3B, F-J**) and promoted a slower clustering of mitochondria after short depolarization

(**Figure 4A-D**). Moreover, while CC3 ubiquitination was proven not essential for TAX1BP1

translocation to mitochondria (**Figure 4A**), a lack of it did significantly reduce the

distribution of the autophagy receptor on CCCP- and OA-depolarized mitochondria (**Figure**

4E-F and Figure 9A-B). Thus, suggesting that the delayed mitochondrial degradation and

clustering of TAX1BP1ΔCC3 was possibly due to an impairment on depolarized

mitochondrial recognition. One of the first reports describing TAX1BP1 domain structure

and its first links to immunity-related functions, described three different coiled-coil domains

localized between the now known SKITCH and UBA domains [37]. Interestingly, Ling and

Goeddel identified the CC1 and CC2 domains to be essential for TAX1BP1

homodimerization under basal conditions, while the region flanking the CC3 domain seemed

to play a minor role in its assembly [37]. Similarly to TAX1BP1, the autophagy receptor

NDP52 also possesses a central CC region recently identified to be crucial for NDP52

homodimerization in the context of mitophagy [38,39]. Based on this, it is tempting to

speculate that the differences observed in TAX1BP1ΔCC3 distribution on mitochondrial

material might be due to a reduced homodimerization capacities. However, given that the

recognition of depolarized mitochondria was significantly reduced in TAX1BP1ΔCC3

(**Figure 4E-F and 9A-B**) and this phenomenon was accompanied by a reduced formation of

LC3B$^+$ vesicles, as indicated by a reduced LC3B intensity and total LC3B area in

TAX1BP1ΔCC3 (**Figure 9F-G**), it seems possible that the delivery of defective

mitochondria in TAX1BP1ΔCC3 could be compromised or delayed from this step on. Along

these lines, our study also reveals that, even though the lack of ubiquitination at CC3 does

not impair complete mitochondrial degradation (**Figure 5C**), the degradative pathways

governing this process may differ between WT and TAX1BP1ΔCC3, as an increased

presence of enlarged degradation vesicles was observed in TAX1BP1ΔCC3 only (**Figure 5**

and Fig. S5), even under conditions where all other autophagy receptors were present

(**Figure 8 and Fig. S7C**). It is noteworthy that the here identified enlarged TAX1BP1$^+$

vesicles were prominently surrounded by the PD gene product VPS35. These findings may

suggest a functional link between the mitophagy pathway regulated by the most common

recessive gene products PINK1 and parkin together with the rare VPS35 gene product,

suggested to be involved in vesicle trafficking effects of the common dominant PD gene

product LRRK2 (Leucine-rich repeat serine/threonine-protein kinase 2). While this latter

connection in our work may be quite superficial, recent work by the group of Vandenberghe

W. and colleagues has made evident a remarkable functional link between VPS35-LRKK2

in the context of PINK1/parkin mitophagy [40]. The authors identify the novel VPS35

mutation p.D620N to not compromise the initiation steps of PINK1/parkin mitophagy but

significantly impair the recruitment of the autophagy receptor OPTN on depolarized

mitochondria [40]. While at this point it still remains yet to be elucidated if TAX1BP1

ubiquitination may have any impact on this molecular axis, our mass-spectrometry data

seems to confirm defects in vesicular trafficking, as many targets related to vesicle-mediated

transport were downregulated in TAX1BP1ΔCC3 (**Figure 10A**). Nevertheless, we could not

identify a unique endolysosomal marker specific for the enlarged degradative vesicles only

under these conditions, as all enlarged and small vesicles were positive for the same markers

(LC3B, VPS35, RAB7 and CD63) (**Figure 6 and Figure 7**). However, it is important to

emphasize that strong structural differences were clearly observed between small and

enlarged TAX1BP1$^+$ degradative vesicles, being the enlarged ones more prevalent in

TAX1BP1ΔCC3 than in WT. More precisely, the enlarged TAX1BP1ΔCC3$^+$ structures

contained numerous TAX1BP1$^+$ LC3B$^+$ vesicles carrying mitochondrial material (**Figure**

6A-B, Fig. S6 and Supplementary videos 1-4), where globally surrounded by RAB7$^+$ and

VPS35$^+$ vesicles (**Figure 6C-D**) and were additionally positive for CD63 (**Figure 6E**). On

the other hand, small vesicles found in all TAX1BP1 constructs consisted of single

TAX1BP1$^+$ LC3B$^+$ positive vesicles also carrying mitochondrial material and were equally

positive for all the tested endolysosomal markers (RAB7, CD63 and VPS35) (**Figure 7**).

These data seem to indicate that the degradation pathways involved in WT and ubiquitination

deprived TAX1BP1 are possibly of the same nature and the identified enlarged and small

vesicles are the result of autophagosomes fused with endosomes (amphisomes) or lysosomes

(autolysosomes) [22], potentially indicating that the complex enlarged TAX1BP1 structures

might be the result of defects in autophagosome maturation. Autophagosome fusion with

endosomes is known to be necessary for autophagosome maturation as it increases the pH of

the degradative vesicle and is considered a previous step for lysosome fusion [22]. On the

other hand, autophagy flux inhibition via BafA does not specifically hinder the fusion of

autophagosomes with lysosomes [41], thus we cannot discard that the observed enlarged

TAX1BP1 positive vesicles simply correspond to autolysosomes with more complex LC3B-

TAX1BP1-HSP60 cargo. It is important to note that TAX1BP1 has been reported to interact

with the cytoskeletal motor MYO6 (unconventional myosin-VI), known to be involved in

endocytic transport and to facilitate autophagosome maturation by interacting with the UBD

of TAX1BP1 [42]. Interestingly, compromising autophagosome maturation via transient

knock-down of MYO6, induced the formation of immature enlarged autophagosomes

positive for SQSTM1/p62 [42]. Another recent example of the relationship between

degradative vesicle size and endosome/autopagosome maturation was achieved by transient

knockdown of individual HOPS complex subunits, known to play an essential role in

endosome and autophagosome fusion with lysosomes [43]. Based on this, it seems reasonable

to think that the enlarged vesicles observed in TAX1BP1ΔCC3 are the result of impaired

autophagosome maturation driven by TAX1BP1ΔCC3 alone and further accentuated by

BafA treatment.

Additionally, it is important to emphasize that the formation of enlarged degradative vesicles

was not only observed in 5KO HeLa (**Figure 5B-D, F-H and Figure 6**) but also in parental

WT-Parkin HeLa with all complementary autophagy receptors (**Figure 8**). In this case, the

observed TAX1BP1ΔCC3 positive structures were generally smaller as the ones observed in

5KO HeLa. However, amongst the regularly sized TAX1BP1 positive structures, an

increased proportion of enlarged TAX1BP1ΔCC3 positive structures was observed (**Figure**

8C-D). These data indicate that while the presence of complementary autophagy receptors

can partially compensate for the observed phenotype, the TAX1BP1ΔCC3 dependent

formation of enlarged degradative vesicles can be considered of a dominant nature.

Moreover, the mitochondrial material recognized by the TAX1BP1$^+$ enlarged vesicles was

significantly increased in TAX1BP1ΔCC3 and TAX1BP1^{K549R} in comparison to WT (**Figure**

8G), strongly suggesting a decreased efficiency of ΔCC3, also under conditions of

complementary autophagy receptor expression. Thus, as larger TAX1BP1 vesicles were

observed in 5KO HeLa when compared to the ones seen in WT-Parkin HeLa, it seems

reasonable to think that the expression of complementary autophagy receptors can partially

compensate for the TAX1BP1-dependent formation of enlarged degradative vesicles. This is

in agreement with previous reports indicating TAX1BP1 as the least efficient autophagy

receptor to promote PINK1/parkin mitophagy in 5KO HeLa [11]. In addition, recent reports

using proximity-labelling approaches have indicated that TAX1BP1 can interact with other

autophagy receptors during mitophagy [44,45] suggesting the cooperation with more

efficient autophagy receptors could partially compensate for the observed phenotype in 5KO

HeLa.

Upon general GO-Term analysis of TAX1BP1ΔCC3 native pulldown (**Figure 10B**), a

significant decrease in interactions related to general vesicle trafficking was identified and is

in agreement with previous results showing an increased presence of TAX1BP1ΔCC3

enlarged vesicles under these conditions (**Figure 5B-D, F-H, Figure 6 and Fig. S5**). An

upregulated interaction with inner mitochondrial proteins and ubiquitin confirmed a

decreased efficiency of ΔCC3 in mitochondria degradation (**Figure 3C, Figure 4B-D and**

Figure 10A). However, expected TAX1BP1 interactors like RB1CC1/FIP200 (RB1-

inducible coiled-coil protein 1) or MYO6 could not be detected for neither of the mutants or

WT TAX1BP1. The increased interaction of TAX1BP1ΔCC3 with ubiquitin and the lack of

MYO6 co-precipitation may be indicative of delayed ubiquitinated mitochondria recognition,

as MYO6 binding is known to compete with ubiquitin at the UBD of TAX1BP1 [42]. On the

other hand, an upregulated interaction of TAX1BP1ΔCC3 with the components of COPII

vesicle coat complex SEC24A, SEC24D, SEC31A and the receptor-mediated endocytosis

EPN1 were identified [46–48] (**Figure 10A**). Similarly, albeit at a lower-significance level,

an increased interaction of two autophagy-related proteins involved in autophagosome

biogenesis (WDR45B and TBC1D15) was observed (**Figure 10A**). Importantly, COPII

vesicles are known to be important source of membrane lipids for the formation of

autophagosomes and indeed, recent reports have demonstrated that both SEC24 and SEC31

can interact with ATG9 vesicles, directly influencing autophagosome biogenesis [49–51].

Similarly, the Rab GTPase activating protein TBC1D15 is closely implicated with mitophagy

progression via its capacity to modulate active RAB7 levels at the OMM in favor of

autophagosome biogenesis [29,30]. On the other hand, WDR45B/WIPI3 is known to localize

at nascent autophagosomes as well as in late endosomes and thus seems to also play a role in

autophagosomes biogenesis and maturation [31,32]. Thus, given that TAX1BP1ΔCC3

induced (i) a delayed delivery of Ub-TOM20 (**Figure 3J**), (ii) a decrease in mitochondrial

clustering as well as a reduced recognition of defective mitochondria (**Figure 4B-F and**

Figure 9B) and (iii) promoted a reduced sequestration of LC3B$^+$ vesicles upon short

depolarization (**Figure 9G**), it seems reasonable to think that the interaction of

TAX1BP1ΔCC3 with the autophagosome biogenesis-related proteins at such later stage of

the pathway is probably due to a slower capability of the autophagy receptor to recruit

autophagy-related proteins in charge of autophagosome biogenesis. Indeed, decreased

interactions of TAX1BP1ΔCC3 were observed for the Rab GTPases RAB14, RAB1A and

RAB8A, the ER-transmembrane protein TMEM33 and the HOPS complex member VPS33

(**Figure 10A**). Rab GTPases are generally involved in endosomal vesicle trafficking but have

also been described to play a role in modulating autophagy: RAB1A has been described to

be essential for autophagosome formation [52,53] while RAB14 is implicated in

autophagosome maturation [53]. Similarly, VPS33A acts as a tether between

autophagosomes and lysosomes to facilitate their fusion [22,54]. Potential defects in

autophagosome maturation would be in agreement with the identified sustained ubiquitin

interaction, considering that MYO6 interaction competes with ubiquitin at the UBD of

TAX1BP1 [42]. Overall, our data suggest that the vesicle-trafficking defects identified in

TAX1BP1ΔCC3 may be due to impaired autophagosome biogenesis or maturation and likely

derive from a reduced recognition of depolarized mitochondria and a reduced sequestration

of LC3B$^+$ vesicles at earlier stages of mitophagy (**Figure 12**).

Conclusions

In this study, we identified the CC3 of TAX1BP1 as a highly ubiquitinated region during PINK1/parkin mitophagy which is not essential for complete mitochondria degradation. However, while reducing CC3 ubiquitination did not really have an impact on OMM protein degradation, other important defects related to mitophagy progression were identified: (i) reduced clearance of ubiquitinated TOM20, (ii) reduced mitochondrial clustering and (iii) reduced TAX1BP1 recognition of depolarized mitochondria, which translated in a (iv) localized presence of LC3B$^+$ vesicles on depolarized mitochondria. Blocking autophagy under these conditions had a significant impact on vesicular trafficking, as evidenced by the complex architecture of enlarged TAX1BP1$^+$ degradative vesicles which were highly prevalent in the absence of the ubiquitinatable CC3 domain. Our data strongly suggest that the identified vesicular trafficking defects may be driven by a delayed initiation of autophagosome biogenesis as well as an impairment with autophagosome maturation and are a reflection of the identified functional changes of TAX1BP1ΔCC3, which relies on an alternative degradation pathway that is substantially more dependent on the endolysosomal machinery (**Figure 12**).

Limitations of the study

This study provides evidence for an influence of TAX1BP1 ubiquitination on PINK1/parkin-mediated mitophagy in an idealized, designed HeLa cell culture model. To expand the relevance of the findings provided in this manuscript, more physiological cell models should be investigated, which would require sophisticated manipulations to study if and how

767　TAX1BP1 ubiquitination influences the clearance pathways of defective mitochondria.

768　Given that all the experiments carried out in this study relied on the overexpression of parkin

769　or TAX1BP1 (WT or mutants), future studies carried out for validation should also pay

770　attention on physiologically relevant expression levels of both the E3 ligase and the

771　autophagy adaptor.

Materials and methods

<u>Cell culture, treatments and transfections</u>

All cell lines were grown at 37°C and 5% CO_2 with high-glucose Dulbecco's modified eagle

medium (DMEM). WT-Parkin and 5KO-PRKN HeLa were supplemented with 0.7 mg/ml

geneticin (Gibco, 10131-027) to ensure stable parkin expression. HeLa cells stably

expressing WT or inactive mutant (C431A) were previously described [55]. To obtain 5KO

HeLa stably expressing WT-Parkin (5KO-PRKN HeLa), cells were transfected with BglII

linearized pcDNA3.1-3xFlag-Parkin for 24 hours and were further selected with 1.8 mg/ml

of Geneticin for 13 days. Upon selection, clones were screened for homogenous Parkin

expression. Mitophagy induction was achieved with 10 µM CCCP (Sigma, 555-60-2) for the

indicated time points. Proteasome and autophagy inhibition were induced with 10 µM

MG132 (Sigma, C2211) or 400 nM BafA (Chemcruz, sc-201550A), respectively. Inhibition

treatments were initiated half an hour before CCCP treatment and were maintained in

combination with CCCP for the indicated periods. Long CCCP treatment of 5KO-PRKN

HeLa was combined with 60 µM Q-VD-Oph (QVD, Selleckchem, S7311) to prevent cell

death. Transient transfection of the indicated constructs was performed with FuGENE6®

(Promega, E2692), following manufacturer's instructions. HeLa cells were transfected with

5 nM of siRNA for three consecutive days using Hiperfect (Qiagen, 301709). The following

siRNA were used: negative control (AllStars negative, Qiagen, SI03650318); TAX1BP1_7,

5'-CAGATCAATCAGCTAATAATA-3' (Qiagen, SI02781268); TAX1BP1_9, 5'-

AAGGGTCTTACTGAAGTAACA-3' (Qiagen, SI03036124); TAX1BP1_8, 5'-

CAGTCTTTGGCTTATCAATAA-3' (Qiagen, SI02781296).

<u>Expression constructs</u>

The construct pcDNA3.1-3xFlag-Parkin has been described previously [56] and was used to

transiently express parkin in 5KO HeLa and to generate the stable 5KO-PRKN as previously

explained. Full length WT, TAX1BP1^{K549R} and TAX1BP1ΔCC3 cDNA were amplified from

commercial vectors obtained from Origene (WT TAX1BP1, RC208233) or eZyvec on

demand (K549R and ΔCC3, Quote: D2022-10) and were sub-cloned into pCMV-6HIS vector

to ensure equal expression conditions. Obtained constructs were further sequenced prior to

use.

<u>Western blotting</u>

Upon treatment, cells were washed once with phosphate-buffered saline (PBS) and directly

harvested. For whole cell lysate samples, cell pellets were resuspended in urea lysis buffer

(10 mM Tris, pH 8.0, 100 mM NaH_2PO_4, and 8 M urea) after a short centrifugation step

(9727 x g for 1 minute [min] at 4°C). Lysates were passed through a 26-gauge needle to break

the DNA for at least 5 times. Whole cell lysate fractions were obtained after pelleting debris

at 21885 x g for 15 min at 4°C. Protein concentration was measured using the Bradford

Protein Assay Dye Reagent Concentrate (Bio-Rad, 5000006) following manufacturer's

instructions. Proteins were separated using 4-12% NuPage (Invitrogen, NP0329BOX), 3-8%

Tris-Acetate pre-casted (Invitrogen, EA0378BOX) or self-casted polyacrylamide gels and

were run at a fixed voltage of 80-100 V for at least 1,5 hours at room temperature (RT).

Proteins were transferred to polyvinylidene fluoride (PVDF) membranes (Merck,

IPVH00010) at constant voltage of 25 V for an overnight run or at 100 V for 2 hours,

respectively. Upon transfer, membranes were blocked with 5% non-fat dried milk or 5%

bovine serum albumin in Tris-buffered saline containing 0.1% Tween-20 (TBS-T).

Membranes were incubated with primary antibodies for 2 h at RT or overnight at 4ºC. After

three consecutive TBT-T washes of 5 min, membranes were incubated with the

corresponding secondary antibodies diluted in 5% of non-fat dried milk for 1 hour at RT.

Upon additional three TBT-T washes of 5 min, protein bands were visualized using

chemiluminescent-horseradish peroxidase (HRP) substrate (Millipore; WBKLS0500) and

Ultracruz autoradiography films (Santa Cruz Biotechnology; sc-201697). The following

primary antibodies were used: Actin (1:50000, Sigma Aldrich, A5441), Calnexin (1:10000,

Enzo Life Sciences, ADI-SPA-860), CS (1:6000, GeneTex, GTX110624), Flag-tag

(1:10000, Sigma-Aldrich, F1804), LAMP1 (1:10000, DSHB Hybridoma bank, Clone

H4A3), LC3B (1:10000, GeneTex, GT3612), LC3B (1:10000, Cell signaling, 835065),

MFN1 (1:10000, Abnova, H00055669-M04), NBR1 (1:10000, ProteinTech, 16004-1-AP),

NDP52 (1:10000, ProteinTech, 12229-1-AP), OPTN (1:10000, Atlas antibodies,

HPA003360), SQSTM1/p62 (1:10000, BD Biosciences, 610832), Parkin (1:50000, Cell

signaling, 4211S), phospho-Ubiquitin (Ser65) (1:1000, Cell Signaling, 52802), TAX1BP1

(1:1000, Novus Biologicals, NBP1-86662), TOM20 (1:30000, Santa Cruz, sc-17764),

TOM20 (1:6000, Proteintech, 11802-1-AP), Total Ubiquitin (1:10000, Millipore,

MAB1510), VDAC (1:10000, ProteinTech, 66345-1), VDAC (1:30000, Millipore,

AB10527), Vinculin (1:50000, Sigma Aldrich, V9131). The following secondary antibodies:

Anti-mouse HRP (1:10000, Amersham Pharmacia, 115-035-003), Anti-rabbit HRP

secondray (1:10000, Amersham Pharmacia, 111-035-003). Quantification of relative protein

levels was obtained via densitometry using ImageJ software. Briefly, individual protein

bands were normalized to Vinculin or Actin protein levels.

<u>Immunofluorescence</u>

HeLa cells were seeded in 0,1 mg/ml PDL (Sigma-Aldrich, P6407) pre-coated round

coverslips (Ø: 12 mm). After treatment, cells were fixed with 4% PFA for 20 min. After

fixation and three consecutive washes with PBS, permeabilization was performed with 1%

Triton-X for 5 min at room temperature (RT). After three washing steps with PBS, blocking

was performed at RT for 1 h with 10% fetal bovine serum or donkey serum (blocking buffer).

Afterwards, coverslips were incubated with the corresponding primary antibodies diluted in

blocking buffer for 2 h at RT. After three washing steps with PBS, coverslips were incubated

with secondary antibodies diluted in blocking buffer for 1 h at RT. Following three additional

washing steps with PBS, nuclei were stained with 2 µg/ml of Hoechst 33342 in PBS for 5

min at RT. After incubation, three additional washing steps with PBS were performed.

Coverslips were finally mounted with fluorescent mounting media (Agligent Dako, S3023).

Imaging was performed using an AxioImager microscope equipped with Apotome imaging

system (Carl Zeiss, Germany). Images were processed with the ZEN 2.3 lite software and

Image J. For LC3B immunostaining, cells were fixed with ice-cold methanol for 2-3 min.

Further immunostaining steps were performed as previously described. Image acquisition of

endogenous LC3B was realized with a Nikon Ti2e spinning disk confocal microscope. In

this case, images were processed using the NIS-Elements software (Nikon, Germany). The

following primary antibodies were used: CD63 (1:250, Antibodies online, ABIN94214),

COX4 (1:10000, Cell signaling, 4850), HIS-tag (1:500, Addgene, 184180), HSP60 (1:1000,

Novus Biologicals, NBP3-05536), LC3B (1:500, Cell signaling, 835065), Parkin(1:10000,

Cell signaling, 4211S), RAB7 (1:200, Cell signaling, 95746), TAX1BP1 (1:500, Novus

Biologicals, NBP1-86662), VPS35 (1:250, Novus Biologicals, NB100-1397). The following

secondary antibodies were used: Donkey anti-sheep Alexa Fluor 647 (1:1000, Invitrogen, A-

21448), Donkey anti-mouse Alexa Fluor 488 (1:1000, Invitrogen, A-21202), Donkey anti-

mouse Alexa Fluor 568 (1:1000, Invitrogen, A-10037), Donkey anti-rabbit Alexa Fluor 488

(1:1000, Invitrogen, A-21206), Donkey anti-rabbit Alexa Fluor 568 (1:1000, Invitrogen, A-

10042), Donkey anti-chicken Alexa Fluor 647 (1:1000, Invitrogen, A78952), Donkey anti-

goat Alexa Fluor 648 (1:1000, Invitrogen, A21447), Donkey anti-goat Alexa Fluor 488

(1:1000, Invitrogen, A11055).

Eye-based quantification of TAX1BP1 colocalization on COX4 was performed by taking

into account only TAX1BP1$^+$ cells with COX4 clumps after 4 h CCCP treatment, to account

for parkin-positive cells. TAX1BP1 behavior was classified between complete or localized

apposition of COX4 clump depending on its distribution. In situations where both TAX1BP1

behaviors were observed in the same TAX1BP1$^+$ cell, localized apposition of TAX1BP1 with

COX4 cluster was considered. Quantification of complete mitochondria elimination upon

TAX1BP1 transfection was achieved by only taking into account parkin-positive cells

(empty control) or cells positive for both parkin and TAX1BP1 (TAX1BP1 transfection) after

CCCP treatment. Cells were considered to be cleared of mitochondria where no SSBP1

puncta were observed. Quantification of small or enlarged TAX1BP1 positive structures after

extended depolarization combined with BafA inhibition was performed by taking into

account only parkin-positive cells. Cells were classified between containing small or

enlarged TAX1BP1 structures according size. In situations where both enlarged and small

TAX1BP1 structures were observed in the same TAX1BP1$^+$ cell, the cell was quantified as

containing enlarged TAX1BP1 structures.

<u>Mitochondria fractionation</u>

Mitochondria purification was performed as described in [57] with some modifications. Cell

pellets were first resuspended in mitochondrial buffer (MB) (210 mM mannitol, 70 mM

sucrose, 10 mM HEPES, 1 mM EDTA at pH 7.5) and further cell lysis was performed

mechanically using a glass homogenizer. After a first lysing series of 30 mechanical stokes,

cell homogenates were centrifuged at 600 x g for 10 min 4ºC. The remaining pellet was

resuspended with the remaining supernatant and two additional lysing series were performed,

with an in-between centrifugation step (600 x g for 10 min 4ºC). After the first and last

centrifugation, the remaining supernatant was recovered (corresponds to F1). The remaining

pellet was resuspended in 800 μl MB and was centrifuged at 700 x g for 10 min at 4ºC to

obtain the crude mitochondria pellet and the second supernatant fraction (F2). Crude

mitochondria pellets were resuspended in 250 μl MB and layered into a discontinuous

sucrose gradient formed by a 6 ml layer of 1.2 M sucrose over a 5 ml layer of 1.6 M sucrose

using ultra-clear tubes (14 x 89mm, Beckmann). Loaded gradients were centrifuged using a

Beckman ultracentrifuged equipped with a SW40Ti rotor at 96000 x g for 2 hours at 4ºC

(acceleration and deceleration parameters were set to 3). Upon centrifugation, bands of

interest were recovered. On the lighter sucrose layer, the band corresponding to membrane

fraction (MMF) was collected and at the layer interface, the band corresponding to enriched

mitochondria fraction (MTF) was collected. Upon collection, sucrose was diluted with four

volumes of MB and all fractions were pelleted at 13000 x g for 10 min at 4ºC. Pellets were

resuspended in urea buffer (10 mM Tris, pH 8.0, 100 mM NaH2PO4, and 8 M urea). Equal

protein amounts of each fraction were subjected to western blot analysis.

<u>Native Ni-NTA pulldown</u>

Cells were pelleted at 9727 x g for 1 min at 4°C. Pellets were lysed with native lysis buffer

(50 mM NaH_2PO_4, 300 mM NaCl, 0.2% NP-40, pH 8.0) complemented with 25 mM

Imidazole, protease (Roche, 04693116001) and phosphatase inhibitors (Roche,

4906845001), respectively. Lysed pellets were incubated on ice for 30 min. After incubation,

lysed pellets were centrifuged at 12000 x g for 15 min at 4°C and supernatant was collected

for total protein quantification. Ni-NTA spin columns (Qiagen, 31014) were used to purify

6HIS-tagged TAX1BP1 and its interacting proteins following provider instructions. Briefly,

Ni-NTA spin columns were first equilibrated using native lysis buffer with 25 mM Imidazole

and centrifuged for 2 min at 890 x g. After equilibration, 250 µg of total protein were loaded

on the columns and lysates were centrifuged at 270 x g for 10 min. Ni-NTA columns were

washed three times with lysis buffer containing 50 mM imidazole, centrifuged for 2 min at

890 x g. After washing, pulldown fractions were collected with elution buffer (50 mM

NaH_2PO_4, 300 mM NaCl, 0.2% NP-40, 500 mM Imidazole, pH 8.0) at 890 x g for 2 min

twice. Samples were sent directly in elution buffer for mass spectrometry analysis.

<u>Ubiquitin pulldown</u>

Cells were regularly washed with PBS before harvesting. After harvesting, cell pellets were

centrifuged at 9727 x g for 1 min at 4°C. Pellets were resuspended in RIPA buffer (50 mM

Tris/HCl, 150 mM NaCl, 1% NP-40, 0,5% deoxycholate, 0,2% SDS at pH 7.5) containing

phosphatase and protease inhibitors and left on ice. After 30 min, homogenates were

centrifuged at 18.000 x g for 15 min at 4°C and supernatant was recovered. Protein

concentration was measurement using the Pierce BCA assay kit (ThermoFischer). Ubiquitin

affinity beads (#UBA01B, Cytoskeleton) were used to pulldown ubiquitinated proteins. First,

both ubiquitin affinity beads and control beads (#CUB02B, Cytoskelton) were reconstituted

following manufacturer's instructions. For each pulldown reaction, 20 µl of ubiquitin or

control beads were aliquoted in a tube on ice and further washed twice with TBS-T.

Following bead wash, 200 µg of total protein were loaded on the beads. Pulldown reaction

tubes were filled with dilution buffer (50 mM Tris/HCl, 150 mM NaCl) containing protease

and phosphatase inhibitors to reach 1.5 ml of volume and were left in a rotating platform for

2 h at 4°C. After incubation, beads were collected and washed with dilution buffer. Three

washing steps were performed as follows: after buffer addition, tubes were allowed to rotate

for 5 min at 4°C, further pelleted at 5.000 x g for 1 minute at 4°C and supernatant was

removed. After the final wash, supernatant was completely removed with a Hamilton needle

and 50 µl of reducing buffer was added instead. Samples were boiled for 10 min at 95°C.

Bead pellets were further centrifuged at 5.000 x g for 1 minute at 4°C and supernatant was

collected. Samples were further analyzed via Western Blot.

<u>Mass spectrometry</u>

Eluates were mixed with 4x LDS sample buffer and proteins were separated by short SDS-

PAGE runs using NuPAGE NovexR Bis-Tris 4–12% Mini Gels (Invitrogen, NP0335BOX).

Protein bands were stained using QC Colloidal Coomassie Stain (Bio Rad Laboratories,

1610803) and whole sample lanes were excised. Gel pieces were further cut into small pieces

prior to in-gel digestion. Gel pieces were destained by incubation with 5 mM ABC in 50%

(v/v) ACN for 20 min three times and dehydrated using acetonitrile. Cysteine bonds were

reduced by incubation with 10 mM DTT for 45 min at 65°C before carbamidomethylation

by incubation with 55 mM iodoacetamide for 45 min at RT in the dark. Gel pieces were

washed two times and dehydrated as before, followed by digestion with trypsin overnight at

37°C. Peptides were extracted in three consecutive incubation steps with Extraction buffer A

(50% (v/v) ACN, 3% trifluoroacetic acid [TFA]), Extraction buffer B (80% (v/v) ACN, 0.5%

(v/v) TFA) and acetonitrile for 30 min each. Supernatants of each extraction were pooled,

and acetonitrile was evaporated by vacuum centrifugation. Peptides were purified using

StageTips [58]. Mass spectrometric analysis was performed using an Orbitrap Exploris 480

online (Thermo Fisher Scientific, Germany) coupled to an EASY nLC-1200 system (Thermo

Fisher Scientific, Germany). Peptides were separated on a 20 cm HPLC column with 75 μm

inner diameter (CoAnn Technologies, ICT36007508F-50) in-house packed with 1.9 μm

ReproSil-Pur C18 -AQ silica beads (Dr. Maisch HPLC GmbH, r119.aq.0001) and eluted by

solvent B in a 60 min linear gradient from 5–33% at a flow rate of 200 nL/min. Carry-over

was minimized by 30 min wash runs after each sample. Ionization was performed by ESI and

the mass spectrometer was operated in positive ion mode controlled by XCalibur (Thermo

Fisher Scientific). Acquisition of spectra was performed with a scan range of 350 – 950 m/z

and a resolution on the MS1 level of 60 000. MS2 scans were performed using DIA isolation

windows of 8 m/z with an overlap of 1 m/z (75 isolation windows), HCD collision energy of

30% and resolution of 15 000. The mass spectrometry proteomics data have been deposited

to the ProteomeXchange Consortium via the PRIDE partner repository [59] with the dataset

identifier PXD063982.

Mass spectrometry analysis

Raw files were processed using Spectronaut version 19 in directDIA+ mode using optimized identification settings as described before [60]. Spectra were predicted using a Uniprot *Homo sapiens* database (104 556 entries, downloaded on 2024/01/30). The data were exported using the protein pivot report using PG.ProteinGroups as a unique protein ID and the PG.Quantity as a quantification column. Data analysis was performed using Perseus version 2.0.5.0 [61], Microsoft Excel and R. Proteins were annotated functionally and by compartment using Gene Ontology Biological Processes and Gene Ontology Cellular Compartment databases (both downloaded on 2024/10/29) as well as for mitochondrial sub-compartments using MitoCarta3.0 (downloaded on 2021/02/24). To assess differences between WT and mutant TAX1BP1, student's T-tests were performed on protein intensities normalized to the protein quantities of the empty vector for both CCCP only and CCCP and BafA treated samples. Significantly up- and downregulated proteins ($p < 0.05$) were further analyzed for statistical overrepresentation on GO biological processes using the panther database (pantherdb.org, version 19.0) [23].

CellProfiler-based image analysis

The version 4.2.1 of the CellProfiler software was used [62]. All analyses rely on pooled single cell values obtained from at least three independent biological replicates. In general, individual cells were initially recognized with the nuclear staining and a cytoplasm area of 80-200 pixels (px) around the nucleus was further categorized as the cytoplasm, unless otherwise noted. Areas of interest per cell correspond to the established cytoplasm without the nucleus area.

For the analysis of mitochondrial clustering in WT-Parkin HeLa transfected with His-empty

or HIS-tagged TAX1BP1, mitochondrial material was recognized using SSBP1 signal as

primary object. SSBP1 objects were identified using a robust background two-class

thresholding method with lower and upper threshold bounds of 0.0-1.0, an adaptive window

of 50 and a size range of 6-500 px. Next, single SSBP1 objects that were in close contact

were merged (distance between objects=0 px). To account for only positively TAX1BP1

transfected cells, TAX1BP1 was recognized as primary objects using a minimum cross

entropy two-class thresholding method with a lower and upper threshold bounds of 0.0-1.0,

an adaptive window of size 30 and a size range of 10-40 px. Only cells with positive

TAX1BP1 signal (count TAX1BP1 objects>0) were taken into account for the TAX1BP1

overexpressing (OE) condition.

For the analysis of mitochondria clustering in 5KO HeLa co-expressing parkin as well as

WT-, K549R and TAX1BP1ΔCC3, the analysis was performed similarly as previously

described for the WT-Parkin HeLa with some variations. Mitochondrial material was

recognized as SSBP1 signal using a roust background two-class thresholding method with

lower and upper threshold bounds of 0.002-1.0, an adaptive window of size 50 and a size

range of 4-500 px. TAX1BP1 was recognized as primary objects using a minimum cross-

entropy two-class thresholding method with lower and upper threshold bounds of 0.0022-

1.0, an adaptive window of size 80 and a size range of 6-40 px. Similarly, Parkin expression

was monitored by recognizing parkin signal as primary object using a minimum cross-

entropy two-class thresholding method with lower and upper threshold bounds of 0.0015-

1.0, an adaptive window of size 50 and a size range of 10-40 px. Parkin objects were further

merged if they were in close contact (object distance=0 px). Only cells with positive

TAX1BP1 and parkin signal (mean object area > 0) were taken into account for this analysis.

Values of total mean SSBP1 area and mitochondrial area or perimeter normalized by cell size

(area or perimeter) are shown. For the analysis of TAX1BP1 distribution on SSBP1 clusters,

only cells with a mean TAX1BP1 intensity of $\geq$10 aiu (arbitrary intensity units) were taken

into account and the recognition method of TAX1BP1 and parkin signals was optimized. In

this case, TAX1BP1 and parkin were recognized as primary objects using a robust

background two-class thresholding method with lower and upper threshold bounds of

0.0035-1.0 (for TAX1BP1) or 0.002-1.0 (for parkin) with a size range recognition of 3-40 px

in both cases. Mitochondria were recognized using SSBP1 signal as described before for

5KO HeLa. To measure the overlapping area between TAX1BP1-SSBP1 and parkin-SSBP1,

total SSBP1 area was masked with TAX1BP1 or parkin merged objects and the percentage

of SSBP1 overlapped by each protein was calculated (% SSBP1 recognized). Similarly,

percentages of TAX1BP1 or parkin on mitochondria were calculated by considering which

portion of the total TAX1BP1 or parkin area was engaged with mitochondrial material (%

TAX1BP1 or parkin on SSBP1).

For the analysis of size classification of TAX1BP1$^+$ puncta and TAX1BP1-HSP60 material

overlap in WT-Parkin (a) and 5KO-PRKN HeLa (b), cell area was recognized using a

watershed-image method with nuclei as a reference point and TAX1BP1 or Parkin signal to

identify cell contours (minimum cross entropy method, threshold bounds: 0.00001-1).

TAX1BP1 puncta were recognized as primary objects using robust background method with

lower and upper threshold bounds of 0.000001-1.0 in (a) and 0.0015-1.0 in (b), an adaptive

window of size 15 and a size range of 4-40 px. In WT-Parkin Hela, only cells with TAX1BP1

mean intensity of 0.004 aiu were considered. To optimize the recognition of individual

puncta, clumped objects were distinguished by shape but dividing lines between clumped

objects were intensity-based. TAX1BP1 objects were merged when in contact with each

other (distance = 0 px), this step allowed to merge TAX1BP1 puncta that were in close

contact with other TAX1BP1 puncta and were particularly difficult to recognize as single

objects. Upon merging TAX1BP1 single objects, objects with an area form factor of 0.68-

0.8 were kept (form factor=1 for a perfectly circular object). This step allowed to keep

relatively circular-shaped objects (individual puncta) and discard merged objects that were

the result of multiple TAX1BP1 puncta. Thus, not all TAX1BP1 puncta were taken into

account for this analysis. After filtering, TAX1BP1 puncta were classified based on their

size: enlarged puncta (area>150 px^2), intermediate puncta (area≤150 px^2) and small puncta

(area<50 px^2). Mitochondrial material (HSP60) was recognized as primary object using a

robust background method with lower and upper threshold bounds of 0.001-1 and a size range

of 1-40 px in (a) and 0.0018-1.0 and a size range of 4-200 px in (b) and a size. HSP60 objects

were used to mask enlarged, intermediate and small TAX1BP1 puncta. In WT-Parkin Hela,

all TAX1BP1 puncta sizes were classified between HSP60-positive or HSP60-negative

according to their overlap. Percentages of HSP60 puncta co-localizing with overall or

enlarged TAX1BP1 puncta are shown as well as the corresponding HSP60 puncta area co-

localizing with enlarged TAX1BP1 puncta. For each analyzed image of this analysis, output

images were generated, showing the localization of the classified TAX1BP1 puncta, the cell

limits. Colocalization analysis in 5KO-PRKN was performed using the

MeasureColocalization module in CellProfiler, where Pearson's correlation coefficient of

TAX1BP1 and HSP60 signals were computed per cell.

Colocalization analysis of TAX1BP1-HSP60 and LC3B-HSP60 in 5KO-PRKN HeLa was

performed similarly as described before with some variations. To ensure the analysis only

included cells with similar TAX1BP1 expression, cells with a minimum mean integrated

intensity of 7 aiu were considered. A TAX1BP1 background region was established to filter

out cells with excessive TAX1BP1 levels (max. integrated intensity of 20 aiu overlapping

nucleus). HSP60 objects were recognized as previously described. TAX1BP1 objects were

recognized using robust background method with lower and upper threshold bounds of

0.0022-1.0 and a size of 1-100 px. LC3B was recognized using a three-class Otsu method.

Co-localization parameters were measured per each recognized cells for all three channels

and Pearson's coefficient as well as thresholded Mander's coefficient were obtained.

<u>Statistics</u>

All data is derived from at least three independent biological replicates unless otherwise

indicated. Statistical analysis and graph depiction of the data was performed using GraphPad

Prism 8.0.2. Statistical significance was established at $p<0.05$ unless otherwise indicated.

Significance was assessed by using a one-way ANOVA for multiple comparisons and

subsequent Tukey's post-hoc test unless otherwise indicated. For the correlation analysis of

TAX1BP1 binding to depolarized mitochondria, a Pearson correlation test was performed.

For analysis in which only two conditions were compared, significance was assessed with an

unpaired t-test analysis. The following significance indicators were used: ****: $p<0.001$,

***: $p<0.005$, **: $p<0.005$, *: $p<0.05$

1082

1083

Acknowledgements

PentaKO (5KO) HeLa cells were kindly provided by Dr. Chunxin (Black) Wang and Dr.

Richard J. Youle, NINDS. This work was supported by the DFG Research Training Group

2364 "MOMbrane", the Hertie Foundation and the German Center for Neurodegenerative

Diseases. We would also like to acknowledge Prof. Dr. Tassula Proikas-Cezanne for valuable

discussion and feedback. Lastly, we would also like to acknowledge Henrietta Lacks and her

family.

1091

1092   **Conflict of interest**

1093   The authors declare no conflict of interest.

1094

1095   **Data availability**

1096   All mass spectrometry proteomics data have been deposited to the ProteomeXchange

1097   Consortium via the PRIDE partner repository and are openly available in

1098   http://www.ebi.ac.uk/pride with the dataset identifier PXD063982.

1099

1100   **References**

1101   [1]   Narendra DP, Kane LA, Hauser DN, et al. p62/SQSTM1 is required for Parkin-

1102   induced mitochondrial clustering but not mitophagy; VDAC1 is dispensable for

1103   both. Autophagy 2010; 6(8):1090–106. doi: 10.4161/auto.6.8.13426

1104   [2]   Geisler S, Holmström KM, Skujat D, et al. PINK1/Parkin-mediated mitophagy is

1105   dependent on VDAC1 and p62/SQSTM1. Nat Cell Biol 2010; 12(2):119–31. doi:

1106   10.1038/ncb2012

1107   [3]   Matsuda N, Sato S, Shiba K, et al. PINK1 stabilized by mitochondrial depolarization

1108   recruits Parkin to damaged mitochondria and activates latent Parkin for mitophagy. J

1109   Cell Biol 2010; 189(2):211–21. doi: 10.1083/jcb.200910140

1110   [4]   Narendra DP, Jin SM, Tanaka A, et al. PINK1 is selectively stabilized on impaired

1111   mitochondria to activate Parkin. PLoS Biol 2010; 8(1). doi:

1112   10.1371/journal.pbio.1000298

1113   [5]   Kondapalli C, Kazlauskaite A, Zhang N, et al. PINK1 is activated by mitochondrial

1114   membrane potential depolarization and stimulates Parkin E3 ligase activity by

1115   phosphorylating Serine 65. Open Biol 2012; 2(MAY). doi: 10.1098/rsob.120080

1116   [6]   Shiba-Fukushima K, Imai Y, Yoshida S, et al. PINK1-mediated phosphorylation of

1117   the Parkin ubiquitin-like domain primes mitochondrial translocation of Parkin and

1118   regulates mitophagy. Sci Rep 2012; 2:1–8. doi: 10.1038/srep01002

1119   [7]   Koyano F, Okatsu K, Kosako H, et al. Ubiquitin is phosphorylated by PINK1 to

1120   activate parkin. Nature 2014; 510(7503):162–6. doi: 10.1038/nature13392

1121   [8]   Ordureau A, Heo JM, Duda DM, et al. Defining roles of PARKIN and ubiquitin

1122   phosphorylation by PINK1 in mitochondrial quality control using a ubiquitin

1123   replacement strategy. Proc Natl Acad Sci U S A 2015; 112(21):6637–42. doi:

1124   10.1073/pnas.1506593112

[9]   Lechado-Terradas A, Schepers S, Zittlau KI, et al. Parkin-dependent mitophagy

occurs via proteasome-dependent steps sequentially targeting separate mitochondrial

sub-compartments for autophagy. Autophagy Reports 2022; 1(1):576–602. doi:

10.1080/27694127.2022.2143214

[10] Heo JM, Ordureau A, Paulo JA, et al. The PINK1-PARKIN Mitochondrial

Ubiquitylation Pathway Drives a Program of OPTN/NDP52 Recruitment and TBK1

Activation to Promote Mitophagy. Mol Cell 2015; 60(1):7–20. doi:

10.1016/j.molcel.2015.08.016

[11] Lazarou M, Sliter DA, Kane LA, et al. The ubiquitin kinase PINK1 recruits

autophagy receptors to induce mitophagy. Nature 2015; 524(7565):309–14. doi:

10.1038/nature14893

[12] Richter B, Sliter DA, Herhaus L, et al. Phosphorylation of OPTN by TBK1 enhances

its binding to Ub chains and promotes selective autophagy of damaged

mitochondria. Proc Natl Acad Sci U S A 2016; 113(15):4039–44. doi:

10.1073/pnas.1523926113

[13] Yamano K, Kikuchi R, Kojima W, et al. Critical role of mitochondrial ubiquitination

and the OPTN ATG9A axis in mitophagy. J Cell Biol 2020; 219(9). doi:

10.1083/jcb.202001120

[14] Nguyen TN, Sawa-makarska J, Khuu G, et al. Unconventional initiation of

PINK1/Parkin mitophagy by Optineurin. Mol Cell 2023; 83(10):1693–709. doi:

10.1016/j.molcel.2023.04.021

[15] Turco E, Savova A, Gere F, et al. Reconstitution defines the roles of p62, NBR1 and

TAX1BP1 in ubiquitin condensate formation and autophagy initiation. Nat Commun

2021; 12(1):1–16. doi: 10.1038/s41467-021-25572-w

[16] Yamano K, Sawada M, Kikuchi R, et al. Optineurin provides a mitophagy contact

site for TBK1 activation. EMBO J 2024; 43(5):754–79. doi: 10.1038/s44318-024-

00036-1

[17] Zittlau KI, Lechado-Terradas A, Nalpas N, et al. Temporal Analysis of Protein

Ubiquitylation and Phosphorylation During Parkin-Dependent Mitophagy. Mol Cell

Proteomics 2022; 21(2):100191. doi: 10.1016/j.mcpro.2021.100191

[18] Rose CM, Isasa M, Ordureau A, et al. Highly Multiplexed Quantitative Mass

Spectrometry Analysis of Ubiquitylomes. Cell Syst 2016; 3(4):395-403.e4. doi:

10.1016/j.cels.2016.08.009

[19] Narendra DP, Youle RJ. The role of PINK1–Parkin in mitochondrial quality control.

Nat Cell Biol 2024; 26(October). doi: 10.1038/s41556-024-01513-9

[20] Adriaenssens E, Ferrari L, Martens S. Orchestration of selective autophagy by cargo

receptors. Curr Biol 2022; 32(24):R1357–71. doi: 10.1016/j.cub.2022.11.002

[21] Sharma K, Kishore A, Lechado-Terradas A, et al. A Novel PINK1 p.F385S Loss-of-

Function Mutation in an Indian Family with Parkinson's Disease. Mov Disord 2024;

39(7):1217–25. doi: 10.1002/mds.29792

[22] Zhao YG, Codogno P, Zhang H. Machinery, regulation and pathophysiological

implications of autophagosome maturation. Nat Rev Mol Cell Biol 2021;

22(11):733–50. doi: 10.1038/s41580-021-00392-4

[23] Mi H, Muruganujan A, Csagrande JT, et al. Large-scale gene function analysis with

PANTHER Classification System. Nat Protoc 2013; 8(8):1551–66. doi:

10.1038/nprot.2013.092

[24] Wang J, Davis S, Zhu M, et al. Autophagosome formation: Where the secretory and

autophagy pathways meet. Autophagy 2017; 13(5):973–4. doi:

10.1080/15548627.2017.1287657

[25] Bhattacharjee S, Lee Y, Zhu B, et al. Epsins in vascular development, function and

disease. Cell Mol Life Sci 2021; 78(3):833–42. doi: 10.1007/s00018-020-03642-4

[26] Hida Y, Ohtsuka T. CAST and ELKS proteins: Structural and functional

determinants of the presynaptic active zone. J Biochem 2010; 148(2):131–7. doi:

10.1093/jb/mvq065

[27] Ramella M, Ribolla LM, de Curtis I. Liquid–Liquid Phase Separation at the Plasma

Membrane–Cytosol Interface: Common Players in Adhesion, Motility, and Synaptic

Function. J Mol Biol 2022; 434(1):167228. doi: 10.1016/j.jmb.2021.167228

[28] Jackson TR, Brown FD, Nie Z, et al. ACAPs are arf6 GTPase-activating proteins

that function in the cell periphery. J Cell Biol 2000; 151(3):627–38. doi:

10.1083/jcb.151.3.627

[29] Yamano K, Fogel AI, Wang C, et al. Mitochondrial Rab GAPs govern

autophagosome biogenesis during mitophagy. Elife 2014; 3:1–24. doi:

10.7554/elife.01612

[30] Yamano K, Wang C, Sarraf SA, et al. Endosomal Rab cycles regulate Parkin-

mediated mitophagy. 2018; :1–32.

[31] Bakula D, Müller AJ, Zuleger T, et al. WIPI3 and WIPI4 β-propellers are scaffolds

for LKB1-AMPK-TSC signalling circuits in the control of autophagy. Nat Commun

2017; 8(May). doi: 10.1038/ncomms15637

[32] Ji C, Zhao H, Chen D, et al. β-propeller proteins WDR45 and WDR45B regulate

autophagosome maturation into autolysosomes in neural cells. Curr Biol 2021;

31(8):1666-1677.e6. doi: 10.1016/j.cub.2021.01.081

[33] Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy.

Nat Rev Mol Cell Biol 2018; 19(6):349–64. doi: 10.1038/s41580-018-0003-4

[34] Lechado-Terradas A, Zittlau KI, Macek B, et al. Regulation of mitochondrial cargo-

selective autophagy by posttranslational modifications. J Biol Chem 2021;

297(5):101339. doi: 10.1016/j.jbc.2021.101339

[35] Sarraf SA, Raman M, Guarani-Pereira V, et al. Landscape of the PARKIN-

dependent ubiquitylome in response to mitochondrial depolarization. Nature 2013;

496(7445):372–6. doi: 10.1038/nature12043

[36] Jumper J, Evans R, Pritzel A, et al. Highly accurate protein structure prediction with

AlphaFold. Nature 2021; 596(7873):583–9. doi: 10.1038/s41586-021-03819-2

[37] Ling L, Goeddel D V. T6BP , a TRAF6-interacting protein involved in IL-1

signaling. 2000; :2–7.

[38] Johansen T, Lamark T. Selective autophagy mediated by autophagic adapter

proteins. Autophagy 2011; 7(3):279–96. doi: 10.4161/auto.7.3.14487

[39] Kataura T, Otten EG, Rabanal-Ruiz Y, et al. NDP52 acts as a redox sensor in

PINK1/Parkin-mediated mitophagy. EMBO J 2023; 42(5):1–16. doi:

10.15252/embj.2022111372

[40]   Manders L, Heyninck T, Imberechts D, et al. VPS35 mutation inhibits

PINK1/parkin-mediated mitophagy via increased LRRK2 kinase activity. Brain

2025; (October):1–18. doi: 10.1093/brain/awaf414

[41]   Mauthe M, Orhon I, Rocchi C, et al. Chloroquine inhibits autophagic flux by

decreasing autophagosome-lysosome fusion. Autophagy 2018; 14(8):1435–55. doi:

10.1080/15548627.2018.1474314

[42]   Tumbarello DA, Waxse BJ, Arden SD, et al. Autophagy receptors link myosin VI to

autophagosomes to mediate Tom1-dependent autophagosome maturation and fusion

with the lysosome. Nat Cell Biol 2012; 14(10):1024–35. doi: 10.1038/ncb2589

[43]   der Beek J van, de Heus C, Sanza P, et al. Loss of the HOPS complex disrupts early-

to-late endosome transition, impairs endosomal recycling and induces accumulation

of amphisomes. Mol Biol Cell 2024; 35(3):1–24. doi: 10.1091/MBC.E23-08-0328

[44]   Zellner S, Schifferer M, Behrends C. Systematically defining selective autophagy

receptor-specific cargo using autophagosome content profiling. Mol Cell 2021;

81(6):1337-1354.e8. doi: 10.1016/j.molcel.2021.01.009

[45]   Heo JM, Harper NJ, Paulo JA, et al. Integrated proteogenetic analysis reveals the

landscape of a mitochondrial-autophagosome synapse during PARK2-dependent

mitophagy. Sci Adv 2019; 5(11). doi: 10.1126/sciadv.aay4624

[46]   Morinaka K, Koyama S, Nakashima S, et al. Epsin binds to the EH domain of POB1

and regulates receptor-mediated endocytosis. Oncogene 1999; 18(43):5915–22. doi:

10.1038/sj.onc.1202974

[47]   Stankewich MC, Stabach PR, Morrow JS. Human Sec31B: A family of new

mammalian orthologues of yeast Sec31p that associate with the COPII coat. J Cell

Sci 2006; 119(5):958–69. doi: 10.1242/jcs.02751

[48]   Mancias JD, Goldberg J. Structural basis of cargo membrane protein discrimination

by the human COPII coat machinery. EMBO J 2008; 27(21):2918–28. doi:

10.1038/emboj.2008.208

[49]   Hurley JH, Young LN. Mechanisms of autophagy initiation. Annu Rev Biochem

2017; 86:225–44. doi: 10.1146/annurev-biochem-061516-044820

[50]   Davis S, Wang J, Zhu M, et al. Sec24 phosphorylation regulates autophagosome

abundance during nutrient deprivation. Elife 2016; 5(NOVEMBER2016):1–22. doi:

10.7554/eLife.21167

[51]   Nie J, Ma S, Wu L, et al. SEC31a-ATG9a Interaction Mediates the Recruitment of

COPII Vesicles for Autophagosome Formation. Adv Sci 2024; 2405127:1–15. doi:

10.1002/advs.202405127

[52]   Gyurkovska V, Murtazina R, Zhao SF, et al. Dual function of Rab1A in secretion

and autophagy: hypervariable domain dependence. Life Sci Alliance 2023; 6(5):1–

16. doi: 10.26508/lsa.202201810

[53]   Haga K, Fukuda M. Comprehensive knockout analysis of the RAB family small

GTPases reveals an overlapping role of RAB2 and RAB14 in autophagosome

maturation. Autophagy 2024; 21(1):21–36. doi: 10.1080/15548627.2024.2374699

[54]   Wartosch L, Günesdogan U, Graham SC, et al. Recruitment of VPS33A to HOPS by

VPS16 Is Required for Lysosome Fusion with Endosomes and Autophagosomes.

Traffic 2015; 16(7):727–42. doi: 10.1111/tra.12283

[55]   Geisler S, Jäger L, Golombek S, et al. Ubiquitin-specific protease USP36

knockdown impairs Parkin-dependent mitophagy via downregulation of Beclin-1-

associated autophagy-related ATG14L. Exp Cell Res 2019; 384(2):111641. doi:

10.1016/j.yexcr.2019.111641

[56]   Geisler S, Vollmer S, Golombek S, et al. The ubiquitin-conjugating enzymes

UBE2N, UBE2L3 and UBE2D2/3 are essential for Parkin-dependent mitophagy. J

Cell Sci 2014; 127(15):3280–93. doi: 10.1242/jcs.146035

[57]   Alberio T, Pieroni L, Ronci M, et al. Toward the Standardization of Mitochondrial

Proteomics: The Italian Mitochondrial Human Proteome Project Initiative. J

Proteome Res 2017; 16(12):4319–29. doi: 10.1021/acs.jproteome.7b00350

[58]   Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-

fractionation and storage of peptides for proteomics using StageTips. Nat Protoc

2007; 2(8):1896–906. doi: 10.1038/nprot.2007.261

[59]   Perez-Riverol Y, Bandla C, Kundu DJ, et al. The PRIDE database at 20 years: 2025

update. Nucleic Acids Res 2025; 53(D1):D543–53. doi: 10.1093/nar/gkae1011

[60]   Bruderer R, Bernhardt OM, Gandhi T, et al. Optimization of experimental

parameters in data-independent mass spectrometry significantly increases depth and

reproducibility of results. Mol Cell Proteomics 2017; 16(12):2296–309. doi:

10.1074/mcp.RA117.000314

[61]   Tyanova S, Temu T, Sinitcyn P, et al. The Perseus computational platform for

comprehensive analysis of (prote)omics data. Nat Methods 2016; 13(9):731–40. doi:

10.1038/nmeth.3901

[62]   Stirling DR, Swain-Bowden MJ, Lucas AM, et al. CellProfiler 4: improvements in

1280   speed, utility and usability. BMC Bioinformatics 2021; 22(1):1–11. doi:

1281   10.1186/s12859-021-04344-9

1282   [63]   Meng EC, Goddard TD, Pettersen EF, et al. UCSF ChimeraX: Tools for structure

1283   building and analysis. Protein Sci 2023; 32(11):1–13. doi: 10.1002/pro.4792

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

 Figure legends

 Figure 1: Ubiquitination of TAX1BP1 upon CCCP-induced mitophagy in HeLa cells

 stably expressing WT- or C431A-Parkin. (**A**) Schematic representation of TAX1BP1

 domain structure. Identified ubiquitinated lysine residues from the ubiquitylome mass-

 spectrometry screen are indicated in red. (**B**) Normalized Gly-Gly peptide signals of the

 identified ubiquitinated residues in both WT- and C431A-parkin HeLa cell lines during

 mitophagy induction at the indicated time points. (**C**) Western blot detection of the indicated

 proteins in whole cell lysates from HeLa cells stably expressing WT- or catalytically inactive

 C431A-parkin treated with CCCP for the indicated times. TAX1BP1 band shift peaking at 4

 h confirms ubiquitination. (**D**) Western blots showing protein signals of 5% of input or 10%

 of eluted pulldown samples performed in WT-Parkin Hela depolarized with CCCP for 4 h or

 under control conditions (DMSO). Pulldown samples are shown for three independent

 replicate experiments (Rep).

 Figure 2: TAX1BP1 translocation to mitochondria during CCCP-induced mitophagy.

 (**A**) Representative immunofluorescence images showing endogenous TAX1BP1 (yellow),

 and the inner mitochondrial marker SSBP1 (cyan) at basal levels or after mitophagy induction

 (10 μM CCCP, 2h). Scale bars: 5 μm (**B**) Subcellular fractionation of WT-Parkin HeLa cells

 under basal conditions or after short depolarization, where translocation of parkin and

 ubiquitinated TAX1BP1 to mitochondria can be observed. WCL: whole cell lysate, F1:

 fraction 1 corresponds to the supernatant recovered after the first centrifugation series, F2:

 fraction 2 corresponds to the supernatant recovered after the second centrifugation step in

1321 which the crude mitochondria pellet were obtained, CM: crude mitochondria fraction, MMF:

1322 membrane enriched fraction, MTF: mitochondrial enriched fraction.

1323

Figure 3: Over-expression of TAX1BP1ΔCC3 leads to changes in its ubiquitination

pattern and reduces clearance of ubiquitinated TOM20. (**A**) Schematic representation of

the HIS-tagged TAX1BP1 constructs harboring residue-specific or deletion mutations at the

CC3 domain. H: 6xHis-tag. (**B**) Mitophagy time course in 5KO-HeLa cells transiently

expressing WT, TAX1BP1^{K549R} and TAX1BP1ΔCC3 with or without Flag-tagged WT-

Parkin co-transfection. Ø: empty control plasmids. Color-coded legends are shown for each

transfection condition. (**C**) Relative levels of ubiquitinated (Ub-) TAX1BP1 after the

indicated CCCP treatments. Quantification corresponds to the Ub_n-TAX1BP1 smear

indicated in panel (B). (**D**) Ubiquitin pulldown western blots showing protein signals of 5%

of input or 10% of eluted samples performed in 5KO-PRKN HeLa depolarized for 4 h of

CCCP compared to control conditions (DMSO). Red and blue arrowheads indicate presence

or absence of poly-ubiquitinated TAX1BP1 bands, respectively. (**E**) Quantification of Ub-

TAX1BP1 bands obtained in Ubiquitin pulldown shown in (D, lower panel). Ubiquitination

levels were normalized to TAX1BP1 input signal for each transfection condition. Further,

normalization to control beads and to DMSO control was performed. For this comparison,

unpaired t-test was used to assess significance. (**F-I**) Relative protein levels of the following

mitochondrial markers: VDAC (F), TOM20 (G), CS (H) and COX4 (I). Quantifications

correspond to protein levels in 5KO Hela shown in (B). (**J**) Relative levels of ubiquitinated

(Ub-) TOM20 after the indicated CCCP treatment shown in (B). All protein quantifications

are derived from at least three independent experiments. Protein signal ratios in (F-I) were

obtained after normalization of protein signal with loading control (vinculin) and subsequent

normalization with protein values at basal level. Ubiquitination levels shown in panels (C)

and (J) were normalized to loading and to TOM20 or TAX1BP1 levels, respectively. Ratios

relative to empty transfection control are shown for TOM20 while for Ub-TAX1BP1, ratios

are relative to Ub-TAX1BP1-WT levels after 2 h CCCP. One-way ANOVA was used to

assess significance unless otherwise specified. Bars: Mean ± SEM.

1350

Figure 4: Single cell analysis of TAX1BP1 translocation to mitochondria after CCCP-

mediated depolarization. (**A**) Representative immunofluorescence images showing

TAX1BP1 (yellow), parkin (magenta) and SSBP1 (cyan) after mitochondria depolarization

in 5KO HeLa cells transiently co-expressing Parkin and TAX1BP1. Solid arrowheads

indicate TAX1BP1 structures co-localizing with mitochondrial material. Arrows indicate

TAX1BP1 structures not engaged with mitochondrial material. ROI 1 represents > 50 % of

cell population while ROI 2 represents < 50% of cases. Scale bars: 10 μm. Zoom insets for

WT (i), TAX1BP1^{K549R} (ii), TAX1BPΔCC3 (iii and iv) are shown below. (**B**) Mean

mitochondrial size of SSBP1 foci after 4h CCCP. (**C-D**) Normalized SSBP1 (C) area or (D)

perimeter per cell. (**E**) Correlation analysis of the percentage of mitochondrial area

recognized by TAX1BP1 (X axis) in relation to the percentage of total TAX1BP1 found on

mitochondria (Y axis). Significance p-values are indicated in the graph for the each

TAX1BP1 construct. (**F**) Left panel: quantitative analysis of WT, TAX1BP1^{K549R} or

TAX1BP1ΔCC3 localization after mitochondria depolarization in 5KO-PRKN cells after

mitophagy induction (4h CCCP). TAX1BP1 localization is classified between cells where

COX4 clumps are fully recognized by TAX1BP1 (complete apposition) and cells where

COX4 clumps recognition is limited (localized apposition). Bars: mean ± SEM. Right panel:

representative immunofluorescence images stained for TAX1BP1 (yellow) and COX4

(magenta) are shown for each category represented in graph.

1370

Figure 5: Autophagy flux blockage effects in 5KO-PRKN HeLa cells. (**A-B**)

Representative immunofluorescence images of 5KO-PRKN HeLa cells showing TAX1BP1

(yellow), parkin (magenta) and SSBP1 (cyan) after mitochondria depolarization only (A) or

with Bafilomycin A (BafA) (B). In (A), the cell contour of cells lacking SSBP1 is

highlighted. In (B), zoom insets are shown for WT (i), TAX1BP1^{K549R} (ii) and

TAX1BPΔCC3 (iii), where solid arrowheads indicate enlarged TAX1BP1/SSBP1-positive

puncta and thin arrowheads indicate small TAX1BP1/SSBP1-positive puncta**.** (**C-D**) Eye-

based quantification of (C) cells with remaining SSBP1 material after extended

depolarization (16 h CCCP) and (D) cells harboring enlarged or small TAX1BP1 puncta

positive for SSBP1 (16 h CCCP + BafA). (**E**) Changes in TAX1BP1 ubiquitination after 16

h of CCCP treatment combined with BafA. **(F)** Mean TAX1BP1 puncta size after autophagy

blockage with BafA combined with CCCP or Oligomycin and Antimycin A (OA) treatment

for 16 h. **(G)** Pearson correlation indexes for TAX1BP1 and HSP60 signals after CCCP and

OA treatment combined with BafA. In panels (E) and (G), data of individual cells is

represented by single dots. All data come from at least three independent experiments. **(H)**

Percentages of cells showing at least 25% of TAX1BP1 enlarged puncta after the indicated

treatments. Bars: Mean ± SEM. Scale bars: 10 μm. KR: K549R, ΔCC: ΔCC3.

1388

Figure 6: Characterization of enlarged TAX1BP1$^+$ vesicles carrying mitochondrial

material upon autophagy flux inhibition in 5KO-PARK. (**A**) Representative images of

5KO-PRKN HeLa cells showing enlarged TAX1BP1$^+$ HSP60$^+$ structures (magenta and cyan,

respectively) positive for the autophagosome marker LC3B (yellow) after mitochondria

depolarization and BafA for 16 h. Scale bars: 5 μm. (**B**) Z-stack projection of enlarged

vesicles identified in (A) for WT, TAX1BP1^{K549R} and TAX1BP1ΔCC3. Thick arrowheads

indicate LC3B$^+$ TAX1BP1$^+$ cup-shaped structures in each Z-plane. Thickness slice: 0.6 μm.

(**C-E**) Representative images showing enlarged TAX1BP1$^+$ HSP60$^+$ structures positive for

RAB7 (C), VPS35 (D) and CD63 (E). Zoom insets are shown for WT (i), TAX1BP1^{K549R} (ii)

and TAX1BPΔCC3 (iii) for all markers. Scale bars: 5 μm.

1399

Figure 7: Characterization of small TAX1BP1$^+$ vesicles carrying matrix mitochondrial

material upon autophagy flux inhibition in 5KO-PARK. (**A-D**) Representative images of

5KO-PRKN HeLa cells showing small TAX1BP1/HSP60-positive structures (magenta,

cyan) positive for the indicated markers (yellow): LC3B (A), RAB7 (B), VPS35 (C) and

CD63 (D). Zoom insets are shown for WT (i), TAX1BP1^{K549R} (ii) and TAX1BPΔCC3 (iii)

for all markers. Scale bars: 5 μm.

1406

Figure 8: Characterization of TAX1BP1$^+$ structures identified in WT-Parkin Hela cells

upon autophagy flux inhibition and extended depolarization. (**A**) Representative images

upon siRNA mediated knock-down of TAX1BP1 in WT-Parkin HeLa cells showing

TAX1BP1$^+$ structures (magenta) carrying mitochondrial material (HSP60, cyan). Scale bars:

10 μm. Zoom insets are shown for the indicated conditions (right side, i-v). Lower panel of

western blot insets illustrates silencing and re-transfection efficiency (n=2). (**B-G**)

CellProfiler based analysis of TAX1BP1 puncta size or HSP60 material colocalizing with

TAX1BP1 puncta in WT-Parkin HeLa cells after siRNA mediated TAX1BP1 knockdown

and TAX1BP1 re-transfection upon CCCP and BafA treatment. (B) Mean TAX1BP1$^+$ puncta

size. (C) Percentage of intracellular enlarged TAX1BP1$^+$ puncta per cell. (D) Percentage of

total cells harboring enlarged TAX1BP1$^+$ puncta. (E) Percentage of total HSP60 foci

colocalizing with TAX1BP1. (F) Percentage of total HSP60 foci colocalizing with enlarged

TAX1BP1$^+$ puncta. (G) Mean HSP60 foci area colocalizing with enlarged TAX1BP1$^+$

structures.

1421

Figure 9: Co-localization analysis of TAX1BP1 and LC3B on depolarized mitochondria

upon overexpression of WT, TAX1BP1^{K549R} or TAX1BP1ΔCC3. Depolarization was

performed for 4 hours with CCCP or OA, as indicated. (**A)** Representative images showing

TAX1BP1 (yellow), LC3B (magenta) and HSP60 (cyan) after 4 h of depolarization. Scale

bars: 8 μm. (**B-E**) Colocalization analysis of TAX1BP1-HSP60 (B-C) and LC3B-HSP60 (D-

E), represented with thresholded Manders coefficient or Pearson's correlation coefficient for

the respective treatments and TAX1BP1 transfection, as indicated. (**F**) Mean LC3B intensity

observed upon the indicated treatments and TAX1BP1 transfection. i.u: intensity units. (**G**)

LC3B area relative to total cell area. In all panels, data of individual cells is represented by

single dots. All data come from at least three independent experiments. KR: K549R, ΔCC:

ΔCC3.

1433

Figure 10: Mass spectrometry analysis of TAX1BP1 native pulldown indicates defects

in protein trafficking processes in TAX1BP1ΔCC3. (**A**) Statistical analysis of

TAX1BP1ΔCC3 vs. WT TAX1BP1 pulldowns represented by the $\log_{10}$ p-value plotted

against student's T-test difference. Significance lines represent $p \leq 0.05$ and $p \leq 0.1$, as

indicated. Proteins are grouped according to the following GO-Terms: vesicle-mediated

transport in orange (GO:0016192), membrane organization in brown (GO:0061024) and

mitophagy in red (GO:1903146; GO:1901524; GO:1905089; GO:1905091); mitochondria

localization in lilac (based on UniProt annotation) or if they are associated with

PINK1/parkin mitophagy in grey (based on literature). Proteins related to the GO-Term

vesicle-mediated transport (GO:0016192) are classified according to their functionality

(based on UniProt and literature); between endosome budding from Golgi (light green) or

COPII vesicle related (dark green). (**B**) Overrepresentation analysis of biological processes

for significantly regulated proteins by Fisher exact testing.

1447

Figure 11: Structural visualization of TAX1BP1 using the predicted Alphafold

structure. (**A**) Domain organization of the coiled-coil domains of TAX1BP1 according to

Alphafold (top panel). Predicted Alphafold structure of TAX1BP1 with the indicated per-

residue model confidence scores (pLDDT). In the zoom panel, identified lysines in the mass-

spectrometry screen are highlighted in green. Highlighted in magenta are the remaining

lysines of the CC3, which were not identified via MS. (**B**) TAX1BP1, structure in

combination with a ubiquitin molecule at its ubiquitin-binding domain (UBD). Zoom panel

shows lysines of CC3 (residue surface) in close proximity to the UBD. Highlighted lysines

of CC3 are shown in green, if they were identified in the mass-spectrometry screen. The rest

of lysines from CC3 are shown in magenta. Images created with ChimeraX [63].

1458

Figure 12: Graphical summary highlighting how reduced ubiquitination on TAX1BP1

can influence the degradative pathway responsible for eliminating defective

mitochondria in a PINK1/parkin context. (**A**) Upon PINK1/parkin activation, TAX1BP1

is ubiquitinated at the CC3 domain and translocates to mitochondria. Recognition of

ubiquitinated mitochondria via TAX1BP1 promotes autophagosomal delivery. Simple

autophagosomal-dependent degradation occurs after successful engulfment of defective

mitochondria, autophagosome closure and autophagosome maturation, fostering the

degradation of mitochondria. (**B**) Reduced TAX1BP1 ubiquitination or lack of the

ubiquitinatable CC3 domain of TAX1BP1 in a context of PINK1/parkin mitophagy follows

a slightly different behavior. Here, while TAX1BP1 is able to recognize ubiquitinated

mitochondria, its translocation to mitochondria is hampered when compared to WT/normal

conditions. This recognition is able to induce the formation of autophagosomes around

defective mitochondria in this case, albeit at lower level. Recruitment of LC3-positive

vesicles occurs in a more localized manner, which deviates mitophagy dynamics from this

point on. Although this pathway leads to successful degradation of defective mitochondria,

in this case the prevailed degradative pathway is more dependent on the endolysosomal

network rather than on the classical autophagy machinery.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

(a) Normal TAX1BP1 ubiquitination at CC3

TAX1BP1
CC3 region
Lipidated ATG8
TAX1BP1
TAX1BP1
TAX1BP1

$\downarrow\Delta\Psi$

Recognition

Autophagosome biogenesis

Autophagosome closure and maturation

Simple autophagosomal dependent degradation

(b) Reduced TAX1BP1 ubiquitination at CC3

TAX1BP1
TAX1BP1
TAX1BP1

$\downarrow\Delta\Psi$

Recognition
REDUCED

Autophagosome biogenesis
LOCALIZED

DELAYED ??
Autophagosome closure and maturation

Endolysosomal and autophagosomal dependent degradation