1   Interchromosomal Linkage Disequilibrium Analysis Reveals Strong Indications of Sign Epistasis in

2   Wheat Breeding Families

Augusto Tessele[1], Geoffrey Morris[2], Eduard Akhunov[3], Blaine Johnson[4], Marshall Clinesmith[1],

Allan Fritz[1]

[1]Department of Agronomy, Kansas State University, Manhattan, Kansas, USA, 66506-5500

[2]Department of Soil and Crop Sciences, Colorado State University, Fort Collins, Colorado, USA,

80523-1101 [3]Department of Plant Pathology, Kansas State University, Manhattan, Kansas, USA,

66506-5500 [4]Department of Agronomy and Horticulture, University of Nebraska, Lincoln,

Nebraska, USA, 68583-0915 *Corresponding author: Department of Agronomy, Kansas State

University, Manhattan, Kansas, USA, 66506-5500 E-mail: tessele@ksu.edu

Abstract

Additive gene action is assumed to underly quantitative traits, but the eventual poor

performance of elite wheat lines as parents suggests that epistasis could be the underlying

genetic architecture. Sign epistasis is characterized by alleles having either a beneficial or

detrimental effect depending on the genetic background, which can result in elite lines that fail

as parents in certain parental combinations. Hence, the objective of this study were to test the

existence of sign epistasis and examine its consequences to wheat breeding. The presence of sign

epistasis is expected to distort the allele frequency distribution between two interacting genes

compared to neutral sites, creating strong linkage disequilibrium (LD). To test this hypothesis,

analysis of interchromosomal LD in breeding families was performed and detected 19 regions in

strong disequilibrium, whose allele frequency distribution matched the sign epistasis prediction

and falsified the competing hypothesis of additive selection. To validate these candidate

interactions while avoiding the biases of a circular analysis and the confounding effects of genetic

drift, two independent sets of populations were analyzed. Genetic drift was attributed to creating

the sign epistasis patterns observed in eleven interactions, but there was not sufficient evidence

to reject the sign epistasis hypothesis in eight interactions. Sign epistasis may explain the poor

performance of elite lines as parents, as crossing lines with complementary allelic combination

re-establishes epistatic variance in the offspring. Reduction in the effective population size in

certain crosses may also occur when unfavorable sign epistatic combinations are deleterious. The

potential existence of di-genic and higher order epistatic interactions in elite germplasm can

tremendously impact breeding strategies as managing epistasis becomes imperative for success.

Introduction

The infinitesimal model (Fisher, 1918) proposes that additive effects play a primary role
in genetic inheritance, while relegating dominance and epistasis to lesser importance (Turelli,
2017). However, the eventual occurrence of elite lines performing poorly as parents suggests that
epistasis might be the underlying genetic architecture. Sign epistasis, a significant form of gene
action in evolutionary genetics (Wright, 1932), is characterized by alleles exhibiting either
advantageous or detrimental effects depending on the allelic state of the interacting gene
(Weinreich et al., 2005; Poelwijk et al., 2007). Under the influence of sign epistasis, genotypes can
adapt to environmental pressures in various combinations of alleles, often leading to similar
adaptation (Wright, 1988). In breeding, it implies the existence of elite genotypes with different
frameworks of favorable allelic combinations. Hence, crossing contrasting genotypes causes the
favorable allelic combinations to segregate in the offspring and results in mean performance
lower than the midparent value. This clash of epistatic interactions between parental lines may
justify the eventual poor performance of elite lines as parents.

Establishing the importance and extent of epistasis in wheat breeding by modeling
epistatic variance has tremendous challenges because of the lack of orthogonality with additive
effects (Tessele et al., 2024; Raffo et al., 2022; Vitezica et al., 2017). However, the existence of
sign epistasis interactions under selection may leave detectable signatures in the genome. The
influence of sign epistasis in each locus closely resembles the effects of balancing selection, which
maintains genetic diversity at sites under selection and can augment diversity at linked loci
(Lewontin and Hubby, 1966; Hudson and Kaplan, 1988; Charlesworth et al., 1997; Takahata and
Satta, 1998). The polymorphism flanking the loci under balancing selection can span over large or
short regions, depending on local recombination frequency and strength of LD (Charlesworth et
al., 1997; Wiuf et al. 2004; Charlesworth, 2006), the age of the advantageous polymorphism and
the selection intensity (Kreitman and Rienzo, 2004; Tian et al., 2002). Besides affecting each locus
independently, sign epistasis also impacts the relationship between interacting genes because
selection acts on allelic combinations rather than on individual alleles. In a two-way interaction,
the preferential selection of favorable allelic combinations distorts the allele frequency
distribution of interacting genes compared to two neutral sites, creating strong disequilibrium
that can be regarded as a selection signature.

Although selection for sign epistasis may leave signatures in the genome, the mere
existence of LD does not imply sign epistasis. For instance, short-range linkage disequilibrium
(SRLD) primarily arises due to random genetic drift or common ancestry of chromosomal blocks
(Koch et al., 2013). SRLD are perpetuated under low recombination, making it virtually impossible
to disentangle from an eventual sign epistasis signature. Long-range linkage disequilibrium,
however, suggests that a countervailing force to recombination maintains strong LD between

distant sites, which is expected to be even more prominent in interchromosomal sites because

the independent assortment of chromosomes recombines physically unlinked loci in every

meiotic event. If sign epistasis is pervasive in wheat, breeding programs may perpetuate

interchromosomal LD by continuously selecting favorable interactions. Detecting true selection

signatures of epistasis (Lewontin et al., 1960) is challenging as other confounding effects can also

create long-range LD, as population admixture (Maccaferri et al., 2005; Chao et al., 2010), genetic

drift (Koch et al., 2013), artificial selection (Flint-Garcia et al., 2005; Joukhadar et al., 2019), or

structural variation in chromosomes (Zhao et al., 2022). Disentangling signatures of epistasis from

genetic artifacts may be possible through examining the allele frequency of sites in strong LD, as

additive selection is expected to create a different allele frequency distribution than epistasis.

Population structure can be minimized by analyzing breeding families independently. Analyzing

interactions independently from the base population used for discovery may reduce the bias

steaming from genetic drift and prevent statistical inference on a circular analysis (Kriegeskorte

et al., 2009).

The main hypothesis is that sign epistasis exists and has major implications to wheat

breeding. The objectives were to: (i) validate the existence of loci in abnormally strong

interchromosomal linkage disequilibrium; (ii) test if selection for sign epistasis creates the

observed strong interchromosomal LD; (iii) verify the existence of and disentangle confounding

genetic artifacts that could likewise create strong interchromosomal LD.

Material and Methods

Plant Material and Genomic Data

The dataset consisted of genomic information on experimental lines from the Kansas

State Hard Red Winter Wheat breeding program for the years 2011 through 2021, which were

categorized based on the last stage of the breeding pipeline reached by each experimental line.

The four stages in the breeding pipeline were early, preliminary, advanced, and elite yield trials,

named IPSR, PYN, AYN and EYN, respectively. All experimental lines were genotyped in the IPSR

stage using genotype-by-sequencing (GBS) (Poland et al. 2012). As part of the genotyping pipeline

of the K-State wheat breeding program, markers with a MAF <0.01, more than 20% missing values

and more than 20% of heterozygotes were filtered, as described in Jarquin et al (2017). The R

package 'rrBLUP'(Endelman, 2011) was used for marker imputation, where missing markers were

imputed with the mean value among all lines for that marker and resulted in a total of 55,148

SNPs.

Validating the existence of strong interchromosomal linkage disequilibrium

This exploratory stages sought to validate the existence of interchromosomal linkage

disequilibrium in breeding germplasm. To test this hypothesis, experimental lines from the EYN

stage of the breeding program from 2010 to 2021 were considered. A linkage disequilibrium

analysis (r^2) focusing only on interchromosomal interactions was conducted and identified

pairwise markers in strong linkage disequilibrium. Validating the existence of strong

interchromosomal LD in wheat enabled further investigation into the hypothesis of selection

signatures of sign epistasis (Figure 1).

Figure 1 Selection of sign epistasis interactions can create interchromosomal LD. **A** depicts the

levels of adaptation of two genes (A and B) under the effects of sign epistasis, drift and additive.

B shows the expected allele frequency distribution under an ideal scenario for each genetic effect.

C illustrates strong linkage disequilibrium between pairwise interchromosomal sites, where inner

links connect sites in strong LD. **D** represents the allele frequency distribution in five populations

for two genes in strong LD and the respective heatmap of intra- and inter-chromosomal LD for

each genetic effect. **E** represents the allele frequency and LD values in an independent panel from

D, which highlights the consistency of sign epistasis contrasting to the randomness of genetic

drift.

Minimizing the influence of population structure and genetic drift

126   Historical data from 25 breeding families was used for the discovery of selection

signatures indicative of sign epistasis (Figure 1C). These were large families tested in the early

yield trial stage of the breeding pipeline, with sizes ranging from 20 to 45 genotypes. As a data

quality measure, a principal component analysis was conducted for each of the families to identify

and exclude any outlier genotypes detected in the two-dimensional plot defined by the first two

principal components to mitigate the potential introduction of population structure resulting

from genotyping or seed processing errors.

133   An interchromosomal linkage disequilibrium (r^2) analysis was conducted only on markers

with MAF>0.3 to reduce the total number of interactions to screen for and to increase the

likelihood of detecting interactions with allele frequency characteristic of sign epistasis (Figure 1A

and B). To circumvent the effect of population structure, the interchromosomal LD analysis was

conducted in each family individually and all interactions presenting $r^2 > 0.9$ were retained.

Although this approach controls population structure, it was highly influenced by genetic drift

within families. To address the genetic drift effect, the combined data of the 25 families was used

to run another interchromosomal LD analysis, and all interactions presenting $r^2 > 0.8$ were

selected. Interactions selected in both analyses and observed in a minimum of two families were

deemed worthy of further investigation.

143   In parallel, a control interaction was utilized to serve as a null hypothesis. The control

interaction consisted of two randomly selected single nucleotide polymorphisms (SNPs) with MAF

greater than 0.3 and found in chromosomes 1D (S1D_463434894) and 2B (S2B_163607648). To

establish a null control for the candidate three-way interaction, a third SNP was sampled from

the genome, which was found in chromosome 5A (S5A_298295448). The analytical procedures

applied to assess the candidate interactions were also applied in the evaluation of the control

interaction.

The Balancing Selection Nature of Sign Epistasis and Interchromosomal Linkage Blocks

152   A sign epistatic interaction under selection, besides the characteristic intrachromosomal

haplotype block, may also create an "interchromosomal block" (Figure 1D). To investigate this

hypothesis, 40 markers flanking each candidate sign epistatic marker were used to run a linkage

disequilibrium analysis, and the resulting intra and interchromosomal LD values were used to

construct heatmaps of 2D LD plots.

Allelic Frequencies to Differentiate Additive and Sign Epistasis

Selection for sign epistasis is expected to change allele frequencies differently than

additive selection (Figure 1B). To empirically test this hypothesis, the allele frequencies of each

candidate interaction was computed, and the patterns were analyzed (Figure 1D). Since these

families were genotyped in the $F_{5:6}$ generation using GBS, heterozygous loci were still observed.

Independent Family and Breeding Program Validation

To minimize the effects of genetic drift and to avoid a circular analysis, 15 distinct

validation families were selected to reanalyze the heatmaps of 2D LD plots and the distribution

of allele frequencies in order to validate candidate interactions (Figure 1E). Although independent

from the discovery set, the total number of families analyzed was relatively small and prone to

the effects of genetic drift. To diminish the confounding effects of drift in small populations, data

from a time span of 10 years in the wheat breeding program was also utilized to further validate

the consistency of candidate interactions. The breeding program dataset was divided sequentially

from early to elite yield trials and allele frequencies were calculated for each stage. In addition,

the combined data of the breeding program was used to falsify candidate interactions if the

opposite combination (relative to the most common allelic combination) had less than 10% of the

total allelic combinations.

Software and Packages

The linkage disequilibrium analysis was performed using the software PLINK (Purcell et

al., 2007). The R software packages "factoextra" (Kassambara et al., 2017), "circlize" (Gu and Gu,

2022), and "gaston" were utilized to perform the principal component analysis, to plot the SNPs

presenting strong interchromosomal LD, and to plot the heatmap of 2D LD plots, respectively.

The package "ggplot2" was used to plot allele frequency counts (Wickham, 2016). In addition,

built-in functions in R were used during data management and to calculate allele frequencies.

Results

Detection of loci in strong interchromosomal linkage disequilibrium

The goal of the discovery stage was to pinpoint a manageable number of interactions in

strong LD while minimizing the likelihood of identifying artifacts of population structure, genetic

drift, or additive selection. In total, 269 pairwise SNPs, which represent 19 interacting genomic

regions, passed the set thresholds, and were considered candidate selection signatures (Figure

2). Wheat subgenome B had the most candidate interactions with 15, followed by A and D with

13 and 10, respectively. The chromosomes with the most candidate interactions were 1B and 6B,

with four interactions each. In total, 14 out of the 19 interactions were intergenomic, and most

involved subgenomes B and D. Subgenome A and B had candidate intragenomic interactions,

which were absent in subgenome D. In addition, a candidate of a three-way selection signature

was also observed in chromosomes 2D, 3B and 4A (Figure 2). A total of seven candidate

interactions had opposed allelic combinations fixed in different families, and 12 were fixed for

one allelic combination while the opposed allelic combination was observed in segregating

families (Supplemental Table 1).

Figure 2 Candidate selection signatures of sign epistasis. The inner links represent regions (SNPs)

in strong interchromosomal LD following the discovery approach previously described. The outer

histogram plots represent the marker density in each chromosome. The outermost bar

represents the 7 chromosomes and 3 subgenomes of wheat.

Putative balancing selection associated with sign epistasis

The analysis of linkage drag was performed for each of the 19 candidate selection

signatures of sign epistasis. The results for the candidate interaction involving chromosomes 1A

and 4A reveals the presence of "interchromosomal linkage blocks" associated with the candidate

selection signature (Figure 3B). These blocks are primarily attributed to a sizable linkage block

located on chromosome 1A and a smaller block situated on chromosome 4A. The control

interaction, however, displayed some level of intrachromosomal LD on chromosome 2B but no

"interchromosomal LD block" was observed (Figure 3B).

The remaining interactions presented "interchromosomal linkage blocks" of variable

sizes, as the number of flanking SNPs in strong LD with the candidate interaction was variable

(Supplemental Figures 1-18). For example, candidate interactions 5D-7B, 2D-3B, 1BS-1D and 1BL-

1D were in intrachromosomal LD with a few SNPs, resulting in very small "interchromosomal LD

blocks".

Figure 3. Testing independent breeding panels for sign epistasis. Allele frequency distribution (left column) and heatmaps of 2D LD plots (right column) of the candidate epistatic interaction involving segments of chromosomes 1A and 4A and the drift competing hypothesis involving segments of chromosomes 1B and 2D. A and B represents the data from the discovery panel. C and D displays data from the validation panel. E and F consists of data from the last four stages

of the breeding pipeline. **G** displays the expected allele frequency distribution in the entire

breeding program under Hardy-Weinberg Equilibrium and **H** shows the observed allele frequency

distribution.

Allele frequencies disentangle epistasis from additive selection

The allele frequencies of candidate interaction 1A-4A in the discovery stage reveals many

families nearly fixed for the A/T and C/C allelic combinations (Figure 3A). Families segregating for

these two alleles exhibited a high frequency of opposed allelic combinations, suggesting that

selection is actively favoring both the A/T and C/C allelic combinations. Off-combinations

consistently presented very low frequency in most families. The absence of a predominant allelic

combination in high frequency among segregating populations weakens the competing

hypothesis of additive selection. The allele frequency of the control interaction shows a variable

predominance of allelic combinations in different families, which is suggestive of genetic drift in

unselected loci (Figure 3A).

The candidate interactions 6B-6D, 3B-3D, 2B-5B had a similar distribution of opposing

allelic combinations as interaction 1A-4A (Supplemental Figures 1-A, 3-A, 4-A). Candidate

interactions 2A-2D, 2D-3B, 2D-4A, 3B-4A and 5A-7A had a common pattern where one of the

contrasting combinations predominated in most families, followed by the opposed combination

as the second most frequent, and followed by heterozygotes (Supplemental Figures 5-A, 8:11-A).

However, for interactions 5A-6B, 1B-6B, 5D-7B, 6A-7A, 3A-3D, 1A-1B, 1BL-1D,1BS-1D, 2B-7A and

6BL-6D, the most common allelic combination was found at a very high frequency in most

families, and the opposing combination was observed in only a few families (Supplemental Figures

2-A, 6-A, 7-A, 12:18-A). In all cases, the off-combinations were observed at very low frequencies

or absent in many families.

The allele frequencies of the candidate three-way epistatic interaction indicates the

predominance of the C/C/C alleles, which was fixed in many families (Supplemental Figure 19-A).

In segregating families, the opposing allelic combination (A/T/T) was found in high frequency,

suggesting that both combinations could be under positive selection. Off-combinations were less

common than heterozygotes in most families.

Validation Families Support Sign Epistasis Hypothesis

The patterns of LD in the validation families remained consistent with the discovery

families for interaction 1A-4A, where a long linkage block on chromosome 1A and a small block

on 4A created the characteristic "interchromosomal linkage block" (Figure 3-D). The frequency of

the allelic combinations in the validation families presented a similar pattern as the discovery

families (Figure 3-C). Six populations are fixed for the A/T combination, and the remaining 11 are

segregating with both A/T and C/C as the most frequent combinations. The frequency of off-

combinations was also very low. The random interaction (1D-2B) exhibited different allelic

combinations dominating different families as in the discovery families, highlighting the influence

of genetic drift in loci likely not under selection.

Interaction 2B-5B (Figure 4-C) had a similar pattern of allelic combination as interaction

1A-4A, where the validation families had an allele frequency pattern similar to the discovery

families. The interchromosomal linkage analysis for interactions 1B-6B, and 5D-7B did not present

the characteristic "interchromosomal LD block" (Supplemental Figure6-D and 7-D) because most

families were fixed for the major allelic combination (Supplemental Figure 6-C and 7-D).

On the allele frequency analysis, the candidate interactions 5A-7A, 1BL-1D, 1BS-1D, 2B-

7A and 6B-6Db (Supplemental Figures 11-C, 15:18-C) had at least one family with an off-

combination as the most frequent allelic combination. This suggests that these loci are not under

selection, and the sign epistasis hypothesis was rejected in these candidate interactions. For

interaction 6B-6D*a*, family KS120427 had an off-combination as the most frequent allelic

combination. However, the abnormal high frequency of heterozygotes, which could have a

biological meaning or simply be the consequence of genotyping errors, prevents drawing any

definitive conclusions about the interaction (Supplemental Figure 1-C). Similarly, for interaction

3B-3D, the family KS090132 was fixed for an off-combination but the remaining families

presented allelic frequencies matching the sign epistasis hypothesis (Supplemental Figure 3-C). It

is probable that the parental lines of population KS090132 were fixed for the off-combination,

which prevents observing the favorable candidate combinations. Thus, this interaction was kept

for further analysis.

For the remaining interactions, a consistent pattern emerged where one allelic

combination predominated in the majority of families, either fixed or at a very high frequency,

while the opposing allelic combination was observed in a few other families at varying frequencies

(Supplemental Figure 2-C, 5-C, 8:10-C, 12:14-C). The candidate three-way interaction was

segregating in few populations, with the allelic combination C/C/C as the most frequent, followed

by the opposed allelic combination (A/T/T) and a small occurrence of off-combinations

(Supplemental Figure 19-C). The remaining families were fixed for the C/C/C allelic combination.

Breeding Program Validation

To further test if the 12 non-falsified interactions are not the result of strong genetic drift

effects in small population sizes, data from 10 years of the breeding program was used to

calculate the frequency of allelic combinations. The results for interaction 1A-4A (Figure 1-E)
reveal that off-combinations were present at low frequencies in the early stages of yield trials
(IPSR). The frequency of off-combination declined with successive generations of selection,
ultimately leading to complete elimination of off-combinations in the EYN. Notably, the two
opposing allelic combinations, A/T and C/C, exhibited a balanced frequency across all breeding
stages analyzed. For the control interaction, an inconsistent pattern of allele frequencies was
observed across breeding stages.

The same pattern was observed for interactions 6B-6Da, 5A-6B, and 6A-7A (Supplemental
Fig 1-E, 2-E, and 12-E), which exhibited no or very few off-combinations at the EYN stage.
However, the latter two candidate interactions had the opposite combination with less than 10%
of the total allelic combinations, suggesting that the detection of these interactions may have
arisen due to genetic drift. Consequently, the sign epistasis hypothesis for the 5A-6B and 6A-7A
interactions was rejected. Similar allele frequencies were observed for interactions 3A-3D and
1A-1B, leading to the rejection of these combinations as well (Supplemental Figure 13-E and 14-
E).

The six remaining interactions (3B-3D, 2B-5B, 2A-2D, 2D-3B, 2D-4A, and 3B-4A) displayed
a high frequency of both opposing allelic combinations but showed minor variations in off-
combinations across the breeding stages (Supplemental Figures 3:5-E, 8:10-E). Interactions 3B-
3D, 2B-5B, and 2A-2D had an average of 94% of genotypes with the two favorable contrasting
allelic combinations, at an average ratio of 4:1 between the most common allelic combination
and its opposed combination. The interactions 2D-3B, 2D-4A, and 3B-4A comprised the candidate
three-way sign epistasis interaction and had the two favorable interactions in very high
frequencies across the four breeding stages (Figure 19-E). The C/C/C combination added up to
almost 88% of the genotypes across the breeding stages while the A/T/T represented a little over
10%. The remaining 6 off-combinations summed up to less than 2% of the interactions.

Discussion

Strong interchromosomal LD could be created from selection of epistatic interactions

The poor performance of elite lines as parents is hypothesized to be related to epistatic
effects. Although estimating epistatic variance is difficult because of non-orthogonality with
additive effects (Tessele et al., 2024; Raffo et al., 2022; Vitezica et al., 2017), selection signatures
of epistasis are expected to exist in breeding programs. In this study, 25 breeding families were
utilized to identify 19 regions in strong interchromosomal LD which are hypothesized to be
selection signatures of epistasis. Interchromosomal LD patterns were also observed in tetraploid
wheat (Laido et al 2014) and common beans (Rossi et al., 2009; Diniz et al., 2018), but no

inferences regarding the genetic mechanism leading to interchromosomal LD were drawn.

The allele frequency distribution of candidate interactions resembled that expected

under the influence of sign epistasis, which falsifies the additive hypothesis. Similar results were

also observed in wild barley, where the cross of plants based on spatial distance resulted in co-

adapted gene complexes fixed in locally abundant genotypes (Volis et al., 2011). In two diverse

populations of barley, several generations of natural selection resulted in two out of the 16

potential combinations of gametes (4^2) as the most frequent, which presented perfectly

complementary gametic types (Clegg et al. 1972), suggesting natural selection acted on

coadapted (Dobzhansky, 1970) multilocus units.

As an attempt of disentangling epistasis from drift, an independent set of 15 validation

families was utilized to further test the consistency of the candidate interactions, which falsified

seven interactions. As the strength of genetic drift effects is expected to be especially

predominant in smaller populations (Charlesworth et al., 2003; Charlesworth, 2009; Kliman et al.,

2008; Rothammer et al. 2013), the assembly of allelic combinations resembling selection for sign

epistasis in the discovery stage could have been a consequence of genetic drift. Random genetic

drift has also been attributed to the emergence of apparent selection signatures between pairs

of SNPs in simulation studies (Id-Lahoucine et al., 2019; Vilas et al., 2012).

To mitigate the impact of genetic drift, a significantly larger dataset spanning a decade of

breeding records was analyzed, leading to the rejection of four interactions. Among the remaining

eight interactions, 2A-2D, 2D-3B, 2D-4A, and 3B-4A had a single allelic combination representing

close to 90% of the total allelic combinations, contradicting the assumption of comparable

adaptability between opposing allelic combinations (Wight, 1932). However, this observed

phenomenon may relate to founder effects (James, 1971), where a restriction in allele frequency

within the founder germplasm results in the spread of one of allelic combination in the breeding

program. In that sense, the balanced frequency of opposed allelic combinations in interactions

1A-4A, 6B-6D, 3B-3D, and 2B-5B within an elite breeding program is surprising, considering that

recycling lines and extensive utilization of good parental lines is expected to favor one epistatic

combination over generations. For this reason, it is expected that the number and extent of sign

epistatic interactions should be much larger when considering different breeding programs, as a

function of founder effects, aka founder germplasm.

Sign Epistasis Reduces the Effective Population Size of Breeding Populations

The near-complete recovery of opposed allelic combinations as observed for interaction

1A-4A suggests artificial selection alone may not be the primary driver of distorted allele

frequencies. Instead, it is possible that natural selection is responsible for recovering favorable

allelic combinations, as observed in barley and wild barley populations (Volis et al., 2011; Clegg et al. 1972), or through a nearly lethal response to off-combinations, similar to hybrid necrosis in wheat (Tsunewaki, 1960). Anecdotal evidence on this hypothesis has been observed in single seed descent families in the Kansas State wheat breeding program. In this speed breeding pipeline, the quantity of seed recovered after three generations of inbreeding is often significantly lower than expected based solely on poor levels of germination. This could be an indication that a varying numbers of epistatic interactions, as interactions 1A-4A, 6B-6D, and 3B-3D, segregate in the families and plants homozygotes to off-combinations may experience a severe setback on performance which eventually leads to premature death or reproductive incapability. Populations with a higher number of segregating epistatic interactions, if additionally segregating to interactions 2B-5B, 2A-2D, and 2D-3B, for instance, may consequently end up with fewer viable plants. One alternative to overcome this challenge is to constrain the genetic diversity of the breeding program with the fixation of one of the interacting alleles, converting epistatic variance into additive variance (Hill et al., 2008; Technow et al., 2021). In the candidate interaction 1A-4A, fixing the A allele in chromosome 1A causes the allele in chromosome 4A to behave additively, where the T allele is always positive, and C is always negative (Figure 3 A, C, and E). This ensures that the selection pressure is constant for the alleles in the interacting locus and increases the probability of having one favorable combination fixed.

Sign Epistasis Creates a Rugged Surface on the Fitness Landscape

The possible existence of di-genic sign epistasis, as observed in interaction 1A-4A (Figure 4A), implies that transitioning from one superior combination to another requires two allelic substitutions, passing through a transitional state of inferior performance (Wright, 1988; Wade, 2002). The fitness landscape of the potential tri-genic sign epistasis interaction reported in this study has a valley between the two optimum allelic combinations (Figure 4B). And, if extended to higher order epistasis and more gene interactions (Figure 4C), it indicates that different combinations of alleles can present similar levels of adaptation to the same environmental conditions (Wright, 1988). The entire field of allelic combinations and respective levels of adaptation can be greatly simplified through the fitness landscape metaphor (Figure 4D), as proposed in the shifting balance theory (Wright, 1932), where hills and valleys represent high and low levels of adaptation, respectively. The fitness landscape, as a metaphor, should not be taken literally (Wade, 2002), as it is inadequate to represent higher order epistatic interactions (Wright, 1932). It does, however, provides a powerful simplified representation of complex genetic systems and how populations navigate the rugged surface during evolution. The concept of evolution under the shifting balance theory has striking similarities with the structure of breeding

programs (Technow et a., 2021) and could be leveraged to better understand epistasis in the face

of managing breeding germplasm pools.

Figure 4. Rugged fitness landscapes are formed from sign epistasis interactions. A The allele

frequency of interaction 1A-4A across the breeding program indicates a di-genic sign epistasis

interaction. **B** The potential three-gene sign epistasis interaction involving chromosomes 2D, 3B

and 4A creates a valley of adaptation between the two optimum allelic combinations. **C**

Combining a hypothetical four- and the potential three-gene sign epistasis interaction creates a

fitness landscape with regions of high, intermediate, and low adaptation. Factual representation

of the fitness landscape is difficult because of the elevated complexity of the genetic system. **D**

Rugged fitness landscape formed by several sign epistasis interactions with variable levels of gene

interactions. This is an extreme simplification of a highly complex, multi-dimensional genetic

system. Adaptations from Wight (1932).

Breeding Strategies on a Rugged Fitness Landscape

A rugged surface on the fitness landscape has implications for breeding programs as the

strategy used to manage the germplasm pool can lead to higher or lower levels of epistatic

variance in breeding families. The operation of a closed breeding program inevitably reduces the

genetic diversity in the germplasm pool and, under the influence of drift and selection, leads to

the fixation of alleles involved in sign epistatic interactions. This process converts epistatic

variance into additive variance and increases the potential of adaptation of the breeding program

(populations) to local environments (Wade and Goodnight, 1998), as less complex genetic

systems have higher heritability and respond better to selection. This can be visualized in Figure

4D as the *black squares* or *blue circles* genotypes, located in independent hills and assumed to

comprise separate breeding programs. Alternatively, operating a breeding program with a large

genetic base may reintroduce epistatic variance when recycling lines from within the breeding

program, which can be depicted as the *white star* genotypes in Figure 4D. The cross of parental

lines that belong to different adaptive peaks leads to a phenomenon called F_2 breakdown

(Whitlock et al., 1995), where the mean performance of the population is lower than the

midparent value. This is a consequence of disrupting, in the offspring, the contrasting epistatic

interactions that were homozygous in each parental line. Breeding families with high levels of

epistatic variance present a low correlation between early and late generations (Upadhyaya and

Nigam, 1998; Humphrey et al., 1969), indicating a limited response to early generation selection.

If late generation selection is adopted, the quick pace of inbreeding and the commonly small

population size of breeding families causes both favorable and unfavorable combinations to be

fixed, reducing the probability of assembling optimal allelic combinations.

One common feature of breeding programs is the exchange of germplasm, where elite

genotypes are anticipated to transmit superior allelic combinations from one breeding pool to

another (Wright, 1932; Technow et al., 2021). The success of introducing germplasm on a rugged

fitness landscape scenario depends on having both parental lines on the same adaptive peak, that

is, sharing the same founding allelic combinations, otherwise, F_2 breakdown may also be observed

in breeding families. For navigating on multiple adaptive peaks, broad genetic base programs may

be more suitable for introducing external germplasm as there is more than one set of founding

allelic combinations that could match the germplasm to be introduced. Although normally used

as a strategy to introduce a few alleles in the background of a recurrent line (Hospital, 2005),

backcrossing can be leveraged by closed breeding programs to recover the epistatic framework

of the recurrent line. Backcrossing constraints epistatic variance, increasing additive variance and

the response to selection in breeding populations. For a comprehensive discussion of the

implications of sign epistasis from the perspective of the fitness landscape metaphor, view

Tessele et al. (2024) and Holland (2001).

Conclusion

To expand the understanding of how epistasis could be responsible for the poor

performance of elite lines as parents and be of critical importance to breeding, the existence of

selection signatures of sign epistasis in wheat families was investigated in this study. In total, 19

candidate interactions were detected, whose allelic patterns agreed with the sign epistasis

predictions and refuting the competing additive hypothesis. To validate the candidate

interactions while avoiding the biases of a circular analysis and the confounding effects of random

genetic drift, two independent sets of materials were analyzed and falsified 11 interactions. There

was not sufficient evidence to reject the sign epistasis hypothesis for eight candidate interactions

with the current dataset and analysis. The possibility existence of sign epistasis could be

responsible for a reduction in the effective population size in crosses with contrasting sign

epistasis combinations, and for the eventual failure to recycle elite lines in the breeding program
as epistatic variance is reestablished in the offspring. The potential presence of a rugged fitness
landscape underscores the importance of adequately managing epistatic variance within
breeding programs for efficiently developing elite germplasm.
468

469 **Conflict of Interest**
470 None declared.
471

Funding
We acknowledge the Kansas Wheat Alliance and the Kansas Wheat Commission for their essential
financial support, which made this research possible.

Data Availability
Marker information of discovery and validation populations in HapMap format is included in files
'Discovery-Populations.hmp.txt' and 'Validation-Populations.hmp.txt'. The code to run the PCA
analysis to eliminate outliers within families is found in 'PCA.R'. The code used to calculate linkage
disequilibrium is found in 'C-Linkage_Disequilibrium_Loops.bach'. The code used to calculate the
2D-LD plots in discovery and validation families and the control plot is found in the
`*_2D_LDplot.R' files. The 'Circleze.R' file has the code used to plot interchromosomal LD with
chromosomes displayed circularly. Supplemental material is available at Figshare. The raw
genotypic data, available at Figshare, will be deposited at NCBI if/upon the manuscript is accepted
for publication.
486

References

Acosta-Pech, R., Crossa, J., de Los Campos, G., Teyssèdre, S., Claustres, B., Pérez-Elizalde,

S., & Pérez-Rodríguez, P. (2017). Genomic models with genotype× environment interaction for

predicting hybrid performance: an application in maize hybrids. Theoretical and Applied Genetics,

130, 1431-1440.

Alvarez-Castro JM, Carlborg O (2007) A unified model for functional and statistical

epistasis and its application in quantitative trait Loci analysis. Genetics 176:1151–1167.

Bamshad, M., & Wooding, S. P. (2003). Signatures of natural selection in the human

genome. Nature Reviews Genetics, 4(2), 99-110.

Chao, S., Dubcovsky, J., Dvorak, J., Luo, M. C., Baenziger, S. P., Matnyazov, R., ... &

Akhunov, E. D. (2010). Population-and genome-specific patterns of linkage disequilibrium and

SNP variation in spring and winter wheat (Triticum aestivum L.). BMC genomics, 11(1), 1-17.

Charlesworth, B. (2009). Effective population size and patterns of molecular evolution

and variation. Nature Reviews Genetics, 10(3), 195-205.

Charlesworth, B., Charlesworth, D., & Barton, N. H. (2003). The effects of genetic and

geographic structure on neutral variation. Annual Review of Ecology, Evolution, and Systematics,

34(1), 99-125.

Charlesworth, B., Nordborg, M., & Charlesworth, D. (1997). The effects of local selection,

balanced polymorphism and background selection on equilibrium patterns of genetic diversity in

subdivided populations. Genetics Research, 70(2), 155-174.

Charlesworth, D. (2006). Balancing selection and its effects on sequences in nearby

genome regions. PLoS genetics, 2(4), e64.

Cooper, M., Podlich, D. W., Micallef, K. P., Smith, O. S., Jensen, N. M., Chapman, S. C., &

Kruger, N. L. (2002). Complexity, quantitative traits and plant breeding: a role for simulation

modelling in the genetic improvement of crops. In Quantitative genetics, genomics and plant

breeding (pp. 143-166). Wallingford UK: CABI Publishing.

Diniz, A. L., Giordani, W., Costa, Z. P., Margarido, G. R., Perseguini, J. M. K., Benchimol-

Reis, L. L., Chiorato, A.F., Garcia, A.A.F., & Vieira, M. L. C. (2018). Evidence for strong kinship

influence on the extent of linkage disequilibrium in cultivated common beans. Genes, 10(1), 5.

Endelman, J. B. (2011). Ridge regression and other kernels for genomic selection with R

package rrBLUP. The plant genome, 4(3).

Federer, W. T., & Raghavarao, D. (1975). On augmented designs. Biometrics, 29-35.

Flint-Garcia, S. A., Thornsberry, J. M., & Buckler IV, E. S. (2003). Structure of linkage

disequilibrium in plants. Annual review of plant biology, 54(1), 357-374.

Forsberg, S. K., & Carlborg, Ö. (2017). On the relationship between epistasis and genetic

variance heterogeneity. Journal of experimental botany, 68(20), 5431-5438.

Gu, Z., Gu, M. Z., & GlobalOptions, I. (2022). Package 'circlize'.

He, S., Reif, J. C., Korzun, V., Bothe, R., Ebmeyer, E., & Jiang, Y. (2017). Genome-wide

mapping and prediction suggests presence of local epistasis in a vast elite winter wheat

populations adapted to Central Europe. Theoretical and Applied Genetics, 130, 635-647.

Hill, W. G., Goddard, M. E., & Visscher, P. M. (2008). Data and theory point to mainly

additive genetic variance for complex traits. PLoS genetics, 4(2), e1000008.

Hudson, R. R., & Kaplan, N. L. (1988). The coalescent process in models with selection and

recombination. Genetics, 120(3), 831-840.

Id-Lahoucine, S., Molina, A., Cánovas, A., & Casellas, J. (2019). Screening for epistatic

selection signatures: A simulation study. Scientific reports, 9(1), 1-5.

Jarquín, D., Lemes da Silva, C., Gaynor, R. C., Poland, J., Fritz, A., Howard, R., ... & Crossa,

J. (2017). Increasing genomic-enabled prediction accuracy by modeling genotype× environment

interactions in Kansas wheat. The plant genome, 10(2), plantgenome2016-12.

Joshi, R., Meuwissen, T. H., Woolliams, J. A., & Gjøen, H. M. (2020). Genomic dissection

of maternal, additive and non-additive genetic effects for growth and carcass traits in Nile tilapia.

Genetics Selection Evolution, 52(1), 1-13.

Joukhadar, R., Daetwyler, H. D., Gendall, A. R., & Hayden, M. J. (2019). Artificial selection

causes significant linkage disequilibrium among multiple unlinked genes in Australian wheat.

Evolutionary Applications, 12(8), 1610-1625.

Kassambara, A., & Mundt, F. (2017). Package 'factoextra'. Extract and visualize the results

of multivariate data analyses, 76(2).

Kliman, R., B. Sheehy & J. Schultz (2008) Genetic drift and effective population size.

Nature Education 1(3).

Koch, E., Ristroph, M., & Kirkpatrick, M. (2013). Long range linkage disequilibrium across

the human genome. PloS one, 8(12), e80754.

Kreitman, M., & Di Rienzo, A. (2004). Balancing claims for balancing selection. TRENDS in

Genetics, 20(7), 300-304.

Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S., & Baker, C. I. (2009). Circular analysis

in systems neuroscience: the dangers of double dipping. Nature neuroscience, 12(5), 535-540.

Kristensen, P. S., Jahoor, A., Andersen, J. R., Cericola, F., Orabi, J., Janss, L. L., & Jensen, J.

(2018). Genome-wide association studies and comparison of models and cross-validation

strategies for genomic prediction of quality traits in advanced winter wheat breeding lines.

Frontiers in plant science, 9, 69.

Lehermeier, C., de Los Campos, G., Wimmer, V., & Schön, C. C. (2017). Genomic variance
estimates: With or without disequilibrium covariances?. Journal of Animal Breeding and Genetics,
134(3), 232-241.

Lewontin, R. C. & Hubby, J. L. A molecular approach to the study of genetic heterozygosity
in natural populations. II. Amount of variation and degree of heterozygosity in natural populations
of Drosophila pseudoobscura. Genetics 54, 595–609 (1966).

Lewontin, R. C., & Kojima, K. I. (1960). The evolutionary dynamics of complex
polymorphisms. Evolution, 458-472.

Lin, C. S., & Poushinsky, G. R. E. G. (1985). A modified augmented design (type 2) for
rectangular plots. Canadian journal of plant science, 65(3), 743-749.

Liu, G., Zhao, Y., Gowda, M., Longin, C. F. H., Reif, J. C., & Mette, M. F. (2016). Predicting
hybrid performances for quality traits through genomic-assisted approaches in Central European
wheat. PLoS One, 11(7), e0158635.

Maccaferri, M., Sanguineti, M. C., Noli, E., & Tuberosa, R. (2005). Population structure
and long-range linkage disequilibrium in a durum wheat elite collection. Molecular Breeding, 15,
271-290.

Moran, P., & Smith, C. (1918). The correlation between relatives on the supposition of
mendelian inheritance. Transactions of the Royal Society of Edinburgh, 52, 899-438.

Pérez, P., & de los Campos, G. (2014). BGLR: a statistical package for whole genome
regression and prediction. Genetics, 198(2), 483-495.

Poland JA, Brown PJ, Sorrells ME, Jannink J-L (2012) Development of high-density genetic
maps for barley and wheat using a novel two-enzyme genotyping-by-sequencing approach. PLoS
ONE 7.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., ... & Sham,
P. C. (2007). PLINK: a tool set for whole-genome association and population-based linkage
analyses. The American journal of human genetics, 81(3), 559-575.

Raffo, M. A., Sarup, P., Guo, X., Liu, H., Andersen, J. R., Orabi, J., Jahoor, A., & Jensen, J.
(2022). Improvement of genomic prediction in advanced wheat breeding lines by including
additive-by-additive epistasis. Theoretical and Applied Genetics, 135(3), 965-978.

Rossi, M., Bitocchi, E., Bellucci, E., Nanni, L., Rau, D., Attene, G., & Papa, R. (2009). Linkage
disequilibrium and population structure in wild and domesticated populations of Phaseolus
vulgaris L. Evolutionary Applications, 2(4), 504-522.

Rothammer, S., Seichter, D., Förster, M., & Medugorac, I. (2013). A genome-wide scan
for signatures of differential artificial selection in ten cattle breeds. BMC genomics, 14(1), 1-17.

Sorensen, D., Fernando, R., & Gianola, D. (2001). Inferring the trajectory of genetic

variance in the course of artificial selection. Genetical Research, 77, 83–94.

Takahata, N., & Satta, Y. (1998). Footprints of intragenic recombination at HLA loci.
Immunogenetics, 47, 430-441.

Technow, F., Podlich, D., & Cooper, M. (2021). Back to the future: implications of genetic
complexity for the structure of hybrid breeding programs. G3, 11(7), jkab153.

Tian, D., Araki, H., Stahl, E., Bergelson, J., & Kreitman, M. (2002). Signature of balancing
selection in Arabidopsis. Proceedings of the National Academy of Sciences, 99(17), 11525-11530.

Tsunewaki K (1960) Monosomic and conventional analyses in common wheat. III.
Lethality. Jpn J Genet 35:71–75

Turelli, M. (2017). Commentary: Fisher's infinitesimal model: A story for the ages.
Theoretical population biology, 118, 46-49.

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. Journal of
dairy science, 91(11), 4414-4423.

Vazquez, M. D., Zemetra, R., Peterson, C. J., Chen, X. M., Heesacker, A., & Mundt, C. C.
(2015). Multi-location wheat stripe rust QTL analysis: genetic background and epistatic
interactions. Theoretical and Applied Genetics, 128(7), 1307-1318.

Vilas, A., Pérez-Figueroa, A., & Caballero, A. (2012). A simulation study on the
performance of differentiation-based methods to detect selected loci using linked neutral
markers. *Journal of Evolutionary Biology*, *25*(7), 1364-1376.

Vitezica ZG, Legarra A, Toro MA, Varona L (2017) Orthogonal estimates of variances for
additive, dominance, and epistatic effects in populations. Genetics 206:1297–1307.

Wade, M. J. (2002). A gene's eye view of epistasis, selection and speciation. Journal of
Evolutionary Biology, 15(3), 337-346.

Wade, M. J., & Goodnight, C. J. (1998). Perspective: the theories of Fisher and Wright in
the context of metapopulations: when nature does many small experiments. Evolution, 52(6),
1537-1553.

Weinreich, D. M., Watson, R. A., & Chao, L. (2005). Perspective: sign epistasis and genetic
costraint on evolutionary trajectories. Evolution, 59(6), 1165-1174.

Whitlock, M. C., Phillips, P. C., Moore, F. B. G., & Tonsor, S. J. (1995). Multiple fitness
peaks and epistasis. Annual review of ecology and systematics, 601-629.

Wickham H (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York. ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.

Wiuf, C., Zhao, K., Innan, H., & Nordborg, M. (2004). The probability and chromosomal
extent of trans-specific polymorphism. Genetics, 168(4), 2363-2372.

Wright, S. (1932). The roles of mutation, inbreeding, crossbreeding, and selection in

evolution.

Wright, S. (1988). Surfaces of selective value revisited. The American Naturalist, 131(1),

115-123.

Zhao, J., Zheng, X., Qiao, L., Yang, C., Wu, B., He, Z., ... & Qi, Z. (2022). Genome-wide

association study reveals structural chromosome variations with phenotypic effects in wheat

(Triticum aestivum L.). The Plant Journal, 112(6), 1447-1461.

Volis, S., Shulgina, I., Zaretsky, M., & Koren, O. (2011). Epistasis in natural populations of

a predominantly selfing plant. Heredity, 106(2), 300-309.

Clegg, M. T., Allard, R. W., & Kahler, A. L. (1972). Is the gene the unit of selection?

Evidence from two experimental plant populations. Proceedings of the National Academy of

Sciences, 69(9), 2474-2478.

Dobzhansky, T. (1970). Genetics of the evolutionary process (Vol. 139). Columbia

University Press. James, J. W. (1970). The founder effect and response to artificial selection.

Genetics Research, 16(3), 241-250.

James, J. W. (1970). The founder effect and response to artificial selection. Genetics

Research, 16(3), 241-250.