Progressive neuroinflammation and deficits in motor function in a mouse model with an

***Epg5* pathogenic variant of Vici syndrome**

Bradley T. Thornton[1], Alexandra G. Hardinger[2], Laramie Pence[3], Priyanka Prem Kumar[1],

Nikolas Connolly[4], Scott J. Weir[5,6], Jay L. Vivian[1,3,7,8]

[1]Division of Clinical Genetics, Department of Pediatrics, Children's Mercy Kansas City, Kansas

City MO USA

[2]Bioengineering Graduate Program, University of Kansas, Lawrence KS USA

[3]Genetically Engineered Mouse Models Core, Children's Mercy Research Institute, Children's

Mercy Kansas City, Kansas City MO USA

[4]Interdisciplinary Graduate Program in Biomedical Sciences, University of Kansas Medical

Center, Kansas City, KS, USA

[5]Department of Pharmacology and Toxicology, and Therapeutics, University of Kansas Medical

Center, Kansas City, KS, USA

[6]Institute for Advancing Medical Innovation, University of Kansas Medical Center, Kansas City,

KS, USA

[7]University of Missouri-Kansas City, Kansas City MO USA

[8]Author for correspondence:

jvivian@cmh.edu

Abstract

Background: Vici syndrome (VS) is a rare pediatric genetic disorder characterized by profound developmental delay, seizures, immune deficits, cardiomyopathy, and progressive motor dysfunction, with a median survival of approximately 42 months. This devastating condition is caused by pathogenic variants in the *EPG5* gene, which encodes a key regulator of autophagy. Loss of EPG5 function leads to the accumulation of toxic intracellular material and widespread cellular dysfunction. Currently, there are no effective treatments for VS. The underlying cellular mechanisms driving the progressive neurological decline in VS remain poorly understood. Although previous studies using *Epg5* knockout models have demonstrated severe neurological phenotypes, these models have not been comprehensively characterized for molecular and cellular deficits within the central nervous system.

Methods: We report the generation and analysis of novel genetically engineered mouse models of VS, including a strain harboring a truncating mutation that recapitulates a pathogenic variant identified in a VS patient and a strain with an *Epg5* null allele. These mutant models were used to further understand the underlying molecular deficits of the nervous system.

Results: These novel *Epg5* mutant mouse models exhibited incompletely penetrant perinatal lethality and neurological deficits were detectable by six weeks of age, which worsen over time. Histological analysis revealed widespread expansion of microglia and astrocytes throughout the central nervous system (CNS). Transcriptomic profiling of CNS tissue revealed robust neuroinflammatory signatures, implicating the activation of both microglia and astrocytes. Notably, the molecular profiles of *Epg5*-deficient mice share features with disease-associated microglia observed in other models of neurological disease and injury.

Conclusions: The analysis of these novel mouse models of Vici syndrome suggest a critical role

for neuroglial activation in the pathogenesis of VS. These novel in vivo models will be a

valuable platform for preclinical evaluation of therapeutic strategies targeting autophagy-related

neurodegeneration in congenital disorders of autophagy.

Keywords:

Vici syndrome

EPG5

Autophagy

Microglia

Astrocyte

Neurodegeneration

Congenital disorders of autophagy

Pathogenic variant

Background

Vici syndrome (VS; OMIM 242840; also known as EPG5-related disorder) is a rare,

autosomal recessive, childhood-onset multisystem disorder that primarily affects the central

nervous system, cardiac function, and immune response, among other organ systems (1, 2).

Clinical manifestations typically emerge within the first year of life and include failure to thrive,

global developmental delay, recurrent infections, and cardiomyopathy. Neurological features

include hypotonia, seizures, progressive motor decline, and structural brain abnormalities.

Affected children are typically nonverbal and unable to ambulate. VS follows a relatively

progressive course, with a median survival of approximately 42 months; however, atypical cases

with survival into early adulthood have been reported (unpublished observations). Mortality is

commonly attributed to a combination of progressive cardiomyopathy, recurrent infections, and

respiratory compromise due to neuromuscular decline. Neurological deterioration is a nearly

universal hallmark of this disease and significantly impairs quality of life. Currently, no disease-

modifying therapies exist, and clinical management remains largely supportive and palliative.

The rarity of VS, coupled with limited access to patient-derived samples and the absence of

well-characterized disease models, has hindered both mechanistic understanding and

therapeutic development.

A definitive diagnosis of VS is established through genetic testing, which reveals

pathogenic variants in the *EPG5* gene (3). *EPG5* (Ectopic P-granules protein 5) encodes a

critical regulator of autophagy, facilitating the fusion of autophagosomes with lysosomes(4).

Most pathogenic variants identified in VS are nonsense or frameshift mutations resulting in loss

of function, although some missense mutations have also been reported(5). Patient-derived

cells exhibit impaired autophagic flux, characterized by the accumulation of undegraded

autophagic material due to defective *EPG5*-mediated vesicle fusion (6, 7). VS is now recognized

as part of a broader group of rare disorders(8) termed congenital disorders of autophagy

(CDAs). Deficits in autophagy may drive a range of neurological disorders and other diseases

(9, 10), suggesting new avenues for treatment using targeted therapies that modulate

autophagy.

Previous studies using *Epg5* knockout models have demonstrated severe neurological

phenotypes(11); however, the molecular and cellular mechanisms underlying the progressive

motor deficits in VS remain poorly understood. Moreover, no existing mouse models that

recapitulate patient-specific *EPG5* mutations have been reported. To address this gap, we

generated and characterized two novel *Epg5* mutant mouse strains: one harboring a truncating

W860X pathogenic variant identified in a patient (case report in preparation) and another

carrying a newly developed null allele. Both models exhibit early-onset and progressive

neurological phenotypes, with subtle but significant differences between alleles. Whole animal

deficits in motor function were are evident by six weeks of age and worsen over time.

Transcriptomic analyses of brain tissue from these models revealed robust neuroinflammatory

signatures, implicating glial activation, particularly of microglia and astrocytes, in disease

progression. Notably, the molecular profiles observed in *Epg5*-deficient mice share features with

disease-associated microglia (12) identified in models of neurodegenerative disorders such as

Alzheimer's disease and amyotrophic lateral sclerosis, suggesting potential mechanistic

overlap. These findings enhance our understanding of VS pathogenesis and provide candidate

biomarkers for disease monitoring. Furthermore, these models establish a critical platform for

preclinical evaluation of therapeutic strategies targeting this devastating disorder.

Methods

Animal use

All animal procedures were approved by the Children's Mercy Research Institute Institutional

Animal Care and Use Committee (IACUC). All housing, experimental procedures, and

monitoring fully conformed to the Guide for the Care and Use of Laboratory Animals, and all

euthanasia procedures conformed with the American Veterinary Medical Association's

guidelines for euthanasia. The mice were generated and maintained on a C57BL/6J genetic

background, and the mutant stocks were maintained as heterozygotes by backcrossing to

vendor-derived animals (Jackson Laboratories, Bar Harbor ME). Homozygous mutant mice

were closely monitored for animal welfare concerns related to neurological dysfunction and

removed from the study via approved euthanasia methods if any effects on welfare were

identified. The animals were housed in a specific pathogen-free vivarium operating as a barrier

facility in single-use disposable mouse cages (Innovive, San Diego CA) with a 12-hour light/dark

cycle and maintained at 22°C +/-1.0°C and 40%+/-5% humidity. Pregnant dams and preweaning

pups were fed high-fat chow and water ad libitum, and postweaning aged mice were fed

standard laboratory chow and water ad libitum.

Generation of mouse strains with *Epg5 W860X* and *Epg5* knockout alleles.

The design and validation of the CRISPR reagents used for in vivo genome editing were

performed as described(13). To generate the *Epg5-W860X* allele, a CRISPR target site

(CCCGAATCACAGCTATCTCT) was identified in exon 14 of the mouse *Epg5* locus. To generate

the *Epg5*-KO allele, a CRISPR target site (CGGGCTCTCCTGCTCGCGAC) was identified at

the 5' end of exon 2 of the mouse *Epg5* locus. CRISPR reagents were synthesized as crRNAs

and annealed to universal tracrRNAs (Integrated DNA Technologies, Coralville IA). Prior to

production, the activity of the CRISPR reagents was verified via embryo electroporation of the

Cas9/crRNA/tracrRNA ribonucleoprotein (RNP) complex. The *Epg5*-W860X mutant mice were

produced as described(14) via CRISPR-mediated genome editing and homology-directed

repair. Single-stranded donor DNA was designed for each allele (Supplemental Table 1) and

generated via DNA synthesis (Integrated DNA Technologies, Coralville IA). For *Epg5*-W860X,

the donor DNA incorporated a single nucleotide change, resulting in a premature stop codon in

exon 2 flanked by 88 bp and 68 bp of *Epg5* homology for targeting via homology-directed repair.

The CRISPR RNP complexes were coinjected with donor DNA into C57BL/6J zygotes, which

were subsequently transferred to pseudopregnant females for delivery. The resulting founders

were confirmed for correct targeting of the donor DNA via allele-specific PCR and sequencing

(Supplemental Table 1). To produce the *Epg5*-KO strain, the CRISPR reagents were

electroporated into C57BL/6J zygotes, which were subsequently transferred to pseudopregnant

females for delivery. The resulting founders were analyzed for deletions generated via

nonhomologous end-joining via PCR, and the nature of the resulting mutations was confirmed

by next-generation sequencing of PCR amplicons. One founder harboring a 22-nucleotide

deletion in exon 2 resulting in a frameshift mutation was expanded by breeding to generate wild-

type C57Bl/6J mice. Germline transmission of each desired targeted allele was confirmed in F1

generation pups via PCR and sequencing.

Behavior analysis

The analysis of motor coordination, endurance, and fatigue resistance was performed via the

rotarod test (Panlab, Barcelona, Spain). The mice were acclimated and housed in rooms with an

adjusted light–dark cycle to allow for testing during the active dark phase. Test subjects were

given two training days with the apparatus set at 4 rpm for 1 minute for three runs each training

day. One week after training, the mice were tested every two weeks with the apparatus set at a

ramp speed from 4–40 rpm over the course of 300 seconds. The mice were tested for three

trials per testing day with an intertrial interval of 10–15 minutes. Failure to initiate was defined as

a trial in which the animal was unable to remain on the rod for one full revolution at 4 rpm at the

beginning of a test. The latency to fall and speed at fall data from each trial were recorded via

SEDACOM 2.0 software (Panlab, Barcelona, Spain). Data visualization and statistical analysis

were performed via Prism software (GraphPad, Boston, MA). Statistical analysis of data from

wild-type and mutant mice at a given timepoint was performed via an unpaired t test.

Protein analysis

Spinal cord tissues from 6-month-old wild-type and mutant siblings of each of the *Epg5* mutant

strains were dissociated in RIPA buffer (Thermo Fisher, Waltham, MA.) with a TissueRuptor

homogenizer (Qiagen, Hilden, Germany.). The supernatant was collected, and a protease and

phosphatase inhibitor cocktail (Thermo Fisher, Waltham, MA) was added. The protein

concentration of the supernatant was quantified via a Pierce BCA assay (Thermo Fisher,

Waltham, MA) per the manufacturer's protocol. Protein samples were denatured in Laemmli

sample buffer (Bio-Rad Laboratories, Hercules, CA), and 20 µg of protein from each sample

was used for western blot analysis. Antibodies against LC3 (Cell Signaling Technology, Danvers,

MA) and a-Tubulin (Cell Signaling Technology, Danvers, MA) and secondary antibodies (LI-COR

Biosciences, Lincoln, NE) were used to detect each protein. Fluorescent imaging was

performed on the membrane via an Odyssey CLx Imager (LI-COR Biosciences, Lincoln, NE).

Each band was quantified via ImageStudioLite analysis software (LI-COR Biosciences, Lincoln,

NE).

RNA analysis

Gene expression analyses were performed via purified RNA from tissues harvested from 6-

month-old wild-type and homozygous mutants of each of the *Epg5* strains. The tissues were

lysed via Qiazol/chloroform extraction, and the RNA was purified via a Qiagen RNeasy column

purification kit (Qiagen, Hilden, Germany) according to the manufacturer's protocols. The

extracted RNA was then either given to the institutional genomics core for RNA sequencing or

was converted into cDNA via the use of 1 µg of total RNA for quantitative PCR. The qPCRs

were performed using TaqMan Gene Expression Master Mix (Thermo Fisher, Waltham, MA) and

the indicated probes (Supplemental Table 2). The samples were analyzed via the ViiA 7 real-

time PCR system (Applied Biosystems, Waltham, MA) according to the manufacturer's

protocols. For plotting, the expression data were normalized to the data of the wild-type control

siblings.

RNA-Seq bioinformatics analysis

Transcriptomic profiling via bulk RNA sequencing was performed on RNA from the cerebellum

and spinal cord of homozygous W860X mice or their wild-type control siblings. RNA sequencing

was performed with stranded total RNA with a RiboZero Gold library preparation kit (Illumina,

San Diego CA) per the manufacturer's protocols. The RNA sequences were processed for gene

expression quantification. Standard two-group differential expression analysis of a set of RNA-

seq samples processed with Kallisto(15). Output files were processed with tximport(16) to

collapse transcript counts to gene counts via the transcript-to-gene mapping table. The mouse

reference genome GRCm39 was used in the analysis. To generate a volcano plot, the

processed average differential expression data from the RNA-seq data consisting of 13,991

genes in the cerebellum and 25,527 genes in the spinal cord were inputted into Prism

(GraphPad, Boston, MA). For comparative differential gene expression analysis between

tissues, genes that had a $\log_2$(fold change) of +1/-1 and a false discovery rate (adj. p value) of

less than 0.01 were used. Overlapping DEGs between tissues were identified via Venny

(https://bioinfogp.cnb.csic.es/tools/venny/index.html). The overlapping genes were then input

into Ingenuity Pathway Analysis (Qiagen, Hilden, Germany) for pathway analysis and to

g:Profiler(17) for Gene Ontology (GO) analysis. Altered pathways and associated p values were

visualized via Prism (GraphPad, Boston, MA).

Histology

For histology, the tissues were processed as previously described(18). In brief, 6-month-old

mutant or control mice were placed under anesthesia (isofluorane) for transcardial perfusion via

4% paraformaldehyde in PBS. CNS tissues were then removed and fixed overnight. The tissues

were then washed two times in 1X PBS prior to preparation for histology.

For immunohistochemistry, the tissues were dehydrated and embedded in paraffin. The

10 μm sections were rehydrated via citrate antigen retrieval buffer for use in staining to detect

GFAP (HRP/DAB IHC Detection Kit, Abcam, Cambridge, UK) per the manufacturer's protocol. A

colorimetric reaction was performed using DAB.

For immunofluorescence imaging, the tissues were frozen for cryosectioning as

described(19). In brief, 20 μm sections were cut. Direct conjugated antibodies to detect GFAP or

IBA-1 were used to detect astrocytes and microglia, respectively, and costained with DAPI to

detect nuclei. The sections were mounted with a coverslip with Fluormount G mounting media

(Thermo Fisher). Immunofluorescence imaging was performed on a Zeiss Axio Observer 7

microscope (Zeiss, Oberkochen, Germany). For quantification, images of distinct regions of the

spinal cord were imported into the Imaris software package (Oxford Instruments, Oxfordshire

UK). For each fluorophore, an absolute intensity threshold value was adjusted via the histogram

and standardized for each image. A new surface was created to measure the total area of the

image. The summed value of the combined area from each surface was divided by the total

area to obtain the percentage value of positivity for each image.

Results

To develop mouse models that recapitulate a pathogenic *EPG5* variant associated with

Vici syndrome, a novel mouse strain was generated via CRISPR-mediated zygotic mutagenesis

(ref) on an inbred C57Bl/6J genetic background (Table 1; Supplemental Table 1). One strain

was engineered to carry the *Epg5 W860X* mutation, which mirrors a recently identified

pathogenic variant in a child with severe Vici syndrome (manuscript in preparation). For

comparison, a knockout (*KO*) strain was also generated by introducing a 22 bp interstitial

deletion near the start of exon 2, resulting in a frameshift and predicted early truncation of

the EPG5 protein. Following generation and colony expansion, each strain was intercrossed to

produce homozygous mutant mice.

Genotypic analysis of offspring at postnatal day 14 (P14) revealed an

underrepresentation of homozygous mutants for both the *W860X* and *KO* alleles (Table 2). Chi-

square analysis indicated that this loss was due to a reduced number of surviving males (p =

0.043 for *KO*; p = 0.035 for *W860X*). A small number of deceased pups at P1 were found to be

homozygous mutants, suggesting perinatal lethality associated with these alleles.

Surviving homozygous mutants were monitored longitudinally postweaning. Mice

homozygous for either the *W860X* or *KO* mutation were generally smaller than their wild-type

littermates, although only *W860X* mutants consistently exhibited reduced body size throughout

life (Figure 1A, B; Supplemental Figure 1). All the homozygous mutants displayed marked motor

deficits, particularly in hindlimb coordination. Compared with age-matched controls, mutant mice

exhibited abnormal hindlimb posture, characterized by a splayed stance and inability to support

body weight (Figure 1C, D). A moderate ataxic phenotype was evident, with wobbly, halting, and

slow gaits (Supplemental Movie 1). Owing to progressive motor impairment, most of the mice

were removed from the study by 7 months of age for animal welfare reasons (Figure 1E). No

sex differences were observed in the severity of motor deficits.

To quantitatively assess motor function, rotarod testing was conducted longitudinally

beginning at 8 weeks of age. Compared with their control littermates, both *W860X* and *KO*

homozygous mutants presented significantly impaired performance (Figure 1F-I). By 12 weeks,

some mutants were unable to remain on the rotating cylinder, and by 14 weeks, all *W860X*

homozygous mutants and, by 16 weeks, all KO homozygous mutants were unable to initiate the

test. Both male and female mutants were equally affected. Owing to the severity of motor

impairment, mutants were unable to perform additional coordination tests, such as beam

walking. These findings of deficits in motor function were similar to previous analysis of other

Epg5 mutant mouse models(11). These studies indicate early-onset and progressive deficits in

survival, growth, and motor function in both *Epg5* mutant strains, with modest differences in

severity between the two mutant alleles analyzed.

Tissues from each mutant strain were collected for transcript and autophagy analyses.

Compared with those of wild-type controls, RNA extracted from the spinal cords of *W860X*

homozygous mice presented significantly lower *Epg5* transcript levels (Figure 2A). Although not

statistically significant, a similar trend was observed in the cerebellum (Figure 2B), suggesting

tissue-specific transcript instability, potentially due to nonsense-mediated decay. In contrast, the

transcript levels in the *KO* mice were not reduced. Western blot analysis of the LC3 protein

(microtubule-associated protein 1 light chain 3 alpha) was used to assess autophagy (Figure

2C,D; Supplemental Figure 2). Analysis of LC3 via western blotting distinguishes the

cytoplasmic form of LC3 (LC3-I) from the autophagosome-localized LC3 (LC3-II). LC3-II is

subject to autophagy-mediated degradation; this the steady state levels of LC3-II allow for an

assessment of autophagy. The autophagosome-associated LC3-II was markedly elevated in

spinal cord lysates from both mutant strains compared to tissues from wildtype controls,

indicating that autophagic flux was impaired and autophagosome accumulation.

To characterize the neurological phenotype further, bulk RNA sequencing was performed

on cerebellar and spinal cord tissues from 6-month-old *W860X* homozygous mice and control

littermates. These CNS tissues were selected because of their roles in motor control. The

transcriptomic profiles (n=4 of each genotype, split evenly by sex) were consistent between the

sexes and were combined for this analysis. A substantial number of DEGs were identified, with

more genes upregulated than downregulated (Figure 3A, B). Notably, genes associated with

glial activation, including microglial markers (*Gpnmb*, *Itgax*) and the astrocytic marker *Gfap* were

upregulated, suggestive of a gliosis in these tissues. Several Toll-like receptor genes, key

components of innate immune signaling, were also upregulated. Among the downregulated

genes, *Myoc* (Myocilin) was nearly absent in the spinal cord of *W860X* mutants but unchanged

in the cerebellum. This gene is specifically expressed in the *glia limitans superficialis*, a

specialized astrocyte population localized to the thin surface layer throughout the CNS.

Previous studies have identified distinct molecular signatures associated with microglial

responses to neurological injury in various mouse disease models (12) and in human

neurodegenerative disorders (20). Among these, a disease-associated microglia (DAM)

signature, also referred to as the neurodegenerative microglial phenotype (MGnD), has been

characterized by the upregulation of genes such as *Clec7a*, *Cst7*, and *Trem2*(12).

Transcriptomic analysis of CNS tissues from *Epg5 W860X* homozygous mutant mice revealed

upregulation of many DAM-associated genes (Figure 3D), suggesting a glial response

reminiscent of other CNS injury models.

Although distinct sets of differentially expressed genes (DEGs) were identified in the

cerebellum and spinal cord of *Epg5 W860X* mutants, a shared gene expression signature was

also observed across both tissues (Figure 3C; Supplemental Table 3). Analysis of these

overlapping DEGs highlighted key cellular pathways affected by the *W860X* mutation in CNS

tissues (Figure 3E, F). Gene ontology and pathway enrichment analyses confirmed that these

shared DEGs were significantly associated with inflammatory and immune responses.

Pathways identified in this analysis include cytokine, interferon, and complement inflammatory

pathways. Cellular pathways identified as neutrophil degranulation and phagosome formation

may suggest components of a microglial activation. Although several pathways were identified

with pathogen response, this analysis likely does not represent an inflammatory or cellular

response to microbial insult, given the specific pathogen-free status of the vivarium within which

these animals were housed. Collectively, these transcriptomic data define a novel and complex

neuroinflammatory and glial activation signature in mice carrying the *Epg5 W860X* pathogenic

variant that is reminiscent of other neurological disease or damage models involving neuroglia.

To further evaluate the expression of selected DEGs, RNA was isolated from the CNS

tissues of 7-month-old mutant and control mice for quantitative PCR (qPCR) analysis (Figure 4).

Genes were selected on the basis of the transcriptomic findings, with a focus on glial markers

(astrocytic and microglial), inflammatory mediators, interferon response genes, and other

relevant targets. All DEGs identified in the *W860X* model were also dysregulated in

the *Epg5 KO* mice, although with varying magnitudes (Figure 4A–D; Supplemental Figure 3,4).

For example, *Gpnmb* and *Gfap* expression was more severely elevated in *W860X* mutants than

in *KO* mice, whereas other genes presented similar expression levels across both models.

Notably, the severity of gene expression changes varied by tissue. *Tlr7* expression, for example,

was more severely affected in the cerebellum of *W860X* mutants, whereas spinal cord

expression levels did not differ significantly between the two models. These molecular findings

parallel the phenotypic severity observed in vivo, with similar levels of severity between the

mutant alleles, with the *W860X* mutants exhibiting somewhat more pronounced deficits than *KO*

mice.

To assess the temporal progression of the gene expression deficits associated with the

Epg5 mutations, the expression of a subset of glial-specific genes was evaluated in 2- and 6-

month-old *W860X* and *KO* mutant mice compared with wild-type controls (Figure 5;

Supplemental Figure 5). The selected genes included the astrocyte marker *Gfap* and the

microglial markers *Gpnmb* and *Itgax*. Many of these genes were already significantly

upregulated at 2 months of age, particularly in the spinal cord of both mutant models (Figure

5A–D), and their expression increased further by 6 months. In contrast, the changes in gene

expression in the cerebellum of the *KO* mice were relatively modest at 2 months but became

more pronounced with age. These results indicate that glial phenotypes in *Epg5 W860X*

mutants were already apparent at early adulthood stages and progress with age, which is

consistent with the age of observed deficits in motor function.

Given the strong glial signature observed via transcriptomic and qPCR analyses,

histological studies were performed to assess astrocyte and microglial distribution and

phenotype in CNS tissues from 7-month-old *Epg5 W860X* mutants and control wildtype

littermates. Immunohistochemistry revealed a substantial increase in GFAP-positive astrocytes

across all CNS regions examined, which was consistent with elevated *Gfap* transcript levels

(Figure 6A, B; Figure 7A–D). Immunofluorescence staining for GFAP and the microglial marker

IBA1 confirmed widespread astrocyte and microglial expansion in the spinal cord (Figure 6C–N)

and cerebellum (Figure 7E–G) in *Epg5* mutant mice. Microglia exhibited a hypertrophic, spider-

like morphology with extended processes (Figure 6G, arrowhead), indicative of activation. In the

cerebellum, increased numbers of astrocytes and microglia were observed across multiple

layers, including the molecular layer (ML), internal granule layer (IGL), and white matter (WM)

(Figure 7F, G). Despite these cellular changes, the overall cerebellar architecture remained

intact.

While glial expansion was broadly distributed, certain white matter regions presented

particularly dense microglia and astrocyte accumulation. In the spinal cord, microglia were

notably concentrated in the dorsal funiculus (Figure 6I–K), a region containing descending

corticospinal and ascending sensory tracts. Additionally, both astrocytes and microglia were

enriched along the surface of the spinal cord (Figure 6C, D, L–N). While astrocytes are normally

present in this region, their numbers and process extension were markedly increased in mutants

(Figure 6D, M). Microglia, which are typically sparse at the surface, were also significantly

expanded (Figure 6M). These findings demonstrate a robust and progressive glial response in

the CNS of *Epg5 W860X* mutant mice, which is correlated with the observed motor deficits and

supports a role for glial dysfunction in the pathogenesis of Vici syndrome.

Discussion

The complex etiology of Vici syndrome (VS) remains incompletely understood,

contributing to the current lack of effective therapeutic strategies. In particular, the molecular

and cellular basis for the neurological manifestations of VS, which are universally present in VS

individuals, remain poorly characterized. This study presents the first published analysis of a

novel mouse model harboring a pathogenic variant in *Epg5* that mirrors a mutation identified in

VS patients. Specifically, the *Epg5-W860X* allele recapitulates the *EPG5-W865X* nonsense

mutation found in a clinically severe VS patient (manuscript in preparation). Homozygous

W860X mutant mice exhibit profound motor deficits that are comparable in severity to those

observed in both the *Epg5* knockout (*KO*) mice analyzed in this study and the previously

reported *Epg5*-deficient models(11).

Focusing on the molecular and cellular underpinnings of neurological dysfunction, our

analysis revealed previously unreported CNS phenotypes in these models. Using established

molecular markers, we observed widespread and progressive expansion of astrocytes and

microglia throughout the CNS in *Epg5* mutant mice. This glial expansion increases with age and

correlates with the severity of motor dysfunction, with significant motor deficits and glial marker

upregulation evident by two months of age. Notably, a molecular signature of

neuroinflammation, characterized by astrocyte and microglial expansion, suggests a

contributory role for glial cells in VS pathogenesis and disease progression. Previous studies

using *Epg5* mutant mice revealed inflammatory and immune-related phenotypes (21, 22),

including gastrointestinal and peripheral immune dysfunction. In conjunction with these prior

studies, the findings in this study suggest the presence of an inflammatory phenotype across

multiple organ systems in *Epg5*-mutant mice, underscoring the potential for inflammation

potentially contributing to the multisystemic nature of the disease. Specific CNS molecular

markers may thus serve as indicators of disease progression and severity.

Comparative analysis of the mutant alleles confirmed that the *W860X* and *KO* alleles

presented severe loss-of-function mutations. This aligns with the clinical severity observed in

the patient harboring the corresponding *W865X* mutation, who succumbed to the disease at 16

months of age (clinical report in preparation). The reduced *Epg5* transcript levels in *W860X*

mutant mice suggest that the mRNA is subject to nonsense-mediated decay. Owing to the lack

of a high-quality antibody for mouse EPG5, protein levels associated with the mutations

generated in this report could not be assessed via western blot. However, unpublished data

from patient-derived cells indicate that the *W860X* pathogenic variant does not produce a stable

truncated protein, further supporting its classification as a null allele.

A novel and striking observation in our study was the reduced survival of male mice

homozygous for either the W860X or KO allele, an outcome not previously reported in other

Epg5 mutant models. This discrepancy may reflect differences in experimental design, such as

the depth of analysis of progeny from heterozygous intercrosses or potential residual activity of

the allele used in earlier studies. In humans, VS is associated with early-onset hypotonia and

failure to thrive. The reduced perinatal survival of homozygous mutant mouse pups may mirror

this clinical phenotype. Further investigations are needed to elucidate the mechanisms

underlying the observed male-specific lethality. Interestingly, previous studies reported sex-

dependent differences in microglial abundance and activity during both embryonic development

and adulthood(23). Future analyses of *Epg5* mutant tissues at late prenatal and early postnatal

stages will be critical for understanding the origins of this lethality and defining the earliest

manifestations of glial dysregulation associated with the defective autophagy in this model.

One of the most striking transcriptional changes observed in the spinal cord of *Epg5*

mutant mice was the downregulation of *myocilin (Myoc)*, a marker of a specialized astrocyte

subtype known as the *glia limitans superficialis* (GLS)(24). GLS astrocytes are localized to the

pial surface of the brain and spinal cord and are thought to contribute to barrier function and

immune surveillance. *Myoc* expression is initiated postnatally in these cells, suggesting Myoc is

a marker for developmental maturation of GLS cells. The substantial downregulation of *Myoc* in

Epg5 mutant mice implicates a requirement for autophagy, either directly or indirectly, in the

survival or maturation of GLS astrocytes. Interestingly, our histological analysis revealed an

expansion of astrocytes and microglia at the surface of the spinal cord, suggesting that reactive

gliosis may interfere with GLS maturation. Further studies are needed to determine whether this

represents a cell-autonomous defect in GLS astrocytes or a secondary effect of microglial

activation and inflammation. Understanding the role of GLS astrocytes in CNS homeostasis and

their potential involvement in VS pathogenesis could provide new insights into how autophagy

deficits disrupt glial development and function.

Astrocytes and microglia are essential for maintaining central nervous system

homeostasis and play increasingly recognized roles in modulating disease progression.

Astrocyte homeostatic functions include synaptic maintenance, neurotransmitter recycling,

integrity of the blood–brain barrier, and metabolic support of neurons. Microglia, the resident

innate immune cells of the CNS, are responsible for the phagocytosis of cellular debris,

apoptotic cells, and pathogens. These glial cells not only are passive support cells but are also

highly dynamic, responding to environmental cues and pathological states with complex

molecular and functional changes. In disease or injury contexts, the functional plasticity of these

cell types, which can adopt a range of activation states, including proinflammatory and

neuroprotective phenotypes, has been revealed. Importantly, these cell types can adopt

phenotypes that may be either neuroprotective or neurotoxic depending on the context and

duration of activation. Both astrocytes and microglia can secrete cytokines and chemokines

depending on the activation state, which influences the local microenvironment and contributes

to neuroinflammatory cascades. Microglia play a critical role in synaptic pruning during

development and in response to injury.

Whole-transcriptome analysis of CNS tissues from *Epg5* mutant mice revealed a

signature of activated microglia resembling other mouse CNS injury or disease models (12, 25).

This distinct transcriptional profile, known as disease-associated microglia (DAM), also referred

to as the neurodegenerative microglial phenotype (MGnD), has been identified in multiple

models of neurodegeneration or injury. This phenotype is characterized by the upregulation of

genes such as *Apoe*, *Trem2*, and *Cst7*, which are not typically induced in classical inflammatory

responses such as lipopolysaccharide (LPS) exposure. DAM microglia are thought to play

complex roles in disease progression, being initially protective via microglial phagocytic activity

by clearing debris, but there is evidence that they may contribute to disease progression if

chronically activated. Future studies involving single cell analysis of microglia in the *Epg5*

mutant mice will be needed to uncover the spectrum of the microglial phenotypes in these mice

and the regional differences in microglial states within the CNS, and how these may change

over the course of the lifespan. There is growing appreciation for the dual roles of glial cells in

contributing to the pathogenesis of neurological disorders. Dysregulation of glial function has

been implicated in a wide range of neurodegenerative diseases(26-28), including Alzheimer's

disease (AD), Parkinson's disease (PD), and amyotrophic lateral sclerosis (ALS). In models of

AD, PD, and ALS, microglial activation has been shown to be correlated with disease

progression. For example, *APOE* alleles associated with increased AD risk have been linked to

altered microglial responses(29), whereas *SOD1* mutations in ALS models have demonstrated

that glial dysfunction can exacerbate motor neuron degeneration(30). These findings

underscore the importance of glial cells not only as responders to neuronal injury but also as

active participants in disease mechanisms.

The complex neuroinflammatory signature seen in *Epg5* mutant mice suggest that glial

dysfunction may be a key driver of disease progression in Vici syndrome. The localized

accumulation of microglia in specific regions of the spinal cord in *Epg5* mutant mice, specifically

in white matter, may point to possible cellular mechanisms for a role of microglia in neuronal

function in this model. Notably, microglial localization to specific CNS regions, such as the

corticospinal tract in models of ALS(31), has been associated with region-specific vulnerability.

Whether spatial differences in glial activation in *Epg5* mutant mice or VS remains to be

determined. The neurological phenotypes observed in *Epg5* mutant mice likely arise from a

combination of neuronal and glial dysfunction driven by impaired autophagy. Future studies

using primary cell cultures from these mice, as well as human induced pluripotent stem cell

(iPSC) models derived from VS patients(32), will be essential for dissecting the cell type-specific

consequences associated with EPG5 loss. The development of new mouse and cellular models

that allow for tissue-specific manipulation of autophagy will be critical for determining the extent

to which glial dysfunction contributes to disease progression in VS and other congenital

disorders of autophagy. Ultimately these models will be critical to determine the molecular

mechanisms that give rise to the complex cellular deficits seen in Vici syndrome. These studies

will have major implications into potential therapeutic targets.

Conclusions

This study presents a detailed analysis of novel mouse models of Vici syndrome, including a

strain harboring a pathogenic variant in *Epg5* that mirrors a mutation identified in VS patients.

These mutant strains displayed an incomplete penetrant perinatal lethality, and significant motor

deficits of surviving mice were evident by two months of age. Our transcriptomic analysis of

CNS tissues revealed a complex molecular signature of neuroinflammation. Widespread and

progressive expansion of astrocytes and microglia was observed throughout the CNS in these

mutant mice, and this glial activation correlated with age and the severity of motor dysfunction.

The detailed analysis of these mouse models suggests a contributory role for glial cells in the

pathogenesis and disease progression of Vici syndrome. These studies validate the utility of

these mouse models for study the progression of VS, and for use in therapeutic assessment for

this devastating disorder, including the potential for targeting neuroglia as therapeutic strategy.

Abbreviations

EPG5: Ectopic P Granules 5

VS: Vici syndrome

AD: Alzheimer's disease

PD: Parkinson's disease

ALS: Amyotrophic lateral sclerosis

iPSC: induced pluripotent stem cell

CNS: central nervous system

GLS: glia limitans superficialis

DAM: Disease-associated microglia

MGnD: neurodegenerative microglia

Data availability

The RNA sequencing datasets supporting the conclusions of this study will be deposited in the

NCBI Gene Expression Omnibus. GEO accession numbers will be provided upon completion of

the deposit. All other data can be found within the article and its supplementary information.

Competing interests

The authors declare that they have no competing or financial interests.

Funding

This study was supported by Mercy Research Partners of Children's Mercy Kansas City, the Vici

Syndrome Foundation, and Vici syndrome families.

Author contributions

All the authors read and approved the final manuscript.

Conceptualization: JLV, SJW

Investigation: BTT, LP, AGH, PPK, NC, JLV

Data curation: BTT,LP, AGH,PPK,JLV

Formal analysis: LP, JLV

Writing – original draft: JLV

Writing – review and editing: BTT,LP, AGH,PPK,JLV

Visualization: BTT, AGH, and JLV

Funding acquisition: JLV

Acknowledgments

The generation of the animal models was performed by the Genetically Engineered Mouse

Models Core of Children's Mercy Research Institute and the Transgenic and Gene Targeting

Institutional Facility of the University of Kansas Medical Center as Shared Resources of the

University of Kansas Cancer Center (NIH P30 CA168524). We acknowledge the efforts of Jing

Huang and Sarah E Tague in the Histology and Integrative Imaging of the Kansas Intellectual

and Developmental Disabilities Resource Center at the University of Kansas Medical Center.

We wish to acknowledge the efforts of cores of the Children's Mercy Research Institute,

including RNA sequencing and analysis were performed by the Genomic Core Service Center of

the Children's Mercy Research Institute, and tissue preparation and sectioning were performed

by the Children's Mercy Pathology Core.

Table 1. Summary of mutant mouse strains
524

Mouse allele	*Epg5* exon	Mutation[1]	human pathogenic variant[2]	human gnomAD variant ID	human RefSNP ID
KO	2	p.R41QfsTer8 (c.122_143del)	n/a	n/a	n/a
W860X	14	p.W860X	W865X (c. 2595G>A)	SNV : 18-45925861 - C-T	rs1160786631

525
526   [1]mouse accession number: NM_001195633.2
527   [2]human accession number: NM_020964.3

Table 2. Summary of the genotypes of offspring from intercrosses of heterozygous *Epg5* mutant mice

		Epg5 allele					
		KO			**W860X**		
		Total	M	F	Total	M	F
genotype	expected	107	56	51	249	126	123
wt	25%	31.8%	32.1%	31.4%	30.1%	29.4%	30.9%
het	50%	55.1%	57.1%	52.9%	51.8%	55.6%	48.0%
mut	25%	13.1%	10.7%	15.7%	18.1%	15.1%	21.1%
p value		0.014	0.043	0.261	0.023	0.035	0.280
significance		*	*	NS	*	*	NS

Total: total number of mice analyzed from heterozygous intercrosses
M: number of male mice
F: number of female mice
*: $p<0.05$ according to the chi square test
NS: not significant

References

1.   Byrne S, Dionisi-Vici C, Smith L, Gautel M, Jungbluth H. Vici syndrome: a review. Orphanet J Rare Dis. 2016;11:21.

2.   Dafsari HS, Ebrahimi-Fakhari D, Saffari A, Deneubourg C, Fanto M, Jungbluth H. EPG5-Related Disorder. In: Adam MP, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, et al., editors. GeneReviews((R)). Seattle (WA)1993.

3.   Cullup T, Kho AL, Dionisi-Vici C, Brandmeier B, Smith F, Urry Z, et al. Recessive mutations in EPG5 cause Vici syndrome, a multisystem disorder with defective autophagy. Nat Genet. 2013;45(1):83-7.

4.   Wang Z, Miao G, Xue X, Guo X, Yuan C, Wang Z, et al. The Vici Syndrome Protein EPG5 Is a Rab7 Effector that Determines the Fusion Specificity of Autophagosomes with Late Endosomes/Lysosomes. Mol Cell. 2016;63(5):781-95.

5.   Kane MS, Vilboux T, Wolfe LA, Lee PR, Wang Y, Huddleston KC, et al. Aberrant splicing induced by the most common EPG5 mutation in an individual with Vici syndrome. Brain. 2016;139(Pt 9):e52.

6.   Hori I, Otomo T, Nakashima M, Miya F, Negishi Y, Shiraishi H, et al. Defects in autophagosome-lysosome fusion underlie Vici syndrome, a neurodevelopmental disorder with multisystem involvement. Sci Rep. 2017;7(1):3552.

7.   Piano Mortari E, Folgiero V, Marcellini V, Romania P, Bellacchio E, D'Alicandro V, et al. The Vici syndrome protein EPG5 regulates intracellular nucleic acid trafficking linking autophagy to innate and adaptive immunity. Autophagy. 2018;14(1):22-37.

8.   Ebrahimi-Fakhari D, Saffari A, Wahlster L, Lu J, Byrne S, Hoffmann GF, et al. Congenital disorders of autophagy: an emerging novel class of inborn errors of neuro-metabolism. Brain. 2016;139(Pt 2):317-37.

9.   Stamatakou E, Wrobel L, Hill SM, Puri C, Son SM, Fujimaki M, et al. Mendelian

neurodegenerative disease genes involved in autophagy. Cell Discov. 2020;6:24.

10.   Klionsky DJ, Petroni G, Amaravadi RK, Baehrecke EH, Ballabio A, Boya P, et al.

Autophagy in major human diseases. EMBO J. 2021;40(19):e108863.

11.   Zhao H, Zhao YG, Wang X, Xu L, Miao L, Feng D, et al. Mice deficient in Epg5 exhibit

selective neuronal vulnerability to degeneration. J Cell Biol. 2013;200(6):731-41.

12.   Barclay KM, Abduljawad N, Cheng Z, Kim MW, Zhou L, Yang J, et al. An inducible

genetic tool to track and manipulate specific microglial states reveals their plasticity and roles in

remyelination. Immunity. 2024;57(6):1394-412 e8.

13.   Larson MA, Gibson KA, Vivian JL. In Vivo Validation of CRISPR Reagents in

Preimplantation Mouse Embryos. Methods Mol Biol. 2020;2066:47-57.

14.   Larson MA. Pronuclear Microinjection of One-Cell Embryos. Methods Mol Biol.

2020;2066:27-33.

15.   Bray NL, Pimentel H, Melsted P, Pachter L. Erratum: Near-optimal probabilistic RNA-seq

quantification. Nat Biotechnol. 2016;34(8):888.

16.   Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. F1000Res. 2015;4:1521.

17.   Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g:Profiler-interoperable

web service for functional enrichment analysis and gene identifier mapping (2023 update).

Nucleic Acids Res. 2023;51(W1):W207-W12.

18.   Bhattacherjee A, Mu Y, Winter MK, Knapp JR, Eggimann LS, Gunewardena SS, et al.

Neuronal cytoskeletal gene dysregulation and mechanical hypersensitivity in a rat model of Rett

syndrome. Proc Natl Acad Sci U S A. 2017;114(33):E6952-E61.

19.   Yang FC, Draper J, Smith PG, Vivian JL, Shapiro SM, Stanford JA. Short Term

Development and Fate of MGE-Like Neural Progenitor Cells in Jaundiced and Non-Jaundiced

Rat Brain. Cell Transplant. 2018;27(4):654-65.

20.   Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R, Ulland TK, et al. A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. Cell. 2017;169(7):1276-90 e17.

21.   Lu Q, Yokoyama CC, Williams JW, Baldridge MT, Jin X, DesRochers B, et al. Homeostatic Control of Innate Lung Inflammation by Vici Syndrome Gene Epg5 and Additional Autophagy Genes Promotes Influenza Pathogenesis. Cell Host Microbe. 2016;19(1):102-13.

22.   Lee S, Kalugotla G, Ingle H, Rodgers R, Wu C, Wang Y, et al. Intestinal antiviral signaling is controlled by autophagy gene Epg5 independent of the microbiota. Autophagy. 2022;18(5):1062-77.

23.   Han J, Fan Y, Zhou K, Blomgren K, Harris RA. Uncovering sex differences of rodent microglia. J Neuroinflammation. 2021;18(1):74.

24.   Hasel P, Cooper ML, Marchildon AE, Rufen-Blanchette U, Kim RD, Ma TC, et al. Defining the molecular identity and morphology of glia limitans superficialis astrocytes in vertebrates. Cell Rep. 2025;44(3):115344.

25.   Hammond TR, Dufort C, Dissing-Olesen L, Giera S, Young A, Wysoker A, et al. Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. Immunity. 2019;50(1):253-71 e6.

26.   Heppner FL, Ransohoff RM, Becher B. Immune attack: the role of inflammation in Alzheimer disease. Nat Rev Neurosci. 2015;16(6):358-72.

27.   Trainor AR, MacDonald DS, Penney J. Microglia: roles and genetic risk in Parkinson's disease. Front Neurosci. 2024;18:1506358.

28.   Clarke BE, Patani R. The microglial component of amyotrophic lateral sclerosis. Brain. 2020;143(12):3526-39.

29.   Yin Z, Rosenzweig N, Kleemann KL, Zhang X, Brandao W, Margeta MA, et al. APOE4 impairs the microglial response in Alzheimer's disease by inducing TGFbeta-mediated checkpoints. Nat Immunol. 2023;24(11):1839-53.

30.   Bond S, Saxena S, Sierra-Delgado JA. Microglia in ALS: Insights into Mechanisms and Therapeutic Potential. Cells. 2025;14(6).

31.   Migliarini S, Scaricamazza S, Valle C, Ferri A, Pasqualetti M, Ferraro E. Microglia Morphological Changes in the Motor Cortex of hSOD1(G93A) Transgenic ALS Mice. Brain Sci. 2021;11(6).

32.   Mitchell MW, Grandizio C, Turan N, Requesens DV. An induced pluripotent stem cell line (CIMRi001-A) from a Vici syndrome donor with a homozygous recessive c.1007A>G (p.Q336R) mutation in the EPG5 gene. Stem Cell Res. 2022;63:102833.

Figure 1. Summary of whole animal deficits in *Epg5* mutant mouse models. A,B. Weight comparison of mutant homozygous mutants with sibling control wild-type mice at 4 weeks (**A**) and 8 weeks of age (**B**). **C-E.** Compared with control mice (**C**), *W860X* homozygous mutant mice (**D**) display poorly placed hindlimbs when attempting to bear weight at 4 months of age. Hindlimb function in homozygous *W860X* mice progressively worsened by 7 months of age (**E**). **F-I.** Assessment of the motor function of *W860X* (**F, G**) and KO (**H, I**) mutant mice via the rotarod test. Compared with age-matched control siblings, homozygous mutant mice have a poor capacity to navigate the rotarod test of coordination. Mutant mice fail the test at a lower speed of rotation ('speed at fall'; **F, H**) and at a shorter time ('latency to fall'; **G, I**) on the apparatus. At 12 weeks of age, some homozygous mutant animals are unable to initiate the rotarod test, all *W860X* mice fail at 14 weeks, and all *KO* mice fail at 16 weeks. *wt*; homozygous wild-type, *KO*; homozygous mutant for the *KO* allele; *W860X*; homozygous mutant for the *W860X* allele. Analysis of data from wild-type and mutant mice at a given timepoint was performed via an unpaired t test.

 ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$. *, $p < 0.05$.

Figure 1

Figure 2. **Molecular deficits in *Epg5* transcript stability and markers of autophagy in tissues derived from *Epg5* mutant mice. A.** Assessment of *Epg5* transcript stability in control and *Epg5* mutant mice at 6 months of age via an RT–qPCR primer pair that amplifies the 3' end of the *Epg5* transcript. Unstable *Epg5* transcript in spinal cord tissue (A) from mice with mutations that recapitulate pathogenic variants. *Epg5* transcript levels in cerebellar tissues (B) were reduced but not significantly. **B,C**. Western blot analysis of the spinal cord tissue of *Epg5* mutant mice at 7 months of age was performed to detect the expression of LC3, a marker of autophagy. Note the increased accumulation of autophagosome-associated LC3-II compared with that of cytoplasmic LC3 (**C**). *wt*; homozygous wild-type, *KO*; homozygous mutant for the *KO* allele; *W860X*; homozygous mutant for the *W860X* allele. Analysis of gene expression levels between wild-type and mutant mice was performed via an unpaired t test. ***, $p < 0.001$.

Figure 2

A

***Epg5* transcript, 3' probe**

cerebellum

spinal cord

B

C

LC3 II
LC3 I

Figure 3. Transcriptomic analysis of CNS tissues revealed a range of molecular deficits associated with *Epg5* pathogenic variants. A,B. Volcano plot from whole-tissue transcriptomic analysis of the anterior spinal cord (**A**) and cerebellum (**B**) showing differentially expressed genes (DEGs) *in Epg5 W860X* homozygous mutant mice compared with control age-matched siblings. Select genes of interest are marked. **C.** Venn diagram of overlapping differentially expressed genes in *Epg5 W860X* mutant CNS tissues. For this analysis, genes with both 2-fold or greater up- or downregulation and adjusted P values less than 0.01 **D were identified.** Heatmap of genes differentially expressed from the spinal cord and cerebellum of *Epg5 W860X* from transcriptomic analysis of select genes associated with a previously identified disease-associated microglial signature in other mouse neurological damage models. Genes that were absent from the transcriptomic analysis of the cerebellum (*Clec7a* and *Ccl4*) were assessed via Taqman qPCR. Notably, a substantial number of these genes are upregulated in the CNS of *Epg5 W860X* mutant mice **E, F.** Gene Ontology (GO; E) and pathway analyses of the overlapping CNS DEG signature in *Epg5 W860X* mutant mice.

Figure 3

Figure 4. Comparison of gene expression defects between *Epg5* mutant alleles. The expression of select differentially expressed genes identified from the transcriptomic analysis of *Epg5* mutant mice was examined via qPCR via TaqMan assays. **A-D**. Comparison of gene expression deficits in the spinal cord (A, B) and cerebellum (C, D) of the three *Epg5* homozygous mutant mice and age-matched sibling wild-type control mice at 6 months of age. Compared with the wild-type controls, all the mutants presented significant deficits in gene expression of these selected markers at 6 months of age. *wt*; homozygous wild-type, *KO*; homozygous mutant for the *KO* allele; *W860X*; homozygous mutant for the *W860X* allele. Analysis of gene expression levels between wild-type and mutant mice via one-way ANOVA. ****, $p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *, $p<0.05$.

Figure 4

Figure 5. Comparison of gene expression defects between *Epg5* mutant alleles and effects on age. Age effects on gene expression deficits in *Epg5 W860X* (**A, B**) and *Epg5 KO* (**C, D**) CNS tissues. Tissues from the anterior spinal cord (**A, C**) and cerebellum (**B, D**) were isolated from the indicated mutants between 2 months and 6 months of age for gene expression analysis. Relative expression is based on comparisons with wild-type mice at the 2-month timepoint. *wt*; homozygous wild-type, *KO*; homozygous mutant for the *KO* allele; *W860X*; homozygous mutant for the *W860X* allele. Analysis of gene expression levels between wild-type and mutant mice in a given cohort via one-way ANOVA. ****, $p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *, $p<0.05$.

Figure 5

Figure 6. Glial expansion in the spinal cord of *Epg5* mutant mice. Anterior spinal cord tissues, corresponding to the region around cervical vertebra 6 C6, were collected from 7-month-old control (**A, C, F, I, L**) or *Epg5 W860X* homozygous mutant mice (**B, D, G, J, M**) for histological analysis. **A-B.** Histochemical analysis to detect GFAP in the ventral spinal cord. A substantial increase in GFAP-positive cells was observed throughout the spinal cord in the *Epg5 W860X* homozygous mutant mice. Scale bar, 100 µm. **C–N**. Immunofluorescence was used to detect astrocytes (GFAP; green), microglia (IBA-1; red), and nuclei (DAPI; blue) in the spinal cord. **C,D.** Expanded microglia and astrocytes were observed throughout the spinal cord in the *Epg5 W860X* homozygous mutant mice. Scale bar, 300 µm. **E–N**. High-magnification assessment of microglia and astrocytes in specific regions of the spinal cord (**E**; diagram generated with Biorender). **F,G.** High-magnification image of the ventral gray matter of the spinal cord showing expanded astrocytes and microglia, with microglia demonstrating morphological changes, including thickened and more extensive processes and a large size, suggesting an activated phenotype (arrowheads). Scale bar, 100 µm. **H.** Quantification of the increase in microglia in the ventral horn gray matter of the spinal cord as a percentage of the total image area, demonstrating a quantitative increase in microglial content. **I,J.** Region of the dorsal funiculus of the spinal cord. Scale bar, 150 µm. **K.** Substantial expansion of microglia in the dorsalmost region of the dorsal funiculus. **L,M.** Details of the ventral border of the spinal cord. The top region of each image shows microglia, which are substantially increased in this region. The bottom portion of each image shows astrocytes, including the *glia limitans superficialis*, which are substantially expanded in the border region of the spinal cord. Scale bar, 100 µm. **N.** Quantification of astrocyte content in the ventral spinal cord and ventral border. Analysis of cell quantification between wild-type and mutant mice was performed via an unpaired t test. ****, $p < 0.0001$.

Figure 6

Figure 7. Glial expansion in the brain and cerebellum of *Epg5* mutant mice. Whole brains

were collected from 7-month-old *Epg5W860X* homozygous mutant mice for histological

analysis. **A-D.** Histochemical analysis to detect GFAP, a marker of astrocytes in the brain (**A, B;**

scale bar, 1 mm), and focus on the cerebellum (**C, D**) in control (**A, C**) or *Epg5 W860X*

homozygous mutant mice (**B, D**). A substantial increase in GFAP-positive cells was observed

throughout all regions of the brain, including the cerebellum. **E-G**. Immunofluorescence was

used to detect astrocytes (GFAP; green), microglia (IBA-1; red), and nuclei (DAPI; blue) in the

indicated regions of the cerebellum (E; diagram generated with Biorender) in control (**F**) or *Epg5*

W860X homozygous mutant mice (**G**). Expanded microglia and astrocytes are observed in all

layers of the cerebellum of *Epg5W860X* homozygous mutant mice. Scale bar, 200 µm. Layers

of the cerebellum are indicated. ML, molecular layer; PCL, Purkinje cell layer; IGL, internal

granule layer; WM, white matter.

Figure 7

Supplemental Figure 1. Longitudinal weight data of *Epg5* mutant mice. Mice were weighed weekly to monitor weight and for monitoring animal welfare. **A, B.** Weight data for control wildtype littermates or homozygous mutant *W860X* males (**A**) or females (**B**). **C,D.** Weight data for control wildtype littermates or homozygous mutant *KO* males (**C**) or females (**D**).

Supplemental Figure 1

Supplemental Figure 2. Uncropped western image. Full blot of lanes presented in Figure 2B. Middle portion of the blot was removed to eliminate samples not presented in this study.

Supplemental Figure 2

Supplemental Figure 3. Additional gene expression analysis of spinal cord tissues from *W860X* and *KO* homozygous mutant mice compared to wildtype control sibling mice.

Supplemental Figure 3

Supplemental Figure 4. Additional gene expression analysis of cerebellum tissues from *W860X* and *KO* homozygous mutant mice compared to wildtype control sibling mice.

Supplemental Figure 4

Supplemental Figure 5. Additional gene expression analysis of spinal cord and cerebellum tissues at 2 months and 6 months of age from *W860X* and *KO* homozygous mutant mice compared to wildtype control sibling mice.

Supplemental Figure 5

Supplemental Movie 1. Ataxic locomotion associated with *Epg5 W860X* homozygous mutant mice. W860X homozygous mutant male ('Mutant) or wildtype control littermate ('Wildtype') at 7 months of age were monitored in their home cage for capacity to walk and navigate cage. Note ataxic movement in W860X homozygous mutant mouse.

Supplemental video 1 screen capture

Supplemental Table 1. Summary of *Epg5* mutant alleles

Mouse allele	MGI allele nomenclature	mutation	Epg5 exon	crRNA target sequence	Donor DNA sequence	Forward Primer	Reverse Primer	Final allele sequence
KO	em1Vvn	p.R41QfsTer8	2	CGGGCTCTCCTGCTCGCGAC	n/a	AACTCCTCATGCTGAAACCACT	CCTCTCCCCTTTCTTTTAGGTCC	AACTCCTCATGCTGAAACCACTAGCTCTTTTATATCTCCTGCCTTGCTTGTAGGAAAA GAAGAAGCATGAAGCCCTTCAGACTTGTGATGCAGGTCCCCTGCCAGAGACCTG TCAGCTTCTGAACTCAAAGGAGATGACCTGAAGTCGTCTGCTGACCCCCAGCTC CACAGTGATGTGTGTGGATGGAATGAAAGTGAGATGTTTGATATACCACTCACCTCC TTGACTATAGGGGATGAAGGTCCCCCGGTACAGGACACAGAGGACCTAAAAGAAA GGGGAGAGG
W860X mouse acces NM_001195633.2	em2Vvn	W860X	14	CCCGAATCACAGCTATCTCT	ATAATGTTCTTGTCTTAAATTTAATTATTATTGTTTTCTTTTTAAAAAGGTTTCTT TATACTTGTTTAAAGAACTGCCTTTGTATCTTTGAAGAATTCCGACCTTCT GCCCTAGAGATAGCTGTGATTCGGGACTGGTTATTGAATAATAATCTGA CAGCAGTGAAGAACAAGTTGGCCT	CATCTTTTCCAAGGAAATTCTGCAA	AGGCTCCTACTTCTCGGAAATG	CATCTTTTCCAAGGAAATTCTGCAAGAGTTTAATAATGTTCTTGTCTTAAATTTAATTATTA TTGTTTTCTTTTTAAAAAGGTTTCTTTATACTTGTTTAAAGAACTGCCTTTGTATCTTTGAA GAATTCCGACCTTCTGCCCTAGCTGTGATTCGGGACTGGTTATTGAATAATAATCTG ACAGCAGTGAAGAACAAGTTGGCCTGTGTGATCCTAGAAGGACTGAATTGGGGATT TACTGAACAGGTATGCCATTTCCGAGAAGTAGGAGCCT

Supplemental Table 2. ThermoFisher TaqMan Gene Expression Assays used in this study.

Gene	Assay ID
C4b	Mm00437890_m1
Cst7	Mm00438351_m1
Gapdh	Mm99999915_g1
Gfap	Mm01253033_m1
Gpnmb	Mm01328587_m1
Irf7	Mm00516788_m1
Itgax	Mm00498701_m1
Lilrb4a	Mm01614371_m1
Epg5	Mm01257763_m1
Mpeg1	Mm01222137_g1
Myoc	Mm00447900_m1
Nkx2-9	Mm00435145_m1
Oas2	Mm00460961_m1
Tlr2	Mm00442346_m1
Tlr7	Mm00446590_m1

Supplemental Table 3. Overlapping differentially expressed genes between spinal cord and cereb

		Spinal cord		Cerebellum		
		Log2 (FC)	padj	Log2 (FC)	padj	
9930111J21F	9930111J21F	1.09323696	6.98E-16	1.33038402	0.00322291	
Abcc3	Abcc3	2.12550858	1.73E-07	1.56327374	0.00217837	
Adgre1	Adgre1	3.08041469	5.13E-165	2.33614275	2.62E-26	
Apobec1	Apobec1	2.43876645	2.34E-63	1.4175323	1.27E-06	
Arhgap25	Arhgap25	1.75621845	7.46E-22	1.32557032	0.00448986	
Arhgap30	Arhgap30	1.56735543	9.94E-58	1.41747037	0.00015443	
Arhgap45	Arhgap45	1.31513497	5.85E-43	1.04870243	1.24E-07	
Aspg	Aspg	2.72973722	3.93E-158	1.70045847	1.19E-20	
Atf3	Atf3	2.70013561	1.17E-30	1.02539951	0.00363607	
B2m	B2m	1.99016329	1.24E-134	2.13836775	1.95E-152	
B430306N03	B430306N03	3.26968387	1.11E-26	4.50528311	6.23E-08	
Blnk	Blnk	1.56527998	1.92E-28	1.66418325	5.93E-05	
Bst2	Bst2	2.32672814	4.18E-50	3.03118447	4.73E-55	
C1qa	C1qa	2.83889106	1.02E-181	2.42318726	5.95E-140	
C1qb	C1qb	2.4770435	3.95E-130	1.98774015	7.96E-73	
C1qc	C1qc	2.48082661	6.62E-158	2.03478932	2.60E-86	
C1ra	C1ra	3.11896874	3.44E-62	1.24464176	0.0079316	
C3	C3	2.97293034	1.85E-226	3.6966202	1.87E-67	
C3ar1	C3ar1	2.77905432	1.79E-99	2.49205263	4.79E-20	
C4b	C4b	3.24945218	0	4.097653	0	
Capn11	Capn11	-1.6181141	1.95E-05	-1.8847395	3.56E-06	*
Casp12	Casp12	2.06220661	3.80E-90	1.10847482	1.82E-07	
Casp4	Casp4	1.12751752	5.53E-07	2.39660891	0.00043959	
Ccl12	Ccl12	3.36680058	1.67E-09	3.76551008	0.00014058	
Ccl3	Ccl3	3.96950057	1.91E-29	3.51187528	2.67E-05	
Ccl5	Ccl5	1.99112676	0.00042323	3.45421804	6.25E-05	
Cd109	Cd109	2.233577	2.89E-81	1.56588797	1.52E-07	
Cd180	Cd180	1.745439	3.00E-19	2.11347131	1.90E-13	
Cd22	Cd22	2.453873	1.62E-05	3.74602405	2.95E-13	
Cd300c2	Cd300c2	2.10913094	4.18E-50	1.98900905	2.17E-08	
Cd300lf	Cd300lf	1.42525364	8.90E-19	2.20563695	0.00023274	
Cd33	Cd33	1.40523312	5.06E-15	1.50917871	5.72E-10	
Cd37	Cd37	1.39699777	7.46E-16	1.3810384	9.34E-08	
Cd4	Cd4	2.36723969	4.51E-05	2.36629778	0.00365418	
Cd44	Cd44	1.62018019	8.61E-54	1.11814697	1.45E-22	
Cd48	Cd48	3.81773186	7.06E-87	2.48453573	3.16E-07	
Cd52	Cd52	3.18278585	9.67E-80	2.23261055	2.65E-07	
Cd53	Cd53	1.64481494	1.52E-51	1.4575883	5.46E-08	
Cd68	Cd68	3.3192545	8.49E-293	2.30429753	9.08E-61	
Cd84	Cd84	2.07652747	3.81E-143	2.55015394	1.13E-22	

Cd86	Cd86	1.29260542	1.09E-10	1.69325869	0.00075391
Cebpd	Cebpd	3.04244063	1.24E-110	1.3648677	1.33E-07
Cp	Cp	2.3877424	4.68E-209	1.01235372	1.34E-21
Csf2rb	Csf2rb	2.23505878	1.63E-34	2.00183887	9.18E-06
Csf3r	Csf3r	1.77824849	1.91E-47	1.78261377	8.94E-18
Cst7	Cst7	7.08358303	1.27E-73	5.59410643	2.30E-13
Ctsh	Ctsh	1.66905581	2.22E-18	1.04331553	1.96E-12
Ctss	Ctss	2.04824464	1.34E-99	2.03825339	1.25E-99
Cxcl10	Cxcl10	3.20387296	6.57E-07	4.49244054	3.57E-26
Cxcl16	Cxcl16	2.4554051	7.76E-27	1.5013931	1.33E-07
Cxcl5	Cxcl5	1.95920013	2.92E-15	2.81624929	3.48E-06
Cxcr4	Cxcr4	1.7517299	1.19E-42	1.71181145	1.42E-06
Cyba	Cyba	2.55570313	3.96E-93	1.21737819	1.63E-06
Cybb	Cybb	3.31367843	1.11E-185	2.58867483	9.72E-15
Cybrd1	Cybrd1	1.07973224	2.89E-16	1.63721629	8.61E-13
Cyp1b1	Cyp1b1	1.3389824	3.25E-27	1.05310163	9.07E-05
Cyth4	Cyth4	1.17195832	1.11E-23	1.32300354	1.78E-10
Ddx60	Ddx60	2.130275	1.54E-28	3.99673794	9.12E-49
Dock2	Dock2	1.89784946	3.35E-11	1.57202733	2.57E-07
Dram1	Dram1	1.06683997	0.00036719	1.35714395	0.0081225
Elf4	Elf4	1.93386756	6.40E-19	1.41327999	6.53E-05
Emp3	Emp3	1.79141591	0.00077939	1.22367421	0.00075711
Endou	Endou	1.87109118	1.98E-31	1.92810235	6.80E-08
Epsti1	Epsti1	1.95196933	1.74E-12	2.81745101	4.02E-06
Fcer1g	Fcer1g	2.57228129	9.12E-20	2.14359677	4.88E-23
Fcgr2b	Fcgr2b	3.38787	2.57E-210	2.74587265	1.89E-34
Fcgr3	Fcgr3	1.656828	3.43E-53	1.45545798	1.96E-13
Fcrls	Fcrls	2.9554464	8.90E-256	2.21805027	1.39E-42
Fermt3	Fermt3	1.41069548	2.83E-24	1.28014374	4.38E-06
Fgr	Fgr	3.2053797	3.17E-53	2.65315916	0.00585116
Flnc	Flnc	1.56141239	1.81E-99	1.24577877	2.69E-19
Ftl1	Ftl1	1.8844304	9.92E-185	1.01330848	5.80E-75
Fyb	Fyb	2.17330256	1.02E-63	1.82447571	4.27E-12
Gbp10	Gbp10	3.96787969	3.50E-05	4.44224147	3.47E-30
Gbp2	Gbp2	2.03551463	2.58E-77	1.98818105	8.70E-40
Gbp3	Gbp3	1.59621172	1.35E-41	2.68173003	2.94E-74
Gbp5	Gbp5	1.60396385	1.75E-31	1.64147129	1.04E-15
Gbp6	Gbp6	2.53213722	6.28E-119	3.88841536	2.62E-89
Gbp7	Gbp7	1.13764387	1.23E-20	2.15340475	4.09E-39
Gbp9	Gbp9	1.2293972	2.92E-13	1.75609935	3.68E-30
Gfap	Gfap	1.859965	2.41E-187	2.42280036	0
Gm4951	Gm4951	1.62456055	3.88E-12	4.42584264	2.69E-20
Gna15	Gna15	1.11235084	3.95E-05	1.17244283	0.00265629

Gpnmb	Gpnmb	6.96937547	0	3.11687788	2.36E-107
Gpsm3	Gpsm3	1.31165269	4.37E-06	1.09670299	0.00227162
Gpx3	Gpx3	1.751674	1.79E-44	2.11788307	1.07E-05
Gsdmd	Gsdmd	1.08427717	3.99E-10	1.07954077	1.92E-05
H2-K1	H2-K1	1.53826538	4.01E-43	1.87352754	4.21E-12
H2-Q4	H2-Q4	1.17785503	3.74E-27	2.23186312	1.13E-08
H2-Q6	H2-Q6	1.35268617	6.68E-41	3.32350483	1.51E-05
H2-Q7	H2-Q7	1.43095142	1.40E-23	2.65803239	7.67E-13
H2-T23	H2-T23	1.17789121	5.25E-17	1.9377008	1.70E-51
Hacd4	Hacd4	2.48283111	3.56E-78	1.05527605	0.00017758
Havcr2	Havcr2	1.43021189	3.17E-23	1.55293502	0.00011274
Hck	Hck	1.63157853	9.90E-30	1.89681402	6.73E-06
Hcls1	Hcls1	1.72310648	8.01E-20	1.21891204	6.07E-10
Herc6	Herc6	1.17841776	6.75E-18	2.04998062	2.02E-21
Hmox1	Hmox1	2.0949206	4.74E-101	1.15696662	9.42E-07
Hpgds	Hpgds	1.10793338	1.30E-32	1.35408527	2.63E-10
Hpse	Hpse	4.04279887	2.57E-119	3.89997715	2.63E-17
Hvcn1	Hvcn1	2.87627013	9.57E-97	2.2310342	5.67E-16
Icam1	Icam1	2.33021974	4.98E-100	1.53898644	1.62E-11
Ifi204	Ifi204	1.74784008	1.44E-16	4.26484537	1.32E-25
Ifi206	Ifi206	3.76255694	4.23E-06	6.37423246	5.88E-16
Ifi207	Ifi207	4.75417028	1.07E-32	4.31043902	6.63E-25
Ifi209	Ifi209	2.15622861	1.10E-12	3.83016174	1.08E-07
Ifi211	Ifi211	3.25123308	3.75E-11	5.54861776	1.40E-05
Ifi213	Ifi213	1.20489433	1.03E-06	4.60156466	1.35E-24
Ifi27l2a	Ifi27l2a	3.09415307	8.32E-33	4.5272449	4.88E-12
Ifi35	Ifi35	1.01853263	1.28E-11	1.78594362	2.75E-10
Ifi44	Ifi44	2.1088134	2.13E-20	3.43396532	1.41E-33
Ifi47	Ifi47	1.71500301	7.11E-11	2.71436936	1.40E-10
Ifih1	Ifih1	1.15167678	7.44E-15	1.98325284	4.53E-48
Ifit1	Ifit1	1.80210266	9.03E-36	4.33450463	7.10E-221
Ifit3	Ifit3	1.58082475	3.87E-12	3.43150581	3.34E-60
Ifit3b	Ifit3b	1.52220083	4.06E-38	3.29314397	7.43E-147
Ifitm3	Ifitm3	1.82946232	4.97E-70	2.43416081	6.56E-56
Igsf6	Igsf6	2.52037471	1.25E-32	2.07354986	0.00037128
Igtp	Igtp	1.00030597	2.69E-15	2.89889032	1.23E-62
Iigp1	Iigp1	1.24035482	6.42E-13	2.34398713	3.97E-24
Ikzf1	Ikzf1	1.3934617	8.33E-18	1.29298109	5.95E-06
Inpp5d	Inpp5d	1.27190344	6.54E-29	1.24737539	2.20E-10
Irf5	Irf5	1.21467096	1.59E-09	1.45034757	3.46E-06
Irf7	Irf7	1.79885208	3.39E-14	4.27563941	1.19E-189
Irf8	Irf8	1.75961838	1.86E-43	1.469239	5.83E-10
Irf9	Irf9	1.41117651	1.74E-42	2.19934043	1.40E-127

Irgm1	Irgm1	1.29359346	1.36E-42	2.84748724	7.70E-135
Irgm2	Irgm2	1.40746868	7.76E-31	2.35813783	4.33E-12
Isg15	Isg15	1.97567196	5.42E-11	3.53013719	2.10E-37
Itgam	Itgam	1.39487586	1.08E-50	1.4785927	0.00015012
Itgax	Itgax	4.86246051	1.09E-90	2.94781726	2.42E-07
Itgb2	Itgb2	2.65340641	2.09E-134	2.12792947	6.59E-25
Kcns1	Kcns1	1.82837515	3.09E-56	1.193473	5.00E-06
Laptm5	Laptm5	1.71192945	2.20E-115	1.55676397	3.90E-41
Lcn2	Lcn2	8.23824511	6.01E-175	6.98157203	9.14E-17
Lcp2	Lcp2	1.15918121	8.83E-08	1.57088278	1.98E-05
Lgals3	Lgals3	3.21109024	6.56E-05	2.6015204	9.07E-32
Lgals3bp	Lgals3bp	2.58888466	4.35E-115	3.20785225	7.22E-23
Lgals9	Lgals9	1.23594525	4.94E-12	2.31785955	2.22E-61
Lilrb4a	Lilrb4a	6.11679115	1.24E-207	3.64229431	2.99E-10
Ly86	Ly86	1.31532624	6.32E-28	1.28221394	1.69E-07
Ly9	Ly9	6.23861607	1.88E-116	4.55341404	2.71E-23
Lyz2	Lyz2	5.91046957	0	2.76446151	6.99E-121
Mafb	Mafb	1.22837804	1.23E-34	1.19402998	5.03E-11
Met	Met	1.51339207	1.19E-21	1.81739235	0.00015166
Mmp19	Mmp19	2.39631242	9.66E-56	1.14488893	0.00961152
Mmrn2	Mmrn2	1.07996509	1.31E-08	1.16303462	8.17E-12
Mndal	Mndal	1.18258563	6.07E-07	2.01059136	2.11E-14
Mpeg1	Mpeg1	3.24940321	1.39E-278	2.6585749	2.95E-163
Ms4a6b	Ms4a6b	2.29815961	7.68E-19	2.64518093	1.96E-06
Ms4a6c	Ms4a6c	4.30058984	6.35E-89	2.67681065	9.55E-06
Ms4a6d	Ms4a6d	3.04188719	9.87E-61	2.78686839	4.98E-08
Ms4a7	Ms4a7	3.53125441	9.59E-82	2.87197894	0.00019914
Myo1f	Myo1f	2.57091034	2.31E-80	1.86017073	3.91E-15
Naip2	Naip2	1.72924888	4.53E-44	2.0926213	1.45E-06
Naip5	Naip5	1.63477516	6.45E-25	1.44817403	0.00402905
Ncf1	Ncf1	1.80737296	1.29E-63	1.56973261	7.66E-11
Nckap1l	Nckap1l	1.50529251	3.41E-59	1.39158974	4.56E-09
Nfam1	Nfam1	1.580226	1.67E-31	1.24620575	0.00304052
Nkx2-9	Nkx2-9	-1.7853189	6.56E-24	-1.5363872	0.0012498 *
Nlrc5	Nlrc5	2.25657049	3.13E-28	3.02177144	1.84E-08
Oas1a	Oas1a	2.55551833	8.45E-23	3.89118159	2.50E-32
Oas1g	Oas1g	3.42955597	1.06E-18	4.80820977	1.60E-17
Oas2	Oas2	3.18867506	2.22E-26	4.02275279	2.65E-20
Oasl1	Oasl1	1.33293957	0.00199826	3.64436117	3.60E-23
Oasl2	Oasl2	1.59728013	8.59E-15	4.5741957	0
Osgin1	Osgin1	3.00950338	2.73E-18	1.597678	0.00047451
Osmr	Osmr	2.60423198	4.54E-106	1.29034216	8.51E-14
P2ry6	P2ry6	1.60154361	3.59E-15	1.28417266	0.00361603

Parp10	Parp10	1.15464847	1.49E-13	1.75256971	3.62E-14
Parp14	Parp14	1.62123279	6.98E-33	2.7252473	2.93E-94
Phf11b	Phf11b	1.43054	0.00062	3.52445404	2.60E-06
Phf11d	Phf11d	1.301265	4.97E-07	3.07515079	8.32E-12
Pik3ap1	Pik3ap1	1.75375416	4.19E-62	1.61840688	2.94E-10
Pik3cg	Pik3cg	1.41647866	5.57E-35	1.5509373	1.51E-07
Pik3r5	Pik3r5	1.34886588	7.37E-24	1.47278377	0.00056886
Pirb	Pirb	3.47346512	1.90E-77	3.17641247	7.93E-11
Pla2g5	Pla2g5	1.0454165	2.22E-09	1.5300915	1.75E-08
Plau	Plau	2.44026133	1.07E-53	1.38022124	0.00867992
Plaur	Plaur	2.46744804	1.08E-17	1.31527775	0.00258336
Plcg2	Plcg2	1.56947193	2.78E-72	1.08154153	2.09E-06
Pld4	Pld4	1.2023639	4.28E-15	1.21704964	1.15E-07
Plscr2	Plscr2	1.34768233	4.24E-22	1.09168062	1.70E-05
Psmb8	Psmb8	2.21546109	2.41E-09	2.12590838	1.46E-21
Psmb9	Psmb9	1.37956227	7.48E-09	2.09133326	3.24E-10
Ptafr	Ptafr	1.86952626	8.90E-28	2.52990537	3.34E-07
Ptgs2	Ptgs2	1.77551623	4.95E-14	2.09768419	0.00051199
Ptpn6	Ptpn6	1.90240483	4.78E-16	1.89198975	1.44E-18
Ptprc	Ptprc	1.66898916	5.19E-67	1.98218167	3.05E-12
Pycard	Pycard	1.67769381	9.20E-25	1.24016306	0.00014135
Qprt	Qprt	1.07912428	0.0003234	1.06267953	0.0026013
Rac2	Rac2	1.25024663	7.28E-12	1.39076245	0.00047451
Rasal3	Rasal3	1.38812073	4.09E-11	1.15441901	0.00133359
Rnf213	Rnf213	1.02936303	4.53E-44	2.92297347	1.60E-269
Rsad2	Rsad2	2.39190388	8.86E-67	4.05477845	1.16E-111
Rtp4	Rtp4	1.33458926	6.19E-22	4.02740414	1.78E-81
S1pr3	S1pr3	2.15344914	4.62E-95	1.06766959	2.29E-07
Samsn1	Samsn1	1.888985	6.79E-22	1.54356173	0.00908748
Sash3	Sash3	1.6070489	2.65E-16	1.2284922	0.00995937
Sctr	Sctr	2.03428616	2.22E-18	2.18036048	0.00979115
Serpina3n	Serpina3n	4.31870614	0	2.69367338	4.53E-62
Siglec1	Siglec1	4.15375252	1.88E-116	3.07308979	4.05E-20
Slamf9	Slamf9	2.381156	6.04E-33	1.56869667	0.00839036
Slc11a1	Slc11a1	3.338065	6.41E-25	2.25945412	1.75E-20
Slc15a3	Slc15a3	2.202387	2.80E-39	1.03871766	0.00248646
Slc43a3	Slc43a3	3.391145	1.20E-78	2.20339326	4.14E-11
Slfn2	Slfn2	2.44927594	2.69E-37	2.30845658	4.57E-07
Slfn5	Slfn5	1.22815019	4.24E-23	1.34205719	5.41E-39
Slfn8	Slfn8	1.81435195	1.09E-19	3.45993299	1.17E-31
Sp100	Sp100	1.21268615	4.56E-17	1.69431912	1.39E-36
Sp110	Sp110	1.26909943	4.11E-13	1.89702199	3.59E-10
Sp140	Sp140	2.31027467	1.84E-10	1.85005001	0.00063355

Spi1	Spi1	1.64679025	1.61E-07	1.30648573	3.36E-06
Stab1	Stab1	1.412411	1.78E-13	1.13929771	6.53E-19
Syk	Syk	1.70671096	3.47E-42	1.61450849	7.14E-10
Tap1	Tap1	1.04311893	2.30E-09	1.90580866	7.45E-23
Tent5c	Tent5c	1.90704566	3.33E-61	1.09685559	0.00014035
Tgtp1	Tgtp1	1.14313856	0.00238102	2.27638666	9.39E-07
Themis2	Themis2	1.947071	2.96E-25	1.88186106	1.23E-06
Timp1	Timp1	6.72298999	9.38E-96	2.13374215	0.00279707
Tlr13	Tlr13	2.32753293	2.72E-91	1.91674908	1.42E-06
Tlr2	Tlr2	2.58309712	4.29E-120	2.25703879	2.65E-14
Tlr7	Tlr7	2.23676878	2.73E-31	2.12004101	2.58E-11
Tmem106a	Tmem106a	1.84723862	2.65E-15	1.83369932	0.00023074
Tmem140	Tmem140	1.09860098	4.14E-08	1.40385403	9.85E-06
Tmem176a	Tmem176a	2.23273212	5.05E-27	1.29958885	1.91E-11
Tnfrsf1b	Tnfrsf1b	1.36348076	0.00263848	1.70122722	1.29E-11
Trem2	Trem2	2.58713712	8.40E-195	2.18780217	2.09E-44
Treml2	Treml2	2.42613098	2.81E-22	3.41506429	6.92E-07
Trim12c	Trim12c	1.058529	3.19E-17	1.71640508	7.22E-15
Trim30a	Trim30a	1.81813707	9.97E-35	3.16077017	2.39E-72
Tubb6	Tubb6	1.25448578	8.35E-16	1.140325	0.00350475
Tyrobp	Tyrobp	2.75002265	8.32E-96	1.84897965	8.43E-34
Uba7	Uba7	1.08654111	9.84E-19	1.98942687	6.54E-31
Unc93b1	Unc93b1	1.28069625	1.58E-28	1.29184662	1.94E-21
Usp18	Usp18	2.26246116	9.81E-20	4.36206577	1.25E-99
Vav1	Vav1	1.78895641	8.18E-26	1.2688305	6.99E-05
Wdfy4	Wdfy4	1.22520348	4.49E-06	1.53619436	0.00016596
Xdh	Xdh	1.71104141	4.05E-39	1.21164822	3.28E-25
Zbp1	Zbp1	4.02148415	1.45E-19	4.63812852	5.92E-34
Zc3hav1	Zc3hav1	1.00055704	1.01E-36	2.02025225	3.05E-75