Microbiome-mediated resilience and cross-generational consequences in male

***Drosophila* exposed to combined environmental stressors**

Komal Maggu*, Abhishek Meena, Alessio N. De Nardo, Benjamin Eggs, Sonja H. Sbilordo,

Rawaa Al Toma Sho & Stefan Lüpold*

Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

Correspondence: komal024maggu@gmail.com, stefan.luepold@uzh.ch

Abstract

Environmental stressors like heat extremes and pesticide exposure can significantly threaten

insect reproduction, yet the interplay of these stressors and the potential mitigating role of the

gut microbiome remain poorly understood, particularly across generations. This study

investigated the interactive effects of acute heat stress and sublethal imidacloprid exposure on

male *Drosophila melanogaster* reproductive success and the subsequent fitness of their

offspring. We manipulated the gut microbiome of male flies (germ-free or colonized with one

or five bacterial species) and subjected them to individual or combined stress. Our findings

reveal that combined stress synergistically impairs male fitness traits, an effect partially

buffered by higher microbiome diversity. We further demonstrate cross-generational

consequences, with paternal stress exposure also impacting offspring fitness. Our results

highlight the crucial role of the gut microbiome in mediating resilience to environmental stress

and underscore the importance of considering multi-stressor and intergenerational effects in

ecological risk assessments.

Keywords- Microbiota, Thermal fertility, Pesticide exposure, Inter-generational, fruit fly

Introduction

Coping with the increasing prevalence of anthropogenic stressors is a central challenge for

organisms under the ongoing global change (Navarro-Ortega et al., 2015; Piggott, Niyogi, et

al., 2015; Piggott, Townsend, et al., 2015; Przeslawski et al., 2015; Segner et al., 2014). Among

these stressors, climate change and agricultural pesticides are key drivers of ecosystem-wide

biodiversity loss (Beketov et al., 2013; Heino et al., 2009; Stehle & Schulz, 2015), with their

interaction posing unique risks. Rising temperatures are projected to intensify pesticide use in

agriculture as warmer climates favor pest outbreaks (Deutsch et al., 2018), while

simultaneously altering how organisms, particularly ectotherms, respond to chemical exposure

(Fournier-Level et al., 2016; Hooper et al., 2013a; Noyes et al., 2009). For ectotherms, which

cannot regulate internal temperatures (Deutsch et al., 2008; Paaijmans et al., 2013), warming

elevates metabolic rates and feeding activity (and so potentially pesticide uptake), forcing

energetic trade-offs between detoxification and essential functions like growth or reproduction

(Jutfelt et al., 2021; Pörtner et al., 2017).

While the individual effects of temperature and pesticides are well-documented (Dougherty et

al., 2024; Wan et al., 2025), their combined impacts remain poorly resolved. Studies report

additive (Arambourou & Stoks, 2015), antagonistic (Meng et al., 2022), or synergistic (Wang

et al., 2019) interactions, with synergism being most concerning for non-target species as

combined harm exceeds the sum of individual detrimental effects (Folt et al., 1999; Siviter et

al., 2021; Verheyen & Stoks, 2019, 2023). Such synergisms may arise through physiological

trade-offs, for example if a diversion of resources away from detoxification by heat stress (Dai

et al., 2023) is combined with a depletion of antioxidants needed for thermal resilience through

pesticide exposure (A. P. N. Silva et al., 2025). Such dual depletion might disproportionately

damage gonads and gametes that are particularly prone to oxidative stress due to their high cell

proliferation, metabolic rate and generation of reactive oxygen species (Simmons et al. 2018).

By eroding reproductive resilience, synergisms threaten to accelerate biodiversity loss in ways

single stressors would not (Shahid et al., 2024).

Beyond direct physiological impacts, host-associated microbial communities (microbiomes)

are increasingly recognized as mediators for environmental stress tolerance (Feldhaar, 2011;

Osborne et al., 2023; Sepulveda & Moeller, 2020b; Shokal et al., 2016). Commensal microbes

can enhance host thermal performance (Renoz et al., 2019; Sepulveda & Moeller, 2020a),

either as obligate or facultative symbionts (Dunbar et al., 2007; Gruntenko et al., 2017;

Montllor et al., 2002; J. A. Russell & Moran, 2006; Zhang et al., 2008). Similarly, a healthy

microbiome can bolster insect resistance to insecticides through direct and indirect pathways.

For example, symbionts like *Citrobacter* in the oriental fruit fly *Bactrocera dorsalis*, or

Burkholderia in the bean bug *Riptortus pedestris*, directly metabolize pesticides using encoded

degradation enzymes, whereas others, such as *Arsenophonus* in *Nilaparvata lugens* or

Lactobacillus in *Drosophila melanogaster*, can induce the host's own detoxification gene

expression (Cheng et al., 2017; Jones et al., 2015; Kikuchi et al., 2012; Pang et al., 2018). The

gut microbiome thus contributes to insect survival and fitness by providing nutrients or

mitigating biotic and abiotic stresses (Hector et al., 2022; Perry et al., 2008). However, this

protective capacity of the gut microbiota is context-dependent; stressor-induced dysbiosis can

impair host fitness, creating feedback loops that may exacerbate vulnerability (Akbar et al.,

2022; Evariste et al., 2019; Khosravi & Mazmanian, 2013; Kumar & Kumar, 2019; R. J.

Russell et al., 2011; Sepulveda & Moeller, 2020a; Swelam et al., 2022). Despite progress in

studying individual stressors, how microbiomes modulate *combined* effects remains unknown.

This gap is critical, as microbiome disruption by one stressor could amplify susceptibility to

another, potentially explaining observed synergisms in some species (Karimzadeh et al., 2020;

Mao et al., 2019; Zhang et al., 2008; Zhang, Cai, et al., 2021).

A parallel knowledge gap concerns stress effects on male reproduction, as much research on

environmental stress in invertebrates focuses on acute toxicity or sterility thresholds (de Franca

et al., 2017; Desneux et al., 2007; B. S. Walsh, Parratt, Hoffmann, et al., 2019; Zwoinska et al.,

2020), overlooking the significant fitness consequences of *sublethal* exposures. Sublethal

effects on female reproductive function are relatively well characterized, but the corresponding

consequences for males have only recently come into focus and revealed that male fertility can

be far more vulnerable than female fecundity, at least to heat stress (Iossa, 2019; Sales et al.,

2018b; Zwoinska et al., 2020). Heat and pesticides have been shown to impair spermatogenesis,

sperm DNA integrity, and ejaculate quality more broadly (Aulsebrook, Bertram, Martin,

Aulsebrook, Brodin, Evans, Hall, O'Bryan, Pask, & Tyler, 2020; Dougherty et al., 2024; B. S.

Walsh, Parratt, Atkinson, et al., 2019). Reductions in ejaculate quality are particularly

impactful in promiscuous species, in which the competitive fertilization success depends on

the relative ejaculate quantity and quality among rival males (Fitzpatrick & Lüpold, 2014;

Parker & Pizzari, 2010; Pizzari & Parker, 2009; Simmons & Fitzpatrick, 2012).

Damage to sperm also has the potential of extending the effects of environmental stress across

generations through epigenetic modifications or sperm-borne small RNAs (Bonduriansky et

al., 2012; Morgan & Watkins, 2019), with profound consequences for population stability and

evolutionary trajectories (Aulsebrook, Bertram, Martin, Aulsebrook, Brodin, Evans, Hall,

O'Bryan, Pask, Tyler, et al., 2020; Crean & Immler, 2021). However, although parental

exposure to combined thermal and chemical stress has been shown to reduce offspring

performance in some species (DeCourten & Brander, 2017; Fathy, 2024), other studies find no

clear effects (Massot et al., 2021; Meng et al., 2022) or even reduced offspring vulnerability

(Tran et al., 2018), indicating interspecific variation potentially due to differences in stressor

intensity (including the thermal sensitivity of pesticides themselves), species-specific

acclimation, or compensatory maternal provisioning. Understanding how environmental

stressors impact not only directly exposed individuals but also subsequent generations is crucial

for predicting long-term ecological consequences and evolutionary adaptation potential.

To address these knowledge gaps, we investigated the interactive effects of acute heat stress

and sublethal pesticide exposure on the reproductive fitness of male *D. melanogaster*, with a

focus on microbiome mediation and cross-generational consequences. We combined sublethal

acute heat stress (4 hours at 36°C), previously shown to negatively affect testicular function

and the fertilizing capacity of sperm (Meena, De Nardo, et al., 2024; Meena, Maggu, et al.,

2024), with exposure to imidacloprid, a widely used systemic insecticide targeting nicotinic

acetylcholine receptors (nAChRs) in the central nervous system (Ihara et al., 2017). Sublethal

imidacloprid impacts have been documented in diverse insects, affecting enzyme activities,

biological performance, and gene expression (He et al., 2011; Kang et al., 2018; W. Li et al.,

2018), and ultimately impairing fitness (De França et al., 2017; He et al., 2011; Xiao et al.,

2016; Zhang, Xu, et al., 2021; Zhou et al., 2017). To isolate the role of the microbiome, we

compared males that were germ-free (axenic) with males associated with defined bacterial

communities: either mono-association of two common gut bacteria (*Acetobacter tropicalis* or

Lactobacillus brevis) or a more complex, yet defined, five-species consortium combining

Acetobacter and *Lactobacillus* species that are commonly found in *Drosophila* (Broderick &

Lemaitre, 2012; Gould et al., 2018; Newell & Douglas, 2014; Ridley et al., 2013). This

gnotobiotic approach (i.e., using defined microbiotas) enabled precise assessment of microbial

contributions to stress tolerance. Furthermore, we assessed the cross-generational

consequences of paternal stress exposure by examining offspring fitness, explicitly testing

whether the father's microbiome status influences these cross-generational effects. Our design

minimized bacterial transmission to offspring, focusing the analysis on direct paternal effects

related to parental condition (Marshall et al., 2010; Moore et al., 2019; Mousseau et al., 2009).

Based on the evidence outlined above, we tested the following five predictions: (1) Combined

sublethal exposure to heat stress and imidacloprid will synergistically reduce male reproductive

fitness more than the individual effects of each stressor. (2) The presence of gut microbes will

partially mitigate the negative impacts of individual and combined stressors on male

reproductive fitness compared to germ-free males. (3) Different bacterial compositions (mono-

associations vs. defined complex community) will vary in their protective capacity, with the

more complex community offering greater resilience than either mono-association. (4) Paternal

exposure to individual and combined stressors will negatively impact offspring fitness (cross-

generational effects), with synergistic interactions between stressors potentially amplifying this

fitness loss. (5) The composition of the paternal microbiome will influence the magnitude of

these cross-generational effects.

Materials and Methods

Experimental populations

Throughout this study, we used an outbred *D. melanogaster* population (LH$_m$ wild-type strain),

maintained under constant conditions of 25°C, 60% humidity, and a 12h:12h light:dark cycle.

First, we generated axenic (germ-free) focal flies by collecting eggs (within six hours of

oviposition) from four large population containers supplied with Petri dishes containing grape-

juice agar medium (300 mL water, 670 mL grape juice, 22g agar, 20g sugar, 19g yeast, 20 mL

nipagin per litre) supplemented with a smear of yeast paste. Eggs were washed off the plates

into a strainer, pooled, and dechorionated by immersion in 3.5% bleach for 2 minutes, followed

by a rinse in 70% ethanol and three rinses in sterile water.

Next, eggs were aseptically transferred to sterile glass vials containing autoclaved standard fly

food (75g glucose, 100g fresh yeast, 55g corn, 10g flour, and 15mL nipagin antimicrobial agent

per litre food media) supplemented with antibiotics [a cocktail of Ampicillin (50mg/mL),

Kanamycin (50mg/mL), Erythromycin (10mg/mL), Tetracycline (10 mg/mL)]. Vials were then

sealed with sterile plugs and incubated under standard laboratory conditions until adult

eclosion. All steps of vial preparation were conducted in a laminar flow chamber to avoid

contamination.

Upon eclosion, axenic status was confirmed using two methods on a subsample of flies. First,

homogenated flies were plated on Lysogeny Broth (LB) and de Man, Rogosa and Sharpe

(MRS) agar plates, incubated, and checked for bacterial colony growth. Second, genomic DNA

was extracted from pools of 3-5 putatively germ-free and conventional flies (positive control),

respectively, using Quick-DNA midiprep plus kit following the instructions provided by the

manufacturer. Universal primers (27F and 1492R) were used as forward and reverse primer,

respectively. To prepare the PCR master mix, 10μL of nuclease-free water, 0.5μL of forward

and reverse primer, 1μL of template DNA and 13μL Taq green master mix were mixed in a

sterile eppendorf tube. PCR conditions were: initial denaturation at 95°C for 90s, followed by

35 cycles of denaturation at 95°C for 30s, annealing at 55°C for 30s and extension at 72°C for

90s, and final extension at 72°C for 5 minutes). PCR products were visualized on a 1% agarose

gel under ultraviolet light to confirm the absence of bacterial DNA amplification in axenic

samples and its presence in controls.

Bacterial cultures and gnotobiotic associations

Axenic males were associated with defined compositions of microbial species, all originally

isolated from the gut of wild-type flies (Obadia et al., 2017): *Acetobacter tropicalis* (*At*),

and *Lactobacillus brevis* (*Lb*) for mono-associations based on their contrasting effects on male

reproductive investments in our previous studies (Maggu et al. in prep.). For comparison, we

also included axenic males and males inoculated with a defined five-species community

consisting of common and abundant *Drosophila* gut bacteria (*A. tropicalis*, *A. orientalis, A.*

pasteurianus, L. plantarum, L. brevis), resulting in four distinct microbial treatments.

Bacterial strains were cultured overnight in an incubator at 30°C: *Acetobacter* in MRS +

mannitol medium with shaking, and *Lactobacillus* in MRS broth under stationary conditions.

The following morning, cultures were checked for growth before measuring their optical

densities (OD) at 600 nm and centrifuging at 7000 rpm for 5 minutes. Bacterial pellets were

then resuspended in sterile phosphate-buffered saline (PBS) to a final concentration of 10^8

cells/mL, following established protocols (Koyle et al., 2016; Newell & Douglas, 2014).

To inoculate vials, 50µL of bacterial suspension (approx. 5×10^6 CFUs) was added to sterile

food vials, with equal representation of bacterial strains in the 5-species consortia. This

bacterial density is comparable to previous reports for conventionally reared flies (Gould et al.,

2018; Newell & Douglas, 2014). Inoculated food vials were then allowed to air-dry in a laminar

flow cell before introducing ten 1-day-old axenic males to each to generate gnotobiotic flies.

Males were transferred to freshly inoculated vials with the same bacterial treatment every 48h

for a total of four days to ensure stable colonization. Successful colonization with the intended

bacteria was confirmed by plating homogenates of males from each treatment and checking for

characteristic colony morphologies (see *Bacterial abundance of males*).

Stress treatments

Following the 4-day colonization period, males were transferred to vials containing sterile food

and exposed to pesticide and heat stress in a fully factorial design. Specifically, half of the

males from each microbial group were subjected to food supplemented with imidacloprid,

whilst the remaining half were maintained on uncontaminated standard food. We used a

concentration of 1 ppm (parts per million) of imidacloprid as concentrations of 0.04 to 20 ppm

have been used effectively in studies on larval or adult *D. melanogaster*, depending on the

exposure regime (Charpentier et al., 2014a; Frantzios et al., 2008; Martelli et al., 2020). These

concentrations are low enough to avoid immediate lethality but sufficient to induce sublethal

effects on behavior, reproduction, and physiology. Males were exposed to imidacloprid for an

acute window of 18 h, following an additional thermal treatment of 4-hour in a hot water bath.

To this end, half of the males from each microbiome and pesticide treatment were subjected to

acute heat stress by transferring them to vials containing only sterile 1% agar (to prevent

dehydration) and placing these vials in a water at 36°C for 4 hours. The remaining males

(thermal controls) were treated identically but at benign 24.5°C. All males were then allowed

to recover overnight in standard food vials under standard laboratory conditions.

Hatching success and fertility

To quantify the fitness consequences of heat and pesticide exposure and their potential

mitigation by the microbiome on the morning after the stress treatments, experimental males

were individually mated to standard females of the same population. All females had been

collected as virgins (within 7 hours of eclosion) and reared under axenic conditions along with

the experimental males to minimize confounding contributions of female microbes to the

measured outcomes. After mating ended, females were allowed to lay eggs for 24 h, and

another 30 h later (allowing time for embryonic development), oviposition vials were frozen

for later quantification of the proportion of eggs hatching as a proxy of fertilization success,

and the total number of larvae estimating male-mediated fecundity. This assay was performed

in two blocks, each with 25 males per treatment combination, totaling 800 experimental males

(4 bacterial treatments × 2 pesticide treatments × 2 temperature treatments × 2 blocks × 25

males).

Sperm competitiveness

To assess male sperm competitiveness, the above experimental design was repeated, but this

time using standard females that had mated once with standard competitor males two days

before the focal mating. These standard competitors were of the same strain but genetically

modified to ubiquitously express green fluorescent protein (GFP), permitting paternity

assignment against the focal wild-type males (Lüpold et al., 2012, 2020). Pairs were monitored

for remating for up to 8 h. Remated females were then isolated in fresh vials for oviposition,

while pairs not mating were separated and given another 8-hour mating opportunity on the

following day. Females still not remating were excluded. Offspring paternity (focal vs.

competitor) was assessed via GFP fluorescence to calculate P_2 as the proportion of offspring

sired by the second (focal) male. The treatment design of this assay was identical to the non-

competitive experiment above, but this time with 30 trials per treatment combination in each

of the two experimental blocks (i.e., 960 trials in total).

Bacterial abundance of males

To investigate the impact of stress treatments on the bacterial load, colony forming units

(CFUs) were compared between ten random males from each experimental treatment, surface-

sterilized with 70% ethanol, and individually homogenized in 1.5 mL microcentrifuge tubes

containing sterile PBS using a pestle. Homogenates were serially diluted to different dilutions

at a final volume of 100 μL, then plated on MRS media for quantification of CFUs at

appropriate dilutions (approx. 100 CFUs/plate). For the control groups with both *Acetobacter*

and *Lactobacillus* species, MRS agar was supplemented with 10 µg/mL ampicillin to avoid the

growth of *Lactobacillus,* whilst replicate plates were incubated in a CO_2 atmosphere to exclude

the growth of *Acetobacter*. The presence of only target bacterial species on plates was

confirmed based on colony morphologies, with *A. tropicalis* producing small tan-brown

colonies and *L. brevis* producing small yellowish colonies. CFUs per fly were calculated as:

$CFU = C \times D/P \times V$, where C = number of colonies counted, D = dilution, P = volume plated,

V = volume of fly homogenate (Koyle et al., 2016). For control males, with all the five bacterial

species, bacterial colonies were identified by their morphology and total CFUs per fly was

obtained by adding CFUs from all the five bacterial species.

Cross-generational effects

To assess cross-generational impacts of paternal (F_0) stress exposure, we quantified hatching

success and fecundity in both male and female offspring (F_1) of our experimental males. Within

each experimental block, up to 20 freshly hatched larvae from each of ten randomly selected

vials per parental treatment combination (i.e., 16 combinations) were transferred to rearing

vials containing sterilized standard fly food. This immediate transfer minimized the potential

acquisition of parental microbes via consumption of the egg chorion or contaminated food.

These larvae developed under standard laboratory conditions.

Upon eclosion, 4-5 virgin F_1 male and female offspring were collected from each rearing vial,

separated by sex. At 2-3 days of age, three males and three females from each lineage were

individually mated with a virgin standard fly of the opposite sex reared under axenic conditions

(to avoid potential confounding effects of their microbiota). Following copulation, hatching

success and fecundity were measured for these F_1 flies as in their parents (see above). These

offspring themselves were not subjected to any experimental treatment, thus assessing solely

the cross-generational effects. This experiment was carried out in two independent blocks of

480 offspring per sex from 160 F_0 males (i.e. total of 1,920 mating trials).

Statistical analysis

All statistical analyses were performed in R version 4.4.0 (R core team, 2024), using

generalized linear mixed-effects models (GLMMs) and generalized linear models (GLMs) to

assess the male responses to heat stress and/or pesticide exposure under different microbiome

conditions. Each model included microbiome composition (4-level factor) along with the

binary thermal and pesticide treatments with all two- and three-way interactions as fixed

effects, and experimental block as a fixed or random effect. Pairwise comparisons were

performed using the emmeans package (Searle et al., 1980) (Estimated Marginal Means) to

assess the significance of differences between levels of each factor.

Hatching success (i.e., proportion of all eggs that hatched) was analyzed in a quasibinomial

GLMM (as the binomial distribution resulted in under-dispersion), with block as a fixed effect.

Male-mediated female fecundity (i.e., total number of eggs laid) was modeled using a truncated

Poisson model with a zero-inflation component as implemented in the package glmmTMB

(Brooks et al., 2017), also including block as a fixed effect. Finally, for sperm competitiveness

(P_2), we fitted a GLMM (glmmTMB) with a Tweedie distribution and a log link function to

account for skewness and under-dispersion in the data. Experimental block and male ID were

included as random effects.

The impact of the different experimental treatments was also analyzed for the total bacterial

abundances (CFUs) and, separately, for compositional shifts in the five-species consortia. For

the former, we used a linear mixed-effects model with block as a random effect. For

microbiome community composition in the five-species consortium, we quantified the species-

specific abundances per sample, then applied a Hellinger transformation to these count data

and computed Bray-Curtis distances using the *vegdist* function in the package vegan (Oksanen

et al., 2023). We assessed treatment effects (temperature, pesticide, and their interaction) in a

permutational multivariate analysis of variance (PERMANOVA) using *adonis2* in the same

package. To visualize compositional shifts, we further ran non-metric multidimensional scaling

(NMDS) (*metaMDS*) on the Bray-Curtis dissimilarities and fitted species vectors onto the

NMDS ordination with *envfit*, which indicates the direction and strength of each species-

specific association with the ordination space. We also compared the relative abundances at

the genus level, using a binomial GLMM with block as a random effect and accounting for

overdispersion by including an additional observation-level random effect.

Finally, the effects of paternal (F_0) treatments on offspring (F_1) fitness were quantified

separately for male and female progeny using GLMMs. Each model included F_0 treatments

and their interactions as fixed effects (as above), with experimental block and sire ID as random

effects. GLMMs for hatching success included a zero-inflation component (for females), whilst

those for hatching success used a Tweedie distribution to account for under-dispersion.

We investigated the performance of all models using the DHARMa package (Florian, 2016).

Results

Hatching success

To assess the combined effects of heat stress, pesticide exposure, and microbiome composition

on male reproductive function, we first investigated fertilization success, estimated as the

proportion of eggs hatched out of all eggs laid by females. A GLM revealed a significant three-

way interaction between microbiome composition, temperature, and pesticide presence (N =

650, $\chi^2 = 16.07$, df = 3, p = 0.0011; Fig. 1, Supplementary Tables S1, S2), highlighting that the

combined effects of these environmental stressors depended on the microbial composition of

the focal males. Overall, males associated with all five bacterial strains exhibited the highest

hatching success, followed by *At-* and *Lb*-associated males, with axenic males being most

severely affected. These general patterns were also confirmed by the significant two-way

interactions of temperature × microbiome ($\chi^2 = 12.13$, df = 3, p = 0.007) and temperature ×

pesticide ($\chi^2 = 43.32$, df = 1, p < 0.0001), as well as by all main effects (all $\chi^2 \geq 520.76$, df ≥ 3,

p ≤ 0.001; Supplementary Tables S1, S2).

Male-mediated fecundity

Next, we investigated male-mediated fecundity, quantified as the total number of larvae

produced per female. In contrast to hatching success, the three-way interaction was not

significant ($\chi^2 = 0.319$, df = 3, p = 0.956; Supplementary Table S3, S4) for this metric and was

removed from the final model. The simplified model revealed significant two-way interactions

between microbiome and temperature (N = 650, $\chi^2 = 7.85$, df = 3, p = 0.049) and between

temperature and pesticide ($\chi^2 = 5.41$, df = 1, p = 0.020; Supplementary Tables S3, S4). All main

effects were again significant (all $\chi^2 \geq 5.41$, p ≤ 0.001). The relative differences in fecundity

between males with different microbiome statuses largely mirrored the patterns observed for

hatching success (Fig. 2).

Sperm competitiveness

Since sperm competitive ability is a crucial determinant of male reproductive fitness in

polygamous species and there are reports of adverse effects of increased temperatures and

pesticide exposure on sperm quality and function (Chirault et al., 2015; Meena, De Nardo, et

al., 2024; D. R. Moreira et al., 2022; S. Moreira et al., 2021; Roy et al., 2017; Sales et al.,

2018a, 2024), we evaluated the combined effects of our treatments on the proportion of

progeny sired by focal males relative to their standard competitors (i.e., P_2). Although the three-

way interaction was not statistically significant (N = 331, χ^2 = 1.003, df = 3, p = 0.800), all

two-way interactions were (Supplementary Tables S5, S6): temperature × microbiome (χ^2 =

22.92, df = 3, p < 0.001), microbiome × pesticide (χ^2 = 11.89, df = 3, p < 0.001), and

temperature × pesticide (χ^2 = 27.33, df = 1, p < 0.001). Similarly, all main effects on P_2 were

significant (all $\chi^2 \geq$ 472.33, p ≤ 0.001). These results indicate strong interdependent effects of

the treatments on sperm competitiveness, with the relative performance across different

microbiome groups again showing similar patterns to those observed for hatching success and

fecundity (Fig. 3).

Bacterial abundance in males in response to heat stress and pesticide exposure

We also analyzed the effect of heat stress and pesticide exposure and their interactions on the

growth and viability of commensal bacteria inside their host under the different stress

treatments by quantifying their colony forming units in homogenates of focal males

immediately after mating. For total CFU counts, we found a significant three-way interaction

(χ^2 = 5.06, df = 2, p = 0.008; Supplementary Tables S7, S8), suggesting that the bacterial strains

differed in their accumulation patterns in adult flies depending on the combined effects of

environmental stress factors. Additionally, we documented significant two-way interactions

and main effects of all three factors (all $\chi^2 \geq$ 42.31, p ≤ 0.001), except for the two-way

interaction between the two stressors (χ^2 = 2.36, df = 2, p = 0.127). Among these and relative

to the control microbiome, higher temperature had a stronger negative effect on the total

number of CFUs than pesticide exposure (Fig. 4; Supplementary Tables S7, S8).

A PERMANOVA controlling for block effects ($F_{1,35}$ = 0.20, p = 0.874, Supplementary Table

S9) revealed that temperature was also the dominant driver of variation in the relative

abundances of the five bacterial species constituting the consortium, explaining 26.3% of the

variation in community composition ($F_{1,35} = 13.88$, p = 0.0001; Supplementary Table S9). In

contrast, the main effect of pesticide was not statistically significant (2.2% of variation, $F_{1,35} =$

1.16, p = 0.314), and the temperature × pesticide interaction showed a marginal effect (4.8%

of variation, $F_{1,35} = 2.55$, p = 0.068). NMDS based on Bray-Curtis dissimilarities provided a

visual representation of community shifts across treatments. The NMDS ordination (Fig. 5)

indicated that the first axis (NMDS1) predominantly tracked the temperature gradient, while

the second axis (NMDS2) aligned more, but only weakly, with pesticide exposure. Overlaying

species vectors revealed notable differences in species associations (all $R^2 \geq 0.54$, $p \leq 0.0001$),

in that both *Lactobacillus* species showed positive (≥ 0.68), and all *Acetobacter* exhibited

negative (≤ -0.38), loadings on NMDS1 (Supplementary Table S10), suggesting an increase in

the relative abundance of *Acetobacter* in the high-temperature regimes. By contrast, loadings

on NMDS2 were negative for *A. tropicalis* (-0.92) and *L. brevis* (-0.44) but positive for all

remaining species (≥ 0.56). This axis tended to be biased toward negative values in the absence

of imidacloprid, but more ambiguous in its presence.

Finally, in a binomial GLMM focusing only on the relative abundances of acetic acid versus

lactic acid bacteria, where the three *Acetobacter* species should theoretically constitute 60% of

the total CFUs among the five species, temperature and pesticide presence interacted

significantly (N = 40, z = 2.15, p = 0.032). Specifically, the relative representation of

Acetobacter was increased to around 66% under heat stress (z = 2.84, p = 0.013), whereas

pesticide exposure reduced its representation (z = -2.32, p = 0.021), although only at the benign

temperature (Fig. 6).

Cross-generational effects on offspring fitness

Finally, we investigated whether the effects of paternal stress exposure carried over to the next

generation by measuring the hatching success and fecundity of F_1 offspring produced by F_0

males from each treatment group.

We found a significant interactive effect between paternal temperature and pesticide treatments

on the hatching success of both male (N = 726, χ^2 = 19.39, df = 1, p < 0.001; Supplementary

Tables S11,12) and female offspring (N = 750, χ^2 = 10.084, df = 1, p < 0.001; Supplementary

Tables S13,14). This indicates that the combined effects of heat and pesticide stress

experienced by fathers carried over to the next generation. However, the composition of the

paternal gut microbiome did not significantly influence offspring hatching success ($\chi^2 \leq 4.76$,

df = 3, p ≥ 0.19). The interactive effect of paternal temperature and pesticide exposure was not

significant for the fecundity induced by sons (χ^2 = 0.36, df = 1, p = 0.55; Supplementary Tables

S15,16). In contrast, the interaction between these two paternal stressors showed significant

effect on fecundity of F_1 daughters (N = 750, χ^2 = 3.990, df = 1, p = 0.006; Supplementary

Tables S17,18). There was also a non-significant trend for a main effect of paternal microbiome

composition on offspring fecundity in daughters (χ^2 = 7.26, df = 3, p = 0.06; Supplementary

Tables S17,18).

Discussion

We investigated the role of the gut microbiome in modulating male reproductive success in *D.*

melanogaster under individual and combined exposure to acute heat stress and the pesticide

imidacloprid. Our results confirm and extend previous work that demonstrated the detrimental

effects of sublethal temperatures (Canal Domenech & Fricke, 2022; David et al., 2005; Fasolo

& Krebs, 2004; Gandara & Drummond-Barbosa, 2022, 2023; Meena, De Nardo, et al., 2024;

Pilakouta & Baillet, 2022; Sales et al., 2024; B. S. Walsh, Parratt, Atkinson, et al., 2019) or

imidacloprid (Charpentier et al., 2014b; Xiao et al., 2016; Zhang, Xu, et al., 2021; Zhou et al.,

2017) on different reproductive functions in diverse organisms. However, our study adds a

crucial layer of complexity by revealing both the combined effects of these stressors and the

critical role of the microbiome in mediating the response of stressed males. Specifically, our

findings highlight the importance of the gut microbiome in mitigating fitness losses under

stress. *Acetobacter* generally offered stronger protection than *Lactobacillus*, while the complex

microbial community provided the highest benefits. Furthermore, we show that paternal stress

exposure imposes fitness costs on the next generation, although these cross-generational effects

were not significantly mediated by the paternal microbiome in our experimental design aimed

at minimizing direct microbial transmission.

Interactive effects of combined stressors on male reproduction

Our study revealed that initial exposure to imidacloprid followed by heat shock reduced male

reproductive functions more than either stressor alone. This observation aligns with previous

studies indicating complex, often order-dependent interactions between thermal stress and

pesticide toxicity (Ahmed et al., 2023; Deng et al., 2016; Meng et al., 2022; Ricupero et al.,

2020; Y.-C. Wang, Chang, Bai, Zhang, Iqbal, Lu, Gong, et al., 2021). For example, heat stress

has been shown to increase the tolerance to subsequent insecticide exposure in diverse insect

pests, such as the serpentine leaf miner, *Liriomyza trifoli* (Y.-C. Wang, Chang, Bai, Zhang,

Iqbal, Lu, Hu, et al., 2021a); diamond back moth, *Plutella xylostella* (Gu et al., 2010); wheat

aphid, *Sitobion avenae* (Cao et al., 2021), or brown planthopper, *Nilaparvata lugens* (Ge et al.,

2013). The order of stress treatments, however, has had an opposing effect in other insects

(Delnat et al., 2019; Y.-C. Wang, Chang, Bai, Zhang, Iqbal, Lu, Gong, et al., 2021). In *L. trifoli*,

for example, where both orders in stress exposure were examined, elevated temperatures

improved survival under exposure to the insecticide abamectin, whereas the reverse stress

sequence reduced survival (Y.-C. Wang, Chang, Bai, Zhang, Iqbal, Lu, Gong, et al., 2021),

indicating the importance of stress sequence in cross-protection between multiple stressors.

Pre-exposure to insecticides can also decrease heat stress tolerance, suggesting that warmer

temperatures may exacerbate insecticide toxicity depending on the exposure sequence.

Increased imidacloprid toxicity at higher temperatures has been documented in mayflies

(Macaulay et al., 2020), earthworms *Eisenia andrei* and collembolans *Folsomia candida*

(Bandeira et al., 2020). In the terrestrial ectotherm *Hypogastrura viatica*, sublethal

concentrations of imidacloprid became lethal on increasing the temperatures resulting in

reduced survival, growth and egg production (A. S. J. Silva et al., 2023). Our study corroborates

these findings, with decreased male reproductive success when pesticide exposure preceded

heat stress. These multi-stressor studies highlight the complex interplay between temperature

and toxicity, which can be positive or negative depending on the mode of action of the

insecticide, the route of exposure, and the insect species. Therefore, the sequence of exposure

must be carefully considered when evaluating combined stressor effects on insect performance.

The mechanisms underlying these sequence-dependent interactions likely involve intricate

physiological adjustments. For example, exposure to one stressor might compromise pathways

needed to cope with the subsequent stressor (lack of cross-tolerance) or trigger shared signaling

pathways that amplify susceptibility (negative cross-talk). Resource allocation trade-offs are

also probable (Sokolova, 2013), in that coping with pesticide toxicity may deplete energy

reserves or specific molecules (e.g., antioxidants, chaperones) needed to subsequently mount

an effective heat stress response. Generally, stressful environments may alter resource

allocation among essential processes like growth, reproduction, or immune defense to optimize

fitness outcomes (Kooijman, 2000). As such, elevated temperature may increase somatic

maintenance costs (Verberk et al., 2016), potentially reducing resources available for defense

and repair (Hallman & Brooks, 2015).

While the precise mechanisms underlying increased toxicity at high temperatures remain

unclear, they may overlap with those explaining the increased toxicity of organophosphate

pesticides at higher mean temperatures (Noyes et al., 2009; Noyes & Lema, 2015a). Increased

temperatures can enhance the activity of detoxification enzymes like cytochrome P450s (Y.-C.

Wang, Chang, Bai, Zhang, Iqbal, Lu, Hu, et al., 2021b), accelerating the breakdown of

pesticides like pyrethroids (Ahmad, 2007). Increased uptake of chemicals at higher

temperatures (Holmstrup et al., 2010), possibly due to higher basal metabolism (Hooper et al.,

2013b), increased membrane permeability (Noyes & Lema, 2015b), and or smaller body size

(Gardner et al., 2011) has also been observed. Stressful conditions can promote the

accumulation of reactive oxygen species (ROS) due to reduced respiration and increased water

loss, thereby negatively affecting development and reproduction (Dampc et al., 2021). For

example, low doses of imidacloprid have been shown to cause ROS formation and metabolic

impairment in *D. melanogaster* (Martelli et al., 2020). Both pesticide tolerance (Bagchi et al.,

1996) and heat stress (Neven, 2000) are also associated with overexpression of heat-shock

proteins like Hsp70 and Hsp90. In the honeybee, for example, these two Hsps followed similar

expression patterns in response to both thermal stress and imidacloprid exposure (Manzi et al.,

2020). This thermal stress-mediated chaperone activation could also be a relevant mechanism

in *D. melanogaster* males exposed to imidacloprid, for cellular protection against combined

stressors. If so, the observed fitness loss following heat and pesticide stress might at least in

part be attributable to the costs associated with enhanced Hsp expression (Hoekstra et al., 2013;

S. Li et al., 2015), which may become overwhelming under combined stress.

Microbiome as a modulator of stress resilience

A key finding of our study was the significant role of the gut microbiome in mitigating the

detrimental reproductive consequences of heat and pesticide stress. Across all measured fitness

components (hatching success, fecundity, sperm competitiveness), axenic males consistently

performed worse than males colonized with bacteria, highlighting the baseline importance of

the microbiome for reproductive function even under benign conditions, and its crucial role

under stress. The reduced fitness of axenic males under combined stress suggests an impaired

ability to metabolize and excrete imidacloprid, potentially leading to higher internal toxin loads

and thus increased vulnerability to subsequent heat stress.

Interestingly, the degree of protection conferred depended on the microbial composition. While

both *Lactobacillus brevis* and *Acetobacter tropicalis* mono-associations offered benefits over

the axenic state, the 5-species community consistently provided the greatest resilience,

resulting in the highest reproductive fitness across most conditions. This underscores the

potential importance of microbial diversity and inter-species interactions for robust host

function (Coyte et al., 2015; Freilich et al., 2011; Levy & Borenstein, 2013; Magnúsdóttir et

al., 2017; Noecker et al., 2019; Zelezniak et al., 2015). Mono-associations might lack the

metabolic versatility or synergistic cross-feeding capabilities present in more complex

consortia, limiting their ability to buffer the host against multiple stressors.

Our results partially contrast with Daisley et al. (2017), who reported that *Lactobacillus*

plantarum (Lp39) supplementation rescued *D. melanogaster* from harmful imidacloprid

effects, accompanied by increased *Acetobacter* and *Lactobacillus* abundance. We did not

observe such a strong rescue with *L. brevis*, nor an increased abundance of either bacterial

genus. This discrepancy could stem from strain-specific functions, similar to opposing effects

on chlorpyrifos (CP) toxicity between *L. plantarum* ISO and *Lactobacillus ATCC* in *D.*

melanogaster (Daisley et al., 2017). Alternatively, differences in experimental protocols could

also contribute to the contrasting results between our and Daisley's studies (e.g., our acute 18h

imidacloprid exposure versus Daisley et al.'s (2017) continued exposure).

We also observed shifts in the relative abundance of *Acetobacter* and *Lactobacillus* within the

host, aligning with studies showing heat-induced microbiome disruptions (Jaramillo &

Castañeda, 2021; Kokou et al., 2018; Moghadam et al., 2018a). Apparently, in mono-

associations, *Lactobacillus* was more abundant and less affected by treatments than

Acetobacter, however, the latter seemed to be overrepresented in 5-species consortium relative

to the expected proportion. Generally, we found a higher proportional representation of

Acetobacter following acute heat stress, consistent with prior work reporting increased

Acetobacter (but fewer *Leuconostoc* bacteria) in heat- compared to cold-acclimated flies

(Moghadam et al., 2018b). While this previous study exposed individuals to thermal treatments

throughout development, our findings demonstrate that such compositional shifts can occur

rapidly (within hours).

The mechanisms by which the microbiome enhances stress tolerance likely involve multiple

pathways. Beyond direct detoxification (Cheng et al., 2017; Jones et al., 2015; Kikuchi et al.,

2012; Pang et al., 2018), gut microbes significantly impact host nutrition and metabolism

(Douglas, 2009; Ridley et al., 2012), which are intrinsically linked to thermal tolerance in

ectotherms (Henry et al., 2020; Moeller et al., 2020; Moghadam et al., 2018a; Semsar-

Kazerouni et al., 2020). It has been shown in *D. melanogaster* that the gut microbiota influences

nutrient allocation, including levels of the thermo-protectant trehalose (Overgaard et al., 2007;

Ridley et al., 2012; Schwasinger-Schmidt et al., 2012). They can also modulate host immune

responses and potentially Hsp expression in the gut epithelium (Arnal & Lalles, 2016; Liu et

al., 2014). Given the critical role of Hsps in mitigating thermal stress in *D. melanogaster*

(Calabria et al., 2012; Sørensen et al., 2003), and their rapid induction after heat exposure

(Hoekstra et al., 2013), the microbiota may enhance heat tolerance in conventional and mono-

associated flies by promoting higher Hsp levels. Disrupting the microbiome can alter these

processes. For instance, heat-induced reductions in the abundance of the bacterial

symbiont *Buchnera aphidicola* in pea aphids lowered heat resistance, with fitness costs for the

host due to a deficiency of *Buchnera*-derived essential amino acids (Dunbar et al.,

2007). Similarly, temperature-driven microbiome shifts have been reported in mice (Chevalier

et al., 2015) and *Nilaparvata lugens*, with the disrupted microbiome assembly in the latter case

being harmful by increasing imidacloprid susceptibility (Zhang, Cai, et al., 2021).

Cross-generational impacts of parental stress

We found clear evidence for cross-generational fitness costs of paternal stress exposure, with

reduced hatching success and fecundity in offspring of fathers exposed to heat and/or pesticide

stress. Cross-generational effects can either increase (Pölkki et al., 2012; Schultz et al., 2016)

or decrease (Brausch & Salice, 2011; Kim et al., 2014; Reátegui-Zirena et al., 2017) offspring

vulnerability to stressors compared to the parental generation. Our results are consistent with

studies reporting detrimental cross- or transgenerational effects of paternal heat stress (Ferrer

et al., 2013; Guillaume et al., 2016; Shama et al., 2014; M. R. Walsh et al., 2014) or toxicant

exposure (Bhandari et al., 2015; Kimberly & Salice, 2015; Schultz et al., 2016; Yu & Liao,

2016), possibly through epigenetic processes (Schultz et al., 2016). The observed synergism

between paternal heat and pesticide exposure in offspring fitness underscores the potential for

long-lasting consequences of multi-stressor events.

Offspring from stressed fathers exhibited reduced fecundity and hatching success and did not

demonstrate increased tolerance to the same stressors, indicating that paternal exposure to

warming and/or pesticide does not buffer against these environmental stressors. These results

contrast with a growing number of studies suggesting that cross-generational effects can buffer

offspring against rapid environmental change (Donelson et al., 2018; Munday, 2014), with

parental heat exposure even being considered as a strategy to enhance species resilience

(Chakravarti et al., 2017). Instead, our findings align with previous work revealing negative

effects of heat and pesticide stress on offspring survival (Tran et al., 2018) and support the

view that cross-generational effects of stress are often non-adaptive or costly for offspring

(Uller et al., 2013), highlighting the potential for cumulative fitness declines across generations

in populations facing recurrent multi-stressor events.

Intriguingly, the paternal microbiome did not significantly mediate these cross-generational

effects in our study. This could be attributable to our experimental design, which deliberately

minimized vertical transmission or environmental acquisition of microbes by using axenic

mothers and transferring these and their offspring to sterile food vials immediately after mating

or hatching, respectively. This allowed us to isolate direct paternal effects (potentially

epigenetic or related to gamete quality) based on the sire's microbial status. However, we

cannot rule out a role for the paternal microbiome if transmission occurs naturally (e.g., during

copulation) (Guilhot et al., 2025), which then triggers differential maternal effects or microbial

deposition on the chorion of eggs (direct paternal transmission of microbes to their offspring

seems rather unlikely as males have little control over female oviposition). Future studies

explicitly allowing or manipulating vertical transmission would be needed to explore this.

Conclusions and broader implications

Our study demonstrates that acute heat stress and sublethal exposure to pesticides (e.g.,

imidacloprid) interact, often synergistically, to negatively impact male reproductive fitness in

D. melanogaster, with consequences extending to offspring viability and fecundity. Crucially,

we show that the gut microbiome is a key determinant of resilience, offering significant

protection against these stressors, with benefits enhanced by microbial complexity. However,

the microbiome itself is vulnerable to disruption by these same stressors.

Our findings underscore the critical need to consider multi-stressor interactions and the role of

host-associated microbes when assessing the ecological risks posed by climate change and

agrochemical use. Predicting population responses requires moving beyond single-stressor

studies to understand these complex interdependencies. The observed negative cross-

generational effects further suggest that transient stress events could have prolonged

demographic consequences. From a conservation perspective, these results highlight the

potential vulnerability of insect populations, particularly those with depleted or altered

microbiomes, to the combined pressures of warming climates and continued pesticide

application. Future research should explore these interactions across diverse species,

environments, and stressor combinations to improve ecological forecasting and develop

strategies for mitigating biodiversity loss in the Anthropocene.

Acknowledgements

We thank T. Ratz and G. Raina for valuable comments on our manuscript. This project was

funded by the Swiss National Science Foundation (grant PP00P3_202662 to S.L.) and a

University of Zurich Postdoctoral Fellowship (to K.M.).

References

Ahmad, M. (2007). Insecticide resistance mechanisms and their management in *Helicoverpa*

armigera (Hübner)–A review. *J. Agric. Res*, *45*(4), 319–335.

Ahmed, T., Masood, H. A., Noman, M., Al-Huqail, A. A., Alghanem, S. M., Khan, M. M.,
Muhammad, S., Manzoor, N., Rizwan, M., & Qi, X. (2023). Biogenic silicon
nanoparticles mitigate cadmium (Cd) toxicity in rapeseed (*Brassica napus L.*) by
modulating the cellular oxidative stress metabolism and reducing Cd translocation.
Journal of Hazardous Materials, *459*, 132070.

Akbar, R., Khan, I. A., Alajmi, R. A., Ali, A., Faheem, B., Usman, A., Ahmed, A. M., El-
Shazly, M., Farid, A., & Giesy, J. P. (2022). Evaluation of insecticidal potentials of
five plant extracts against the stored grain pest, *Callosobruchus maculatus*
(Coleoptera: Bruchidae). *Insects*, *13*(11), 1047.

Arambourou, H., & Stoks, R. (2015). Combined effects of larval exposure to a heat wave and
chlorpyrifos in northern and southern populations of the damselfly *Ischnura elegans*.
Chemosphere, *128*, 148–154.

Arnal, M.-E., & Lalles, J.-P. (2016). Gut epithelial inducible heat-shock proteins and their
modulation by diet and the microbiota. *Nutrition Reviews*, *74*(3), 181–197.

Aulsebrook, L. C., Bertram, M. G., Martin, J. M., Aulsebrook, A. E., Brodin, T., Evans, J. P.,
Hall, M. D., O'Bryan, M. K., Pask, A. J., & Tyler, C. R. (2020). Reproduction in a
polluted world: Implications for wildlife. *Reproduction*, *160*(2), R13–R23.

Aulsebrook, L. C., Bertram, M. G., Martin, J. M., Aulsebrook, A. E., Brodin, T., Evans, J. P.,
Hall, M. D., O'Bryan, M. K., Pask, A. J., Tyler, C. R., & Wong, B. B. M. (2020).
Reproduction in a polluted world: Implications for wildlife. *Reproduction*, *160*(2),
R13–R23. https://doi.org/10.1530/REP-20-0154

Bagchi, D., Bhaitacharya, G., & Stohs, S. (1996). Production of reactive oxygen species by
gastric cells in association with *Helicobacter pylori. Free Radical Research*, *24*(6),
439–450.

Bandeira, F. O., Alves, P. R. L., Hennig, T. B., Toniolo, T., Natal-da-Luz, T., & Baretta, D. (2020). Effect of temperature on the toxicity of imidacloprid to *Eisenia andrei* and *Folsomia candida* in tropical soils. *Environmental Pollution*, *267*, 115565.

Beketov, M. A., Kefford, B. J., Schäfer, R. B., & Liess, M. (2013). Pesticides reduce regional biodiversity of stream invertebrates. *Proceedings of the National Academy of Sciences*, *110*(27), 11039–11043.

Bhandari, R. K., Vom Saal, F. S., & Tillitt, D. E. (2015). Transgenerational effects from early developmental exposures to bisphenol A or 17α-ethinylestradiol in medaka, Oryzias latipes. *Scientific Reports*, *5*(1), 9303.

Bonduriansky, R., Crean, A. J., & Day, T. (2012). The implications of nongenetic inheritance for evolution in changing environments. *Evolutionary Applications*, *5*(2), 192–201. https://doi.org/10.1111/j.1752-4571.2011.00213.x

Brausch, J. M., & Salice, C. J. (2011). Effects of an environmentally realistic pesticide mixture on *Daphnia magna* exposed for two generations. *Archives of Environmental Contamination and Toxicology*, *61*, 272–279.

Broderick, N. A., & Lemaitre, B. (2012). Gut-associated microbes of *Drosophila melanogaster*. *Gut Microbes*, *3*(4), 307–321.

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling.

Calabria, G., Dolgova, O., Rego, C., Castañeda, L., Rezende, E., Balanyà, J., Pascual, M., Sørensen, J. G., Loeschcke, V., & Santos, M. (2012). Hsp70 protein levels and thermotolerance in *Drosophila subobscura*: A reassessment of the thermal co-adaptation hypothesis. *Journal of Evolutionary Biology*, *25*(4), 691–700.

Canal Domenech, B., & Fricke, C. (2022). Recovery from heat-induced infertility—A study

of reproductive tissue responses and fitness consequences in male *Drosophila*

melanogaster. *Ecology and Evolution*, *12*(12), e9563.

Cao, J., Xing, K., & Zhao, F. (2021). Complex delayed and transgenerational effects driven

by the interaction of heat and insecticide in the maternal generation of the wheat

aphid, *Sitobion avenae*. *Pest Management Science*, *77*(10), 4453–4461.

Chakravarti, L. J., Beltran, V. H., & van Oppen, M. J. (2017). Rapid thermal adaptation in

photosymbionts of reef-building corals. *Global Change Biology*, *23*(11), 4675–4688.

Charpentier, G., Louat, F., Bonmatin, J.-M., Marchand, P. A., Vanier, F., Locker, D., &

Decoville, M. (2014a). Lethal and sublethal effects of imidacloprid, after chronic

exposure, on the insect model *Drosophila melanogaster*. *Environmental Science &*

Technology, *48*(7), 4096–4102.

Charpentier, G., Louat, F., Bonmatin, J.-M., Marchand, P. A., Vanier, F., Locker, D., &

Decoville, M. (2014b). Lethal and Sublethal Effects of Imidacloprid, After Chronic

Exposure, On the Insect Model *Drosophila melanogaster*. *Environmental Science &*

Technology, *48*(7), 4096–4102. https://doi.org/10.1021/es405331c

Cheng, D., Guo, Z., Riegler, M., Xi, Z., Liang, G., & Xu, Y. (2017). Gut symbiont enhances

insecticide resistance in a significant pest, the oriental fruit fly *Bactrocera dorsalis*

(Hendel). *Microbiome*, *5*, 1–12.

Chevalier, C., Stojanović, O., Colin, D. J., Suarez-Zamorano, N., Tarallo, V., Veyrat-

Durebex, C., Rigo, D., Fabbiano, S., Stevanović, A., Hagemann, S., Montet, X.,

Seimbille, Y., Zamboni, N., Hapfelmeier, S., & Trajkovski, M. (2015). Gut

Microbiota Orchestrates Energy Homeostasis during Cold. *Cell*, *163*(6), 1360–1374.

https://doi.org/10.1016/j.cell.2015.11.004

Chirault, M., Lucas, C., Goubault, M., Chevrier, C., Bressac, C., & Lécureuil, C. (2015). A

combined approach to heat stress effect on male fertility in *Nasonia vitripennis*: From

the physiological consequences on spermatogenesis to the reproductive adjustment of

females mated with stressed males. *PLoS One*, *10*(3), e0120656.

Coyte, K. Z., Schluter, J., & Foster, K. R. (2015). The ecology of the microbiome: Networks,

competition, and stability. *Science*, *350*(6261), 663–666.

Crean, A. J., & Immler, S. (2021). Evolutionary consequences of environmental effects on

gamete performance. *Philosophical Transactions of the Royal Society B*, *376*(1826),

20200122. https://doi.org/10.1098/rstb.2020.0122

Dai, W., Holmstrup, M., Slotsbo, S., Bakker, R., Damgaard, C., & van Gestel, C. A. (2023).

Heat stress delays detoxification of phenanthrene in the springtail *Folsomia candida.*

Chemosphere, *311*, 137119.

Daisley, B. A., Trinder, M., McDowell, T. W., Welle, H., Dube, J. S., Ali, S. N., Leong, H.

S., Sumarah, M. W., & Reid, G. (2017). Neonicotinoid-induced pathogen

susceptibility is mitigated by *Lactobacillus plantarum* immune stimulation in a

Drosophila melanogaster model. *Scientific Reports*, *7*(1), 2703.

Dampc, J., Mołoń, M., Durak, T., & Durak, R. (2021). Changes in aphid—Plant interactions

under increased temperature. *Biology*, *10*(6), 480.

David, J., Araripe, L., Chakir, M., Legout, H., Lemos, B., Pétavy, G., Rohmer, C., Joly, D., &

Moreteau, B. (2005). Male sterility at extreme temperatures: A significant but

neglected phenomenon for understanding *Drosophila* climatic adaptations. *Journal of*

Evolutionary Biology, *18*(4), 838–846.

De França, S. M., Breda, M. O., Barbosa, D. R., Araujo, A., Guedes, C. A., & Shields, V.

(2017). The sublethal effects of insecticides in insects. *Biological Control of Pest and*

Vector Insects, *10*, 66461.

DeCourten, B. M., & Brander, S. M. (2017). Combined effects of increased temperature and

endocrine disrupting pollutants on sex determination, survival, and development

across generations. *Scientific Reports*, *7*(1), 9310. https://doi.org/10.1038/s41598-017-

09631-1

Delnat, V., Janssens, L., & Stoks, R. (2019). Whether warming magnifies the toxicity of a

pesticide is strongly dependent on the concentration and the null model. *Aquatic*

Toxicology, *211*, 38–45.

Deng, Z., Zhang, F., Wu, Z., Yu, Z., & Wu, G. (2016). Chlorpyrifos-induced hormesis in

insecticide-resistant and-susceptible Plutella xylostella under normal and high

temperatures. *Bulletin of Entomological Research*, *106*(3), 378–386.

Desneux, N., Decourtye, A., & Delpuech, J.-M. (2007). The sublethal effects of pesticides on

beneficial arthropods. *Annual Review of Entomology*, *52*(1), 81–106.

https://doi.org/10.1146/annurev.ento.52.110405.091440

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D.

C., & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms

across latitude. *Proceedings of the National Academy of Sciences of the United States*

of America, *105*(18), 6668–6672. https://doi.org/10.1073/pnas.0709472105

Deutsch, C. A., Tewksbury, J. J., Tigchelaar, M., Battisti, D. S., Merrill, S. C., Huey, R. B.,

& Naylor, R. L. (2018). Increase in crop losses to insect pests in a warming climate.

Science, *361*(6405), 916–919.

Donelson, J. M., Salinas, S., Munday, P. L., & Shama, L. N. S. (2018). Transgenerational

plasticity and climate change experiments: Where do we go from here? *Global*

Change Biology, *24*(1), 13–34. https://doi.org/10.1111/gcb.13903

Dougherty, L. R., Frost, F., Maenpaa, M. I., Rowe, M., Cole, B. J., Vasudeva, R., Pottier, P.,

Schultner, E., Macartney, E. L., & Lindenbaum, I. (2024). A systematic map of

studies testing the relationship between temperature and animal reproduction.

Ecological Solutions and Evidence, *5*(1), e12303.

Douglas, A. E. (2009). The microbial dimension in insect nutritional ecology. *Functional*

Ecology, *23*(1), 38–47.

Dunbar, H. E., Wilson, A. C. C., Ferguson, N. R., & Moran, N. A. (2007). Aphid thermal

tolerance is governed by a point mutation in bacterial symbionts. *PLoS Biology*, *5*(5),

e96.

Evariste, L., Barret, M., Mottier, A., Mouchet, F., Gauthier, L., & Pinelli, E. (2019). Gut

microbiota of aquatic organisms: A key endpoint for ecotoxicological studies.

Environmental Pollution, *248*, 989–999.

Fasolo, A. G., & Krebs, R. A. (2004). A comparison of behavioural change in *Drosophila*

during exposure to thermal stress. *Biological Journal of the Linnean Society*, *83*(2),

197–205.

Fathy, R. F. (2024). Divergent perspectives on the synergistic impacts of thermal-chemical

stress on aquatic biota within the framework of climate change scenarios.

Chemosphere, *355*, 141810. https://doi.org/10.1016/j.chemosphere.2024.141810

Feldhaar, H. (2011). Bacterial symbionts as mediators of ecologically important traits of

insect hosts. *Ecological Entomology*, *36*(5), 533–543. https://doi.org/10.1111/j.1365-

2311.2011.01318.x

Ferrer, A., Dorn, S., & Mazzi, D. (2013). Cross-generational effects of temperature on flight

performance, and associated life-history traits in an insect. *Journal of Evolutionary*

Biology, *26*(11), 2321–2330.

Fitzpatrick, J. L., & Lüpold, S. (2014). Sexual selection and the evolution of sperm quality.

Molecular Human Reproduction, *20*(12), 1180–1189.

Florian, H. (2016). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. *CRAN: Contributed Packages*.

https://doi.org/10.32614/cran.package.dharma

Folt, C., Chen, C., Moore, M., & Burnaford, J. (1999). Synergism and antagonism among

multiple stressors. *Limnology and Oceanography*, *44*(3part2), 864–877.

Fournier-Level, A., Neumann-Mondlak, A., Good, R., Green, L., Schmidt, J. M., & Robin, C.

(2016). Behavioural response to combined insecticide and temperature stress in

natural populations of *Drosophila melanogaster*. *Journal of Evolutionary Biology*,

29(5), 1030–1044.

Frantzios, G., Paptsiki, K., Sidiropoulou, B., Lazaridis, I., Theophilidis, G., & Mavragani-

Tsipidou, P. (2008). Evaluation of insecticidal and genotoxic effects of imidacloprid

and acetochlor in *Drosophila melanogaster*. *Journal of Applied Entomology*, *132*(7),

583–590.

Freilich, S., Zarecki, R., Eilam, O., Segal, E. S., Henry, C. S., Kupiec, M., Gophna, U.,

Sharan, R., & Ruppin, E. (2011). Competitive and cooperative metabolic interactions

in bacterial communities. *Nature Communications*, *2*(1), 589.

Gandara, A. C. P., & Drummond-Barbosa, D. (2022). Warm and cold temperatures have

distinct germline stem cell lineage effects during *Drosophila* oogenesis. *Development*,

149(5), dev200149.

Gandara, A. C. P., & Drummond-Barbosa, D. (2023). Chronic exposure to warm temperature

causes low sperm abundance and quality in *Drosophila melanogaster*. *Scientific*

Reports, *13*(1), 12331.

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body

size: A third universal response to warming? *Trends in Ecology & Evolution*, *26*(6),

285–291.

Ge, L., Huang, L., Yang, G., Song, Q., Stanley, D., Gurr, G., & Wu, J. (2013). Molecular

basis for insecticide-enhanced thermotolerance in the brown planthopper *Nilaparvata*

lugens S tål (H emiptera: D elphacidae). *Molecular Ecology*, *22*(22), 5624–5634.

Gould, A. L., Zhang, V., Lamberti, L., Jones, E. W., Obadia, B., Korasidis, N., Gavryushkin,

A., Carlson, J. M., Beerenwinkel, N., & Ludington, W. B. (2018). Microbiome

interactions shape host fitness. *Proceedings of the National Academy of Sciences*,

115(51), E11951–E11960.

Gruntenko, N. E., Ilinsky, Y. Yu., Adonyeva, N. V., Burdina, E. V., Bykov, R. A.,

Menshanov, P. N., & Rauschenbach, I. Yu. (2017). Various Wolbachia genotypes

differently influence host *Drosophila* dopamine metabolism and survival under heat

stress conditions. *BMC Evolutionary Biology*, *17*(2), 252.

https://doi.org/10.1186/s12862-017-1104-y

Gu, X., Tian, S., Wang, D., Fei, G., & Wei, H. (2010). Interaction between short-term heat

pretreatment and fipronil on 2nd instar larvae of diamondback moth, *Plutella*

xylostella (Linn). *Dose-Response*, *8*(3), dose-response.

Guilhot, R., Xuéreb, A., & Fellous, S. (2025). Transmission of yeast and bacterial symbionts

between sexual partners in *Drosophila suzukii* and *Drosophila melanogaster*. *Royal*

Society Open Science, *12*(2), 241149.

Guillaume, A. S., Monro, K., & Marshall, D. J. (2016). Transgenerational plasticity and

environmental stress: Do paternal effects act as a conduit or a buffer? *Functional*

Ecology, *30*(7), 1175–1184.

Hallman, T. A., & Brooks, M. L. (2015). The deal with diel: Temperature fluctuations,

asymmetrical warming, and ubiquitous metals contaminants. *Environmental Pollution*

(Barking, Essex: 1987), *206*, 88–94. https://doi.org/10.1016/j.envpol.2015.06.005

He, Y., Zhao, J., Wu, D., Wyckhuys, K. A., & Wu, K. (2011). Sublethal effects of

imidacloprid on *Bemisia tabaci* (Hemiptera: Aleyrodidae) under laboratory

conditions. *Journal of Economic Entomology*, *104*(3), 833–838.

Hector, T. E., Hoang, K. L., Li, J., & King, K. C. (2022). Symbiosis and host responses to

heating. *Trends in Ecology & Evolution*, *37*(7), 611–624.

Heino, J., Virkkala, R., & Toivonen, H. (2009). Climate change and freshwater biodiversity:

Detected patterns, future trends and adaptations in northern regions. *Biological*

Reviews, *84*(1), 39–54.

Henry, Y., Overgaard, J., & Colinet, H. (2020). Dietary nutrient balance shapes phenotypic

traits of *Drosophila melanogaster* in interaction with gut microbiota. *Comparative*

Biochemistry and Physiology Part A: Molecular & Integrative Physiology, *241*,

110626.

Hoekstra, L. A., Siddiq, M. A., & Montooth, K. L. (2013). Pleiotropic effects of a

mitochondrial–nuclear incompatibility depend upon the accelerating effect of

temperature in *Drosophila. Genetics*, *195*(3), 1129–1139.

Holmstrup, M., Bindesbøl, A.-M., Oostingh, G. J., Duschl, A., Scheil, V., Köhler, H.-R.,

Loureiro, S., Soares, A. M., Ferreira, A. L., & Kienle, C. (2010). Interactions between

effects of environmental chemicals and natural stressors: A review. *Science of the*

Total Environment, *408*(18), 3746–3762.

Hooper, M. J., Ankley, G. T., Cristol, D. A., Maryoung, L. A., Noyes, P. D., & Pinkerton, K.

E. (2013a). Interactions between chemical and climate stressors: A role for

mechanistic toxicology in assessing climate change risks. *Environmental Toxicology*

and Chemistry, *32*(1), 32–48.

Hooper, M. J., Ankley, G. T., Cristol, D. A., Maryoung, L. A., Noyes, P. D., & Pinkerton, K.

E. (2013b). Interactions between chemical and climate stressors: A role for

809   mechanistic toxicology in assessing climate change risks. *Environmental Toxicology*

810   *and Chemistry*, *32*(1), 32–48.

811 Ihara, M., D. Buckingham, S., Matsuda, K., & B. Sattelle, D. (2017). Modes of action,

812   resistance and toxicity of insecticides targeting nicotinic acetylcholine receptors.

813   *Current Medicinal Chemistry*, *24*(27), 2925–2934.

814 Iossa, G. (2019). Sex-specific differences in thermal fertility limits. *Trends in Ecology &*

815   *Evolution*, *34*(6), 490–492. https://doi.org/10.1016/j.tree.2019.02.016

816 Jaramillo, A., & Castañeda, L. E. (2021). Gut microbiota of *Drosophila subobscura*

817   contributes to its heat tolerance and is sensitive to transient thermal stress. *Frontiers*

818   *in Microbiology*, *12*, 654108.

819 Jones, R. M., Desai, C., Darby, T. M., Luo, L., Wolfarth, A. A., Scharer, C. D., Ardita, C. S.,

820   Reedy, A. R., Keebaugh, E. S., & Neish, A. S. (2015). *Lactobacilli* modulate

821   epithelial cytoprotection through the Nrf2 pathway. *Cell Reports*, *12*(8), 1217–1225.

822 Jutfelt, F., Norin, T., Åsheim, E. R., Rowsey, L. E., Andreassen, A. H., Morgan, R., Clark, T.

823   D., & Speers-Roesch, B. (2021). 'Aerobic scope protection'reduces ectotherm growth

824   under warming. *Functional Ecology*, *35*(7), 1397–1407.

825 Kang, Z.-W., Liu, F.-H., Pang, R.-P., Tian, H.-G., & Liu, T.-X. (2018). Effect of Sublethal

826   Doses of Imidacloprid on the Biological Performance of Aphid Endoparasitoid

827   *Aphidius gifuensis* (Hymenoptera: Aphidiidae) and Influence on Its Related Gene

828   Expression. *Frontiers in Physiology*, *9*, 1729.

829   https://doi.org/10.3389/fphys.2018.01729

830 Karimzadeh, R., Javanshir, M., & Hejazi, M. J. (2020). Individual and combined effects of

831   insecticides, inert dusts and high temperatures on *Callosobruchus maculatus*

832   (coleoptera: *Chrysomelidae*). *Journal of Stored Products Research*, *89*, 101693.

833   Khosravi, A., & Mazmanian, S. K. (2013). Disruption of the gut microbiome as a risk factor

834   for microbial infections. *Current Opinion in Microbiology*, *16*(2), 221–227.

835   Kikuchi, Y., Hayatsu, M., Hosokawa, T., Nagayama, A., Tago, K., & Fukatsu, T. (2012).

836   Symbiont-mediated insecticide resistance. *Proceedings of the National Academy of*

837   *Sciences*, *109*(22), 8618–8622.

838   Kim, H. Y., Yu, S., Jeong, T., & Kim, S. D. (2014). Relationship between trans-generational

839   effects of tetracycline on *Daphnia magna* at the physiological and whole organism

840   level. *Environmental Pollution*, *191*, 111–118.

841   Kimberly, D. A., & Salice, C. J. (2015). Multigenerational contaminant exposures produce

842   non-monotonic, transgenerational responses in *Daphnia magna. Environmental*

843   *Pollution (Barking, Essex: 1987)*, *207*, 176–182.

844   https://doi.org/10.1016/j.envpol.2015.09.020

845   Kokou, F., Sasson, G., Nitzan, T., Doron-Faigenboim, A., Harpaz, S., Cnaani, A., & Mizrahi,

846   I. (2018). Host genetic selection for cold tolerance shapes microbiome composition

847   and modulates its response to temperature. *Elife*, *7*, e36398.

848   Kooijman, S. A. L. M. (2000). Dynamic energy and mass budgets in biological systems.

849   Cambridge university press.

850   Koyle, M. L., Veloz, M., Judd, A. M., Wong, A. C.-N., Newell, P. D., Douglas, A. E., &

851   Chaston, J. M. (2016). Rearing the fruit fly *Drosophila melanogaster* under axenic

852   and gnotobiotic conditions. *JoVE (Journal of Visualized Experiments)*, *113*, e54219.

853   Kumar, V., & Kumar, P. (2019). Pesticides in agriculture and environment: Impacts on

854   human health. *Contaminants in Agriculture and Environment: Health Risks and*

855   *Remediation*, *1*, 76.

Levy, R., & Borenstein, E. (2013). Metabolic modeling of species interaction in the human

microbiome elucidates community-level assembly rules. *Proceedings of the National*

Academy of Sciences, *110*(31), 12804–12809.

Li, S., Tan, H.-Y., Wang, N., Zhang, Z.-J., Lao, L., Wong, C.-W., & Feng, Y. (2015). The

role of oxidative stress and antioxidants in liver diseases. *International Journal of*

Molecular Sciences, *16*(11), 26087–26124.

Li, W., Lu, Z., Li, L., Yu, Y., Dong, S., Men, X., & Ye, B. (2018). Sublethal effects of

imidacloprid on the performance of the bird cherry-oat aphid *Rhopalosiphum padi*.

PLOS ONE, *13*(9), e0204097. https://doi.org/10.1371/journal.pone.0204097

Liu, H., He, J., Chi, C., & Shao, J. (2014). Differential HSP70 expression in *Mytilus coruscus*

under various stressors. *Gene*, *543*(1), 166–173.

Lüpold, S., de Boer, R. A., Evans, J. P., Tomkins, J. L., & Fitzpatrick, J. L. (2020). How

sperm competition shapes the evolution of testes and sperm: A meta-analysis.

Philosophical Transactions of the Royal Society B, *375*(1813), 20200064.

Lüpold, S., Manier, M. K., Berben, K. S., Smith, K. J., Daley, B. D., Buckley, S. H., Belote,

J. M., & Pitnick, S. (2012). How multivariate ejaculate traits determine competitive

fertilization success in *Drosophila melanogaster*. *Current Biology*, *22*(18), 1667–

1672.

Macaulay, S. J., Buchwalter, D. B., & Matthaei, C. D. (2020). Water temperature interacts

with the insecticide imidacloprid to alter acute lethal and sublethal toxicity to mayfly

larvae. *New Zealand Journal of Marine and Freshwater Research*, *54*(1), 115–130.

Magnúsdóttir, S., Heinken, A., Kutt, L., Ravcheev, D. A., Bauer, E., Noronha, A.,

Greenhalgh, K., Jäger, C., Baginska, J., & Wilmes, P. (2017). Generation of genome-

scale metabolic reconstructions for 773 members of the human gut microbiota. *Nature*

Biotechnology, *35*(1), 81–89.

Manzi, C., Vergara-Amado, J., Franco, L. M., & Silva, A. X. (2020). The effect of

882   temperature on candidate gene expression in the brain of honey bee *Apis mellifera*

883   (Hymenoptera: Apidae) workers exposed to neonicotinoid imidacloprid. *Journal of*

884   *Thermal Biology*, *93*, 102696. https://doi.org/10.1016/j.jtherbio.2020.102696

885   Mao, K., Jin, R., Li, W., Ren, Z., Qin, X., He, S., Li, J., & Wan, H. (2019). The influence of

886   temperature on the toxicity of insecticides to *Nilaparvata lugens* (Stål). *Pesticide*

887   *Biochemistry and Physiology*, *156*, 80–86.

888   Marshall, D. J., Heppell, S. S., Munch, S. B., & Warner, R. R. (2010). The relationship

889   between maternal phenotype and offspring quality: Do older mothers really produce

890   the best offspring? *Ecology*, *91*(10), 2862–2873. https://doi.org/10.1890/09-0156.1

891   Martelli, F., Zhongyuan, Z., Wang, J., Wong, C.-O., Karagas, N. E., Roessner, U.,

892   Rupasinghe, T., Venkatachalam, K., Perry, T., & Bellen, H. J. (2020). Low doses of

893   the neonicotinoid insecticide imidacloprid induce ROS triggering neurological and

894   metabolic impairments in *Drosophila*. *Proceedings of the National Academy of*

895   *Sciences*, *117*(41), 25840–25850.

896   Massot, M., Bagni, T., Maria, A., Couzi, P., Drozdz, T., Malbert-Colas, A., Maïbèche, M., &

897   Siaussat, D. (2021). Combined influences of transgenerational effects, temperature

898   and insecticide on the moth *Spodoptera littoralis*. *Environmental Pollution*, *289*,

899   117889.

900   Meena, A., De Nardo, A. N., Maggu, K., Sbilordo, S. H., Roy, J., Snook, R. R., & Lüpold, S.

901   (2024). Fertility loss and recovery dynamics after repeated heat stress across life

902   stages in male *Drosophila melanogaster*: Patterns and processes. *Royal Society Open*

903   *Science*, *11*(10), 241082.

904   Meena, A., Maggu, K., De Nardo, A. N., Sbilordo, S. H., Eggs, B., Al Toma Sho, R., &

905   Lüpold, S. (2024). Life stage-specific effects of heat stress on spermatogenesis and

oogenesis in *Drosophila melanogaster*. *Journal of Thermal Biology*, *125*, 104001.

https://doi.org/10.1016/j.jtherbio.2024.104001

Meng, S., Tran, T. T., Van Dinh, K., Delnat, V., & Stoks, R. (2022). Acute warming

increases pesticide toxicity more than transgenerational warming by reducing the

energy budget. *Science of The Total Environment*, *805*, 150373.

Moeller, A. H., Ivey, K., Cornwall, M. B., Herr, K., Rede, J., Taylor, E. N., & Gunderson, A.

R. (2020). The lizard gut microbiome changes with temperature and is associated with

heat tolerance. *Applied and Environmental Microbiology*, *86*(17), e01181-20.

Moghadam, N. N., Thorshauge, P. M., Kristensen, T. N., de Jonge, N., Bahrndorff, S.,

Kjeldal, H., & Nielsen, J. L. (2018b). Strong responses of *Drosophila melanogaster*

microbiota to developmental temperature. *Fly*, *12*(1), 1–12.

https://doi.org/10.1080/19336934.2017.1394558

Montllor, C. B., Maxmen, A., & Purcell, A. H. (2002). Facultative bacterial endosymbionts

benefit pea aphids *Acyrthosiphon pisum* under heat stress. *Ecological Entomology*,

27(2), 189–195. https://doi.org/10.1046/j.1365-2311.2002.00393.x

Moore, M. P., Whiteman, H. H., & Martin, R. A. (2019). A mother's legacy: The strength of

maternal effects in animal populations. *Ecology Letters*, *22*(10), 1620–1628.

https://doi.org/10.1111/ele.13351

Moreira, D. R., de Souza, T. H. S., Galhardo, D., Puentes, S. M. D., Figueira, C. L., Silva, B.

G. da, Chagas, F. das, Gigliolli, A. A. S., de Toledo, V. de A. A., & Ruvolo-

Takasusuki, M. C. C. (2022). Imidacloprid induces histopathological damage in the

midgut, ovary, and spermathecal stored spermatozoa of queens after chronic colony

exposure. *Environmental Toxicology and Chemistry*, *41*(7), 1637–1648.

Moreira, S., Pereira, S. C., Seco-Rovira, V., Oliveira, P. F., Alves, M. G., & Pereira, M. de L.

(2021). Pesticides and male fertility: A dangerous crosstalk. *Metabolites*, *11*(12), 799.

Morgan, H. L., & Watkins, A. J. (2019). Transgenerational Impact of Environmental Change.

In P. Comizzoli, J. L. Brown, & W. V. Holt (Eds.), *Reproductive Sciences in Animal*

Conservation (Vol. 1200, pp. 71–89). Springer International Publishing.

https://doi.org/10.1007/978-3-030-23633-5_4

Mousseau, T. A., Uller, T., Wapstra, E., & Badyaev, A. V. (2009). Evolution of maternal

effects: Past and present. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *364*(1520), 1035–1038. https://doi.org/10.1098/rstb.2008.0303

Munday, P. L. (2014). Transgenerational acclimation of fishes to climate change and ocean

acidification. *F1000prime Reports*, *6*.

Navarro-Ortega, A., Acuña, V., Bellin, A., Burek, P., Cassiani, G., Choukr-Allah, R.,

Dolédec, S., Elosegi, A., Ferrari, F., Ginebreda, A., Grathwohl, P., Jones, C., Rault, P.

K., Kok, K., Koundouri, P., Ludwig, R. P., Merz, R., Milacic, R., Muñoz, I., …

Barceló, D. (2015). Managing the effects of multiple stressors on aquatic ecosystems

under water scarcity. The GLOBAQUA project. *Science of The Total Environment*,

503–504, 3–9. https://doi.org/10.1016/j.scitotenv.2014.06.081

Neven, L. G. (2000). Physiological responses of insects to heat. *Postharvest Biology and*

Technology, *21*(1), 103–111.

Newell, P. D., & Douglas, A. E. (2014). Interspecies interactions determine the impact of the

gut microbiota on nutrient allocation in *Drosophila melanogaster*. *Applied and*

Environmental Microbiology, *80*(2), 788–796.

Noecker, C., Chiu, H.-C., McNally, C. P., & Borenstein, E. (2019). Defining and evaluating

microbial contributions to metabolite variation in microbiome-metabolome

association studies. *MSystems*, *4*(6), 10–1128.

Noyes, P. D., & Lema, S. C. (2015a). Forecasting the impacts of chemical pollution and

climate change interactions on the health of wildlife. *Current Zoology*, *61*(4), 669–

689.

Noyes, P. D., & Lema, S. C. (2015b). Forecasting the impacts of chemical pollution and

climate change interactions on the health of wildlife. *Current Zoology*, *61*(4), 669–

689.

Noyes, P. D., McElwee, M. K., Miller, H. D., Clark, B. W., Van Tiem, L. A., Walcott, K. C.,

Erwin, K. N., & Levin, E. D. (2009). The toxicology of climate change:

Environmental contaminants in a warming world. *Environment International*, *35*(6),

971–986.

Obadia, B., Güvener, Z. T., Zhang, V., Ceja-Navarro, J. A., Brodie, E. L., William, W. J., &

Ludington, W. B. (2017). Probabilistic invasion underlies natural gut microbiome

stability. *Current Biology*, *27*(13), 1999–2006.

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'hara, R.,

Solymos, P., Stevens, M., & Szoecs, E. (2023). *Vegan: Community Ecology Package.*

R package version 2.6-4. 2022.

Osborne, B., Siboni, N., Seymour, J. R., Ralph, P., & Pernice, M. (2023). Exploring the

potential of algae-bacteria interactions in the biocontrol of the marine pathogen *Vibrio*

parahaemolyticus. Journal of Applied Phycology, *35*(6), 2731–2743.

https://doi.org/10.1007/s10811-023-03071-3

Overgaard, J., Malmendal, A., Sørensen, J. G., Bundy, J. G., Loeschcke, V., Nielsen, N. C.,

& Holmstrup, M. (2007). Metabolomic profiling of rapid cold hardening and cold

shock in *Drosophila melanogaster. Journal of Insect Physiology*, *53*(12), 1218–1232.

Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock, C. C., &

Thomas, M. B. (2013). Temperature variation makes ectotherms more sensitive to

climate change. *Global Change Biology*, *19*(8), 2373–2380.

https://doi.org/10.1111/gcb.12240

Pang, R., Chen, M., Yue, L., Xing, K., Li, T., Kang, K., Liang, Z., Yuan, L., & Zhang, W.

(2018). A distinct strain of Arsenophonus symbiont decreases insecticide resistance in

its insect host. *PLoS Genetics*, *14*(10), e1007725.

Parker, G. A., & Pizzari, T. (2010). Sperm competition and ejaculate economics. *Biological*

Reviews, *85*(4), 897–934. https://doi.org/10.1111/j.1469-185X.2010.00140.x

Perry, T., Heckel, D. G., McKenzie, J. A., & Batterham, P. (2008). Mutations in Dα1 or Dβ2

nicotinic acetylcholine receptor subunits can confer resistance to neonicotinoids in

Drosophila melanogaster. *Insect Biochemistry and Molecular Biology*, *38*(5), 520–

528.

Piggott, J. J., Niyogi, D. K., Townsend, C. R., & Matthaei, C. D. (2015). Multiple stressors

and stream ecosystem functioning: Climate warming and agricultural stressors

interact to affect processing of organic matter. *Journal of Applied Ecology*, *52*(5),

1126–1134. https://doi.org/10.1111/1365-2664.12480

Piggott, J. J., Townsend, C. R., & Matthaei, C. D. (2015). Climate warming and agricultural

stressors interact to determine stream macroinvertebrate community dynamics. *Global*

Change Biology, *21*(5), 1887–1906. https://doi.org/10.1111/gcb.12861

Pilakouta, N., & Baillet, A. (2022). Effects of temperature on mating behaviour and mating

success: A meta-analysis. *Journal of Animal Ecology*, *91*(8), 1642–1650.

Pizzari, T., & Parker, G. A. (2009). Sperm competition and sperm phenotype. In *Sperm*

Biology (pp. 207–245). Elsevier. https://doi.org/10.1016/B978-0-12-372568-4.00006-

9

Pölkki, M., Kangassalo, K., & Rantala, M. J. (2012). Transgenerational effects of heavy

metal pollution on immune defense of the blow fly *Protophormia terraenovae*. *PLoS*

One, *7*(6), e38832.

Pörtner, H.-O., Bock, C., & Mark, F. C. (2017). Oxygen-and capacity-limited thermal

tolerance: Bridging ecology and physiology. *Journal of Experimental Biology*,

220(15), 2685–2696.

Przeslawski, R., Byrne, M., & Mellin, C. (2015). A review and meta-analysis of the effects of

multiple abiotic stressors on marine embryos and larvae. *Global Change Biology*,

21(6), 2122–2140. https://doi.org/10.1111/gcb.12833

Reátegui-Zirena, E. G., Fidder, B. N., Olson, A. D., Dawson, D. E., Bilbo, T. R., & Salice, C.

J. (2017). Transgenerational endpoints provide increased sensitivity and insight into

multigenerational responses of *Lymnaea stagnalis* exposed to cadmium.

Environmental Pollution, *224*, 572–580.

Renoz, F., Pons, I., & Hance, T. (2019). Evolutionary responses of mutualistic insect–

bacterial symbioses in a world of fluctuating temperatures. *Current Opinion in Insect*

Science, *35*, 20–26. https://doi.org/10.1016/j.cois.2019.06.006

Ricupero, M., Abbes, K., Haddi, K., Kurtulus, A., Desneux, N., Russo, A., Siscaro, G.,

Biondi, A., & Zappalà, L. (2020). Combined thermal and insecticidal stresses on the

generalist predator *Macrolophus pygmaeus*. *Science of the Total Environment*, *729*,

138922.

Ridley, E. V., Wong, A. C., & Douglas, A. E. (2013). Microbe-dependent and nonspecific

effects of procedures to eliminate the resident microbiota from *Drosophila*

melanogaster. *Applied and Environmental Microbiology*, *79*(10), 3209–3214.

Ridley, E. V., Wong, A. C., Westmiller, S., & Douglas, A. E. (2012). Impact of the resident

microbiota on the nutritional phenotype of *Drosophila melanogaster*. *PloS One*, *7*(5),

e36765.

Roy, P., Phukan, P. K., Changmai, D., & Boruah, S. (2017). Pesticides, insecticides and male

infertility. *Int. J. Reprod. Contracept. Obs. Gynecol*, *6*, 3387–3391.

Russell, J. A., & Moran, N. A. (2006). Costs and benefits of symbiont infection in aphids:

Variation among symbionts and across temperatures. *Proceedings. Biological*

Sciences, *273*(1586), 603–610. https://doi.org/10.1098/rspb.2005.3348

Russell, R. J., Scott, C., Jackson, C. J., Pandey, R., Pandey, G., Taylor, M. C., Coppin, C. W.,

Liu, J., & Oakeshott, J. G. (2011). The evolution of new enzyme function: Lessons

from xenobiotic metabolizing bacteria versus insecticide-resistant insects.

Evolutionary Applications, *4*(2), 225–248.

Sales, K., Thomas, P., Gage, M. J., & Vasudeva, R. (2024). Experimental heatwaves reduce

the effectiveness of ejaculates at occupying female reproductive tracts in a model

insect. *Royal Society Open Science*, *11*(5), 231949.

Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, Ł.,

Hebberecht, L., Thomas, P., Franco, A., & Gage, M. J. G. (2018b). Experimental

heatwaves compromise sperm function and cause transgenerational damage in a

model insect. *Nature Communications*, *9*(1), 4771. https://doi.org/10.1038/s41467-

018-07273-z

Schultz, C. L., Wamucho, A., Tsyusko, O. V., Unrine, J. M., Crossley, A., Svendsen, C., &

Spurgeon, D. J. (2016). Multigenerational exposure to silver ions and silver

nanoparticles reveals heightened sensitivity and epigenetic memory in *Caenorhabditis*

elegans. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1832),

20152911.

Schwasinger-schmidt, T. E., Kachman, S. D., & Harshman, L. G. (2012). Evolution of starvation resistance in *Drosophila melanogaster*: Measurement of direct and correlated responses to artificial selection. *Journal of Evolutionary Biology*, *25*(2), 378–387.

Searle, S. R., Speed, F. M., & Milliken, G. A. (1980). Population Marginal Means in the Linear Model: An Alternative to Least Squares Means. *The American Statistician*, *34*(4), 216–221. https://doi.org/10.1080/00031305.1980.10483031

Segner, H., Schmitt-Jansen, M., & Sabater, S. (2014). Assessing the Impact of Multiple Stressors on Aquatic Biota: The Receptor's Side Matters. *Environmental Science & Technology*, *48*(14), 7690–7696. https://doi.org/10.1021/es405082t

Semsar-Kazerouni, M., Boerrigter, J. G., & Verberk, W. C. (2020). Changes in heat stress tolerance in a freshwater amphipod following starvation: The role of oxygen availability, metabolic rate, heat shock proteins and energy reserves. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *245*, 110697.

Sepulveda, J., & Moeller, A. H. (2020b). The Effects of Temperature on Animal Gut Microbiomes. *Frontiers in Microbiology*, *11*. https://doi.org/10.3389/fmicb.2020.00384

Shahid, N., Siddique, A., & Liess, M. (2024). Synergistic interaction between a toxicant and food stress is further exacerbated by temperature. *Environmental Pollution*, *363*, 125109.

Shama, L. N., Strobel, A., Mark, F. C., & Wegner, K. M. (2014). Transgenerational plasticity in marine sticklebacks: Maternal effects mediate impacts of a warming ocean. *Functional Ecology*, *28*(6), 1482–1493.

1074 Shokal, U., Yadav, S., Atri, J., Accetta, J., Kenney, E., Banks, K., Katakam, A., Jaenike, J., &

1075   Eleftherianos, I. (2016). Effects of co-occurring Wolbachia and Spiroplasma

1076   endosymbionts on the *Drosophila* immune response against insect pathogenic and

1077   non-pathogenic bacteria. *BMC Microbiology*, *16*(1), 16.

1078   https://doi.org/10.1186/s12866-016-0634-6

1079 Silva, A. P. N., Andrade, E. S., Nascimento, V. L., & Haddi, K. (2025). Thermal modulation

1080   of insecticide-induced hormetic and oxidative responses in insect pests. *Chemosphere*,

1081   *370*, 143920.

1082 Silva, A. S. J., Kristiansen, S. M., Sengupta, S., van Gestel, C. A., Leinaas, H. P., & Borgå,

1083   K. (2023). Using dietary exposure to determine sub-lethal effects from imidacloprid

1084   in two springtail (*Collembola*) species. *Ecotoxicology*, *32*(10), 1209–1220.

1085 Simmons, L. W., & Fitzpatrick, J. L. (2012). Sperm wars and the evolution of male fertility.

1086   *Reproduction*, *144*(5), 519–534. https://doi.org/10.1530/REP-12-0285

1087 Siviter, H., Bailes, E. J., Martin, C. D., Oliver, T. R., Koricheva, J., Leadbeater, E., & Brown,

1088   M. J. (2021). Agrochemicals interact synergistically to increase bee mortality. *Nature*,

1089   *596*(7872), 389–392.

1090 Sokolova, I. M. (2013). Energy-limited tolerance to stress as a conceptual framework to

1091   integrate the effects of multiple stressors. *Integrative and Comparative Biology*,

1092   *53*(4), 597–608.

1093 Sørensen, J. G., Kristensen, T. N., & Loeschcke, V. (2003). The evolutionary and ecological

1094   role of heat shock proteins. *Ecology Letters*, *6*(11), 1025–1037.

1095 Stehle, S., & Schulz, R. (2015). Agricultural insecticides threaten surface waters at the global

1096   scale. *Proceedings of the National Academy of Sciences*, *112*(18), 5750–5755.

Swelam, B. A., Verswijveren, S. J., Salmon, J., Arundell, L., & Ridgers, N. D. (2022).

Exploring activity compensation amongst youth and adults: A systematic review.

International Journal of Behavioral Nutrition and Physical Activity, *19*(1), 25.

Tran, T. T., Janssens, L., Dinh, K. V., & Stoks, R. (2018). Transgenerational interactions

between pesticide exposure and warming in a vector mosquito. *Evolutionary*

Applications, *11*(6), 906–917.

Uller, T., Nakagawa, S., & English, S. (2013). Weak evidence for anticipatory parental

effects in plants and animals. *Journal of Evolutionary Biology*, *26*(10), 2161–2170.

https://doi.org/10.1111/jeb.12212

Verberk, W. C. E. P., Overgaard, J., Ern, R., Bayley, M., Wang, T., Boardman, L., &

Terblanche, J. S. (2016). Does oxygen limit thermal tolerance in arthropods? A

critical review of current evidence. *Comparative Biochemistry and Physiology Part*

A: Molecular & Integrative Physiology, *192*, 64–78.

https://doi.org/10.1016/j.cbpa.2015.10.020

Verheyen, J., & Stoks, R. (2019). Current and future daily temperature fluctuations make a

pesticide more toxic: Contrasting effects on life history and physiology.

Environmental Pollution, *248*, 209–218.

Verheyen, J., & Stoks, R. (2023). Thermal performance curves in a polluted world: Too cold

and too hot temperatures synergistically increase pesticide toxicity. *Environmental*

Science & Technology, *57*(8), 3270–3279.

Walsh, B. S., Parratt, S. R., Atkinson, D., Snook, R. R., Bretman, A., & Price, T. A. (2019).

Integrated approaches to studying male and female thermal fertility limits. *Trends in*

Ecology & Evolution, *34*(6), 492–493.

Walsh, B. S., Parratt, S. R., Hoffmann, A. A., Atkinson, D., Snook, R. R., Bretman, A., &

Price, T. A. R. (2019). The impact of climate change on fertility. *Trends in Ecology &*

Evolution, *34*(3), 249–259. https://doi.org/10.1016/j.tree.2018.12.002

Walsh, M. R., Whittington, D., & Funkhouser, C. (2014). Thermal transgenerational

plasticity in natural populations of *Daphnia*.

Wan, N.-F., Fu, L., Dainese, M., Kiær, L. P., Hu, Y.-Q., Xin, F., Goulson, D., Woodcock, B.

A., Vanbergen, A. J., & Spurgeon, D. J. (2025). Pesticides have negative effects on

non-target organisms. *Nature Communications*, *16*(1), 1360.

Wang, X., Ma, D., Jin, Q., Deng, S., Stančin, H., Tan, H., & Mikulčić, H. (2019). Synergistic

effects of biomass and polyurethane co-pyrolysis on the yield, reactivity, and heating

value of biochar at high temperatures. *Fuel Processing Technology*, *194*, 106127.

Wang, Y.-C., Chang, Y.-W., Bai, J., Zhang, X.-X., Iqbal, J., Lu, M.-X., Gong, W.-R., & Du,

Y.-Z. (2021). Temperature affects the tolerance of *Liriomyza trifolii* to insecticide

abamectin. *Ecotoxicology and Environmental Safety*, *218*, 112307.

Wang, Y.-C., Chang, Y.-W., Bai, J., Zhang, X.-X., Iqbal, J., Lu, M.-X., Hu, J., & Du, Y.-Z.

(2021a). High temperature stress induces expression of CYP450 genes and

contributes to insecticide tolerance in *Liriomyza trifolii*. *Pesticide Biochemistry and*

Physiology, *174*, 104826.

Wang, Y.-C., Chang, Y.-W., Bai, J., Zhang, X.-X., Iqbal, J., Lu, M.-X., Hu, J., & Du, Y.-Z.

(2021b). High temperature stress induces expression of CYP450 genes and

contributes to insecticide tolerance in *Liriomyza trifolii*. *Pesticide Biochemistry and*

Physiology, *174*, 104826.

Xiao, D., Zhao, J., Guo, X., Chen, H., Qu, M., Zhai, W., Desneux, N., Biondi, A., Zhang, F.,

& Wang, S. (2016). Sublethal effects of imidacloprid on the predatory seven-spot

ladybird beetle *Coccinella septempunctata*. *Ecotoxicology*, *25*, 1782–1793.

Yu, C.-W., & Liao, V. H.-C. (2016). Transgenerational Reproductive Effects of Arsenite Are

Associated with H3K4 Dimethylation and SPR-5 Downregulation in *Caenorhabditis*

elegans. *Environmental Science & Technology*, *50*(19), 10673–10681.

https://doi.org/10.1021/acs.est.6b02173

Zelezniak, A., Andrejev, S., Ponomarova, O., Mende, D. R., Bork, P., & Patil, K. R. (2015).

Metabolic dependencies drive species co-occurrence in diverse microbial

communities. *Proceedings of the National Academy of Sciences*, *112*(20), 6449–6454.

Zhang, X., Yu, X., & Chen, J. (2008). High temperature effects on yeast-like endosymbiotes

and pesticide resistance of the small brown planthopper, *Laodelphax striatellus*. *Rice*

Science, *15*(4), 326–330.

Zhang, Y., Cai, T., Ren, Z., Liu, Y., Yuan, M., Cai, Y., Yu, C., Shu, R., He, S., & Li, J.

(2021). Decline in symbiont-dependent host detoxification metabolism contributes to

increased insecticide susceptibility of insects under high temperature. *The ISME*

Journal, *15*(12), 3693–3703.

Zhang, Y., Xu, G., Jiang, Y., Ma, C., & Yang, G. (2021). Sublethal effects of Imidacloprid on

fecundity, apoptosis and virus transmission in the small brown planthopper

Laodelphax striatellus. *Insects*, *12*(12), 1131.

Zhou, C., Liu, L., Yang, H., Wang, Z., Long, G., & Jin, D. (2017). Sublethal effects of

imidacloprid on the development, reproduction, and susceptibility of the white-backed

planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae). *Journal of Asia-Pacific*

Entomology, *20*(3), 996–1000.

Zwoinska, M. K., Rodrigues, L. R., Slate, J., & Snook, R. R. (2020). Phenotypic responses to

and genetic architecture of sterility following exposure to sub-lethal temperature

during development. *Frontiers in Genetics*, *11*, 573.

https://doi.org/10.3389/fgene.2020.00573

1170

1171

Figure 1 Proportion of eggs fertilized by males under varying conditions of thermal treatments

(low: 24.5°C; high: 36°C), microbiome type (Axenic, *Acetobacter tropicalis*, *Lactobacillus*

brevis, Control), and pesticide exposure (present and absent). Each colour represents a different

microbiome treatment and two different panels correspond to presence/absence of pesticide

treatment with two temperature treatments within each panel. Each circle represents the mean

of all the replicates in that treatment combination with error bars corresponding to 95%

confidence intervals.

1179

Figure 2 Mean fecundity (±95% CI) of females mated to males exposed to different combinations of two thermal treatments (low: 24.5°C; high: 36°C), microbiome type (Axenic, *Acetobacter tropicalis*, *Lactobacillus brevis*, and Control), and pesticide exposure (No pesticide, Imidacloprid). Each circle represents the mean of replicates per treatment group. The left and right panels correspond to pesticide absence and presence, respectively, with temperature conditions shown on the x-axis. Colours represent different microbiome treatments.

1187

1188

1189

Figure 3 Proportion of progenies sired by the focal males (2nd in order to mate with the female)

under varying conditions of thermal treatments (low: 24.5°C; high: 36°C), microbiome type

(Axenic, *Acetobacter tropicalis*, *Lactobacillus brevis*, Control), and pesticide exposure

(present and absent). Each colour represents a different microbiome treatment and two different

panels correspond to presence/absence of pesticide treatment with two temperature conditions

(low/high) within each panel. Each circle represents the mean of all replicates in that treatment

combination with error bars corresponding to 95% confidence intervals.

1197

1198

1199

1200

1201

1202

1203

Figure 4 Bacterial abundance/colony forming units of different microbial treatment

(*Acetobacter tropicalis*, *Lactobacillus brevis*, Control) in response to different conditions of

thermal treatments (low: 24.5°C; high: 36°C), and pesticide exposure (present and absent).

Each colour represents a different microbiome treatment and two different panels correspond

to presence/absence of pesticide treatment with two temperature conditions (low/high) within

each panel. Each circle represents the mean of all the replicates in that treatment combination

with error bars corresponding to 95% confidence intervals.

1211

1212

1213

1214

1215

Figure 5. Non-metric multidimensional scaling (NMDS) ordination (Bray-Curtis distance, k =

4, stress = 0.046) of microbial community composition across temperature and pesticide

treatments. Each point represents one male, colored by temperature treatment (red = Low, blue

= High) and shaped by pesticide exposure (circle = Absent, triangle = Present). Shaded convex

hulls denote 95% confidence areas for each temperature × pesticide group. Arrows represent

the direction and strength of species (Hellinger-transformed) most strongly correlated with the

ordination space, as identified by envfit. Species abbreviations: Ao = *Acetobacter orientalis*,

Ap = *A. pasteuriensis*, At = *A. tropicalis*, Lb = *Lactobacillus brevis*, Lp = *L. plantarum*.

1224

1225

1226

1227 **Figure 6:** Proportion of acetic acid bacteria in the five-species consortium in response to heat

1228 and pesticide stress. Each circle represents the mean of all replicates in that treatment

1229 combination with error bars corresponding to 95% confidence intervals.

1230

Figure 7. Proportion of eggs hatched out of total number of eggs fertilized by male offspring produced by fathers under different conditions of thermal treatments (low: 24.5°C; high: 36°C), microbiome type (Axenic, *Acetobacter tropicalis*, *Lactobacillus brevis*, Control), and pesticide presence (present and absent). Each colour represents a different F_0 microbiome treatment and two different panels correspond to presence/absence of F_0 pesticide treatment with two F_0 temperature conditions(low/high) within each panel. Each circle represents the mean of all the replicates in that treatment combination with error bars corresponding to 95% confidence intervals.

1239

1240

1241

1242

1243

1244

1245

1246

1247

Figure 8. Proportion of hatched eggs out of total number of eggs laid by female offsprings

produced by fathers under varying conditions of thermal treatments (low: 24.5°C; high: 36°C),

microbiome type (Axenic, *Acetobacter tropicalis*, *Lactobacillus brevis*, Control), and pesticide

presence (present and absent). Each colour represents a different F_0 microbiome treatment and

two different panels correspond to presence/absence of F_0 pesticide treatment with two F_0

temperature conditions(low/high) within each panel. Each circle represents the mean of all the

replicates in that treatment combination with error bars corresponding to 95% confidence

intervals.

1256

1257

Figure 9. Total number of fertilised eggs laid by females mated to male offsprings produced

by fathers under varying conditions of thermal treatments (low: 24.5°C; high: 36°C),

microbiome type (Axenic, *Acetobacter tropicalis*, *Lactobacillus brevis*, Control), and pesticide

presence (present and absent). Each colour represents a different F_0 microbiome treatment and

two different panels correspond to presence/absence of F_0 pesticide treatment with two F_0

temperature conditions (low/high) within each panel. Each circle represents the mean of all the

replicates in that treatment combination with error bars corresponding to 95% confidence

intervals.

Figure 10. Total number of fertilised eggs laid by female offsprings produced by fathers

under varying conditions of thermal treatments (low: 24.5°C; high: 36°C), microbiome type

(Axenic, *Acetobacter tropicalis*, *Lactobacillus brevis*, Control), and pesticide presence

(present and absent). Each colour represents a different F_0 microbiome treatment and two

different panels correspond to presence/absence of F_0 pesticide treatment with two F_0

temperature conditions (low/high) within each panel. Each circle represents the mean of all

the replicates in that treatment combination with error bars corresponding to 95% confidence

intervals.