1

Variation in responses to temperature across admixed genotypes of *Populus trichocarpa* × *P. balsamifera* predict geographic shifts in regions where hybrids are favored

Alayna Mead*, Joie R. Beasley-Bennett, Andrew Bleich, Dylan Fischer, Shelby Flint, Julie Golightly, Sara K. Klopf, Mason W. Kulbaba, Jesse R. Lasky, Jared M. LeBoldus, David B. Lowry, Nora Mitchell, Emily Moran, Jason P. Sexton, Kelsey L. Søndreli, Baxter Worthing, Michelle Zavala-Paez, Matthew C. Fitzpatrick, Jason Holliday, Stephen Keller, Jill A. Hamilton

*Corresponding author: alaynamead@psu.edu

Abstract

In a rapidly changing environment, predicting changes in the growth and survival of local populations can inform conservation and management. Plastic responses vary as a result of genetic differentiation within and among species, so accurate rangewide predictions require characterization of genotype-specific reaction norms across the continuum of historic and future climate conditions comprising a species' range. Natural hybrid zones can give rise to novel recombinant genotypes associated with high phenotypic variability, further increasing the variance of plastic responses within the ranges of the hybridizing species. Experiments that plant replicated genotypes across a range of environments can characterize genotype-specific reaction norms; identify genetic, geographic, and climatic factors affecting variation in climate responses; and make predictions of climate responses across complex genetic and geographic landscapes. The North American hybrid zone of *Populus trichocarpa* and *P. balsamifera* represents a natural system in which reaction norms are likely to vary with underlying genetic variation that has been shaped by climate, geography, and introgression. Here, we leverage a dataset containing 45 clonal genotypes of varying ancestry from this natural hybrid zone, planted across 17 replicated common garden experiments spanning a broad climatic range, including sites warmer than the natural species ranges. Growth and mortality were measured over two years, enabling us to model reaction norms for each genotype across these tested environments. Genomic variation associated with species ancestry and northern/southern regions significantly influenced growth across environments, with genotypic variation in reaction norms reflecting a trade-off between cold tolerance and growth. Using modeled reaction norms for each genotype, we predicted that genotypes with more *P. trichocarpa* ancestry may gain an advantage under warmer climates. Spatial shifts of the hybrid zone could facilitate the spread of beneficial alleles into novel climates. These results highlight that genotypic variation in responses to temperature will have landscape-level effects.

Introduction

Predicting the phenotypic responses of populations to changing climates and how they vary across
species ranges is essential for conserving and restoring native ecosystems. Predicted changes in fitness
between current and future climates can be used to identify populations most at risk, or to identify
populations resilient to change which could be valuable seed sources for assisted gene flow and
restoration (Aitken and Whitlock 2013; Aitken and Bemmels 2016). Phenotypes expressed in the field
arise from the genotypic variation underlying phenotypic traits (G), the plastic response to the
environment (E), and genotypic variation in response to the environment (GxE) (Des Marais et al.
2013). Each of these factors vary across complex and changing landscapes. By characterizing reaction
norms, the set of phenotypes expressed across a range of environments, it is possible to predict
organisms' responses to the variable environments they inhabit and understand the selective forces that
may be shaping plasticity (Via et al. 1995; Arnold et al. 2019). Reaction norms can vary within species,
resulting in part from selection imposed by spatially varying climates (Des Marais et al. 2013; Ikeda et
al. 2017; Rehfeldt et al. 2018; Patsiou et al. 2020). Predicting genotype-specific responses will be
particularly important for species with large ranges, which may exhibit variation in reaction norms due
to the combined influence of geography, population demography, and climate (Cooper et al. 2019,
2022; Van Nuland et al. 2020). This is particularly true where glacial refugia have shaped species'
demographic history and connectivity (Woolbright et al. 2014; Love et al. 2023; Bolte et al. 2024).
However, because of the difficulty of phenotyping multiple genotypes across many environments,
many methods for modeling a species' response to climate change ignore genotypic variation, and
instead consider only the climate envelope of the extant species range (Capblancq et al. 2020).
Furthermore, regions within a species' range will vary in the rate, magnitude, and nature of climate
change (e.g, increases or decreases in precipitation), making it important to understand localized
climate responses. To accurately predict local phenotypic responses to changing climates, it is
necessary to characterize genotype-specific reaction norms across the continuum of historic and future
climate conditions comprising a species' range.

Common garden experiments and provenance trials are invaluable tools for quantifying intraspecific
variation in phenotypic plasticity and for predicting phenotypic responses to current and future
environments (O'Neill et al. 2008; Wang et al. 2010; Leites et al. 2012; Fischer et al. 2014a; Grady et al.
2015; Browne et al. 2019; Leites and Benito Garzón 2023; Ye et al. 2023; Hord et al. 2025). Capturing
variation in reaction norms varies across genetic and climatic gradients is critical and can be
accomplished by repeated planting of genotypes across multiple environments. Genotype-specific
reaction norms modeled across continuous environments can be used to identify genotypes or loci
associated with optimized fitness or yield across varying environmental conditions; however, such
studies have been limited to a handful of species (Gray et al. 2011; Arnold et al. 2019; Lowry et al.

2019; Patsiou et al. 2020; VanWallendael et al. 2022; Li et al. 2024). Because plasticity results from adaptation to climate as well as demographic history, it may be possible to predict the reaction norms of unmeasured genotypes if they vary as a function of climate of origin (O'Neill et al. 2008; Wang et al. 2010; Rehfeldt et al. 2018) or genetic variation. When genomic data are available in addition to phenotypic and climate data collected in common garden experiments, predictions of phenotypic responses can be improved (Mahony et al. 2020; Archambeau et al. 2022; Putra et al. 2023; Li et al. 2024). Therefore, incorporating genetic information into characterization of reaction norms will enable more accurate predictions of rangewide climate responses.

Natural hybrid zones are ideal systems for disentangling the effects of adaptation and demographic history on phenotypes, including reaction norms, because multiple generations of backcrossing can result in novel recombinant genotypes associated with high phenotypic variability (Janes and Hamilton 2017). Furthermore, hybrid zones can increase the genetic variation available for adaptation to rapidly changing climates and allow movement of adaptive loci from one species into another via introgression (Janes and Hamilton 2017; Suarez-Gonzalez et al. 2018b; Kremer and Hipp 2020; Buck et al. 2023; Hord et al. 2025). Comparing responses to the environment across admixed genotypes can inform how existing genetic variation within hybrid zones underlies fitness differences across environments. Hybrid zones can also be used to monitor responses to climate change (Taylor et al. 2015), with documented geographic shifts in some zones in response to changing environments (Billerman et al. 2016; Wielstra 2019; Alexander et al. 2022). If reaction norms depart from the intraspecific pattern across a hybrid zone, climate change may alter regions where particular species, genotypes, or even genes are favored (Hord et al. 2025).

North American *Populus* is a model system for forest trees due to their small genomes, ease of clonal propagation, and development as a biofuel feedstock (Jansson and Douglas 2007; Sannigrahi et al. 2010; Porth and El-Kassaby 2015), yet like many non-model trees, its genetic variation is shaped by interactions between climate, geography, and interspecific introgression. *Populus* species occupy heterogeneous landscapes, often forming multi-species hybrid zones that are a source of novel recombinant genetic variation (Suarez-Gonzalez et al. 2016, 2018a; Chhatre et al. 2018; Bolte et al. 2024). These factors make *Populus* an ideal system for quantifying intraspecific variation in climate responses and predicting landscape-scale changes in fitness. Here, we focus on the North American hybrid zone between *P. trichocarpa*, a western species spanning latitudes from Alaska to California, and *P. balsamifera*, which occurs transcontinentally throughout the boreal regions of the contiguous United States, Canada and Alaska (Figure 1A). Populations within the hybrid zone exhibit genetic and phenotypic differentiation, likely resulting from varying demographic history as well as adaptation to wide-ranging biotic and abiotic environments (Keller et al. 2010, 2011, 2012; Slavov et al. 2012; Evans et al. 2014; Geraldes et al. 2014; McKown et al. 2014a, c,b; Zhou et al. 2014; Fitzpatrick and Keller

2015; Holliday et al. 2016; Chhetri et al. 2019; Zhang et al. 2019; Fitzpatrick et al. 2021). Furthermore,

rates of gene flow across the hybrid zone vary in part due to geographic barriers, resulting in spatial

differences in population differentiation (Bolte et al. 2024).

We leveraged the *P. trichocarpa* × *P. balsamifera* hybrid zone to determine how genetic and

environmental parameters interact to influence genotype-specific reaction norms, ultimately using our

predictions to model changes to hybrid zone composition in the context of global change. We used a

series of replicated provenance trials of clonal genotypes of *P. trichocarpa*, *P. balsamifera*, and their

hybrids, which we planted in 17 gardens across the United States that span a wide range of

environments, including many warmer than the climate of origin. We tested three hypotheses: 1)

Populus hybrids display heritable differences in adaptation to climate, which manifest as genotype-

specific responses to climatic variation. We measured fitness-related traits in each garden to determine

how genetic background and the environment interact to determine fitness in this system, quantifying

the effect of genetic structure (genotype effect, G), climate (environmental effect, E) and how the

response to climate varies by genotype (G×E). A significant genotype × environment interaction could

suggest that the response to climate is mediated by local adaptation to the climate of origin. 2)

Variation in reaction norms are predictable based on a continuous gradient of genomic ancestry,

enabling us to predict where the hybrid zone is favored to move under future climates. We used whole-

genome sequence data to characterize multivariate genetic structure, including both species ancestry

and intraspecific variation, and predicted its effect on phenotypic responses among genotypes. Finally,

we used genotype-specific reaction norms to predict phenotypic responses to climate change across the

landscape of the *P. trichocarpa* x *P. balsamifera* hybrid zone. 3) Inclusion of genetic information in

phenotypic response models improves their predictions due to inherent genetic structure contributing

to variation in reaction norms. We tested whether climate of origin can serve as a proxy for adaptive

genetic structure, circumventing the need for genetic data. Using this experiment, we show that

increased temperatures could shift where genotypes with different levels of admixture are favored,

illustrating the importance of combining genomic data with common garden experiments to predict

rangewide responses to temperature.

Methods

Field collections and propagation

In 2020 we established 17 common gardens containing clonal replicates of 48 poplar genotypes,

including *P. trichocarpa*, *P. balsamifera*, and admixed individuals. Genotypes were established from

dormant branches collected between October 2019 and March 2020 from natural populations

spanning five transects that traverse the natural hybrid zone of *P. trichocarpa* and *P. balsamifera* and

span latitudes from Alaska to southern British Columbia and Alberta (Figure 1). Dormant vegetative
cuttings were transported to Virginia Tech (Critz, VA, USA) for propagation, as described in Bolte et
al. (2024). Briefly, cuttings were exposed to a 30-second ZeroTol 2.0 fungal dip treatment, dipped in
Garden Safe Take Root hormone (0.1% indole-3-butyric acid), rooted into a standard rooting mix, and
placed on a mist bench for 38 days following which clonal genotypes were shipped and planted into
common garden sites.

Common garden design

Seventeen common gardens were established at colleges, universities, and arboreta across the United
States in fall 2020, including: The Evergreen State College, WA; University of Idaho; Lockerly
Arboretum, GA; Morton Arboretum, IL; Michigan State University; North Dakota State University;
Northwest Missouri State Arboretum; Oregon State University; Our Lady of the Lake University, TX;
Pennsylvania State University; Salisbury University, MD; Southwest Minnesota State University;
University of California, Merced; Virginia Tech University; University of Wisconsin - Eau Claire;
Washington State University; and University of Wyoming (Figure 1A). Garden sites span a range of
environments and include the southern range of both *P. trichocarpa* and *P. balsamifera*, as well as
warmer regions to the south of their native ranges, enabling us to predict responses to novel climates
(Figure 1B). Each garden included two blocks, with each genotype replicated in each block. In total,
the design included 48 genotypes x 17 gardens x 2 blocks x 1 genotype/block. Each garden had a
minimum of 94 trees, for a total of 1656 individuals included across all gardens. Phenotypic data were
collected during the growing season for three years: 2021-2023. All 17 gardens were evaluated in the
first year, but high mortality and changing local phenotyping capacity at some sites impacted the total
number of gardens assessed, with fifteen and twelve sites evaluated in 2022 and 2023, respectively.

As part of the same experiment, three additional common gardens were established in March 2020.
These gardens included 544 genotypes, each with three replicated individuals planted in three blocks
(1 replicate per block), located at North Dakota State University (NDSU), Virginia Tech University
(VA), and the University of Vermont (VT). We designated the 17 smaller gardens as "mini" gardens,
and the three larger gardens as "maxi" gardens. All 48 genotypes included in the mini gardens were also
planted in the maxi gardens. In this study, our analysis focuses on the mini gardens, which allows us to
compare phenotypes and fitness metrics across a wide range of environments while controlling for
genetic effects by using replicated clonal genotypes; we use the maxi gardens for model evaluation (see
statistical methods below).

Phenotypic data

Height was measured each year (2021-2023) prior to bud flush and after budset. Using these data,
annual growth increment was calculated per year based on the height accumulated during the growing
season (pre-bud flush height subtracted from post-budset height). Some gardens experienced
herbivory, disease, and accidental mechanical damage leading to height measurements decreasing from
bud flush to bud set in some individuals. As these negative growth increments likely represent the

consequences of measurement error or herbivory, they were removed prior to analyses. Given the
frequency of herbivory, it is likely that some growth increment values are underestimates of potential
cumulative growth.

Climate data

Climate data were extracted from ClimateNA (Wang et al. 2016; AdaptWest Project 2022) for
genotype provenances (locations of origin) and common garden environments using terra 1.8-21
(Hijmans 2025) in R version 4.4.2 (R Core Team 2024). For provenances, we extracted historical
climate data for the 30-year period from 1961-1990. We focused on average historical climate data
because variation in climate response is partially the result of selection associated with climate of origin
over the lifespan of the collected adult trees. For garden climates, we extracted the average climate data
associated with each year of data collection, starting when the seedlings were planted in 2020. To
identify climatic variables contributing to variation in growth across gardens, we fit LOESS curves of
each climate variable against growth increment measurements using the `loess` function in base R.
We selected mean coldest month temperature (MCMT) for use in statistical modeling because it
corresponded to variation in performance across gardens and because the range of MCMT values at
the common garden sites encompassed climates for most natural populations (see Results), enabling us
to predict growth and mortality responses to MCMT within much of the native range of the two
poplar species. We did not use precipitation variables in statistical modeling because most gardens were
irrigated during the first year and some drier gardens (OLLU, SWMN, and UCM) continued
irrigation in later years to ensure survival, limiting the selective effect of natural precipitation variation
across gardens.

Genomic data

Whole-genome sequence data associated with each genotype and previously described in Bolte et al.
(2024) was used in phenotypic prediction across common gardens. DNA extraction, sequencing, and
variant filtration are described in Supplementary Methods and filtering scripts are available at
https://github.com/alaynamead/poplar_hybrid_vcf_filtering. A total of 334,657 variable sites were
used to characterize genetic variation and admixture. Bolte et al. (2024) identified three distinct
lineages in this hybrid zone: *P. balsamifera*, coastal *P. trichocarpa*, and a separate interior *P.
trichocarpa* lineage originating from an ancient admixture event between the two species. We
incorporated this genetic variation into our predictions using the PC scores of each genotype from a
PCA of the 334,657 SNPs created with vegan 2.6-8 in R (Oksanen et al. 2024), defining genetic
structure as variation in genomic PC space. PC1 separates *P. balsamifera* and the two *P. trichocarpa*
lineages, PC2 separates all three lineages, and PC3 separates the northern (Alaska and Cassiar) and
southern (Chilcotin, Jasper, and Crowsnest) transects (**Figure S4**). To visualize how responses varied
by species ancestry across the repeated contact zones, we assigned species ancestry for each genotype
using the proportions based on K=2 from the ancestry estimation software ADMIXTURE
(Alexander et al. 2009). Across 48 genotypes, three genotypes were genetic outliers, possibly as a result

of mixed ancestry with additional *Populus* species. These individuals were removed from the analyses,
leaving a total of 45 genotypes for statistical modeling.

Statistical model

We fit a linear mixed-effects model to evaluate the effects of garden climate, home climate, genetic
structure and their interactions (G×E) on growth across the common gardens and to predict responses
to future climates. We defined the environment as the MCMT of the common garden climate for the
year of measurement, and we defined home climate as the historical average of MCMT at the
provenance of each genotype. Genotypes may vary in their responses to climate as a result of local
adaptation to their climate of origin and as a result of neutral genetic variation. Given this, we
accounted for these sources of variation using different model variables. Previous studies have tested
how climate of origin determines response to the environment within a species, essentially using
climate of origin as a proxy for genotype to model G×E, assuming that much of the response is
explained by adaptation to the local environment of origin (O'Neill et al. 2008; Wang et al. 2010).
However, this approach does not account for variation in responses that result from other processes,
such as neutral evolution resulting from historic demographic processes, selection for recent but not
long-term climate patterns, or novel recombinants due to hybridization. We include genetic structure
and climate of origin in the model to account for these multiple interacting factors that may influence
the response to the environment.

We included data from the first two years of measurement (2021 and 2022) across 17 and 14 gardens,
respectively, analyzing 2768 measurements of 1610 individual trees. We excluded 2023 data because
gardens at the upper and lower temperature extremes were lost due to high mortality (Figure S1), and
climatic extremes are important to provide "anchor points" to produce biologically realistic response
curves (Wang et al. 2006). However, we did compare the model fit for each year individually and
observed similar effect directions across all three years (Supplementary Figure S10). Individuals with a
negative growth increment, attributed to herbivory or measurement error, were excluded from the
statistical modeling (137 individuals in 2021 and 97 in 2022).

We included survival and yearly growth increment as two measures of performance within the same
model by fitting a zero-inflated Gaussian model using glmmTMB 1.1.11 (Brooks et al. 2017) in R
version 4.5.0 (R Core Team 2024). We set the growth of all trees marked dead at the end of each
growing season as 0, resulting in a large number of zero values (381 trees in 2021 and 343 trees in 2022,
27% of the total). This zero-inflated model assumes that values of zero can have two origins:
"sampling" zeros that are part of the distribution of growth increment (i.e., some individuals survived
but had growth at or near zero) and "structural" zeros due to mortality (Hu et al. 2011; Brooks et al.
2017). This allowed us to evaluate how two separate fitness-associated traits, growth and mortality,
respond to MCMT. Together their combined effect can be considered a proxy for overall fitness,
incorporating both the probability of mortality and the predicted growth for surviving individuals.
We report phenotypic predictions for three model components: the zero-inflated model representing
mortality, the conditional model representing growth, and the overall model combining the two.

The full model included the fixed effects of garden MCMT, home MCMT, their square terms (to
account for the observed quadratic response associated with a fitness optimum, Figure S2 and S3), and
all interactions between garden and home MCMT and their square terms (indicating that the response
to garden temperature varies based on temperature of origin). We also included genetic information as
a fixed effect using the values for genetic PC1, PC2, and PC3 (Figure S4). MCMT was negatively
correlated with genetic PC1 (R = -0.69), consistent with the climate niches associated with the two
species, but was only weakly correlated with PC2 (R=-0.22) and PC3 (-0.40), so including multiple
PCs enables us to identify variation in responses resulting from neutral genetic structure or adaptation
to other climate variables. To test for G×E, we included interaction effects between garden climate and
its square term with all genetic PCs. We also included the random effects of individual, genotype, year,
and block nested within garden to account for non-independence of multiple observations of the same
individual, genotype and garden site. The same formula was used for the zero-inflated portion of the
model to test the effect of each factor on mortality. To improve model convergence, we scaled the
numeric variables using the R function `scale` without centering them and retained the scaling factor
used for each variable to back-transform scaled values to actual values. Growth increment was log-
transformed to account for a distribution skewed towards lower values. Using each individual as a
separate observation, we fit the following model in using the `glmmTMB` function, accounting for
home MCMT, garden MCMT and their interactions and square terms; as well as genetic PCs 1-3 and
their interaction terms with garden MCMT and its square term.

Model 1:

`log(growth increment + 1) ~ garden MCMT × home MCMT + garden MCMT`^{2}` × home`
`MCMT`^{2}`  + garden MCMT`^{2}` × home MCMT  + garden MCMT × home MCMT`^{2}` + genetic PC1 ×`
`garden MCMT + genetic PC2 × garden MCMT + genetic PC3 × garden MCMT + genetic`
`PC1 × garden MCMT`^{2}` + genetic PC2 × garden MCMT`^{2}` + genetic PC3 × garden MCMT`^{2}` +`
`(1 | genotype) + (1 | garden/block) + (1 | year) + (1 | individual)`

The random intercepts of genotype, block nested within garden, year, and individual tree were
included and are indicated by parentheses. The significance of each term was evaluated using the
`tab_model` function from the R package sjPlot 2.8.17 (Lüdecke 2024), which uses a type II Wald
chi square test.

Model evaluation

To evaluate whether the model can accurately predict poplar phenotypic response to environmental
conditions, we compared its predictions of growth and mortality to actual measures in two ways. First,
we removed one of the garden environments from the dataset, trained the model on the remaining 16
gardens, and used the fitted model to predict height in the garden that was excluded (hereafter referred
to as leave-one-out cross validation). Second, we trained the model using all 17 gardens and then
predicted growth increment of 544 genotypes in the three maxi gardens, two of which also have a mini
garden site that was included in the training data (Virginia and North Dakota) and one that does not
have a mini garden (University of Vermont). 500 of these genotypes were novel genotypes not

evaluated within the mini gardens. We predicted heights for individuals based on the full model using
the `predict` function from the glmmTMB package. To quantify the model's predictive ability for
each year, we calculated the Pearson correlation between the actual heights and the predicted heights
(conditional model) as well as the predicted heights when the probability of mortality was included
(overall model). To calculate predictive ability for the conditional model, we removed the individuals
with a measured height of zero (which is accounted for in the zero-inflated portion of the model). We
also evaluated model predictive ability with and without the random effects of garden, block, and
genotype. Including random effects (using option re.form=NULL in the predict function) accounts
for the varying intercepts associated with each group, while excluding them and setting all random
effects to zero (re.form = NA) makes predictions based on the fixed effects of climate and genetics
only. When the model was used to predict growth in genotypes or gardens not included in the training
set, a new random effect was predicted.

We also used leave-one-out cross-validation to compare the performance of three different models with
different combinations of genetic and climatic information. If most of the variation among genotypes
in response to garden environments as estimated by MCMT is explained by local adaptation,
provenance MCMT could serve as a proxy for genetic structure, enabling genotype-specific
predictions of responses to temperature without the need for genetic data. To determine whether
genetic PCs or home temperature increased the predictive power of the model, we tested simplified
versions of the full model described above: a genetics-only model with home temperature excluded to
test the predictive power of the genetic PCs, and a temperature-only model with genetic information
excluded to test the predictive power of home MCMT (Table 1). We trained the three models on
datasets with each garden excluded and predicted responses in the excluded garden for 2021 and 2022,
evaluating models using the Pearson correlation between actual and predicted heights as described
above.

Table 1. Factors included in each model used for testing whether home climate can be a proxy for
genetic structure in modeling the response to environment

	E: Garden MCMT (quadratic)	**G:** Genetic PCs 1-3	**GxE** (genotype x garden climate interaction)	Home MCMT as a proxy for **G** (quadratic)	Home MCMT x **E**
Full	x	x	x	x	x
Genetics only	x	x	x		
Climate only	x			x	x

Predicting the response to temperature

We used the above model to predict each genotype's norm of reaction; i.e. the response to a

continuous temperature gradient, allowing us to characterize responses to temperature across the

hybrid zone. We predicted the height and mortality probability of each genotype across a range of 100

equally-spaced MCMT values from -23.9 to 9.8 °C, encompassing both historic 30-year home climates

(-23.9 to -3.8 °C) and yearly garden climates (-16.5 to 9.8 °C). Phenotypic predictions should be more

reliable for the warmer climatic range included in the common gardens, where phenotypic data was

collected, than for the coldest parts of the home range. If so, this would increase confidence in our

ability to predict responses to warming climates. We predicted the phenotypic response of each

genotype across this climatic range of MCMT, accounting for genotype-specific responses with the

genetic PCs and the home MCMT of that genotype. The glmmTMB `predict` function was used,

ignoring the random effects of genotype, block, and garden and predicting the overall response to

temperature rather than the genotype- or garden-specific response (option re.form = NA). Predictions

were made for three phenotypes: the conditional model reflecting growth increment, the zero-inflated

model predicting the probability of mortality, and the overall model incorporating both.

We estimated fitness changes under future values of MCMT using the norm of reaction generated

from the model to predict the growth, mortality, and combined fitness of each genotype at its

provenance under historic (1961-1980) and future values of MCMT. Future MCMT values were

predicted from an ensemble containing 13 general circulation models (GCMs) from the CMIP6

database, generated from ClimateNA (Wang et al. 2016) and available at AdaptWest (AdaptWest

Project 2022). We used the time period of 2041-2070 under the shared socioeconomic pathway (SSP)

2-4.5, an intermediate scenario in which emissions rise until mid-century, then decline (IPCC 2023).

To predict how the relative fitness of *P. balsamifera*, *P. trichocarpa*, and hybrid genotypes may change

under future values of MCMT, we predicted which genotypes would have the best performance

(highest overall fitness combining growth and survival) across the species ranges and regions with

common gardens. For each grid cell, we extracted MCMT values for past and future climates, and

identified the genotype with the highest predicted fitness at that temperature.

Results

Variation in growth among gardens

Growth increment varied widely among gardens (Figure S1), with the highest growth occurring at

WSU (Washington) and SU (Maryland) gardens, and the lowest growth occurring at the upper and

lower winter temperature extremes, particularly at OLLU (Texas), UCM (California), OSU (Oregon),

and NDSU (North Dakota). Genotypes typically reached their greatest height at intermediate

temperatures warmer than those of their climate of origin, with height decreasing as MCMT increased

or decreased (Figure S1 and S2), suggesting that MCMT exerts selection pressure on poplars, as
previously found in high-latitude tree species (Leites et al. 2012, 2019; Yeaman et al. 2016; Rehfeldt et
al. 2018; Mahony et al. 2020).

Factors predicting growth and mortality

We evaluated the effect size and significance of each model factor for the conditional model,
representing growth, and for the zero-inflated model, representing the probability of mortality (**Figure
2** and **Table S1**). The two models had similar effects, indicating that the same factors associated with
increased growth were also associated with low mortality. MCMT of the common garden site and its
square term had the strongest effect on growth (**Figure 2** and **Table S1**). Both had negative effects,
indicating decreased growth in climates warmer or colder than the optimum temperature. Intra- and
interspecific genetic structure represented by PC1 and PC3 both had significant effects on growth,
although with a smaller effect size than garden temperature. Individuals with greater *P. trichocarpa*
ancestry (PC1) and from the three southernmost transects (PC3) had higher growth on average. PC2,
which explains genetic structure within *P. trichocarpa* (**Figure S4**), was not statistically significant,
suggesting that genetic differences between interior and coastal *P. trichocarpa* did not significantly
affect growth. Home temperature and its interaction with garden temperature did not significantly
affect growth, suggesting that responses to MCMT were better explained by genetic structure than by
MCMT of origin. While no genotype × environment interaction terms were significant at the p<0.05
level, Garden MCMT^2 × Genetic PC2 was near significant (p=0.052, **Table S1**) and was significant
when only measurements from 2021 were included in the model (**Figure S10**), suggesting a weak
effect of variation in the temperature response curve between two *P. trichocarpa* lineages during first
year growth (**Figure S4**). Overall, our results show that growth is primarily determined by
temperature effects. Genetics also contributed to variation in growth, which suggests the response to
temperature is partially determined by species ancestry and latitudinal variation within both species.
Similarly, in the zero-inflated model, garden temperature significantly affected the probability of
mortality, with increased mortality in sites at the higher and lower temperature extremes. Genetic
structure, home MCMT, and their interactions with garden MCMT were not significant, suggesting
that mortality probability was primarily driven by planting site temperature, with limited variation
among genotypes.

Evaluation of model predictive ability

We evaluated the predictive ability of the full model that included garden climate, climate of origin,
genetic structure, and their interactions (GxE) (**Table 1**) using the Pearson correlation between the
actual and predicted heights for all individuals. Both models generally performed well in predicting the
response to climate: correlation between actual and predicted heights when random effects were
included in the prediction was 0.795 for the overall model, and 0.809 for the conditional model
(**Figure S6**). When random effects were not incorporated, predictive ability decreased to 0.631 and
0.538 for the overall and conditional model, respectively (**Figure S6**), indicating there were garden-

specific effects on plant response that were not accounted for by garden MCMT, and/or genotype-

specific effects that were not accounted for by home MCMT or the genetic PCs, and year-specific

effects. However, the model consistently underpredicted actual growth increment values, particularly

for the tallest trees. For this reason we focus on the relative differences in growth among genotypes and

gardens rather than predicting specific yearly growth increments.

We also evaluated models by testing their predictive power for environments and genotypes not

included in the input "training" dataset, using two methods. First, leave-one-out cross validation was

performed for each garden; we report prediction ability for each garden for the conditional model with

random effects of garden and genotype incorporated into the predictions. Prediction ability (Pearson's

correlation between observed and predicted heights, with dead individuals excluded) varied widely

among gardens (**Figure S7**), with the average being 0.49 for 2021 and 0.38 for 2022, the highest

correlation being 0.67 for OSU in 2022 (Oregon) and the lowest correlation being -0.8 for NDSU in

2022 (North Dakota). Most sites with poor predictions were sites with high mortality (NDSU,

OLLU, LOCK, UCM, and OSU had >40% mortality in the first year). Other sites with low

prediction ability included SWMN (r=0.058 in 2021,) and WI (r=0.4 in 2021, -0.09 in 2022), which

are two sites having low MCMT (**Figure S2**) and therefore may provide unique information on

responses to lower temperatures.

Second, we predicted growth for our three 'maxi' common garden experiments, which included 500

novel genotypes and one novel site using the model trained on all 17 mini gardens (**Figure 3**).

Prediction ability was low for NDSU, likely due to high mortality (Pearson's r=0.04 for the

conditional model). However, prediction ability for the VA and VT predictions was relatively high.

For the conditional model VA had correlation between actual and predicted values of 0.45 and VT

0.65 (**Figure 3**), and when dead individuals were included in the overall model VA had an prediction

ability of 0.48 and VT 0.53 (**Figure S8**). However, as with the leave-one-garden-out tests, the model

underpredicted height. Taken together, these results suggest that for environments with low mortality,

the model can predict relative performance for both novel genotypes and novel environments outside

of the training dataset, but is limited in its prediction of specific growth increments.

The role of genetics in phenotypic prediction across environments

When predicting growth and overall performance for each garden and year using leave-one-out cross

validation, the climate-only model that excluded genetic information had worse performance than the

genetics-only model (p=0.0039) and on average had lower performance than the full model, but did

not significantly differ from it (p = 0.061) (**Table 1, Figure 4**). These results show that genetic

information improved predictions when climate data was excluded, but that models with either

genetics or climate data missing performed similarly to the full model. When random effects of garden

and genotype were included in the predictions, all models performed equally well (**Figure S9**). This is

likely because all genotypes were included in the training datasets allowing the intercept of each

genotype to be estimated, improving predictions of their growth in a novel environment without

explicitly including genetic or climate information.

Using the full model trained on all 17 gardens, we predicted the response of each genotype to a range

of temperatures (**Figure 5**). Our model predicts that genotypes with more *P. trichocarpa* ancestry have

higher growth and overall performance in warmer environments (MCMT ≈ -14 to 10 °C) compared

to *P. balsamifera* genotypes, but that genotypes with more *P. balsamifera* ancestry have higher

growth in colder climates, consistent with the climatic preferences of the two species (**Figure 5A-B**).

Two majority-*P. balsamifera* genotypes were predicted to have biologically unrealistic reaction norms,

with growth increasing exponentially with decreased winter temperatures (**Figure 5B**). These

modeled growth responses likely reflect a preference for very cold environments that are not well-

represented in our gardens, and when the effects of growth and mortality are combined, these

genotypes show realistic response curves with optima at relatively low temperatures. Across species

ancestries, mortality is predicted to increase at the colder and warmer extremes, with a greater

probability of mortality under extreme cold climates than warmer climates (**Figure 5C**). As with

predictions of growth, mortality responses predict that *P. balsamifera* is more cold tolerant, with

increased probability of mortality *P. trichocarpa* genotypes occurring at less extreme cold temperatures

than for *P. balsamifera* (**Figure 5C**). Likewise, the probability of mortality for *P. balsamifera*

generally increases with warmer temperatures more than *P. trichocarpa*. Combining the growth and

mortality predictions as an overall fitness proxy, genotypes with a majority *P. trichocarpa* ancestry have

consistently higher fitness except in the coldest climates, where majority-ancestry *P. balsamifera*

genotypes are predicted to outperform them (**Figure 5A**). The overall maximum growth of *P.*

balsamifera genotypes is lower than that of *P. trichocarpa*, which has previously been reported to be a

faster-growing species. Admixed genotypes with majority *P. trichocarpa* ancestry had maximum

heights intermediate to the two parental species, while admixed majority-*P. balsamifera* genotypes had

responses more similar to parental *P. balsamifera* genotypes. For most genotypes, the highest growth

and lowest mortality occurred in climates warmer than their climate of origin (**Figure 5**).

Rangewide projections of future fitness changes

Our model predicts increased growth and survival under climates with MCMT temperatures warmer

than the climate of origin (**Figure 5D-F** and **Figure S11**). However, we are limited in our ability to

predict responses to multivariate changes in climate. Here, we focus on relative performance of

genotypes rather than absolute changes in performance metrics. Predictions of the best-performing

genotype under historic values of MCMT generally follows previously-described species ranges; with

P. trichocarpa occurring along the west coast of North America and *P. balsamifera* occurring in the

more northern and interior regions of the continent (**Figure 6**). Hybrid genotypes are predicted to outcompete parental species in intermediate regions of the contact zone, particularly in central British Columbia, where sampled genotypes exhibited high levels of admixture. Under future climates (2041-2070 for a "middle-of-the-road" emissions scenario), the model predicts shifts in the best-performing genotype following the two species' climatic preferences: the regions suitable for *P. trichocarpa* and its backcrosses will shift northward as climate warms (**Figure 6**). For example, interior sites where majority-*P. balsamifera* genotypes were sampled are predicted to become more favorable to increased *P. trichocarpa* ancestry, which could favor introgression of *P. trichocarpa* genes into these regions.

Figures

Figure 1. (A) Map of sampled genotype localities (circles) and common garden sites (triangles) in
relation to the ranges of both species (Little 1971); color of sampled genotypes indicates species
ancestry calculated from ADMIXTURE at K=2. (B-C) PCAs of past and future climates across
collection sites and garden sites. Circles represent the historic climate (1961-1990) of each genotype's
provenance, colored by species ancestry, and arrowheads represent the predicted climate for 2041-
2070 under an ensemble of 13 GCMs. Triangles and squares represent the climate at common garden
sites for 2021 and 2021. Text indicates the loadings of climate and geographic variables on each axis,
with abbreviations as follows: CMD, climatic moisture deficit; lat, latitude; lon, longitude; MAP,
mean annual precipitation, MAT, mean annual temperature; MCMT, mean coldest month
temperature; MWMT, mean warmest month temperature, PAS, precipitation as snow; RH, relative
humidity; TD, temperature difference, or continentality. Loadings of climate variables have been
downscaled to allow visibility of sites.

Annual Growth Increment (log transformed)

Figure 2. Forest plot of model coefficients using sjPlot (Lüdecke 2024); with an estimate of one indicating no effect, and estimates significantly greater or less than one indicating a positive and negative effect, respectively. Coefficients are standardized to allow comparison when interaction effects are included. Top plot shows results for the conditional model (growth increment) and bottom plot shows the estimates for the zero-inflated model (mortality); the effects for the two models are reversed

Figure 3. Performance of model predictions in the maxi gardens, predicting growth across 544
genotypes (500 novel genotypes) and one novel environment (VT) using a model trained on all 17
mini gardens. Models were evaluated by comparing predicted and actual height for the conditional
model with no random effects for garden and genotype. As the conditional portion of the model
predicts growth rather than mortality, individuals that did not survive were removed from the model.
R values and p-values are given for the Pearson correlation between actual and predicted height.

Figure 4. Comparison of model performance among the full model and those excluding provenance
MCMT or genetic PCs (Table 1), estimated as the correlation between predicted and observed growth
increments for each garden and year predicted using the leave-one-out models, in which heights were
predicted for a single year and garden using a model trained on other gardens. The model including
genetic data without home temperature data resulted in better predictive ability than the model with
temperature data alone. Other comparisons with the overall model and with random effects included
are shown in Figure S9. ** indicates p ≤0.01, "ns" indicates not significant.

Figure 5. Model predictions for each genotype's growth and mortality response to temperature, with each genotype indicated as a separate line colored by species ancestry at K=2, with green indicating *P. trichocarpa* and blue indicating *P. balsamifera*. Predictions for the overall model (A and D) incorporate both growth and the probability of mortality, predictions for the conditional model (B and E) only predict growth, ignoring the probability of zeros arising from other processes (mortality), and predictions for the zero-inflated model (C and F) predict the probability of zeros arising from mortality. A-C: Responses across the range of mean coldest month temperature (MCMT) values at garden and home climates. Actual values of home climates (|) and garden climates (circle) used for model training are shown on the x-axis. D-E. Responses based on distance from climate of origin (garden - home MCMT); positive values indicate a warmer climate (higher coldest month temperature) and negative values indicate a colder climate. If genotypes perform best in environments similar to their home environment, growth should be highest and mortality should be lowest when the transfer distance is 0.

Figure 6. Maps showing the species ancestry of the studied genotype which is predicted to have highest fitness (as measured by growth and mortality) in that location under historic (A and B) and future climate (C and D), indicated by the color of the base layer. A and C show predictions for the species ranges and studied locations, and B and D show the same predictions for the sampled hybrid zone region. Regions with MCMT values outside of the range measured in the common gardens

(-13.05 to 10.85 °C) are masked and colored white. Actual ancestry of collected genotypes are shown
as circles, and locations of common gardens are shown as triangles. Species ranges are shown as dark
blue and green outlines (Little 1971). As MCMT increases, we predict that genotypes with higher *P.*
trichocarpa ancestry may be able to outcompete genotypes with higher *P. balsamifera* ancestry in some
portions of the *P. balsamifera range*, favoring a northeastern shift of the *P. trichocarpa* range and the
hybrid zone and into historically colder, more continental regions.

Discussion
Using 17 common gardens with *Populus* genotypes which originated from a hybrid zone spanning a
broad temperature range, we evaluated the genotype-specific response to warming winter temperatures
and predicted future responses across the hybrid zone. We found that 1) Fitness metrics and reaction
norms varied among genotypes due to species ancestry and region of origin, consistent with a trade-off
between cold tolerance and growth potential in warmer environments, 2) hybrids displayed reaction
norms and temperature optima intermediate between their parental species, suggesting that warming
temperatures could favor movement of the hybrid zone, and 3) a significant variance in growth
explained by genetic structure illustrates the importance of genetic variation in predicting responses to
climate change.

Question 1: Do genotypes from different climates vary in fitness responses across environments?
Local adaptation to climate can shape not only traits, but their reaction norms across environments,
contributing to variation in plastic responses within and across species (Patsiou et al. 2020). We found
that poplar growth and survival across common garden environments was predicted by species
ancestry and by genetic differences associated with a northern and southern region within each species
(**Figure 2**), illustrating that genetic variation will contribute to spatially varying responses to warming
temperatures (Patsiou et al. 2020; Martinez del Castillo et al. 2024). Performance traits varied along a
continuous ancestry gradient between the two parental species, consistent with their climatic niches.
In general, individuals with greater *P. trichocarpa* ancestry had greater overall growth, but these
genotypes experienced lower growth and increased mortality in cold, continental environments typical
of the *P. balsamifera* range (**Figure 5**). In addition, genotypes from the three southernmost transects
had higher maximum heights than those from the two northern transects (as described by genetic
PC3). This is consistent with a tradeoff between growth potential and cold tolerance often observed in
temperate or boreal tree species, likely enabling *P. balsamifera* to outcompete *P. trichocarpa* in colder
regions (Leites et al. 2012; Menon et al. 2015; Rehfeldt et al. 2018). While no G×E terms were
significant in the model based on two years of data, the effect of garden $MCMT^2$ × genetic PC2 was

near significant (p = 0.052) (**Figure 2, Table S1**) and was significant when only measurements from
2021 were included in the model (Figure S10). This result suggests a weak effect of genetic variation
between two *P. trichocarpa* lineages on the plastic response to winter temperatures, which may be
partially explained by maternal effects expressed in the first year of growth. Additionally, optimum
temperatures varied for each genotype, with *P. trichocarpa* individuals reaching maximum heights in
warmer environments than *P. balsamifera* **(Figure 5)**. This suggests that selection associated with
minimum temperatures may be acting to produce different plastic responses across the range of both
species and their hybrid zone. While the majority of *P. balsamifera* genotypes had lower overall
growth, their climatic range appeared to be wider (**Figure 5**), perhaps reflecting adaptation to more
continental climates with larger annual temperature ranges (**Figure 1**). Generally, the response of
hybrid genotypes was intermediate between the parental species; however, some hybrid genotypes have
reduced probability of mortality at lower temperatures (**Figure 5**), suggesting that introgression could
promote increased cold hardiness (Hamilton et al. 2013). However, the effects of species ancestry and
geography may be difficult to disentangle, as genotypes from the coldest part of the sampled range
(Alaska and northern British Columbia) are admixed (**Figure S4**).

In this study, we find that most genotypes reach their fitness optima in environments that are warmer
than their climate of origin (**Figure 6**), and that genotypes planted at sites having MCMT values
similar to their home site did not always perform better than other, nonlocal genotypes (**Figure S12**).
Taken together, these results suggest genotypes may not be locally adapted to winter temperatures
alone when considering either the "home vs away" or "local vs foreign" criteria (Kawecki and Ebert
2004). This decoupling between the physiological optimum, the climate in which genotypes reach
their maximum growth rate, and the ecological optimum, the climate in which they exist in the
ecosystem, has been previously observed for temperate tree species (Rehfeldt et al. 2018). Higher
growth rates in non-local environments does not necessarily indicate a lack of local adaptation
(Kawecki and Ebert 2004), but provides important context for predicting responses to climate change,
which often assume optimal performance in local climates. Instead, we found that most genotypes,
when in gardens with temperatures similar to their climate of origin, were outperformed by a small
number of primarily *P. trichocarpa* genotypes. These genotypes may be ideal for planting as part of
restoration projects using assisted gene flow or for biofuel production across diverse environments,
particularly under increasingly variable and unpredictable climate futures (Sannigrahi et al. 2010;
Porth and El-Kassaby 2015; Mahoney et al. 2019). However, if increased growth corresponds to a lack
of cold tolerance or lower survival and fecundity (Leites et al. 2012, 2019; Yeaman et al. 2016; Rehfeldt
et al. 2018; Mahony et al. 2020), caution is warranted when planting in locations susceptible to cold
stress. Further work comparing the cold tolerance of these genotypes could determine the lower
temperature limits where they are likely to be successful. Regardless of the mechanisms underlying

genetic variation in fitness proxies, the large differences in responses to temperature illustrate the

importance and benefit of measuring reaction norms across wide geographic regions within and

among species.

Our predictions of increased growth under warming climates for these *Populus* genotypes raises the

question: is active conservation and management of poplar in this region necessary, or should we

prioritize species more vulnerable to increased temperatures? However, predictions of increased

growth should be treated with caution, as they only represent aboveground height and mortality in the

first two years of growth and exclude reproductive measures of fitness and belowground biomass

(Fischer et al. 2007), which could vary between the species given the presence of *P. balsamifera* in drier

environments (**Figure 1B**). Additionally, we only modeled growth over two growing seasons, in

relatively warm climates and with irrigation during the first year, limiting the selective response to cold

or drought stresses that genotypes may experience in their home environments. Major selective

weather events may occur only rarely; for example, an unusually cold winter could select against

genotypes with high growth and low cold tolerance, favoring cold-tolerant genotypes over longer

timescales (Rehfeldt et al. 2018; Lowry et al. 2019). In this case, our dataset may underestimate the

benefit that *P. balsamifera* has in colder regions. Similarly, if warming winters are accompanied by

summer heat waves, the benefits of warming we observe may not persist long-term in wild

populations. We also did not consider biotic factors such as competition with more cold-hardy

conifers or pathogen pressure that may exclude *Populus* from warmer climates (La Mantia et al. 2013;

Suarez-Gonzalez et al. 2018a), or interactions with belowground communities (Fischer et al. 2014b). If

biotic interactions shift along with climate change, overall fitness could decrease even if higher

temperatures favor faster aboveground growth. For example, pressure from leaf rust (*Melampsora*

spp.) may be higher in warmer, more southern regions of the *P. trichocarpa* range (La Mantia et al.

2013; McKown et al. 2014; Suarez-Gonzalez et al. 2018). Lastly, our predictions of the response to

climate are limited by the training data used to create the model – in this case, the climatic range of the

common gardens (Rogers and Holland 2021). Because the gardens were largely planted to the south

and in warmer climates than the climate of origin, most temperature variables (including multivariate

PC axes) had little overlap between home and garden climates, limiting our ability to make predictions

for growth and mortality in the colder historic climates of origin. We choose to use MCMT as the

predictor variable because of its biological importance in past studies (Leites et al. 2012, 2019; Yeaman

et al. 2016; Rehfeldt et al. 2018; Mahony et al. 2020) and because garden climates included

temperatures matching historic home climates for both *P. balsamifera* and *P. trichocarpa* genotypes

(Figure 1B-C, Figure 5). However, we lack common gardens matching the coldest collection sites,

limiting our predictions for the multivariate home environments experienced by natural populations,

including some genotypes studied here. Given these limitations, testing how native poplar populations

are responding to climate change *in situ* could complement common garden studies (Sharma et al.
2022; Astigarraga et al. 2024; Martinez del Castillo et al. 2024).

Question 2: How will warming temperatures alter the geography of the two species and their hybrid zone?
Geographic variation in the magnitude of climate change and differences in phenotypic plasticity
among genotypes could alter regions where certain genomic backgrounds outcompete others,
potentially leading to range shifts for the two species and their hybrid zone. While we predict that local
poplar populations across the sampled region could experience increased growth and decreased
mortality from increased winter temperatures, they must contend with competition from non-local
genotypes that may have greater fitness under novel climates than local genotypes. Species ranges are
already beginning to shift in response to climate change (Parmesan 2006; Bell et al. 2014; Astigarraga et
al. 2024) and multi-species hybrid zones may also shift (Taylor et al. 2014, 2015; Billerman et al. 2016;
Hamilton and Miller 2016; Wielstra 2019; Alexander et al. 2022). If dispersal of poplar genotypes is
fast enough to track changes in climate, regions where particular genomic backgrounds dominate may
shift under climate change. For example, under higher minimum winter temperatures, *P. balsamifera*
may be outcompeted by less cold-tolerant but faster growing genotypes with higher *P. trichocarpa*
ancestry (**Figure 6**). However, such shifts in genetic composition would rely on gene flow or
migration being fast enough to track climate change, either through dispersal of genotypes or seeds or
wind-dispersed pollen (Corlett and Westcott 2013). While *Populus* can likely disperse over long
distances via wind dispersal or via river-dispersed vegetative material (Kling and Ackerly 2021),
geographic barriers to gene flow persist across the hybrid zone, particularly in mountainous regions
(Bolte et al. 2024). Our model predicts that parental *P. trichocarpa* genotypes may begin to
outcompete *P. trichocarpa* backcrosses in central British Columbia, a region with high gene flow (Bolte
et al. 2024), so this region may see shifts in the genetic composition of poplars (**Figure 6**). Conversely,
while the regions where *P. trichocarpa* ancestry is favored are predicted to shift to the northeast in
Alberta, the Rocky Mountains are a barrier to gene flow that may prevent their dispersal (Bolte et al.
2024). Our predictions in colder regions, including the *P. balsamifera* range, are likely limited because
our dataset includes few parental *P. balsamifera* genotypes and we did not measure growth in gardens
with cold climates matching this part of the *P. balsamifera* range: as a result, it predicts backcrossed *P.*
balsamifera genotypes would perform better than parental *P. balsamifera* in northern, interior North
America within the core of the species distribution (**Figure 6**). Because the reaction norms of parental
and backcrossed *P. balsamifera* are very similar (**Figure 5A**), we interpret these as regions where
increased *P. balsamifera* ancestry is beneficial, not necessarily that backcrosses will be significantly
more successful than parental genotypes.

Spatial changes in the location of the hybrid zone could provide an opportunity for adaptive
introgression to occur, enabling individual alleles and novel allelic combinations to track climate

change. Ancient and contemporary introgression can allow species to persist as environments change

(Kremer and Hipp 2020; Leroy et al. 2020; O'Donnell et al. 2021; Buck et al. 2023; Yang et al. 2023).

The presence of admixed genotypes in intermediate climates in this *Populus* hybrid zone and the

reaction norms that are intermediate between the two parental species are consistent with a the

bounded hybrid superiority model, in which hybrids have higher fitness than either parental species in

intermediate climates (Hamilton et al. 2013; De La Torre et al. 2014; Bolte et al. 2024). Movement of

the *P. trichocarpa* x *P. balsamifera* hybrid zone could allow genes to move across species boundaries at

new contact zones, increasing the levels of standing genetic variation in these regions and facilitating

introgression of alleles that are beneficial under novel climates into new regions.

Question 3: What information is needed to predict climate change responses in trees?

Like generations of provenance studies, we find that the environment has a large effect on tree

performance (O'Neill et al. 2008; Wang et al. 2010; Leites et al. 2012; Browne et al. 2019; Leites and

Benito Garzón 2023; Ye et al. 2023). Understanding how growth and survival varies across species,

populations, and environments will be essential to predicting shifts in the performance and carbon

sequestration abilities of trees under changing climates. Because the temperature of the planting site

had the greatest effect in determining growth, common garden experiments will continue to be

essential to predicting phenotypic responses to the environment. We also show that a model including

genetic PCs but not climate of origin significantly improved predictions of growth compared to a

model containing only home and garden climates (Figure 5), suggesting that universal transfer

functions for predicting tree growth (O'Neill et al. 2008; Wang et al. 2010), could benefit from genetic

information, as found by other studies (Mahony et al. 2020; Archambeau et al. 2022; Putra et al. 2023;

Li et al. 2024). Furthermore, the model that included genetic information but not climate of origin

predicted growth equally as well as the full model, suggesting genetic structure may be used to

effectively model responses to climate in this system. Genetic isolation-by-environment is stronger

than isolation-by-distance in this system (Bolte et al. 2024), suggesting environmental adaptation

drives much of the observed genetic structure. Conversely, in species with high gene flow and less

structured populations, adaptation to climate may be the primary factor contributing to intraspecific

variation, and genetic structure may not be as effective in predicting responses.

Conclusions

Across 17 common garden experiments spanning a wide climatic range, the performance of *Populus*

genotypes was explained by genetic variation associated with species ancestry and origin from northern

or southern regions. Genotypes with greater *P. trichocarpa* ancestry and from southern regions had

higher growth potential, but at a cost of decreased growth and increased mortality in cold regions

typical of *P. balsamifera*. By characterizing genotype-specific reaction norms, we were able to predict

which genotypes should have the highest performance in historic and future climates across the hybrid
zone, identifying regions where the contact zone is favored to shift under warming winter
temperatures. If migration and gene flow is able to track climate change, a moving hybrid zone could
facilitate adaptive introgression of alleles beneficial under novel climates, enabling poplar populations
to persist in their ecosystems even as their genetic makeup may change. This study shows the large
effect that environment has on performance, illustrating the continued value that common garden
experiments will have in predicting organisms' responses to climate change. Furthermore, using
genomic data, we were able to characterize how genetic structure shapes varying responses within and
among species, enabling predictions of climate change responses to take into account the genetic
variation in traits that exists in natural populations.

Acknowledgments

This research was supported by NSF PGR 1856450, the Schatz Center for Tree Molecular Genetics,
National Institute of Food and Agriculture Award #PEN04809 to JAH, NIH award R35GM138300
to JRL, and the Life and Environmental Sciences Dept. at UCM. AM was supported by NSF PRFP
2209410. NM was supported by University of Wisconsin– Eau Claire Office of Research and
Sponsored Programs through the Biology Research Scholars Program and Summer Research
Experience for Undergraduate awards.

We would like to thank everyone involved in with sampling, garden establishment and maintenance,
and data collection, including Shane Baumgart, Joseph Braasch, Schaefer Buchanan, Ian Clancy-
Mallue, Emma Collins, Kevin Cowden, Lionel Di Santo, Marcia Fairbanks, Alice Fischer, Abby
Ferson-Mitchell, Linnea Fraser, Carter Guffey, Carri LeRoy, David Hainlen, Alex Hass, Christian
Hernandez, Kristina Hufford, Daria Hutchinson, Erika Jones, Lee Kalcsits, Ben Levitt, Eli Levitt,
James Levitt, Jessica Lindstrom, Abigail Marshall, Luke McCormack, Albert Medel, Chloe Meyer,
Katie Nelson, George Newcombe, Thu Nguyen, Amanda Penn, Jennifer Pollard, Steven Quick,
Michelle Reid, Brendon Reidy, Christy Rollinson, Seth Townsend, Paul Warnick, Rebekah Wells,
Dean Wu, Kate Volk, Lindsey Zakopal, and Thomas Zambiasi. We would also like to thank Tongli
Wang for discussions of climate response functions.

Data Availability

Scripts for filtering genomic data are available at
github.com/alaynamead/poplar_hybrid_vcf_filtering.

Analysis scripts are available at github.com/alaynamead/popup_poplar_reaction_norms and

corresponding R markdown files with script outputs are provided as supplementary material. Github

repositories will be archived at Zenodo upon manuscript acceptance.

Data from this study is included as supplementary material and will be archived in Dryad upon

manuscript acceptance.

References

AdaptWest Project. 2022. Gridded current and projected climate data for North America at 1km
resolution, generated using the ClimateNA v7.30 software (T. Wang et al., 2022). Available at
adaptwest.databasin.org.

Aitken, S. N., and J. B. Bemmels. 2016. Time to get moving: assisted gene flow of forest trees.
Evolutionary Applications 9:271–290.

Aitken, S. N., and M. C. Whitlock. 2013. Assisted gene flow to facilitate local adaptation to climate
change. Annu. Rev. Ecol. Evol. Syst 44:367–88.

Alexander, A., M. B. Robbins, J. Holmes, R. G. Moyle, and A. T. Peterson. 2022. Limited movement
of an avian hybrid zone in relation to regional variation in magnitude of climate change.
Molecular Ecology 31:6634–6648.

Alexander, D. H., J. Novembre, and K. Lange. 2009. Fast model-based estimation of ancestry in
unrelated individuals. Genome Res 19:1655–1664.

Archambeau, J., M. Benito Garzón, F. Barraquand, M. de Miguel, C. Plomion, and S. C. González-
Martínez. 2022. Combining Climatic and Genomic Data Improves Range-Wide Tree Height
Growth Prediction in a Forest Tree. The American Naturalist 200:E141–E159. The
University of Chicago Press.

Arnold, P. A., L. E. B. Kruuk, and A. B. Nicotra. 2019. How to analyse plant phenotypic plasticity in
response to a changing climate. New Phytologist 222:1235–1241.

Astigarraga, J., A. Esquivel-Muelbert, P. Ruiz-Benito, F. Rodríguez-Sánchez, M. A. Zavala, A. Vilà-
Cabrera, M.-J. Schelhaas, G. Kunstler, C. W. Woodall, E. Cienciala, J. Dahlgren, L. Govaere,
L. A. König, A. Lehtonen, A. Talarczyk, D. Liu, and T. A. M. Pugh. 2024. Relative decline in
density of Northern Hemisphere tree species in warm and arid regions of their climate niches.
Proceedings of the National Academy of Sciences 121:e2314899121. Proceedings of the
National Academy of Sciences.

Bell, D. M., J. B. Bradford, and W. K. Lauenroth. 2014. Early indicators of change: divergent climate
envelopes between tree life stages imply range shifts in the western United States. Global
Ecology and Biogeography 23:168–180.

Billerman, S. M., M. A. Murphy, and M. D. Carling. 2016. Changing climate mediates sapsucker
(Aves: Sphyrapicus) hybrid zone movement. Ecology and Evolution 6:7976–7990.

Bolte, C. E., T. Phannareth, M. Zavala-Paez, B. N. Sutara, M. F. Can, M. C. Fitzpatrick, J. A.
Holliday, S. R. Keller, and J. A. Hamilton. 2024. Genomic insights into hybrid zone
formation: The role of climate, landscape, and demography in the emergence of a novel hybrid
lineage. Molecular Ecology 33:e17430.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J. Skaug,
M. Mächler, and B. M. Bolker. 2017. glmmTMB Balances Speed and Flexibility Among
Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal 9:378–400.

Browne, L., J. W. Wright, S. Fitz-Gibbon, P. F. Gugger, and V. L. Sork. 2019. Adaptational lag to
temperature in valley oak (*Quercus lobata*) can be mitigated by genome-informed assisted gene
flow. PNAS 116:25179–25185. National Academy of Sciences.

Buck, R., D. Ortega-Del Vecchyo, C. Gehring, R. Michelson, D. Flores-Rentería, B. Klein, A. V.

Whipple, and L. Flores-Rentería. 2023. Sequential hybridization may have facilitated
ecological transitions in the Southwestern pinyon pine syngameon. New Phytologist
237:2435–2449.
Capblancq, T., M. C. Fitzpatrick, R. A. Bay, M. Exposito-Alonso, and S. R. Keller. 2020. Genomic
Prediction of (Mal)Adaptation Across Current and Future Climatic Landscapes. Annual
Review of Ecology, Evolution, and Systematics 51:245–269.
Chhatre, V. E., L. M. Evans, S. P. DiFazio, and S. R. Keller. 2018. Adaptive introgression and
maintenance of a trispecies hybrid complex in range-edge populations of Populus. Molecular
Ecology 27:4820–4838.
Chhetri, H. B., D. Macaya-Sanz, D. Kainer, A. K. Biswal, L. M. Evans, J.-G. Chen, C. Collins, K.
Hunt, S. S. Mohanty, T. Rosenstiel, D. Ryno, K. Winkeler, X. Yang, D. Jacobson, D.
Mohnen, W. Muchero, S. H. Strauss, T. J. Tschaplinski, G. A. Tuskan, and S. P. DiFazio.
2019. Multitrait genome-wide association analysis of Populus trichocarpa identifies key
polymorphisms controlling morphological and physiological traits. New Phytologist 223:293–
309.
Cooper, H. F., R. J. Best, L. V. Andrews, J. P. M. Corbin, I. Garthwaite, K. C. Grady, C. A. Gehring,
K. R. Hultine, T. G. Whitham, and G. J. Allan. 2022. Evidence of climate-driven selection on
tree traits and trait plasticity across the climatic range of a riparian foundation species.
Molecular Ecology 31:5024–5040.
Cooper, H. F., K. C. Grady, J. A. Cowan, R. J. Best, G. J. Allan, and T. G. Whitham. 2019. Genotypic
variation in phenological plasticity: Reciprocal common gardens reveal adaptive responses to
warmer springs but not to fall frost. Global Change Biology 25:187–200.
Corlett, R. T., and D. A. Westcott. 2013. Will plant movements keep up with climate change? Trends
in Ecology & Evolution 28:482–488. Elsevier.
De La Torre, A. R., T. Wang, B. Jaquish, and S. N. Aitken. 2014. Adaptation and exogenous selection
in a Picea glauca × Picea engelmannii hybrid zone: implications for forest management under
climate change. New Phytologist 201:687–699.
Des Marais, D. L., K. M. Hernandez, and T. E. Juenger. 2013. Genotype-by-environment interaction
and plasticity: exploring genomic responses of plants to the abiotic environment. Annual
Review of Ecology, Evolution, and Systematics 44:5–29.
Evans, L. M., G. T. Slavov, E. Rodgers-Melnick, J. Martin, P. Ranjan, W. Muchero, A. M. Brunner,
W. Schackwitz, L. Gunter, J.-G. Chen, G. A. Tuskan, and S. P. DiFazio. 2014. Population
genomics of *Populus trichocarpa* identifies signatures of selection and adaptive trait
associations. Nature Genetics 46:1089–1096.
Fischer, D. G., S. K. Chapman, A. T. Classen, C. A. Gehring, K. C. Grady, J. A. Schweitzer, and T. G.
Whitham. 2014a. Plant genetic effects on soils under climate change. Plant Soil 379:1–19.
Fischer, D. G., S. K. Chapman, A. T. Classen, C. A. Gehring, K. C. Grady, J. A. Schweitzer, and T. G.
Whitham. 2014b. Plant genetic effects on soils under climate change. Plant Soil 379:1–19.
Fischer, D. G., S. C. Hart, C. J. LeRoy, and T. G. Whitham. 2007. Variation in below-ground carbon
fluxes along a Populus hybridization gradient. New Phytologist 176:415–425.
Fitzpatrick, M. C., V. E. Chhatre, R. Y. Soolanayakanahally, and S. R. Keller. 2021. Experimental

support for genomic prediction of climate maladaptation using the machine learning approach
Gradient Forests. Molecular Ecology Resources 21:2749–2765.

Fitzpatrick, M. C., and S. R. Keller. 2015. Ecological genomics meets community-level modelling of
biodiversity: mapping the genomic landscape of current and future environmental adaptation.
Ecology Letters 18:1–16.

Geraldes, A., N. Farzaneh, C. J. Grassa, A. D. McKown, R. D. Guy, S. D. Mansfield, C. J. Douglas,
and Q. C. B. Cronk. 2014. Landscape Genomics of Populus Trichocarpa: The Role of
Hybridization, Limited Gene Flow, and Natural Selection in Shaping Patterns of Population
Structure. Evolution 68:3260–3280.

Grady, K. C., T. E. Kolb, D. H. Ikeda, and T. G. Whitham. 2015. A bridge too far: cold and pathogen
constraints to assisted migration of riparian forests. Restoration Ecology 23:811–820.

Gray, L. K., T. Gylander, M. S. Mbogga, P. Chen, and A. Hamann. 2011. Assisted migration to
address climate change: recommendations for aspen reforestation in western Canada.
Ecological Applications 21:1591–1603.

Hamilton, J. A., C. Lexer, and S. N. Aitken. 2013. Genomic and phenotypic architecture of a spruce
hybrid zone (Picea sitchensis × P. glauca). Molecular Ecology 22:827–841.

Hamilton, J. A., and J. M. Miller. 2016. Adaptive introgression as a resource for management and
genetic conservation in a changing climate. Conservation Biology 30:33–41.

Hijmans, R. J. 2025. terra: Spatial Data Analysis.

Holliday, J. A., L. Zhou, R. Bawa, M. Zhang, and R. W. Oubida. 2016. Evidence for extensive
parallelism but divergent genomic architecture of adaptation along altitudinal and latitudinal
gradients in Populus trichocarpa. New Phytologist 209:1240–1251.

Hord, A. M., D. G. Fischer, J. A. Schweitzer, C. J. LeRoy, T. G. Whitham, and J. K. Bailey. 2025.
Hybrid introgression as a mechanism of rapid evolution and resilience to climate change in a
riparian tree species. Communications Biology.

Hu, M.-C., M. Pavlicova, and E. V. Nunes. 2011. Zero-Inflated and Hurdle Models of Count Data
with Extra Zeros: Examples from an HIV-Risk Reduction Intervention Trial. The American
Journal of Drug and Alcohol Abuse 37:367–375. Taylor & Francis.

Ikeda, D. H., T. L. Max, G. J. Allan, M. K. Lau, S. M. Shuster, and T. G. Whitham. 2017. Genetically
informed ecological niche models improve climate change predictions. Global Change Biology
23:164–176.

IPCC. 2023. Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III
to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. , doi:
10.59327/IPCC/AR6-978929169164.

Janes, J. K., and J. A. Hamilton. 2017. Mixing It Up: The Role of Hybridization in Forest
Management and Conservation under Climate Change. Forests 8:237. Multidisciplinary
Digital Publishing Institute.

Jansson, S., and C. J. Douglas. 2007. Populus: A Model System for Plant Biology. Annual Review of
Plant Biology 58:435–458. Annual Reviews.

Kawecki, T. J., and D. Ebert. 2004. Conceptual issues in local adaptation. Ecology Letters 7:1225–
1241.

Keller, S. R., N. Levsen, M. S. Olson, and P. Tiffin. 2012. Local Adaptation in the Flowering-Time
Gene Network of Balsam Poplar, Populus balsamifera L. Molecular Biology and Evolution
29:3143–3152.
Keller, S. R., M. S. Olson, S. Silim, W. Schroeder, and P. Tiffin. 2010. Genomic diversity, population
structure, and migration following rapid range expansion in the Balsam Poplar, Populus
balsamifera. Molecular Ecology 19:1212–1226.
Keller, S. R., R. Y. Soolanayakanahally, R. D. Guy, S. N. Silim, M. S. Olson, and P. Tiffin. 2011.
Climate-driven local adaptation of ecophysiology and phenology in balsam poplar, Populus
balsamifera L. (Salicaceae). American Journal of Botany 98:99–108.
Kling, M. M., and D. D. Ackerly. 2021. Global wind patterns shape genetic differentiation,
asymmetric gene flow, and genetic diversity in trees. Proceedings of the National Academy of
Sciences 118:e2017317118. Proceedings of the National Academy of Sciences.
Kremer, A., and A. L. Hipp. 2020. Oaks: an evolutionary success story. New Phytologist 226:987–
1011.
La Mantia, J., J. Klápště, Y. A. El-Kassaby, S. Azam, R. D. Guy, C. J. Douglas, S. D. Mansfield, and R.
Hamelin. 2013. Association Analysis Identifies Melampsora ×columbiana Poplar Leaf Rust
Resistance SNPs. PLOS ONE 8:e78423. Public Library of Science.
Leites, L., and M. Benito Garzón. 2023. Forest tree species adaptation to climate across biomes:
Building on the legacy of ecological genetics to anticipate responses to climate change. Global
Change Biology 29:4711–4730.
Leites, L. P., G. E. Rehfeldt, and K. C. Steiner. 2019. Adaptation to climate in five eastern North
America broadleaf deciduous species: Growth clines and evidence of the growth-cold tolerance
trade-off. Perspectives in Plant Ecology, Evolution and Systematics 37:64–72.
Leites, L. P., A. P. Robinson, G. E. Rehfeldt, J. D. Marshall, and N. L. Crookston. 2012. Height-
growth response to climatic changes differs among populations of Douglas-fir: a novel analysis
of historic data. Ecological Applications 22:154–165.
Leroy, T., Q. Rougemont, J.-L. Dupouey, C. Bodénès, C. Lalanne, C. Belser, K. Labadie, G. Le
Provost, J.-M. Aury, A. Kremer, and C. Plomion. 2020. Massive postglacial gene flow between
European white oaks uncovered genes underlying species barriers. New Phytologist 226:1183–
1197.
Li, F., D. J. Gates, E. S. Buckler, M. B. Hufford, G. M. Janzen, R. Rellán-Álvarez, F. Rodríguez-
Zapata, J. A. R. Navarro, R. J. H. Sawers, S. J. Snodgrass, K. Sonder, M. C. Willcox, S. J.
Hearne, J. Ross-Ibarra, and D. E. Runcie. 2024. The utility of environmental data from
traditional varieties for climate-adaptive maize breeding. bioRxiv 2024.09.19.613351. Cold
Spring Harbor Laboratory.
Little, E. L. 1971. Atlas of United States trees. U.S. Department of Agriculture, Forest Service,
Washington, DC.
Love, S. J., J. A. Schweitzer, S. A. Woolbright, and J. K. Bailey. 2023. Sky Islands Are a Global Tool for
Predicting the Ecological and Evolutionary Consequences of Climate Change. Annual Review
of Ecology, Evolution, and Systematics 54:219–236. Annual Reviews.
Lowry, D. B., J. T. Lovell, L. Zhang, J. Bonnette, P. A. Fay, R. B. Mitchell, J. Lloyd-Reilley, A. R. Boe,

Y. Wu, F. M. Rouquette, R. L. Wynia, X. Weng, K. D. Behrman, A. Healey, K. Barry, A.
Lipzen, D. Bauer, A. Sharma, J. Jenkins, J. Schmutz, F. B. Fritschi, and T. E. Juenger. 2019.
QTL × environment interactions underlie adaptive divergence in switchgrass across a large
latitudinal gradient. Proceedings of the National Academy of Sciences 116:12933–12941.
Proceedings of the National Academy of Sciences.

Lüdecke, D. 2024. sjPlot: Data Visualization for Statistics in Social Science.

Mahoney, S. M., J. B. Mike, J. M. Parker, L. S. Lassiter, and T. G. Whitham. 2019. Selection for
genetics-based architecture traits in a native cottonwood negatively affects invasive tamarisk in
a restoration field trial. Restoration Ecology 27:15–22.

Mahony, C. R., I. R. MacLachlan, B. M. Lind, J. B. Yoder, T. Wang, and S. N. Aitken. 2020.
Evaluating genomic data for management of local adaptation in a changing climate: A
lodgepole pine case study. Evolutionary Applications 13:116–131.

Martinez del Castillo, E., M. C. A. Torbenson, F. Reinig, E. Tejedor, M. de Luis, and J. Esper. 2024.
Contrasting Future Growth of Norway Spruce and Scots Pine Forests Under Warming
Climate. Global Change Biology 30:e17580.

McKown, A. D., R. D. Guy, J. Klápště, A. Geraldes, M. Friedmann, Q. C. B. Cronk, Y. A. El-Kassaby,
S. D. Mansfield, and C. J. Douglas. 2014a. Geographical and environmental gradients shape
phenotypic trait variation and genetic structure in *Populus trichocarpa*. New Phytologist
201:1263–1276.

McKown, A. D., R. D. Guy, L. Quamme, J. Klápště, J. L. Mantia, C. P. Constabel, Y. A. El-Kassaby,
R. C. Hamelin, M. Zifkin, and M. S. Azam. 2014b. Association genetics, geography and
ecophysiology link stomatal patterning in *Populus trichocarpa* with carbon gain and disease
resistance trade-offs. Molecular Ecology 23:5771–5790.

McKown, A. D., J. Klápště, R. D. Guy, A. Geraldes, I. Porth, J. Hannemann, M. Friedmann, W.
Muchero, G. A. Tuskan, J. Ehlting, Q. C. B. Cronk, Y. A. El-Kassaby, S. D. Mansfield, and C.
J. Douglas. 2014c. Genome-wide association implicates numerous genes underlying ecological
trait variation in natural populations of *Populus trichocarpa*. New Phytologist 203:535–553.

Menon, M., W. J. Barnes, and M. S. Olson. 2015. Population genetics of freeze tolerance among
natural populations of *Populus balsamifera* across the growing season. New Phytologist
207:710–722.

O'Donnell, S. T., S. T. Fitz-Gibbon, and V. L. Sork. 2021. Ancient Introgression Between Distantly
Related White Oaks (Quercus sect. Quercus) Shows Evidence of Climate-Associated
Asymmetric Gene Exchange. Journal of Heredity 112:663–670.

Oksanen, J., G. L. Simpson, F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, P.
Solymos, M. H. H. Stevens, E. Szoecs, H. Wagner, M. Barbour, M. Bedward, B. Bolker, D.
Borcard, G. Carvalho, M. Chirico, M. D. Caceres, S. Durand, H. B. A. Evangelista, R.
FitzJohn, M. Friendly, B. Furneaux, G. Hannigan, M. O. Hill, L. Lahti, D. McGlinn, M.-H.
Ouellette, E. R. Cunha, T. Smith, A. Stier, C. J. F. T. Braak, and J. Weedon. 2024. vegan:
Community Ecology Package.

O'Neill, G. A., A. Hamann, and T. Wang. 2008. Accounting for population variation improves
estimates of the impact of climate change on species' growth and distribution. Journal of

Applied Ecology 45:1040–1049.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annual
Review of Ecology, Evolution, and Systematics 37:637–669.

Patsiou, T. S., T. A. Shestakova, T. Klein, G. di Matteo, H. Sbay, M. R. Chambel, R. Zas, and J.
Voltas. 2020. Intraspecific responses to climate reveal nonintuitive warming impacts on a
widespread thermophilic conifer. New Phytologist 228:525–540.

Porth, I., and Y. A. El-Kassaby. 2015. Using Populus as a lignocellulosic feedstock for bioethanol.
Biotechnology Journal 10:510–524.

Putra, A. R., J. D. L. Yen, and A. Fournier-Level. 2023. Forecasting trait responses in novel
environments to aid seed provenancing under climate change. Molecular Ecology Resources
23:565–580.

R Core Team. 2024. R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rehfeldt, G. E., L. P. Leites, D. G. Joyce, and A. R. Weiskittel. 2018. Role of population genetics in
guiding ecological responses to climate. Global Change Biology 24:858–868.

Rogers, A. R., and J. B. Holland. 2021. Environment-specific genomic prediction ability in maize
using environmental covariates depends on environmental similarity to training data. G3
(Bethesda) 12:jkab440.

Sannigrahi, P., A. J. Ragauskas, and G. A. Tuskan. 2010. Poplar as a feedstock for biofuels: A review of
compositional characteristics. Biofuels, Bioproducts and Biorefining 4:209–226.

Sharma, S., R. Andrus, Y. Bergeron, M. Bogdziewicz, D. C. Bragg, D. Brockway, N. L. Cleavitt, B.
Courbaud, A. J. Das, M. Dietze, T. J. Fahey, J. F. Franklin, G. S. Gilbert, C. H. Greenberg, Q.
Guo, J. Hille Ris Lambers, I. Ibanez, J. F. Johnstone, C. L. Kilner, J. M. H. Knops, W. D.
Koenig, G. Kunstler, J. M. LaMontagne, D. Macias, E. Moran, J. A. Myers, R. Parmenter, I. S.
Pearse, R. Poulton-Kamakura, M. D. Redmond, C. D. Reid, K. C. Rodman, C. L. Scher, W.
H. Schlesinger, M. A. Steele, N. L. Stephenson, J. J. Swenson, M. Swift, T. T. Veblen, A. V.
Whipple, T. G. Whitham, A. P. Wion, C. W. Woodall, R. Zlotin, and J. S. Clark. 2022. North
American tree migration paced by climate in the West, lagging in the East. Proceedings of the
National Academy of Sciences 119:e2116691118. Proceedings of the National Academy of
Sciences.

Slavov, G. T., S. P. DiFazio, J. Martin, W. Schackwitz, W. Muchero, E. Rodgers-Melnick, M. F.
Lipphardt, C. P. Pennacchio, U. Hellsten, L. A. Pennacchio, L. E. Gunter, P. Ranjan, K.
Vining, K. R. Pomraning, L. J. Wilhelm, M. Pellegrini, T. C. Mockler, M. Freitag, A. Geraldes,
Y. A. El-Kassaby, S. D. Mansfield, Q. C. B. Cronk, C. J. Douglas, S. H. Strauss, D. Rokhsar,
and G. A. Tuskan. 2012. Genome resequencing reveals multiscale geographic structure and
extensive linkage disequilibrium in the forest tree Populus trichocarpa. New Phytologist
196:713–725.

Suarez-Gonzalez, A., C. A. Hefer, C. Christe, O. Corea, C. Lexer, Q. C. B. Cronk, and C. J. Douglas.
2016. Genomic and functional approaches reveal a case of adaptive introgression from
Populus balsamifera (balsam poplar) in P. trichocarpa (black cottonwood). Molecular Ecology
25:2427–2442.

Suarez-Gonzalez, A., C. A. Hefer, C. Lexer, Q. C. B. Cronk, and C. J. Douglas. 2018a. Scale and
984   direction of adaptive introgression between black cottonwood (Populus trichocarpa) and
985   balsam poplar (P. balsamifera). Molecular Ecology 27:1667–1680.

986   Suarez-Gonzalez, A., C. Lexer, and Q. C. B. Cronk. 2018b. Adaptive introgression: a plant
987   perspective. Biology Letters 14:20170688. Royal Society.

988   Taylor, S. A., E. L. Larson, and R. G. Harrison. 2015. Hybrid zones: windows on climate change.
989   Trends in Ecology & Evolution 30:398–406. Elsevier.

990   Taylor, S. A., T. A. White, W. M. Hochachka, V. Ferretti, R. L. Curry, and I. Lovette. 2014. Climate-
991   Mediated Movement of an Avian Hybrid Zone. Current Biology 24:671–676.

992   Van Nuland, M. E., J. B. Vincent, I. M. Ware, L. O. Mueller, S. L. J. Bayliss, K. K. Beals, J. A.
993   Schweitzer, and J. K. Bailey. 2020. Intraspecific trait variation across elevation predicts a
994   widespread tree species' climate niche and range limits. Ecology and Evolution 10:3856–3867.

995   VanWallendael, A., D. B. Lowry, and J. A. Hamilton. 2022. One hundred years into the study of
996   ecotypes, new advances are being made through large-scale field experiments in perennial plant
997   systems. Current Opinion in Plant Biology 66:102152.

998   Via, S., R. Gomulkiewicz, G. D. Jong, S. M. Scheiner, C. D. Schlichting, and P. H. V. Tienderen.
999   1995. Adaptive phenotypic plasticity: consensus and controversy. Trends in Ecology &
1000   Evolution 10:212–217. Elsevier.

1001   Wang, T., A. Hamann, D. Spittlehouse, and C. Carroll. 2016. Locally downscaled and spatially
1002   customizable climate data for historical and future periods for North America. PLOS ONE
1003   11:e0156720.

1004   Wang, T., A. Hamann, A. Yanchuk, G. A. O'neill, and S. N. Aitken. 2006. Use of response functions
1005   in selecting lodgepole pine populations for future climates. Global Change Biology 12:2404–
1006   2416.

1007   Wang, T., G. A. O'Neill, and S. N. Aitken. 2010. Integrating environmental and genetic effects to
1008   predict responses of tree populations to climate. Ecological Applications 20:153–163.

1009   Wielstra, B. 2019. Historical hybrid zone movement: More pervasive than appreciated. Journal of
1010   Biogeography 46:1300–1305.

1011   Woolbright, S. A., T. G. Whitham, C. A. Gehring, G. J. Allan, and J. K. Bailey. 2014. Climate relicts
1012   and their associated communities as natural ecology and evolution laboratories. Trends in
1013   Ecology & Evolution 29:406–416. Elsevier.

1014   Yang, N., Y. Wang, X. Liu, M. Jin, M. Vallebueno-Estrada, E. Calfee, L. Chen, B. P. Dilkes, S. Gui, X.
1015   Fan, T. K. Harper, D. J. Kennett, W. Li, Y. Lu, J. Ding, Z. Chen, J. Luo, S. Mambakkam, M.
1016   Menon, S. Snodgrass, C. Veller, S. Wu, S. Wu, L. Zhuo, Y. Xiao, X. Yang, M. C. Stitzer, D.
1017   Runcie, J. Yan, and J. Ross-Ibarra. 2023. Two teosintes made modern maize. Science
1018   382:eadg8940. American Association for the Advancement of Science.

1019   Ye, Z., G. A. O'Neill, and T. Wang. 2023. Optimization and validation of universal response functions
1020   for interior spruce (*Picea glauca*, *Picea engelmannii*, and their hybrids). Forest Ecology and
1021   Management 550:121509.

1022   Yeaman, S., K. A. Hodgins, K. E. Lotterhos, H. Suren, S. Nadeau, J. C. Degner, K. A. Nurkowski, P.
1023   Smets, T. Wang, L. K. Gray, K. J. Liepe, A. Hamann, J. A. Holliday, M. C. Whitlock, L. H.

Rieseberg, and S. N. Aitken. 2016. Convergent local adaptation to climate in distantly related
conifers. Science 353:1431–1433.
Zhang, M., H. Suren, and J. A. Holliday. 2019. Phenotypic and Genomic Local Adaptation across
Latitude and Altitude in Populus trichocarpa. Genome Biology and Evolution 11:2256–2272.
Zhou, L., R. Bawa, and J. A. Holliday. 2014. Exome resequencing reveals signatures of demographic
and adaptive processes across the genome and range of black cottonwood (Populus
trichocarpa). Molecular Ecology 23:2486–2499.