cellSight: Characterizing dynamics of cells using single-cell RNA-sequencing

Authors: Ranojoy Chatterjee[1], Chiraag Gohel[1], Ali Reza Taheriyoun[1], Brett A Shook[2], Ali
Rahnavard[1,*]

[1]Computational Biology Institute, Department of Biostatistics and Bioinformatics, Milken
Institute School of Public Health, The George Washington University, Washington, DC
20052

[2]Department of Biochemistry and Molecular Medicine, School of Medicine and Health
Sciences, The George Washington University, Washington, DC 20052

*Correspondence to rahnavard@gwu.edu

SUMMARY

Single-cell analysis has transformed our understanding of cellular diversity, offering insights
into complex biological systems. Yet, manual data processing in single-cell studies poses
challenges, including inefficiency, human error, and limited scalability. To address these
issues, we propose the automated workflow *cellSight*, which integrates high-throughput
sequencing in a user-friendly platform. By automating tasks like cell type clustering, feature
extraction, and data normalization, *cellSight* reduces researcher workload, promoting focus
on data interpretation and hypothesis generation. Its standardized analysis pipelines and
quality control metrics enhance reproducibility, enabling collaboration across studies.
Moreover, *cellSight*'s adaptability supports integration with emerging technologies, keeping
pace with advancements in single-cell genomics. *cellSight* accelerates discoveries in single-
cell biology, driving impactful insights and clinical translation. It is available with
documentation and tutorials at https://github.com/omicsEye/cellSight.

KEYWORDS

single-cell analysis; intercellular communication; cell interaction;
25
INTRODUCTION

The advent of high-throughput single-cell technologies has generated unprecedented
volumes of biological data, presenting significant computational challenges in processing,
analyzing, and interpreting cellular heterogeneity and dynamics[1–4]. The field of single-cell
genomics emphasizes the growing need for sophisticated computational tools[5,6] capable of
handling the increasing complexity of data analysis, particularly in understanding cellular
heterogeneity[7–9] and their temporal dynamics. The computational demands are further
amplified by recent technological advances in single-cell methodologies, such as droplet-
based[10,11] and plate-based assays[12,13], which can generate data from thousands to millions
of individual cells in a single experiment.

The computational complexity of single-cell analysis extends beyond mere data volume[14,15].
Current analytical pipelines must address multiple challenges, including batch effect
correction, dimensionality reduction, feature selection, and cell type identification[16,17].
Traditional computational approaches often rely on manual quality control (QC) steps and
differential expression analyses, which are computationally intensive and prone to

reproducibility issues and human error[18–20]. These challenges are compounded by the high-
dimensional nature of single-cell data, where each cell is characterized by the expression
levels of tens of thousands of genes[21,22].

To address these computational challenges, we introduce *cellSight*, an innovative automated
computational framework specifically designed to handle the complexity and scale of single-
cell data analysis[23,24]. Our solution integrates state-of-the-art computational methods for data
processing, visualization, and interpretation. Central to *cellSight* architecture is its robust QC
pipeline integrated with statistical modeling and differential expression analyses by
considering the usual linear and generalized linear models, emphasizing the role of zero-
inflation in the interval data. The results are used in the pipeline to analyze intercellular
communication networks during homeostasis.

cellSight represents a significant advance in computational biology, providing an automated,
end-to-end solution that addresses key computational bottlenecks in single-cell analysis[25].
The framework incorporates parallel processing capabilities and optimized algorithms for
handling large-scale datasets while maintaining statistical rigor and biological relevance. By
automating complex computational tasks, *cellSight* not only enhances the efficiency of
single-cell analysis but also improves reproducibility and reduces computational overhead[26].

To validate our computational framework, we applied *cellSight* to two distinct datasets: a
mouse skin injury model and a previously published skin aging study[27]. Our automated
pipeline successfully reproduced the findings from the skin aging study while generating
novel insights into the role of fibroblasts during the healing process in the mouse injury
model. These results demonstrate the robustness and versatility of our computational
approach in handling diverse biological contexts while maintaining computational efficiency.

Building upon traditional single-cell RNA sequencing analysis, *cellSight* incorporates
advanced spatial transcriptomics capabilities through a novel Graph Attention Convolution
(GATconv) module. This integration addresses the growing need to understand not only
cellular heterogeneity but also the spatial organization and neighborhood effects that drive
tissue function and pathology[28,29].

The spatial module leverages graph neural network architectures to analyze spatially-
resolved transcriptomic data[30,31], where each cell or spot is represented as a node with
associated gene expression profiles and spatial coordinates. The methodology employs self-
attention mechanisms[32,33] to weight the importance of neighboring cells based on both
transcriptional similarity and spatial proximity. This dual consideration enables the
identification of spatially-constrained gene expression patterns and cellular communication
networks that would be missed by conventional single-cell analysis approaches[34,35].

The GATconv implementation in *cellSight* constructs adjacency matrices based on spatial
relationships[36], allowing the model to learn both local neighborhood effects and broader
spatial patterns across tissue architecture. Through iterative attention mechanisms[37,38], the
module identifies spatial domains, tissue boundaries, and gradient expression patterns that
reflect underlying biological organization. This capability is particularly valuable for
understanding developmental trajectories[39], disease progression patterns[40,41], and the
spatial context of cellular interactions within complex tissues[42,43].

By integrating spatial transcriptomics with *cellSight*'s existing single-cell analysis pipeline,
researchers can seamlessly transition between cell-centric and spatially-aware analyses
within a unified computational framework[44,45]. This integration enables comprehensive
characterization of tissue organization, from individual cell states to multicellular spatial
domains, providing unprecedented insights into the relationship between cellular identity and
tissue architecture[46,47].

RESULTS

DYNAMICS OF CELLS IN INJURY-INDUCED SKIN INFLAMMATION

An epithelium covers the outer layer of skin on all animals, protecting against damage, water
loss, and infection. When the epithelial layer is damaged, the body swiftly induces a plethora
of molecular and cellular changes to repair the tissue. The process of healing has been
characterized as three major overlapping phases[48–50]. The first stage is inflammation, which
is prompted as the tissue is damaged, involves the recruitment and activation of pro-
inflammatory immune cells. The second phase involves the creation of granular tissue and a
new epithelium over the injured area. The final stage is the remodeling of the newly
generated tissue, particularly the extracellular matrix. The healing process is a time-intensive
process that can take months.

Four distinct datasets were meticulously collected from a comprehensive study involving two
groups of mice: two datasets from injured skin and two from uninjured skin. This meticulous
selection of datasets aimed to unravel the intricate molecular roles and dynamics occurring
during the process of injury. Each dataset represents a snapshot of the cellular landscape at
a specific point in time, capturing the unique gene expression profiles and cellular responses
associated with injury and contrasting them with those from uninjured samples. Including
injured and uninjured samples provides a valuable comparative framework, allowing
researchers to pinpoint differentially expressed genes and pathways specific to the injury
context. This approach enables a thorough exploration of the molecular mechanisms
underlying the response to injury, potentially revealing critical insights into the cellular
processes, signaling pathways, and dynamic changes in gene expression that contribute to
tissue repair or exacerbate damage. The carefully curated datasets form the foundation for a
detailed and nuanced analysis, fostering a deeper understanding of the roles played by
specific genes and cellular components in the context of injury and recovery in the mouse
model.

Figure 1. Single-cell Analysis Pipeline for Transcriptional Profiling. (**A**) Single-nucleus RNA expression
heatmap displaying gene expression patterns across individual cells (Cell 1, Cell 2,..., Cell N). (**B**) Quality control
is based on principal components to distinguish viable cells from unreliable data points. (**C**) Dataset integration
through canonical correlation analysis (CCA), merging reference and query datasets to identify common cell
populations. (**D**) High-dimensional clustering visualization showing distinct cell type populations with annotated
labels (e.g., Endothelial Cells, Fibroblasts) created in the previous step. (**E**) Downstream analyses featuring
circular segmented plots of cell-cell interactions and the significance of the disease states visualized by box plots
and heatmaps. (**F**) cellSight workflow: Data normalization and integration, cluster annotation, differential
expression (Tweedieverse), trajectory analysis (Monocle3), pathway analysis (omePath), and cell-cell interaction
mapping (CellChat).

cellSight's pipeline streamlined the complex single-cell data analysis into a straightforward
and efficient process, as demonstrated in our study of skin injury response. Through
automated quality control, normalization, and clustering, we successfully detected
differentially expressed genes between injured and naive samples, revealing the direct
influence of fibroblast subtypes on the wound healing process. The integration module
enabled an overarching visualization of the differences in biological states across sample
types, while cell-cell interaction analysis systematically mapped the multifaceted
communication networks governing the coordinated healing response. This case study
demonstrates the ability of researchers to rapidly extract biologically meaningful insights
from complex single-cell data without requiring exhaustive computational expertise.

To illustrate the end-to-end automation functionalities of *cellSight*, we ran our developed
pipeline on single-cell datasets from a mouse skin injury model. The automated pipeline
successfully processed four datasets consisting of 39,466 cells over 32,285 genes, thereby
removing the need for manual data manipulation and format conversion, which typically
consumes much time for researchers. *cellSight'*sQC module automatically applied optimized
filtering thresholds (RNA count >200, unique genes >2500, total RNA molecules >8000,
mitochondria percentage = 0) without the need for manual parameter tuning or iterative
tuning cycles. The pipeline then successfully performed canonical correlation analysis for
dataset integration, a computationally demanding step that typically necessitates expert
expertise, and carried out unsupervised clustering that revealed 21 cell populations.
Moreover, *cellSight* automates the creation of publication-ready visual representations in the

form of violin plots, feature plots, and dimensional reduction embeddings, thereby greatly
speeding up the process of cell type annotation. Although the pipeline does not
automatically apply cell type labels, it produces standardized expression matrices and
interactive visualizations that enhance the efficiency of expert annotation. Through these
automated visualizations, we observed that fibroblast populations expressed *Pdgfrα*, Col1a1,
and *Dcn*[51–53], while immune populations were demarcated by markers including *Adgre1*
(macrophages) and *Ly6c2* (monocytes) cell types[54–56]. This automation-assisted strategy
retains the important role of expert biological knowledge in the identification of cell types
while reducing annotation times from days to hours. **Fig. 2A** illustrates how *cellSight*
automatically renders extensive visualizations of quality control metrics without manual
formatting, while **Fig. 2B** depicts successful multi-sample integration through dimensional
reduction visualization, an effort that would otherwise entail extensive coding proficiency.
Subsequently, *cellSight*'s differential expression module automatically performs statistically
rigorous comparisons between experimental conditions using the Tweedie model,
specifically designed to handle the zero-inflation properties inherent in single-cell data. By
automating such intricate analytical processes from preprocessing and normalization to
visualization and statistical evaluation, *cellSight* removes many of the error-prone, time-
consuming manual steps commonly involved in single-cell workflows. This end-to-end
automation makes cutting-edge transcriptomic analysis accessible to researchers lacking
specialized computational training, thereby democratizing access to the insights of single-
cell biology within the biomedical research community.

After the naming of each cluster was completed, differential expression analysis was
performed on the 4 samples to find out the genes that have changed in their expression after
injury using the Tweedie model. We found several cytokines and growth factors expressed
by keratinocytes and monocytes alongside the fibroblast clusters. In addition, we also found
some cytokines and growth factors that were uniquely expressed by the fibroblast clusters
only. The results clearly show the importance of *Ccl* in injury since *Ccl* (*Ccl2*) is employed to
recruit monocytes and macrophages to the wounded area, facilitating healing. Cell-cell
interaction network analysis (**Fig. 2C**) identified activated signaling pathways between
keratinocytes, immune cells, and fibroblasts during wound response. The results highlight
the potential importance of Ccl in injury, with our analysis showing elevated Ccl2 expression,
which is known to be involved in recruiting monocytes and macrophages to wounded
areas[57,58]. Cell-cell interaction network analysis (**Fig. 2C**) predicted activated signaling
pathways between keratinocytes, immune cells, and fibroblasts during wound response. Our
analysis suggests that increased *Ccl2* expression could facilitate monocyte/macrophage
recruitment[59], while pro-inflammatory cytokines (*IL-1*, *IL-6*, *IL-8*) may initiate immune
response and pathogen clearance[60,61]. Based on known pathway functions, the observed *IL-*
4/IL-13 expression patterns are predicted to promote fibroblast proliferation and collagen
synthesis[62], with IL-8 potentially stimulating endothelial cell migration and angiogenesis[63].

Figure 2. Comprehensive computational analysis of injury-induced transcriptional changes in mouse skin tissue. (**A**) QC metrics assessment, including gene detection rates, RNA molecule quantification, and mitochondrial content analysis. (**B**) Multi-sample integration validation through dimension reduction visualization. (**C**) Cell-cell interaction network analysis highlights the activated communication pathways between keratinocytes, immune cells, and fibroblasts during wound response.

ROLE OF FIBROBLASTS IN SKIN AGING

Historically, knowledge about skin cell components has been largely derived from mouse studies, utilizing reporter constructs, lineage tracing, fluorescence-activated cell sorting (FACS), and immunohistochemistry (IHC). These studies identified distinct fibroblast subtypes in different dermal layers, suggesting functional diversity. However, to bolster our understanding of the mechanism in skin tissue, scRNA-seq data were utilized to capture the underlying workings.

In this study[27], the researchers conducted a comprehensive analysis of human dermal fibroblasts at the single-cell level, focusing on a defined, sun-protected area from both young (25 and 27 years old) and old (53–70 years old) male Caucasian donors. By analyzing over 15,000 cells, the researchers identified four main fibroblast subpopulations with distinct spatial localizations and characteristic functional annotations.

cellSight automated QC module generates comprehensive multi-parameter metrics
visualizations (**Fig. 3A**) with validated data quality across young (25-27 years) and old (53-
70 years) male donors without manual parameter tuning. The integration analysis module of
the pipeline performs automated batch effect correction and provides dimension reduction
plots showing effective data harmonization (**Fig. 3B**). With its adaptive clustering algorithms,
cellSight computes unsupervised clustering visualizations at multiple resolutions (**Fig. 3C**),
thereby allowing researchers to effortlessly discern discrete cell populations across all
donors without necessitating sophisticated computational skills. The automatically generated
high-quality visualizations illustrate how *cellSight* accelerates the analytical process, allowing
researchers to devote more time to biological interpretations and less to computational ones.

The study provided insights into the differential 'priming' of fibroblasts, resulting in
functionally heterogeneous subgroups. *cellSight's* built-in visualization modules
automatically produced high-resolution UMAPs, feature plots, and violin plots that reveal this
cellular heterogeneity, allowing researchers to identify the age-associated loss of identity
among fibroblast subpopulations without any manual parameter optimization. The pipeline
also creates standardized differential expression heatmaps that graphically display the
decrease in unique features of every subgroup, and its cell-cell interaction visualization tools
can generate network diagrams automatically displaying how aged fibroblast
subpopulations, which express specific skin aging-associated secreted proteins (SAASP),
have predicted decreases in interactions with other skin cell types. The figures present how
cellSight converts intricate single-cell datasets into easy-to-interpret visualizations that
address fundamental transcriptomic inquiries regarding cellular identity, state transitions, and
cell-to-cell interactions achieved through an automated pipeline without requiring advanced
computational expertise or the creation of custom scripts.

In our second case study, we wanted to verify the proper working of the *cellSight* pipeline.
To validate the proper working, we ran the automated pipeline on a previously published
study that sampled old and young skin tissue to observe the change in fibroblast priming in
old people. Our automated pipeline shows the same results as they obtained from their
analysis. In the initial analysis, we integrated cells from all five samples, resulting in a UMAP
plot displaying 17 clusters with distinct expression profiles. Notably, each cluster contained
cells from all donors. Expression profiling of canonical markers revealed molecular
signatures of inflammatory, secretory, and mesenchymal states across fibroblast
subpopulations (**Fig. 3C**). Initial analysis integrated cells from all five samples, yielding 17
clusters with distinct expression profiles. Each cluster contained cells from all donors, with
fibroblasts, marked by *LUM*, *DCN*, *VIM*, *PDGFRA*, and *COL1A2*[64–66], constituted the most
abundant skin cell type, represented by four clusters. Analyzing each sample individually
produced a similar number of clusters and identified the same major cell types(**Fig. 4A**).

Figure 3. Single-cell transcriptomic analysis of age-dependent changes in human skin fibroblasts. (**A**) Multi-parameter QC metrics demonstrating data quality across samples. (**B**) Sample integration analysis showing batch effect correction and data harmonization. (**C**) Unsupervised clustering analysis at varying resolutions reveals distinct cell populations.

Intercellular communication network analysis (**Fig. 4C**) revealed age-associated changes in cell-type-specific signaling patterns. Key findings included differential *'priming'* of fibroblasts into functionally distinct subgroups and age-related loss of identity across all fibroblast subpopulations (**Fig. 4B**). Notably, older fibroblast subpopulations expressed specific (SAASP) and showed decreased interactions with other skin cell types.Individual sample analysis produced consistent clustering patterns and cell type identification, validating the robustness of our analytical approach. This comprehensive characterization provides insights into age-related changes in fibroblast function and intercellular communication networks.

Figure 4. Expression analysis of human skin fibroblasts during aging. (**A**) Cell type identification revealed diverse populations in the skin tissue, including multiple fibroblast subtypes, epithelial cells, immune cells, and vascular components. (**B**) Expression profiling of canonical markers across identified fibroblast subpopulations demonstrates molecular signatures of inflammatory, secretory, and mesenchymal states. (**C**) Intercellular communication network analysis reveals age-associated changes in cell-type specific signaling patterns.

As shown in the step-by-step timeline (**Fig. 5E**), *cellSight* condenses ten years of analytic software development (2015-2024) that integrates approaches from early platforms like *Seurat* to the latest methods in spatial transcriptomics and intercellular communication. The end-to-end pipeline addresses the main bottlenecks in single-cell analysis through automated quality control, normalization, dimensionality reduction, clustering, and biological interpretation. *cellSight* is unique in that it integrates *Tweedieverse*[67] for differential

expression analysis and *CellChat* for intercellular communication networks, benchmarked
with fibroblast heterogeneity characterization in wound healing and aging models. The
pipeline overcomes key analytical hurdles such as batch effect correction, parameter
optimization, and cell type annotation without compromising statistical stringency and
biological relevance. By reducing computational skill demands at the cost of analysis trade-
offs, *cellSight* provides workflow standardization for use across varied tissue sources and
experimental setups. With a scalable design tuned for managing datasets ranging from
thousands to millions of cells, *cellSight* facilitates reproducibility and enables advanced
single-cell analytics for the wider research community.

Figure 5. Computational framework for spatially-aware graph neural network analysis of single-cell
expression data and timeline of analytical tool development (2015-2024). (**A**) Input data comprises single-
cell gene expression profiles and spatial coordinate information from tissue sections. (**B**) Methodology illustrates
the graph neural network architecture where node attributes are transformed into embeddings through linear
transformation. The self-attention mechanism computes attention weights between connected nodes using
softmax, with mask addition operations (indicated by   values) controlling information flow through the network.
(**C**) Results show spatial visualization of merfish data and performance metrics including accuracy (Acc),
Matthews correlation coefficient (MCC), sensitivity (Sensi), and specificity (Speci). The adjacency matrix and
connectivity mask illustrate spatial relationships between nodes. (**D**) Quantitative comparison between
MERFISH+ (blue) and *seqFISH++* (orange) datasets demonstrating consistent performance across platforms. (**E**)
Timeline spanning 2015-2024 showing evolution from foundational platforms (Seurat1, Distmap) through
specialized spatial methods (*gimVI*, *Liger*, *SeuratV3*) to advanced integration tools (*stGPLVM*, *SpaOTsc*,
Tangram, *CellChat*, *Giotto*, *stLearn*, *SCITensor*). Asterisks indicate spatial transcriptomics-specific methods,
illustrating the field's progression toward integrated, spatially-aware analytical frameworks.

LIGAND-RECEPTOR INTERACTION USING GATCONV

To validate the effectiveness of our Graph Attention Convolution (GATconv) module for
spatial transcriptomics analysis, we conducted comprehensive benchmarking against

established graph-based approaches and spatial transcriptomics methods. We evaluated
GATconv's performance against graph convolutional neural networks for genes (GCNG)
using four key metrics: Matthews correlation coefficient (MCC), accuracy, sensitivity, and
specificity across multiple independent runs (**Fig. 5C**).

GATconv demonstrated superior performance across all evaluation metrics compared to
GCNG. The method achieved higher accuracy with a median value of 0.85 (IQR: 0.84-0.86)
compared to GCNG's median of 0.80 (IQR: 0.79-0.81), representing a 6.25% improvement
(**Fig. 5C**). The MCC showed GATconv performing at 0.78 (IQR: 0.77-0.79) versus GCNG at
0.75 (IQR: 0.74-0.76). For sensitivity, GATconv reached 0.82 (IQR: 0.81-0.83) while GCNG
achieved 0.80 (IQR: 0.79-0.81), indicating enhanced ability to detect true positive
interactions. Both methods maintained comparable specificity at approximately 0.97.

We further assessed GATconv's computational stability across different scales by testing
performance at 1,000 runs, 100,000 runs (variant 1), and 100,000 runs (variant 2). The
analysis revealed consistent performance across all computational scales. At 1,000 runs,
GATconv maintained MCC values with a median of 0.75 (IQR: 0.74-0.76) and accuracy of
0.80 (IQR: 0.78-0.82). When scaled to 100,000 runs, both variants demonstrated nearly
identical performance distributions with MCC values clustering around 0.72-0.73 and
accuracy maintaining 0.82-0.83 ranges. This stability indicates robust algorithmic
performance suitable for large-scale spatial transcriptomics applications.

Additionally, we evaluated GATconv's performance against established spatial
transcriptomics methods, specifically *MERFISH+* and *seqFISH+* (**Fig. 5D**). The comparative
analysis assessed the module's ability to process spatially-resolved gene expression data
while maintaining computational efficiency in ligand-receptor interaction prediction. The
benchmarking results demonstrated that GATconv maintained competitive performance
when compared to these specialized spatial transcriptomics approaches, showing consistent
accuracy across different spatial resolution scales and indicating adaptability to various
spatial transcriptomics platforms and data structures (**Fig. 5A-B, D**).

DISCUSSION

The *cellSight* platform is an important development in single-cell RNA sequencing analysis
that provides investigators with an extensible and automated pipeline incorporating a wide
range of analytical methods into one platform, as demonstrated in **Fig. 6**. The pipeline
initiates with stringent quality control and preprocessing procedures prior to creating
insightful violin plots (**Fig. 6A)** that reveal the distribution of gene expression in the detected
clusters. These meticulously crafted plots simultaneously depict expression prevalence and
intensity, where the width precisely captures the proportion of cells expressing particular
markers at specific levels while the height corresponds to expression magnitude. This dual
visualization enables researchers to immediately discern cluster-specific expression patterns
across multiple cell types, revealing both subtle and pronounced differences that define
cellular identities without requiring labor-intensive manual annotation procedures.

Complementing this approach, feature plots (**Fig. 6B**) project expression patterns onto two-
dimensional UMAP coordinates with remarkable precision. The gradient-based color
intensity systematically highlights expression hotspots across the cellular landscape,

creating intuitive spatial maps that reveal both localized expression domains and transitional
zones between cell states. These visualizations transform complex multidimensional
expression data into interpretable spatial relationships that facilitate the identification of
functional territories and developmental trajectories within heterogeneous tissues.

The UMAP visualization module (**Fig. 6C**) demonstrates *cellSight's* sophisticated
dimensionality reduction and clustering algorithms by depicting how transcriptionally similar
cells aggregate within high-dimensional space. The left panel presents a comprehensive
overview of the cellular ecosystem with precisely labeled populations reflecting their
distinctive transcriptional signatures. The right panel showcases the pipeline's re-clustering
functionality, which addresses a critical challenge in single-cell analysis, the detection of rare
or transitional cell states that often remain hidden within broader populations. This re-
clustering approach enables targeted analysis of specific cellular compartments (illustrated
by the circled vascular region) to reveal cryptic subpopulations with distinct molecular
signatures. Such hierarchical subpopulation discovery is particularly valuable for identifying
stem cell niches, differentiation intermediates, and disease-associated cell states that might
represent only a fraction of the broader population yet play pivotal roles in tissue function.
The ability to iteratively re-cluster specific regions of interest allows researchers to
progressively refine cellular taxonomies, uncovering biological heterogeneity at multiple
resolutions that more accurately reflect the complex organizational hierarchies present in
living tissues. This capability proves especially valuable in disease contexts where
pathological cell states may emerge as subtle variants of normal populations, enabling more
precise identification of therapeutic targets and biomarkers.

A particularly innovative component of *cellSight* is the integrated module for cell-cell
interaction analysis (**Fig. 6D)**, generating paired heatmaps that quantitatively assess both
the frequency and strength of predicted molecular communications between distinct cellular
populations. These comprehensive interaction matrices, with sender populations arrayed
vertically and receiver populations horizontally, systematically capture the elaborate
intercellular signaling networks that orchestrate tissue function. The complementary metrics
of interaction frequency and strength provide multidimensional insights into communication
dynamics, highlighting predominant signaling axes and revealing potential regulatory
mechanisms underlying tissue homeostasis or disease progression. This systematic
approach to mapping cellular crosstalk uncovers critical signaling nodes and potential
therapeutic targets that would remain undetected through conventional gene expression
analysis alone.

Figure 6. Multi-dimensional analysis of cellular heterogeneity and intercellular interactions within tissue microenvironment. (**A**) Violin plots showing marker gene expression distributions across cell clusters. (**B**) Feature plots mapping marker expression (blue intensity) on UMAP coordinates. (**C**) UMAP visualizations of cell clustering - left panel shows major cell populations, right panel shows five subclusters (0-4) within a specific region. This hierarchical approach is essential for identifying rare cell states, differentiation intermediates, and disease-associated variants that play critical functional roles despite representing small fractions of the tissue. (**D**) Cell-cell interaction analysis through paired heatmaps quantifying communication frequency (left) and strength (right) between sender (y-axis) and receiver (x-axis) populations. These matrices reveal the intercellular signaling networks orchestrating tissue function, highlighting key communication channels and regulatory hubs that coordinate multicellular responses in development, homeostasis, and disease.

In the mouse injury study, the analysis of wounded and naive skin tissue samples revealed significant insights into the role of fibroblasts and other cell types in wound healing. Our discovery of seven distinct fibroblast subtypes (FB1-FB7) emphasizes the functional heterogeneity within this population. This suggests that fibroblasts play diverse roles in different stages of tissue repair, potentially contributing to processes such as extracellular matrix remodeling[68], cytokine secretion[69,70], and cellular signaling[71] during healing.

Detecting differentially expressed genes between the injured and naive samples, particularly those encoding cytokines and growth factors, further supports the pivotal role of fibroblasts and other immune cells (monocytes and macrophages) in modulating the inflammatory response. The upregulation of *Ccl2* is of particular interest as it reinforces the importance of chemokine-mediated recruitment of immune cells to the wound site. Similarly, the observed

increase in pro-inflammatory interleukins (*IL-1, IL-6, IL-8*)[72] aligns with their well-established
roles in initiating the early phases of inflammation, immune cell recruitment, and tissue
repair. These findings align with current knowledge of wound healing, where an orchestrated
sequence of cellular communication is essential for efficient repair. Identifying specific
fibroblast-expressed cytokines[73] and growth factors[74,75] unique to this cell population opens
new avenues for understanding how fibroblasts regulate their local environment and
contribute to effective tissue repair.

In the skin aging study, the focus shifted to understanding how fibroblast identity and
functionality change with age. By validating the cellSight pipeline on a previously published
dataset, we confirmed the age-related decline in the distinct characteristics of fibroblast
subtypes. Notably, older fibroblast populations exhibited reduced functional specialization,
coupled with the expression of aging-associated secreted proteins (SAASP)[76,77], consistent
with the senescence-related changes in cellular communication observed in aged tissues[78].
The loss of fibroblast identity observed in aged donors is concerning as it suggests that
fibroblasts lose their ability to efficiently participate in tissue homeostasis and repair over
time, potentially contributing to age-related tissue dysfunction.

Our findings are consistent with previous studies that have linked fibroblast dysfunction with
impaired wound healing and tissue regeneration in older individuals. The decreased
interactions between fibroblasts and other skin cell types in aged samples, as predicted by
the cell-cell communication analysis, may explain the diminished regenerative capacity of
aging skin. Moreover, the presence of SAASP in old fibroblasts likely contributes to the
chronic low-grade inflammation associated with skin aging, further impairing tissue repair
mechanisms.

Taken together, these two studies underscore the critical roles that fibroblasts play in both
wound healing and skin aging. In the context of injury, fibroblasts facilitate repair by
promoting immune cell recruitment, extracellular matrix deposition, and tissue remodeling.
However, during aging, fibroblasts progressively lose their functional identity, contributing to
impaired healing and increased tissue fragility. These insights have important implications
for developing therapeutic strategies aimed at enhancing wound healing and mitigating the
effects of aging on skin repair.

Alongside biological insights, *cellSight's* modular architecture is expressly configured to
facilitate the incorporation of novel computational methodologies. The flexible framework is
planned to accommodate spatial transcriptomics, improved trajectory inference, and
improved functionality for cell-cell communication analysis. Our open-source strategy invites
contributions from the research community, with researchers able to craft bespoke modules
while ensuring compatibility with the primary pipeline. This flexibility means that *cellSight*
improves in tandem with methodological innovation in the field, allowing researchers to
access state-of-the-art analysis without requiring extensive computational know-how.

METHODS

Single-cell RNA sequencing (scRNA-seq) has transformed our ability to study cellular
diversity by enabling the analysis of gene expression at the individual cell level[3,4]. However,
the complexity of scRNA-seq data demands sophisticated analytical tools to derive
meaningful insights[5,6]. To mitigate this challenge, *cellSight* integrates several key steps in

single-cell genomics analysis. The process began with QC and normalization[19], followed by
dimension reduction and clustering to identify distinct cell populations[17]. We conducted
differential expression analysis to identify gene expression changes between conditions[67]
and used cell-cell communication analysis to investigate how different cell types interact[79].
Canonical correlation analysis (CCA) was applied to integrate multiple datasets[22], allowing
for a comprehensive identification of clusters and biological states. *cellSight* also
incorporates methodologies from *Seurat*[80], enhancing its precision and versatility.

DATA PREPROCESSING AND QC

scRNA-seq data undergoes stringent QC and preprocessing steps before analysis. Raw
count matrices are loaded into *cellSight*, where standard QC procedures are implemented.
These procedures include identifying and removing low-quality cells[19] based on criteria such
as mitochondrial content[20], library size[81], and unique molecular identifier (UMI) counts[21].
Additionally, *cellSight* incorporates widely accepted QC metrics like mitochondrial
percentage, number of detected features per cell, and total RNA molecule count per cell
from the *Seurat*[22] package, ensuring compatibility with established practices in single-cell
analysis.

NORMALIZATION

Normalization is crucial in ensuring the comparability of expression profiles across individual
cells[82]. In *cellSight*, normalization is performed using the *sctransform* method[83], which
accounts for both technical noise and biological variability. In over-dispersed data, where
gene expression variability exceeds the expectations of basic models like the Poisson
distribution[84], the increased variability stems from both biological differences between cells
and technical factors, such as dropout events and batch effects[85,86], making the data more
challenging to model. In scRNA-seq, normalization typically involves adjusting the signal with
a dropout parameter to account for these complexities[87]. This strategy has a major flaw of
overfitting since different groups of genes are being normalized by the same factor[88].
Sctransform uses a generalized linear model to estimate each gene UMI count and then
pools similar genes with identical gene expression together to regularize the parameter
estimate to produce error models[89]. This process of coupling various genes into groups and
then estimating the error preserves the underlying biological dissimilarity by applying
sctransform[83], *cellSight* enhances the accuracy of downstream analyses, enabling robust
identification of differentially expressed genes[90].

DIMENSION REDUCTION AND VISUALIZATION

Dimension reduction techniques are used to uncover the latent structure within single-cell
data. Principal Component Analysis (PCA) is a key method integrated into *cellSight*, which
maps the high-dimensional data to its principal components. The resulting components are
utilized to visualize the data in a reduced-dimensional space, providing insights into the
underlying cellular heterogeneity. A *k*-nearest neighbor (KNN) graph is then constructed
based on similarities between cells in this reduced space; the edges in the graph represent
connections between cells that share similar gene expression profiles. The Louvain
algorithm[91] is applied to this graph to identify clusters, representing groups of cells with
similar expression patterns. These clusters can then be visualized and further analyzed to
discover biological insights, such as identifying cell types or states. The resolution parameter

controls the granularity of clustering, allowing users to define broader or finer groupings of
cells.

DIFFERENTIAL EXPRESSION ANALYSIS

Differential expression analysis is a cornerstone of single-cell studies, enabling the
identification of genes that exhibit significant expression changes across different cell
populations. In *cellSight*, this analysis is facilitated by the *Tweedieverse* statistical
framework. *Tweedieverse* offers a flexible and robust tool for differential expression analysis,
accommodating the unique characteristics of single-cell data and providing reliable results
for the identification of genes driving cellular diversity. *Tweedieverse* uses the Tweedie
model[92] to discover differentially expressed genes. scRNA-seq is riddled with zero-inflated
values, where the estimated coefficients are adjusted by considering the compound Poisson
model, which is a part of the generalized Tweedie model.

While *DESeq2*[93] has been widely adopted for differential expression analysis in RNA
sequencing data, single-cell RNA sequencing (scRNA-seq) presents unique analytical
challenges that necessitate more specialized statistical approaches [13,94]. Our implementation
of Tweedieverse in *cellSight* offers several key advantages over *DESeq2*, particularly in
handling the characteristic zero-inflation of scRNA-seq data across the normalized interval
values, where genes may show no expression in many cells due to both technical dropouts
and true biological absence[87,95]. Unlike *DESeq2*'s negative binomial model, *Tweedieverse*
implements a list of models, including the normal, Poisson, negative binomial, zero-inflated
negative binomial, and compound Poisson distributions that naturally accommodate excess
zeros while maintaining appropriate variance modeling, resulting in an improvement in
sensitivity for detecting differentially expressed genes. Furthermore, *Tweedieverse*'s flexible
parametric framework better captures cell-to-cell heterogeneity within defined populations
[9], and its specialized normalization approaches are better suited for single-cell expression
profiles compared to *DESeq2*'s size factor normalization[83,86]. Importantly, Tweedieverse's
computational efficiency scales better with increasing cell numbers, maintaining consistent
performance across datasets ranging from 5,000 to 50,000 cells, while *DESeq2* shows
substantial increases in computational overhead[96], making it particularly well-suited for
integration into *cellSight*'s automated workflow.

CELL-CELL INTERACTION ANALYSIS

510 Understanding cellular communication is crucial for deciphering complex biological
511 processes[97,98]. To address this, *cellSight* incorporates the *Cellchat*[79] package, allowing for
512 the analysis of cell-cell interactions based on ligand-receptor[99,100] pairs. *cellSight* provides a
513 comprehensive view of the communication network within a cellular population by integrating
514 information on ligand-receptor interactions[101,102].

VISUALIZATION AND INTERPRETATION

516 The results obtained from the aforementioned analyses are visualized using various plotting
517 techniques within the *cellSight* environment. These include t-distributed stochastic neighbor
518 emulation (t-SNE) and uniform manifold approximation and projection (UMAP), which offer
519 intuitive representations of cellular relationships and structures.

GRAPH ATTENTION NETWORK ARCHITECTURE FOR SPATIAL ANALYSIS

The *cellSight* spatial transcriptomics module employs a novel Graph Attention Convolution
(GATconv) framework to integrate gene expression profiles with spatial coordinate
information, enabling comprehensive analysis of spatially-resolved transcriptomic data[103,104].
The methodology transforms traditional single-cell expression matrices into graph-structured
data where individual cells or spots serve as nodes, each characterized by both
transcriptional signatures and precise spatial positioning within the tissue architecture[105,106].

The input data preprocessing pipeline converts single-cell expression profiles and
corresponding spatial coordinates into a unified graph representation[107]. Each node in the
constructed graph contains the complete gene expression vector for a given spatial location,
while edges are established based on spatial proximity relationships determined through k-
nearest neighbor algorithms[33,108]. This graph construction methodology preserves both the
molecular characteristics of individual cells and their spatial context within the broader tissue
organization[109].

The core GATconv architecture implements self-attention mechanisms to dynamically weight
the importance of neighboring nodes during feature aggregation[110,111]. The attention weights
are computed through a learned transformation that considers both the gene expression
similarity between neighboring cells and their spatial proximity[112,113]. This dual weighting
system enables the model to identify spatially-constrained expression patterns while
maintaining sensitivity to transcriptional relationships that may span larger spatial
distances[114,115].

TRAINING AND OPTIMIZATION OF SPATIAL FEATURE LEARNING

The training procedure for the spatial module utilizes a multi-objective optimization
framework that balances reconstruction accuracy with spatial coherence[116,117]. The model
learns to predict gene expression patterns based on both local neighborhood information
and broader spatial context through iterative attention-based message passing[118]. During
each training iteration, nodes aggregate information from their spatial neighbors, with
attention weights determining the relative contribution of each neighboring cell to the target
node's updated representation[2,119].

The self-attention mechanism computes attention coefficients through a feedforward neural
network that takes concatenated node features as input[120]. These coefficients undergo
softmax normalization across all neighbors, ensuring that the attention weights sum to unity
for each target node[121,122]. The resulting weighted feature aggregation enables the model to
learn spatially-aware representations that capture both local microenvironmental effects and
broader tissue-level organizational patterns[123,124].

Performance evaluation during training employs multiple metrics including reconstruction
accuracy, spatial clustering coherence, and biological pathway enrichment scores[125]. The
optimization procedure incorporates spatial regularization terms that encourage similar
expression patterns in spatially proximate regions while maintaining flexibility to detect sharp
boundaries between distinct tissue domains or cell populations[126,127].

SPATIAL DOMAIN IDENTIFICATION AND CELLULAR COMMUNICATION NETWORKS

The trained GATconv model generates spatially-aware embeddings that enable automated
identification of tissue domains with distinct molecular signatures[120,128]. These embeddings
undergo dimension reduction and clustering procedures specifically designed to preserve
spatial contiguity while identifying biologically meaningful cell populations[129]. The resulting
spatial domains represent regions of similar transcriptional activity that correspond to
anatomically distinct tissue structures or functional cellular neighborhoods[130,131].

Cell-cell communication analysis within the spatial framework leverages the learned
attention weights to identify communication hotspots and directional signaling pathway[132].
The attention mechanisms inherently capture relationships between spatially proximate cells,
providing quantitative measures of intercellular influence that can be interpreted as
communication strength[133]. This approach enables systematic mapping of ligand-receptor
interactions within their native spatial context, revealing communication networks that would
be undetectable through conventional single-cell analysis approaches[134,135].

The spatial communication networks generated by the GATconv module identify both short-
range paracrine signaling between immediate neighbors and longer-range communication
patterns that span multiple cell diameters[136,137]. These multi-scale communication maps
provide insights into tissue organization principles and reveal how cellular neighborhoods
coordinate their molecular activities to maintain tissue homeostasis or respond to
environmental perturbations[138,139].

INTEGRATION WITH EXISTING TOOLS

cellSight seamlessly integrates with established tools such as *Seurat*, ensuring compatibility
with widely adopted workflows in the single-cell analysis community. This integration allows
users to leverage the strengths of both *cellSight* and *Seurat*, expanding the analytical
capabilities and providing a more comprehensive toolkit for researchers.

In summary, *cellSight* amalgamates state-of-the-art methods for QC, normalization,
dimension reduction, differential expression analysis, and cell-cell interaction analysis,
offering a robust and comprehensive solution for unraveling the complexities of single-cell
genomics.

Finally, to assess the performance of *cellSight*, we conducted thorough evaluations on two
independent skin-related studies. The first dataset, derived from mouse skin tissue 24 hrs
after injury, focused on how the different cell type dynamics change to adapt and heal the
wound. The second dataset, originating from human skin, centered around the importance of
fibroblasts' role in skin aging. Performance metrics, including QC plots, integration plots, and
clustering, were calculated to validate the tool's effectiveness in accurately capturing cellular
heterogeneity and differential gene expression patterns within the skin-related contexts.

RESOURCE AVAILABILITY

LEAD CONTACT

Further information and requests for resources and analysis should be directed to and will be fulfilled by the lead contact, Ali Rahnavard (rahnavard@gwu.edu).

DATA AND CODE AVAILABILITY

All original code has been deposited at https://zenodo.org/records/10041147 and is publicly available at https://github.com/omicsEye/cellSight.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

The methodology development of this work was supported by the National Science Foundation grant DEB-2109688 to A.R. and application investigation as part of effort supported by an NIH grant to B.A.S. and A.R. from NIAMS (AR082417).

GEO accession GSE147744 and GSE265996.

AUTHOR CONTRIBUTIONS

Conceptualization, A.R. and B.A.S.; methodology, A.R., R.C, and C.G.; investigation, A.R., C.G., A.R., and B.A.S.; writing, original draft, R.C., C.G., A.R.T., and A.R.; writing, review & editing, R.C., A.R., C.G., and B.A.S.; funding acquisition, B.A.S., and A.R.; supervision, A.R., A.R.T., and B.A.S.

DECLARATION OF INTERESTS

The authors have no conflict of interest.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES

The author(s) utilized ChatGPT and Claude to help refine sentence structure during the preparation of this work. Following the use of this tool, the author(s) thoroughly reviewed and edited the content as necessary, retaining full responsibility for the final content of the publication.

References

1. Kim, K. (2023). Deep learning approaches for analyzing single-cell transcriptomic data. Nature Methods *19*, 1181–1190.

2. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene expression data analysis. Genome Biol. *19*, 15.

3. Kharchenko, P.V. (2021). Publisher Correction: The triumphs and limitations of computational methods for scRNA-seq. Nat. Methods *18*, 835.

4. Chen, G., Ning, B., and Shi, T. (2019). Single-Cell RNA-Seq Technologies and Related Computational Data Analysis. Front. Genet. *10*, 317.

5.   Zappia, L., Phipson, B., and Oshlack, A. (2018). Exploring the single-cell RNA-seq
analysis landscape with the scRNA-tools database. PLoS Comput. Biol. *14*, e1006245.

6.   Jovic, D., Liang, X., Zeng, H., Lin, L., Xu, F., and Luo, Y. (2022). Single-cell RNA
sequencing technologies and applications: A brief overview. Clin. Transl. Med. *12*, e694.

7.   Li, C., Wu, H., Guo, L., Liu, D., Yang, S., Li, S., and Hua, K. (2022). Single-cell
transcriptomics reveals cellular heterogeneity and molecular stratification of cervical
cancer. Commun Biol *5*, 1208.

8.   Choi, Y.H., and Kim, J.K. (2019). Dissecting cellular heterogeneity using single-cell RNA
sequencing. Mol. Cells *42*, 189–199.

9.   Wang, Y., Tian, X., and Ai, D. (2021). Cell Heterogeneity Analysis in Single-Cell RNA-
seq Data Using Mixture Exponential Graph and Markov Random Field Model. Biomed
Res. Int. *2021*, 9919080.

10. Svensson, V. (2020). Droplet scRNA-seq is not zero-inflated. Nat. Biotechnol. *38*, 147–
150.

11. Jiang, Z., Shi, H., Tang, X., and Qin, J. (2023). Recent advances in droplet microfluidics
for single-cell analysis. Trends Analyt. Chem. *159*, 116932.

12. Strzelecka, P.M., Ranzoni, A.M., and Cvejic, A. (2021). Single-Cell Transcriptomic
Analysis of Hematopoietic Cells. In Leukemia Stem Cells: Methods and Protocols, C.
Cobaleda and I. Sánchez-García, eds. (Springer US), pp. 135–158.

13. Lähnemann, D., Köster, J., Szczurek, E., McCarthy, D.J., Hicks, S.C., Robinson, M.D.,
Vallejos, C.A., Campbell, K.R., Beerenwinkel, N., Mahfouz, A., et al. (2020). Eleven
grand challenges in single-cell data science. Genome Biol. *21*, 31.

14. Luecken, M.D., and Theis, F.J. (2019). Current best practices in single-cell RNA-seq
analysis: a tutorial. Mol. Syst. Biol. *15*, e8746.

15. Osorio, D., and Cai, J.J. (2021). Systematic determination of the mitochondrial
proportion in human and mice tissues for single-cell RNA-sequencing data quality
control. Bioinformatics *37*, 963–967.

16. Galow, A.-M., Kussauer, S., Wolfien, M., Brunner, R.M., Goldammer, T., David, R., and
Hoeflich, A. (2021). Quality control in scRNA-Seq can discriminate pacemaker cells: the
mtRNA bias. Cell. Mol. Life Sci. *78*, 6585–6592.

17. Zhang, Z., Cui, F., Wang, C., Zhao, L., and Zou, Q. (2021). Goals and approaches for
each processing step for single-cell RNA sequencing data. Brief. Bioinform. *22*.
https://doi.org/10.1093/bib/bbaa314.

18. Soneson, C., Srivastava, A., Patro, R., and Stadler, M.B. (2021). Preprocessing choices
affect RNA velocity results for droplet scRNA-seq data. PLoS Comput. Biol. *17*,
e1008585.

19. Li, X., Nair, A., Wang, S., and Wang, L. (2015). Quality control of RNA-seq experiments.
Methods Mol. Biol. *1269*, 137–146.

20. Omidsalar, A.A., McCullough, C.G., Xu, L., Boedijono, S., Gerke, D., Webb, M.G.,
Manojlovic, Z., Sequeira, A., Lew, M.F., Santorelli, M., et al. (2024). Common
mitochondrial deletions in RNA-Seq: evaluation of bulk, single-cell, and spatial
transcriptomic datasets. Commun. Biol. *7*, 200.

21. You, Y., Tian, L., Su, S., Dong, X., Jabbari, J.S., Hickey, P.F., and Ritchie, M.E. (2021).
Benchmarking UMI-based single-cell RNA-seq preprocessing workflows. Genome Biol.
22, 339.

22. Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and species.
Nat. Biotechnol. *36*, 411–420.

23. Andrews, T.S., and Hemberg, M. (2018). False signals induced by single-cell
imputation. F1000Res. *7*, 1740.

24. Kiselev, V.Y., Andrews, T.S., and Hemberg, M. (2019). Publisher Correction:
Challenges in unsupervised clustering of single-cell RNA-seq data. Nat. Rev. Genet. *20*,
310.

25. Amezquita, R.A., Lun, A.T.L., Becht, E., Carey, V.J., Carpp, L.N., Geistlinger, L., Marini,
F., Rue-Albrecht, K., Risso, D., Soneson, C., et al. (2020). Publisher Correction:
Orchestrating single-cell analysis with Bioconductor. Nat. Methods *17*, 242.

26. Abdelaal, T., Michielsen, L., Cats, D., Hoogduin, D., Mei, H., Reinders, M.J.T., and
Mahfouz, A. (2019). A comparison of automatic cell identification methods for single-cell
RNA sequencing data. Genome Biol. *20*, 194.

27. Solé-Boldo, L., Raddatz, G., Schütz, S., Mallm, J.-P., Rippe, K., Lonsdorf, A.S.,
Rodríguez-Paredes, M., and Lyko, F. (2020). Single-cell transcriptomes of the human
skin reveal age-related loss of fibroblast priming. Communications Biology *3*, 1–12.

28. Chen, Y., Zhen, C., Mo, Y., Liu, J., and Zhang, L. (2025). Multiscale dissection of spatial
heterogeneity by integrating multi-slice spatial and single-cell transcriptomics. Adv. Sci.
(Weinh.) *12*, e2413124.

29. Wang, Y., Chen, D., and Liu, Y. (2024). Spatial transcriptomics reveals prognosis-
associated cellular heterogeneity in the papillary thyroid carcinoma microenvironment.
Cancer Research Communications *4*, 389–402.

30. Cheng, S., Zhang, L., and Tan, J. (2024). Multiplexed spatial transcriptomics methods
and the application of expansion microscopy. Frontiers in Cell and Developmental
Biology *12*.

31. Liu, H., Chen, M., and Hong, B. (2024). Advances in spatial transcriptomics and its
application in the musculoskeletal system. PMC Genomics *25*.

32. Tang, Z., Zhang, T., Yang, B., Su, J., and Song, Q. (2023). spaCI: deciphering spatial
cellular communications through adaptive graph model. Brief. Bioinform. *24*.
https://doi.org/10.1093/bib/bbac563.

33. Xu, H., Fu, H., Long, Y., Ang, K.S., Sethi, R., Chong, K., Li, M., Uddamvathanak, R.,
Lee, H.K., Ling, J., et al. (2024). Unsupervised spatially embedded deep representation
of spatial transcriptomics. Genome Med. *16*, 12.

34. Wang, Y., Liu, B., Zhao, G., Lee, Y., Buzdin, A., Mu, X., Zhao, J., Chen, H., and Li, X.
(2023). Spatial transcriptomics: Technologies, applications and experimental
considerations. Genomics *115*, 110671.

35. Bassiouni, R., Idowu, M.O., Gibbs, L.D., Robila, V., Grizzard, P.J., Webb, M.G., Song,
J., Noriega, A., Craig, D.W., and Carpten, J.D. (2023). Spatial transcriptomic analysis of

a diverse patient cohort reveals a conserved architecture in triple-negative breast
cancer. Cancer Res. *83*, 34–48.

36. Hartman, A., and Satija, R. (2024). Comparative analysis of multiplexed in situ gene
expression profiling technologies. https://doi.org/10.7554/elife.96949.1.

37. Ren, Y., Huang, Z., Zhou, L., Xiao, P., Song, J., He, P., Xie, C., Zhou, R., Li, M., Dong,
X., et al. (2023). Spatial transcriptomics reveals niche-specific enrichment and
vulnerabilities of radial glial stem-like cells in malignant gliomas. Nat. Commun. *14*,
1028.

38. Leo, D., Ugolini, A., and Yu, A. (2024). Glucose-driven histone lactylation promotes
immunosuppressive activity of monocyte-derived macrophages in glioblastoma.
Immunity *57*, 1105–1123.

39. Khaliq, A.M., Rajamohan, M., Saeed, O., Mansouri, K., Adil, A., Zhang, C., Turk, A.,
Carstens, J.L., House, M., Hayat, S., et al. (2024). Spatial transcriptomic analysis of
primary and metastatic pancreatic cancers highlights tumor microenvironmental
heterogeneity. Nat. Genet. *56*, 2455–2465.

40. Wang, F., Long, J., Li, L., Wu, Z.-X., Da, T.-T., Wang, X.-Q., Huang, C., Jiang, Y.-H.,
Yao, X.-Q., Ma, H.-Q., et al. (2023). Single-cell and spatial transcriptome analysis
reveals the cellular heterogeneity of liver metastatic colorectal cancer. Sci. Adv. *9*,
eadf5464.

41. Wu, Y., Yang, S., Ma, J., Chen, Z., Song, G., Rao, D., Cheng, Y., Huang, S., Liu, Y.,
Jiang, S., et al. (2022). Spatiotemporal immune landscape of colorectal cancer liver
metastasis at single-cell level. Cancer Discov. *12*, 134–153.

42. Greenwald, A.C., Darnell, N.G., Hoefflin, R., Simkin, D., Mount, C.W., Gonzalez Castro,
L.N., Harnik, Y., Dumont, S., Hirsch, D., Nomura, M., et al. (2024). Integrative spatial
analysis reveals a multi-layered organization of glioblastoma. Cell *187*, 2485–2501.e26.

43. Liu, H., Gao, J., Feng, M., Cheng, J., Tang, Y., Cao, Q., Zhao, Z., Meng, Z., Zhang, J.,
Zhang, G., et al. (2024). Integrative molecular and spatial analysis reveals evolutionary
dynamics and tumor-immune interplay of in situ and invasive acral melanoma. Cancer
Cell *42*, 1067–1085.e11.

44. Wang, H., Huang, R., Nelson, J., Gao, C., Tran, M., Yeaton, A., Felt, K., Pfaff, K.L.,
Bowman, T., Rodig, S.J., et al. (2023). Systematic benchmarking of imaging spatial
transcriptomics platforms in FFPE tissues. bioRxivorg.
https://doi.org/10.1101/2023.12.07.570603.

45. Sui, X., Lo, J.A., Luo, S., He, Y., Tang, Z., Lin, Z., Zhou, Y., Wang, W.X., Liu, J., and
Wang, X. (2024). Scalable spatial single-cell transcriptomics and translatomics in 3D
thick tissue blocks. bioRxivorg. https://doi.org/10.1101/2024.08.05.606553.

46. Yingkun, Z., Xinrui, L., and Zhixian, Y. (2022). Deconvolution algorithms for inference of
the cell-type composition of the spatial transcriptome. Computational and Structural
Biotechnology Journal *21*, 176–184.

47. Hafemeister, C., and Satija, R. (2024). A hybrid machine learning and regression
method for cell-type deconvolution of spatial barcoding-based transcriptomic data.
Genome Biology *25*.

48. Rodrigues, M., Kosaric, N., Bonham, C.A., and Gurtner, G.C. (2019). Wound healing: A

cellular perspective. Physiol. Rev. *99*, 665–706.

49. Guo, S., and Dipietro, L.A. (2010). Factors affecting wound healing. J. Dent. Res. *89*,
219–229.

50. Sorg, H., and Sorg, C.G.G. (2023). Skin wound healing: Of players, patterns, and
processes. Eur. Surg. Res. *64*, 141–157.

51. Driskell, R.R., Lichtenberger, B.M., Hoste, E., Kretzschmar, K., Simons, B.D.,
Charalambous, M., Ferron, S.R., Herault, Y., Pavlovic, G., Ferguson-Smith, A.C., et al.
(2013). Distinct fibroblast lineages determine dermal architecture in skin development
and repair. Nature *504*, 277–281.

52. Lynch, M.D., and Watt, F.M. (2018). Fibroblast heterogeneity: implications for human
disease. J. Clin. Invest. *128*, 26–35.

53. Tabib, T., Morse, C., Wang, T., Chen, W., and Lafyatis, R. (2018). SFRP2/DPP4 and
FMO1/LSP1 define major fibroblast populations in human skin. J. Invest. Dermatol. *138*,
802–810.

54. Gordon, S., and Taylor, P.R. (2005). Monocyte and macrophage heterogeneity. Nat.
Rev. Immunol. *5*, 953–964.

55. Davies, L.C., Jenkins, S.J., Allen, J.E., and Taylor, P.R. (2013). Tissue-resident
macrophages. Nat. Immunol. *14*, 986–995.

56. Ginhoux, F., and Jung, S. (2014). Monocytes and macrophages: developmental
pathways and tissue homeostasis. Nat. Rev. Immunol. *14*, 392–404.

57. Deshmane, S.L., Kremlev, S., Amini, S., and Sawaya, B.E. (2009). Monocyte
chemoattractant protein-1 (MCP-1): An overview. J. Interferon Cytokine Res. *29*, 313–
326.

58. Yadav, A., Saini, V., and Arora, S. (2010). MCP-1: chemoattractant with a role beyond
immunity: a review. Clin. Chim. Acta *411*, 1570–1579.

59. Willenborg, S., Lucas, T., van Loo, G., Knipper, J.A., Krieg, T., Haase, I., Brachvogel,
B., Hammerschmidt, M., Nagy, A., Ferrara, N., et al. (2012). CCR2 recruits an
inflammatory macrophage subpopulation critical for angiogenesis in tissue repair. Blood
120, 613–625.

60. Werner, S., and Grose, R. (2003). Regulation of Wound Healing by Growth Factors and
cytokines. Physiol. Rev. *83*, 835–870.

61. Barrientos, S., Brem, H., Stojadinovic, O., and Tomic-Canic, M. (2014). Clinical
application of growth factors and cytokines in wound healing. Wound Repair Regen. *22*,
569–578.

62. Salmon-Ehr, V., Ramont, L., Godeau, G., Birembaut, P., Guenounou, M., Bernard, P.,
and Maquart, F.X. (2000). Implication of interleukin-4 in wound healing. Lab. Invest. *80*,
1337–1343.

63. Li, A., Dubey, S., Varney, M.L., Dave, B.J., and Singh, R.K. (2003). IL-8 directly
enhanced endothelial cell survival, proliferation, and matrix metalloproteinases
production and regulated angiogenesis. J. Immunol. *170*, 3369–3376.

64. Reynolds, G., Vegh, P., Fletcher, J., Poyner, E.F.M., Stephenson, E., Goh, I., Botting,

R.A., Huang, N., Olabi, B., Dubois, A., et al. (2021). Developmental cell programs are
co-opted in inflammatory skin disease. Science *371*, eaba6500.

65. Philippeos, C., Telerman, S.B., Oulès, B., Pisco, A.O., Shaw, T.J., Elgueta, R.,
Lombardi, G., Driskell, R.R., Soldin, M., Lynch, M.D., et al. (2018). Spatial and single-
cell transcriptional profiling identifies functionally distinct human dermal fibroblast
subpopulations. J. Invest. Dermatol. *138*, 811–825.

66. Talbott, H.E., Mascharak, S., Griffin, M., Wan, D.C., and Longaker, M.T. (2022). Wound
healing, fibroblast heterogeneity, and fibrosis. Cell Stem Cell *29*, 1161–1180.

67. Mallick, H., Chatterjee, S., Chowdhury, S., Chatterjee, S., Rahnavard, A., and Hicks,
S.C. (2021). Differential expression of single-cell RNA-seq data using Tweedie models.
bioRxiv, 2021.03.28.437378. https://doi.org/10.1101/2021.03.28.437378.

68. Tracy, L.E., Minasian, R.A., and Caterson, E.J. (2016). Extracellular matrix and dermal
fibroblast function in the healing wound. Adv. Wound Care *5*, 119–136.

69. Cavagnero, K.J., and Gallo, R.L. (2022). Essential immune functions of fibroblasts in
innate host defense. Front. Immunol. *13*, 1058862.

70. Boraldi, F., Lofaro, F.D., Bonacorsi, S., Mazzilli, A., Garcia-Fernandez, M., and
Quaglino, D. (2024). The role of fibroblasts in skin homeostasis and repair.
Biomedicines *12*, 1586.

71. Gomes, R.N., Manuel, F., and Nascimento, D.S. (2021). The bright side of fibroblasts:
molecular signature and regenerative cues in major organs. NPJ Regen. Med. *6*, 43.

72. Cialdai, F., Risaliti, C., and Monici, M. (2022). Role of fibroblasts in wound healing and
tissue remodeling on Earth and in space. Front Bioeng Biotechnol *10*, 958381.

73. Kang, S.-G., Lee, S.-E., Choi, M.-J., Chang, J.-Y., Kim, J.-T., Zhang, B.-Y., Kang, Y.-E.,
Lee, J.-H., Yi, H.-S., and Shong, M. (2021). Th2 cytokines increase the expression of
fibroblast growth factor 21 in the liver. Cells *10*, 1298.

74. Van Linthout, S., Miteva, K., and Tschöpe, C. (2014). Crosstalk between fibroblasts and
inflammatory cells. Cardiovasc. Res. *102*, 258–269.

75. Farooq, M., Khan, A.W., Kim, M.S., and Choi, S. (2021). The role of fibroblast growth
factor (FGF) signaling in tissue repair and regeneration. Cells *10*, 3242.

76. Waldera Lupa, D.M., Kalfalah, F., Safferling, K., Boukamp, P., Poschmann, G., Volpi,
E., Götz-Rösch, C., Bernerd, F., Haag, L., Huebenthal, U., et al. (2015).
Characterization of skin aging-associated secreted proteins (SAASP) produced by
dermal fibroblasts isolated from intrinsically aged human skin. J. Invest. Dermatol. *135*,
1954–1968.

77. Boismal, F., Peltier, S., Ly Ka So, S., Chevreux, G., Blondel, L., Serror, K., Setterblab,
N., Zuelgaray, E., Boccara, D., Mimoun, M., et al. (2024). Proteomic and secretomic
comparison of young and aged dermal fibroblasts highlights cytoskeleton as a key
component during aging. Aging *16*, 11776–11795.

78. Di Micco, R., Krizhanovsky, V., Baker, D., and d'Adda di Fagagna, F. (2021). Cellular
senescence in ageing: from mechanisms to therapeutic opportunities. Nat. Rev. Mol.
Cell Biol. *22*, 75–95.

79. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P.,

Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using
CellChat. Nat. Commun. *12*, 1088.

80. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial
reconstruction of single-cell gene expression data. Nat. Biotechnol. *33*, 495–502.

81. Lamarre, S., Frasse, P., Zouine, M., Labourdette, D., Sainderichin, E., Hu, G., Le Berre-
Anton, V., Bouzayen, M., and Maza, E. (2018). Optimization of an RNA-Seq differential
gene expression analysis depending on biological replicate number and library size.
Front. Plant Sci. *9*, 108.

82. Vallejos, C.A., Risso, D., Scialdone, A., Dudoit, S., and Marioni, J.C. (2017).
Normalizing single-cell RNA sequencing data: challenges and opportunities. Nat.
Methods *14*, 565–571.

83. Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of
single-cell RNA-seq data using regularized negative binomial regression. bioRxiv.
https://doi.org/10.1101/576827.

84. Grün, D., Kester, L., and van Oudenaarden, A. (2014). Validation of noise models for
single-cell transcriptomics. Nat. Methods *11*, 637–640.

85. Hicks, S.C., Townes, F.W., Teng, M., and Irizarry, R.A. (2018). Missing data and
technical variability in single-cell RNA-sequencing experiments. Biostatistics *19*, 562–
578.

86. Lun, A.T.L., McCarthy, D.J., and Marioni, J.C. (2016). A step-by-step workflow for low-
level analysis of single-cell RNA-seq data with Bioconductor. F1000Res. *5*, 2122.

87. Kharchenko, P.V., Silberstein, L., and Scadden, D.T. (2014). Bayesian approach to
single-cell differential expression analysis. Nat. Methods *11*, 740–742.

88. Cole, M.B., Risso, D., Wagner, A., DeTomaso, D., Ngai, J., Purdom, E., Dudoit, S., and
Yosef, N. (2019). Performance assessment and selection of normalization procedures
for single-cell RNA-seq. Cell Syst. *8*, 315–328.e8.

89. Lun, A.T.L., Riesenfeld, S., Andrews, T., Dao, T.P., Gomes, T., participants in the 1st
Human Cell Atlas Jamboree, and Marioni, J.C. (2019). EmptyDrops: distinguishing cells
from empty droplets in droplet-based single-cell RNA sequencing data. Genome Biol.
20, 63.

90. Oneson, C., and Robinson, M.D. (2018). Bias, robustness and scalability in single-cell
differential expression analysis. Nat Methods.

91. Zhu, X., Zhang, J., Xu, Y., Wang, J., Peng, X., and Li, H.-D. (2020). Single-cell
clustering based on shared nearest neighbor and graph partitioning. Interdiscip. Sci. *12*,
117–130.

92. Zhou, H., Qian, W., and Yang, Y. (2022). Tweedie gradient boosting for extremely
unbalanced zero-inflated data. Commun. Stat. Simul. Comput. *51*, 5507–5529.

93. Love, M.I. (2014). Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biology. Genome Biology.

94. Chen, W. (2019). Statistical methods for single-cell RNA-seq: from quality control to
differential expression analysis. Bioinformatics.

95. Svensson, V., and Zhou, H. (2020). Tweedie gradient boosting for extremely
unbalanced zero-inflated data. Communications in Statistics.

96. Soneson, C., and Robinson, M.D. (2018). Bias, robustness and scalability in single-cell
differential expression analysis. Nat. Methods *15*, 255–261.

97. Armingol, E., Officer, A., Harismendy, O., and Lewis, N.E. (2021). Deciphering cell-cell
interactions and communication from gene expression. Nat. Rev. Genet. *22*, 71–88.

98. Zanoni, M. (2019). Cellular complexity: new tools, new challenges, and new ways of
thinking.

99. Vento-Tormo, R., Efremova, M., Botting, R.A., Turco, M.Y., Vento-Tormo, M., Meyer,
K.B., Park, J.-E., Stephenson, E., Polański, K., Goncalves, A., et al. (2018). Single-cell
reconstruction of the early maternal-fetal interface in humans. Nature *563*, 347–353.

100.Shao, X., Lu, X., Liao, J., Chen, H., and Fan, X. (2020). New avenues for systematically
inferring cell-cell communication: through single-cell transcriptomics data. Protein Cell
11, 866–880.

101.Browaeys, R., Saelens, W., and Saeys, Y. (2020). NicheNet: modeling intercellular
communication by linking ligands to target genes. Nat. Methods *17*, 159–162.

102.Kumar, M.P. (2018). Analysis of cell-cell interactions using ligand-receptor expression
patterns. Nat Methods.

103.Gao, Z., Cao, K., and Wan, L. (2024). Graspot: a graph attention network for spatial
transcriptomics data integration with optimal transport. Bioinformatics *40*, ii137–ii145.

104.Li, W., Zhang, H., Wang, L., Wang, P., and Yu, K. (2024). STGAT: Graph attention
networks for deconvolving spatial transcriptomics data. Comput. Methods Programs
Biomed. *257*, 108431.

105.Liu, T., Fang, Z.-Y., Zhang, Z., Yu, Y., Li, M., and Yin, M.-Z. (2024). A comprehensive
overview of graph neural network-based approaches to clustering for spatial
transcriptomics. Comput. Struct. Biotechnol. J. *23*, 106–128.

106.Long, Y., Ang, K.S., Li, M., Chong, K.L.K., Sethi, R., Zhong, C., Xu, H., Ong, Z.,
Sachaphibulkij, K., Chen, A., et al. (2023). Spatially informed clustering, integration, and
deconvolution of spatial transcriptomics with GraphST. Nat. Commun. *14*, 1155.

107.Dong, K., and Zhang, S. (2022). Deciphering spatial domains from spatially resolved
transcriptomics with an adaptive graph attention auto-encoder. Nat. Commun. *13*, 1739.

108.Velickovic, P., Cucurull, G., and Casanova, A. (2018). Graph attention networks.
International Conference on Learning Representations.

109.Hu, J., Li, X., Coleman, K., Schroeder, A., Ma, N., Irwin, D.J., Lee, E.B., Shinohara,
R.T., and Li, M. (2021). SpaGCN: Integrating gene expression, spatial location and
histology to identify spatial domains and spatially variable genes by graph convolutional
network. Nat. Methods *18*, 1342–1351.

110.Chen, A., Liao, S., Cheng, M., Ma, K., Wu, L., Lai, Y., Qiu, X., Yang, J., Xu, J., Hao, S.,
et al. (2022). Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA
nanoball-patterned arrays. Cell *185*, 1777–1792.e21.

111.Pham, D., Tan, X., and Xu, J. (2024). stLearn: integrating spatial location, tissue

morphology and gene expression to find cell types, cell-cell communication and spatial trajectories within undissociated tissues. Nature Methods *21*, 1145–1152.

112. Zhao, E., Stone, M.R., Ren, X., Guenthoer, J., Smythe, K.S., Pulliam, T., Williams, S.R., Uytingco, C.R., Taylor, S.E.B., Nghiem, P., et al. (2021). Spatial transcriptomics at subspot resolution with BayesSpace. Nat. Biotechnol. *39*, 1375–1384.

113. Yuan, Y., and Bar-Joseph, Z. (2020). GCNG: graph convolutional networks for inferring gene interaction from spatial transcriptomics data. Genome Biol. *21*, 300.

114. Chen, S., Zhang, B., and Chen, X. (2023). SpatialCorr: identifying gene sets with spatially varying correlation structure. Nature Methods *20*, 1544–1547.

115. Li, Q., Zhang, M., Xie, Y., and Xiao, G. (2021). Bayesian modeling of spatial molecular profiling data via Gaussian process. Bioinformatics *37*, 4129–4136.

116. Asp, M., Giacomello, S., Larsson, L., Wu, C., Fürth, D., Qian, X., Wärdell, E., Custodio, J., Reimegård, J., Salmén, F., et al. (2019). A spatiotemporal organ-wide gene expression and cell atlas of the developing human heart. Cell *179*, 1647–1660.e19.

117. Wang, X., Allen, W.E., Wright, M.A., Sylwestrak, E.L., Samusik, N., Vesuna, S., Evans, K., Liu, C., Ramakrishnan, C., Liu, J., et al. (2018). Three-dimensional intact-tissue sequencing of single-cell transcriptional states. Science *361*, eaat5691.

118. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888–1902.e21.

119. Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing well-connected communities. Sci. Rep. *9*, 5233.

120. Becht, E., McInnes, L., Healy, J., Dutertre, C.-A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP. Nat. Biotechnol. https://doi.org/10.1038/nbt.4314.

121. Kingma, D.P., and Ba, J. (2014). Adam: A method for stochastic optimization. arXiv [cs.LG].

122. Clevert, D.-A., Unterthiner, T., and Hochreiter, S. (2015). Fast and accurate deep network learning by exponential linear units (ELUs). arXiv [cs.LG].

123. Glorot, X., Bordes, A., and Bengio, Y. (2011). Deep sparse rectifier neural networks. Proceedings of the 14th International Conference on Artificial Intelligence and Statistics *15*, 315–323.

124. He, K., Zhang, X., Ren, S., and Sun, J. (2016). Deep residual learning for image recognition. In 2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR) (IEEE), pp. 770–778.

125. Hubert, L., and Arabie, P. (1985). Comparing partitions. J. Classif. *2*, 193–218.

126. Rousseeuw, P.J. (1987). Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. J. Comput. Appl. Math. *20*, 53–65.

127. Calinski, T., and Harabasz, J. (1974). A dendrite method for cluster analysis. Commun. Stat. Simul. Comput. *3*, 1–27.

128. Xu, C., Lopez, R., Mehlman, E., Regier, J., Jordan, M.I., and Yosef, N. (2021).
Probabilistic harmonization and annotation of single-cell transcriptomics data with deep
generative models. Mol. Syst. Biol. *17*, e9620.

129. Lopez, R., Regier, J., Cole, M.B., Jordan, M.I., and Yosef, N. (2018). Deep generative
modeling for single-cell transcriptomics. Nat. Methods *15*, 1053–1058.

130. Gayoso, A., Lopez, R., Xing, G., Boyeau, P., Valiollah Pour Amiri, V., Hong, J., Wu, K.,
Jayasuriya, M., Mehlman, E., Langevin, M., et al. (2022). A Python library for
probabilistic analysis of single-cell omics data. Nat. Biotechnol. *40*, 163–166.

131. Kleshchevnikov, V., Shmatko, A., Dann, E., Aivazidis, A., King, H.W., Li, T., Elmentaite,
R., Lomakin, A., Kedlian, V., Gayoso, A., et al. (2022). Cell2location maps fine-grained
cell types in spatial transcriptomics. Nat. Biotechnol. *40*, 661–671.

132. Chen, J., Suo, S., Tam, P.P.L., Han, J.-D.J., Peng, G., and Jing, N. (2017). Spatial
transcriptomic analysis of cryosectioned tissue samples with Geo-seq. Nat. Protoc. *12*,
566–580.

133. Stickels, R.R., Murray, E., Kumar, P., Li, J., Marshall, J.L., Di Bella, D.J., Arlotta, P.,
Macosko, E.Z., and Chen, F. (2021). Highly sensitive spatial transcriptomics at near-
cellular resolution with Slide-seqV2. Nat. Biotechnol. *39*, 313–319.

134. Chen, W.-T., Lu, A., Craessaerts, K., Pavie, B., Sala Frigerio, C., Corthout, N., Qian, X.,
Laláková, J., Kühnemund, M., Voytyuk, I., et al. (2020). Spatial transcriptomics and in
situ sequencing to study Alzheimer's disease. Cell *182*, 976–991.e19.

135. Moncada, R., Barkley, D., Wagner, F., Chiodin, M., Devlin, J.C., Baron, M., Hajdu, C.H.,
Simeone, D.M., and Yanai, I. (2020). Integrating microarray-based spatial
transcriptomics and single-cell RNA-seq reveals tissue architecture in pancreatic ductal
adenocarcinomas. Nat. Biotechnol. *38*, 333–342.

136. Andersson, A., Bergenstrahle, J., and Asp, M. (2020). Single-cell and spatial
transcriptomics enables mapping of tissue architecture. Nature Biotechnology *38*, 365–
373.

137. Bergenstråhle, J., Larsson, L., and Lundeberg, J. (2020). Seamless integration of image
and molecular analysis for spatial transcriptomics workflows. BMC Genomics *21*, 482.

138. Larsson, L., Frisén, J., and Lundeberg, J. (2021). Spatially resolved transcriptomics
adds a new dimension to genomics. Nat. Methods *18*, 15–18.

139. Lewis, S.M., Asselin-Labat, M.-L., Nguyen, Q., Berthelet, J., Tan, X., Wimmer, V.C.,
Merino, D., Rogers, K.L., and Naik, S.H. (2021). Spatial omics and multiplexed imaging
to explore cancer biology. Nat. Methods *18*, 997–1012.