***cellSight*: Characterizing dynamics of cells using single-cell RNA-sequencing**

Authors: Ranojoy Chatterjee[1], Chiraag Gohel[1], Brett A Shook[2], Ali Rahnavard[1,*]

[1]Computational Biology Institute, Department of Biostatistics and Bioinformatics, Milken
Institute School of Public Health, The George Washington University, Washington, DC 20052

[2]Department of Biochemistry and Molecular Medicine, School of Medicine and Health
Sciences, The George Washington University, Washington, DC 20052

*Correspondence to rahnavard@gwu.edu

SUMMARY

Single-cell analysis has transformed our understanding of cellular diversity, offering insights
into complex biological systems. Yet, manual data processing in single-cell studies poses
challenges, including inefficiency, human error, and limited scalability. To address these
issues, we propose the automated workflow *cellSight*, which integrates high-throughput
sequencing in a user-friendly platform. By automating tasks like cell type clustering, feature
extraction, and data normalization, *cellSight* reduces researcher workload, promoting focus
on data interpretation and hypothesis generation. Its standardized analysis pipelines and
quality control metrics enhance reproducibility, enabling collaboration across studies.
Moreover, *cellSight*'s adaptability supports integration with emerging technologies, keeping
pace with advancements in single-cell genomics. *cellSight* accelerates discoveries in single-
cell biology, driving impactful insights and clinical translation. It is available with documentation
and tutorials at https://github.com/omicsEye/cellSight.

KEYWORDS

Single-cell analysis; Intercellular communication; Cell interaction; Software; Bioinformatics
23

INTRODUCTION

The advent of high-throughput single-cell technologies has generated unprecedented volumes
of biological data, presenting significant computational challenges in processing, analyzing,
and interpreting cellular heterogeneity and dynamics[1–4]. The field of single-cell genomics
emphasizes the growing need for sophisticated computational tools[5,6] capable of handling the
increasing complexity of data analysis, particularly in understanding cellular heterogeneity[7–9]
and their temporal dynamics. The computational demands are further amplified by recent
technological advances in single-cell methodologies, such as droplet-based[10,11] and plate-
based assays[12,13], which can generate data from thousands to millions of individual cells in a
single experiment.

The computational complexity of single-cell analysis extends beyond mere data volume[14,15].
Current analytical pipelines must address multiple challenges, including batch effect
correction, dimensionality reduction, feature selection, and cell type identification[16,17].
Traditional computational approaches often rely on manual quality control(QC) steps and
differential expression analyses, which are computationally intensive and prone to
reproducibility issues and human error[18–20]. These challenges are compounded by the high-

dimensional nature of single-cell data, where each cell is characterized by the expression
levels of tens of thousands of genes[21,22].

To address these computational challenges, we introduce *cellSight*, an innovative automated
computational framework specifically designed to handle the complexity and scale of single-
cell data analysis[23,24]. Our solution integrates state-of-the-art computational methods for data
processing, visualization, and interpretation. Central to *cellSight's* architecture is its robust QC
pipeline integrated with statistical modeling and differential expression analyses by
considering the usual linear and generalized linear models, emphasizing the role of zero-
inflation in the interval data. The results are used in the pipeline to analyze intercellular
communication networks during homeostasis.

cellSight represents a significant advance in computational biology, providing an automated,
end-to-end solution that addresses key computational bottlenecks in single-cell analysis[25].
The framework incorporates parallel processing capabilities and optimized algorithms for
handling large-scale datasets while maintaining statistical rigor and biological relevance. By
automating complex computational tasks, *cellSight* not only enhances the efficiency of single-
cell analysis but also improves reproducibility and reduces computational overhead[26].

To validate our computational framework, we applied *cellSight* to two distinct datasets: a
mouse skin injury model and a previously published skin aging study[27]. Our automated
pipeline successfully reproduced the findings from the skin aging study while generating novel
insights into the role of fibroblasts during the healing process in the mouse injury model. These
results demonstrate the robustness and versatility of our computational approach in handling
diverse biological contexts while maintaining computational efficiency.

62 RESULTS

63 DYNAMICS OF CELLS IN INJURY-INDUCED SKIN INFLAMMATION

An epithelium covers the outer layer of skin on all animals, protecting against damage, water
loss, and infection. When the epithelial layer is damaged, the body swiftly induces a plethora
of molecular and cellular changes to repair the tissue. The process of healing has been
characterized as three major overlapping phases[28–30]. The first stage is inflammation, which is
prompted as the tissue is damaged, involves the recruitment and activation of pro-
inflammatory immune cells. The second phase involves the creation of granular tissue and a
new epithelium over the injured area. The final stage is the remodeling of the newly generated
tissue, particularly the extracellular matrix. The healing process is a time-intensive process
that can take months.

Four distinct datasets were meticulously collected from a comprehensive study involving two
groups of mice: two datasets from injured skin and two from uninjured skin. This meticulous
selection of datasets aimed to unravel the intricate molecular roles and dynamics occurring
during the process of injury. Each dataset represents a snapshot of the cellular landscape at
a specific point in time, capturing the unique gene expression profiles and cellular responses
associated with injury and contrasting them with those from uninjured samples. Including
injured and uninjured samples provides a valuable comparative framework, allowing
researchers to pinpoint differentially expressed genes and pathways specific to the injury
context. This approach enables a thorough exploration of the molecular mechanisms

underlying the response to injury, potentially revealing critical insights into the cellular
processes, signaling pathways, and dynamic changes in gene expression that contribute to
tissue repair or exacerbate damage. The carefully curated datasets form the foundation for a
detailed and nuanced analysis, fostering a deeper understanding of the roles played by
specific genes and cellular components in the context of injury and recovery in the mouse
model.

Figure 1. Single-cell Analysis Pipeline for Transcriptional Profiling. (**A**) Single-nucleus RNA expression
heatmap displaying gene expression patterns across individual cells (Cell 1, Cell 2,..., Cell N). (**B**) Quality control
is based on principal components to distinguish viable cells from unreliable data points. (**C**) Dataset integration
through canonical correlation analysis (CCA), merging reference and query datasets to identify common cell
populations. (**D**) High-dimensional clustering visualization showing distinct cell type populations with annotated
labels (e.g., Endothelial Cells, Fibroblasts) created in the previous step. (**E**) Downstream analyses featuring circular
segmented plots of cell-cell interactions and the significance of the disease states visualized by box plots and
heatmaps. (**F**) cellSight workflow: Data normalization and integration, cluster annotation, differential expression
(Tweedieverse), trajectory analysis (Monocle3), pathway analysis (omePath), and cell-cell interaction mapping
(CellChat).

cellSight's pipeline streamlined the complex single-cell data analysis into a straightforward
and efficient process, as demonstrated in our study of skin injury response. Through
automated quality control, normalization, and clustering, we successfully detected
differentially expressed genes between injured and naive samples, revealing the direct
influence of fibroblast subtypes on the wound healing process. The integration module
enabled an overarching visualization of the differences in biological states across sample
types, while cell-cell interaction analysis systematically mapped the multifaceted
communication networks governing the coordinated healing response. This case study
demonstrates the ability of researchers to rapidly extract biologically meaningful insights from
complex single-cell data without requiring exhaustive computational expertise.

To illustrate the end-to-end automation functionalities of cellSight, we ran our developed
pipeline on single-cell datasets from a mouse skin injury model. The automated pipeline
successfully processed four datasets consisting of 39,466 cells over 32,285 genes, thereby
removing the need for manual data manipulation and format conversion, which typically

consumes much time for researchers. cellSight's quality control module automatically applied
optimized filtering thresholds (RNA count >200, unique genes >2500, total RNA molecules
>8000, mitochondria percentage = 0) without the need for manual parameter tuning or iterative
tuning cycles. The pipeline then successfully performed canonical correlation analysis for
dataset integration—a computationally demanding step that typically necessitates expert
expertise—and carried out unsupervised clustering that revealed 21 cell populations.
Moreover, cellSight automates the creation of publication-ready visual representations in the
form of violin plots, feature plots, and dimensional reduction embeddings, thereby greatly
speeding up the process of cell type annotation. Although the pipeline does not automatically
apply cell type labels, it produces standardized expression matrices and interactive
visualizations that enhance the efficiency of expert annotation. Through these automated
visualizations, we observed that fibroblast populations expressed *Pdgfra*, Col1a1, and *Dcn*[31–
33], while immune populations were demarcated by markers including *Adgre1* (macrophages)
and *Ly6c2* (monocytes) cell types[34–36]. This automation-assisted strategy retains the important
role of expert biological knowledge in the identification of cell types while reducing annotation
times from days to hours. Figure 2A illustrates how cellSight automatically renders extensive
visualizations of quality control metrics without manual formatting, while (**Fig. 2B**) depicts
successful multi-sample integration through dimensional reduction visualization—an effort that
would otherwise entail extensive coding proficiency. Subsequently, cellSight's differential
expression module automatically performs statistically rigorous comparisons between
experimental conditions using the Tweedie model, specifically designed to handle the zero-
inflation properties inherent in single-cell data. By automating such intricate analytical
processes—from preprocessing and normalization to visualization and statistical evaluation—
cellSight removes many of the error-prone, time-consuming manual steps commonly involved
in single-cell workflows. This end-to-end automation makes cutting-edge transcriptomic
analysis accessible to researchers lacking specialized computational training, thereby
democratizing access to the insights of single-cell biology within the biomedical research
community.

After the naming of each cluster was completed, differential expression(D.E.) analysis was
performed on the 4 samples to find out the genes that have changed in their expression after
injury using the Tweedie model. We found several cytokines and growth factors expressed by
keratinocytes and monocytes alongside the fibroblast clusters. In addition, we also found some
cytokines and growth factors that were uniquely expressed by the fibroblast clusters only. The
results clearly show the importance of *Ccl* in injury since *Ccl*(*Ccl2*) is employed to recruit
monocytes and macrophages to the wounded area, facilitating healing. Cell-cell interaction
network analysis (**Fig. 2C**) identified activated signaling pathways between keratinocytes,
immune cells, and fibroblasts during wound response. The results highlight the potential
importance of Ccl in injury, with our analysis showing elevated Ccl2 expression, which is
known to be involved in recruiting monocytes and macrophages to wounded areas[37,38]. Cell-
cell interaction network analysis (**Fig. 2C**) predicted activated signaling pathways between
keratinocytes, immune cells, and fibroblasts during wound response. Our analysis suggests
that increased *Ccl2* expression could facilitate monocyte/macrophage recruitment[39], while pro-
inflammatory cytokines (*IL-1*, *IL-6*, *IL-8*) may initiate immune response and pathogen
clearance[40,41]. Based on known pathway functions, the observed *IL-4/IL-13* expression
patterns are predicted to promote fibroblast proliferation and collagen synthesis[42], with IL-8
potentially stimulating endothelial cell migration and angiogenesis[43].

Figure 2. Comprehensive computational analysis of injury-induced transcriptional changes in mouse skin tissue. (A) Quality control metrics assessment, including gene detection rates, RNA molecule quantification, and mitochondrial content analysis. (B) Multi-sample integration validation through dimensional reduction visualization. (C) Cell-cell interaction network analysis highlights the activated communication pathways between keratinocytes, immune cells, and fibroblasts during wound response.

ROLE OF FIBROBLASTS IN SKIN AGING

Historically, knowledge about skin cell components has been largely derived from mouse studies, utilizing reporter constructs, lineage tracing, fluorescence-activated cell sorting (FACS), and immunohistochemistry (IHC). These studies identified distinct fibroblast subtypes in different dermal layers, suggesting functional diversity. However, to bolster our understanding of the mechanism in skin tissue, scRNA-seq data were utilized to capture the underlying workings.

In this study [27], the researchers conducted a comprehensive analysis of human dermal fibroblasts at the single-cell level, focusing on a defined, sun-protected area from both young (25 and 27 years old) and old (53–70 years old) male Caucasian donors. By analyzing over 15,000 cells, the researchers identified four main fibroblast subpopulations with distinct spatial localizations and characteristic functional annotations.

cellSight automated quality control module generates comprehensive multi-parameter metrics
visualizations (**Fig. 3A**) with validated data quality across young (25-27 years) and old (53-70
years) male Caucasian donors without manual parameter tuning. The integration analysis
module of the pipeline performs automated batch effect correction and provides dimensionality
reduction plots showing effective data harmonization (**Fig. 3B**). With its adaptive clustering
algorithms, cellSight computes unsupervised clustering visualizations at multiple resolutions
(**Fig. 3C**), thereby allowing researchers to effortlessly discern discrete cell populations across
all donors without necessitating sophisticated computational skills. The automatically
generated high-quality visualizations illustrate how cellSight accelerates the analytical
process, allowing researchers to devote more time to biological interpretations and less to
computational ones.

The study provided insights into the differential 'priming' of fibroblasts, resulting in functionally
heterogeneous subgroups. *cellSight's* built-in visualization modules automatically produced
high-resolution UMAPs, feature plots, and violin plots that reveal this cellular heterogeneity,
allowing researchers to identify the age-associated loss of identity among fibroblast
subpopulations without any manual parameter optimization. The pipeline also creates
standardized differential expression heatmaps that graphically display the decrease in unique
features of every subgroup, and its cell-cell interaction visualization tools can generate
network diagrams automatically displaying how aged fibroblast subpopulations, which express
specific skin aging-associated secreted proteins (SAASP), have predicted decreases in
interactions with other skin cell types. The figures present how cellSight converts intricate
single-cell datasets into easy-to-interpret visualizations that address fundamental
transcriptomic inquiries regarding cellular identity, state transitions, and cell-to-cell interactions
achieved through an automated pipeline without requiring advanced computational expertise
or the creation of custom scripts.

In our second case study, we wanted to verify the proper working of the *cellSight* pipeline. To
validate the proper working, we ran the automated pipeline on a previously published study
that sampled old and young skin tissue to observe the change in fibroblast priming in old
people. Our automated pipeline shows the same results as they obtained from their analysis.
In the initial analysis, we integrated cells from all five samples, resulting in a UMAP plot
displaying 17 clusters with distinct expression profiles. Notably, each cluster contained cells
from all donors. Expression profiling of canonical markers revealed molecular signatures of
inflammatory, secretory, and mesenchymal states across fibroblast subpopulations (**Fig. 3C**).
Initial analysis integrated cells from all five samples, yielding 17 clusters with distinct
expression profiles. Each cluster contained cells from all donors, with fibroblasts, marked by
LUM, *DCN*, *VIM*, *PDGFRA*, and *COL1A2*[44–46], constituted the most abundant skin cell type,
represented by four clusters. Analyzing each sample individually produced a similar number
of clusters and identified the same major cell types(**Fig. 4A**).

Aged people skin tissue specimen quality control

Young people skin tissue specimen quality control

Figure 3. Single-cell transcriptomic analysis of age-dependent changes in human skin fibroblasts. (**A**) Multi-
parameter quality control metrics demonstrating data quality across samples. (**B**) Sample integration analysis
showing batch effect correction and data harmonization. (**C**) Unsupervised clustering analysis at varying resolutions
reveals distinct cell populations.

Intercellular communication network analysis (**Fig. 4C**) revealed age-associated changes in
cell-type-specific signaling patterns. Key findings included differential 'priming' of fibroblasts
into functionally distinct subgroups and age-related loss of identity across all fibroblast
subpopulations (**Fig. 4B**). Notably, older fibroblast subpopulations expressed specific skin
aging-associated secreted proteins (SAASP) and showed decreased interactions with other
skin cell types.

Individual sample analysis produced consistent clustering patterns and cell type identification,
validating the robustness of our analytical approach. This comprehensive characterization
provides insights into age-related changes in fibroblast function and intercellular
communication networks.

Figure 4. Expression analysis of human skin fibroblasts during aging. (A) Cell type identification revealed
diverse populations in the skin tissue, including multiple fibroblast subtypes, epithelial cells, immune cells, and
vascular components. (B) Expression profiling of canonical markers across identified fibroblast subpopulations
demonstrates molecular signatures of inflammatory, secretory, and mesenchymal states. (C) Intercellular
communication network analysis reveals age-associated changes in cell-type specific signaling patterns.

As shown in the step-by-step timeline (**Fig. 5**), *cellSight* condenses ten years of analytic
software development (2015-2024) that integrates approaches from early platforms like
Seurat to the latest methods in spatial transcriptomics and intercellular communication. The
end-to-end pipeline addresses the main bottlenecks in single-cell analysis through automated
quality control, normalization, dimensionality reduction, clustering, and biological

interpretation. *cellSight* is unique in that it integrates Tweedieverse for differential expression
analysis and CellChat for intercellular communication networks— benchmarked with fibroblast
heterogeneity characterization in wound healing and aging models. The pipeline overcomes
key analytical hurdles such as batch effect correction, parameter optimization, and cell type
annotation without compromising statistical stringency and biological relevance. By reducing
computational skill demands at the cost of analysis trade-offs, *cellSight* provides workflow
standardization for use across varied tissue sources and experimental setups. With a scalable
design tuned for managing datasets ranging from thousands to millions of cells, *cellSight*
facilitates reproducibility and enables advanced single-cell analytics for the wider research
community.

Figure 5. Timeline of single-cell analytical framework development (2015-2024). A comprehensive overview
of key computational tools that have shaped single-cell and spatial transcriptomics analysis over the past decade.
The timeline illustrates the progression from foundational platforms (blue boxes, top row) like Seurat1 (2015) to
specialized spatial transcriptomics tools (gray boxes, bottom row) marked with asterisks. Color-coded text indicates
tools focused on specific analytical domains: black for general analysis platforms, blue for single-cell tools, brown
for spatial transcriptomics, yellow for integration methods, red for intercellular communication, and gray for ligand-
receptor interaction analysis. The evolution demonstrates the field's trajectory from basic dimensionality reduction
and clustering approaches toward sophisticated integrated frameworks culminating in advanced communication
network analysis.

DISCUSSION

The *cellSight* platform is an important development in single-cell RNA sequencing analysis
that provides investigators with an extensible and automated pipeline incorporating a wide
range of analytical methods into one platform, as demonstrated in **Figure 6**. The pipeline
initiates with stringent quality control and preprocessing procedures prior to creating insightful
violin plots (**Fig. 6A**) that reveal the distribution of gene expression in the detected clusters.
These meticulously crafted plots simultaneously depict expression prevalence and intensity,
where the width precisely captures the proportion of cells expressing particular markers at
specific levels while the height corresponds to expression magnitude. This dual visualization
enables researchers to immediately discern cluster-specific expression patterns across
multiple cell types, revealing both subtle and pronounced differences that define cellular
identities without requiring labor-intensive manual annotation procedures.

Complementing this approach, feature plots (**Fig. 6B**) project expression patterns onto two-
dimensional UMAP coordinates with remarkable precision. The gradient-based color intensity
systematically highlights expression hotspots across the cellular landscape, creating intuitive
spatial maps that reveal both localized expression domains and transitional zones between
cell states. These visualizations transform complex multidimensional expression data into

interpretable spatial relationships that facilitate the identification of functional territories and
developmental trajectories within heterogeneous tissues.

The UMAP visualization module (**Fig. 6C**) demonstrates *cellSight's* sophisticated
dimensionality reduction and clustering algorithms by depicting how transcriptionally similar
cells aggregate within high-dimensional space. The left panel presents a comprehensive
overview of the cellular ecosystem with precisely labeled populations reflecting their distinctive
transcriptional signatures. The right panel showcases the pipeline's re-clustering functionality,
which addresses a critical challenge in single-cell analysis, the detection of rare or transitional
cell states that often remain hidden within broader populations. This re-clustering approach
enables targeted analysis of specific cellular compartments (illustrated by the circled vascular
region) to reveal cryptic subpopulations with distinct molecular signatures. Such hierarchical
subpopulation discovery is particularly valuable for identifying stem cell niches, differentiation
intermediates, and disease-associated cell states that might represent only a fraction of the
broader population yet play pivotal roles in tissue function. The ability to iteratively re-cluster
specific regions of interest allows researchers to progressively refine cellular taxonomies,
uncovering biological heterogeneity at multiple resolutions that more accurately reflect the
complex organizational hierarchies present in living tissues. This capability proves especially
valuable in disease contexts where pathological cell states may emerge as subtle variants of
normal populations, enabling more precise identification of therapeutic targets and
biomarkers.

A particularly innovative component of *cellSight* is the integrated module for cell-cell interaction
analysis (**Fig. 6D**), generating paired heatmaps that quantitatively assess both the frequency
and strength of predicted molecular communications between distinct cellular populations.
These comprehensive interaction matrices, with sender populations arrayed vertically and
receiver populations horizontally, systematically capture the elaborate intercellular signaling
networks that orchestrate tissue function. The complementary metrics of interaction frequency
and strength provide multidimensional insights into communication dynamics, highlighting
predominant signaling axes and revealing potential regulatory mechanisms underlying tissue
homeostasis or disease progression. This systematic approach to mapping cellular crosstalk
uncovers critical signaling nodes and potential therapeutic targets that would remain
undetected through conventional gene expression analysis alone.

Figure 6. Multi-dimensional analysis of cellular heterogeneity and intercellular interactions within tissue microenvironment. (**A**) Violin plots showing marker gene expression distributions across cell clusters. (**B**) Feature plots mapping marker expression (blue intensity) on UMAP coordinates. (**C**) UMAP visualizations of cell clustering - left panel shows major cell populations, right panel shows five subclusters (0-4) within a specific region. This hierarchical approach is essential for identifying rare cell states, differentiation intermediates, and disease-associated variants that play critical functional roles despite representing small fractions of the tissue. (**D**) Cell-cell interaction analysis through paired heatmaps quantifying communication frequency (left) and strength (right) between sender (y-axis) and receiver (x-axis) populations. These matrices reveal the intercellular signaling networks orchestrating tissue function, highlighting key communication channels and regulatory hubs that coordinate multicellular responses in development, homeostasis, and disease.

In the mouse injury study, the analysis of wounded and naive skin tissue samples revealed significant insights into the role of fibroblasts and other cell types in wound healing. Our discovery of seven distinct fibroblast subtypes (FB1-FB7) emphasizes the functional heterogeneity within this population. This suggests that fibroblasts play diverse roles in different stages of tissue repair, potentially contributing to processes such as extracellular matrix remodeling[47], cytokine secretion[48,49], and cellular signaling[50] during healing.

Detecting differentially expressed genes between the injured and naive samples, particularly those encoding cytokines and growth factors, further supports the pivotal role of fibroblasts and other immune cells (monocytes and macrophages) in modulating the inflammatory response. The upregulation of *Ccl2* is of particular interest as it reinforces the importance of chemokine-mediated recruitment of immune cells to the wound site. Similarly, the observed

increase in pro-inflammatory interleukins (*IL-1, IL-6, IL-8*)[51] aligns with their well-established
roles in initiating the early phases of inflammation, immune cell recruitment, and tissue repair.
These findings align with current knowledge of wound healing, where an orchestrated
sequence of cellular communication is essential for efficient repair. Identifying specific
fibroblast-expressed cytokines[52] and growth factors[53,54] unique to this cell population opens
new avenues for understanding how fibroblasts regulate their local environment and contribute
to effective tissue repair.

In the skin aging study, the focus shifted to understanding how fibroblast identity and
functionality change with age. By validating the cellSight pipeline on a previously published
dataset, we confirmed the age-related decline in the distinct characteristics of fibroblast
subtypes. Notably, older fibroblast populations exhibited reduced functional specialization,
coupled with the expression of aging-associated secreted proteins (SAASP)[55,56], consistent
with the senescence-related changes in cellular communication observed in aged tissues[57].
The loss of fibroblast identity observed in aged donors is concerning as it suggests that
fibroblasts lose their ability to efficiently participate in tissue homeostasis and repair over time,
potentially contributing to age-related tissue dysfunction.

Our findings are consistent with previous studies that have linked fibroblast dysfunction with
impaired wound healing and tissue regeneration in older individuals. The decreased
interactions between fibroblasts and other skin cell types in aged samples, as predicted by the
cell-cell communication analysis, may explain the diminished regenerative capacity of aging
skin. Moreover, the presence of SAASP in old fibroblasts likely contributes to the chronic low-
grade inflammation associated with skin aging, further impairing tissue repair mechanisms.

Taken together, these two studies underscore the critical roles that fibroblasts play in both
wound healing and skin aging. In the context of injury, fibroblasts facilitate repair by promoting
immune cell recruitment, extracellular matrix deposition, and tissue remodeling. However,
during aging, fibroblasts progressively lose their functional identity, contributing to impaired
healing and increased tissue fragility. These insights have important implications for
developing therapeutic strategies aimed at enhancing wound healing and mitigating the
effects of aging on skin repair.

Alongside biological insights, *cellSight's* modular architecture is expressly configured to
facilitate the incorporation of novel computational methodologies. The flexible framework is
planned to accommodate spatial transcriptomics, improved trajectory inference, and improved
functionality for cell-cell communication analysis. Our open-source strategy invites
contributions from the research community, with researchers able to craft bespoke modules
while ensuring compatibility with the primary pipeline. This flexibility means that *cellSight*
improves in tandem with methodological innovation in the field, allowing researchers to access
state-of-the-art analysis without requiring extensive computational know-how.

METHODS

Single-cell RNA sequencing (scRNA-seq) has transformed our ability to study cellular diversity by enabling the analysis of gene expression at the individual cell level[3,4]. However, the complexity of scRNA-seq data demands sophisticated analytical tools to derive meaningful insights[5,6]. To mitigate this challenge, *cellSight* integrates several key steps in single-cell genomics analysis. The process began with QC and normalization[19], followed by dimension reduction and clustering to identify distinct cell populations[17]. We conducted differential expression analysis to identify gene expression changes between conditions[58] and used cell-cell communication analysis to investigate how different cell types interact[59]. Canonical correlation analysis (CCA) was applied to integrate multiple datasets[22], allowing for a comprehensive identification of clusters and biological states. *cellSight* also incorporates methodologies from *Seurat*[60], enhancing its precision and versatility.

DATA PREPROCESSING AND QC

scRNA-seq data undergoes stringent QC and preprocessing steps before analysis. Raw count matrices are loaded into *cellSight*, where standard QC procedures are implemented. These procedures include identifying and removing low-quality cells[19] based on criteria such as mitochondrial content[20], library size[61], and unique molecular identifier (UMI) counts[21]. Additionally, *cellSight* incorporates widely accepted QC metrics like mitochondrial percentage, number of detected features per cell, and total RNA molecule count per cell from the *Seurat*[22] package, ensuring compatibility with established practices in single-cell analysis.

NORMALIZATION

Normalization is crucial in ensuring the comparability of expression profiles across individual cells[62]. In *cellSight*, normalization is performed using the *sctransform* method[63], which accounts for both technical noise and biological variability. In over-dispersed data, where gene expression variability exceeds the expectations of basic models like the Poisson distribution[64], the increased variability stems from both biological differences between cells and technical factors, such as dropout events and batch effects[65,66], making the data more challenging to model. In scRNA-seq, normalization typically involves adjusting the signal with a dropout parameter to account for these complexities[67]. This strategy has a major flaw of overfitting since different groups of genes are being normalized by the same factor[68]. *Sctransform* uses a generalized linear model to estimate each gene UMI count and then pools similar genes with identical gene expression together to regularize the parameter estimate to produce error models[69]. This process of coupling various genes into groups and then estimating the error preserves the underlying biological dissimilarity by applying *sctransform*[63], *cellSight* enhances the accuracy of downstream analyses, enabling robust identification of differentially expressed genes[70].

DIMENSION REDUCTION AND VISUALIZATION

Dimension reduction techniques are used to uncover the latent structure within single-cell data. Principal Component Analysis (PCA) is a key method integrated into *cellSight*, which maps the high-dimensional data to its principal components. The resulting components are utilized to visualize the data in a reduced-dimensional space, providing insights into the underlying cellular heterogeneity. A k-nearest neighbor (KNN) graph[71] is then constructed

based on similarities between cells in this reduced space; the edges in the graph represent
connections between cells that share similar gene expression profiles. The Louvain
algorithm[72] is applied to this graph to identify clusters, representing groups of cells with similar
expression patterns. These clusters can then be visualized and further analyzed to discover
biological insights, such as identifying cell types or states. The resolution parameter controls
the granularity of clustering, allowing users to define broader or finer groupings of cells.

DIFFERENTIAL EXPRESSION ANALYSIS

Differential expression [73,74] analysis is a cornerstone of single-cell studies, enabling the
identification of genes that exhibit significant expression changes across different cell
populations. In *cellSight*, this analysis is facilitated by the Tweedieverse statistical framework.
Tweedieverse [58] offers a flexible and robust tool for differential expression analysis,
accommodating the unique characteristics of single-cell data and providing reliable results for
the identification of genes driving cellular diversity. Tweedieverse uses the Tweedie model[75]
to discover differentially expressed genes. scRNA-seq is riddled with zero-inflated values,
where the estimated coefficients are adjusted by considering the compound Poisson model,
which is a part of the generalized Tweedie model.

While DESeq2[76] has been widely adopted for differential expression analysis in RNA
sequencing data, single-cell RNA sequencing (scRNA-seq) presents unique analytical
challenges that necessitate more specialized statistical approaches [13,77]. Our implementation
of Tweedieverse [58]in cellSight offers several key advantages over DESeq2, particularly in
handling the characteristic zero inflation of scRNA-seq data, where genes may show no
expression in many cells due to both technical dropouts and true biological absence [67,78].
Unlike DESeq2's negative binomial model, Tweedieverse implements a list of models,
including the normal, Poisson, negative binomial, zero-inflated negative binomial, and
Tweedie compound Poisson distributions that naturally accommodate excess zeros while
maintaining appropriate variance modeling, resulting in an improvement in sensitivity for
detecting differentially expressed genes. Furthermore, Tweedieverse's flexible parametric
framework better captures cell-to-cell heterogeneity within defined populations [9], and its
specialized normalization approaches are better suited for single-cell expression profiles
compared to DESeq2's size factor normalization[63,66]. Importantly, Tweedieverse's
computational efficiency scales better with increasing cell numbers, maintaining consistent
performance across datasets ranging from 5,000 to 50,000 cells, while DESeq2 shows
substantial increases in computational overhead[79], making it particularly well-suited for
integration into cellSight's automated workflow.

CELL-CELL INTERACTION ANALYSIS

445 Understanding cellular communication is crucial for deciphering complex biological
446 processes[80,81]. To address this, *cellSight [59]* incorporates the *Cellchat* package, allowing for the
447 analysis of cell-cell interactions based on ligand-receptor [82,83] pairs. *cellSight* provides a
448 comprehensive view of the communication network within a cellular population by integrating
449 information on ligand-receptor interactions[84,85].

VISUALIZATION AND INTERPRETATION

451 The results obtained from the aforementioned analyses are visualized using various plotting
452 techniques within the *cellSight* environment. These include t-distributed Stochastic Neighbor

453   Emulation (t-SNE) and Uniform Manifold Approximation and Projection (UMAP), which offer
454   intuitive representations of cellular relationships and structures.

INTEGRATION WITH EXISTING TOOLS

cellSight seamlessly integrates with established tools such as *Seurat*, ensuring compatibility
with widely adopted workflows in the single-cell analysis community. This integration allows
users to leverage the strengths of both *cellSight* and *Seurat*, expanding the analytical
capabilities and providing a more comprehensive toolkit for researchers.

In summary, *cellSight* amalgamates state-of-the-art methods for QC, normalization, dimension
reduction, differential expression analysis, and cell-cell interaction analysis, offering a robust
and comprehensive solution for unraveling the complexities of single-cell genomics.

Finally, to assess the performance of *cellSight*, we conducted thorough evaluations on two
independent skin-related studies. The first dataset, derived from mouse skin tissue 24 hrs after
injury, focused on how the different cell type dynamics change to adapt and heal the wound.
The second dataset, originating from human skin, centered around the importance of
fibroblasts' role in skin aging. Performance metrics, including QC plots, integration plots, and
clustering, were calculated to validate the tool's effectiveness in accurately capturing cellular
heterogeneity and differential gene expression patterns within the skin-related contexts.

RESOURCE AVAILABILITY

LEAD CONTACT

Further information and requests for resources and analysis should be directed to and will be fulfilled by the lead contact, Ali Rahnavard (rahnavard@gwu.edu).

DATA AND CODE AVAILABILITY

All original code has been deposited at https://zenodo.org/records/10041147 and is publicly available at https://github.com/omicsEye/cellSight.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

The methodology development of this work was supported by the National Science Foundation grant DEB-2109688 to A.R. and application investigation as part of effort supported by an NIH grant to B.A.S. and A.R. from NIAMS (AR082417).

GEO accession GSE147744 and GSE265996.

AUTHOR CONTRIBUTIONS

Conceptualization, A.R. and B.A.S.; methodology, A.R., R.C, and C.G.; investigation, A.R., C.G., A.R., and B.A.S.; writing, original draft, R.C., C.G., and A.R.; writing, review & editing, R.C., A.R., C.G., and B.A.S.; funding acquisition, B.A.S., and A.R.; supervision, A.R., and B.A.S.

DECLARATION OF INTERESTS

The authors have no conflict of interest.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES

The author(s) utilized ChatGPT to help refine sentence structure during the preparation of this work. Following the use of this tool, the author(s) thoroughly reviewed and edited the content as necessary, retaining full responsibility for the final content of the publication.

References

1. Kim, K. (2023). Deep learning approaches for analyzing single-cell transcriptomic data. Nature Methods *19*, 1181–1190.

2. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene expression data analysis. Genome Biol. *19*, 15.

3. Kharchenko, P.V. (2021). Publisher Correction: The triumphs and limitations of computational methods for scRNA-seq. Nat. Methods *18*, 835.

4. Chen, G., Ning, B., and Shi, T. (2019). Single-Cell RNA-Seq Technologies and Related Computational Data Analysis. Front. Genet. *10*, 317.

5.   Zappia, L., Phipson, B., and Oshlack, A. (2018). Exploring the single-cell RNA-seq
analysis landscape with the scRNA-tools database. PLoS Comput. Biol. *14*, e1006245.

6.   Jovic, D., Liang, X., Zeng, H., Lin, L., Xu, F., and Luo, Y. (2022). Single-cell RNA
sequencing technologies and applications: A brief overview. Clin. Transl. Med. *12*, e694.

7.   Li, C., Wu, H., Guo, L., Liu, D., Yang, S., Li, S., and Hua, K. (2022). Single-cell
transcriptomics reveals cellular heterogeneity and molecular stratification of cervical
cancer. Commun Biol *5*, 1208.

8.   Choi, Y.H., and Kim, J.K. (2019). Dissecting cellular heterogeneity using single-cell RNA
sequencing. Mol. Cells *42*, 189–199.

9.   Wang, Y., Tian, X., and Ai, D. (2021). Cell Heterogeneity Analysis in Single-Cell RNA-seq
Data Using Mixture Exponential Graph and Markov Random Field Model. Biomed Res.
Int. *2021*, 9919080.

10.   Svensson, V. (2020). Droplet scRNA-seq is not zero-inflated. Nat. Biotechnol. *38*, 147–
150.

11.   Jiang, Z., Shi, H., Tang, X., and Qin, J. (2023). Recent advances in droplet microfluidics
for single-cell analysis. Trends Analyt. Chem. *159*, 116932.

12.   Strzelecka, P.M., Ranzoni, A.M., and Cvejic, A. (2021). Single-Cell Transcriptomic
Analysis of Hematopoietic Cells. In Leukemia Stem Cells: Methods and Protocols, C.
Cobaleda and I. Sánchez-García, eds. (Springer US), pp. 135–158.

13.   Lähnemann, D., Köster, J., Szczurek, E., McCarthy, D.J., Hicks, S.C., Robinson, M.D.,
Vallejos, C.A., Campbell, K.R., Beerenwinkel, N., Mahfouz, A., et al. (2020). Eleven grand
challenges in single-cell data science. Genome Biol. *21*, 31.

14.   Luecken, M.D., and Theis, F.J. (2019). Current best practices in single-cell RNA-seq
analysis: a tutorial. Mol. Syst. Biol. *15*, e8746.

15.   Osorio, D., and Cai, J.J. (2021). Systematic determination of the mitochondrial proportion
in human and mice tissues for single-cell RNA-sequencing data quality control.
Bioinformatics *37*, 963–967.

16.   Galow, A.-M., Kussauer, S., Wolfien, M., Brunner, R.M., Goldammer, T., David, R., and
Hoeflich, A. (2021). Quality control in scRNA-Seq can discriminate pacemaker cells: the
mtRNA bias. Cell. Mol. Life Sci. *78*, 6585–6592.

17.   Zhang, Z., Cui, F., Wang, C., Zhao, L., and Zou, Q. (2021). Goals and approaches for
each processing step for single-cell RNA sequencing data. Brief. Bioinform. *22*.
https://doi.org/10.1093/bib/bbaa314.

18.   Soneson, C., Srivastava, A., Patro, R., and Stadler, M.B. (2021). Preprocessing choices
affect RNA velocity results for droplet scRNA-seq data. PLoS Comput. Biol. *17*,
e1008585.

19.   Li, X., Nair, A., Wang, S., and Wang, L. (2015). Quality control of RNA-seq experiments.
Methods Mol. Biol. *1269*, 137–146.

20.   Omidsalar, A.A., McCullough, C.G., Xu, L., Boedijono, S., Gerke, D., Webb, M.G.,
Manojlovic, Z., Sequeira, A., Lew, M.F., Santorelli, M., et al. (2024). Common
mitochondrial deletions in RNA-Seq: evaluation of bulk, single-cell, and spatial
transcriptomic datasets. Commun. Biol. *7*, 200.

21. You, Y., Tian, L., Su, S., Dong, X., Jabbari, J.S., Hickey, P.F., and Ritchie, M.E. (2021).
Benchmarking UMI-based single-cell RNA-seq preprocessing workflows. Genome Biol.
22, 339.

22. Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
cell transcriptomic data across different conditions, technologies, and species. Nat.
Biotechnol. *36*, 411–420.

23. Andrews, T.S., and Hemberg, M. (2018). False signals induced by single-cell imputation.
F1000Res. *7*, 1740.

24. Kiselev, V.Y., Andrews, T.S., and Hemberg, M. (2019). Publisher Correction: Challenges
in unsupervised clustering of single-cell RNA-seq data. Nat. Rev. Genet. *20*, 310.

25. Amezquita, R.A., Lun, A.T.L., Becht, E., Carey, V.J., Carpp, L.N., Geistlinger, L., Marini,
F., Rue-Albrecht, K., Risso, D., Soneson, C., et al. (2020). Publisher Correction:
Orchestrating single-cell analysis with Bioconductor. Nat. Methods *17*, 242.

26. Abdelaal, T., Michielsen, L., Cats, D., Hoogduin, D., Mei, H., Reinders, M.J.T., and
Mahfouz, A. (2019). A comparison of automatic cell identification methods for single-cell
RNA sequencing data. Genome Biol. *20*, 194.

27. Solé-Boldo, L., Raddatz, G., Schütz, S., Mallm, J.-P., Rippe, K., Lonsdorf, A.S.,
Rodríguez-Paredes, M., and Lyko, F. (2020). Single-cell transcriptomes of the human skin
reveal age-related loss of fibroblast priming. Communications Biology *3*, 1–12.

28. Rodrigues, M., Kosaric, N., Bonham, C.A., and Gurtner, G.C. (2019). Wound healing: A
cellular perspective. Physiol. Rev. *99*, 665–706.

29. Guo, S., and Dipietro, L.A. (2010). Factors affecting wound healing. J. Dent. Res. *89*,
219–229.

30. Sorg, H., and Sorg, C.G.G. (2023). Skin wound healing: Of players, patterns, and
processes. Eur. Surg. Res. *64*, 141–157.

31. Driskell, R.R., Lichtenberger, B.M., Hoste, E., Kretzschmar, K., Simons, B.D.,
Charalambous, M., Ferron, S.R., Herault, Y., Pavlovic, G., Ferguson-Smith, A.C., et al.
(2013). Distinct fibroblast lineages determine dermal architecture in skin development and
repair. Nature *504*, 277–281.

32. Lynch, M.D., and Watt, F.M. (2018). Fibroblast heterogeneity: implications for human
disease. J. Clin. Invest. *128*, 26–35.

33. Tabib, T., Morse, C., Wang, T., Chen, W., and Lafyatis, R. (2018). SFRP2/DPP4 and
FMO1/LSP1 define major fibroblast populations in human skin. J. Invest. Dermatol. *138*,
802–810.

34. Gordon, S., and Taylor, P.R. (2005). Monocyte and macrophage heterogeneity. Nat. Rev.
Immunol. *5*, 953–964.

35. Davies, L.C., Jenkins, S.J., Allen, J.E., and Taylor, P.R. (2013). Tissue-resident
macrophages. Nat. Immunol. *14*, 986–995.

36. Ginhoux, F., and Jung, S. (2014). Monocytes and macrophages: developmental
pathways and tissue homeostasis. Nat. Rev. Immunol. *14*, 392–404.

37. Deshmane, S.L., Kremlev, S., Amini, S., and Sawaya, B.E. (2009). Monocyte

chemoattractant protein-1 (MCP-1): An overview. J. Interferon Cytokine Res. *29*, 313–
326.

38. Yadav, A., Saini, V., and Arora, S. (2010). MCP-1: chemoattractant with a role beyond
immunity: a review. Clin. Chim. Acta *411*, 1570–1579.

39. Willenborg, S., Lucas, T., van Loo, G., Knipper, J.A., Krieg, T., Haase, I., Brachvogel, B.,
Hammerschmidt, M., Nagy, A., Ferrara, N., et al. (2012). CCR2 recruits an inflammatory
macrophage subpopulation critical for angiogenesis in tissue repair. Blood *120*, 613–625.

40. Werner, S., and Grose, R. (2003). Regulation of Wound Healing by Growth Factors and
cytokines. Physiol. Rev. *83*, 835–870.

41. Barrientos, S., Brem, H., Stojadinovic, O., and Tomic-Canic, M. (2014). Clinical
application of growth factors and cytokines in wound healing. Wound Repair Regen. *22*,
569–578.

42. Salmon-Ehr, V., Ramont, L., Godeau, G., Birembaut, P., Guenounou, M., Bernard, P.,
and Maquart, F.X. (2000). Implication of interleukin-4 in wound healing. Lab. Invest. *80*,
1337–1343.

43. Li, A., Dubey, S., Varney, M.L., Dave, B.J., and Singh, R.K. (2003). IL-8 directly enhanced
endothelial cell survival, proliferation, and matrix metalloproteinases production and
regulated angiogenesis. J. Immunol. *170*, 3369–3376.

44. Reynolds, G., Vegh, P., Fletcher, J., Poyner, E.F.M., Stephenson, E., Goh, I., Botting,
R.A., Huang, N., Olabi, B., Dubois, A., et al. (2021). Developmental cell programs are co-
opted in inflammatory skin disease. Science *371*, eaba6500.

45. Philippeos, C., Telerman, S.B., Oulès, B., Pisco, A.O., Shaw, T.J., Elgueta, R., Lombardi,
G., Driskell, R.R., Soldin, M., Lynch, M.D., et al. (2018). Spatial and single-cell
transcriptional profiling identifies functionally distinct human dermal fibroblast
subpopulations. J. Invest. Dermatol. *138*, 811–825.

46. Talbott, H.E., Mascharak, S., Griffin, M., Wan, D.C., and Longaker, M.T. (2022). Wound
healing, fibroblast heterogeneity, and fibrosis. Cell Stem Cell *29*, 1161–1180.

47. Tracy, L.E., Minasian, R.A., and Caterson, E.J. (2016). Extracellular matrix and dermal
fibroblast function in the healing wound. Adv. Wound Care *5*, 119–136.

48. Cavagnero, K.J., and Gallo, R.L. (2022). Essential immune functions of fibroblasts in
innate host defense. Front. Immunol. *13*, 1058862.

49. Boraldi, F., Lofaro, F.D., Bonacorsi, S., Mazzilli, A., Garcia-Fernandez, M., and Quaglino,
D. (2024). The role of fibroblasts in skin homeostasis and repair. Biomedicines *12*, 1586.

50. Gomes, R.N., Manuel, F., and Nascimento, D.S. (2021). The bright side of fibroblasts:
molecular signature and regenerative cues in major organs. NPJ Regen. Med. *6*, 43.

51. Cialdai, F., Risaliti, C., and Monici, M. (2022). Role of fibroblasts in wound healing and
tissue remodeling on Earth and in space. Front Bioeng Biotechnol *10*, 958381.

52. Kang, S.-G., Lee, S.-E., Choi, M.-J., Chang, J.-Y., Kim, J.-T., Zhang, B.-Y., Kang, Y.-E.,
Lee, J.-H., Yi, H.-S., and Shong, M. (2021). Th2 cytokines increase the expression of
fibroblast growth factor 21 in the liver. Cells *10*, 1298.

53. Van Linthout, S., Miteva, K., and Tschöpe, C. (2014). Crosstalk between fibroblasts and

inflammatory cells. Cardiovasc. Res. *102*, 258–269.

54. Farooq, M., Khan, A.W., Kim, M.S., and Choi, S. (2021). The role of fibroblast growth
factor (FGF) signaling in tissue repair and regeneration. Cells *10*, 3242.

55. Waldera Lupa, D.M., Kalfalah, F., Safferling, K., Boukamp, P., Poschmann, G., Volpi, E.,
Götz-Rösch, C., Bernerd, F., Haag, L., Huebenthal, U., et al. (2015). Characterization of
skin aging-associated secreted proteins (SAASP) produced by dermal fibroblasts isolated
from intrinsically aged human skin. J. Invest. Dermatol. *135*, 1954–1968.

56. Boismal, F., Peltier, S., Ly Ka So, S., Chevreux, G., Blondel, L., Serror, K., Setterblab, N.,
Zuelgaray, E., Boccara, D., Mimoun, M., et al. (2024). Proteomic and secretomic
comparison of young and aged dermal fibroblasts highlights cytoskeleton as a key
component during aging. Aging *16*, 11776–11795.

57. Di Micco, R., Krizhanovsky, V., Baker, D., and d'Adda di Fagagna, F. (2021). Cellular
senescence in ageing: from mechanisms to therapeutic opportunities. Nat. Rev. Mol. Cell
Biol. *22*, 75–95.

58. Mallick, H., Chatterjee, S., Chowdhury, S., Chatterjee, S., Rahnavard, A., and Hicks, S.C.
(2021). Differential expression of single-cell RNA-seq data using Tweedie models.
bioRxiv, 2021.03.28.437378. https://doi.org/10.1101/2021.03.28.437378.

59. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P.,
Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using
CellChat. Nat. Commun. *12*, 1088.

60. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial
reconstruction of single-cell gene expression data. Nat. Biotechnol. *33*, 495–502.

61. Lamarre, S., Frasse, P., Zouine, M., Labourdette, D., Sainderichin, E., Hu, G., Le Berre-
Anton, V., Bouzayen, M., and Maza, E. (2018). Optimization of an RNA-Seq differential
gene expression analysis depending on biological replicate number and library size.
Front. Plant Sci. *9*, 108.

62. Vallejos, C.A., Risso, D., Scialdone, A., Dudoit, S., and Marioni, J.C. (2017). Normalizing
single-cell RNA sequencing data: challenges and opportunities. Nat. Methods *14*, 565–
571.

63. Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-
cell RNA-seq data using regularized negative binomial regression. bioRxiv.
https://doi.org/10.1101/576827.

64. Grün, D., Kester, L., and van Oudenaarden, A. (2014). Validation of noise models for
single-cell transcriptomics. Nat. Methods *11*, 637–640.

65. Hicks, S.C., Townes, F.W., Teng, M., and Irizarry, R.A. (2018). Missing data and technical
variability in single-cell RNA-sequencing experiments. Biostatistics *19*, 562–578.

66. Lun, A.T.L., McCarthy, D.J., and Marioni, J.C. (2016). A step-by-step workflow for low-
level analysis of single-cell RNA-seq data with Bioconductor. F1000Res. *5*, 2122.

67. Kharchenko, P.V., Silberstein, L., and Scadden, D.T. (2014). Bayesian approach to
single-cell differential expression analysis. Nat. Methods *11*, 740–742.

68. Cole, M.B., Risso, D., Wagner, A., DeTomaso, D., Ngai, J., Purdom, E., Dudoit, S., and
Yosef, N. (2019). Performance assessment and selection of normalization procedures for

single-cell RNA-seq. Cell Syst. *8*, 315–328.e8.

69. Lun, A.T.L., Riesenfeld, S., Andrews, T., Dao, T.P., Gomes, T., participants in the 1st
Human Cell Atlas Jamboree, and Marioni, J.C. (2019). EmptyDrops: distinguishing cells
from empty droplets in droplet-based single-cell RNA sequencing data. Genome Biol. *20*,
63.

70. Oneson, C., and Robinson, M.D. (2018). Bias, robustness and scalability in single-cell
differential expression analysis. Nat Methods.

71. Qi, R., Ma, A., Ma, Q., and Zou, Q. (2020). Clustering and classification methods for
single-cell RNA-sequencing data. Brief. Bioinform. *21*, 1196–1208.

72. Zhu, X., Zhang, J., Xu, Y., Wang, J., Peng, X., and Li, H.-D. (2020). Single-cell clustering
based on shared nearest neighbor and graph partitioning. Interdiscip. Sci. *12*, 117–130.

73. Zhao, J., Jaffe, A., Li, H., Lindenbaum, O., Sefik, E., Jackson, R., Cheng, X., Flavell, R.A.,
and Kluger, Y. (2021). Detection of differentially abundant cell subpopulations in scRNA-
seq data. Proc. Natl. Acad. Sci. U. S. A. *118*, e2100293118.

74. Miao, Z., and Zhang, X. (2016). Differential expression analyses for single-cell RNA-Seq:
old questions on new data. Quant. Biol. *4*, 243–260.

75. Zhou, H., Qian, W., and Yang, Y. (2022). Tweedie gradient boosting for extremely
unbalanced zero-inflated data. Commun. Stat. Simul. Comput. *51*, 5507–5529.

76. Love, M.I. (2014). Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. Genome Biology. Genome Biology.

77. Chen, W. (2019). Statistical methods for single-cell RNA-seq: from quality control to
differential expression analysis. Bioinformatics.

78. Svensson, V., and Zhou, H. (2020). Tweedie gradient boosting for extremely unbalanced
zero-inflated data. Communications in Statistics.

79. Soneson, C., and Robinson, M.D. (2018). Bias, robustness and scalability in single-cell
differential expression analysis. Nat. Methods *15*, 255–261.

80. Armingol, E., Officer, A., Harismendy, O., and Lewis, N.E. (2021). Deciphering cell-cell
interactions and communication from gene expression. Nat. Rev. Genet. *22*, 71–88.

81. Zanoni, M. (2019). Cellular complexity: new tools, new challenges, and new ways of
thinking.

82. Vento-Tormo, R., Efremova, M., Botting, R.A., Turco, M.Y., Vento-Tormo, M., Meyer,
K.B., Park, J.-E., Stephenson, E., Polański, K., Goncalves, A., et al. (2018). Single-cell
reconstruction of the early maternal-fetal interface in humans. Nature *563*, 347–353.

83. Shao, X., Lu, X., Liao, J., Chen, H., and Fan, X. (2020). New avenues for systematically
inferring cell-cell communication: through single-cell transcriptomics data. Protein Cell *11*,
866–880.

84. Browaeys, R., Saelens, W., and Saeys, Y. (2020). NicheNet: modeling intercellular
communication by linking ligands to target genes. Nat. Methods *17*, 159–162.

85. Kumar, M.P. (2018). Analysis of cell-cell interactions using ligand-receptor expression
patterns. Nat Methods.