**Divergent paths, convergent heads: morphological adaptation of head shape to**

**ecological niches in snakes.**

David Hudry[1,2]* and Anthony Herrel[1,3,4,5]

1. Mécanismes Adaptatifs et Evolution, UMR 7179, Muséum national d'Histoire naturelle

CNRS, Paris, France

2. University of Amsterdam, Netherlands

3. Department of Biology, Evolutionary Morphology of Vertebrates, Ghent University, Ghent,

Belgium

4. Department of Biology, University of Antwerp, Wilrijk, Belgium

5. Naturhistorisches Museum Bern, Bern, Switzerland

*Corresponding author

David Hudry: hudry.david@yahoo.com

ORCID D. Hudry: 0009-0001-0322-7409

ORCID A. Herrel: 0000-0003-0991-4434

**Abstract**

Morphological convergence—where distantly related species evolve similar traits in

response to shared ecological pressures—is a hallmark of adaptive evolution. In snakes, head

shape is a key functional trait linked to habitat use and diet, yet the extent and drivers of its

convergence across habitat types and diet remain poorly understood. Here, we used geometric

morphometric analyses to analyze dorsal head shape from photographs of over 266 snake

species, representing ~5% of global snake diversity. Our analyses reveal that head shape

variation is only weakly structured by phylogeny and allometry, but shaped by ecological

specialization. Morphospace patterns reflect distinct differences in species with different

ecologies: fossorial species exhibit compact, broadened heads suited for burrowing; aquatic

species show streamlined profiles for hydrodynamic efficiency; and arboreal snakes tend to

possess elongated skulls for maneuvering in complex habitats. Terrestrial and semi-fossorial

snakes display broad morphospace overlap and an elevated shape disparity highlighting

morphological versatility. While diet covaried significantly with head shape, species with

similar diets did not exhibit strong morphological convergence. These findings underscore the

dominant role of ecology over phylogeny in shaping the evolution of head shape in gape-

limited predators like snakes.

**Keywords:** diet; convergence; head shape; habitat use; snakes

**Introduction**

The environment shapes animal morphology and selects for traits that increase survival and reproductive success. When facing similar ecological pressures, species often develop similar phenotypes (Esquerré & Keogh, 2016). However, some traits appear more impacted than others by ecology (Deepak et al., 2022). Different factors can explain head shape variation including feeding behavior, sexual selection, and habitat use (Arnold & Pough, 1988; Moon et al., 2019). Species with specialized diets, such as mollusks feeders, may, for example evolve asymmetric mandibles in response to whether snails are left of right wound (Pandelis et al., 2023). Sexual selection may also impact head shape, especially if it impacts mate attraction or competitive ability (Fabre et al., 2016). However, most often a multitude of sometimes conflicting selective pressures drive the evolution of any given trait (Huey & Kingsolver, 1989). Consequently, species inhabiting similar niches may follow distinct evolutionary and morphological pathways (Losos et al., 1998). This is especially true for aquatic snakes facing similar environmental pressures developing different body plans and head shapes (Fabre et al., 2016; Segall et al., 2020; Sheratt et al., 2018). Moreover, despite similar ecological pressures, head shapes may diverge across different clades because other factors such as evolutionary history and niche-specific behaviors also shape morphology (Speed & Arbuckle, 2017; Grundler & Rabosky, 2014).

Snakes are an excellent model system to study morphological convergence because of their ecological versatility (Pyron & Burbrink, 2012) and their reliance on the head to feed as gape-limited predators (Moon et al., 2019). Previous studies have suggested that the evolutionary success of snakes is in large part due to their feeding strategies and trophic specialization (Title et al., 2024). For example, feeding-related morphological evolution is evident in sea snakes, where selective pressures related to foraging mode have resulted in

distinct head shapes (Sherratt et al., 2018). Similarly, homalopsid snakes have evolved unique

head shapes driven by the functional demands imposed by the prey they eat (Fabre et al.,

2016). Habitat-related selective pressures also influence snake morphology. *Boidae* and

*Pythonidae*, for instance, exhibit head shape convergence driven by habitat use (Esquerré &

Keogh, 2016).

Previous studies on head shape convergence in snakes have primarily focused on

aquatic and fossorial snakes. For instance, aquatic species frequently develop narrow heads

with upward-facing eyes and nostrils, features that improve their ability to capture prey, move

efficiently through water and spot predators while remaining submerged (Hibbitts and

Fitzgerald, 2005; Vincent et al., 2009; Segall et al., 2016, 2020; Deepak et al., 2022).

Similarly, burrowing snakes possess robust skulls adapted for digging and streamlined bodies

with reduced ventral scales, which help them burrow with minimal friction (Fabre et al., 2016;

Herrel et al., 2008).  Meanwhile, microcephalic sea snakes exhibit distinct skull shapes linked

to developmental changes, representing niche specialization rather than direct functional

convergence (Sherratt et al., 2018; Segall et al., 2020).

Our overarching prediction is that snakes inhabiting similar environments or eating

similar prey will show morphological convergence. Particularly, we predict that fossorial

snakes will exhibit a narrow, V-shaped head facilitating substrate penetration and minimizing

the energetic cost of burrowing (Navas et al., 2004). Head shape in fossorial snakes should,

however, not be influenced by feeding habits as they mostly feed on small soil invertebrates.

Arboreal snakes are predicted to have slender heads to blend in with the environment (i.e.

camouflage), except for mollusk specialists that are expected to have blunt heads. Aquatic

snakes are expected to express disparity in head shape as they feed on a variety of prey in a

variety of underwater habitats. For species that effectively move in different environments

(semi-aquatic, semi-arboreal, semi-fossorial) we predict that they will show transitional

features. We also predict that greater morphological variation will be observed in snakes that

live in transitional habitats and that have non-specialized diets.

**Materials and Methods**

*Sample and ecological classification*

We collected data from 266 species used for habitat use (508 specimens) and 204

species for diet (397 specimens). Animals were housed in three European Natural History

Museums: the National Museum of Natural History in Paris (97 specimens), the Alexander

Koenig Museum in Bonn (316 specimens), and the Museum of Natural Sciences in Brussels

(95 specimens). Our measurements were taken from adult specimens whenever possible.

However, in some cases, only young individuals were available. For some species (i.e. large

pythons and boas), only subadults were measured for practical reasons and the lack of

availability of adults (Table S1). Our sampling aimed to encompass the phylogenetic and

ecological diversity of snakes, while remaining limited by the availability of specimens in

these collections.

Based on the literature, eight habitat use categories were determined (Spawls et al.,

2002; Whitaker & Captain, 2004; Campbell & Lamar, 2004; Marais, 2004; O'Shea, 2005;

Vogel, 2006; Glaw & Vences, 2007; Bartlett & Bartlett, 2009; Das, 2010; O'Shea, 2018;

Allen, 2019; Chippaux, 2019; Pietersen et al., 2021; Egan, 2022; Das, 2023). We classified

the species into 102 terrestrial, 34 arboreal, 19 aquatic, 6 fossorial, 43 semi-arboreal, 33 semi-

aquatic, and 29 semi-fossorial species. We based our ecological categorization on the

biomechanical constraints imposed by the environment (Table S2). For instance, similar

hydrodynamic constraints affect both marine and freshwater snakes leading us to group them

into the same category.

We further use dietary data from the supplementary material of the paper by Title et al.

(2024). Categories were modified and regrouped based on the functional constraints imposed

by the prey. We define a total of ten dietary groups, including 5 generalists, 4 hard egg

feeders, 90 hard prey feeders, 16 invertebrate feeders, 3 mollusk specialists, 2 soft egg

specialists, and 83 soft prey eaters (Table S3). In creating these categories, we grouped

salamanders, lizards, and fish together as they have cylindrical and flexible bodies and as such

should impose similar functional constraints.

*Geometric morphometric data*

For each individual a photo of the head in dorsal view was taken with a Nikon ©

D3100 camera. Snakes were placed on a grid paper for calibration and scale (1 x 1cm squares;

Figure S1A). To perform the geometric morphometric analysis, 25 landmarks and 10 semi-

landmarks were selected (Figure S1B) and defined (Table S4). These landmarks and semi-

landmarks were based on the literature (Manier, 2004; Mangiacotti et al., 2014; Ruane, 2014;

Roth-Monzonetal et al., 2021).

As we based our analysis on landmarks placed on cephalic scales, only species from

the colubrid family, including Colubridae, Elapidae, Lamprophiidae, Homalopsidae, Pareidae,

Psammophiidae, and Pseudoxyrhophiidae were used. A repeatability test was performed using

ten pictures from three individuals of *Lampropeltis californiae* to explore whether the

positioning of the head caused variation in landmark placement. Similarly, landmarks were

placed ten times on these pictures to evaluate error in landmark placement. Variation in

landmark placement due to head positioning as well as landmark placement was limited

(Figure S2). Species were landmarked using the tpsDig software (Rohlf, 2015). The 2D

landmark coordinates were imported into R and subjected to a Generalized Procrustes

Analysis (GPA; Figure S1C). Species means were then computed from Procrustes-aligned

coordinates and used in subsequent analyses. Only species present in both the geometric

morphometric dataset and phylogenetic tree were retained. Geometric morphometric analyses

were conducted using the *geomorph* package (Adams & Otárola-Castillo, 2013). Semi-

landmarks were slid along their tangent directions, minimizing bending energy to optimize

their positions.

*Phylogenetic signal*

To evaluate the extent to which shape variation is structured by phylogeny we used the

*physignal* function in R from the *geomorph* package (Adams & Otárola-Castillo, 2013). This

method is based on a statistical procedure that quantifies the degree of similarity in shape

traits among closely related species relative to what would be expected under a null model of

trait evolution. The function was run with 1000 iterations to generate a null distribution of

phylogenetic signals by randomly reshuffling the phylogenetic tree and comparing the

observed signal to the null distribution. A value close to 1 would indicate a strong

phylogenetic signal, meaning closely related species exhibit more similar shapes than would

be expected by random chance. On the other hand, values significantly less than 1 suggest a

weaker phylogenetic signal, indicating that the variation in shape is less influenced by the

evolutionary history of the species.

*Allometry*

Allometry was quantified using a phylogenetic generalized least squares (PGLS)

model, with shape coordinates as the dependent variable and centroid size as the predictor.

This test aimed to assess size-related variation in shape (i.e., allometry) while accounting for

phylogenetic relatedness among species.

*Phylogenetic principal components analysis*

A phylogenetically-informed principal component analysis (PCA) was performed

using the *phytools* package (Revell, 2012) to explore shape variation in relation to habitat and

diet categories. Eigenvalues were calculated to assess the variance explained by the principal

components, and the first two principal components (PC1 & PC2) were visualized. The

phylomorphospace was visualized by integrating the pruned phylogenetic tree and the

principal component scores, with species colored based on their habitat use and diet using the

*ggphylomorpho* function (Barr, 2017).

*Differences between groups*

A phylogenetic multivariate analysis of variance was performed using Procrustes shape

coordinates as the dependent variables to test whether habitat or diet groups differed in head

shape. The test was run using the *procD.pgls* function from the *geomorph* package (Adams &

Otárola-Castillo, 2013).

*Analyses of covariation*

A phylogenetic Two-Block PLS (Partial Least Squares) analysis was carried out using

the *phylo.integration* function from the *geomorph* package (Adams & Otárola-Castillo, 2013),

to explore whether variation in shape was associated with variation in diet while accounting

for shared evolutionary history. We used continuous proportions of prey items (e.g., hard

prey, soft prey, invertebrates, etc..), allowing for a finer representation of dietary gradients

across species. This matrix was aligned with shape data from Procrustes-aligned landmarks,

and phylogenetic relationships were accounted for.

*Convergence*

To assess morphological convergence among species sharing similar ecological

niches, we employed the *search.conv* function from the *RRphylo* package (Castiglione et al.,

2019). This method evaluates whether species occupying the same habitat or dietary category

exhibit greater phenotypic similarity than expected under a Brownian motion model of trait

evolution. The analysis was conducted on a time-calibrated phylogeny and focused on the

first two principal components of head shape variation (PC1 and PC2). To ensure statistical

robustness, only categories represented by at least three species were retained for analysis

(e.g., soft eggs dietary was not included in some analyses as they were less than 3 individuals

in this group). The function quantifies convergence by calculating the mean angular distance

in trait space between focal taxa and their nearest phenotypic neighbors, both in terms of trait

values (ang.state) and their estimated evolutionary timing (ang.state.time). Significant

convergence is inferred when observed similarity is greater than expected under neutral

evolutionary expectations.

*Disparity analysis*

We performed a morphological disparity analysis to assess the relationship between

shape variation and ecological niches or diet groups while accounting for phylogenetic

relationships, using the *morphol.disparity* function from the *geomorph* package (Adams &

Otárola-Castillo, 2013). We used Procrustes-aligned shape coordinates (GPA) and tested for

differences in morphological disparity across ecological niches and diet categories. A

phylogenetically informed disparity analysis was conducted using a permutation-based

resampling method (1000 iterations), enabling robust comparisons of shape variance between

habitat types and diet groups.

**Results**

*Phylogenetic signal*

The observed value of K is low for shape (K = 0.0964), indicating a weak

phylogenetic signal (Table 1). Furthermore, the result was not statistically significant,

suggesting that the morphological shape variation among species is not strongly structured by

phylogenetic relationships. This implies that closely related species do not resemble each

other more than expected by chance in terms of overall shape.

*Allometry*

The phylogenetic generalized least squares (PGLS) analysis revealed a statistically

significant relationship between shape and centroid size (Table 2). Despite a relatively low

proportion of variance explained ($R^2 = 0.027$), the allometric signal in head shape variation

was significant ($P < 0.005$).

*Variation in head shape*

In our principal component analysis, 97% of the variation in head shape is explained

by the first two principal components (PC1: 73.39%; PC2: 23.99%; Figure 1). The

phylomorphospace of head landmarks reveals structuring of head shape variation among

snakes. Fossorial species, characterized by low PC1 and PC2 values, exhibit elongated and

broad posterior skulls (Figure 1A). Arboreal and semi-arboreal snakes, by contrast, tend to

occupy higher PC1 values and display slender, elongated skulls. Aquatic and semi-aquatic

species are more dispersed along PC1, with skulls often appearing streamlined. Terrestrial

snakes are widely spread across the morphospace a and semi-fossorial and semi-arboreal

species tend to fall in intermediate regions of the morphospace.

In terms of diet, species consuming soft-bodied prey (e.g., snakes, lizards) generally

exhibit higher PC1 values, associated with elongated and narrower heads (Figure 1B; Figure

2). Conversely, species feeding on harder prey (e.g., rodents, birds, mollusks) are found in the

lower range of PC1 and often exhibit broader, more robust skulls. Mollusk specialists and

other durophagous feeders show this trend most strongly. PC2 variation also reflects dietary

adaptations: species with lower PC2 values, such as soft-prey specialists and generalists, tend

to have narrower skulls, while invertebrate feeders cluster around broader and more robust

head shapes. Deformation grids show that many species exhibit a combination of longer,

narrower snouts with posterior skull widening—contributing to the shared V-shaped

morphology in fossorial, semi-aquatic, and some hard-prey specialists (Figure 1). Overall,

generalist feeders and species with diverse prey types occupy wide, overlapping regions of the

morphospace, indicating that head shape in snakes is shaped by a complex interplay between

habitat use and dietary specialization.

*Differences between habitat groups*

The MANOVA indicates significant differences between habitat groups in head shape

(Table 3). When adding $Log_{10}$-transformed centroid size as a covariate, we found significant

effects of habitat, centroid size and the interaction between the two (Table 3). The interaction

between habitat and centroid size was also significant. Moreover, post-hoc pairwise

comparisons further revealed that fossorial snakes, as well as semi-fossorial, differ

significantly from all ecological groups (Table 4). Additionally, semi-aquatic and semi-

arboreal differ from terrestrial snakes in head shape.

*Differences between diet groups*

There was a trend toward head shape differentiation among diet groups, based on dominant dietary regimes, as revealed by the phylogenetic MANOVA (Table 5). When adding $Log_{10}$-transformed centroid size as a covariate, this tendency remained. Centroid size did differ between dietary groups, while interactions were not significant (Table 5). Post hoc pairwise comparisons indicated a significant difference in head shape between species consuming hard prey and those consuming soft prey ($P < 0.05$; Table 6). There were trends observed between hard prey and soft eggs feeders ($P = 0.090$), as well as between invertebrates and soft eggs feeders ($P = 0.092$).

*Analyses of covariance*

Our results revealed a significant covariation between head shape and habitat use (r-PLS = 0.38, Z = 2.67, $P < 0.01$), as well as between head shape and diet (r-PLS = 0.59, Z = 3.82, $P < 0.05$) (Table 7). This indicates that species occupying different habitat types or relying on different dietary resources differ in head shape. Specifically, the distribution of species along the first PLS axis suggests that morphological variation is aligned with ecological gradients (Figure 3). Aquatic and fossorial species appear to cluster separately from arboreal or terrestrial taxa, reflecting adaptive morphological specializations for swimming, burrowing, or climbing. In the case of diet, species that consume hard prey tend to separate from those feeding primarily on soft-bodied prey or eggs, suggesting that cranial shape variation may be driven by biomechanical demands related to prey processing (Figure 4).

*Convergence*

The convergence test results revealed contrasting patterns in morphological evolution

depending on whether species were grouped by habitat use or diet (Table 8). Among habitat

groups, strong evidence for morphological convergence was found within aquatic and

fossorial species. These two groups showed significantly lower mean angles of phenotypic

distance (ang.state = 58.46 and 11.29, respectively; $P < 0.01$ for both). These results were also

supported by the timing-based convergence metric, indicating that the convergent traits

evolved synchronously within these clades (ang.state.time = 331.51 and 98.38 respectively; $P$

$< 0.01$ for both groups). In contrast, no significant convergence was detected within the other

habitat groups, such as terrestrial, arboreal, semi-arboreal, semi-aquatic, or semi-fossorial

species (Table 8). While some of these groups (e.g., arboreal or semi-aquatic) showed

moderately low angles of phenotypic distance, the results were not statistically significant.

No dietary group showed statistically significant morphological convergence, although

Mollusks approached the lowest angle values (ang.state = 53.65, $P = 0.165$), hinting at

possible trends that did not reach significance, potentially due to the very small sample size.

All other dietary groups—including hard prey, soft prey, invertebrates, and hard egg

specialists—exhibited non-significant convergence metrics. Timing-based convergence

likewise yielded no significant patterns, indicating that when morphological similarities were

observed, they did not evolve at the same time across lineages.

*Disparity analysis*

Our findings indicate that semi-fossorial species exhibit the highest morphological

disparity (0.0122; Table 9). Aquatic species displayed the second highest score in

morphological disparity (0.0116). Moderate levels of disparity were observed in the following

categories: terrestrial species (0.0112) and semi-arboreal species (0.0111). Conversely,

fossorial species (0.0108), semi-aquatic species (0.0108) and arboreal species (0.0087)

demonstrated the lowest non-zero variances. Pairwise absolute differences reveal that the

greatest differences occur between aquatic and arboreal species (0.003), as well as between

arboreal and semi-fossorial species (Table S5). However, pairwise absolute differences

among the other habitat types remain small and statistically non-significant (Table S6).

Our findings further indicated that species consuming soft prey exhibit the highest

morphological disparity (0.018; Table 9). Species consuming hard prey and mollusk displayed

notable morphological disparity (0.011 and 0.0085 respectively). Moderate levels of disparity

were observed in generalist species (0.0071), invertebrate feeders (0.0088) and hard-egg

consumers (0.0086). Conversely, soft-egg consumers (0.0025) demonstrated the lowest non-

zero variances. Pairwise absolute differences confirm that soft-prey consumers exhibit

significantly higher morphological disparity compared to most other dietary categories,

particularly soft-egg, mollusk, and hard-egg consumers (Table S7). Yet, pairwise absolute

differences and significance levels among the other diet categories remain minimal and non-

significant, except marginally for the comparison between soft-prey and soft-egg consumers

(Table S8).

**Discussion**

Our analyses demonstrate that snake head shape variation is not primarily governed by

phylogenetic relatedness or body size, but rather reflects adaptations to ecological contexts

and feeding strategies. The weak and statistically non-significant phylogenetic signal (K =

0.0964) indicates that closely related species do not resemble each other in head shape more

than expected by chance. This supports the idea that ecological and functional demands may

override phylogenetic constraints, enabling repeated and independent morphological

adaptations across lineages. Similar patterns have been observed by da Silva et al. (2018),

who reported convergent skull shapes in aquatic and fossorial snakes regardless of

phylogenetic affiliation, emphasizing the influence of habitat-driven functional demands.

Likewise, da Silva et al. (2017) found that trophic specialization drove significant cranial

divergence among Neotropical snakes with limited phylogenetic signal, suggesting that diet,

like habitat, can supersede phylogenetic inertia. On the other hand, Esquerré and Keogh

(2016) found a moderate phylogenetic influence on head shape in Australian elapids, though

ecological divergence still played a key role—highlighting potential clade-specific variation

in the balance between constraint and adaptation. Although our allometric test revealed a

statistically significant relationship between centroid size and shape, the effect was small ($R^2$

= 0.025), suggesting that body size alone explains little of the head shape variation. This

decoupling indicates that selection acts more directly on cranial morphology than on ody size

(da Silva et al., 2018).

The distribution of species in morphospace reinforces this interpretation. Ecological

specialization corresponds to distinct regions of cranial variation, especially among fossorial,

arboreal, and semi-aquatic species. Fossorial snakes, with their V-shaped and posteriorly

broadened skulls, occupy a unique morphospace region—consistent with mechanical

requirements for head-first burrowing and maneuvering through confined spaces (da Silva et

al., 2018; Strong et al., 2020; Pandelis et al., 2023). Arboreal and semi-arboreal species

exhibit slender, elongated skulls, which likely facilitate navigation and prey capture in

structurally complex habitats (Herrel et al., 2008; Klaczko et al., 2016). Similarly, aquatic and

semi-aquatic species often show streamlined cranial shapes associated with reduced

hydrodynamic drag (Segall et al., 2016 – 2020; Deepak et al., 2022).

The phylogenetic MANOVA confirmed significant overall differences in head shape

among habitat groups, with fossorial and semi-fossorial species showing distinct

morphologies compared to most others. This pattern highlights the strong biomechanical

constraints imposed by a subterranean lifestyle. Interestingly, semi-fossorial species exhibited

the highest disparity among all habitat types. This elevated variation may stem from their

intermediate ecological position, requiring adaptations to both surface and subsurface

environments. Variability in substrate type, time spent underground, and degrees of cranial

reinforcement or kinesis could contribute to this morphological diversity (Herrel et al., 2008;

da Silva et al., 2018; Tingle et al., 2024).

Aquatic species displayed relatively high morphological disparity, likely reflecting the

broad ecological diversity of aquatic environments and the wide array of prey types these

snakes exploit. This is consistent with findings from Segall et al. (2016, 2020) and Deepak et

al. (2022) who showed that aquatic snakes adapted to different flow regimes, prey sizes, and

foraging strategies often evolve distinct cranial morphologies. Water acts both as a

biomechanical constraint and a driver of morphological diversification (Sherratt et al., 2018).

In contrast, fossorial species displayed more moderate disparity levels, which may reflect

stronger functional constraints linked to a fully subterranean lifestyle. Such species often

converge toward compact, reinforced head shapes adapted to head-first burrowing, thereby

limiting morphological variation. In arboreal snakes, stability during strikes and secure

navigation along narrow, flexible supports may favor streamlined, stable cranial morphologies

(Klaczko et al., 2016; Alencar et al., 2017). Similarly, semi-aquatic species may face

conflicting selective pressures between aquatic locomotion and terrestrial maneuverability,

potentially resulting in conservative morphologies shaped by functional compromise (Sherratt

et al., 2018; Segall et al., 2020).

Dietary specialization also contributes meaningfully to head shape variation. Species

feeding on soft-bodied prey such as snakes, amphibians, or fish often display elongated,

narrow heads—likely facilitating the consumption of these prey. In contrast, snakes that

consume hard or bulky prey, including mammals, birds, or shelled invertebrates, possess

broader, more robust skulls adapted to gripping and processing resistant prey. In 2019,

Perkins & Eason showed that sympatric water snake species exhibited head shapes matching

their dietary preferences: narrow heads were associated with fish diets, and broad heads with

amphibians. Generalists spanned a wide morphospace, reflecting cranial versatility and

behavioral plasticity (Deepak et al., 2022; Pandelis et al., 2023). Soft-prey consumers

exhibited the highest morphological disparity, likely reflecting the variety of functional

solutions for handling elongate, mobile prey (Segall et al., 2020; Pandelis et al., 2023). In

contrast, mollusk and soft-egg specialists clustered tightly in morphospace, suggesting

convergence toward a limited set of biomechanically effective forms for durophagy (Pandelis

et al., 2023).

Convergence analyses further illuminate the strong influence of habitat-driven

functional constraints. Significant morphological convergence was detected among fossorial

and aquatic species, likely reflecting parallel adaptations to movement through compact

substrates or fluid environments (Herrel et al., 2008; Segall et al., 2016). Similarly, our

detection of convergence among fossorial taxa supports the view that dealing with resistant

media—whether water or soil—can lead to similar head shapes across distantly related

lineages. In contrast, we did not detect significant convergence among dietary groups.

Although species consuming similar prey types often exhibited analogous head shapes, these

patterns lacked statistical support for convergence. This contrasts with findings such as

Perkins and Eason (2019), who documented strong diet-morphology correlations in sympatric

species. The broader inconsistency may be due to variability in foraging behavior, strike

mechanics, or ecological context across clades, limiting the emergence of shared

morphological optima. Furthermore, lineages may face developmental or functional

constraints that shape divergent cranial solutions to similar dietary challenges (Segall et al.,

2020; Pandelis et al., 2023). The exclusion of underrepresented groups, such as mollusk

feeders or egg specialists, from our convergence analyses may also limit our ability to detect

certain trends.

Taken together, our findings reinforce the growing consensus that habitat use and

dietary specialization play significant but distinct roles in shaping the evolution of head shape

in snakes. Environments imposing strong biomechanical constraints, such as arboreal and

fossorial habitats, clearly shape cranial form. However, diet—especially differences in prey

hardness and body shape—exerts an even stronger and more consistent influence on cranial

morphology across lineages, as highlighted by the PLS analysis. The pronounced covariation

among head shape, habitat, and diet underscores that ecological specialization involves a

dynamic interplay of selective pressures, with both habitat use and feeding ecology jointly

driving morphological evolution in snakes.

Beyond analytical methods, a number of biological and data-driven limitations should

be acknowledged. First, the categorization of ecological niches remains somewhat coarse and

potentially subjective. For example, semi-fossorial species may share more ecological traits

with terrestrial than fully fossorial snakes, and semi-aquatic species vary widely in their

dependence on aquatic environments (Esquerré & Keogh, 2016; Silva et al., 2017).

Incorporating quantitative ecological data—such as habitat features (e.g., branch density, soil

compaction, water salinity) or substrate use (e.g., frequency of use of forest soil, mud, or tree

species)—could refine these categories and better capture the ecological continuum

(Manjarrez et al., 2017; Tingle et al., 2024). Second, although our dietary classifications are

grounded in empirical data, they could be enhanced by integrating quantitative prey traits—

such as size and hardness—that more accurately represent the functional challenges faced by

predators, rather than relying solely on prey shape (Vincent et al., 2009; Da Silva et al., 2018).

For instance, both birds and frogs may be categorized as soft prey, yet their ecologies and

physical properties differ markedly, potentially requiring distinct feeding adaptations (Segall

et al., 2020; Deepak et al., 2022). Understanding the functional properties of prey would thus

provide more nuanced insights into head shape evolution. Furthermore, the inclusion of

behavioral variables—such as foraging strategy or strike kinematics—would significantly

strengthen the link between morphology and ecological function (Herrel et al., 2008; Segall et

al., 2016; Moon et al., 2019). These traits are likely to interact with head shape in complex

ways that are not fully captured by static ecological or dietary categories alone. Additionally,

focusing on taxonomically or ecologically specialized clades (e.g., snail-eaters, egg

specialists) could shed light on extreme morphological adaptations that illustrate more clearly

how head shape evolves under strong selective pressure (Pandelis et al., 2023; Palci et al.,

2024).

Our study underscores the complex and multifactorial nature of head shape evolution

in snakes, driven more by ecological function than phylogenetic history or allometry.

Building on this, Moon et al. (2019) emphasize the remarkable adaptability of the snake

feeding system, noting that head morphology reflects not only dietary and ecological

demands, but also performance constraints like gape limitation, strike kinematics, and prey

handling strategies. They highlight how head shape plays a central role in functional

performance—such as strike accuracy, hydrodynamics, and force generation—yet also

caution that our understanding remains limited by the narrow taxonomic and ecological scope

of most studies. To address this, future research should aim for a more integrative approach—

linking head morphology with detailed data on strike mechanics, bite force, and prey-handling

behaviors across ecological contexts. For instance, striking behaviors vary between aquatic

and terrestrial environments, and bite performance has only been measured in a few species,

suggesting that much remains to be explored (Bartolon et al., 2013; Penning, 2017; Moon et

al., 2019). Prey manipulation could have a significant effect on the overall morphology of

snakes, and could be of great interest in understanding the adaptability of certain niches to

different prey items (Crotty & Jane, 2017). Furthermore, intraspecific variation in feeding

plasticity (Manjarrez et al., 2017; Bonnet et al., 2020; Ortiz-Medina et al., 2022), as well as

ontogenetic (Lopez et al., 2013; Patterson et al., 2022) and sexual dimorphism (Borczyk et al.,

2021; Shine & Goiran, 2021) in head shape and diet, could further illuminate the selective

pressures shaping head form. Expanding datasets to include internal morphology, muscle

architecture, and functional performance metrics would not only validate the ecological

relevance of head shape variation but also test longstanding hypotheses about convergence

and specialization in snake evolution. In doing so, we can build a more mechanistic

understanding of how morphology translates into ecological success.

**Conclusion**

Our findings reveal that snake head shape is shaped less by phylogenetic relatedness

or body size and more by ecological function and dietary specialization. Morphological

variation in cranial form reflects repeated, and often convergent, adaptations to environmental

demands such as burrowing, climbing, or swimming, as well as to prey capture and

processing. This is especially apparent in fossorial and semi-aquatic snakes, where head

shapes align with biomechanical requirements like substrate penetration or hydrodynamic

efficiency, and in durophagous species, which exhibit robust skulls for handling hard prey.

Despite a weak phylogenetic signal and minimal allometric influence, we detected strong

associations between head shape and ecological context, reinforced by disparity patterns and

PLS analyses. Intriguingly, high morphological disparity within narrow ecological

categories—such as fossorial and aquatic niches—suggests the evolution of multiple

functional solutions to similar environmental challenges, likely shaped by fine-scale habitat

variation, prey diversity, or biomechanical constraints.

Our use of cephalic scale-based morphometrics offers a pragmatic and ecologically relevant

proxy for head shape in large comparative datasets, but we recognize its limitations. Internal

cranial architecture, muscle arrangement, and soft tissue all contribute to ecological

performance and remain largely unexplored in this context. Moving forward, a more

integrative approach—incorporating performance data (e.g., bite force, strike dynamics),

behavioral traits (e.g., foraging strategy, prey handling), and finer-scale ecological metrics—

will be essential for linking form to function. Investigating intraspecific variation, ontogeny,

and sexual dimorphism can further illuminate how head shape evolves under diverse selective

pressures. By bridging morphology with performance and behavior, future research can

deepen our understanding of how ecological demands shape evolutionary trajectories, not

only in snakes but across vertebrate lineages navigating similar functional landscapes.

**Acknowledgments**

The authors would like to thank Nicolas Vidal from the Muséum National d'Histoire
Naturelle in Paris, Dennis Rödder and Morris Fleck from the Museum Alexander Koenig in
Bonn, and Olivier Pauwels from the Museum of Natural Sciences of Brussels for access to
specimens.

**References**

Adams, D. C., & Otárola-Castillo, E. (2013). Geomorph: An r package for the collection and
analysis of geometric morphometric shape data. *Methods in Ecology and Evolution*, *4*(4),
393–399.

Alencar, L. R. V., Martins, M., Burin, G., & Quental, T. B. (2017). Arboreality constrains
morphological evolution but not species diversification in vipers.

Allen, L. R. (2019). *Venomous Snakes of Australia and Oceania*. CSIRO Publishing.

Arnold, S. J., & Pough, F. H. (1983). Allometry and the evolution of gape-limited predation
in snakes. *American Naturalist*, 121(2), 151–170.

Barr , W. A. (2017). ggphylomorpho, version 0.2.

Bartlett, R. D., & Bartlett, P. P. (2009). *Guide and Reference to the Snakes of Western North*

*America (North of Mexico) and Hawaii*. University Press of Florida

Bartolon, G., Stepp-Bolling, C., Jackson, K., Kley, N., & Brainerd, E. (2013). To Crush or to

Engulf Your Victim: a biomechanical trade-off in lizards versus snakes. *American Naturalist*

(Vol. 66).

Bonnet, X., Brischoux, F., Briand, M., & Shine, R. (2021). Plasticity matches phenotype to

local conditions despite genetic homogeneity across 13 snake populations. *Proceedings of the*

*Royal Society B: Biological Sciences*, *288*(1943).

Borczyk, B., Paśko, Ł., Kusznierz, J., & Bury, S. (2021). Sexual dimorphism and skull size

and shape in the highly specialized snake species, *Aipysurus eydouxii* (Elapidae:

Hydrophiinae). *PeerJ*, *9*.

Campbell, J. A., & Lamar, W. W. (2004). *The Venomous Reptiles of the Western Hemisphere*.

Cornell University Press.

Castiglione, S., Serio, C., Tamagnini, D., Melchionna, M., Mondanaro, A., di Febbraro, M.,

Profico, A., Piras, P., Barattolo, F., & Raia, P. 2019. A new, fast method to search for

morphological convergence with shape data. *PLoS ONE*, *14*(12).

Chippaux, J.-P. (2019). *Snakes of Central and Western Africa*. Johns Hopkins University

Press.

Crotty, T. L., & Jayne, B. C. (2014). Trade-offs between eating and moving: what happens to

the locomotion of slender arboreal snakes when they eat big prey? *Biological Journal of the*

*Linnean Society*, 114, 446-458.

da Silva, F. O., Fabre, A. C., Savriama, Y., Ollonen, J., Mahlow, K., Herrel, A., Müller, J., &

Di-Poï, N. (2018). The ecological origins of snakes as revealed by skull evolution. *Nature*

*Communications*, *9*(1).

Das, I. (2010). *A Field Guide to the Reptiles of South-East Asia*. New Holland Publishers.

Das, I. (2023). *Snakes of South-East Asia*. Bloomsbury Publishing.

Deepak, V., Gower, D. J., & Cooper, N. (2022). Diet and habit explain head-shape

convergences in natricine snakes. *Journal of Evolutionary Biology*, *36*(2), 399–411.

Egan, D. (2022). *Snakes of the Middle East*. Bloomsbury Wildlife.

Esquerré, D., & Scott Keogh, J. (2016). Parallel selective pressures drive convergent

diversification of phenotypes in pythons and boas. *Ecology Letters*, *19*(7), 800–809.

Fabre, A.-C., Bickford, D., Segall, M., & Herrel, A. (2016). The impact of diet, habitat use,

and behaviour on head shape evolution in homalopsid snakes. *Biological Journal of the*

*Linnean Society,* 118, 634-647.

Glaw, F., & Vences, M. (2007). *A Field Guide to the Amphibians and Reptiles of*

*Madagascar*. Vences & Glaw Verlag.

Grundler, M. C., & Rabosky, D. L. (2014). Trophic divergence despite morphological

convergence in a continental radiation of snakes. *Proceedings of the Royal Society B:*

*Biological Sciences*, 281(1795), 20140413.

Herrel, A., Huyghe, K., Vanhooydonck, B., Backeljau, T., Breugelmans, K., Grbac, I., van

Damme, R., & Irschick, D. J. (2008). Rapid large-scale evolutionary divergence in

morphology and performance associated with exploitation of a different dietary resource.

Hibbitts, T. J., & Fitzgerald, L. A. (2005). Morphological and ecological convergence in two

natricine snakes.

Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm

performance. *Trends in Ecology & Evolution*, 4(5), 131-135.

Klaczko, J., Sherratt, E., & Setz, E. Z. F. (2016). Are diet preferences associated to skulls

shape diversification in xenodontine snakes? *PLoS ONE*, *11*(2).

López, M. S., Manzano, A. S., & Prieto, Y. A. (2013). Ontogenetic variation in head

morphology and diet in two snakes (Viperidae) from Northeastern Argentina. *Journal of*

*Herpetology*, *47*(3), 406–412.

Lettoof, D. C., Parkin, T., Jolly, C. J., de Laive, A., & von Takach, B. (2023). Snake life
history traits and their association with urban habitat use in a tropical city. *Urban Ecosystems*,
*26*(2), 433–445.

Losos, J. B., Jackman, T. R., Larson, A., de Queiroz, K., & Rodríguez-Schettino, L. (1998).
Contingency and determinism in replicated adaptive radiations of island lizards. *Science*,
279(5359), 2115–2118.

Mangiacotti, M., Giansanti, A., & Sacchi, R. (2014). Geometric morphometrics reveals inter-
and intraspecific shape variation in head morphology of European lacertid lizards.
*Zoologischer Anzeiger*, 253(2), 83–93.

Manier, M. K. (2004). Geographic variation in the long-nosed snake, *Rhinocheilus lecontei*
(*Colubriade*): beyond the subspecies debate. In *Biological Journal of the Linnean Society*, 83,
65-85.

Manjarrez, J., Macías Garcia, C., & Drummond, H. (2017). Morphological convergence in a
Mexican garter snake associated with the ingestion of a novel prey.

Manjarrez, J., Pacheco-Tinoco, M., & Venegas-Barrera, C. S. (2017). Intraspecific variation
in the diet of the Mexican garter snake Thamnophis eques. *PeerJ*, *2017*(11).

Marais, J. (2004). *A Complete Guide to the Snakes of Southern Africa*. Struik Publishers.

Moon, B. R., Penning, D. A., Segall, M., & Herrel, A. (2019). Feeding in Snakes: Form,
Function, and Evolution of the Feeding System (pp. 527–574).

Navas, C. A., Antoniazzi, M. M., Jared, C., & Kohlsdorf, T. (2004). Morphological and
physiological specialization for digging in amphisbaenians, an ancient lineage of fossorial
vertebrates. *Journal of Experimental Biology*, 207(14), 2433–2441.

Ortiz-Medina, J. A., Cedeño-Vázquez, J. R., González-Solís, D., & Mata-Silva, V. (2022).
Intraspecific Variation in Diet Composition and Morphology of Yucatecan Cantils
(*Agkistrodon russeolus* Gloyd 1972 [Serpentes: Viperidae]). *Herpetologica*, *78*(4), 244–252.

O'Shea, M. (2005). *Reptiles et Amphibiens*. Delachaux et Niestlé.

O'Shea, M. (2018). *The Book of Snakes: A Life-Size Guide to Six Hundred Species from*

*Around the World*. University of Chicago Press.

Palci, A., Lee, M. S. Y., Aubret, F., & Sherratt, E. (2024). Multiple Paths to a Larger Mouth:

Morphological Divergence in Two Island Populations of Australian Tiger Snakes (Notechis

scutatus). *Evolutionary Biology*.

Pandelis, G. G., Grundler, M. C., & Rabosky, D. L. (2023). Ecological correlates of cranial

evolution in the megaradiation of dipsadine snakes. *BMC Ecology and Evolution*, *23*(1).

Patterson, M., Wolfe, A. K., Fleming, P. A., Bateman, P. W., Martin, M. L., Sherratt, E., &

Warburton, N. M. (2022). Ontogenetic shift in diet of a large elapid snake is facilitated by

allometric change in skull morphology. *Evolutionary Ecology*, *36*(4), 489–509.

Penning, D. A. (2017). The scaling of bite force and constriction pressure in kingsnakes

(*Lampropeltis getula): *Proximate determinants and correlated performance. *Integrative*

*Zoology*, *12*(2), 121–131.

Perkins, M. W., & Eason, P. K. (2019). The relationship of head morphology and diet among

three sympatric watersnake species. *Amphibia Reptilia*, *40*(1), 65–77.

Pietersen, D. W., Conradie, W., & Branch, W. (2021). *Snakes and Other Reptiles of Zambia*

*and Malawi*. Struik Nature.

Pyron, R. A., & Burbrink, F. T. (2012). Extinction, ecological opportunity, and the origins of

global snake diversity. *Evolution*, *66*(1), 163–178.

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology & Evolution, 3, 217–223.

Rohlf, F. J. (2015). tpsDig, version 2.0

Roth-Monzón, A. J., Devitt, T. J., Ríos-Muñoz, C. A., Parra-Olea, G., Campbell, J. A., &

Flores-Villela, O. (2021). Geographic Variation in the Green Rat Snake *Senticolis triaspis*

(Squamata: Colubridae): Evidence from Mitochondrial DNA, Morphology, and Niche

Modeling. *Ichthyology and Herpetology*, *109*(4), 1026–1035.

Ruane, S. (2015). Using geometric morphometrics for integrative taxonomy: An examination
of head shapes of milksnakes (genus Lampropeltis). *Zoological Journal of the Linnean*
*Society*, *174*(2), 394–413.

Segall, M., Cornette, R., Fabre, A. C., Godoy-Diana, R., & Herrel, A. (2016). Does aquatic
foraging impact head shape evolution in snakes? *Proceedings of the Royal Society B:*
*Biological Sciences*, *283*(1837).

Segall, M., Cornette, R., Godoy-Diana, R., & Herrel, A. (2020). Exploring the functional
meaning of head shape disparity in aquatic snakes. *Ecology and Evolution*, *10*(14), 6993–
7005.

Sherratt, E., Rasmussen, A. R., & Sanders, K. L. (2018). Trophic specialization drives
morphological evolution in sea snakes. *Royal Society Open Science*, *5*(3).

Shine, R., & Goiran, C. (2021). Sexual dimorphism in size and shape of the head in the sea
snake *Emydocephalus annulatus* (Hydrophiinae, Elapidae). *Scientific Reports*, *11*(1).

Silva, F. M., Prudente, A. L. da C., Machado, F. A., Santos, M. M., Zaher, H., & Hingst-
Zaher, E. (2017). Aquatic adaptations in a Neotropical coral snake: A study of morphological
convergence. *Journal of Zoological Systematics and Evolutionary Research*, *56*(3), 382–394.

Spawls, S., Howell, K., Drewes, R., & Ashe, J. (2002). *A Field Guide to the Reptiles of East*
*Africa: Kenya, Tanzania, Uganda, Rwanda, and Burundi*. A&C Black Publishers.

Speed, M. P., & Arbuckle, K. (2017). Quantification provides a conceptual basis for
convergent evolution. *Biological Reviews*, *92*(2), 815–829.

Strong, C. R. C., Palci, A., & Caldwell, M. W. (2020). Insights into skull evolution in
fossorial snakes, as revealed by the cranial morphology of *Atractaspis irregularis* (Serpentes:
Colubroidea). *Journal of Anatomy*, *238*(1), 146–172.

Title, P. O., Singhal, S., Grundler, M. C., Costa, G. C., Alexander Pyron, R., Colston, T. J.,
Grundler, M. R., Prates, I., Stepanova, N., H Jones, M. E., Q Cavalcanti, L. B., Colli, G. R.,
Di-Poï, N., Donnellan, S. C., Moritz, C., Mesquita, D. O., Pianka, E. R., Smith, S. A., Vitt, L.
J., & Rabosky, D. L. (n.d.). The macroevolutionary singularity of snakes. *Science,* 383, 918-
923.

Vincent, S. E., Brandley, M. C., Herrel, A., & Alfaro, M. E. (2009). Convergence in trophic

morphology and feeding performance among piscivorous natricine snakes. *Journal of*

*Evolutionary Biology*, *22*(6), 1203–1211.

Vogel, G. (2006). *Venomous Snakes of Asia*. Edition Chimaira.

Whitaker, R., & Captain, A. (2004). *Snakes of India: The Field Guide*. Draco Books.

Table 1: Blomberg's K values and significance.

|       | K      | *P*    |
|-------|--------|--------|
| Shape | 0.0964 | 0.9071 |

620  Table 2: Summary of the allometric test results using a phylogenetic generalized least squares

621  (PGLS) model with log-transformed centroid size as the predictor of shape. The table reports

622  the degrees of freedom (Df), sums of squares (SS), F statistic, effect size (Z), and associated

623  p-value (*P*) for the regression.

|  | Df | SS | $R^2$ | F | Z | r | *P* |
|---|---|---|---|---|---|---|---|
| Centroid size | 1 | 4.12 | 0.027 | 6.82 | 2.63 | 0.16 | *P* < 0.05 |

625 Table 3: Results of the phylogenetic MANCOVA and MANOVA testing the effect of habitat

626 use on shape with and without accounting for allometry. The table presents the degrees of

627 freedom (Df), sum of squares (SS), mean squares (MS), proportion of explained variance

628 $(R^2)$, F-statistic (F), and p-value.

| | Df | SS | MS | $R^2$ | $F$ | $P$ |
|---|---|---|---|---|---|---|
| MANCOVA | | | | | | |
| Habitat | 6 | 24.94 | 4.16 | 0.13 | 6.72 | $P < 0.05$ |
| Centroid size | 1 | 2.91 | 2.91 | 0.01 | 4.71 | $P < 0.05$ |
| Interaction | 6 | 11.46 | 1.91 | 0.06 | 2.73 | $P < 0.05$ |
| MANOVA | | | | | | |
| Habitat | 6 | 24.94 | 4.16 | 0.13 | 6.33 | $P < 0.05$ |

630    Table 4: Results of the Tukey post hoc test for pairwise comparisons of morphological

631    disparity across niche groups. The table presents the effect size (d), upper confidence limit at

632    95% (UCL 95%), Z-score (Z), and p-value for each comparison.

| Comparison | d | UCL (95%) | Z | P |
|---|---|---|---|---|
| Fossorial – aquatic | 0.17 | 0.13 | 2.20 | *P* < 0.05 |
| Fossorial – arboreal | 0.17 | 0.13 | 2.17 | *P* < 0.05 |
| Fossorial – semi-aquatic | 0.18 | 0.13 | 2.28 | *P* < 0.05 |
| Fossorial – semi-arboreal | 0.17 | 0.13 | 2.25 | *P* < 0.05 |
| Fossorial – semi-fossorial | 0.20 | 0.13 | 2.40 | *P* < 0.05 |
| Fossorial – terrestrial | 0.18 | 0.13 | 2.23 | *P* < 0.05 |
| Semi-fossorial - arboreal | 0.07 | 0.06 | 2.12 | *P* < 0.05 |
| Semi-fossorial – semi-aquatic | 0.09 | 0.07 | 2.34 | *P* < 0.05 |
| Semi-fossorial – semi-arboreal | 0.06 | 0.05 | 1.94 | *P* < 0.05 |
| Semi-fossorial – terrestrial | 0.09 | 0.04 | 2.91 | *P* < 0.05 |
| Semi-aquatic – terrestrial | 0.08 | 0.06 | 2.35 | *P* < 0.05 |
| Semi-arboreal – terrestrial | 0.05 | 0.05 | 1.89 | *P* < 0.05 |

634    Table 5: Results of the phylogenetic MANCOVA and MANOVA testing the effect of diet on

635    head shape. The table presents the degrees of freedom (Df), sum of squares (SS), mean

636    squares (MS), proportion of explained variance ($R^2$), F-statistic (F), and p-value.

| | Df | SS | MS | $R^2$ | *F* | *P* |
|---|---|---|---|---|---|---|
| MANCOVA | | | | | | |
| Diet | 5 | 6.78 | 1.36 | 0.04 | 1.99 | 0.083 |
| Centroid size | 1 | 5.85 | 5.85 | 0.04 | 8.59 | $P < 0.05$ |
| Interaction | 5 | 5.75 | 1.15 | 0.04 | 1.69 | 0.141 |
| MANOVA | | | | | | |
| Diet | 5 | 6.78 | 1.36 | 0.05 | 1.89 | 0.089 |

638    Table 6: Results of the Tukey post hoc test for pairwise comparisons of morphological

639    disparity across diet groups. The table presents the effect size (d), upper confidence limit at

640    95% (UCL 95%), Z-score (Z), and p-value (p.value) for each comparison. Significant p-

641    values are denoted by an asterisk (*).

| Comparison | d | UCL (95%) | Z | $P$ |
|---|---|---|---|---|
| Hard prey – Soft prey | 0.04 | 0.04 | 2.08 | $P < 0.05$ |
| Hard prey – Soft egg | 0.16 | 0.21 | 1.46 | 0.090 |
| Invertebrates – Soft eggs | 0.16 | 0.20 | 1.40 | 0.092 |

643 Table 7: Results of the two-block partial least squares (PLS) analysis assessing the

644 relationship between morphology and habitat use and morphology and diet while accounting

645 for phylogeny. The table reports the correlation coefficient from the PLS analysis (r-PLS), the

646 standardized effect size (Z-score), and the associated p-value.

| | r-PLS | Effect size | $P$ |
|---|---|---|---|
| Habitat | 0.38 | 2.67 | $P < 0.01$ |
| Diet | 0.59 | 3.82 | $P < 0.01$ |

648 Table 8: Results of the convergence test. State1 and state2 correspond to pair of ecological

649 groups being compared. The mean angles of phenotypic distance are reported using ang.state

650 (state) and the test of convergence with timing of diversification using ang.state.time (time),

651 with the corresponding p-values.

| | State | *P* | Time | *P* |
|---|---|---|---|---|
| **Niche** | | | | |
| terrestrial | 90.31 | 0.736 | 518.63 | 0.984 |
| arboreal | 85.40 | 0.087 | 478.54 | 0.271 |
| semi-arboreal | 91.41 | 0.886 | 507.65 | 0.715 |
| aquatic | 58.46 | *P < 0.01* | 331.51 | *P < 0.01* |
| semi-aquatic | 85.42 | 0.078 | 538.57 | 0.937 |
| fossorial | 11.29 | *P < 0.01* | 98.38 | *P < 0.01* |
| semi-fossorial | 90.73 | 0.563 | 525.32 | 0.850 |
| **Diet** | | | | |
| Hard eggs | 102.42 | 0.640 | 532.97 | 0.604 |
| Hard prey | 89.35 | 0.443 | 491.36 | 0.290 |
| Invertebrates | 87.85 | 0.304 | 508.23 | 0.621 |
| Mollusks | 53.65 | 0.165 | 448.20 | 0.372 |
| Soft prey | 88.04 | 0.155 | 501.86 | 0.616 |

654 Table 9: Results of the morphological disparity analysis across ecological niches and diets.

655 The table presents Procrustes variances for each niche and diet, indicating the level of

656 morphological variation within groups.

| Comparison | Procrustes variance |
|---|---|
| Habitat type | |
| Aquatic | 0.011605613 |
| Arboreal | 0.008724903 |
| Fossorial | 0.010839093 |
| Semi-aquatic | 0.010766939 |
| Semi-arboreal | 0.011107773 |
| Semi-fossorial | 0.012223323 |
| Terrestrial | 0.011249285 |
| Diet | |
| Generalist | 0.007111103 |
| Hard eggs | 0.008578330 |
| Hard prey | 0.010988433 |
| Invertebrates | 0.008819887 |
| Mollusks | 0.008497849 |
| Soft eggs | 0.002473988 |
| Soft prey | 0.017594851 |

Phylogenetic PCA with Dietary Proportions

Phylogenetic Two-Block PLS: Morphology vs Niche

**Phylogenetic Two-Block PLS: Morphology vs Diet**