Title

Transcriptional Repression of *reaper* by Stand Still Safeguards Female

Germline Development in *Drosophila*

Authors

Masaya Matsui[1], Shinichi Kawaguchi[1], and Toshie Kai[1]*

*Corresponding author

E-mail: kai.toshie.fbs@osaka-u.ac.jp (KT)

Affiliations

[1] Graduate School of Frontier Biosciences, The University of Osaka, Osaka

565-0871, Japan

Abstract

Apoptosis plays a central role in shaping tissues and preserving cellular integrity across developmental stages. In the germline, its precise regulation is critical to ensure both the elimination of aberrant cells and the maintenance of reproductive capacity. However, the molecular mechanisms that control apoptotic susceptibility in germline cells remain poorly defined. Here, we identify *stand still* (*stil*) as a female germline-specific regulator of apoptosis in *Drosophila*. Loss of *stil* leads to near-complete depletion of germline cells at the time of eclosion, associated with upregulation of the pro-apoptotic gene *reaper* (*rpr*) and activation of caspase-dependent cell death. Reporter assays in S2 cells show that Stil directly represses *rpr* transcription through its N-terminal BED-type zinc finger domain. Despite the absence of *stil*, undifferentiated germline cells remain resistant to apoptosis. Analysis of publicly available chromatin data reveals that the *rpr* locus in these cells resides in a closed, H3K9me3-enriched chromatin state, suggesting a Stil-independent mode of transcriptional silencing. Together, our findings uncover a dual mechanism that protects the female germline from *rpr*-dependent apoptosis: active repression by Stil in differentiating cells and chromatin-mediated silencing in undifferentiated cells. This work provides new insights into the transcriptional and chromatin-based mechanisms that maintain germline cell identity and survival.

Introduction

Apoptosis, a form of programmed cell death (PCD), is a highly regulated biological process essential for animal development and tissue homeostasis. *Drosophila* is an excellent model organism for studying apoptotic cell death, due to its powerful genetic and molecular tools. In Drosophila, apoptosis is often initiated by the expression of IAP (inhibitor of apoptosis) antagonists, including members of the RHG gene family: *reaper* (*rpr*), *hid* (*head involution defective*), *grim*, and *sickle* (*skl*) (Abbott & Lengyel, 1991; P. Chen et al., 1996; Christich et al., 2002; White et al., 1994). Among these, the pro-apoptotic gene *rpr* plays a central role in PCD and stress-induced apoptosis (Yan et al., 2004). The *Drosophila* IAP protein DIAP1 is degraded via auto-ubiquitination, triggered by the physical interaction of Rpr through its conserved IAP-binding motif (IBM), leading to caspase activation and apoptotic cell death (Holley et al., 2002; Sandu et al., 2010). *rpr* expression is regulated by transcription factors and histone modifications, such as H3K27 and H3K9 trimethylation, in response to developmental and environmental signals (Beira et al., 2014; Brodsky et al., 2000; Jiang et al., 2000; Lohmann et al., 2002; Zhang et al., 2008).

For reproductive success, apoptosis is meticulously regulated to control cell numbers and maintain germline cell quality (Almeida et al., 2013; Chaudhry et al., 2021; Huang et al., 2023; Knudson et al., 1995). *Drosophila* adult female has a pair of ovaries, each containing 16–20 ovarioles. At the apical end of each ovariole lies the germarium, which houses germline stem cells (GSCs) that divide asymmetrically to produce a self-renewing GSC and a cystoblast. Cystoblast undergoes four mitotic divisions to form a 16-cell cyst

interconnected by ring canals. Of these 16 cells, one differentiates into the

oocyte and enters meiosis, while the remaining 15 become nurse cells that

support the oocyte by transporting proteins and RNAs. Encapsulated by

somatic follicle cells, the cysts develop into egg chambers that progress

through 14 stages, culminating in mature, fertilizable eggs (Gleason et al.,

2018; Spradling et al., 2001).

To safeguard the fidelity of this process, apoptosis operates at two

key checkpoints —within the germarium and during mid-oogenesis —ensuring

that only healthy cells contribute to egg development. In the germarium, the

first checkpoint triggers apoptotic cell death in response to DNA damage,

mediated by p53-induced *hid* expression rather than *rpr* (Hou et al., 2008;

Park et al., 2019). At the second checkpoint during mid-oogenesis, germline

cells undergo caspase-regulated apoptotic and autophagic cell death in

nutrient deprived condition, even though RHG genes are dispensable.

(Baonza et al., 2022; Jeanne S. Peterson et al., 2007; T L Pritchett & McCall,

2012; Xing et al., 2015). Although apoptotic cell death is intricately regulated

in response to various stimuli during oogenesis, the involvement of *rpr* in

germline apoptosis remains largely unknown. Previous studies have identified

stand still (*stil*) as a critical regulator of *Drosophila* germline cell survival

(Schüpbach & Wieschaus, 1989). Genetic interactions with female germline-

specific genes, such as *ovo* and *ovarian tumor* (*otu*), along with the female

germline-specific requirement of *stil*, suggested its involvement in sex

determination (G & D., 1997; Sahut-Barnola & Pauli, 1999). However, how Stil

controls germline cell fate remains unknown. In this study, we identify *stil* as a

repressor of *rpr* in female germline cells. Loss of *stil* function led to apoptosis-

mediated depletion of ovarian germline cells, associated with *rpr* upregulation.

Stil was enriched at the *rpr* locus and its upstream transcriptional control

region, suggesting a direct role in *rpr* regulation. Reporter assays in

Drosophila S2 cells revealed that Stil represses *rpr* transcription via its N-

terminal BED-type zinc finger domain, indicating its role as a repressive

transcription factor. In addition, we found that undifferentiated germline stem

cell (GSC)-like cells, in which the *rpr* locus exhibits a closed chromatin

structure, were resistant to apoptosis caused by *stil* loss. Our findings reveal a

molecular mechanism by which Stil and chromatin accessibility repress *rpr*,

providing novel insights into the regulation of germline cell fate.

Results

Loss of *stand still* (*stil*) leads to female germline cell loss

Previous studies have demonstrated the necessity of *standstill* (*stil*) function in female germline cells; however, its molecular role remains elusive (G & D., 1997; PK et al., 1996). To address this, we analyzed a loss of function allele, *stil^{EY16156}*, which harbors a P-element insertion near the 3'end of the second exon of *stil* gene, potentially disrupting the splicing of *stil* transcripts (Fig 1A) (Bellen et al., 2004, 2011). RT-PCR revealed that *stil* transcripts were undetectable in homozygous *stil^{EY16156}* ovaries and in *stil^{EY16156}* testes, indicating a complete loss of *stil* expression in both gonads (Fig. 1B).

Similar to the previously reported loss-of-function allele *stil^3*, which exhibited the most severe phenotype (Fig. 1A; Fig. S1A, B) (PK et al., 1996), ovaries of both *stil^{EY16156}* homozygous and *stil^{EY16156}/Df(2R)vg-C* trans-heterozygote, in which the deficiency uncovers the entire *stil* locus, were severely reduced in size and completely lacked germline cells (Fig. 1C, D). Likewise, *stil^3/stil^{EY16156}* trans-heterozygote females exhibited more severe germline cell loss than *stil^3/Df(2R)vg-C* (Fig. S1A, B). However, male germline cell in *stil^{EY16156}* testes progressed to differentiate into mature sperms (Fig.S1A). Notably, fertility assayed by egg production and hatching rate, was comparable between *stil^{EY16156}* and *stil^{EY16156}/CyO* males (Fig. S1B), indicating that *stil* is specifically required for female germline cells. The loss of germline cell in *stil^{EY16156}* ovaries was fully rescued by overexpression of the *UASp-6×Myc-Stil* transgene, driven by the germline-specific *NGT40; Nos-Gal4-VP16* driver (Fig. 1C, D) (Grieder et al., 2000). Mutant females expressing 6×Myc-Stil,

germline cells were differentiated and developed into mature eggs that

hatched comparably to those of heterozygous control (Fig. 1E). Collectively,

these results confirm the pivotal role of *stil* in maintaining female germline

cells and ensuring proper oogenesis.

stil-deficient germline cells undergo apoptotic cell death

To investigate the cause of female germline cell loss, we knocked down *stil*

function by expressing short hairpin RNA targeting the 4th exon of the *stil*

transcript (GL01230) using a germline driver (Fig. 1A). Germline-specific

knockdown (*stil*-GLKD) reduced *stil* expression in ovaries by approximately

90% compared to control (Fig. 2A). *stil*-GLKD females contained significantly

smaller ovaries and did not lay any eggs, resembling to *stil* loss-of-function

mutant females (Fig. 2B and Fig. S2A). Unlike the complete germline loss

observed in *stil* mutants, however, most ovarioles in *stil*-GLKD retained

germline cells, indicating a milder phenotype (Fig. 2C). Notably, the number of

ovarioles containing at least two egg chambers was remarkably fewer (27.4%

versus 100% in the control), suggesting that oogenesis was perturbed before

mid-oogenesis (Fig. 2B, D). While *stil*-GLKD exhibited severe defects similar

to its loss-of function alleles, its somatic knockdown using a tj-Gal4 driver (*stil*-

STKD) did not affect egg production and the hatching rate at all (Fig. S2A).

These observations suggest that *stil* functions in germline cells both for the

survival of female germline cells and egg development.

Previous studies have suggested that stil may play a role in sex

determination, particularly through its genetic interaction with *ovo* and the key

sex determination regulator *otu*, which represses the male identity factor

Tdrd5l (G & D., 1997; Primus et al., 2019; Sahut-Barnola & Pauli, 1999) . To

investigate whether *stil* is involved in sex determination, we examined the

transcripts of *sxl*, a master regulator of sex determination of germline cells in

stil mutant ovaries (Hashiyama et al., 2011). *sxl* pre-mRNA undergoes

alternative splicing to generate sex-specific isoforms, producing a functional

Sxl protein essential for female development, or a truncated non-functional

male-specific isoform (Bell et al., 1991) (Moschall et al., 2019). RT-PCR

revealed an enrichment of the female-specific *sxl* isoform in both control and

stil-GLKD ovaries (Fig. S2B). In addition, male-specific *sxl* transcripts,

detected in both control and *stil*-GLKD testes, were absent in *stil*-GLKD

ovaries, indicating that *sxl* splicing patterns remain unaffected by *stil*

depletion. These results suggest that *stil* functions independently of sex

identity determination in germline cells.

Damaged or abnormal cells are typically removed through regulated cell

death mechanisms, including apoptosis, autophagy, and necrosis (Fuchs &

Steller, 2011, 2015). In *Drosophila* ovaries, such cell death occurs at two

distinct developmental checkpoints during oogenesis: in the germarium and

during mid-oogenesis, in response to either developmental cues or

environmental stressors (McCall, 2004; Tracy L. Pritchett et al., 2009). To

determine whether germline cell loss in *stil*-deficient ovaries was due to

apoptosis, we performed a TUNEL (Terminal deoxynucleotidyl Transferase

Biotin dUTP Nick End Labeling) assay, which detects DNA fragmentation

indicative of apoptotic cell death. In region 2A and 2B of the germarium, the

first developmental checkpoint in oogenesis (Drummond-Barbosa &

Spradling, 2001), the number of TUNEL-positive cells significantly increased

in *stil*-GLKD ovaries compared to controls (Fig. 2E, F). These results suggest

that *stil*-GLKD germline cells undergoes apoptotic cell death.

To further confirm the involvement of *stil* in apoptosis, we overexpressed

inhibitor of apoptosis proteins (IAPs) in *stil* mutant germline cells. Both the

Drosophila protein DIAP1 (Death-associated inhibitor of apoptosis 1) and the

baculovirus-derived P35 are well-characterized inhibitors that can effectively

suppress apoptosis in *Drosophila*. DIAP1 inhibits the activity of initiator

caspase, Dronc (Death regulator Nedd2-like caspase) which activates effector

caspase, while P35 directly inhibits effector caspase (Baum et al., 2007; J.S

Peterson et al., 2003; Werz et al., 2005). Germline-specific overexpression of

P35 fully rescued the germline cell loss observed in *stil*^{EY16156} mutants (Fig.

2G, H). However, germline development remained arrested, and egg

chambers degenerated at the mid-oogenesis (Fig. S3). Likewise, DIAP1

overexpression partially restored germline cells (~50% of ovarioles retained

germline cells), but did not fully rescue oogenesis defects (Fig. 2G, H).

Altogether, these findings indicate that the loss of *stil* function not only triggers

apoptosis that can be suppressed by apoptosis inhibitors but also causes

defects in oogenesis progression that are not rescued by blocking cell death.

Stil regulates proapoptotic genes to suppress apoptosis in female

germline cells

Stil protein harbors a characteristic DNA-binding domain, BED-type zinc

finger, containing highly conserved aromatic residues followed by a shared

pattern of cysteines and histidines (Fig. S4A) (Aravind, 2000). Given that

many of the known BED-type zinc fingers including the founder members,

BEAF and DREF, function as transcription factors or chromatin regulators (McKowen et al., 2022; Sokolova et al., 2023; Tue et al., 2017), Stil may play a role in transcriptional regulation. To examine this possibility, we analyzed the transcriptional profiles of ovarian RNA of *stil* mutant and control. We took advantage of the ectopic expression of P35 to overcome cell death defect of germline cells due to the loss of in *stil* function. The RNA-seq analysis identified 3,569 (30.7%) differentially expressed genes (DEGs) (log2FC>|1| and p-adj<0.01) out of 11,608 annotated genes (PRJDB20709; Supplementary Table 1). Among these, 2,044 genes were up-regulated (17.6 %, red dots) and 1,525 genes were down-regulated (13.1 %, blue dots) in *stil* mutant ovaries (Fig. S4B). Consistent with the observation of developmental arrest and egg chambers degeneration at the mid-stage of oogenesis (Fig. S3), genes known to be highly expressed in late-stage oogenesis, such as proteins composing the chorion, were significantly downregulated in *stil* mutant ovaries (Fig. 3A). Of those upregulated, notably, the proapoptotic gene *rpr* (*reaper*) ranked among the top 10 upregulated DEGs (Fig. 3A). Given the apoptotic phenotype of *stil* mutant germline cells, 67 apoptosis-related genes out of 224 annotated in the dataset (GO:0006915) were identified as DEGs (Fig. 3B). Among the top 10 genes list of apoptosis-related DEGs, proapoptotic genes, *rpr* and *hid* (*head involution defective*), were significantly upregulated, while the hedgehog signaling genes expressed in later oogenesis such as *ptc*, *duox*, and *rdx* were significantly downregulated (Fig. 3C). *rpr* and *hid* are the key members of IAP antagonists, known for their central role in initiating apoptosis (Abbott & Lengyel, 1991; White et al., 1994). Their activation through various signaling pathways leads to the ubiquitin-

mediated degradation and translational inhibition of DIAP1, which

subsequently results in the activation of Caspase proteins (Freel et al., 2008;

Holley et al., 2002). Further validation by RT-qPCR demonstrated that *rpr*

expression increased approximately 700-fold, while *hid* expression increased

15-fold in the rescued *stil* mutant ovaries by P35 expression (Fig. 3D). To

exclude a possibility that egg chamber degeneration in germline-rescued *stil*

mutants affects proapoptotic gene expression, we isolated RNAs from early-

stage ovaries (up to stage 5-6 of oogenesis) by manual removal of later

stages and re-examined their expression levels. *rpr* expression remained

elevated in RNAs prepared from early-stage ovaries, while *hid* expression did

not (Fig. S4B). Furthermore, RT-qPCR analysis using *stil*-GLKD ovaries

driven by three different germline drivers, *NGT40*; *NosGal4-VP16*, *Bam-Gal4*,

and *matα-Tub-Gal4*, consistently exhibited significant upregulation of *rpr* but

not *hid* across all conditions (Fig. S5A). Conversely, *rpr* was not prominently

upregulated in *stil*-GLKD testes with *NGT40*; *NosGal4-VP16* or *Bam-Gal4*,

consistent with the phenotypic observations in *stil* mutant testes (Fig. S5C).

These results suggest that *rpr* is robustly upregulated in the absence of *stil*

function in female germline cells.

To determine whether proapoptotic genes upregulation directly contributes to

apoptosis of *stil* mutant female germline cells, we generated double mutants

of *stil* and the proapoptotic gene, *rpr*. Strikingly, the germline cell death and

differentiation defects observed in *stil* mutant ovaries were fully rescued in the

double mutant ovaries (Fig. 3E, F). Germline development proceeded,

resulting in the formation of mature eggs and partial restoration of fertility;

mutant females laid a few eggs and approximately 40 % of those hatched

(Fig. 3G, H). In contrast, *stil-hid* double mutants showed a slight increase in

germline cell survival, but no developing egg chambers were observed (Fig.

S5D, E), indicating that *hid* does not play a central role in apoptosis caused by

the loss of *stil* function. These findings highlight that *rpr* regulation by *stil* is

critical for suppressing apoptosis in female germline cells, establishing *rpr* as

a key downstream effector in this pathway.

Stil represses *rpr* expression by targeting the regulatory regions via its

BED-type zinc finger

The transcription of *rpr* is governed by a complex network of

developmental, environmental, and stress-related signals, each employing

distinct regulatory mechanisms. These signals modulate transcription factors

such as p53, Dfd, the EcR/USP complex, AP-1, Shn, leading to either the

activation or suppression of apoptosis (Beira et al., 2014; Brodsky et al., 2000;

Jiang et al., 2000; Lohmann et al., 2002). These transcription factors interact

with upstream enhancer and promoter regions of the *rpr* locus (*rpr-*

transcriptional control region) to either activate or suppress its transcription,

ensuring precise spatial and temporal control of apoptosis (Fuchs & Steller,

2011). To examine whether Stil also directly interacts with *rpr*-transcriptional

control region, we performed DNA adenine methyltransferase identification

(DamID)-seq analysis (Aughey et al., 2019). A fusion of Stil with *E. coli* DNA

adenine methyltransferase (Dam-Stil) or Dam alone as a control were

expressed specifically in germline cells (Díaz-Torres et al., 2021; Southall et

al., 2013). Genomic DNA from ovaries was digested with DpnI and subjected

to NGS to detect GATC sites methylated by Dam-Stil near Stil-bound regions

(PRJDB20722). Genome-wide DamID-seq analysis revealed that ~80% of

Stil-enriched peaks were located in promoter regions, supporting a role for Stil

in transcriptional regulation (Fig. 4A). At higher resolution, prominent peaks

were observed at the *rpr* locus and its upstream region (5–6 kb), suggesting

direct binding of Stil to the *rpr* regulatory element (Fig. 4B).Next, we examined

the role of Stil's BED-type zinc finger in *rpr* repression by generating four Stil

variants: Stil-FL (full-length), Stil-NT (N-terminal domain containing the BED-

type zinc finger and nuclear localization signal), Stil-CT (C-terminal domain

lacking these two motifs), and Stil-AAYA (full-length Stil with a mutated BED-

type zinc finger substituting four residues including two critical cysteines) (Fig.

4C). In nurse cells, despite lacking the predicted nuclear localization signal,

Stil-CT localized to the nucleus and overlapped with chromosomes, as did

Stil-FL and Stil-AAYA, whereas Stil-NT showed diffuse nuclear distribution

without being enriched onto chromosomes (Fig. S6A). Collectively, these data

suggest that Stil's nuclear and chromatin localization are mediated by the C-

terminal domain, independently of the zinc finger motif. To assess the

functional significance of these domains *in vivo*, we expressed the variants in

stil^{EY16156} germline cells. Stil-FL fully rescued the defects of both germline cell

survival and development, leading to the production of mature eggs (Fig. 4D,

E, F). Remarkably, Stil-NT partially rescued germline defects despite its

diffuse nuclear localization, while neither Stil-CT nor Stil-AAYA, both lacking

functional BED-type zinc fingers, restored germline cell survival or

differentiation. These findings suggest that the BED-type zinc finger is

essential for Stil function in germline cells possibly as a transcription regulator,

while the C-terminal domain mediates nuclear and chromatin localization.

To validate whether the BED-type zinc finger functions in *rpr* repression

through interaction with the *rpr* transcriptional control region, we performed a

reporter assay using *Drosophila* S2 cells. We constructed a *rpr*-GFP reporter

containing the 5-kb upstream region of *rpr*, where DamID-seq signals were

detected, fused to GFP (Fig. 4G) and transfected with Stil variants. Stil

variants exhibited localization patterns in S2 cells similar to those in nurse

cells (Fig. S6B), supporting their suitability for functional analysis in S2 cells.

Consistent with the *in vivo* rescue experiment (Fig.4 D-F), co-expression of

Stil-FL with the *rpr*-GFP reporter significantly repressed GFP expression. Stil-

NT containing the BED-type zinc finger also reduced GFP expression, albeit

partially. In contrast, Stil-CT and Stil-AAYA failed to exhibit any detectable

repressive effects (Fig. 4H). To confirm the specificity of this Stil-mediated

repression, we utilized a *tubP*-GFP reporter containing the *αTub84B* gene

promoter fused to GFP (Fig. S6C)(Brennecke et al., 2003). Co-expression of

Stil-FL with the *tubP*-GFP reporter led to a modest reduction in GFP

expression, while other variants showed no discernible effect (Fig. S6D). This

effect was markedly weaker than the repression observed with the *rpr*-GFP

reporter, suggesting that Stil preferentially targets the *rpr* upstream region.

These results demonstrate that Stil interacts with the *rpr* transcriptional control

region via its BED-type zinc finger, which is critical for transcriptional

repression, while the chromatin association through the C-terminal region is

required for full repressive activity in germline cells.

Undifferentiated Germline Cells Exhibit Resistance to Apoptosis in *stil*

loss

Loss of *stil* function resulted in severe depletion of germline cells by the 3rd

instar larval stage (120 hours after egg laying, AEL), leaving only a few

surviving germline cells in *stil* mutant (Fig. 5A, B). Notably, these surviving

germline cells were predominantly found near the terminal filaments and

future cap cells, the niche that supports germline stem cell (GSC) (Fig.

5A)(Spradling et al., 2001). This observation suggested that GSCs within the

niche may be resistant to cell death caused by *stil* loss. To test whether GSCs

are inherently resistant to cell death caused by *stil* deficiency, we induced

GSC-like properties by overexpressing a constitutively active the

Decapentaplegic (Dpp) receptor, Thickveins (Tkv.CA) in germline cells, or the

loss of function of *bag of marbles* (*bam*), a key differentiation factor

(Casanueva & Ferguson, 2004; D. Chen & McKearin, 2003a). These

manipulations resulted in the tumorigenic proliferation of undifferentiated

GSC-like cells or pre-cystoblast, respectively (Tiwari et al., 2019).

Remarkably, Tkv.CA overexpression and *bam* loss rescued germline cell

depletion in *stil* mutant ovaries, leading to the formation of tumorous germaria

containing undifferentiated germline cells comparable to the *stil* heterozygous

control (Fig. 5C, D). Indeed, TUNEL assay showed no increase in apoptotic

cells in such tumorous germaria (Fig. 5E, F), indicating that the

undifferentiated state of germline cells confers resistance to apoptosis in the

absence of *stil*.

Consistent with this observation, RNA-seq and gene Ontology (GO)

analysis of those tumorous ovaries revealed that the expression levels of

most apoptosis-related genes (GO:0006915), including *rpr*, remain unaffected

(PRJDB20709; Supplementary Table 2) (Fig. 5G). Furthermore, RT-qPCR

confirmed that *rpr* expression was not significantly upregulated in the GSC-

like tumor *stil* mutant ovaries compared to p35-rescued *stil* mutant ones (Fig.

3D, 5H), suggesting that apoptosis-related genes, including *rpr*, are

transcriptionally repressed in GSC-like cells even in the absence of *stil*

function.

** *rpr* Silencing in Undifferentiated Germline Cells Is Associated with**

Chromatin States

Previous studies in mammals have established that stem cells are

generally resistant to apoptosis. In mouse mammary stem cells and cancer

stem cells, this resistance is attributed to several mechanisms, including

enhanced DNA repair and overexpression of anti-apoptotic proteins (Chang et

al., 2015; Nathansen et al., 2021; Safa, 2016). Similarly, in *Drosophila*,

intestinal stem cells were shown to resist cell death through a two-tiered

regulation of *rpr*, involving chromatin state and spatial control of its expression

(Sulekh et al., 2024). To explore whether chromatin dynamics contribute to

GSC apoptosis resistance in *stil* mutants, we analyzed the chromatin status at

the *rpr* locus using Assay for Transposase-Accessible Chromatin-seq (ATAC-

seq) data of Drosophila ovarian cells (Pang et al., 2023). This dataset was

generated using fluorescence-activated cell sorting (FACS) of GSCs (2C–4C)

and nurse cells (NCs) at various stages of egg chamber maturation, with

ploidy levels ranging from 4C to 512C due to endoreplication (Dej &

Spradling, 1999). In early mid-stage NCs where *stil* deficient-germline cells

undergo apoptosis, ATAC-seq signals were enriched at three upstream

regions of *rpr* (-0.5 to 0.5 kb, 1.5 to 2.0 kb, and 3.5 to 5.0 kb relative to the

transcription start site [TSS] of *rpr*) (Fig. 6A). Conversely, signals at these

three regions were significantly lower in 2C-4C GSCs and 64C-512C NCs.

These results suggest that the chromatin state upstream of *rpr* is closed in

GSCs and late-stage NCs (64C-512C) but open in early mid-stage NCs (4C-

32C). Notably, despite harboring the same ploidy, 4C GSCs and 4C NCs

displayed distinct chromatin accessibility, reflecting regulatory differences

likely responsible for their differential responses to apoptosis (Fig. 6A). We

further analyzed epigenetic marks such as H3K9me3 and H3K27me3 at the

rpr locus using publicly available data sets (DeLuca et al., 2020; Pang et al.,

2023). H3K9me3, a histone modification associated with transcriptional

repression, was observed around the *rpr* locus and its transcriptional control

region in GSCs, whereas only minimal signals were detected in 4C NCs (Fig.

6B). In contrast, H3K27me3, another repression mark exhibited no significant

difference between GSC and st5 NC (Fig. 6C) (DeLuca et al., 2020). This

suggests that H3K9me3, rather than H3K27me3, may establish a

heterochromatic state at the *rpr* loci in GSCs, contributing to transcriptional

silencing. These findings imply that chromatin dynamics play a crucial role in

regulating *rpr* expression during germline development.

Discussion

In this study, we investigated the role of the Drosophila gene *stand still* (*stil*), which is required for the survival and maturation of female germline cells. Loss of *stil* leads to apoptosis of germline cells via upregulation of the pro-apoptotic gene *rpr* (Fig. 3E), identifying *rpr* as the primary effector of cell death in this context. GFP-reporter assays with Stil variants in S2 cells suggest that Stil represses *rpr* expression, likely through direct binding to its upstream regulatory region via the BED-type zinc finger domain (Fig. 4H).

The transcription of *rpr* is regulated by multiple transcription factors that bind to cis-regulatory elements located upstream of the gene (Fuchs & Steller, 2011). For example, the tumor suppressor protein p53 binds to a well-characterized p53 response element (p53RE) located 4.8 kb upstream of *rpr*, which is responsive to intrinsic and extrinsic signals during embryogenesis and in germline cells (Lu et al., 2010; Wylie et al., 2014). However, p53 binding to this element has only been observed in the anterior germarium, and no direct link between p53 and apoptosis has been established in the female germline (Park et al., 2019). Other known regulators, such as the homeobox transcription factor Deformed (Dfd) and the JNK-responsive complex AP-1, also bind upstream *rpr* enhancers in somatic tissues, but neither is active in the female germline (Beira et al., 2014; Lohmann et al., 2002). Thus, factors that suppress *rpr* transcription specifically in germline cells have remained unidentified. Our study identifies Stil as a female germline-specific repressor

of *rpr*. DamID profiling revealed Stil binding to the *rpr* locus to 5–6 kb

upstream, covering regulatory elements associated with known transcription

factors (Fig. 4B). Consistently, Stil repressed *rpr*-GFP reporter containing the

5 kb upstream regulatory region in S2 cells, and this repression required Stil's

BED-type zinc finger domain (Fig. 4H). This domain is known to mediate DNA

binding and contribute to transcriptional regulation and chromatin remodeling

(Aravind, 2000). These findings support a model in which Stil directly occupies

the *rpr* regulatory region and blocks activator access to maintain

transcriptional repression in germline cells.

Whereas DIAP1 overexpression only partially rescued germline loss

in *stil* mutants, P35 restored germline survival in nearly all cases (Fig. 2G, H),

highlighting a functional distinction in how these inhibitors act in the context of

rpr-induced apoptosis. Both inhibit apoptosis through distinct mechanisms:

DIAP1 blocks caspase activation by inhibiting initiator and effector caspases,

while P35, a baculoviral protein, acts as a pseudo-substrate that irreversibly

inhibits effector caspases such as DrICE and Dcp-1 (Li et al., 2024; Nair &

Baker, 2024). *rpr* antagonizes DIAP1 through two independent mechanisms—

by promoting its auto-ubiquitination and proteasomal degradation, and by

inhibiting its translation (Freel et al., 2008; Holley et al., 2002). Therefore, high

rpr levels in *stil* mutant ovaries may deplete functional DIAP1, limiting the

rescue effect of its overexpression. By contrast, P35 directly targets

downstream caspases, bypassing DIAP1 loss and producing stronger

inhibition. These results support the model that Stil maintains the apoptotic

threshold by regulating *rpr* levels and balancing Rpr–DIAP1 interactions

(Sandu et al., 2010; Yin & Thummel, 2004).

Despite the loss of *stil*, undifferentiated female germline cells (GSCs

and pre-cyst cells) remain resistant to apoptosis (Fig. 5E, F), correlating with

closed chromatin and H3K9me3 enrichment at the *rpr* locus (Fig. 6A, B).

Similar chromatin-based repression of *rpr* has been reported in *Drosophila*

intestinal stem cells (Sulekh et al., 2024) and in late-stage embryos, where

heterochromatin silences *rpr* expression (Zhang et al., 2008). Together with

our findings, these studies suggest that chromatin-mediated silencing

provides an additional layer of protection against apoptosis in female GSCs,

independent of *stil*. By contrast, *stil* mutant male germline cells show no

phenotype and no *rpr* upregulation (Fig. S5B), suggesting that that rpr-

mediated apoptosis is not activated in the male germline. This points to a sex-

specific mechanism of *rpr* repression, potentially involving chromatin-based

silencing, although this possibility has yet to be examined.

Together, our findings reveal two mechanisms that protect female

germline cells from *rpr*-dependent apoptosis: Stil-mediated transcriptional

repression in differentiating germline cells and chromatin-based silencing in

undifferentiated germline cells. Although the upstream regulation of *stil* is

unknown, its selective role in female germline cells suggests it functions as a

developmental checkpoint that prevents premature apoptosis during

oogenesis. A key question is how *stil* integrates intrinsic or environmental

cues to modulate the apoptotic threshold during germline development.

Materials and Methods

Fly stock and cultures

All *Drosophila* stocks and crosses were reared at room temperature (25ºC) on standard food [5% (w/v) dry yeast, 5% (w/v) corn flower, 2% (w/v) rice bran, 10% (w/v) glucose, 0.7% (w/v) agar, 0.2% (v/v) propionic acid and 0.05% (w/v) p-hydroxy butyl benzoic acid]. The following stocks were used in this study: *yellow white* (*y w*), *attP2* (BL#25710), *stil^EY16156* (BL#21191), *stil^3* (BL#533, splicing donor site of the second intron was mutated from GTATG to ATATG at 12570993), *Df(2R)vg-C* (BL#754, deficiency for *stil*), *Df(3L)X14* (BL#600368, deficiency for *hid*) *UASp-P35* (a gift from Dr. Bergmann), *UASp-DIAP1* (BL#63818), *rpr^87* (BL#83150), *hid^05014* (BL#83349), *UASp-Tkv.CA* (a gift from Dr. Dahua Chen), *bam^Δ86* (BL#5427), *NGT40-Gal4*; *nosGal4-VP16* (Grieder et al., 2000), *mat α-tub-Gal4* (BL#7063), *bam-Gal4* (D. Chen & McKearin, 2003b) , and *tj-Gal4* (DGRC#104055). The mutation site of *stil^3* was identified by sanger sequencing of the PCR-amplified fragment.

Generation of transgenic fly lines

Transgenic fly lines expressing Stil variants (6×Myc-Stil-FL, 6×Myc-Stil-NT, 6×Myc-Stil-CT, and 6×Myc-Stil-AAYA) and DamID constructs (Dam-alone and Dam-Stil) were generated using the PhiC31 integrase-mediated transgenesis system. DNA fragments encoding Stil variant were amplified using cDNAs obtained by reverse transcription of ovarian RNA extracted from *y w* flies.

DamID constructs were designed such that a primary open reading frame

(ORF) encoding GFP was followed by two stop codons and a secondary ORF

encoding either Dam or Dam-Stil (Southall et al., 2013). The DNA fragment

encoding Dam was amplified from the pUASTattB-LT3-NDam vector (a gift

from Dr. Andrea H. Brand). The resulting DNA fragments were cloned into the

pUASp-K10-attB plasmid backbone (Koch et al., 2009). These transgenic

constructs were injected into embryos of attP-containing strains (Stil variants

into P40, BDSC #25709, Dam and Dam-Stil into P2, BDSC #25710,

respectively), and transformants expressing *mini-white* were selected.

Immunofluorescence staining

Immunofluorescence staining was performed as previously described (Xu et

al., 2024). Ovaries and testis from adults and the 3rd instar larva were

dissected in cold phosphate buffer saline (PBS) and fixed with PBS containing

5.3% (w/v) paraformaldehyde for 10 min. Fixed samples were washed in PBX

[PBS with 0.2% (v/v) Triton-X (Nacalai)] for at least 30 min with several

change of solution, then blocked with PBX containing 4% (w/v) bovine serum

albumin (BSA) for at least 30 min at room temperature. After rinsing once with

PBX, the samples were incubated with the primary antibodies diluted in PBX

containing 0.4% (w/v) BSA for more than 4 h at room temperature or

overnight at 4°C. After several washes with PBX for at least 1 h, the samples

were incubated in secondary antibodies diluted in PBX containing 0.4% (w/v)

BSA for more than 2 h at room temperature. The samples were washed

several times with PBX for at least 1 h and stained with 1 µg/ml of 4',6-

diamidino-2phenylindole (DAPI) (Sigma) diluted in PBX for 10 min at room

temperature. Following two rinses with PBX and PBS, tissues were mounted

in Fluoro-KEEPER Antifade Reagent (Nacalai). The following primary

antibodies were used at the indicated dilutions: guinea pig anti-Vas (1:1000)

(Patil & Kai, 2010), mouse anti c-Myc [Wako (9E10), 1:500], and rabbit anti-

FLAG [MBL, PM020, 1:1000]. Secondary antibodies were Alexa Fluor 488-

and 555-conjugated goat antibodies against rabbit, mouse and guinea pig IgG

(Invitrogen, A11029, A11034, A11073, A21424, A21429 and A21435) at a

1:200 dilution in PBX containing 0.4% (w/v) BSA. Images were captured by

Zeiss LSM780 Confocal Microscope with 10x, 20x, and 40x lens and

processed with Zen (Zeiss) and Fiji application (Schindelin et al., 2012) .

TUNEL assay

TUNEL assay was conducted using ApopTag® Kits (Millipore), labelling the

free 3'-OH DNA termini in situ with fluorescein-nucleotide by terminal

deoxynucleotidyl transferase (TdT). Dissected ovaries in cold PBS were fixed

with PBS containing 5.3% (w/v) paraformaldehyde and rinsed 3 times with

PBX. Ovaries were washed with PBX for 10 min twice and further washed

with PBST [PBS with 0.1% (v/v) Tween 20 (Nacalai)] twice. After rinsing with

Equilibration Buffer twice, the samples were incubated with TdT reagent [70%

(v/v) Reaction Buffer and 30% (v/v) TdT Enzyme] for 1 h at 37ºC. After

incubation with 1/35 diluted Stop/Wash Buffer for 15 min, samples were

rinsed 3 times with PBX and then washed 3 times with PBX for 10 min. The

tissue mounting and microscopic observation were as described in the

"Immunofluorescence staining" section.

RT-PCR and quantitative RT-PCR

Total RNA was isolated from ovaries using TRIzol LS Reagent (Invitrogen),

following the manufacturer's protocol. RNA samples (1 μg each) were treated

with DNase I (NEB) to digest genomic DNA, followed by inactivation of the

enzyme with 5 mM EDTA at 75ºC for 10 min. Reverse transcription was

performed using the DNase I-treated RNA with oligo(dT)20 primers and

SuperScript III Reverse Transcriptase (Invitrogen), following the

manufacturer's protocol. cDNAs were subjected to 30 cycles of RT-PCR with

GoTaq DNA polymerase (Promega).

Quantitative PCR (qPCR) was performed with Fast SYBR Green regents

(Invitrogen) using at least 3 biological replicates. Deltal-delta-CT method was

used to obtain the relative transcript levels. *α-Tub84B (tubulin)* was used as

the reference. The sequences of the primers are listed in Supplementary

Table 3.

mRNA-seq analysis

Total RNAs were extracted from 100 ovaries of control and *stil* mutant of

Tkv.CA OE background and 40 ovaries of control and *stil* mutant of *P35 OE*

background using TRIzol (ThermoFisher) following the manufacture's

instruction. Extracted RNAs were treated with DNase I for 10 min at 37ºC and

reactions were inhibited by 5 mM EDTA at 70ºC for 10 min, following that

RNAs were purified using phenol/ chloroform/ isoamyl alcohol (mixed with a

ratio of 25: 24: 1, pH 5.2). Purified RNAs were prepared for mRNA-seq

libraries with NEBNext Ultra II RNA Library Prep Kit for Illumina (NEB), and

deep-sequencing was conducted using NovaSeq6000 (Illumina). Sequencing

analysis was conducted using bioinformatic software, CLC workbench 12

(QIAGEN). Adaptor sequences of total reads obtained from mRNA-seq were

trimmed and reads were mapped to *Drosophila* genomes, r6.32 from Flybase

(ftp://ftp.flybase.net/genomes/Drosophila_melanogaster/). After mapping the

sequence data, Differential expression analysis was performed between

control and *stil* mutant with *P35 OE* or*Tkv.CA OE* and calculated the

expression level of each gene.

DamID-seq and data processing

UASp-Dam and UASp-Dam-Stil flies were crossed with *NGT40; NosGal4-*

VP16 driver and maintained at 25°C. For each replicate, 100 ovaries of 50

flies were dissected in *Drosophila* Schneider's medium. Genomic DNA was

extracted using the QIAamp Fast DNA Tissue Kit (QIAGEN cat. no. 51404),

and methylated DNA was processed and amplified as described (Díaz-Torres

et al., 2021). Briefly, genomic DNA was digested overnight with *Dpn*I (NEB),

which cuts methylated GATC sites, and DamID adaptors were ligated to the

resulting fragments. A subsequent digestion with *Dpn*II (NEB), which

selectively cuts unmethylated GATC sites, was performed, and fragments

containing consecutive methylated GATCs were then amplified by PCR using

ligated adaptor-specific primers and PrimeSTAR® HS DNA Polymerase

(Takara).

The amplified DNA was sonicated using a Bioruptor (Diagenode) to obtain an

average fragment size of 300 bp. DamID adaptors were removed by digestion

with *Alw*I, and the DNA was purified using AMPure XP beads (Beckman

Coulter). For next-generation sequencing, the DNA was subjected to end-

repair, A-tailing, Illumina adaptor ligation, and PCR amplification. Single-end

100 bp reads were subsequently generated on a NovaSeq 6000 platform

(Illumina). Illumina NGS reads were aligned to the *Drosophila melanogaster*

reference genome (r6.32), and enrichment profiles were generated using the

damidseq_pipeline with default settings (Marshall & Brand, 2015), with

replicate libraries subsequently averaged. The Dam-Stil enrichment profile

was normalized by the Dam control profile and Dam-Stil peaks were identified

using find_peaks and annotated with the ChIPseeker pipeline (Marshall &

Brand, 2015) .

Fertility test

For female fertility assays, three virgin females were mated with three *y w*

males for 3 days per replicate. For male fertility assays, three males were

mated with three *y w* virgin females under the same conditions. After the

mating period, eggs were collected for 24 hours, followed by incubation for an

additional 24 hours. The total number of eggs and the number of hatched

eggs were then counted for each replicate.

S2 cell culture experiments

Drosophila Schneider S2 cells were cultured at 28°C in Schneider's medium

(Thermo) supplemented with 10% (v/v) fetal bovine serum (FBS), 100 U/mL

penicillin, and 100 µg/mL streptomycin (Thermo). Plasmids encoding Stil

variants were generated using the Gateway cloning system with (Life

Technologies) with pAFW, which drives the expression of N-terminal FLAG-

tagged fusion proteins under the control of the Actin5C promoter. *rpr*-GFP

reporter plasmid was constructed by replacing 2.4-kb tubulin promoter of

the *tubP*-GFP, a gift from Dr, Steve Cohen (Brennecke et al., 2003), with the

PCR-amplified 5 kb promoter region of *rpr*.

For the reporter assay, a total of 4 μg of plasmid DNA was used for each

transfection. Transgenes and reporter plasmids were included at 1 μg each as

needed, and the remaining amount was supplemented with the control

plasmid pAW without attR sites to reach 4 μg. The plasmid mixture was

combined with 20 μL of Hilymax transfection reagent (Dojindo) and

transfected into 1 mL of S2 cells seeded one day prior, following the

manufacturer's instructions. Two days of post-transfection, cells were

collected, sonicated, and followed by centrifugation. The supernatant was

mixed with 20% volume of 5× sample buffer. Proteins were then analyzed by

Western blotting using the following antibodies and Signal Enhancer Reagent

HIKARI (Nacalai): mouse anti-α-Tubulin [Santa Cruz, DM1A, 1:1000] rabbit

anti-GFP [Clonetech, 632569, 1:1000], and rabbit anti-FLAG [MBL, PM020,

1:1000]. Chemiluminescence detection was performed using the Chemi-Lumi

One reagent kit (Nacalai), and immunoreactive bands were visualized with a

ChemiDoc Touch imaging system (Bio-Rad) and analyzed using Fiji

(Schindelin et al., 2012).

For immunostaining, transfected S2 cells were collected at 2-days post-

transfection and plated onto poly-L-Lysine-coated coverslips for at least 30

min to promote adhesion. Cells were fixed in 5.3% (w/v) paraformaldehyde for

15 min, permeabilized with PBX for 10 min, and washed three times with

PBX. After blocking with PBX containing 4% (w/v) BSA for 30 min, cells were

incubated with primary and secondary antibodies for at least 1 hour each,

followed by nuclear staining with 1 μg/ml of DAPI for 10 min. After rinsing with

PBS, samples were mounted with Fluoro-KEEPER Antifade Reagent

(Nacalai). Images were acquired using a Zeiss LSM780 and 900 confocal

microscopes with a 63× objective lens and processed using Zen and ImageJ

software.

Competing Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgments

We thank Dr. Andrea H. Brand (The Gurdon Institute, University of Cambridge), Dr. Andreas Bergmann (UMass Chan Medical School, Massachusetts), and Dr. Dahua Chen (Yunnan University Institute of Biomedical Research, Kunming) for their generous gifts of the Dam CDS plasmid, the UASp-P35 transgenic fly line, and the UASp-Tkv.CA transgenic fly line, respectively. We acknowledge Bloomington Drosophila Stock Centre for the fly stocks. Confocal images and Western blotting data were acquired with LSM780, LSM900 and ChemiDoc Touch, respectively, at FBS Core Facility in The University of Osaka. We appreciate the insightful discussion and suggestions from all the members of KT's laboratory.

Author Contributions

All authors contributed to the experiment design. MM, KS, and KT performed all experiments. MM and KS performed the computational analyses. MM, KS, and KT wrote the paper. KS and KT supervised the project.

References

Abbott, M. K. & Lengyel, J. A. (1991). Embryonic head involution and rotation of male terminalia require the Drosophila locus head involution defective. *Genetics*, *129*(3), 783–789. https://doi.org/10.1093/genetics/129.3.783

Almeida, C., Correia, S., Rocha, E., Alves, Â., Ferraz, L., Silva, J., Sousa, M. & Barros, A. (2013). Caspase signalling pathways in human spermatogenesis. *Journal of Assisted Reproduction and Genetics*, *30*(4), 487–495. https://doi.org/10.1007/s10815-013-9938-8

Aravind, L. (2000). The BED finger, a novel DNA-binding domain in chromatin-boundary-element-binding proteins and transposases. *Trends in Biochemical Sciences*, *25*(9), 421–423. https://doi.org/10.1016/s0968-0004(00)01620-0

Aughey, G. N., Cheetham, S. W. & Southall, T. D. (2019). DamID as a versatile tool for understanding gene regulation. *Development*, *146*(6), dev173666. https://doi.org/10.1242/dev.173666

Baonza, A., Tur-Gracia, S., Pérez-Aguilera, M. & Estella, C. (2022). Regulation and coordination of the different DNA damage responses in Drosophila. *Frontiers in Cell and Developmental Biology*, *10*, 993257. https://doi.org/10.3389/fcell.2022.993257

Baum, J. S., Arama, E., Steller, H. & McCall, K. (2007). The Drosophila caspases Strica and Dronc function redundantly in programmed cell death during oogenesis. *Cell Death & Differentiation*, *14*(8), 1508–1517. https://doi.org/10.1038/sj.cdd.4402155

Beira, J. V., Springhorn, A., Gunther, S., Hufnagel, L., Pyrowolakis, G. & Vincent, J.-P. (2014). The Dpp/TGFβ-Dependent Corepressor Schnurri Protects Epithelial Cells from JNK-Induced Apoptosis in Drosophila Embryos. *Developmental Cell*, *31*(2), 240–247. https://doi.org/10.1016/j.devcel.2014.08.015

Bell, L. R., Horabin, J. I., Schedl, P. & Cline, T. W. (1991). Positive autoregulation of Sex-lethal by alternative splicing maintains the female determined state in Drosophila. *Cell*, *65*(2), 229–239. https://doi.org/10.1016/0092-8674(91)90157-t

Bellen, H. J., Levis, R. W., He, Y., Carlson, J. W., Evans-Holm, M., Bae, E., Kim, J., Metaxakis, A., Savakis, C., Schulze, K. L., Hoskins, R. A. & Spradling, A. C. (2011). The Drosophila Gene Disruption Project: Progress

Using Transposons With Distinctive Site Specificities. *Genetics*, *188*(3), 731–743. https://doi.org/10.1534/genetics.111.126995

Bellen, H. J., Levis, R. W., Liao, G., He, Y., Carlson, J. W., Tsang, G., Evans-Holm, M., Hiesinger, P. R., Schulze, K. L., Rubin, G. M., Hoskins, R. A. & Spradling, A. C. (2004). The BDGP Gene Disruption Project. *Genetics*, *167*(2), 761–781. https://doi.org/10.1534/genetics.104.026427

Brennecke, J., Hipfner, D. R., Stark, A., Russell, R. B. & Cohen, S. M. (2003). bantam Encodes a Developmentally Regulated microRNA that Controls Cell Proliferation and Regulates the Proapoptotic Gene hid in Drosophila. *Cell*, *113*(1), 25–36. https://doi.org/10.1016/s0092-8674(03)00231-9

Brodsky, M. H., Nordstrom, W., Tsang, G., Kwan, E., Rubin, G. M. & Abrams, J. M. (2000). Drosophila p53 Binds a Damage Response Element at the reaper Locus. *Cell*, *101*(1), 103–113. https://doi.org/10.1016/s0092-8674(00)80627-3

Casanueva, M. O. & Ferguson, E. L. (2004). Germline stem cell number in the Drosophila ovary is regulated by redundant mechanisms that control Dpp signaling. *Development*, *131*(9), 1881–1890. https://doi.org/10.1242/dev.01076

Chang, C.-H., Zhang, M., Rajapakshe, K., Coarfa, C., Edwards, D., Huang, S. & Rosen, J. M. (2015). Mammary Stem Cells and Tumor-Initiating Cells Are More Resistant to Apoptosis and Exhibit Increased DNA Repair Activity in Response to DNA Damage. *Stem Cell Reports*, *5*(3), 378–391. https://doi.org/10.1016/j.stemcr.2015.07.009

Chaudhry, S. R., Lopes, J., Levin, N. K., Kalpage, H. & Tainsky, M. A. (2021). Germline mutations in apoptosis pathway genes in ovarian cancer; the functional role of a TP53I3 (PIG3) variant in ROS production and DNA repair. *Cell Death Discovery*, *7*(1), 62. https://doi.org/10.1038/s41420-021-00442-y

Chen, D. & McKearin, D. (2003a). Dpp Signaling Silences bam Transcription Directly to Establish Asymmetric Divisions of Germline Stem Cells. *Current Biology*, *13*(20), 1786–1791. https://doi.org/10.1016/j.cub.2003.09.033

Chen, D. & McKearin, D. M. (2003b). A discrete transcriptional silencer in the bam gene determines asymmetric division of the Drosophila germline stem cell. *Development*, *130*(6), 1159–1170. https://doi.org/10.1242/dev.00325

Chen, P., Nordstrom, W., Gish, B. & Abrams, J. M. (1996). grim, a novel cell death gene in Drosophila. *Current Biology*, *6*(7), 767. https://doi.org/10.1016/s0960-9822(02)00588-2

Christich, A., Kauppila, S., Chen, P., Sogame, N., Ho, S.-I. & Abrams, J. M. (2002). The Damage-Responsive Drosophila Gene sickle Encodes a Novel IAP Binding Protein Similar to but Distinct from reaper, grim, and hid. *Current Biology*, *12*(2), 137–140. https://doi.org/10.1016/s0960-9822(01)00658-3

Dej, K. J. & Spradling, A. C. (1999). The endocycle controls nurse cell polytene chromosome structure during Drosophila oogenesis. *Development*, *126*(2), 293–303. https://doi.org/10.1242/dev.126.2.293

DeLuca, S. Z., Ghildiyal, M., Pang, L.-Y. & Spradling, A. C. (2020). Differentiating Drosophila female germ cells initiate Polycomb silencing by regulating PRC2-interacting proteins. *ELife*, *9*, e56922. https://doi.org/10.7554/elife.56922

Díaz-Torres, A., Rosales-Nieves, A. E., Pearson, J. R., Mateos, C. S.-C., Marín-Menguiano, M., Marshall, O. J., Brand, A. H. & González-Reyes, A. (2021). Stem cell niche organization in the Drosophila ovary requires the ECM component Perlecan. *Current Biology*, *31*(8), 1744-1753.e5. https://doi.org/10.1016/j.cub.2021.01.071

Drummond-Barbosa, D. & Spradling, A. C. (2001). Stem Cells and Their Progeny Respond to Nutritional Changes during Drosophila Oogenesis. *Developmental Biology*, *231*(1), 265–278. https://doi.org/10.1006/dbio.2000.0135

Freel, C. D., Richardson, D. A., Thomenius, M. J., Gan, E. C., Horn, S. R., Olson, M. R. & Kornbluth, S. (2008). Mitochondrial Localization of Reaper to Promote Inhibitors of Apoptosis Protein Degradation Conferred by GH3 Domain-Lipid Interactions*. *Journal of Biological Chemistry*, *283*(1), 367–379. https://doi.org/10.1074/jbc.m708931200

Fuchs, Y. & Steller, H. (2011). Programmed Cell Death in Animal Development and Disease. *Cell*, *147*(4), 742–758. https://doi.org/10.1016/j.cell.2011.10.033

Fuchs, Y. & Steller, H. (2015). Live to die another way: modes of programmed cell death and the signals emanating from dying cells. *Nature Reviews Molecular Cell Biology*, *16*(6), 329–344. https://doi.org/10.1038/nrm3999

G, P. & D., P. (1997). stand still, a Drosophila gene involved in the female
germline for proper survival, sex determination and differentiation.
Genetics, *145*(4), 975–987. https://doi.org/10.1093/genetics/145.4.975

Gleason, R. J., Anand, A., Kai, T. & Chen, X. (2018). Protecting and
Diversifying the Germline. *Genetics*, *208*(2), 435–471.
https://doi.org/10.1534/genetics.117.300208

Grieder, N. C., Cuevas, M. de & Spradling, A. C. (2000). The fusome
organizes the microtubule network during oocyte differentiation in
Drosophila. *Development*, *127*(19), 4253–4264.
https://doi.org/10.1242/dev.127.19.4253

Hashiyama, K., Hayashi, Y. & Kobayashi, S. (2011). Drosophila Sex lethal
Gene Initiates Female Development in Germline Progenitors. *Science*,
333(6044), 885–888. https://doi.org/10.1126/science.1208146

Holley, C. L., Olson, M. R., Colón-Ramos, D. A. & Kornbluth, S. (2002).
Reaper eliminates IAP proteins through stimulated IAP degradation and
generalized translational inhibition. *Nature Cell Biology*, *4*(6), 439–444.
https://doi.org/10.1038/ncb798

Hou, Y.-C. C., Chittaranjan, S., Barbosa, S. G., McCall, K. & Gorski, S. M.
(2008). Effector caspase Dcp-1 and IAP protein Bruce regulate starvation-
induced autophagy during Drosophila melanogaster oogenesis. *The
Journal of Cell Biology*, *182*(6), 1127–1139.
https://doi.org/10.1083/jcb.200712091

Huang, C., Zhao, S., Yang, Y., Guo, T., Ke, H., Mi, X., Qin, Y., Chen, Z.-J. &
Zhao, S. (2023). TP63 gain-of-function mutations cause premature ovarian
insufficiency by inducing oocyte apoptosis. *Journal of Clinical Investigation*,
133(5), e162315. https://doi.org/10.1172/jci162315

Jiang, C., Lamblin, A.-F. J., Steller, H. & Thummel, C. S. (2000). A Steroid-
Triggered Transcriptional Hierarchy Controls Salivary Gland Cell Death
during Drosophila Metamorphosis. *Molecular Cell*, *5*(3), 445–455.
https://doi.org/10.1016/s1097-2765(00)80439-6

Knudson, C. M., Tung, K. S. K., Tourtellotte, W. G., Brown, G. A. J. &
Korsmeyer, S. J. (1995). Bax-Deficient Mice with Lymphoid Hyperplasia
and Male Germ Cell Death. *Science*, *270*(5233), 96–99.
https://doi.org/10.1126/science.270.5233.96

Koch, R., Ledermann, R., Urwyler, O., Heller, M. & Suter, B. (2009).
Systematic Functional Analysis of Bicaudal-D Serine Phosphorylation and

Intragenic Suppression of a Female Sterile Allele of BicD. *PLoS ONE*, *4*(2),
e4552. https://doi.org/10.1371/journal.pone.0004552

Li, J., Jordana, L., Mehsen, H., Wang, X. & Archambault, V. (2024). Nuclear
reassembly defects after mitosis trigger apoptotic and p53-dependent
safeguard mechanisms in Drosophila. *PLOS Biology*, *22*(8), e3002780.
https://doi.org/10.1371/journal.pbio.3002780

Lohmann, I., McGinnis, N., Bodmer, M. & McGinnis, W. (2002). The
Drosophila Hox Gene Deformed Sculpts Head Morphology via Direct
Regulation of the Apoptosis Activator reaper. *Cell*, *110*(4), 457–466.
https://doi.org/10.1016/s0092-8674(02)00871-1

Lu, W.-J., Chapo, J., Roig, I. & Abrams, J. M. (2010). Meiotic Recombination
Provokes Functional Activation of the p53 Regulatory Network. *Science*,
328(5983), 1278–1281. https://doi.org/10.1126/science.1185640

Marshall, O. J. & Brand, A. H. (2015). damidseq_pipeline: an automated
pipeline for processing DamID sequencing datasets. *Bioinformatics*,
31(20), 3371–3373. https://doi.org/10.1093/bioinformatics/btv386

McCall, K. (2004). Eggs over easy: cell death in the Drosophila ovary.
Developmental Biology, *274*(1), 3–14.
https://doi.org/10.1016/j.ydbio.2004.07.017

McKowen, J. K., Avva, S. V. S. P., Maharjan, M., Duarte, F. M., Tome, J. M.,
Judd, J., Wood, J. L., Negedu, S., Dong, Y., Lis, J. T. & Hart, C. M. (2022).
The Drosophila BEAF insulator protein interacts with the polybromo subunit
of the PBAP chromatin remodeling complex. *G3*, *12*(11), jkac223.
https://doi.org/10.1093/g3journal/jkac223

Moschall, R., Rass, M., Rossbach, O., Lehmann, G., Kullmann, L., Eichner,
N., Strauss, D., Meister, G., Schneuwly, S., Krahn, M. P. & Medenbach, J.
(2019). Drosophila Sister-of-Sex-lethal reinforces a male-specific gene
expression pattern by controlling Sex-lethal alternative splicing. *Nucleic*
Acids Research, *47*(5), 2276–2288. https://doi.org/10.1093/nar/gky1284

Nair, S. & Baker, N. E. (2024). Extramacrochaetae regulates Notch signaling
in the Drosophila eye through non-apoptotic caspase activity. *ELife*, *12*,
RP91988. https://doi.org/10.7554/elife.91988

Nathansen, J., Meyer, F., Müller, L., Schmitz, M., Borgmann, K. & Dubrovska,
A. (2021). Beyond the Double-Strand Breaks: The Role of DNA Repair
Proteins in Cancer Stem-Cell Regulation. *Cancers*, *13*(19), 4818.
https://doi.org/10.3390/cancers13194818

Pang, L.-Y., DeLuca, S., Zhu, H., Urban, J. M. & Spradling, A. C. (2023).
Chromatin and gene expression changes during female Drosophila
germline stem cell development illuminate the biology of highly potent stem
cells. *ELife*, *12*(6), RP90509. https://doi.org/10.7554/elife.90509

Park, J.-H., Nguyen, T. T. N., Lee, E.-M., Castro-Aceituno, V., Wagle, R., Lee,
K.-S., Choi, J. & Song, Y.-H. (2019). Role of p53 isoforms in the DNA
damage response during Drosophila oogenesis. *Scientific Reports*, *9*(1),
11473. https://doi.org/10.1038/s41598-019-47913-y

Patil, V. S. & Kai, T. (2010). Repression of Retroelements in Drosophila
Germline via piRNA Pathway by the Tudor Domain Protein Tejas. *Current
Biology*, *20*(8), 724–730. https://doi.org/10.1016/j.cub.2010.02.046

Peterson, Jeanne S., Bass, B. P., Jue, D., Rodriguez, A., Abrams, J. M. &
McCall, K. (2007). Noncanonical cell death pathways act during Drosophila
oogenesis. *Genesis*, *45*(6), 396–404. https://doi.org/10.1002/dvg.20306

Peterson, J.S, Barkett, M. & McCall, K. (2003). Stage-specific regulation of
caspase activity in drosophila oogenesis. *Developmental Biology*, *260*(1),
113–123. https://doi.org/10.1016/s0012-1606(03)00240-9

PK, M., AR, C. & JR, J. (1996). Mutations in the gene stand still disrupt germ
cell differentiation in Drosophila ovaries. *Developmental Genetics*, *18*(4),
316–324. https://doi.org/10.1002/(sici)1520-6408(1996)18:4<316::aid-
dvg5>3.0.co;2-c

Primus, S., Pozmanter, C., Baxter, K. & Doren, M. V. (2019). Tudor-domain
containing protein 5-like promotes male sexual identity in the Drosophila
germline and is repressed in females by Sex lethal. *PLoS Genetics*, *15*(7),
e1007617. https://doi.org/10.1371/journal.pgen.1007617

Pritchett, T L & McCall, K. (2012). Role of the insulin/Tor signaling network in
starvation-induced programmed cell death in Drosophila oogenesis. *Cell
Death & Differentiation*, *19*(6), 1069–1079.
https://doi.org/10.1038/cdd.2011.200

Pritchett, Tracy L., Tanner, E. A. & McCall, K. (2009). Cracking open cell
death in the Drosophila ovary. *Apoptosis*, *14*(8), 969.
https://doi.org/10.1007/s10495-009-0369-z

Safa, A. R. (2016). Resistance to Cell Death and Its Modulation in Cancer
Stem Cells. *Critical Reviews in Oncogenesis*, *21*(3–4), 203–219.
https://doi.org/10.1615/critrevoncog.2016016976

Sahut-Barnola, I. & Pauli, D. (1999). The Drosophila gene stand still encodes
a germline chromatin-associated protein that controls the transcription of
the ovarian tumor gene. *Development*, *126*(9), 1917–1926.
https://doi.org/10.1242/dev.126.9.1917

Sandu, C., Ryoo, H. D. & Steller, H. (2010). Drosophila IAP antagonists form
multimeric complexes to promote cell death. *Journal of Cell Biology*,
190(6), 1039–1052. https://doi.org/10.1083/jcb.201004086

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M.,
Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez,
J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A.
(2012). Fiji: an open-source platform for biological-image analysis. *Nature*
Methods, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Schüpbach, T. & Wieschaus, E. (1989). Female sterile mutations on the
second chromosome of Drosophila melanogaster. I. Maternal effect
mutations. *Genetics*, *121*(1), 101–117.
https://doi.org/10.1093/genetics/121.1.101

Sokolova, O., Morgunova, V., Sizova, T. V., Komarov, P. A., Olenkina, O. M.,
Babaev, D. S., Mikhaleva, E. A., Kwon, D. A., Erokhin, M. & Kalmykova, A.
(2023). The insulator BEAF32 controls the spatial-temporal expression
profile of the telomeric retrotransposon TART in the Drosophila germline.
Development, *150*(12). https://doi.org/10.1242/dev.201678

Southall, T. D., Gold, K. S., Egger, B., Davidson, C. M., Caygill, E. E.,
Marshall, O. J. & Brand, A. H. (2013). Cell-Type-Specific Profiling of Gene
Expression and Chromatin Binding without Cell Isolation: Assaying RNA
Pol II Occupancy in Neural Stem Cells. *Developmental Cell*, *26*(1), 101–
112. https://doi.org/10.1016/j.devcel.2013.05.020

Spradling, A., Drummond-Barbosa, D. & Kai, T. (2001). Stem cells find their
niche. *Nature*, *414*(6859), 98–104. https://doi.org/10.1038/35102160

Sulekh, S., Ikegawa, Y., Naito, S., Oji, A., Hiratani, I. & Yoo, S. K. (2024). A
feedback loop that drives cell death and proliferation and its defect in
intestinal stem cells. *Life Science Alliance*, *7*(4), e202302238.
https://doi.org/10.26508/lsa.202302238

Tiwari, M. D., Zeitler, D. M., Meister, G. & Wodarz, A. (2019). Molecular
profiling of stem cell-like female germ line cells in Drosophila delineates
networks important for stemness and differentiation. *Biology Open*, *8*(11),
bio046789. https://doi.org/10.1242/bio.046789

Tue, N. T., Yoshioka, Y., Mizoguchi, M., Yoshida, H., Zurita, M. & Yamaguchi,
M. (2017). DREF plays multiple roles during Drosophila development.
Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms,
1860(6), 705–712. https://doi.org/10.1016/j.bbagrm.2017.03.004

Werz, C., Lee, T. V., Lee, P. L., Lackey, M., Bolduc, C., Stein, D. S. &
Bergmann, A. (2005). Mis-specified cells die by an active gene-directed
process, and inhibition of this death results in cell fate transformation in
Drosophila. *Development*, *132*(24), 5343–5352.
https://doi.org/10.1242/dev.02150

White, K., Grether, M. E., Abrams, J. M., Young, L., Farrell, K. & Steller, H.
(1994). Genetic Control of Programmed Cell Death in Drosophila. *Science*,
264(5159), 677–683. https://doi.org/10.1126/science.8171319

Wylie, A., Lu, W.-J., D'Brot, A., Buszczak, M. & Abrams, J. M. (2014). p53
activity is selectively licensed in the Drosophila stem cell compartment.
ELife, *3*, e01530. https://doi.org/10.7554/elife.01530

Xing, Y., Su, T. T. & Ruohola-Baker, H. (2015). Tie-mediated signal from
apoptotic cells protects stem cells in Drosophila melanogaster. *Nature
Communications*, *6*(1), 7058. https://doi.org/10.1038/ncomms8058

Xu, F., Suyama, R., Inada, T., Kawaguchi, S. & Kai, T. (2024). HemK2
functions for sufficient protein synthesis and RNA stability through eRF1
methylation during Drosophila oogenesis. *Development*, *151*(14).
https://doi.org/10.1242/dev.202795

Yan, N., Wu, J.-W., Chai, J., Li, W. & Shi, Y. (2004). Molecular mechanisms of
DrICE inhibition by DIAP1 and removal of inhibition by Reaper, Hid and
Grim. *Nature Structural & Molecular Biology*, *11*(5), 420–428.
https://doi.org/10.1038/nsmb764

Yin, V. P. & Thummel, C. S. (2004). A balance between the diap1 death
inhibitor and reaper and hid death inducers controls steroid-triggered cell
death in Drosophila. *Proceedings of the National Academy of Sciences*,
101(21), 8022–8027. https://doi.org/10.1073/pnas.0402647101

Zhang, Y., Lin, N., Carroll, P. M., Chan, G., Guan, B., Xiao, H., Yao, B., Wu,
S. S. & Zhou, L. (2008). Epigenetic Blocking of an Enhancer Region
Controls Irradiation-Induced Proapoptotic Gene Expression in Drosophila
Embryos. *Developmental Cell*, *14*(4), 481–493.
https://doi.org/10.1016/j.devcel.2008.01.018

Legends

Figure 1. Disruption of *stil* induces female germline cell depletion

(A) Schematic representation of the *Drosophila stil* gene locus showing the position of P-element insertion, *EY16156*, mutation site of *stil³*, and the target site of shRNA for knockdown. Black and blue arrows represent primers for RT-PCR and RT-qPCR, respectively. (B) Expression of *stil* and the control *α-tub* transcripts in *stil^EY16156^* mutant and the heterozygous control ovaries or testes analyzed by RT-PCR. (C) Representative images showing entire ovarian morphology (left panels) and ovarioles immunostained with with a germline marker, Vasa (red), and nuclear staining with DAPI (blue) (right panels) from *y w*, *stil^EY16156^/CyO*, *stil^EY16156^*, *stil^EY16156^/Df*, and *stil^EY16156^* expressing *6×Myc-Stil-FL* transgene driven by the germline-specific driver, *NGT40; NosGal4-VP16*. Scale bar: 1 mm for whole ovaries in the left panels, 50 µm for ovariole in the right panels. (D) Quantification of the percentage of germline cell-containing ovarioles in 2-3-days old females. Genotypes are indicated below the graph and the numbers of ovarioles assessed are shown above each bar. Error bars represent standard deviation (s.d.). (E) Analysis of egg laying and hatching rate. The egg numbers and their hatching rate by three females indicated genotypes corresponding to those in (C) mated with three *y w* males were measured daily (*n*=3). Error bars indicate s.d.

Figure 2. *stil* loss induces apoptotic cell death in female germline cells

(A) Quantitative RT-PCR confirming the knockdown efficiency in ovaries of *stil*-germline knockdown (*stil*-GLKD) where shRNA targeting stil transcript was driven by *NGT40; NosGal4-VP16*, and the control with shRNA alone. Expression levels of an internal control, *rp49*, and *stil* are normalized to that of *α-tub*. Error bar indicates s.d. (*n*=3). (B) Representative images showing entire ovarian morphology (left panels) and ovarioles immunostained with a germline marker, Vasa (red), and nuclear staining with DAPI (blue) (right panels) from *stil*-GLKD and the control with shRNA alone. Scale bar: 1 mm for whole ovaries and 50 µm for ovarioles, respectively. (C) Quantification of the percentage of germline cell-containing ovarioles in 2-3-days old females. Genotypes are indicated below the graph and the number of germarium assessed is noted above each bar. Error bars represent standard deviation (s.d.). (D) Quantification of the percentage of developed ovarioles defined as ovarioles that contain at least two egg chambers beyond the germarium. Genotypes are indicated below the graph and the number of germarium assessed is noted above each bar. Error bars represent standard deviation (s.d.). (E) Representative images of TUNEL-positive apoptotic cells (green), Vasa (red), and DAPI (blue) in control with shRNA alone and *stil*-GLKD germaria (white dotted). Scale bar: 20 µm. (F) Quantification of apoptosis in germarium with TUNEL assay in control and *stil*-GLKD. (G) Representative images of germaria (white dotted) from *stil^{EY16156}*, *stil^{EY16156}; P35 OE* (*NGT40; NosGal4-VP16> P35*), and *stil^{EY16156}; DIAP1 OE* (*NGT40; NosGal4-VP16> DIAP1*) immunostained

with Vasa (red) with DAPI (blue). Scale bar: 20 μm. (H) Quantification of the

percentage of germline cell-containing ovarioles in 2–3-day old females.

Genotypes are indicated below the graph and the number of germarium

assessed is noted above each bar. Error bars indicate standard deviation (s.d.).

Figure 3. *stil* regulates *rpr* expression in female germline cells

(A) Top 10 upregulated and downregulated genes in *stil*-deficient ovaries. Bars

indicate fold changes in *stil* mutant ovaries overexpressing (OE) of the

apoptosis inhibitor, P35 in germline cells (*stil^{EY16156}, NGT40; NosGal4-VP16>*

P35), compared to control with P35 OE in a heterozygous background

(*stil^{EY16156}/CyO, NGT40; NosGal4-VP16> P35*). Bars are according to adjusted

p-values (p-adj). (B) MA plot of apoptosis-related genes (GO:0006915) showing

their mean gene expression (CPM: counts per million) and fold change in

stil^{EY16156}; P35 OE comparing *stil^{EY16156}/CyO; P35 OE*. Each dot represents an

individual gene, with dot size inversely correlated with p-value adjusted (p-adj).

Larger dots highlight genes with higher statistical significance. The red dots

(upregulated genes) and blue dots (downregulated genes) represent the

differentially expressed RNAs with fold threshold>2.0, and p-adj<0.01. (C) Top

10 upregulated and downregulated genes related to apoptosis (GO:0006915).

Bars represent fold changes and are colored by adjusted p-values (p-adj), as

in (A). (D) Quantitative RT-PCR analysis of proapoptotic genes, *rpr* and *hid* in

stil^{EY16156}; P35 OE and *stil^{EY16156}/CyO; P35 OE* ovaries. Expression levels were

normalized to *αtub* with *rp49* as an internal control. Error bars indicate s.d. (*n*=3).

(E) Immunofluorescence of Vasa (red) and DAPI (blue) in ovarioles from

stil^{EY16156}/CyO; rpr^{87}/+, *stil^{EY16156}/CyO; rpr^{87}*, *stil^{EY16156}; rpr^{87}/+*, and *stil^{EY16156};*

rpr^{87}. Scale bar: 50 µm. (F) Quantification of the percentage of germline cell-

containing ovarioles in 2-3-days old females. Genotypes of females and the

number of germarium assessed are noted above each bar. Error bars represent

standard deviation (s.d.). (G) Quantification of the percentage of developed

ovarioles defined as ovarioles that contain at least two egg chambers beyond

the germarium. Genotypes females and the number of germarium assessed

are noted above each bar. Error bars represent standard deviation (s.d.). (H)

Analysis of egg laying and hatching rate. Daily egg laying by three females

indicated genotypes mated with three *y w* male and their hatching rate are

shown (*n*=3). Error bar indicates s.d.

1022

Figure 4. Stil interacts with *rpr*-transcriptional control region to repress

its expression

(A) Distribution of DamID peaks across entire genome categorized by genomic

features including promoter (≤1, 1–2, and 2–3 kb), UTR, exon, intron, distal

intergenic, and downstream. (B) The plot showing the log_2 ratio of Dam-Stil

peaks to Dam-alone control across the *rpr* genomic region. The blue bar on the

bottom indicates the *rpr* gene body, and the arrow shows the direction of *rpr*

transcription. Three independent biological replicates were performed for both

Dam-Stil and Dam-alone and data were merged for visualization. (C) Schematic

representation of the Stil full-length (Stil-FL), the truncated variants: Stil-NT (1st

to 169th aa) and Stil-CT (169th to the end), and amino acid substitutions of the

C2H2 zinc finger CCYC: Stil-AAYA. (D) *stil^{EY16156}* ovaries expressing Myc-

tagged Stil variants corresponding to C in germline cells. Ovaries are

immunostained with a germline marker, Vasa (red), and nuclear staining with

DAPI (blue). Scale bar: 50 µm. (E) Quantification of the percentage of germline

cell-containing ovarioles in 2-3 days-old females. Genotypes are indicated

below the graph and the number of germarium assessed is noted above each

bar. Error bars represent standard deviation (s.d.). (F) Quantification of the

percentage of developed ovarioles defined as ovarioles that contain at least two

egg chambers beyond the germarium. Genotypes are indicated below the

graph and the number of germarium assessed is noted above each bar. Error

bars indicate standard deviation (s.d.). (G) Schematic representation of *rpr*-

GFP reporter containing the 5 kb upstream region of the *rpr* gene fused to GFP

CDS. (H) Reporter assay in S2 cells followed by western blot analysis to detect

GFP and α-Tubulin (αTub) as a loading control. Co-transfection of the *rpr*-GFP

reporter and FLAG-tagged Stil variants were performed in S2 cells. The lower

panel shows the signal ratio of GFP expression to αTub, normalized to that in

cells transfected with the reporter alone. Error bars represent the standard

deviation (s.d.) (*n* = 3).

1052

Figure 5. *stil* mutant germline stem cells resist apoptosis

(A) Fluorescent immunostaining with anti-Engrailed (red), anti-Vasa (green),

and DAPI (blue) for terminal filaments of late 3rd instar larval (LL3) ovaries from

the heterozygous control, *stil^{EY16156}/CyO* and *stil* mutant, *stil^{EY16156}*. Scale bar:

50 μm. (B) Quantification of surviving germline cells in 3rd instar larval ovaries

from *stil^{EY16156}/CyO* and *stil^{EY16156}*. (C) Immunostaining of Vasa (red) and DAPI

(blue) in control, *Tkv.CA* (constitutive active Thickveins) *OE* (*NGT40; NosGal4-*

VP16>Tkv.CA), and *bam* mutant (*bam^{Δ86}*) germaria (white dotted) in the control

heterozygous (*stil^{EY16156}/CyO*) or *stil* mutant (*stil^{EY16156}*) background. Scale bar:

20 μm. (D) Quantification germline cell-containing germaria per ovary in 2-3

days-old females. Genotypes and the number of germarium assessed are

indicated above each bar. Error bars represent standard deviation (s.d.). (E)

Representative images of germaria (white dashed lines) showing TUNEL-

positive apoptotic cells (green), Vasa (red), and DAPI (blue) in the control

tumorous germaria (*stil^{EY16156}/CyO; Tkv.CA OE*) and tumorous germaria of *stil*

mutant ovaries (*stil^{EY16156}; Tkv.CA OE*). White arrowheads indicate TUNEL-

positive apoptotic cells. Scale bar: 20 μm. (F) Quantification of apoptosis in

germaria from the control tumorous ovaries (*stil^{EY16156}/CyO; Tkv.CA OE*) and

stil mutant ovaries (*stil^{EY16156}; Tkv.CA OE*). (G) MA plot of apoptosis-related

genes showing the mean gene expression (CPM: counts per million) and fold

change in *stil* mutant ovaries (*stil^EY16156; Tkv.CA OE*) compared to the control

tumorous ovaries (*stil^EY16156/CyO; Tkv.CA OE*). Each dot represents an

individual gene, with dot size inversely correlated with p-adj. The blue dots

represent the downregulated genes with fold change<-2.0, and p-adj<0.01. (H)

Quantitative RT-PCR analysis of *rpr* in the control tumorous ovaries

(*stil^EY16156/CyO; Tkv.CA OE*) and *stil* mutant ovaries (*stil^EY16156; Tkv.CA OE*).

Expression levels of an internal control, *rp49*, and *rpr* are normalized to that of

α-tub. Error bars indicate s.d. (*n*=3).

1081

Figure 6. chromatin state changes in female germline cells during

oogenesis

(A) ATAC-seq signal intensity (RPM-normalized) at the *rpr* locus in germline

stem cells (GSCs: 2N and 4N) and nurse cells (NCs) at different ploidy levels

(4C to 512C). The Y-axis is fixed at 0-30 PRM for all tracks. For 64C, 128C,

256C, and 512C NCs, data from three biological replicates are averaged; all

other samples are based on single replicates.

(B) H3K9me3 ChIP-seq signal at the *rpr* locus in 4C GSCs and 4C NCs. (C)

H3K27me3 ChIP-seq signal at the *rpr* locus in GSCs and stage 5 (st5) NCs.

1091

1092

Supplementary Legends

Figure S1. *stil* function is indispensable in female germline cells not male germline cells

(A) Immunostaining of ovaries from *y w*, *stil³/CyO*, *stil³/Df*, and *stil³/stil^{EY16156}* with antibody against Vasa (red) with DAPI (blue). Scale bar: 50 μm. (B) Quantification of the percentage of germline cell-containing ovarioles in 2-3 days old females. (C) Immunostaining of the apical end of testes from *stil^{EY16156}/CyO* and *stil^{EY16156}* with anti-Vasa antibody (red) and DAPI (blue) (top panels). Seminal vesicles (SV) harboring sperms are also stained with DAPI (blue) (bottom panels). Asterisk denotes the apex of testis. Scale bar: 50μm (apical end of testes, top) and 20 μm (SV, bottom). (D) The numbers of egg laying and hatching rate. Daily egg laying by three *y w* females mated with three males of the indicated genotypes are shown (*n*=3). The number of ovarioles assessed is noted above each bar. Error bar indicates s.d.

Figure S2. Germline knockdown of *stil*, but not somatic knockdown, specifically disrupts egg development

(A) Analysis of egg laying and hatching rates. The number of laid eggs and their hatching rates are measured daily for three females of the indicated genotypes: control, *stil*-germline knockdown (*stil*-GLKD) driven by *NGT40; NosGal4-VP16*, and *stil*-somatic knockdown (*stil*-STKD) driven by *tj-Gal4*, each mated with three *y w* males (*n* = 3). Error bars indicate standard deviation (s.d.). (B) RT-

1115   PCR analysis of *sxl* and the control, *α-tub* transcripts in ovaries and testes from

1116   control and *stil*-GLKD flies, respectively.

1117

Figure S3. Egg development is disrupted at the mid-stage of oogenesis

in *stil* mutants rescued by P35 or DIAP

Immunostaining of ovarioles from *y w, stil^EY16156^/CyO; P35 OE* (*NGT40;*

NosGal4-VP16> P35), *stil^EY16156^; P35 OE* and *stil^EY16156^; DIAP OE* (*NGT40;*

NosGal4-VP16> DIAP1) flies with antibody against Vasa (red) and DAPI (blue).

The enlarged image highlights a degenerated egg chamber at the mid-stage of

oogenesis in *stil^EY16156^; P35 OE*. Scale bar: 50 μm.

1125

Figure S4. BED-type zinc finger of Stil and transcriptome analysis of

germline-rescued *stil* mutant ovaries

(A) A schematic presentation of Stil protein, highlighting the BED-type zinc

finger (BED, green) and the nuclear localization signal (NLS, blue). A multiple

sequence alignment of Stil with three other proteins containing BED-type zinc

finger motif is shown. The BED-type zinc finger motif consists of two cysteine

and histidine residues (red) and aromatic residues (yellow) in the N-terminal

region. (B) Transcriptome analysis of germline-rescued *stil* mutant ovaries

expressing P35. Differential expressed genes (DEGs) are obtained with

DEseq2. Volcano plot displays the adjusted *p*-value (*p*-adj) versus log2-fold

change (log2FC) of expression level in *stil^EY16156^; P35 OE* compared to that in

stil^EY16156^/CyO; P35 OE. Significantly upregulated genes (log2FC > 1, *p*-adj <

0.01) and downregulated genes (log2FC < -1, *p*-adj < 0.01) are highlighted in

red and blue, respectively, while non-DEGs are shown in black.

Figure S5. *rpr*, but not *hid*, plays a central role in *stil*-deficient female germline cell death

(A-C) Quantitative RT-PCR are performed on ovaries from germline-rescued

stil mutants, as well as ovaries and testes with germline-specific *stil* knockdown

using distinct drivers. Expression levels are normalized to *αtub* with *rp49*

serving as an internal control. (A) Early-stage ovaries are isolated by dissecting

stage 4–5 oocytes from *stil^EY16156^/CyO; P35 OE* and *stil^EY16156^; P35 OE* flies (B)

stil knockdown in female germline cells is achieved using *NGT40-Gal4;*

NosGal4-VP16, *Bam-Gal4*, and *matα-tub-Gal4* drivers. (C) *stil* knockdown in

male germline cells is achieved using *NGT40-Gal4; NosGal4-VP16* and *Bam-*

Gal4 drivers. Error bars represent standard deviation (s.d.). (D) Immunostaining

of *stil^EY16156^; hid^05014^/+* and *stil^EY16156^; hid^05014^/Df(3L)X14* ovaries with antibody

against Vasa (red) with DAPI (blue). Scale bar: 50 μm. (E) Quantification of the

percentage of germline cell-containing ovarioles in 2-3-days old females. The

number of ovarioles assessed is noted above each bar.

Figure S6. Stil variants localization and control validation in S2 reporter assay

(A) Immunofluorescence images showing Myc-tagged Stil variants (green) and DNA (DAPI, blue) in stage 3~4 egg chambers of the ovaries. Insets show enlarged views of nurse cell nuclei. Scale bar: 50 μm (egg chamber) and 20 μm (nurse cell nuclei). (B) Immunostaining of S2 cells expressing FLAG-tagged Stil variants with an anti-FLAG (green) antibody and DAPI (blue). (C) Schematic representation of the *tubP*-GFP reporter construct, in which the 2.4-kb upstream region of the *αTub84B* gene is fused to GFP. (D) Reporter assay in S2 cells followed by western blot analysis to detect GFP expression. Arrowheads indicate Stil-variant proteins. *α*-Tubulin (αTub) is used as a loading control. The lower panel quantifies the GFP/αTub signal ratio, normalized to that in S2 cells transfected with the reporter alone. Error bars indicate standard deviation (s.d.) ($n = 3$).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

ATAC

GSC 2C
GSC 4C
NC 4C
NC 8C
NC 16C
NC 32C
NC 64C
NC 128C
NC 256C
NC 512C

rpr

B

H3K9me3

GSC 4C
NC 4C

rpr

C

H3K27me3

GSC 4C
St5 NC

rpr