Comparing analytical protocols for identifying causes of population changes

Abstract

1. Understanding the factors driving population dynamics is critical for conservation efforts, enabling e.g. the implementation of effective recovery strategies. In this study, we analysed common methods for their ability to correctly identify factors contributing to population changes.

2. We compared time series (TS) and species distribution models (SDM), alongside process-oriented models (POM), which integrate mechanistic components with a correlative approach. To achieve this, we used a virtual ecologist approach to generate synthetic data where the true underlying drivers were known. We then fitted all three models to the data, mimicking the TS and SDM by aggregating the data within a specific domain (spatial or temporal).

3. Our results show that POM outperforms the other methods in all performance measures used, exhibiting the highest accuracy (0.88), sensitivity (0.84), and specificity (0.93) in terms of identifying factors that contribute to population changes. SDM demonstrated medium (0.68), while TS had the lowest accuracy (0.50). Furthermore, our results indicate that variable selection marginally improved the performance of suboptimal models (TS) but significantly reduced the accuracy of alternative methods (SDM and POM).

4. Our findings highlight the importance of incorporating mechanistic aspects into

species distribution modelling to more accurately identify the causes of population

changes. These results have the potential to be applicable to a range of ecological and

conservation studies and would facilitate the selection of analytical protocol for

developing specific indicators to monitor population changes.

Key words: population change, time-series models, species distribution models, process-

oriented models, generalized additive modelling, simulation study, virtual species.

1. Introduction

Identifying the causes of population change is a central issue in ecology and conservation

biology. This problem is particularly relevant today, as anthropogenic environmental

impacts are causing rapid changes in the populations of many species (Hutchings & Reynolds,

2004; Rigal et al., 2023). This leads to dramatic declines in some species, often culminating

in extinction and irreversible loss of biodiversity (Ceballos et al., 2017). At the same time,

biological invasions lead to human-nature conflicts, with measurable economic losses

(Knowler, 2005; Nentwig et al., 2018; Perrings et al., 2002). Accurate identification of the

underlying causes is essential in order to create specific indicators for reporting population

change. These can then be used in an indicator-based monitoring framework, thus enabling

the formulation of effective responses and enhancing our ability to prevent harmful change.

A broad range of analytical approaches could be used to identify the causes of population

changes (Green, 1995; T. F. Johnson et al., 2024; Rhodes et al., 2011; Zachariah Peery et al.,

2004). For example, demographic modelling can provide profound insights into the

intrapopulation dynamics that contribute to population declines (Allen et al., 2022; Chiquet

et al., 2013; Enneson & Litzgus, 2008; H. E. Johnson et al., 2010). However, such models

require specific biological data that are difficult to obtain and often unavailable.

Consequently, when only abundance data are accessible, ecologists typically rely on

statistical methods based on the correlative approach (Green, 1995; Zachariah Peery et al.,

2004). Two of the most commonly used statistical modelling approaches for this purpose are

time series (Shumway & Stoffer, 2005; Wauchope et al., 2021) and species distribution

modelling (Elith & Leathwick, 2009; Guisan & Thuiller, 2005).

Time series modelling is used for census and survey data to analyse trends and seasonal

variations over time. Numerous ecological studies have been conducted using temporal data

across a wide range of taxa, including predictors such as weather or environmental variables

(Di Fonzo et al., 2016; Kölzsch et al., 2007). Conventional time series analysis, at least in the

context of identifying factors influencing population changes, does not utilise spatially

explicit abundance data. Instead, population size and the corresponding environmental data

are aggregated to the specific spatial scale at which the analysis is conducted (Hsieh et al.,

2005; Huthings 2000; Knape & de Valpine, 2011). Importantly, time series analysis

inherently treats a single unit of time (usually a year) as a single data point. As a result, it is

challenging to make inferences with this approach using typical ecological datasets (multi-

year time series). In the case of 'gold standard' datasets (time series >20 years), such

spatiotemporal compression, together with the lack of spatial replication, reduces statistical

power and makes it significantly more difficult to detect multidimensional or non-linear

relationships.

Species distribution models are very popular and have demonstrated remarkable versatility

in ecological applications, serving as valuable tools for understanding species-environment

relationships, predicting potential distributions, and informing conservation strategies

(Zurell et al., 2020). Typically, they combine environmental predictors with information on

the species occurrence or abundance, without considering the temporal dimension (Elith &

Leathwick, 2009; Guisan & Thuiller, 2005), which leads to several limitations. Firstly, they

assume that populations are in a state of equilibrium with their environment, which is not

usually the case (Araújo & Guisan, 2006; Guisan & Thuiller, 2005; Pagel & Schurr, 2012; Zurell

et al., 2009). Secondly, this approach ignores ecological dynamism such as density-dependent

demographic processes, biotic interactions, dispersal, and other fundamental range-shaping

phenomena that lead to forecast mismatches (Franklin, 2010; Holt, 2009; Normand et al.,

2014; Zurell et al., 2009), and have difficulty in extrapolating models to a novel environment

(Briscoe et al., 2019). However, it is worth mentioning that classical SDMs are undergoing

extensive development with several novel approaches integrating temporal aspects, such as

phenology, with these models (Zurell et al., 2024). Furthermore, there is considerable

potential in the application of demographic SDM, which incorporates demographic processes

or multi-species models (joint-SDM) that facilitate the modelling of species communities

(Zurell, 2017).

In this study, we compare two widely used protocols for the identification of factors

influencing population abundance and, consequently, the causes of population change,

namely time series (TS), and species distribution model (SDM), with a process-oriented

model (POM). The latter constitutes our proposal for an easily implementable spatio-

temporal model incorporating mechanistic elements such as, density dependence, spatial

and temporal autocorrelation (acting as a proxy for both dispersal and large-scale

environmental gradients), random effects for space and time (absorbing environmental

noise), and random effects for observers (mimicking the observational process). In order to

achieve this objective, a virtual species was generated using large-scale and long-term bird

monitoring data, thereby providing a known reference. The effectiveness of TS, SDM and the

proposed POM in identifying the biotic and abiotic factors responsible for shaping population

dynamics was then tested. We hypothesise that POM will outperform both TS and SDM in

identification of the factors driving population dynamics as its additional components will

increase the overall effectiveness of the models.

98   2. Material and Methods

99   2.1 Study design

The aim of our work was to compare the performance of analytical approaches typically used

to identify factors influencing population change from abundance data. One is based on

inference from time series correlations, while the other is based on the use of spatial data

and the construction of species distribution models. In addition, we propose a more

sophisticated approach, which we call 'process-oriented', and test its performance against

other methods.

Comparing different methods requires knowledge of the reference. Therefore, we employed

an approach to generate a virtual species (VS) based on predefined relationships between

environmental factors and population abundance, and then test the effectiveness of the

selection of variables actually used in VS generation (Fig. 1) (Malinowska et al., 2023).

Figure 1. The study design. First, we fitted a model to the spatio-temporal bird abundance

data using full set of predictors. We then selected half of the most important predictors from

the full model to fit a reduced model. Multiple predictions were then generated from this

reduced model, forming many instances of virtual species. By introducing a realistic level of

randomness at this stage, each virtual species was different, even though it was generated

from the same model. The sampling procedure, which mimicked data collection, produced

virtual ecologist data. These data, which provided a known reference, were then used as input

to test the three analytical approaches – time series (TS), species distribution model (SDM),

and process oriented approach (POM). Each of these models was fitted with the full set of

predictors and then the variables were classified as "significant" using the $p<0.05$ criterion.

As the true relevance of the predictors was known, it was possible to evaluate the three

methods in terms of their effectiveness in identifying the real factors influencing population

change.

To test the effectiveness of the analytical methods, we used abundance data for 10 pairs of

western Palearctic passerine bird species (Supplementary File 1, Table S1). For each pair,

100 virtual instances of the species were generated, resulting in a total of 1000 replications

in this computational experiment. For detailed algorithm of the study design see

Supplementary File 2.

2.2 Bird abundance data

The study uses data from 14 common western Palearctic passerine birds organised into 10

pairs (see Table S1 for details). Bird data were obtained from the Common Breeding Birds

Survey in Poland (MPPL), and span 20 years (2001-2020). The monitoring programme

strictly follows the methodology of the British Breeding Bird Survey (Gregory et al., 1996)

and is also included in the Pan-European Common Bird Monitoring Scheme (PECBMS, Brlík

et al., 2021). Specific details on data collection are presented in Supplementary File 1. The

MPPL programme provides counts that are used to calculate species densities (pairs/km^2)

using the distance sampling method (Buckland et al., 2001, 2004).

2.3 Environmental data

In the study we used dynamic environmental data, such as annual land cover data that were

derived from ESA CCI Land Cover (Defourny et al., 2017, 2021) and monthly climate data

from TerraClimate (Abatzoglou et al., 2018), which were obtained for each MPPL survey site

(see Supplementary File 1 and Table S2 for full details). The environmental variables

mentioned above are bionomic, which means that they are neither interacting nor

consumable (Soberón & Nakamura, 2009). Therefore, we also incorporated biotic interaction

into the model, as it can influence species-environmental relationships and species co-

occurrence patterns (Wisz et al., 2013). We used the example of social information transfer,

which is a mechanism of obtaining, using, and sharing environmental information from

conspecifics or heterospecifics (Szymkowiak, 2013). Social cues may provide useful

information during the habitat selection process, potentially leading to various population-

level consequences (Szymkowiak, 2013). To integrate this mechanism into models, we

selected a single variable representing a biotic component, which was the abundance of an

interacting species, defined as a species for which there is external, empirical evidence that

it interacts biotically with the target species. The set of potential interacting species (Table

S1) was restricted to positively interacting species only, due to the mechanism of information

use during habitat selection (Szymkowiak, 2013).

2.4 Virtual species and virtual ecologist

The virtual data, which serve as a known reference, were generated using a framework

described in Malinowska et. al (2023), which could be summarised with the following steps:

1) **Fitting a model to the observed data**. In this step we used Generalised Additive Mixed

Models (GAMM) (Wood, 2017), as they offer a range of distributions relevant to population

modelling, and are capable of handling zero-inflated and overdispersed data. Flexible

functions can also be used, which is important as species-environment relationships tend

to be non-linear and complex. Hierarchical structures typical of biodiversity monitoring

data can be incorporated via random effects, and autocorrelation structures allow for

increased biological realism.

The model was fitted with 20 variables, including the interacting species. Half of the

variables were then removed by backward elimination. In this way, the final (reduced)

model, which was later used to generate the VS, was built using only part of the available

variables. In this way it was possible to create a scenario in which certain factors had an

effect on species abundance and others did not.

2) **Generating the VS.** Predictions from the above reduced model were computed and

treated as a virtual species abundance. Although they were generated from the same

model, the randomness required for statistical inference was achieved by assigning

random intercepts to sites and then generating random realisations from the original

distribution (Malinowska et al. 2023). As a result, virtual species abundance were

obtained for each site i and year t.

3) **Sampling**. Virtual ecologist data was obtained by simulating the observational process

applied to the VS. For each virtual observer, we simulated the sampling error p by

drawing a random value from a lognormal distribution with a mean of 1 and a variance

which was estimated by fitting the GAMM (this was the variance of the random intercepts

for the observer identifiers). The virtual ecologist data were than obtained by

multiplying the VS densities by a relevant p.

4) **Calibration**. VE data generated in this way tend to overestimate prevalence and

underestimate variance (Malinowska et al., 2023). Therefore, to increase ecological

realism, we calibrated the VE data using a procedure known as quantile mapping. The

virtual data were mapped so that the frequency distribution of the transformed data

matched the frequency distribution of the data observed. Simulations show that this

procedure is capable of generating virtual data with distributional properties that are

indistinguishable from the real data (Malinowska et al., 2023).

We repeated the described process (steps 2-4) 100 times to generate multiple instances

of virtual ecologist data for every 10 case studies. These virtual data were then used as a

known reference to test the effectiveness of variable selection for the three methods

considered.

2.5 Models tested

Many methods can be used to make inferences about the effects of environmental factors on

abundance, provided they allow specific properties of these data to be modelled. The obvious

choice seems to be Generalized Linear Models (GLM). However, the relationships between

environmental factors and population densities very often are non-monotonic because

organisms have certain optimal preferences along a gradient of environmental

characteristics, which are best described by unimodal functions. This implies that it is not

feasible to describe a species' preference for a particular environmental factor with a single

parameter. Consequently, it is usually not possible to ascertain unambiguously the direction

or magnitude of the effect. For this reason, we do not attempt to reconstruct effect sizes, but

rather test the effectiveness of a given statistical method in terms of a binary decision as to

whether or not a given variable actually affects population numbers.

It is evident that this creates problems of interpretation, but the solution seems relatively

straightforward and may involve exploratory analysis and visual evaluation of response

curves. Thus, the approach we propose is to first assess whether a given environmental factor

is important and has an effect on population densities, and then to analyse the direction and

strength of the effect exploratively. This can also be done formally, using methods that can

infer non-monotonic effects.

For these reasons, we used GAM as a flexible way of testing the models for their ability to

select the true drivers of population change. An additional advantage of this method is that it

not only allows non-linear relationships to be modelled, but also allows the smoothness of

those relationships to be reduced until linearity is achieved. Thus, if the patterns in the data

are linear, or the data are so sparse that more complex functions cannot be estimated, GAM

is reduced to GLM.

2.5.1 Time series

To mimic the time series data, we averaged both population density information and each of

the m environmental variables across space and then built a GAM model:

$$222 \qquad\qquad\qquad N_t \sim T_\xi(\mu_t, \phi)$$

$$223 \qquad\qquad\qquad \ln(\mu_t) = \alpha + \sum_{j=1}^{m} f_j\left(X_{t,j}\right)$$

224   Here, population density N_t, which is assumed to come from the Tweedie distribution with

225   index parameter ξ and dispersion parameter ϕ, is indexed by the year identifier t. The

226   natural logarithm of the expected population density is a sum of the intercept α and smooth

227   functions f_j, each fitted to predictor X_j.

228   2.5.2. Species distribution model

229   To mimic the static spatial data, we averaged both population densities and each of the m

230   environmental variables across time (i.e. by calculating arithmetic means for each of 1x1 km

231   MPPL squares). Then, the following SDM was build:

$$232 \qquad\qquad\qquad N_i \sim T_\xi(\mu_i, \phi)$$

$$233 \qquad\qquad\qquad \ln(\mu_i) = \alpha + \sum_{j=1}^{m} f_j\left(X_{i,j}\right)$$

234   The model structure is analogous to the time series model (above) except for the population

235   density N is varying in space, i.e. it is indexed by sampling site identifiers denoted as i.

236   2.5.3 Process-oriented model

In this approach both species and environmental data are dynamic, i.e. they vary in space and

time. Moreover, the model was extended by explicitly embedding the following processes

into the model structure: density-dependence, temporal and spatial autocorrelation, random

temporal and spatial variation and observational process.

The density dependence was modelled using Gompertz dynamics based on the following

concept: for every site i, the population growth rate between time $t-1$ and time t can be

expressed as $r = \ln(N_t/N_{t-1})$, which implies $\ln(N_t) = \ln(N_{t-1}) + r$. Density-dependence

takes the form of self-regulation, where r is a function of N_{t-1}. Thus, we can substitute r by

an arbitrary function of N_{t-1} to obtain:

$$\ln(N_t) = \ln(N_{t-1}) + f_d\big(\ln(N_{t-1})\big),$$

where f_d is a function determining how population growth rate changes with changing

population density. Traditionally, this function is assumed to be monotonically decreasing,

however it can take many forms, including non-monotonic (Allee effect). Here, to allow for

flexibility and generality in density-dependent effects, we allowed the f_d function to be

smooth and non-parametric. To do so, we fitted the following GAMM model:

$$N_{i,t} \sim T_\xi\big(\mu_{i,t}, \phi\big)$$

$$\ln\big(\mu_{i,t}\big) = \alpha + \ln\big(N_{i,t-1}\big) + f_d\left(\ln\big(N_{i,t-1}\big)\right) +$$

$$+ \sum_{j=1}^{m} f_j\big(X_{i,t,j}\big) + \mathcal{GP}\big(f_t(t), u(t,t')\big) + \mathcal{GP}\big(f_s(x), v(x,x')\big) + a_i + b_t + o_{i,t}$$

$$u(t,t') = c_0(\| \, t - t' \, \|, \rho_t), \qquad v(x,x') = c_0(\| \, x - x' \, \|, \rho_x)$$

$$c_0(d, \rho) = (1 + d/\rho)\exp(-d/\rho)$$

257
$$a_i \sim N(0, \sigma_a^2), \qquad b_t \sim N(0, \sigma_b^2), \qquad o_{i,t} \sim N(0, \sigma_o^2)$$

Population density $N_{i,t}$, which is assumed to come from the Tweedie distribution, is indexed

by the site identifier i and the year identifier t. As in the previous models, the natural

logarithm of expected density $\mu_{i,t}$ is the sum of smooth functions fitted to m predictors.

However, in this model, all predictors vary in both space and time.

In addition to the density-dependence, this model has been extended to include the following

processes:

1. Temporal autocorrelation, modelled as a Gaussian process $\mathcal{GP}$ with the expectation f_t

which itself is modelled as a smooth function of time and the covariance matrix u

describing temporal autocorrelation structure. The elements of matrix u are modelled

as a function of a between-year distance using the Matérn correlation function c_0 with

the parameter ρ_t.

2. Spatial autocorrelation, modelled as a Gaussian process $\mathcal{GP}$ with the expectation f_s

which is a smooth, 2-dimensional function of geographic coordinates (denoted as x) and

the covariance matrix v describing a spatial autocorrelation structure. The elements of

matrix v are modelled as a function of geographic distance between sampling sites using

the Matérn correlation function c_0 with the parameter ρ_x.

3. Random spatial variation, modelled as a normally distributed random intercept for each

sampling site i with the mean zero and the variance σ_a^2.

4. Random temporal variation, modelled as a normally distributed random intercept for

each year t with the mean zero and the variance σ_b^2.

5.   Observational process, modelled as a normally distributed random intercept for each

observer, with the mean zero and the variance σ_o^2.

2.6 Evaluation of models

After generating virtual data simulating population densities, we were able to make model

comparisons. Since the virtual data were replicated multiple times (100 simulations for each

of the 10 virtual species pairs), formal statistical inference was possible.

A commonly used approach to improve the interpretability of a model while increasing its

predictive accuracy is variable selection. In the case of GAM, Marra & Wood (2011) used an

extensive simulation study to demonstrate that, in terms of predictive power, shrinkage

approaches to variable selection perform significantly better than competing methods. To

test this in the context of our study, and identify which variables were 'significant', the models

were fitted in two configurations: one in which variable selection was not applied and one in

which it was. In the latter case, the double penalty shrinkage was used. In this approach, in

addition to the standard penalty that controls for the shape of smooth functions and thus

prevents them from overfitting, an additional penalty is constructed that penalises the size

of functions. In this way, the model can effectively shrink the entire function for a variable to

zero if its contribution is negligible, thus performing automatic variable selection (Marra &

Wood, 2011). Technically, the double penalty is invoked by using `gam(…, select = TRUE)`

from the `mgcv` package (Wood, 2017).

For each of the fitted models, we classified variables as 'significant' if the estimated p-value

was below the threshold of 0.05. This is, of course, an arbitrary criterion, but the objective of

this study is to provide recommendations on the use of analytical methods for their

subsequent application. Therefore, it seems reasonable to adopt a criterion that is commonly

used in the practice of ecological and conservation research.

After classifying variables as 'significant' or not, we compared them to the variables that were

actually used to generate the virtual data. We then calculated the following measures:

1) Accuracy (correct classification rate) - the probability of detecting an effect if there is

one or detecting no effect if there is none, i.e., whether the model correctly classified

the variable as included in the set used to generate virtual data or not.

2) True positive rate (sensitivity) – the power of a test, that is the probability of detecting

an effect if there is one. More specifically, in this study it is the probability that a

variable classified as significant in the model was actually included in the set used to

generate the virtual data.

3) True negative rate (specificity) – the probability of detecting no effect when there is

none, i.e., the probability that a variable classified as non-significant in the model was

actually excluded from the variable set used to generate the virtual data.

A binomial Generalised Linear Mixed Model using the R package "glmmTMB" (Brooks et al.,

2017) was used to quantitatively assess the effect of method, variable selection strategy and

their interaction on the above measures of classification performance. Species was entered

into the model as a random intercept. Estimated marginal means were used to assess

contrasts using the R package "emmeans" (Searle et al., 1980). The analyses were performed

using R 4.5.0 software (R Core Team, 2025). The full code is available on the GitHub

repository: https://anonymous.4open.science/r/Causes-2EB1.

3. Results

3.1 Accuracy

TS performed most poorly of the approaches tested. Its accuracy was 0.50 (95% confidence

interval: 0.48-0.53) when no variable selection was applied and 0.55 (CI: 0.52-0.57) when

variable selection was used (Fig. 2, Table S3), which improved accuracy (odds ratio: 0.83,

p<0.0001) (Table S4). SDMs performed better: the mean accuracy was 0.68 (CI: 0.66–0.70)

without variable selection and 0.63 (CI: 0.61-0.65) when it was applied. Variable selection

reduced the accuracy in this case (odds ratio: 1.24, p<0.0001). SDM was significantly more

accurate than TS under both variable selection strategies (odds ratio: 0.47, p<0.0001 and

0.71, p<0.0001) (Table S5). The POM approach outperformed both TS and SDM, its accuracy

was 0.88 (CI: 0.87-0.89), or 0.77 (CI:0.76-0.79) depending of the variable selection strategy

(odds ratio: 2.22, p<.0001). POM was significantly more accurate than both SDM (odds ratio:

0.28, p<0.0001 and 0.50, p<0.0001), and TS (odds ratio: 0.13, p<0.0001 and 0.36, p<0.0001).

3.2 Sensitivity

Sensitivity (true positive rate) was zero for the TS approach with no selection (in all

simulations none of the variables that actually influenced population abundance were

selected as significant) and 0.14 (CI: 0.11-0.17) when variable selection was used (Table S3).

For the SDM approach, it was 0.68 (CI: 0.62-0.73) and 0.73 (CI: 0.68-0.78), respectively, with

variable selection improving sensitivity (odds ratio: 0.77, p<0.0001) (Table S7). In the case

of POM, the sensitivity was the highest of the methods tested: 0.84 (CI: 0.81-0.88), and there

was no difference regarding the variable selection strategy (odds ratio: 1.0, p=0.79). POM

outperformed SDM in both cases (odds ratio: 0.38, p<0.0001 and 0.50, p<0.0001) and TS

when the variable selection was applied (odds ratio: 0.03, p<0.0001) (Table S8).

3.3 Specificity

The specificity (true negative rate) for the TS approach was 1.00 (in all simulations the

variables identified as non-significant were correctly classified as such) in the case of no

variable selection, and 0.95 (CI: 0.95-0.96) in the case when it was used (Table S3). For the

SDM, these estimates were 0.69 (CI: 0.66-0.72) and 0.54 (CI: 0.51-0.57), respectively, and

variable selection reduced the specificity (odds ratio: 2.00, p<0.0001) (Table S10). For the

POM approach, the specificity was 0.93 (CI: 0.92-0.94) and 0.71 (CI: 0.68-0.74), depending on

the selection strategy. The difference between these estimates was significant, the use of

variable selection noticeably produced poorer results (odds ratio: 5.00, p<0.0001). POM

outperformed both SDM in terms of specificity, regardless of selection strategy (odds ratio

for both contrasts: 0.00, p<0.0001), and TS when variable selection was applied (odds ratio:

8.0, p<0.0001) (Table S11).

Figure 2. Performance measures for the classification of variables using the following methods: TS - time series, SDM - species distribution model, POM – process-oriented model, with different variable selection strategies: blue indicates no variable selection, whereas yellow indicates that variable selection was applied. Box plots show the distribution of classification measures over 1000 replications. Whiskers represent 95% confidence intervals, lower and upper hinges are 50% confidence intervals and the middle line is the mean. The baseline at 1/2 indicates expected value if the variables were classified randomly.

4. Discussion

This study compares two analytical protocols for inferring causes of population changes with

a process-oriented approach. Our findings demonstrate that time series data are unable to

detect factors influencing population abundance. In the simulations carried out, all variables

were classified as non-significant, regardless of their actual impact. SDM performed

significantly better than TS across all performance measures and achieved satisfactory

accuracy (0.68) and balanced sensitivity and specificity. Finally, the most complex approach,

POM showed the best performance overall, with the highest accuracy (0.88) and well-

balanced sensitivity and specificity.

The choice of modelling method, although constrained by data availability and analytical

skills, depends first and primarily on the research question posed. The consequences of this

choice are generally underestimated and can drastically affect the results obtained and the

conclusions drawn. For example, Gould et al. (2025) show, that a single research problem,

when analysed using different approaches, can provide a different set of answers. When

examining the factors influencing population dynamics the choice of analytical approach also

depends on the nature of the data and the research hypotheses. For instance, when working

with demographic data, time series models may be a better tool to capture fluctuations in

apparent survival, as this parameter shows more variation across years than between sites.

Conversely, species distribution models may be better suited to analysing productivity,

which tends to vary more spatially than temporally (Morrison et al., 2022).

4.1 Efficiency of time series and species distribution models

Our results demonstrate that although these two popular methods are often used to identify

the drivers of population change, they do not always provide satisfactory results and may be

of limited use in conservation decision making. The main reason for their poor performance

appears to be data aggregation, which fundamentally alters the statistical properties of the

data, reducing variance and decreasing degrees of freedom. As a result, the signal-to-noise

ratio is lowered, reducing the validity of statistical inference and introducing the bias of

variable selection. This affects both sensitivity (true positive rate) and specificity (true

negative rate) in identifying relevant predictors. The higher the level of data aggregation, the

greater the loss of granularity and the greater the attenuation of effect sizes - this explains

the poor performance of TS, slightly better for SDM and quite satisfactory for POM, where the

original fine-scaled patterns are retained. Data aggregation could lead to biased results

across a wide range of statistical models (Abrahamowicz et al., 2004; Pollack et al., 2022;

Pollet et al., 2015; Schad et al., 2024), and time series tend to demonstrate a weaker covariate

effect if the dataset has been previously averaged over inadequate time scales (Ferguson et

al., 2017).

4.2 Properties of process-oriented models

The strong performance of the process-oriented models is most likely due not only to the use

of spatio-temporal data, but also to the structure of the model itself. Including at least some

elements of the data-generating mechanism, such as density dependence, autocorrelation,

random effects or observer effects, is likely to significantly improve the ability to correctly

identify relevant factors and reject spurious ones. Similar findings are recognised in the

literature (Johnson et al., 2024). Ignoring the hierarchical structure of the data can lead to

underestimation of population trends, and complex models provide more accurate

inferences about ecological systems (Johnson et al., 2024). It is acknowledged that dynamic

and process-orientated models are highly promising tools in ecology for a better and more

effective understanding of population dynamics, e.g., in the light of climate change (Franklin,

2010; Zurell, 2017), especially considering that static models, such as classic SDM, are

characterized by a number of drawbacks (Guisan & Thuiller, 2005).

However, POMs are subject to certain limitations. The increasing complexity of a model can

have a negative impact on its feasibility and tractability, which affects its subsequent

understanding. Furthermore, spatio-temporal models require large amounts of data. In

addition to data availability requirements, higher complexity can lead to higher

computational costs, including longer simulation times and more time spent on data

preparation and analysis. Whether this approach will always be more profitable than less

advanced models will therefore depend on the specific problem being addressed and the data

available.

4.4 Usability of variable selection

The results of our study contribute to the ongoing debate about the validity of variable

selection strategies (Tredennick et al., 2021; Whittingham et al., 2006). We have shown that

in the case of TS, variable selection improves accuracy; however, the difference is without

practical importance due to the method's inherent flaws. Furthermore, in our study variable

selection actually reduces the efficiency of more advanced approaches, such as SDM and POM.

Many variable selection strategies are now used in ecology, although each method, such as

the double shrinkage penalty used here, may have potential weaknesses that limit its

usefulness. Alternative approaches can therefore be used, or the full model can be fitted to

obtain the best outcomes (Tredennick et al., 2021; Whittingham et al., 2006).

4.4 Practical implications

The results of this study also have practical implications for conservation ecology and management practice in terms of population and ecosystem monitoring. We emphasise the need to collect fine-grained, long-term and large-scale monitoring data, such as that from the Pan-European Common Bird Monitoring Scheme (Brlík et al., 2021). This enables the use of complex and effective analytical methods, which are crucial for identifying reliable biodiversity indicators for monitoring global change (Clavero et al., 2011).

4.5 Overall conclusions

In conclusion, our study provides a comparison of methods frequently used to infer the drivers of population change, and we acknowledge that these do not exhaust all the possibilities available. However, our aim was not to provide a detailed comparison of multiple methods or different selection strategies, but rather to show that the commonly used analytical approaches applied to the problem of identifying the causes of population change may not be very effective and may lead to erroneous conclusions. Instead, we offer an easy-to-implement method that not only accounts for ecological mechanisms (such as density dependence or the observational process), but is also much more effective at correctly identifying the factors driving population change. We hope that this will contribute to a more effective use of large-scale and long-term monitoring data in a time of biodiversity crisis.

References

Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958-2015. *Scientific Data*, *5*, 170191. https://doi.org/10.1038/sdata.2017.191

Abrahamowicz, M., Du Berger, R., Krewski, D., Burnett, R., Bartlett, G., Tamblyn, R. M., & Leffondré, K. (2004). Bias due to aggregation of individual covariates in the Cox regression model. *American Journal of Epidemiology*, *160*(7), 696–706. https://doi.org/10.1093/aje/kwh266

Allen, A. M., Jongejans, E., van de Pol, M., Ens, B. J., Frauendorf, M., van der Sluijs, M., & de Kroon, H. (2022). The demographic causes of population change vary across four decades in a long-lived shorebird. *Ecology*, *103*(4). https://doi.org/10.1002/ecy.3615

Araújo, M. B., & Guisan, A. (2006). Five (or so) challenges for species distribution modelling. *Journal of Biogeography*, *33*(10), 1677–1688. https://doi.org/10.1111/j.1365-2699.2006.01584.x

Briscoe, N. J., Elith, J., Salguero-Gómez, R., Lahoz-Monfort, J. J., Camac, J. S., Giljohann, K. M., Holden, M. H., Hradsky, B. A., Kearney, M. R., McMahon, S. M., Phillips, B. L., Regan, T. J., Rhodes, J. R., Vesk, P. A., Wintle, B. A., Yen, J. D. L., & Guillera-Arroita, G. (2019). Forecasting species range dynamics with process-explicit models: matching methods to applications. *Ecology Letters*, *22*(11), 1940–1956. https://doi.org/10.1111/ele.13348

Brlík, V., Šilarová, E., Škorpilová, J., Alonso, H., Anton, M., Aunins, A., Benkö, Z., Biver, G., Busch, M., Chodkiewicz, T., Chylarecki, P., Coombes, D., de Carli, E., del Moral, J. C., Derouaux, A., Escandell, V., Eskildsen, D. P., Fontaine, B., Foppen, R. P. B., … Klvaňová, A. (2021). Long-term and large-scale multispecies dataset tracking population changes of common European breeding birds. *Scientific Data*, *8*(1). https://doi.org/10.1038/s41597-021-00804-2

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, *9*(2), 378–400. https://doi.org/10.32614/RJ-2017-066

Buckland, S. T., Anderson, D., Burnham, K., Laake, J., Borchers, D., & Thomas, L. (2001). *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford University Press.

Buckland, S. T., Anderson, D., Burnham, K., Laake, J., Borchers, D., & Thomas, L. (2004).
Advanced Distance Sampling. Oxford University Press.

Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing sixth
mass extinction signaled by vertebrate population losses and declines. *Proceedings of*
the National Academy of Sciences of the United States of America, *114*(30), E6089–E6096.
https://doi.org/10.1073/pnas.1704949114

Chiquet, R. A., Ma, B., Ackleh, A. S., Pal, N., & Sidorovskaia, N. (2013). Demographic analysis of
sperm whales using matrix population models. *Ecological Modelling*, *248*, 71–79.
https://doi.org/10.1016/j.ecolmodel.2012.09.023

Clavero, M., Villero, D., & Brotons, L. (2011). Climate change or land use dynamics: Do we
know what climate change indicators indicate? *PLoS ONE*, *6*(4).
https://doi.org/10.1371/journal.pone.0018581

Defourny, P., Lamarche, C., Bontemps, S., De Maet, T., Van Bogaert, E., Moreau, I., Brockmann,
C., Boettcher, M., Kirches, G., Wevers, J., Santoro, M., Ramoino, F., & Arino, O. (2017). *Land*
Cover Climate Change Initiative - Product User Guide v2. Issue 2.0.

Defourny, P., Lamarche, C., Marissiaux, Q., Brockmann, C., Boettcher, M., & Kirches, G. (2021).
Product User Guide and Specification ICDR Land Cover 2016-2020. Tech. Rep.

Di Fonzo, M. M. I., Collen, B., Chauvenet, A. L. M., & Mace, G. M. (2016). Patterns of mammalian
population decline inform conservation action. *Journal of Applied Ecology*, *53*(4), 1046–
1054. https://doi.org/10.1111/1365-2664.12659

Elith, J., & Leathwick, J. R. (2009). Species distribution models: Ecological explanation and
prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*,
40, 677–697. https://doi.org/10.1146/annurev.ecolsys.110308.120159

Enneson, J. J., & Litzgus, J. D. (2008). Using long-term data and a stage-classified matrix to
assess conservation strategies for an endangered turtle (Clemmys guttata). *Biological*
Conservation, *141*(6), 1560–1568. https://doi.org/10.1016/j.biocon.2008.04.001

Ferguson, J. M., Reichert, B. E., Fletcher, R. J., & Jager, H. I. (2017). Detecting population–
environmental interactions with mismatched time series data. *Ecology*, *98*(11), 2813–
2822. https://doi.org/10.1002/ecy.1966

Franklin, J. (2010). Moving beyond static species distribution models in support of
conservation biogeography. *Diversity and Distributions*, *16*(3), 321–330.
https://doi.org/10.1111/j.1472-4642.2010.00641.x

Gould, E., Fraser, H. S., Parker, T. H., Nakagawa, S., Griffith, S. C., Vesk, P. A., Fidler, F., Hamilton, D. G., Abbey-Lee, R. N., Abbott, J. K., Aguirre, L. A., Alcaraz, C., Aloni, I., Altschul, D., Arekar, K., Atkins, J. W., Atkinson, J., Baker, C. M., Barrett, M., … Zitomer, R. A. (2025). Same data, different analysts: variation in effect sizes due to analytical decisions in ecology and evolutionary biology. *BMC Biology*, *23*(1), 35. https://doi.org/10.1186/s12915-024-02101-x

Green, R. E. (1995). Diagnosing causes of bird population declines. *Ibis*, *137*, S47–S55. https://doi.org/10.1111/j.1474-919X.1995.tb08457.x

Gregory, R. D., Bashford, R. I., Balmer, D. E., Marchant, J. H., Wilson, A. M., & Baillie, S. R. (1996). The Breeding Bird Survey Report 1994-1995. In *British Trust for Ornithology, Joint Nature Conservation Committee and Royal Society for the Protection of Birds*.

Guisan, A., & Thuiller, W. (2005). Predicting species distribution: Offering more than simple habitat models. *Ecology Letters*, *8*(9), 993–1009. https://doi.org/10.1111/j.1461-0248.2005.00792.x

Holt, R. D. (2009). Bringing the Hutchinsonian niche into the 21st century: Ecological and evolutionary perspectives. *Proceedings of the National Academy of Sciences*, *106*(Suppl 2), 19659–19665. https://doi.org/10.1073/pnas.0905137106

Hsieh, C. H., Glaser, S. M., Lucas, A. J., & Sugihara, G. (2005). Distinguishing random environmental fluctuations from ecological catastrophes for the North Pacific Ocean. *Nature*, *435*(7040), 336–340. https://doi.org/10.1038/nature03553

Hutchings, J. A. (2000). Collapse and recovery of marine fishes. *Nature*, *406*(6798), 882–885. https://doi.org/10.1038/35022565

Hutchings, J. A., & Reynolds, J. D. (2004). Marine Fish Population Collapses: Consequences for Recovery and Extinction Risk. *BioScience*, *54*(4), 297–309. https://doi.org/10.1641/0006-3568(2004)054[0297:MFPCCF]2.0.CO;2

Johnson, H. E., Mills, L. S., Stephenson, T. R., & Wehausen, J. D. (2010). Population-specific vital rate contributions influence management of an endangered ungulate. *Ecological Applications*, *20*(6), 1753–1765. https://doi.org/10.1890/09-1107.1

Johnson, T. F., Beckerman, A. P., Childs, D. Z., Webb, T. J., Evans, K. L., Griffiths, C. A., Capdevila, P., Clements, C. F., Besson, M., Gregory, R. D., Thomas, G. H., Delmas, E., & Freckleton, R. P. (2024). Revealing uncertainty in the status of biodiversity change. *Nature*, *628*(8009), 788–794. https://doi.org/10.1038/s41586-024-07236-z

Knape, J., & de Valpine, P. (2011). Effects of weather and climate on the dynamics of animal
population time series. *Proceedings of the Royal Society B*, *278*(1708), 985–992.
https://doi.org/10.1098/rspb.2010.1333

Knowler, D. (2005). Reassessing the costs of biological invasion: Mnemiopsis leidyi in the
Black sea. *Ecological Economics*, *52*(2), 187–199.
https://doi.org/10.1016/j.ecolecon.2004.06.013

Kölzsch, A., Sæther, S. A., Gustafsson, H., Fiske, P., Höglund, J., & Kålås, J. A. (2007). Population
fluctuations and regulation in great snipe: A time-series analysis. *Journal of Animal*
Ecology, *76*(4), 740–749. https://doi.org/10.1111/j.1365-2656.2007.01246.x

Malinowska, K., Markowska, K., & Kuczyński, L. (2023). Making virtual species less virtual by
reverse engineering of spatiotemporal ecological models. *Methods in Ecology and*
Evolution, *14*(9), 2376–2389. https://doi.org/10.1111/2041-210X.14176

Marra, G., & Wood, S. N. (2011). Practical variable selection for generalized additive models.
Computational Statistics & Data Analysis, *55*(7), 2372–2387.
https://doi.org/10.1016/J.CSDA.2011.02.004

Morrison, C. A., Butler, S. J., Clark, J. A., Arizaga, J., Baltà, O., Cepák, J., Nebot, A. L., Piha, M.,
Thorup, K., Wenninger, T., Robinson, R. A., & Gill, J. A. (2022). Demographic variation in
space and time: implications for conservation targeting. *Royal Society Open Science*, *9*(3).
https://doi.org/10.1098/rsos.211671

Nentwig, W., Bacher, S., Kumschick, S., Pyšek, P., & Vilà, M. (2018). More than "100 worst"
alien species in Europe. *Biological Invasions*, *20*(6), 1611–1621.
https://doi.org/10.1007/s10530-017-1651-6

Normand, S., Zimmermann, N. E., Schurr, F. M., & Lischke, H. (2014). Demography as the basis
for understanding and predicting range dynamics. *Ecography*, *37*(12), 1149–1154.
https://doi.org/10.1111/ecog.01490

Pagel, J., & Schurr, F. M. (2012). Forecasting species ranges by statistical estimation of
ecological niches and spatial population dynamics. *Global Ecology and Biogeography*,
21(2), 293–304. https://doi.org/10.1111/j.1466-8238.2011.00663.x

Perrings, C., Williamson, M., Barbier, E. B., Delfino, D., Dalmazzone, S., Shogren, J., Simmons,
P., & Watkinson, A. (2002). Biological Invasion Risks and the Public Good: an Economic
Perspective. *Ecology*, *6*(1). https://doi.org/10.5751/ES-00396-060101

Pollack, A. B., Sue Wing, I., & Nolte, C. (2022). Aggregation bias and its drivers in large-scale
flood loss estimation: A Massachusetts case study. *Journal of Flood Risk Management*,
15(4). https://doi.org/10.1111/jfr3.12851

Pollet, T. V., Stulp, G., Henzi, S. P., & Barrett, L. (2015). Taking the aggravation out of data
aggregation: A conceptual guide to dealing with statistical issues related to the pooling
of individual-level observational data. *American Journal of Primatology*, *77*(7), 727–740.
https://doi.org/10.1002/ajp.22405

R Core Team. (2025). *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing. https://www.R-project.org/

Rhodes, J. R., Ng, C. F., de Villiers, D. L., Preece, H. J., McAlpine, C. A., & Possingham, H. P. (2011).
Using integrated population modelling to quantify the implications of multiple
threatening processes for a rapidly declining population. *Biological Conservation*,
144(3), 1081–1088. https://doi.org/10.1016/j.biocon.2010.12.027

Rigal, S., Dakos, V., Alonso, H., Auniņš, A., Benkő, Z., Brotons, L., Chodkiewicz, T., Chylarecki,
P., de Carli, E., del Moral, J. C., Domşa, C., Escandell, V., Fontaine, B., Foppen, R., Gregory,
R., Harris, S., Herrando, S., Husby, M., Ieronymidou, C., … Devictor, V. (2023). Farmland
practices are driving bird population decline across Europe. *Proceedings of the National*
Academy   of   Sciences,   *120*(21),   e2216573120.
https://doi.org/10.1073/pnas.2216573120

Schad, D. J., Nicenboim, B., & Vasishth, S. (2024). Data Aggregation Can Lead to Biased
Inferences in Bayesian Linear Mixed Models and Bayesian Analysis of Variance.
Psychological Methods. https://doi.org/10.1037/met0000621

Searle, S. R., Speed, F. M., & Milliken, G. A. (1980). Population marginal means in the linear
model: An alternative to least squares means. *American Statistician*, *34*(4), 216–221.
https://doi.org/10.1080/00031305.1980.10483031

Shumway, R. H., & Stoffer, D. S. (2005). *Time Series Analysis and Its Applications. With R*
Examples. Springer.

Soberón, J., & Nakamura, M. (2009). Niches and distributional areas: Concepts, methods, and
assumptions. *Proceedings of the National Academy of Sciences*, *106*(Suppl. 2), 19644–
19650. https://doi.org/10.1073/pnas.0901637106

Szymkowiak, J. (2013). Facing Uncertainty: How Small Songbirds Acquire and Use Social
Information in Habitat Selection Process? *Springer Science Reviews*, *1*, 115–131.
https://doi.org/10.1007/s40362-013-0012-9

Tredennick, A. T., Hooker, G., Ellner, S. P., & Adler, P. B. (2021). A practical guide to selecting
models for exploration, inference, and prediction in ecology. *Ecology*, *102*(6).
https://doi.org/10.1002/ecy.3336

Wauchope, H. S., Amano, T., Geldmann, J., Johnston, A., Simmons, B. I., Sutherland, W. J., &
Jones, J. P. G. (2021). Evaluating Impact Using Time-Series Data. *Trends in Ecology and*
Evolution, *36*(3), 196–205. https://doi.org/10.1016/j.tree.2020.11.001

Whittingham, M. J., Stephens, P. A., Bradbury, R. B., & Freckleton, R. P. (2006). Why do we still
use stepwise modelling in ecology and behaviour? *Journal of Animal Ecology*, *75*(5),
1182–1189. https://doi.org/10.1111/j.1365-2656.2006.01141.x

Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C. F.,
Forchhammer, M. C., Grytnes, J. A., Guisan, A., Heikkinen, R. K., Høye, T. T., Kühn, I., Luoto,
M., Maiorano, L., Nilsson, M. C., Normand, S., Öckinger, E., Schmidt, N. M., … Svenning, J.
C. (2013). The role of biotic interactions in shaping distributions and realised
assemblages of species: Implications for species distribution modelling. *Biological*
Reviews, *88*(1), 15–30. https://doi.org/10.1111/j.1469-185X.2012.00235.x

Wood, S. N. (2017). *Generalized Additive Models: An Introduction with R* (2nd ed.). Chapman
and Hall/CRC. https://doi.org/10.1201/9781315370279

Zachariah Peery, M., Beissinger, S. R., Newman, S. H., Burkett, E. B., & Williams, T. D. (2004).
Applying the Declining Population Paradigm: Diagnosing Causes of Poor Reproduction
in the Marbled Murrelet. *Conservation Biology*, *18*(4), 1088–1098.
https://doi.org/10.1111/j.1523-1739.2004.00134.x

Zurell, D. (2017). Integrating demography, dispersal and interspecific interactions into bird
distribution models. *Journal of Avian Biology*, *48*(12), 1505–1516.
https://doi.org/10.1111/jav.01225

Zurell, D., Franklin, J., König, C., Bouchet, P. J., Dormann, C. F., Elith, J., Fandos, G., Feng, X.,
Guillera-Arroita, G., Guisan, A., Lahoz-Monfort, J. J., Leitão, P. J., Park, D. S., Peterson, A. T.,
Rapacciuolo, G., Schmatz, D. R., Schröder, B., Serra-Diaz, J. M., Thuiller, W., … Merow, C.
(2020). A standard protocol for reporting species distribution models. *Ecography*, *43*(9),
1261–1277. https://doi.org/10.1111/ecog.04960

Zurell, D., Jeltsch, F., Dormann, C. F., & Schröder, B. (2009). Static species distribution models
in dynamically changing systems: how good can predictions really be? *Ecography*, *32*(5),
733–744. https://doi.org/10.1111/j.1600-0587.2009.05810.x

Zurell, D., Zimmermann, N. E., & Brun, P. (2024). The niche through time: Considering
phenology and demographic stages in plant distribution models. In *Journal of Ecology*
(Vol. 112, Issue 9, pp. 1926–1939). John Wiley and Sons Inc.
https://doi.org/10.1111/1365-2745.14361

Supporting Information

Additional supporting information can be found online in the Supporting Information

section at the end of this article.

Supplementary File 1. Specific details concerning data used in the analysis, data treatment

and variable selection.

Supplementary File 2. Additional results not included in the main article.