1 **Sea-ice microbial community succession and the potential role of parasitoids in the maintenance of**
2 **diversity during the spring bloom**

3 Kyle B. Dilliplaine[1,2]*, Laura M. Whitmore[1,3], Ana Aguilar-Islas[1], Channing Bolt[1,4], Kenneth
4 Dumack[2], Meibing Jin[3], Mette Kaufman[3], Marc Oggier[3], Gwenn M. M. Hennon[1]
5
6 [1]College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Fairbanks, AK, USA
7 [2]Aquatic Ecosystem Analyses, Institute for Integrated Natural Sciences, University of Koblenz,
8 Koblenz, Germany
9 [3]International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, AK, USA
10 [4]Naval Information Warfare Center, Pacific, San Diego, CA, USA
11
12 ***Corresponding author:** kbdilliplaine@alaska.edu
13
14 **Email addresses:** lmwhitmore@alaska.edu; amaguilarislas@alaska.edu;
15 channing.e.bolt2.civ@us.navy.mil; kenneth.dumack@uni-koblenz.de; mjin@alaska.edu;
16 mrkaufman@alaska.edu; moggier@alaska.edu; gmhennon@alaska.edu
17

Acknowledgements: The authors thank the Ukpeaġvik Iñupiat Corporation Science (UICS) for
their logistical support in Utqiaġvik during field work. We thank Hajo Eicken and Rob Rember
who were instrumental to securing funding and planning the field campaign, Anika Pinzer for
assistance with map making, and Drs. Rebecca Duncan and Jozef Wiktor for their expertise
identifying diatoms. This project was supported by research grants funded through the National
Science Foundation (NSF) OPP-1735862 (AAI) and OCE-1937715 (GMH) and the Coastal
Marine Institute M20AC10007 (GMH), a joint institute between the Bureau of Ocean Energy
Management (BOEM) and the University of Alaska Fairbanks. Kenneth Dumack was supported
by the German Research Foundation (DFG) with the grant number 555596351. Micrographs
were captured at the UAF Molecular Imaging Facility, National Institute of General Medical
Sciences (NIGMS) P20GM130443, and sequencing conducted at the UAF Genomic's Core Lab,
NIGMS P20GM103395. We would like to thank our three peer reviewers who volunteered their
time and expertise to improve this manuscript.

Conflict of Interest: The authors declare no conflict of interest.

Data Availability: Biogeochemical data are available through the Arctic Data Center,
doi:10.18739/A21J9793S (Whitmore et al., 2024), and DNA sequencing data are available through
NCBI's SRA under BioProject PRJNA1108783. Taxonomically annotated Raw read counts are
available as a supplementary dataset Supplementary Information (Table S1 & S2). Data
processing and analysis R scripts are publicly available on GitHub at
https://github.com/kbdilliplaine/2021-Utqiagvik-time-series.git.

Author Contributions (CRediT):
Conceptualization: KBD and GMH contributed to the conceptualization of overarching
research goals and aims.
Data curation: LW, KBD and MO contributed to data curation.
Formal analysis: KBD and LW contributed to formal analysis.
Funding acquisition: AAI acquired funding to support the project leading to this publication.
GMH acquired funding to support sample processing, labor, and publication costs.

Investigation: LW, CB, MK and MO conducted field work and processed samples for physical
and chemical properties. KBD processed samples for molecular analysis.
Methodology: AAI, MO, LW and KBD contributed to methodological development.
Project administration: GMH and AAI managed different aspects of this project and its
execution.
Resources: AAI, MK and GMH contributed resources necessary for the completion of this
study.
Visualization: KBD was responsible for the creation of all visual components of this manuscript.
Writing – original draft: KBD was responsible for the creation of the initial draft.
Writing – review & editing: GMH, LW, KD, MJ and AAI provided substantial critical review and
commentary of this manuscript.

Abstract

Sea ice is a crucial, yet declining, habitat in high latitude ecosystems. Here we present a high-temporal resolution amplicon sequence data set collected during the spring ice-algal bloom near Utqiaġvik, Alaska in 2021 to study sea-ice microbial dynamics. The ice-algal bloom peaked on May 8th, reaching 46.6 mg chlorophyll *a* m^{-2}, and thereafter became limited by nitrate availability. A massive bloom of the oil-degrading bacterium, *Oleispira* (>80% relative abundance), coincided with the algal bloom raising questions about hydrocarbon exposure. The sea-ice algal bloom was dominated by diatoms, particularly *Nitzschia spp.*, and transitioned into a flagellate dominated post-bloom community which aligned with melt-associated changes to the physicochemical environment. We explored the relationship between putative parasitoids, Chytridiomycetes, Thecofilosea (Cercozoa), Oomycetes, Syndiniales (Dinoflagellata), and Labyrinthulomycetes (Bigyra), and potential microalgal hosts. Chytrids peaked periodically suggesting synchronized infections and *Cryothecomonas* (Thecofilosea) was observed parasitizing *Nitzschia spp.* for the first time in Arctic sea ice. Co-occurrence analysis suggested that diatoms, especially *Nitzschia*, were the primary hosts of Pacific-Arctic parasitoids, and that top-down parasitoid control may dramatically alter community composition over short timescales, such as days. These results provide important insights into the drivers of spring bloom timing and maintenance of microalgal diversity in sea ice.

Keywords: sea-ice microalgae, parasitoids, *Oleispira*, *Cryothecomonas*, chytrids, landfast sea ice, Utqiaġvik, AK

1. Introduction

Climate change is rapidly impacting the Arctic sea-ice system by shifting the timing of freezeup and melt (Walsh et al., 2022), precipitation patterns (Bintanja and Selten, 2014; Webster et al., 2014), and ice extent and thickness (Babb et al., 2022; Stroeve and Notz, 2018). Sea-ice associated (sympagic) microalgae contribute substantially to Arctic primary production(Gosselin et al., 1997; Mortenson et al., 2020). Ice-algal production is primarily controlled by the availability of light, which is determined by ice and snow thickness (Mcdonald et al., 2015; Veyssière et al., 2022), and inorganic nutrients supplied via underlying water (Cota et al., 1987; Cota and Smith, 1991; Dalman et al., 2019). Sympagic communities have been observed to shift toward heterotrophic protists and dinoflagellates during the spring to summer transition (Rózańska et al., 2009; von Quillfeldt et al., 2003). Irradiance levels likely contribute to species-specific succession and community composition due to differing photophysiology, e.g., centric diatoms (such as *Thalassiosira* spp*.),* dinoflagellates, and certain pennate diatoms (such as *Navicula* spp.), increase with greater light availability while pennate diatoms like *Nitzschia frigida* decline (Croteau et al., 2022; Duncan et al., 2024). Nutrient availability can also influence microalgal community structure based on the resource utilization traits and stoichiometric requirements of different algal taxa (Litchman et al., 2015a, 2015b). Nitrogen is generally the limiting nutrient throughout most of the Arctic (Pineault et al., 2013; Rózańska et al., 2009), though silicic acid, which is required by diatoms, has been found to be limiting regionally (Giesbrecht and Varela, 2021; Gosselin and Legendre, 1990; Smith et al., 1990). Sympagic algae are positioned in the upper ocean, allowing them to take advantage of light early in the spring season, thus providing an early source of nutrition to the broader Arctic food web (Kohlbach et al., 2017). Sea ice melt processes release dense algal material which may be grazed by plankton (Juul-Pedersen et al., 2008), sink rapidly to the seafloor as aggregates (Ambrose et al., 2005; Boetius et al., 2013), or seed a successive phytoplankton bloom (Garrison et al., 1987; Lizotte, 2001; Yan et al., 2020). While much is known about bottom-up environmental control of sympagic microbial communities, top-down biological pressures may also affect community composition and are less well-understood.

The study of parasites in sea ice has been limited to small and focused investigations. While fungi such as chytrids (Hassett and Gradinger, 2016), and a newly discovered thraustochytrid (Labyrinthulomycetes), infect sea-ice algae (Hassett, 2020), Arctic Cryomonadida (Cercozoa, Thecofilosea), have thus far only been observed as phagotrophic grazers (Thaler and Lovejoy, 2012), though the commonly reported genus, *Cryothecomonas,* is currently the only known cryomonad genus to exhibit endoparasitic behavior (Drebes et al., 1996; Kühn et al., 2000). In contrast to parasites, parasitoids ultimately kill their host after infection (Skovgaard, 2014). Parasitoids are capable of terminating phytoplankton blooms and altering species succession (Chambouvet et al., 2008; Tillmann et al., 1999). Despite their potential ecological importance, marine parasitoids are understudied due to ambiguous morphologies that make identification difficult (Käse et al., 2021), though molecular surveys suggest high undescribed diversity (B. T. Hassett et al., 2019; Thaler and Lovejoy, 2012). Identifying potential parasitoid-host interactions is required to better understand bloom dynamics and biogeochemical cycles in sea ice where parasitoid infections of microalgae have the potential to modulate the transfer of carbon between trophic levels (Amundsen et al., 2009).

Despite the importance of sympagic microbes to the Arctic ecosystem, basic information regarding microbial community structure, dynamics, and microalgal bloom phenology is

spatiotemporally limited along the coastal US Arctic. We conducted a high-resolution time series
(April 21 to June 11, 2021, sampled every few days) to determine the timing, magnitude, and
yield-limiting factor of the ice-algal bloom at Utqiaġvik, Alaska, formerly known as Barrow. We
examined the prokaryotic and unicellular eukaryotic community composition and dynamics over
the spring microalgal bloom using 16S and 18S rRNA amplicon sequencing, and leveraged
these data to identify putative parasitoid-host relationships within the sympagic community.

2. Methods

2.1. *Study site and ice sampling*

Field collections were conducted near Utqiaġvik, Alaska (71.375 N, 156.537 W) between
April 21 and June 11, 2021 (Day of Year (DOY) 111 — 162), on a large level pan of landfast sea
ice, measuring ~11 km^2 (Fig. S1). Additional site information and detailed methodology is
provided in Appendix 1. Sea ice cores were collected from a ~900 m^2 area, with each sampling
event constrained to 1 m^2. Ice core samples were collected approximately three times per week
and measurements of temperature, salinity, density, nutrients, chlorophyll *a* (chl *a*), and samples
for DNA were collected (seeAppendix 1 for more details). Cores were retrieved using a 9-cm
inner diameter Kovacs ice corer. Incident and under-ice photosynthetically active radiation
(PAR; 400-700 nm) photon flux densities were measured adjacent to the sampling area using a
LI-COR LI-1400 data logger with planar (incident; 2π) and spherical (under-ice; 4π) quantum
sensors (Lincoln, NE, USA), respectively, and a 4π:2π ratio was used to determine
"pseudotransmissivity" (see Appendix 1 for more details). Biological samples were obtained
from the bottom 10 cm of a single ice core by quickly removing the section with a cleaned
handsaw. The section was sealed in a clean plastic bag and stored in a cooler in the dark until
transport back to the lab. Samples were processed by adding 100 ml of 0.2 μm filtered seawater
cm^{-1} of ice to prevent osmotic shock (Garrison and Buck, 1986) and melted at 4 °C in the dark
overnight. Once entirely melted, 1-2 aliquots (12-171 mL) of each sample were vacuum filtered
($\leq$ 5 psi) for chl *a* using 25 mm glass fiber filters (Whatman GF/F). All duplicated chlorophyll
subsamples were averaged and reported as integrated areal chl *a* concentrations. The
remaining sample volume was processed for DNA by filtering the sample onto a 0.2 μm Sterivex
filter (Millipore Sigma) using a peristaltic pump and stored at -80 °C until further processing.

2.2. *DNA extraction, sequencing, and bioinformatic processing*

DNA was extracted using the NucleoMag DNA/RNA Water kit (Macherey Nagel, Düren, GE)
. The V4 region of both the 16S and 18S rRNA genes were amplified for each sample using the
revised Earth Microbiome Project primers (515FB and 806RB; Caporaso et al. 2012; Apprill et
al. 2015; Parada et al. 2016) and TAReuk454FWD1 and TAReukREV3 (Stoeck et al., 2010),
respectively. Primers were modified with TaggiMatrix indices to enable pooling of samples prior
to adaptation with Illumina sequencing adapters (Glenn et al., 2019). Samples were sequenced
on an Illumina MiSeq with a 2 x 300 paired-end kit at the University of Alaska Fairbanks'
Genomics Core Laboratory. The QIIME2 (version 2023.5) workflow was used to process reads
before further analysis (Caporaso et al., 2010); pre-processing and detailed information can be
found in Appendix 1. Prokaryotic amplicon sequence variants (ASVs) were classified using a
weighted classifier (515f-806r-average-classifier.qza) trained on the Silva 138.1 database
(Bokulich et al., 2018; Kaehler et al., 2019; Quast et al., 2012; Robeson et al., 2021). A classifier
was trained on the PR2 database (version 5.0.0) for eukaryotic ASV classification (Guillou et al.,
2012). ASVs were identified to the lowest taxonomic level possible and ASVs of the same
taxonomic identity were numbered sequentially (Table S1). Taxonomic corrections were made to

the eukaryotic classifications due to classifier uncertainty and incomplete taxonomic
representation in the PR2 database (Table S1). To provide the most up-to-date taxonomic
assignment, each ASV involved with a significant parasitoid correlation was manually checked
via NCBI BLAST alignment. Additional details regarding sequence processing can be found in
Appendix 1.

2.3 Bioinformatic analysis

Community structuring across our time-series was investigated according to Walker et al.
(2023) for rarefied prokaryotic and eukaryotic ASV data. Community structure was visualized
using non-metric multidimensional scaling (nMDS) using Bray-Curtis dissimilarity matrices with
Wisconsin double standardization transformed ASV tables. Outliers were identified using
Mahalanobis distance using 'rstatix' (Kassambara, 2023); outliers were retained on the nMDS
plot but were excluded from cluster polygons. Cluster association indices were calculated for
each ASV to identify which were significantly associated with each of the clusters. The
'indicspecies' package was used to calculate the point biserial correlation coefficient on rarefied
data (Cáceres and Legendre, 2009), and corrected for unequal sample distribution between
clusters (Tichy and Chytry, 2006). Only taxa with $p < 0.001$, correlation coefficients >0.5, and
contributed at least 0.5% of all sequences (16S or 18S) were retained.

Correlations between putative parasitoids and phototrophic hosts were investigated using
Spearman's rank correlation of select ASVs using the 'psych' (Revelle, 2023; v.2.3.9) package in
R with false discovery rate corrections for p values (Benjamini et al., 2009). Unrarefied relative
abundance data were first centered log-ratio transformed using the 'SpiecEasi' package (v.1.1.3;
Kurtz et al., 2017), after removal of rare taxa (occurring as <0.1% in fewer than 50% of
samples). Parasitoids were limited to the classes Chytridiomycetes, Thecofilosea (Cercozoa),
Labyrinthulomycetes (Bigyra), Oomycetes and Syndiniales (Dinoflagellata) while hosts were
limited to the dominant phototrophs, i.e., diatoms (Bacillariophyceae, Mediophyceae, and
Coscinodiscophyceae) and dinoflagellates (Dinophyceae). To assess whether parasitoids are
associated with microalgal α-diversity (limited to the taxa defined above), we regressed the per-
sample Inverse Simpson index on the summed relative abundance of all parasitoid classes.
Correlation matrices were produced at the ASV level, retaining only those with at least a
moderate correlation strength, i.e. |rho| ≥ 0.5; $p < 0.05$). We sampled again in April of 2022 and
used microscopy to validate potential correlations and quantify *Cryothecomonas* infection
prevalence (see Appendix 1for more details).

3. Results

Snow depth ranged from 0 to 6 cm during the study period (Figs. 1c & S2b). Direct
measurements of under-ice photosynthetically active radiation (PAR) ranged from 2.34 – 77.50
μmol photons m^{-2} s^{-1} (Fig. 1c) which was 0.25 – 11.13% of incident PAR (Fig. S2). The air
temperature began warming on DOY 133 and reached above 0 °C on DOY 141 (Fig. S3a). The
temperature measured at 2.5 cm from the bottom of the ice warmed above -1.8 °C by DOY
134, indicating the initiation of bottom ice melt between DOY 132 and 134 (Fig. S3d; ice growth
< DOY 134). Ice thickness was relatively uniform with a thickness of 109 ± 2 cm (mean ± SD)
until melt onset, declining thereafter to a minimum thickness of 95 cm (Fig. S2c). The brine
salinity from the bottom 10 cm decreased substantially between DOYs 132 and 137 (Fig. S4c).
The peak chlorophyll biomass occurred on DOY 128 (May 8th), reaching a maximum of 46.6 mg
chl *a* m^{-2} (Fig. 1). The calculated accumulation and loss rates were 2.64 and -4.57 mg chl *a* m^{2}

d $^{-1}$, respectively. Areal chl a concentration declined below 1 mg m^{-2} on DOY 144 (May 24th);
bloom phases are therefore defined as "bloom" < DOY 144 and "post-bloom" ≥ DOY 144 (Fig.
1a). The average proportion of chlorophyll degradation product (phaeophytin) increased from
31% to 45% after the onset of bottom ice melt. Seawater nutrient concentrations peaked on the
date of bloom termination (DOY 144; Fig. 1b). Bulk nutrient concentrations in the bottom 10 cm
of ice were significantly correlated with chl a; nitrate (adj. r^2=0.45, p =0.0014), nitrite (adj.
r^2=0.17, p =0.0447), ammonium (adj. r^2=0.36, p =0.0038), phosphate (adj. r^2=0.63, p <0.0001)
and silicate (adj. r^2=0.41, p <0.0019; Fig. S5).

3.1.1. *Sympagic community composition patterns*

The total number of prokaryotic ASVs identified from the unrarefied data was 469, representing
124 families from 33 classes of bacteria and archaea. A relative abundance of approximately
48% of the sequence reads belonged to just two ASVs identified to the Genus level, *Oleispira*
sp. and *Paraglaciecola* sp., 30% and 17.5%, respectively. *Polaribacter* was most abundant early
in the time-series before transitioning to a bloom of *Oleispira* that approximately follows the
microalgal bloom (Fig. 2). *Tenacibaculum* and *Colwellia* genera were the most abundant taxa in
the post-bloom period (Fig. 2). The combined relative abundance of all four *Oleispira* ASVs
reached a peak of 84.4% on DOY 120 and was above 5% during a 35-d period (n=15); eight of
those samples were above 50% (Fig. S6a). A positive correlation was found between *Oleispira*
relative abundance and the areal chl a concentration (adj. r^2 = 0.68, p = <0.001; Fig. S6b).
Three prokaryotic community clusters were identified as unique using hierarchical clustering
analysis (adonis2: F = 8.59, p < 0.001), and was supported by a marginally insignificant within-
cluster dispersion (betadisper: p = 0.075). The average silhouette width of 0.47 supports the
separation of these clusters, referred to as clusters P1, P2, and P3 (Figs. 2a & S7a). Prokaryotic
taxa identified by indicator taxa analysis that were significantly associated with the clusters are
used as representatives for describing the major successional patterns and associated
metabolisms (Table S3). *Polaribacter* was the dominant genus associated during the early-
bloom (P1), preceding a bloom of *Oleispira* coincident with the microalgal bloom (P2), before
transitioning to a more even post-bloom (P3) community represented by associated members of
Colwellia, Marinomonas, and *Tenacibaculum*.

The total number of unicellular eukaryotic ASVs identified from the unrarefied data was 751,
representing 14 divisions. The relative abundance of the top five most abundant classes
contributed 75.4% of all sequences reads: Bacillariophyceae (26.5%), Dinophyceae (23.4%),
Thecofilosea (13.8%), Oomycetes (6.8%), and the Imbricatea (4.9%). A total of 119 diatom
ASVs contributed to 28% of the relative read abundance. The diatom genus *Nitzschia*
contributed most to the bloom-phase, before transitioning into a post-bloom phase with a large
contribution by the dinoflagellate genus *Heterocapsa* (Fig. 2d). Hierarchical clustering analysis
identified two unique eukaryotic clusters (adonis2: F = 8.72, p < 0.001). The dendrogram
indicated a substructure within these two major clusters (Fig. S7b). Within-cluster dispersion
was not significant (betadisper: p = 0.292), and the average silhouette width of 0.27 is generally
considered weak. We designate these two clusters as E1 (bloom) and E2 (post-bloom; Fig. 1c).
One outlier (DOY = 113) was identified using Mahalanobis distance and retained in analyses,
although excluded from the nMDS polygon (Fig. 2b). Eukaryotic association indices revealed
taxa significantly associated with each of the two clusters (Table S4). Four diatom, and one
flagellate, ASVs were associated with cluster E1, while seven flagellate ASVs were associated
with cluster E2.

3.1.2. *Parasitoid-host associations*

The putative parasitoids investigated in this study appear to follow successional co-exclusion during the sympagic algal bloom, seen by their relative abundance (Fig. 3). The Chytridiomycetes followed a rhythmic pattern of peaks and troughs, roughly following a single cycle approximately every seven days during the bloom period. The Thecofilosea had genera dependent development during the time series, *Cryothecomonas* peaked during the late-bloom period while *Protaspa* became most prevalent after *Cryothecomonas* declined. The unclassified Cryomonadida (Cercozoa, Thecofilosea), Labyrinthulomycetes and Syndiniales remained relatively low throughout the timeseries, while the relative abundance of Oomycetes dramatically increased during the post-bloom period. Inverse Simpson index showed a weak (r^2 = 0.07), insignificant (p = 0.2314) positive association with summed parasitoid counts (β = 0.15 ± 0.12 SE; Fig. S8). Directed correlations, i.e., parasitoid versus microalgal pairings, were investigated at the ASV level (Fig. 4). Of the 91 significant correlations, 75 were with ASVs of the Thecofilosea (62 of which were with diatoms). Using the combined positive and negative correlations as an indicator of host range, the thecofilosean parasitoids had the largest potential host range while the labyrinthulids were the most limited (Fig. 4). Special attention was given to the correlated ASV pairing of Lobulomycetales Order_1 with *Nitzschia* ASVs (Fig. 5). Subsequent microscope observations support at least some of the correlations that were found; to the best of our knowledge,we provide first-time evidence of internal parasitization of Arctic diatoms by *Cryothecomonas* (Fig. 6). In 2022, 5.1% ± 0.98% (mean ± SD) of *N. arctica* cells were found to be infected by *Cryothecomonas*.

4. Discussion

4.1. *Physicochemical environment and bloom dynamics*

The maximum ice thickness of undeformed sea ice at Utqiaġvik was approximately 0.5 m less than measurements from the early 2000's (Jin et al., 2006; Mahoney et al., 2007; Lee et al., 2008; Manes and Gradinger, 2009; Fig. S2). The maximum annual landfast ice thickness near this location has thinned at a rate of 10 cm decade^{-1}, over twice the rate of several locations in the Canadian Arctic Archipelago (Eicken et al., 2012; Howell et al., 2016; Osborne et al., 2018). Snow depth in 2021 was also thinner than previous years, which often reached ~20 cm (or greater in drifts; Herzfeld et al., 2006; Jin et al., 2006; Manes and Gradinger, 2009). Thinner snow and ice layers provided greater light transmittance than in past decades (Figs. 1c & S2). Pseudotransmitted irradiances remained below photoinhibitory thresholds until the post-bloom period (~50 µmol photons m^{-2} s^{-1}; Arrigo et al., 2010; Juhl and Krembs, 2010; Croteau et al., 2022; Dilliplaine and Hennon, 2023; Fig. S2e), though, excluding chlorophyll from attenuation calculations suggests photoinhibition may have occurred at the top of the dense ice algal layer (Fig. S2h). An overall brighter under-ice light field is likely to become more common as ice thins and less snow falls, which may influence community composition or the vertical distribution of species as they balance light and nutrient requirements and preferences (Aumack et al., 2014; Croteau et al., 2022; Lannuzel et al., 2020).

The underlying seawater nutrient concentration and flux are generally considered important factors controlling the magnitude of ice algal blooms (Cota et al., 1991; Dalman et al., 2019; Gradinger, 2009; Rózańska et al., 2009). During our study, seawater nutrient concentrations (Fig. 1b) and nutrient ratios (Fig. S9) were similar to previous studies in the region during the spring ice-algal bloom (Lee et al., 2008; Manes and Gradinger, 2009). Nutrients remained

replete in the underlying seawater, with resupply likely driven by unmeasured physical
processes such as turbulence at the ice-water interface. We excluded the use of sea-ice brine
nutrient data because it appeared that cell lysis during melting released intracellular nutrient
pools, evident by the strong positive correlation of each macronutrient analyte with chl *a* (Fig.
S5). Brine nutrient concentrations are expected to behave conservatively according to the
equations of Cox and Weeks (1983), though Roukaerts et al. (2021) presented a biofilm-
mediated hypothesis to explain the accumulation of nutrients with algae in the Southern Ocean.
However, recent studies show that sea-ice nutrient-chlorophyll correlations likely reflect the
release of intracellular nutrient pools during melting protocols (Mundy et al., 2025; Ahmed et al.
in review). The positive silicate-chlorophyll relationship in our study suggests that our ~24h
direct melt method released contents of the silicon deposition vesicles. The observed nutrient
ratios in underlying seawater were well below the expected 16:1 (N:P) from Redfield and greater
than 1:1 (Si:N; Fig. S9), indicating nitrate resupply became the limiter to maximum biomass
accumulation (Redfield et al., 1963) prior to the onset of melt. Using the empirical nitrate-
chlorophyll relationship from Rózańska et al. (2009) the average seawater nitrate concentration
(5.6 µmol L^{-1}) predicts a maximum chl *a* concentration of 44.9 mg m^2, closely matching our
maximum observed concentration of 46.6 mg m^{-2} (Fig. 1a).

This study is the first to document the near-complete growth, decay, and fully realized
magnitude of the sea-ice algal bloom at Utqiaġvik, AK, USA. The peak of the algal bloom
occurred on DOY 128 (May 8th) similar to the peak date (DOY 121; May 1st) derived from
regionally compiled data presented in Leu et al. (2015). The 2021 bloom magnitude exceeded
all previously reported values near our study site, but was similar to collated data for landfast ice
in Resolute Bay (Leu et al., 2015), and exceeds that of other landfast ice time-series such as
the Green Edge ice camps in Baffin Bay (Massicotte et al., 2020; ~30 mg m^{-2} 2015, ~7 mg m^{-2}
2016). The highest prior measurement at this location was 36.2 mg chl *a* m□², measured in late
May (Gradinger et al., 2009), while our peak was ~25% greater than this isolated observation,
and nearly 300% greater than the model estimate from compiled data (Leu et al., 2015). It
remains unclear whether this elevated biomass reflects environmental change (e.g., thinning ice
and snow, faster under-ice currents) or is due to higher temporal resolution. Sampling just one
week before or after the actual peak would have underestimated biomass by 26-38%,
respectively. If sea-ice algal biomass is increasing, it could lead to greater secondary
production. Bloom termination and succession were tightly linked to the onset of sea ice melt,
consistent with previous observations (Leu et al., 2015 and references therein; Oziel et al.,
2019). Bottom-ice melt coincided with the increase of phaeophytin proportion (Fig. 1a),
indicating the onset of unfavorable conditions such as freshening, nutrient depletion, grazing, or
infection.

4.2. *Sympagic community dynamics*

We observed distinct patterns of succession in eukaryotic and prokaryotic communities
associated with bloom stage (Fig. 2). The eukaryotic community gradually transitioned between
the bloom (E1) and post-bloom (E2) phases, potentially driven by brine drainage and export of
non-motile taxa. The E1 cluster was enriched in pennate diatoms and transitioned to flagellate
dominance in E2 (Fig. 2; Table S4), a well-established progression in Arctic sea ice (Alou-Font
et al., 2013; Rózańska et al., 2009; von Quillfeldt et al., 2003).

We observed three distinct prokaryotic community clusters, largely following the temporal
evolution of the bloom: early- (P1), mid- (P2), and post-bloom (P3; Figs. 2a). Such changes are

likely driven by shifts in dissolved organic matter availability and composition, as prokaryotic
communities respond to the dynamic concentration and quality of organic matter that evolve
over the course of microalgal bloom development(Zhang et al., 2018; Zhou et al., 2018). The P1
associated *Polaribacter*, and P3 associated *Tenacibaculum* and *Colwellia* ASVs, are
heterotrophic bacteria known for their utilization of microalgal-derived dissolved organic matter
(DOM; Landa et al., 2016; Underwood et al., 2019). *Colwellia* and *Tenacibaculum* are capable of
degrading high molecular weight DOM; additionally, *Tenacibaculum* can utilize complex
extracellular polymeric substances derived from diatom biofilms (Bohórquez et al., 2017;
Underwood et al., 2019). *Marinomonas,* which was also associated with the P3 cluster, has the
capacity to metabolize aromatics from terrestrial peat (Sipler et al., 2017). These results suggest
Polaribacter uses labile carbohydrates, while P3 members degrade recalcitrant high-molecular-
weight substrates remaining after labile material is consumed. *Paraglaciecola* and
Winogradsyella, which have been previously observed associated with sea-ice algal aggregates
(Rapp et al., 2018), were abundant throughout the time series and not associated with any of
the clusters. This may be due to their ability to use a diverse array of carbohydrates (Schultz-
Johansen et al., 2018; Sun et al., 2025), maintaining abundances as the quality and quantity of
algal substrates change.

During the mid-bloom (P2), the prokaryotic community was unexpectedly dominated by the
oil-degrading bacterium *Oleispira* (Fig. S6; Table S3). While many sympagic genera (e.g.,
Colwellia, Polaribacter, Nonlabens, Paraglaciecola) can degrade hydrocarbons, *Oleispira* is an
obligate hydrocarbonoclastic bacterium that rapidly blooms after petroleum exposure (Brakstad
et al., 2008; Lofthus et al., 2018; Netzer et al., 2018; Yakimov et al., 2003; Yang et al., 2016).
Relative abundances >5% typically indicate hydrocarbon input (Krolicka et al., 2019), and
although *Oleispira* exceeded 80% in some samples there was no visible contamination. This
enrichment may reflect advection from a distant spill and subsequent entrainment into bottom
ice via seawater flux and physical sieving (Spindler, 1994; Syvertsen, 1991).However, the close
association of hydrocarbonoclastic bacteria, such as *Oleispira*, with phytoplankton (Thompson
et al., 2020; this study), or decaying microalgae (Cono et al., 2022), and the correlation of
abundance with chl *a*, warrants further evaluation of *Oleispira* as an indicator taxon of fossil
hydrocarbon exposure, as these could be sources of biogenic hydrocarbons.

The eukaryotic cluster, E1, was associated with the diatom genera *Nitzschia* and *Navicula,*
which are among the most abundant algae in Arctic sea ice (Table S4; Leu et al., 2015). An
unidentified cryomonad (Cryomonadida Order_1) was also associated with E1; Cryomonadida
are commonly reported in Arctic sea ice and are generally considered heterotrophic (Thaler and
Lovejoy, 2012). Rhythmic fluctuations of *Nitzschia* relative abundance were seemingly unrelated
to environmental drivers. These fluctuations may be explained by spatial heterogeneity, or
parasitoid dynamics as discussed in section 4.3.1.

The E2 community was associated with heterotrophic flagellates and ciliates (Table S4),
consistent with summer sea ice (Leu et al., 2015; Marquardt et al., 2023). Dinoflagellates
included *Gymnodinium* and *Heterocapsa*with *Gymnodinium* spp. known to feed on small
diatoms (Eddie et al., 2010) and are correlated with higher light conditions (Duncan et al., 2024).
Additional E2 ASV associations included included a Pyramimonodaophyte, *Pyramimonas sp.,*
thecofilosean *Protaspa*, Oomycetes and Cilliophora. The green alga, *Pyramimonas*, sometimes
forms dense blooms in under-ice brackish water ponds associated with melt conditions
(Gradinger, 1996; Massicotte et al., 2020). The heterotrophic flagellate *Protaspa* can feed on

diatoms using pseudopodia (Howe et al., 2011; Schnepf and Kühn, 2000). The Oomycetes can
be parasitic (Thines et al., 2015) or saprotrophic, thus their increased presence in the post-
bloom community may represent an opportunity to feed on light and salinity stressed diatoms or
on the post-bloom exopolymers that are preferentially retained during melt (Juhl et al., 2011;
Meiners et al., 2008). The abundance of putative parasitoids both in bloom and post-bloom
conditions was striking, leading us to further explore their potential role in shaping microalgal
succession. While prokaryotic community composition appears largely shaped by bottom-up
processes, with shifts in dissolved organic matter quality and quantity selecting for taxa with
specialized metabolic capacities, eukaryotic community composition reflects both bottom-up
drivers, e.g., nutrient and light availability and desalination processes, and top-down interactions
such as those from parasitoids and grazers.

401 4.3. *Sea-ice parasitoid dynamics and potential host range*

To explore the relationship of parasitoid groups in shaping sea-ice microalgal communities, we
identified putative parasitoid sequences from those taxonomic groups expected to be found in
relatively high proportion within our samples: Chytridiomycetes (Fungi), Cryomonadida
(Cercozoa, Thecofilosea)*,* Oomycetes*,* Syndiniales (Dinophyceae) and Labyrinthulomycetes
(Bigyra). Within the Cryomonadida, the relationship between *Protaspa* and *Cryothecomonas* are
not fully resolved, so both of these groups were retained in this study despite parasitism
confined only to the *Cryothecomonas*. Our time-series data revealed successional patterns
through the microalgal bloom (Fig. 3). Environmental controls on parasitoid infections remain
poorly understood, aside from a few specific cases. Infections by the cryomonad
Cryothecomonas aestivalis appear inhibited below 4 °C (Catlett et al., 2023; Peacock et al.,
2014). Our observations demonstrate that parasitoid groups periodically spike, suggesting that
infections are occurring throughout the spring bloom at temperatures below -1 °C (Fig. 3). Brine
salinity varies more widely in the spring (Fig. S4c) and likely plays a key role in regulating
parasitoid pathogenicity in sea ice. Chytridiomycetes were abundant during the early bloom but
declined with melt onset and may be stenohaline, while cryomonads and Oomycetes may be
euryhaline and tolerant of fresher meltwater (Fig. 3). However, previous studies report both
cryomonads and chytrids as more common in phytoplankton influenced by sea-ice melt waters
(Kilias et al., 2020; Thaler and Lovejoy, 2012), implying that the physical melting and flushing of
the ice more strongly govern their presence. Stress due to excessive irradiance increases
microalgal susceptibility to chytrid infection in sea ice (Hassett and Gradinger, 2016), and along
with salinity stress, these factors may help shape parasitoid succession as host susceptibility
varies with environmental factors.

We analyzed 18S ASV time series data for significant correlations between putative
parasitoids and potential hosts, assuming correlations could be positive or negative depending
on the infection phase captured. It is important to note that correlation does not infer causality
and experimental validation is required to confirm real parasite-host associations.
Dinoflagellates showed far fewer correlations with putative parasitoids than diatoms, suggesting
they are less likely hosts (Fig. 4). The "kill-the-winner" hypothesis postulates that the most
abundant taxa experience greater infection rates from viruses or parasitoids, particularly when
pathogens are diverse or have a broad host range (Abonyi et al., 2024; Thingstad, 2000).
Navicula and *Nitzschia* were both highly abundant and significantly correlated with parasitoids,
and based on this hypothesis, we would expect more parasitoid associations with these genera.
Simpson Diversity was not significantly related to parasitoid abundance, which may reflect

uncertainty surrounding the assignment of ASVs as parasitoids, temporal lags between infection
and diversity responses, and confounding environmental factors (e.g., brine salinity). A targeted
study is required to confirm parasitoid behavior and further investigate whether the suppression
of dominant strains by parasitoids may promote microalgal coexistence and diversity in sea ice
(Abonyi et al., 2024).

Chytridiomycetes were the most abundant parasitoid group during the algal bloom's growth
phase (Fig. 3a), consistent with previous observations (Donk and Ringelberg, 1983; Ibelings et
al., 2004). These host-specific fungal parasites infect Arctic microalgae (Hassett and Gradinger,
2016) and exert top-down control, shaping bloom dynamics and succession. The chytrid ASV,
Lobulomycetales Order_1, was significantly correlated with a select few diatoms and
dinoflagellates, including *Nitzschia* Genus_4 (Fig. 4). However, their dynamics suggest temporal
separation rather than infection (Fig. 5). *Nitzschia* Genus_1 & _5, though not significant (p adj. >
0.05), showed strong correlations (rho ≥ 0.5) and dynamics resembling Lotka-Volterra patterns.
Nitzschia genera (ASV 1 & 5) were negatively correlated with Lobulomycetales Order_1 during
the bloom phase, but were overall, positively correlated, due to post-bloom dynamics. It is
therefore likely that we are missing true parasitoid-host interactions that were not significantly
correlated while also identifying false positives. Similar inconsistencies in known parasitoid-host
pairings have been difficult to parse from high-frequency long-term metabarcoding time-series
data due to inconsistent dynamics. We suggest that *Nitzschia* spp. were primary chytrid hosts
during our timeseries due to their contrasting peaks and troughs throughout the bloom. Although
lifecycle data for marine chytrids are limited, the amphibian pathogen *Batrachochytrium*
dendrobatidis, completes its lifecycle in ~4-5 days (Berger et al., 2005). In our time series,
periodic chytrid peaks, and *Nitzschia* genera 1 & 5 dips (Figs. 3 & 5), offset by ~2 days for
genus 5, suggest synchronized infection and zoospore release every ~7 days. We propose that
chytrid peaks reflect sporangia attached to algal hosts retained during sampling, while dips
reflect zoospore release and potential loss through brine drainage or dispersal into underlying
water. These waves of infection support a cyclic interaction with *Nitzschia* spp. (Fig. 5), though
microscopy, isolation, and co-culture are needed to confirm this lifecycle and host range.

Few correlations were found within the Labyrinthulomycetes (Fig. 4); though some
thraustochytrids can parasitize diatoms (Hassett, 2020) their typical role is organic matter
decomposition (Hassett and Gradinger, 2018). Oomycetes increased post-bloom, along with low
levels of Syndiniales (Fig. 3d,e), both of which have a broad host range across Kingdoms (Käse
et al., 2021). The order Syndiniales may be more important in underlying waters (Jacquemot et
al., 2022; Kellogg et al., 2019; Marquardt et al., 2016; Terrado et al., 2009). Most oomycete
ASVs (>85%) were unclassifiable beyond class level, as previously reported in Hassett et al.
(2019). Their post-bloom increase may reflect saprotrophic breakdown of ice-retained carbon.
However, a negative correlation with a *Cymbellales* ASV, supported by known *Lagena-Cymbella*
interaction (Thines and Buaya, 2022), and sequence homology to *Diatomophthora drebesii*, a
diatom parasitoid, suggest parasitic roles are probable. Given their relative abundance,
Oomycetes may play a key role in Arctic biogeochemical as both degraders and parasitoids.

Cryomonadida (Thecofilosea) featured prominently in our parasitoid correlation matrix (Fig.
4), with some ASVs showing broad host associations among diatoms. *Cryothecomonas* and
Protaspa, display two trophic modes: heterotrophy of small protists (Thaler and Lovejoy, 2012),
and by internal parasitization of large diatoms, with endoparasitism currently restricted to the
genus *Cryothecomonas* (Drebes et al., 1996; Kühn et al., 2000). Correlated diatom hosts (e.g.:

*Nitzschi*a and *Navicula* spp.) are typically large morphotypes within sea ice, consistent with
Cryothecomonas hosts. Although *C. aestivalis* is a known diatom parasitoid (Drebes et al.,
1996), Arctic *Cryothecomonas* spp. have only been observed as free-living grazers (Thaler and
Lovejoy, 2012). In 2022, we observed an undescribed *Cryothecomonas* sp. actively parasitizing
Nitzschia arctica, consuming the plastids and protoplasm (Fig. 6). *Cryothecomonas* was
identified according to Thomsen et al. (1991) and Schnepf & Kühn (2000), i.e., flagellates
possess two homodynamic flagella inserted apically,   asexual reproduction by binary fission,
flagellates were entirely within the host frustule, and the presence of brown digestive vacuoles
derived from plastids (Fig. 6). Information regarding the differentiation between the closely
related *Protaspa* and *Cryothecomonas* is limited, but current knowledge distinguishes the two
based on their feeding behavior, i.e., *Protaspa* attaches to the host diatom, where it remains
externally, and penetrates the frustule with a pseudopodia used for feeding (Chantangsi and
Leander, 2010; Hoppenrath and Leander, 2006). Endobiotic Oomycetes overtake the diatom
protoplasm with a swollen thallus which later develops into a sporangium that releases small
biflagellate zoospores (Garvetto et al., 2018); all stages are readily distinguished from
Cryothecomonas.

Our correlation data suggests that several genera beyond *Nitzschia* and *Navicula* may also
be hosts for cryomonads, though *C. aestivalis* is known to be host specific (Drebes et al., 1996;
Schnepf and Kühn, 2000) and our observations indicate *N. arctica* was the primary host in 2022
with ~5% of cells found to be infected. *Protaspa*-host correlations were similar to
Cryothecomonas suggesting potential host overlap or similar environmental response. As
Cryomonadida are abundant in polar waters (Thaler and Lovejoy, 2015; Thomsen et al., 1991),
their trophic role in sea ice warrants reevaluation in light of observed parasitization, particularly
their influence on diatom dynamics and succession.

We found compelling evidence that the wide diversity of parasitoids observed in this study
alter microalgal community composition, particularly of the most abundant sea-ice diatoms.
Previous studies have found that, depending on the host species and the parasitoid size
classes, carbon flux may be shunted to, or from, grazers (Rasconi et al., 2014). For example,
cryomonads and chytrids may redirect carbon from large diatoms—typically too big for grazers
like *Calanus glacialis*—into smaller, grazer-accessible forms such as flagellated cells,
zoospores, or sporangia (Cleary et al., 2017; Frenken et al., 2016; Rasconi et al., 2014).
Parasitoid infections in sea-ice may be prolific (pers. obs.) due to the dense microalgal
concentrations in the confined brine channels. Models and experimental investigations indicate
enhanced contact rates between viruses and hosts as brine volume decreases with lower
temperatures (Wells and Deming, 2006) while meiofaunal grazers become limited in their ability
to access brine channels with a diameter less than 200 μm (Krembs et al., 2000). The smaller
size of parasitoids, or of certain life stages, are less likely to be limited by narrowing of brine
channels, increasing their potential reach and influence within brine channels. Epidemics of
parasitoids or viruses may suppress microalgal blooms leading to patchiness, and shift
microalgal composition while increasing diversity (Donk and Ringelberg, 1983; Frenken et al.,
2016). We emphasize that further research into top-down controls of sea-ice algae and
prokaryotes by interactions with parasitoids and viruses is required to better understand
community composition structuring and succession.

5. Conclusions

As Arctic warming continues, monitoring the timing and magnitude of spring ice-algal blooms is
crucial for understanding the ecosystem response to environmental change. High-frequency
sampling of landfast sea ice near Utqiaġvik, Alaska in 2021 captured the full bloom cycle,
revealing a much larger magnitude relative to prior observations, and highlighting the
importance of nitrate limitation in constraining biomass accumulation, and physical forcing (melt
onset) on terminating the bloom. Parasitoids emerged as major contributors to the eukaryotic
microbial community. We report many potential associations of diatoms with *Cryothecomonas,*
including the first observation of parasitic behavior in sea ice, and suggest chytrid-linked
declines in *Nitzschia*. Understanding the complex dynamics of parasitoids within the sea-ice
ecosystem is crucial to elucidate their role in shaping microbial communities, nutrient cycles,
and broader ecological interactions. These findings support the potential role of parasitoids in
regulating dominant microalgae through "kill-the-winner" dynamics (Abonyi et al., 2024;
Thingstad, 2000), promoting coexistence and microbial diversity by freeing space within the
constrained brine channel system for other taxa. As a result, parasitoids may enhance resilience
through increased diversity at the base of the sea-ice food web – a critical component of the
rapidly changing Pacific Arctic ecosystem.

References

Abonyi, A., Fornberg, J., Rasconi, S., Ptacnik, R., Kainz, M.J., Lafferty, K.D., 2024. The chytrid insurance hypothesis: integrating parasitic chytrids into a biodiversity–ecosystem functioning framework for phytoplankton–zooplankton population dynamics. Oecologia 204, 279–288. https://doi.org/10.1007/s00442-024-05519-w

Ahmed, F., Leu, E., Juhl, A.R., Campbell, K., Dilliplaine, K.B., Assmy, P., Niemi, A., Gradinger, R., Alou-Font, E., Torres-Valdés, S., Whitmore, L., Jones, E.M., Fransson, A., Chierici, M., Olsen, L.M., McKay, R.D., Lee, S.H., Oggier, M., Lange, B.A., Tremblay, J.-É., Gosselin, M., Mundy, C.J., n.d. A pan-Arctic perspective on the influence of ice algae on sea-ice nutrient concentrations. Elem. Sci. Anthr. https://doi.org/ELEMENTA-D-25-00059R1

Alou-Font, E., Mundy, C., Roy, S., Gosselin, M., Agustí, S., 2013. Snow cover affects ice algal pigment composition in the coastal Arctic Ocean during spring. Mar. Ecol. Prog. Ser. 474, 89–104. https://doi.org/10.3354/meps10107

Ambrose, W.G., Quillfeldt, C. von, Clough, L.M., Tilney, P.V.R., Tucker, T., 2005. The sub-ice algal community in the Chukchi Sea: large- and small-scale patterns of abundance based on images from a remotely operated vehicle. Polar Biol. 28, 784–795. https://doi.org/10.1007/s00300-005-0002-8

Amundsen, P., Lafferty, K.D., Knudsen, R., Primicerio, R., Klemetsen, A., Kuris, A.M., 2009. Food web topology and parasites in the pelagic zone of a subarctic lake. J. Anim. Ecol. 78, 563–572. https://doi.org/10.1111/j.1365-2656.2008.01518.x

Apprill, A., McNally, S., Parsons, R., Weber, L., 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat. Microb. Ecol. 75, 129–137. https://doi.org/10.3354/ame01753

Arrigo, K.R., Mills, M.M., Kropuenske, L.R., van Dijken, G.L., Alderkamp, A.-C., Robinson, D.H., 2010. Photophysiology in Two Major Southern Ocean Phytoplankton Taxa: Photosynthesis and Growth of *Phaeocystis antarctica* and *Fragilariopsis cylindrus* under Different Irradiance Levels. Integr. Comp. Biol. 50, 950–966. https://doi.org/10.1093/icb/icq021

Aumack, C.F., Juhl, A.R., Krembs, C., 2014. Diatom vertical migration within land-fast Arctic sea ice. J. Marine Syst. 139, 496–504. https://doi.org/10.1016/j.jmarsys.2014.08.013

Babb, D.G., Galley, R.J., Howell, S.E.L., Landy, J.C., Stroeve, J.C., Barber, D.G., 2022. Increasing multiyear sea ice loss in the Beaufort Sea: A new export pathway for the diminishing multiyear ice cover of the Arctic Ocean. Geophys. Res. Lett. 49, 1–11. https://doi.org/10.1029/2021GL097595

Benjamini, Y., Heller, R., Yekutieli, D., 2009. Selective inference in complex research. Philos. Trans. R. Soc. A Math. Phys. Eng. Sci. 367, 4255–4271. https://doi.org/10.1098/rsta.2009.0127

Berger, L., Hyatt, A., Speare, R., Longcore, J., 2005. Life cycle stages of the amphibian chytrid *Batrachochytrium dendrobatidis*. Dis. Aquat. Organ. 68, 51–63. https://doi.org/10.3354/dao068051

Bintanja, R., Selten, F.M., 2014. Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat. Nature 509, 479–482. https://doi.org/10.1038/nature13259

Boetius, A., Albrecht, S., Bakker, K., Bienhold, C., Felden, J., Fernández-Méndez, M.,

Hendricks, S., Katlein, C., Lalande, C., Krumpen, T., Nicolaus, M., Peeken, I., Rabe, B., Rogacheva, A., Rybakova, E., Somavilla, R., Wenzhöfer, F., 2013. Export of algal biomass from the nelting Arctic sea ice. Science 339, 1430–1432. https://doi.org/10.1126/science.1231346

Bohórquez, J., McGenity, T.J., Papaspyrou, S., García-Robledo, E., Corzo, A., Underwood, G.J.C., 2017. Different types of diatom-derived extracellular polymeric substances drive changes in heterotrophic bacterial communities from intertidal sediments. Front. Microbiol. 8. https://doi.org/10.3389/fmicb.2017.00245

Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., Huttley, G.A., Gregory Caporaso, J., 2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6, 90. https://doi.org/10.1186/s40168-018-0470-z

Brakstad, O.G., Nonstad, I., Faksness, L.-G., Brandvik, P.J., 2008. Responses of microbial communities in Arctic sea ice after contamination by crude petroleum oil. Microb. Ecol. 55, 540–552. https://doi.org/10.1007/s00248-007-9299-x

Cáceres, M. De, Legendre, P., 2009. Associations between species and groups of sites: indices and statistical inference. Ecology 90, 3566–3574. https://doi.org/10.1890/08-1823.1

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. Nat. Methods 7, 335–336. https://doi.org/10.1038/nmeth.f.303

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., Knight, R., 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J. 6, 1621–1624. https://doi.org/10.1038/ismej.2012.8

Catlett, D., Peacock, E.E., Crockford, E.T., Futrelle, J., Batchelder, S., Stevens, B.L.F., Gast, R.J., Zhang, W.G., Sosik, H.M., 2023. Temperature dependence of parasitoid infection and abundance of a diatom revealed by automated imaging and classification. Proc. Natl. Acad. Sci. 120, 2017. https://doi.org/10.1073/pnas.2303356120

Chambouvet, A., Morin, P., Marie, D., Guillou, L., 2008. Control of toxic narine dinoflagellate blooms by serial parasitic killers. Science 322, 1254–1257. https://doi.org/10.1126/science.1164387

Chantangsi, C., Leander, B.S., 2010. An SSU rDNA barcoding approach to the diversity of marine interstitial cercozoans, including descriptions of four novel genera and nine novel species. Int. J. Syst. Evol. Microbiol. 60, 1962–1977. https://doi.org/10.1099/ijs.0.013888-0

Cleary, A.C., Søreide, J.E., Freese, D., Niehoff, B., Gabrielsen, T.M., 2017. Feeding by *Calanus glacialis* in a high Arctic fjord: potential seasonal importance of alternative prey. ICES J. Mar. Sci. 74, 1937–1946. https://doi.org/10.1093/icesjms/fsx106

Cono, V. La, Smedile, F., Crisafi, F., Marturano, L., Toshchakov, S. V., Spada, G. La, Bản, N.K., Yakimov, M.M., 2022. Wintertime simulations induce changes in the structure, diversity and function of Antarctic sea ice-associated microbial communities. Microorganisms 10, 623. https://doi.org/10.3390/microorganisms10030623

Cota, G.., Legendre, L., Gosselin, M., Ingram, R.., 1991. Ecology of bottom ice algae: I. Environmental controls and variability. J. Marine Syst. 2, 257–277. https://doi.org/10.1016/0924-7963(91)90036-T

Cota, G.F., Anning, J.L., Watson, N.H.F., Prinsenberg, S.J., Bennett, E.B., Lewis, M.R., 1987. Nutrient fluxes during extended blooms of Arctic ice algae. J. Geophys. Res. 92, 1951. https://doi.org/10.1029/JC092iC02p01951

Cota, G.F., Smith, R.E.H., 1991. Ecology of bottom ice algae: II. Dynamics, distributions and productivity. J. Marine Syst. 2, 279–295. https://doi.org/10.1016/0924-7963(91)90037-U

Cox, G.F.N., Weeks, W.F., 1983. Equations for determining the gas and brine volumes in sea ice samples. J. Glaciol. 29, 306–316. https://doi.org/10.3198/1983JoG29-102-306-316

Croteau, D., Lacour, T., Schiffrine, N., Morin, P.I., Forget, M.H., Bruyant, F., Ferland, J., Lafond, A., Campbell, D.A., Tremblay, J.É., Babin, M., Lavaud, J., 2022. Shifts in growth light optima among diatom species support their succession during the spring bloom in the Arctic. J. Ecol. 110, 1356–1375. https://doi.org/10.1111/1365-2745.13874

Dalman, L.A., Else, B.G.T., Barber, D., Carmack, E., Williams, W.J., Campbell, K., Duke, P.J., Kirillov, S., Mundy, C.J., 2019. Enhanced bottom-ice algal biomass across a tidal strait in the Kitikmeot Sea of the Canadian Arctic. Elem. Sci. Anthr. 7. https://doi.org/10.1525/elementa.361

Dilliplaine, K., Hennon, G., 2023. Impacts of crude oil on Arctic sea-ice diatoms modified by irradiance. Elem Sci Anth 11, 1–21. https://doi.org/10.1525/elementa.2023.00074

Donk, E. van, Ringelberg, J., 1983. The effect of fungal parasitism on the succession of diatoms in Lake Maarsseveen I (The Netherlands). Freshw. Biol. 13, 241–251. https://doi.org/10.1111/j.1365-2427.1983.tb00674.x

Drebes, G., Kühn, S.F., Gmelch, A., Schnepf, E., 1996. *Cryothecomonas aestivalis* sp. nov., a colourless nanoflagellate feeding on the marine centric diatom *Guinardia delicatula* (Cleve) Hasle. Helgoländer Meeresuntersuchungen 50, 497–515. https://doi.org/10.1007/BF02367163

Duncan, R.J., Søreide, J.E., Varpe, Ø., Wiktor, J., Pitusi, V., Runge, E., Petrou, K., 2024. Spatio-temporal dynamics in microalgal communities in Arctic land-fast sea ice. Prog. Oceanogr. 224, 103248. https://doi.org/10.1016/j.pocean.2024.103248

Eddie, B., Juhl, A., Krembs, C., Baysinger, C., Neuer, S., 2010. Effect of environmental variables on eukaryotic microbial community structure of land-fast Arctic sea ice. Environ. Microbiol. 12, 797–809. https://doi.org/10.1111/j.1462-2920.2009.02126.x

Eicken, H., Gradinger, R., Heinrichs, T., Johnson, M.A., Lovecraft, A.L., Kaufman, M., 2012. Automated ice mass balance site data (SIZONET).

Frenken, T., Velthuis, M., de Senerpont Domis, L.N., Stephan, S., Aben, R., Kosten, S., van Donk, E., Van de Waal, D.B., 2016. Warming accelerates termination of a phytoplankton spring bloom by fungal parasites. Glob. Change. Biol. 22, 299–309. https://doi.org/10.1111/gcb.13095

Garrison, D.L., Buck, K.R., 1986. Organism losses during ice melting: A serious bias in sea ice community studies. Polar Biol. 6, 237–239. https://doi.org/10.1007/BF00443401

Garrison, D.L., Buck, K.R., Fryxell, G.A., 1987. Algal assemblages in Antarctic pack ice and in

ice-edge plankton. J. Phycol. 23, 564–572. https://doi.org/10.1111/j.1529-
8817.1987.tb04206.x

Garvetto, A., Nézan, E., Badis, Y., Bilien, G., Arce, P., Bresnan, E., Gachon, C.M.M., Siano, R.,
2018. Novel widespread marine oomycetes parasitising diatoms, including the toxic genus
Pseudo-nitzschia: Genetic, morphological, and ecological characterisation. Front.
Microbiol. 9, 1–19. https://doi.org/10.3389/fmicb.2018.02918

Giesbrecht, K.E., Varela, D.E., 2021. Summertime biogenic silica production and silicon
limitation in the Pacific Arctic region from 2006 to 2016. Global Biogeochem. Cycles 35, 1–
27. https://doi.org/10.1029/2020GB006629

Glenn, T.C., Pierson, T.W., Bayona-Vásquez, N.J., Kieran, T.J., Hoffberg, S.L., Thomas, J.C.,
Lefever, D.E., Finger, J.W., Gao, B., Bian, X., Louha, S., Kolli, R.T., Bentley, K.E.,
Rushmore, J., Wong, K., Shaw, T.I., Rothrock, M.J., McKee, A.M., Guo, T.L., Mauricio, R.,
Molina, M., Cummings, B.S., Lash, L.H., Lu, K., Gilbert, G.S., Hubbell, S.P., Faircloth, B.C.,
2019. Adapterama II: Universal amplicon sequencing on Illumina platforms (TaggiMatrix).
PeerJ 2019. https://doi.org/10.7717/peerj.7786

Gosselin, M., Legendre, L., 1990. Light and nutrient limitation of sea-ice microalgae (Hudson
Bay, Canadian Arctic). J. Phycol. https://doi.org/10.1016/S0198-0254(06)80482-6

Gosselin, M., Levasseur, M., Wheeler, P.A., Horner, R.A., Booth, B.C., 1997. New
measurements of phytoplankton and ice algal production in the Arctic Ocean. Deep-Sea.
Res. Pt II 44, 1623–1644. https://doi.org/10.1016/S0967-0645(97)00054-4

Gradinger, R., 2009. Sea-ice algae: Major contributors to primary production and algal biomass
in the Chukchi and Beaufort Seas during May/June 2002. Deep-Sea Res. Pt II 56, 1201–
1212. https://doi.org/10.1016/j.dsr2.2008.10.016

Gradinger, R., 1996. Occurrence of an algal bloom under Arctic pack ice. Mar. Ecol. Prog. Ser.
131, 301–305. https://doi.org/10.3354/meps131301

Gradinger, R., Kaufman, M., Bluhm, B., 2009. Pivotal role of sea ice sediments in the seasonal
development of near-shore Arctic fast ice biota. Mar. Ecol. Prog. Ser. 394, 49–63.
https://doi.org/10.3354/meps08320

Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de
Vargas, C., Decelle, J., del Campo, J., Dolan, J.R., Dunthorn, M., Edvardsen, B.,
Holzmann, M., Kooistra, W.H.C.F., Lara, E., Le Bescot, N., Logares, R., Mahé, F.,
Massana, R., Montresor, M., Morard, R., Not, F., Pawlowski, J., Probert, I., Sauvadet, A.-
L., Siano, R., Stoeck, T., Vaulot, D., Zimmermann, P., Christen, R., 2012. The Protist
Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit
rRNA sequences with curated taxonomy. Nucleic Acids Res. 41, D597–D604.
https://doi.org/10.1093/nar/gks1160

Hassett, B.T., 2020. A widely distributed thraustochytrid parasite of diatoms isolated from the
Arctic represents a gen. and sp. nov. J. Eukaryot. Microbiol. 67, 480–490.
https://doi.org/10.1111/jeu.12796

Hassett, B. T., Borrego, E.J., Vonnahme, T.R., Rämä, T., Kolomiets, M. V., Gradinger, R., 2019.
Arctic marine fungi: biomass, functional genes, and putative ecological roles. ISME J. 13,
1484–1496. https://doi.org/10.1038/s41396-019-0368-1

Hassett, B.T., Gradinger, R., 2018. New species of saprobic Labyrinthulea

(=Labyrinthulomycota) and the erection of a gen. nov. to resolve molecular polyphyly within the Aplanochytrids. J. Eukaryot. Microbiol. 65, 475–483. https://doi.org/10.1111/jeu.12494

Hassett, B.T., Gradinger, R., 2016. Chytrids dominate arctic marine fungal communities. Environ. Microbiol. 18, 2001–2009. https://doi.org/10.1111/1462-2920.13216

Hassett, Brandon T, Thines, M., Buaya, A., Ploch, S., Gradinger, R., 2019. A glimpse into the biogeography, seasonality, and ecological functions of arctic marine Oomycota. IMA Fungus 10, 6. https://doi.org/10.1186/s43008-019-0006-6

Herzfeld, U.C., Maslanik, J.A., Sturm, M., 2006. Geostatistical characterization of snow-depth structures on sea ice near Point Barrow, Alaska—A contribution to the AMSR-Ice03 field validation campaign. IEEE Trans. Geosci. Remote Sens. 44, 3038–3056. https://doi.org/10.1109/TGRS.2006.883349

Hoppenrath, M., Leander, B.S., 2006. Dinoflagellate, Euglenid, or Cercomonad? The Ultrastructure and molecular phylogenetic position of *Protaspis grandis* n. sp. J. Eukaryot. Microbiol. 53, 327–342. https://doi.org/10.1111/j.1550-7408.2006.00110.x

Howe, A.T., Bass, D., Scoble, J.M., Lewis, R., Vickerman, K., Arndt, H., Cavalier-Smith, T., 2011. Novel cultured protists identify deep-branching environmental DNA clades of Cercozoa: New genera *Tremula*, *Micrometopion*, *Minimassisteria*, *Nudifila*, *Peregrinia*. Protist 162, 332–372. https://doi.org/10.1016/j.protis.2010.10.002

Howell, S.E.L., Laliberté, F., Kwok, R., Derksen, C., King, J., 2016. Landfast ice thickness in the Canadian Arctic Archipelago from observations and models. Cryosph. 10, 1463–1475. https://doi.org/10.5194/tc-10-1463-2016

Ibelings, B.W., De Bruin, A., Kagami, M., Rijkeboer, M., Brehm, M., Donk, E. Van, 2004. Host parasite interactions between freshwater phytoplankton and chytrid fungi (Chytridiomycota). J. Phycol. 40, 437–453. https://doi.org/10.1111/j.1529-8817.2004.03117.x

Jacquemot, L., Vigneron, A., Tremblay, J.-É., Lovejoy, C., 2022. Contrasting sea ice conditions shape microbial food webs in Hudson Bay (Canadian Arctic). ISME Commun. 2, 1–10. https://doi.org/10.1038/s43705-022-00192-7

Jin, M., Deal, C.J., Wang, J., Shin, K.-H., Tanaka, N., Whitledge, T.E., Lee, S.H., Gradinger, R.R., 2006. Controls of the landfast ice–ocean ecosystem offshore Barrow, Alaska. Ann. Glaciol. 44, 63–72. https://doi.org/10.3189/172756406781811709

Juhl, A., Krembs, C., Meiners, K., 2011. Seasonal development and differential retention of ice algae and other organic fractions in first-year Arctic sea ice. Mar. Ecol. Prog. Ser. 436, 1–16. https://doi.org/10.3354/meps09277

Juhl, A.R., Krembs, C., 2010. Effects of snow removal and algal photoacclimation on growth and export of ice algae. Polar Biol. 33, 1057–1065. https://doi.org/10.1007/s00300-010-0784-1

Juul-Pedersen, T., Michel, C., Gosselin, M., Seuthe, L., 2008. Seasonal changes in the sinking export of particulate material under first-year sea ice on the Mackenzie Shelf (western Canadian Arctic). Mar. Ecol. Prog. Ser. 353, 13–25. https://doi.org/10.3354/meps07165

Kaehler, B.D., Bokulich, N.A., McDonald, D., Knight, R., Caporaso, J.G., Huttley, G.A., 2019. Species abundance information improves sequence taxonomy classification accuracy. Nat. Commun. 10, 4643. https://doi.org/10.1038/s41467-019-12669-6

Käse, L., Metfies, K., Neuhaus, S., Boersma, M., Wiltshire, K.H., Kraberg, A.C., 2021. Host-parasitoid associations in marine planktonic time series: Can metabarcoding help reveal them? PLoS One 16, e0244817. https://doi.org/10.1371/journal.pone.0244817

Kassambara, A., 2023. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.

Kellogg, C.T.E., McClelland, J.W., Dunton, K.H., Crump, B.C., 2019. Strong seasonality in Arctic estuarine microbial food webs. Front. Microbiol. 10. https://doi.org/10.3389/fmicb.2019.02628

Kilias, E.S., Junges, L., Šupraha, L., Leonard, G., Metfies, K., Richards, T.A., 2020. Chytrid fungi distribution and co-occurrence with diatoms correlate with sea ice melt in the Arctic Ocean. Commun. Biol. 3, 183. https://doi.org/10.1038/s42003-020-0891-7

Kohlbach, D., Schaafsma, F.L., Graeve, M., Lebreton, B., Lange, B.A., David, C., Vortkamp, M., Flores, H., 2017. Strong linkage of polar cod (*Boreogadus saida*) to sea ice algae-produced carbon: Evidence from stomach content, fatty acid and stable isotope analyses. Prog. Oceanogr. 152, 62–74. https://doi.org/10.1016/j.pocean.2017.02.003

Krembs, C., Gradinger, R., Spindler, M., 2000. Implications of brine channel geometry and surface area for the interaction of sympagic organisms in Arctic sea ice. J. Exp. Mar. Biol. Ecol. 243, 55–80. https://doi.org/10.1016/S0022-0981(99)00111-2

Krolicka, A., Boccadoro, C., Nilsen, M.M., Demir-Hilton, E., Birch, J., Preston, C., Scholin, C., Baussant, T., 2019. Identification of microbial key-indicators of oil contamination at sea through tracking of oil biotransformation: An Arctic field and laboratory study. Sci. Total Environ. 696, 133715. https://doi.org/10.1016/j.scitotenv.2019.133715

Kühn, S., Lange, M., Medlin, L.K., 2000. Phylogenetic position of *Cryothecomonas* inferred from nuclear-encoded small subunit ribosomal RNA. Protist 151, 337–345. https://doi.org/10.1078/S1434-4610(04)70032-2

Kurtz, Z., Mueller, C., Miraldi, E., Bonneau, R., 2017. SpiecEasi: Sparse inverse covariance for ecological statistical inference.

Landa, M., Blain, S., Christaki, U., Monchy, S., Obernosterer, I., 2016. Shifts in bacterial community composition associated with increased carbon cycling in a mosaic of phytoplankton blooms. ISME J. 10, 39–50. https://doi.org/10.1038/ismej.2015.105

Lannuzel, D., Tedesco, L., van Leeuwe, M., Campbell, K., Flores, H., Delille, B., Miller, L., Stefels, J., Assmy, P., Bowman, J., Brown, K., Castellani, G., Chierici, M., Crabeck, O., Damm, E., Else, B., Fransson, A., Fripiat, F., Geilfus, N.-X., Jacques, C., Jones, E., Kaartokallio, H., Kotovitch, M., Meiners, K., Moreau, S., Nomura, D., Peeken, I., Rintala, J.-M., Steiner, N., Tison, J.-L., Vancoppenolle, M., Van der Linden, F., Vichi, M., Wongpan, P., 2020. The future of Arctic sea-ice biogeochemistry and ice-associated ecosystems. Nat. Clim. Chang. 10, 983–992. https://doi.org/10.1038/s41558-020-00940-4

Lee, S.H., Whitledge, T.E., Kang, S.H., 2008. Spring time production of bottom ice algae in the landfast sea ice zone at Barrow, Alaska. J. Exp. Mar. Biol. Ecol. 367, 204–212. https://doi.org/10.1016/j.jembe.2008.09.018

Leu, E., Mundy, C.J., Assmy, P., Campbell, K., Gabrielsen, T.M., Gosselin, M., Juul-Pedersen, T., Gradinger, R., 2015. Arctic spring awakening - Steering principles behind the phenology of vernal ice algal blooms. Prog. Oceanogr. 139, 151–170. https://doi.org/10.1016/j.pocean.2015.07.012

Litchman, E., de Tezanos Pinto, P., Edwards, K.F., Klausmeier, C.A., Kremer, C.T., Thomas, M.K., 2015a. Global biogeochemical impacts of phytoplankton: A trait-based perspective. J. Ecol. 103, 1384–1396. https://doi.org/10.1111/1365-2745.12438

Litchman, E., Edwards, K.F., Klausmeier, C.A., 2015b. Microbial resource utilization traits and trade-offs: Implications for community structure, functioning, and biogeochemical impacts at present and in the future. Front. Microbiol. 6, 1–10. https://doi.org/10.3389/fmicb.2015.00254

Lizotte, M.P., 2001. The contributions of sea ice algae to Antarctic marine primary production 41, 57–73.

Lofthus, S., Netzer, R., Lewin, A.S., Heggeset, T.M.B., Haugen, T., Brakstad, O.G., 2018. Biodegradation of n-alkanes on oil–seawater interfaces at different temperatures and microbial communities associated with the degradation. Biodegradation 29, 141–157. https://doi.org/10.1007/s10532-018-9819-z

Mahoney, A., Eicken, H., Shapiro, L., 2007. How fast is landfast sea ice? A study of the attachment and detachment of nearshore ice at Barrow, Alaska. Cold Reg. Sci. Technol. 47, 233–255. https://doi.org/10.1016/j.coldregions.2006.09.005

Manes, S.S., Gradinger, R., 2009. Small scale vertical gradients of Arctic ice algal photophysiological properties. Photosynth. Res. 53–66. https://doi.org/10.1007/s11120-009-9489-0

Marquardt, M., Goraguer, L., Assmy, P., Bluhm, B.A., Aaboe, S., Down, E., Patrohay, E., Edvardsen, B., Tatarek, A., Smoła, Z., Wiktor, J., Gradinger, R., 2023. Seasonal dynamics of sea-ice protist and meiofauna in the northwestern Barents Sea. Prog. Oceanogr. 218, 103128. https://doi.org/10.1016/j.pocean.2023.103128

Marquardt, M., Vader, A., Stübner, E.I., Reigstad, M., Gabrielsen, T.M., 2016. Strong seasonality of marine microbial eukaryotes in a high-Arctic fjord (Isfjorden, in West Spitsbergen, Norway). Appl. Environ. Microbiol. 82, 1868–1880. https://doi.org/10.1128/AEM.03208-15

Massicotte, P., Amiraux, R., Amyot, M.-P., Archambault, P., Ardyna, M., Arnaud, L., Artigue, L., Aubry, C., Ayotte, P., Bécu, G., Bélanger, S., Benner, R., Bittig, H.C., Bricaud, A., Brossier, É., Bruyant, F., Chauvaud, L., Christiansen-Stowe, D., Claustre, H., Cornet-Barthaux, V., Coupel, P., Cox, C., Delaforge, A., Dezutter, T., Dimier, C., Domine, F., Dufour, F., Dufresne, C., Dumont, D., Ehn, J., Else, B., Ferland, J., Forget, M.-H., Fortier, L., Galí, M., Galindo, V., Gallinari, M., Garcia, N., Gérikas Ribeiro, C., Gourdal, M., Gourvil, P., Goyens, C., Grondin, P.-L., Guillot, P., Guilmette, C., Houssais, M.-N., Joux, F., Lacour, L., Lacour, T., Lafond, A., Lagunas, J., Lalande, C., Laliberté, J., Lambert-Girard, S., Larivière, J., Lavaud, J., LeBaron, A., Leblanc, K., Le Gall, F., Legras, J., Lemire, M., Levasseur, M., Leymarie, E., Leynaert, A., Lopes dos Santos, A., Lourenço, A., Mah, D., Marec, C., Marie, D., Martin, N., Marty, C., Marty, S., Massé, G., Matsuoka, A., Matthes, L., Moriceau, B., Muller, P.-E., Mundy, C.-J., Neukermans, G., Oziel, L., Panagiotopoulos, C., Pangrazi, J.-J., Picard, G., Picheral, M., Pinczon du Sel, F., Pogorzelec, N., Probert, I., Quéguiner, B., Raimbault, P., Ras, J., Rehm, E., Reimer, E., Rontani, J.-F., Rysgaard, S., Saint-Béat, B., Sampei, M., Sansoulet, J., Schmechtig, C., Schmidt, S., Sempéré, R., Sévigny, C., Shen, Y., Tragin, M., Tremblay, J.-É., Vaulot, D., Verin, G., Vivier, F., Vladoiu, A., Whitehead, J., Babin, M., 2020. Green Edge ice camp campaigns: understanding the processes controlling the under-ice Arctic phytoplankton spring bloom. Earth Syst. Sci. Data 12, 151–176. https://doi.org/10.5194/essd-12-151-2020

Mcdonald, S., Koulis, T., Ehn, J., Campbell, K., Gosselin, M., Mundy, C.J., 2015. A functional
regression model for predicting optical depth and estimating attenuation coefficients in sea-
ice covers near Resolute Passage, Canada. Ann. Glaciol. 56, 147–154.
https://doi.org/10.3189/2015AoG69A004

Meiners, K., Krembs, C., Gradinger, R., 2008. Exopolymer particles: microbial hotspots of
enhanced bacterial activity in Arctic fast ice (Chukchi Sea). Aquat. Microb. Ecol. 52, 195–
207. https://doi.org/10.3354/ame01214

Mortenson, E., Steiner, N., Monahan, A.H., Hayashida, H., Sou, T., Shao, A., 2020. Modeled
impacts of sea ice exchange processes on Arctic Ocean carbon uptake and acidification
(1980–2015). J. Geophys. Res-Oceans. 125, 1–20. https://doi.org/10.1029/2019JC015782

Mundy, C.J., Leu, E., Campbell, K., Galindo, V., Levasseur, M., Poulin, M., Tremblay, J.-É.,
Gosselin, M., 2025. Intracellular nutrient storage during ice algal spring blooms in the
Canadian high Arctic. iScience 28, 113148. https://doi.org/10.1016/j.isci.2025.113148

Netzer, R., Henry, I.A., Ribicic, D., Wibberg, D., Brönner, U., Brakstad, O.G., 2018. Petroleum
hydrocarbon and microbial community structure successions in marine oil-related
aggregates associated with diatoms relevant for Arctic conditions. Mar. Pollut. Bull. 135,
759–768. https://doi.org/10.1016/j.marpolbul.2018.07.074

Osborne, E., Richter-Menge, J., Jeffries, M., 2018. Arctic Report Card 2018, Arctic Report Card.

Oziel, L., Massicotte, P., Randelhoff, A., Ferland, J., Vladoiu, A., Lacour, L., Galindo, V.,
Lambert-Girard, S., Dumont, D., Cuypers, Y., Bouruet-Aubertot, P., Mundy, C.-J., Ehn, J.,
Bécu, G., Marec, C., Forget, M.-H., Garcia, N., Coupel, P., Raimbault, P., Houssais, M.-N.,
Babin, M., 2019. Environmental factors influencing the seasonal dynamics of spring algal
blooms in and beneath sea ice in western Baffin Bay. Elem. Sci. Anthr. 7.
https://doi.org/10.1525/elementa.372

Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: Assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. Environ. Microbiol. 18, 1403–1414. https://doi.org/10.1111/1462-
2920.13023

Peacock, E., Olson, R., Sosik, H., 2014. Parasitic infection of the diatom *Guinardia delicatula*, a
recurrent and ecologically important phenomenon on the New England Shelf. Mar. Ecol.
Prog. Ser. 503, 1–10. https://doi.org/10.3354/meps10784

Pineault, S., Tremblay, J., Gosselin, M., Thomas, H., Shadwick, E., 2013. The isotopic signature
of particulate organic C and N in bottom ice: Key influencing factors and applications for
tracing the fate of ice⬚algae in the Arctic Ocean. J. Geophys. Res-Oceans. 118, 287–300.
https://doi.org/10.1029/2012JC008331

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner,
F.O., 2012. The SILVA ribosomal RNA gene database project: improved data processing
and web-based tools. Nucleic Acids Res. 41, D590–D596.
https://doi.org/10.1093/nar/gks1219

Rapp, J.Z., Fernández-Méndez, M., Bienhold, C., Boetius, A., 2018. Effects of ice-algal
aggregate export on the connectivity of bacterial communities in the central Arctic Ocean.
Front. Microbiol. 9. https://doi.org/10.3389/fmicb.2018.01035

Rasconi, S., Grami, B., Niquil, N., Jobard, M., Sime-Ngando, T., 2014. Parasitic chytrids sustain

zooplankton growth during inedible algal bloom. Front. Microbiol. 5, 1–19.
https://doi.org/10.3389/fmicb.2014.00229

Redfield, A., Ketchum, B., Richards, F., 1963. The influence of organisms on the composition of
sea-water, in: The Sea. Interscience Publishers, New York, pp. 26–77.

Revelle, W., 2023. psych: Procedures for psychological, psychometric, and personality
research.

Robeson, M.S., O'Rourke, D.R., Kaehler, B.D., Ziemski, M., Dillon, M.R., Foster, J.T., Bokulich,
N.A., 2021. RESCRIPt: Reproducible sequence taxonomy reference database
management. PLOS Comput. Biol. 17, e1009581.
https://doi.org/10.1371/journal.pcbi.1009581

Roukaerts, A., Deman, F., Van der Linden, F., Carnat, G., Bratkic, A., Moreau, S., Lannuzel, D.,
Dehairs, F., Delille, B., Tison, J.-L., Fripiat, F., 2021. The biogeochemical role of a
microbial biofilm in sea ice. Elem. Sci. Anthr. 9, 1–14.
https://doi.org/10.1525/elementa.2020.00134

Rózańska, M., Gosselin, M., Poulin, M., Wiktor, J., Michel, C., 2009. Influence of environmental
factors on the development of bottom ice protist communities during the winter–spring
transition. Mar. Ecol. Prog. Ser. 386, 43–59. https://doi.org/10.3354/meps08092

Schnepf, E., Kühn, S.F., 2000. Food uptake and fine structure of *Cryothecomonas longipes* sp.
nov., a marine nanoflagellate incertae sedis feeding phagotrophically on large diatoms.
Helgol. Mar. Res. 54, 18–32. https://doi.org/10.1007/s101520050032

Schultz-Johansen, M., Bech, P.K., Hennessy, R.C., Glaring, M.A., Barbeyron, T., Czjzek, M.,
Stougaard, P., 2018. A novel enzyme portfolio for red algal polysaccharide degradation in
the marine bacterium *Paraglaciecola hydrolytica* S66T encoded in a sizeable
polysaccharide utilization locus. Front. Microbiol. 9, 1–15.
https://doi.org/10.3389/fmicb.2018.00839

Sipler, R.E., Kellogg, C.T.E., Connelly, T.L., Roberts, Q.N., Yager, P.L., Bronk, D.A., 2017.
Microbial community response to terrestrially derived dissolved organic matter in the
coastal Arctic. Front. Microbiol. 8, 1–19. https://doi.org/10.3389/fmicb.2017.01018

Skovgaard, A., 2014. Dirty tricks in the plankton: Diversity and role of marine parasitic protists.
Acta Protozool. 53, 51–62. https://doi.org/10.4467/16890027AP.14.006.1443

Smith, R.E.H., Harrison, W.G., Harris, L.R., Herman, A.W., 1990. Vertical fine structure of
particulate matter and nutrients in sea ice of the high Arctic. Can. J. Fish. Aquat. Sci. 47,
1348–1355. https://doi.org/10.1139/f90-154

Spindler, M., 1994. Notes on the biology of sea ice in the Arctic and Antarctic. Polar Biol. 14,
319–324. https://doi.org/10.1007/BF00238447

Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M.D.M., Breiner, H.-W., Richards, T.A.,
2010. Multiple marker parallel tag environmental DNA sequencing reveals a highly complex
eukaryotic community in marine anoxic water. Mol. Ecol. 19, 21–31.
https://doi.org/10.1111/j.1365-294X.2009.04480.x

Stroeve, J., Notz, D., 2018. Changing state of Arctic sea ice across all seasons. Environ. Res.
Lett. 13, 103001. https://doi.org/10.1088/1748-9326/aade56

Sun, A.-Q., Deng-Li, J.-Y., Zhang, H.-Z., Teng, J.-H., Du, Z.-J., Lu, D.-C., 2025. Genomic

characterization and new species identification of *Winogradskyella* with comparative
analysis of polysaccharide utilization loci. bioRxiv 2025.03.31.646394.
https://doi.org/10.1101/2025.03.31.646394

Syvertsen, E.E., 1991. Ice algae in the Barents Sea: types of assemblages, origin, fate and role
in the ice-edge phytoplankton bloom. Polar Res. 10, 277–288.
https://doi.org/10.1111/j.1751-8369.1991.tb00653.x

Terrado, R., Vincent, W., Lovejoy, C., 2009. Mesopelagic protists: diversity and succession in a
coastal Arctic ecosystem. Aquat. Microb. Ecol. 56, 25–39.
https://doi.org/10.3354/ame01327

Thaler, M., Lovejoy, C., 2015. Biogeography of heterotrophic flagellate populations indicates the
presence of generalist and specialist taxa in the Arctic Ocean. Appl. Environ. Microbiol. 81,
2137–2148. https://doi.org/10.1128/AEM.02737-14

Thaler, M., Lovejoy, C., 2012. Distribution and diversity of a protist predator *Cryothecomonas*
(Cercozoa) in Arctic marine waters. J. Eukaryot. Microbiol. 59, 291–299.
https://doi.org/10.1111/j.1550-7408.2012.00631.x

Thines, M., Buaya, A.T., 2022. *Lagena*—an overlooked oomycete genus with a wide range of
hosts. Mycol. Prog. 21, 66. https://doi.org/10.1007/s11557-022-01818-9

Thines, M., Nam, B., Nigrelli, L., Beakes, G., Kraberg, A., 2015. The diatom parasite *Lagenisma*
coscinodisci (Lagenismatales, Oomycota) is an early diverging lineage of the
Saprolegniomycetes. Mycol. Prog. 14, 75. https://doi.org/10.1007/s11557-015-1099-y

Thingstad, T.F., 2000. Elements of a theory for the mechanisms controlling abundance,
diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. Limnol.
Oceanogr. 45, 1320–1328. https://doi.org/10.4319/lo.2000.45.6.1320

Thompson, H.F., Summers, S., Yuecel, R., Gutierrez, T., 2020. Hydrocarbon-degrading bacteria
found tightly associated with the 50–70 µm cell-size population of eukaryotic phytoplankton
in surface waters of a Northeast Atlantic region. Microorganisms 8, 1955.
https://doi.org/10.3390/microorganisms8121955

Thomsen, H.A., Buck, K.R., Bolt, P.A., Garrison, D.L., 1991. Fine structure and biology of
Cryothecomonas gen. nov. (Protista incertae sedis) from the ice biota. Can. J. Zool. 69,
1048–1070. https://doi.org/10.1139/z91-150

Tichy, L., Chytry, M., 2006. Statistical determination of diagnostic species for site groups of
unequal size. J. Veg. Sci. 17, 809–818. https://doi.org/10.1111/j.1654-1103.2006.tb02504.x

Tillmann, U., Hesse, K.-J., Tillmann, A., 1999. Large-scale parasitic infection of diatoms in the
Northfrisian Wadden Sea. J. Sea Res. 42, 255–261. https://doi.org/10.1016/S1385-
1101(99)00029-5

Underwood, G.J.C., Michel, C., Meisterhans, G., Niemi, A., Belzile, C., Witt, M., Dumbrell, A.J.,
Koch, B.P., 2019. Organic matter from Arctic sea-ice loss alters bacterial community
structure and function. Nat. Clim. Chang. 9, 170–176. https://doi.org/10.1038/s41558-018-
0391-7

Veyssière, G., Castellani, G., Wilkinson, J., Karcher, M., Hayward, A., Stroeve, J.C., Nicolaus,
M., Kim, J.-H., Yang, E.-J., Valcic, L., Kauker, F., Khan, A.L., Rogers, I., Jung, J., 2022.
Under-ice light field in the western Arctic Ocean during late summer. Front. Earth Sci. 9, 1–
19. https://doi.org/10.3389/feart.2021.643737

von Quillfeldt, C.H., Ambrose, W.G., Clough, L.M., 2003. High number of diatom species in first-
year ice from the Chukchi Sea. Polar Biol. 26, 806–818. https://doi.org/10.1007/s00300-
003-0549-1

Walker, A.M., Leigh, M.B., Mincks, S.L., 2023. Benthic bacteria and archaea in the North
American Arctic reflect food supply regimes and impacts of coastal and riverine inputs.
Deep-Sea Res. Pt II 207, 105224. https://doi.org/10.1016/j.dsr2.2022.105224

Walsh, J.E., Eicken, H., Redilla, K., Johnson, M., 2022. Sea ice breakup and freeze-up
indicators for users of the Arctic coastal environment. Cryosph. 16, 4617–4635.
https://doi.org/10.5194/tc-16-4617-2022

Webster, M.A., Rigor, I.G., Nghiem, S. V., Kurtz, N.T., Farrell, S.L., Perovich, D.K., Sturm, M.,
2014. Interdecadal changes in snow depth on Arctic sea ice. J. Geophys. Res-Oceans.
119, 5395–5406. https://doi.org/10.1002/2014jc009985

Wells, L.E., Deming, J.W., 2006. Modelled and measured dynamics of viruses in Arctic winter
sea☐ice brines. Environ. Microbiol. 8, 1115–1121. https://doi.org/10.1111/j.1462-
2920.2006.00984.x

Whitmore, L., Oggier, M., Dilliplaine, K., Kaufman, M., Bolt, C., Aguilar-Islas, A., 2024. Sea ice
biogeochemistry time series at Utqiaġvik, Alaska (April - June 2021) [WWW Document].
Arct. Data Cent. https://doi.org/10.18739/A21J9793S

Yakimov, M.M., Giuliano, L., Gentile, G., Crisafi, E., Chernikova, T.N., Abraham, W.-R.,
Lünsdorf, H., Timmis, K.N., Golyshin, P.N., 2003. *Oleispira antarctica* gen. nov., sp. nov., a
novel hydrocarbonoclastic marine bacterium isolated from Antarctic coastal sea water. Int.
J. Syst. Evol. Microbiol. 53, 779–785. https://doi.org/10.1099/ijs.0.02366-0

Yan, D., Yoshida, K., Nishioka, J., Ito, M., Toyota, T., Suzuki, K., 2020. Response to sea ice
melt indicates high seeding potential of the ice diatom *Thalassiosira* to spring
phytoplankton blooms: A laboratory study on an ice algal community from the Sea of
Okhotsk. Front. Mar. Sci. 7. https://doi.org/10.3389/fmars.2020.00613

Yang, T., Nigro, L.M., Gutierrez, T., D′Ambrosio, L., Joye, S.B., Highsmith, R., Teske, A., 2016.
Pulsed blooms and persistent oil-degrading bacterial populations in the water column
during and after the Deepwater Horizon blowout. Deep-Sea Res. Pt II 129, 282–291.
https://doi.org/10.1016/j.dsr2.2014.01.014

Zhang, H., Wang, K., Shen, L., Chen, H., Hou, F., Zhou, X., Zhang, D., Zhu, X., 2018. Microbial
community dynamics and assembly follow trajectories of an early-spring diatom bloom in a
semienclosed bay. Appl. Environ. Microbiol. 84. https://doi.org/10.1128/AEM.01000-18

Zhou, J., Richlen, M.L., Sehein, T.R., Kulis, D.M., Anderson, D.M., Cai, Z., 2018. Microbial
community structure and associations during a marine dinoflagellate bloom. Front.
Microbiol. 9, 1–21. https://doi.org/10.3389/fmicb.2018.01201

1007

1008

1009

Figure 1. Development of biologically relevant measures over time. Temporal evolution of chlorophyll a and phaeophytin proportion (a), seawater concentrations of ammonium (NH_4^+), nitrite + nitrate (NO_2^- + NO_3^-), phosphate (PO_4^{3-}) and silicate ($Si(OH)_4$), (b), and snow thickness and under-ice PAR as measured by a spherical quantum sensor (4π; c). Green shading indicates the ice-algal bloom and the vertical dashed line indicates the bloom termination (<1 mg chlorophyll a m^{-2}).

1015

Figure 2. Sympagic microbial community structure and the most abundant genera over time. Non-metric multidimensional scaling (nMDS) ordination using Bray-Curtis dissimilarities (a & b). Sample groupings were determined by hierarchical clustering and labeled by their relation to the stage of the bloom. For prokaryotes (a), early- (P1), mid- (P2), and post-bloom (P3) clusters are observed. For Eukaryotes (b), bloom (E1) and post-bloom (E2) clusters were distinguished. Relative read contribution of the top 10 most abundant genera of prokaryotes (c) and unicellular eukaryotes (d). Vertical dashed line indicates bloom termination (<1 mg chlorophyll a m^{-2}).

1023

Figure 3. Relative abundance of putative parasitoid groups over time. (a) Chytridiomycota, class Chytridiomycetes, (b) Cercozoa, class Thecofilosea, (c) Bigyra, order Labyrinthulomycetes, (d) Oomycetes, (e) Dinoflagellata order Syndiniales. Green shading indicates the ice-algal bloom and the vertical dashed line indicates the bloom termination (1 mg chlorophyll a m^{-2}).

1028

Figure 4. Correlation of putative parasitoids with putative hosts. Directed spearman correlation coefficient grid of amplicon sequence variants (ASVs) containing correlations between putative parasitoids and putative hosts with |rho| ≥ 0.5. Black dots indicate significant correlation (adj. p < 0.05).

1032

Figure 5. Relative abundance over time of a putative parasitoid (red line, Chytridiomycota) and *Nitzschia* hosts (black, grey, orange and purple lines). Solid lines indicate a significant correlation (p < 0.05); all *Nitzschia* ASVs presented here had a correlation coefficient ≥ |0.5|. Green shading indicates the ice-algal bloom and the vertical dashed line indicates the bloom termination (1 mg chlorophyll a m^{-2}).

1037

Figure 6. A small chain of *Nitzschia arctica* infected by *Cryothecomonas sp.* at different stages of infection. White arrow points to a compromise in the frustule that may be an entry point for the flagellate and black arrows point to flagella. 1) A single flagellate beginning to phagocytize the plastid of a newly infected diatom. 2) Two small daughter cells with digestive vacuoles and fresh, still green, plastid material. Note the partially consumed plastid in the top left of the diatom frustule. 3) A mature flagellate with brown digestive vacuoles and surrounded by several defecated brown fecal bodies. 4) A large flagellate with many brown digestive vacuoles (the missing diatom plastids) with a cleavage furrow developing at the top left indicating the initiation of longitudinal mitotic division.

Chytridiomycetes

Thecofilosea
Cryomonadida Order
Cryothecomonas
Protaspa

Labyrinthulomycetes (Bigyra)

Oomycetes

Syndiniales (Dinophyceae)

Relative abundance (%)

Day of year

20 µm