Viral infection induces oxylipin chemical signaling at the end of a summer upwelling bloom:
implications for carbon cycling.
Bethanie Edwards[1*], Kimberlee Thamatrakoln[2], Chana F. Kranzler[3], Justin Ossolinski[4], Helen Fredricks[4],
Matthew D. Johnson[5], Jeffrey W. Krause[6,7], Kay D. Bidle[2], Ben Van Mooy[4].
1- Department of Earth and Planetary Science, University of California-Berkeley, Berkeley, CA 94720, USA
2-Department of Marine and Coastal Sciences, Rutgers University, New Brunswick, NJ 08901, USA;
3- Mina & Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat Gan, Israel
4-Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Falmouth, MA 02543
5- Department of Biological Oceanography, Woods Hole Oceanographic Institution, Falmouth, MA 02543
6- Dauphin Island Sea Lab, Dauphin Island, AL, USA.
7- Stokes School of Marine and Environmental Sciences, University of South Alabama, Mobile, AL 36608 USA
*Corresponding author: bethanie_edwards@berkeley.edu

Abstract/Summary (298/300-word max)
Diatoms are large phytoplankton that form the base of the marine food web and often bloom first when
nutrients are injected into the surface ocean through upwelling or deep ocean mixing[1,2]. Diatoms
contribute 20% of global photosynthesis[3] while disproportionately representing 40% of carbon export[4],
with most export occurring along the continental margins[5]. Oxylipin chemical signaling by diatoms has
been extensively studied in the Mediterranean Sea where oxylipins are linked to grazing with subsequent
insidious effects on copepod reproduction[6–13]. Culture studies with diatoms have shown that stress,
growth phase, and viral infection also impact oxylipin production[14–16]. This study provides the first
glimpse into the role of oxylipins as biomarkers and chemical signals during diatom viral infection at sea.
Biomarkers for lysis and senescence were identified in laboratory experiments and observed at elevated
concentrations in meta-lipidomes collected in the California Coastal Ecosystem (CCE) where diatoms had
recently been lysed by viruses[17]. Deck-board incubations with natural communities showed that oxylipins
stimulate particle-attached and surface-ocean microbes in a dose and community-dependent manner,
while inhibiting microzooplankton grazing and phytoplankton growth rates. Carbon export was two times
higher at the Post-lytic site than elsewhere along the transect consistent with the viral shuttle, whereby
viruses facilitate carbon export. We previously reported enhanced enzymatic activity at the Post-lytic
site[17], suggestive of the viral shunt, whereby carbon is remineralized or attenuated into non-sinking
dissolved organic matter. Here we layer geochemical evidence to show that lysis of oxylipin producing
diatoms amplified the vertical flux of carbon from the surface ocean even in the presence of viral shunt
processes. The remineralization length scale and community composition have been hypothesized as
controls on shunt vs. shuttle[18–20]; our analysis provides another example of how community interactions
may toggle a system between favoring shunt or shuttle.

Introduction
In 1999, Miralto *et al*. linked chemical signals produced by diatoms to decreased copepod egg viability
during spring blooms in the Adriatic Sea (Northern Mediterranean)[6]. These chemical signals,
polyunsaturated aldehydes (PUAs; Fig 1A), belong to the oxylipin subclass of lipids and resemble grazing
defense molecules produced by higher plants. Subsequent research using diatom cultures explored the
biosynthesis pathways of PUAs and other non-volatile oxylipins (NVOs)[14,15,21,22], which rely on
enzymatic oxidation of polyunsaturated fatty acids, specifically C22:6, C20:5, C20:4, C16:4, and C16:3
in diatoms. The production of oxylipins in diatoms is generally tied to stress with studies showing
relationships between enhanced oxylipin concentrations and nutrient stress, copepod grazing, cell

disruption, microzooplankton grazing, membrane permeability, virus infection and growth phase[8,10,12,16,23–
26].

To better understand the breadth of oxylipin bioactivity in the sea, scientists exposed a wide range of
marine organisms to oxylipins in amendment experiments with cultured isolates and natural
populations[27–35](summarized in Fig. 1B), revealing that oxylipins have a variety of cellular and ecosystem
impacts. Oxylipins are largely deleterious, causing teratogenic effects on the offspring of metazoan
grazers[6,7,9,10,36], deterring grazing by microzooplankton[31–33,37], while being allelopathic and stunting the
growth of eukaryotic phytoplankton competitors[27,31]. Specifically, PUAs are also a fundamental
component of a nitric oxide-based, stress surveillance system whereby diatoms sense and respond to
biotic and abiotic stressors, which can lead to the activation of programmed cell death at threshold
concentrations[29]. At sub-lethal concentrations, PUAs can serve to desensitize diatom cells to
physiological stress and fortify cellular responses for growth and survival[29]. Lastly, PUAs have been
found to be stimulatory to particle-associated bacteria and diatom epibionts in sublethal doses (1-10 μM
in the Atlantic; 13-18 μM in culture) while free-living bacterial isolates in the lab tend to be inhibited at
concentrations as low as 3 μM[28,35]. PUA concentrations high enough to deter microzooplankton grazing
and stimulate particle associated communities have been observed in the water column in the
Mediterranean Sea[7,23,25,27,38] , within sinking particles in the Atlantic[35], and within suspended particles in
the Pearl River Estuary, China[39] (Table S1; Fig. S1).

65
Figure 1. Diatoms in the ocean biosynthesize oxylipins from membrane lipids, which act as chemical signals
impacting their growth, behavior, cell fate, and biogeochemistry as well as that of the surrounding organisms.
A) Oxylipin biosynthesis pathway proposed for the 14-lipoxygenase enzyme acting on eicosapentaenoic acid which
was lipolytically cleaved from the membrane lipid, MGDG 20:5/16:3 (monogalactosyldiacylglycerol). A
hydroperoxy acid is the first compound produced, 14-hydroperoxy eicosapentaenoic acid (14-HpEPE). Enzymes
downstream of LOX act on the hydroperoxy acid to produce a plethora of oxylipin compounds. Here we include
other common lipoxygenases which add a hydroperoxy group (-OOH) at the specified carbon position along the
fatty acid chain. **B)** Assessments of oxylipin bioactivity in surface ocean ecosystems based on culture amendment
experiments and deck-board incubations[6,7,12–16,26–28,31,32,35,40,41]. Arrows denote the directionality of response and +/-
denotes stimulation or inhibition, respectively (depending on concentration). Circled responses indicate interactions
explored in this study. Inset shows the trophic cascade mediated by PUAs in a natural population from the
Chesapeake Bay, where oxylipins decreased grazing of microzooplankton like dinoflagellates (pink flagellate) and
ciliates (purple ciliated cell) on diatoms but increased grazing on cyanobacteria (green chain-forming cells) and
increased copepod grazing on both diatoms and ciliates[31].

In a separate manuscript, we presented a detailed comparison of the suite of oxylipins produced by three
different *Chaetoceros* diatom host-virus systems, highlighting the unique oxylipins produced by each
host-virus pair, and focusing on compounds with known allelopathic effects on meso- and
microzooplankton grazers[16]. *Chaetoceros* species are abundant in natural diatom communities in the
CCE and the most abundant diatom in the global TARA oceans dataset[17,42–45]. Virus infection is an
important loss term for phytoplankton in the marine environment with mortality rates on par with grazing
across the North Atlantic and Southern Ocean[46,47]. Given other lipids have been used as diagnostic as
biomarkers for viral infection of the *Gephyrocapsa huxleyi*[48], including infection of natural *G. huxleyi*
populations across the North Atlantic[49], we sought to 1) explore oxylipins as biomarkers of viral infection
in situ and 2) test the ecological relevance of oxylipin chemical signaling during viral infection of
diatoms. We took a two-pronged approach, using a laboratory model system with three diatom host-virus
pairs[16,50–52]and analysis of meta-lipidomic data collected from diatom assemblages in the CCE.

Results and Discussion
We previously identified three phytoplankton patches with diatoms populations in different stages of the
bloom lifecycle and viral infection (Active Infection, Post-lytic, and Non-blooming) in a 2013 field study
along the CCE and established a link between viral infection and silicon limitation[17]. Dissolved oxylipins
in the CCE meta-lipidome were more abundant at a Post-lytic sites than at Active Infection sites or a Non-
blooming sites dominated by dinoflagellates. Five of the oxylipin biomarkers from control, laboratory
infection experiments were more abundant *in situ* at the Post-lytic site coinciding with higher bacterial
ectoenzyme activity[17] and higher Si-stress[17,43] compared to the Active Infection site. Deck-board
incubations exposing surface and sinking particle-associated microbial communities to exogenous
oxylipins suggest that they impact grazing, growth, and organic matter cycling by marine microbes.
Finally, sediment traps and satellite data confirm the higher oxylipin post-lytic siters also had higher
magnitude and efficiency of carbon export.

Viral Infection and mortality biomarkers identified in culture experiments
Two diatom species (*Chaetoceros socialis* and *Chaetoceros tenuissimus*) served as hosts in laboratory
infection experiments with three taxonomically distinct viruses: a single-stranded DNA (ssDNA) virus
(CtenDNAV) and a single-stranded RNA (ssRNA) virus (CtenRNAV), both infecting *C. tenuissimus;* and
a ssRNA virus (CsfrRNAV), infecting *C. socialis.* Control and infected treatments were sampled for
lipidomics and cell concentration over 3-4 days post infection (dpi; Fig. 2A, S2). samples were
categorized based on the growth curves and dissolved lipidomes were annotated as described in ref[16]. We
evaluated the general lipidomic response of the three host-virus pairs together to elucidate viral infection
and senescence responses shared across the genus *Chaetoceros.*

The dissolved lipidome (<0.2 μm here) represents small and often oxidized lipids that are more soluble
than the hydrophobic storage lipids or polar membrane lipids. High Resolution Accurate Mass-Mass
Spectrometry (HRAM-MS) paired with reverse phase ultra-high performance liquid chromatography
analysis was used to analyze the dissolved lipid extracts and reference standards (Table S2). To address
specific questions about oxylipin abundance and diversity, we focused on the 156 compounds confidently
annotated as free fatty acid (FFA) precursors of oxylipins and NVOs. Equal weight was given to high and
low abundance compounds by mean-centering and log-transforming the peak area data after normalizing

to the volume filtered and the recovery of an internal standard (Supplemental File 1). A comparative -
omics scheme[53,54] was applied to the resulting dissolved oxylipin profiles using hierarchical clustering
analysis and multivariate analysis to determine the significant biomarkers of viral infection, stress, and
death.
128

Figure 2. Virus infection of diatoms leads to the release of oxylipins and free fatty acids into the dissolved
organic matter pool. A) Experimental design showing representative growth curves of the host-only controls and
representative time course of infection for three diatom host-virus pairs: The approximate sampling timepoints are
noted with red asterisks. The actual growth curve of each triplicate treatment can be found in Fig. S2. **B)** Box and
whisker plots of the average peak area of oxylipins and free fatty acids in the dissolved lipidome for replicates
across the *C. tenuissimus* (2-10) and *C. socialis* (L4) host-only control time point and the phases of infection
averaged across all three viruses. The edges of the box indicate the 1st to 3rd quartile range; error bars represent the
standard deviation with outliers noted by dots. **C)** Heatmap of the oxylipidome (Oxylipins + FFA) from the
Chaetoceros viral infection experiments showing the relative abundance of each compound relative to its mean
concentration across all samples with red indicating a relative increase and blue a relative decrease, reported as a log
fold change. The samples on the X-axis and compounds in the Y-axis are ordered based on the dendrogram created
using Ward clustering. Important clusters of samples that are discussed in the text are labeled on the dendrogram.
143

We found that *Chaetoceros* produced the highest absolute abundance of FFAs and NVOs (*i.e.*, non-PUA
oxylipins) during the later stages of viral infection, when cells are undergoing lysis and during the onset
of stationary phase in controls (especially in *C. socialis* controls; Fig. 2B-C). Hierarchical clustering
analysis split the data into two clusters revealing distinct lipidome structures between the cells lysing due
to viral infection (Cluster 2) and the aging, control treatments (Cluster 1; Fig. 2C). Within Cluster 2, 15
of 22 samples are virally infected, and within Cluster 2bi, 10 of 10 samples are *Chaetoceros tenuissimus*
from mid infection and lysis. This suggests the *C. tenuissimus* lipidomes from virocells are more similar
to one another than other species infected with similar viruses.

Triplicates from two controls grouped with the T= 3 and 4 dpi virus samples, stationary phase C. socialis
and declining C. tenuissimus from a different strain (2-6) than the CtenDNAV and CtenRNAV host (Ct 2-
10). The declining host samples grouped with the declining *C. socialis* in Cluster 1. Again *C. tenuissimus*
(2-6) was not used as a host in our virus infection experiments but was included in our metanalysis to
serve as an outgroup. *C. tenuissimus* (2-6) had a unique lipidomic signature in decline phase, grouping
with the stationary phase *C. socialis* controls and the virus infected treatments in Cluster 2 but forming its
own subgroup (Fig. 2C, Cluster 2a.). *C. socialis* showed elevated oxylipins in stationary phase with an
overall lipidome similar to CsfrRNAV infected cells (Fig. 2C, Cluster 2bii samples labeled C. soc 3 dpi).

Within Cluster 1, 23 of 29 samples are controls. The only virus infected treatments in Cluster 1 are early
infection samples (T= 1 dpi are 6 of 29 samples). The declining samples from the *Chaetoceros* hosts (Ct
2-10 and Cs) clustered with the Stationary phase *Ct 2-10* samples showing lower relative abundances of
most oxylipins observed in the experimental dissolved meta-lipidome (Fig. 2C- Cluster 1b).

The observed strain-level variability in the relative abundance and chemical diversity of oxylipins
produced was consistent with Wichard *et al.*, which surveyed PUA production in 50 strains of
phytoplankton (primarily diatoms) and found strain level variation[16,22]. Ribalet *et al.* had previously
established that diatoms from *Skeletonmena* and *Thalassiosira* increase the production of free fatty acid
and PUAs as they reach stationary phase and even more so under nutrient stress, particularly Si-stress[14,15].
Our method was not optimized for PUA quantification but rather for the pool of less volatile compounds
adjacent and upstream of PUA production (Fig. 1A), showing plasticity in the range of non-volatile
oxylipins produced by *Chaetoceros* diatoms under different growth conditions.

Fig 3. Biomarkers for virus infection and general cell death of diatoms A) Loadings of the dissolved lipidome samples on Component 1 and 2 of the sparse PLSDA. Control samples are represented by a colored circle corresponding to the time of sampling (T=0, exponential, early stationary, stationary, and decline) and infected treatments are represented by pink triangles of various shades.   show growth phase and viral infection lead to distinct changes in the dissolved lipidome. **B)** Component 1 separates the viral infection lipidomes from the decline lipidome **C)** Component 2 captured the signal from cell death more generally. **B-C)** The significant compounds assigned to Components 1 and 2 by the sPLS-DA, their loadings on either component are shown as a dot-plot and the heatmap shows the average relative abundance of each compound across each timepoint.

Sparse Partial Least Squares - Discriminant Analysis (sPLS-DA) was used to distill the dataset down to the twenty most distinctive features that best differentiate time-points and treatments from each other, in hopes of identifying discrete biomarkers for diatom bloom decline by viral infection or general cell death (Fig. 3). Compounds that were most abundant during lysis and that increased over the course of infection had the highest loading on Component 1 (Fig. 3B). The strongest of the viral biomarker were the compounds that were only abundant in the control on the transfer day T=0 and increased with viral infection (NVO 20:2 +1O, NVO 22:4 +1O, NVO 20:3 +1O, and NVO 14:2 +1O; Fig. 3B). Compounds that were more associated with a general death response (*i.e.*, high during decline of the control and during lysis of the infected cultures) loaded most significantly on Component 2 (Fig. 3C).

Twelve of the twenty significant oxylipins were close analogs of known allelopathic compounds produced by diatoms that influence copepod reproductive success in the Mediterranean Sea and microzooplankton grazing in the laboratory[8,10,25,33] (Table S3). These results suggest viral infection triggers a pulse of chemical signaling via oxylipins that is distinct from the shared suite of oxylipins produced by dying, uninfected cells. This chemical signaling could impact trophic interactions and the associated biogeochemistry. To test this hypothesis, we analyzed a dissolved lipidome dataset from a field study in the California Coastal Ecosystem (CCE), where we previously identified different stages of viral infection and conducted deck board amendment experiments across different communities using a range of oxylipin concentrations.

Bloom decline and viral infection linked to oxylipin production and efficient export *in situ*

Biogeochemical evidence of bloom decline and enhanced export
The DYEatom cruise (June 27th – July 6th, 2013; *R/V Point Sur*) occupied nine stations within the CCE upwelling zone. The physical oceanography, phytoplankton composition, Si-stress, and infection state at our study site have been previously characterized[17,43]. Briefly, near-shore stations outside Monterey Bay were dominated by diatoms, particularly *Pseudo-nitzschia*, and were characterized as a population in Active Infection at the time of sampling. Near-shore stations by Point Reyes were also diatom-dominated but were more diverse in species composition (see Fig. 1c in ref.[17]); they were further characterized as having undergone a recent mortality event (Post-lytic). Here, we present an analysis of the dissolved meta-lipidome for these two contrasting regions, along with two offshore stations that were dominated by dinoflagellates that serve as functional controls (Fig. 4, 5, S3).

Mean net primary productivity (NPP) in the region during our occupation was derived from satellite data and the CAFE algorithm[55] (Fig. 4B-C). This analysis showed that we sampled an Active Infection at the Monterey Bay sites (Fig. 4E), coincident with a peak in NPP (Fig. 4B). The Post-lytic Point Reyes site had the lowest NPP at the time of sampling, consistent with a decline in productivity and diatom bloom

222 demise. *In situ* measurements of chlorophyll, a common proxy for phytoplankton, further supported the
223 distinction between sampling sites with concentrations between 8-18 μg L^{-1} in Monterey Bay and between
224 3.9- 7.5 μg L^{-1} at Point Reyes[43].
225

226
Figure 4. Diatom-virus dynamics, net primary productivity, and export signatures. A) A map of sampling sites
during the DYEatom cruise (June 27 -July 6, 2013) and associated net primary productivity (NPP; dark green
gradient) integrated over 8-day bins from June 18, 2013, through July 12th, 2013 using the CAFE Model[52] of
MODIS satellite data. Open circle symbols indicate Monterey Bay stations. Open triangle symbols indicate stations
at Point Reyes. The stations marked with an 'x' indicate non-bloom site outside the upwelling filaments. Time
course of the mean ocean NPP within the 1° latitude by 1° longitude rectangles drawn around **B)** Monterey Bay and
C) Point Reyes. Filled symbols represent when Monterey Bay (cyan circle) and Point Reyes (coral triangle) were
sampled. **D)** Particulate organic carbon (POC; black bars) and particulate organic nitrogen (PON; grey bars scaled to
C by multiply by 6.6) flux to 50 m, as determined using sediment traps deployed at four stations over different
stages of bloom decline. The blue text above each black bar indicates the export ratio calculated from the POC data
and the NPP satellite estimates. **E)** Data from Kranzler *et al.* (2019)[17] showing cell-associated diatom virus and
extracellular virus copy numbers for three contigs identified as diatom RNA viruses similar to those in the culture
experiments from Fig. 3[50,51].

Sediment traps were used to quantify carbon export out of the mixed layer (to 50 m) and calculate export
efficiency as an apparent e-ratio (export at $Z_{euphotic}$/NPP; Fig. 4D). The carbon flux in Monterey Bay was
similar to the offshore, non-diatom dominated station (24.7 and 29.3 mg m^{-2} d^{-1}, respectively), although
the latter had an ~2-fold higher e-ratio of 0.032 (compared to 0.017). The highest carbon flux (46 mg m^{-2}
d^{-1}) and highest e-ratio (0.093) were observed at the lytic Pont Reyes site on July 4th (St. 10), suggesting
export efficiency increased as the viral infection and bloom declined progressed. The lowest measured
carbon export flux was two days prior in this location (at St. 8; also a lytic site) reflecting the episodic
nature of carbon export[56–59]. Unlike St. 8, St. 10 still had some indication of intracellular virus (Fig. 4E)
which could also help explain the differences in export. St. 10, nearer to the origin of upwelling, was a
little earlier on the lytic process with higher, more efficient carbon export. St. 8 was further offshore and
further along in bloom demise, presumably post export event, evidenced by the removal of viruses from
the free-living fraction as well as the cellular fraction presumably by adsorption onto particles (a known
sink for viruses from the water column[60]).

We note that the type of sediment trap deployed, a surface tethered net trap, was not as quantitative as a
particle interceptor trap (PIT) array or neutrally buoyant trap in determining quantitative fluxes[61];
consequently, we likely underestimate export. Given our net traps were used across all stations, these
caveats (high precision, low accuracy) should be uniform allowing for intercomparisons of export
between sites. The highest e-value measured (0.093) aligns with regional estimates from Henson *et al.*
(2011) derived as a function of SST[62,63] although it is lower than an estimated global average e-ratio of
0.17[62] and regional estimates from other models based on satellite color and food web dynamics. Again,
we believe the intercomparison to be robust if an underestimation of the absolute magnitude.

Dissolved meta-lipidome reveals abundance of death and virus biomarkers at the Post-lytic sites
Within the dissolved meta-lipidome from the DYEatom cruise, 175 compounds were annotated as
oxylipins or free fatty acid precursors (Supplemental File 2). The post-lytic site had a higher relative
abundance of dissolved organic lipids compared to the Active Infection or non-bloom sites, especially on
July 4th (St 10; Fig. 5A).The Active Infection and Post-lytic lipidomes were more similar to one another
than to the non-bloom lipidomes, clustering together in the sample dendrogram (Fig. 5A). sPLS-DA was
used to identify the most significant features in the dissolved lipidome for differentiating between the
bloom states. Component 1 represented 55.1% and Component 2 represented 19.7% of the respective
variability in the data (Fig. 5B), with the former containing compounds significantly associated with the
Post-lytic stage of infection (Fig. 5C) and the latter compounds more associated with Active Infection
(Fig. 5D).

Three general death biomarkers and two viral lysis biomarkers from the *Chaetoceros* lab experiments
were amongst the most significant compounds associated with the post-lytic site (Fig. 5C, bold and red
compounds, respectively; Table S5). The general death biomarkers 9-hydroxy hexadecadienoic acid (9-
HHDE), 9-hydroxy hexadecaenoic acid (9-HHME), and 9-oxo hexadecaenoic acid (9-oxoHME) all
belong to the 9-LOX pathway acting on 16:n fatty acids previously observed in *Skeletonema marinoi* and
Thalssiosira rotula (Table S3). The viral lysis biomarkers, 20:4 +1O and 22:5 +1O, were too low
abundance to gain positional information from ms^2 fragmentation but our retention time model indicates
these oxylipins are hydroxy acids similar to those made by *Tr*, *Stephanopyxis turris*, and *Lepidocylindrus*
(Table S3; Supplemental File 3).

The most significant compounds in the dissolved lipidome from the Active infection site were eight
highly oxidized C22:n oxylipin isomers, the FFA and oxylipin precursor, hexadecatetraenoic acid
(C16:4), and a C18:4 oxylipin (Fig 5D). While less oxidized DHA oxylipins, 22:5 +1O and 22:4 +1O,
were prominent viral infection biomarkers in the *Chaetoceros* lab infection experiments (Fig. 3B), the
highly oxidized +3O and +4O compounds associated with the Active infection site are unique for marine
systems (Fig 5D). However, C22:n oxylipins alone may not be specific enough to be indicators of viral
infection. C22:n biosynthesis pathways have been studied in *Lepidocylindrus* diatoms showing that, in
response to sonication, 15- and 18-lipoxygenases are employed to make +1O and +2O C22:n oxylipins[64]
which constitute about 20% of the total NVO pool from January to December 2017 at the LTER-
MareChiara in the Mediterranean Sea[25]. We previously observed C22:6 +2O oxylipins produced in
culture experiments feeding *Phaeodactylum tricornutum* to the dinoflagellate *Oxyrrhis marina*[33]. The
highly oxidized nature (+3O and +4O) of the Active Infection C22:n oxylipins may be correlated to the
stress-level of the community and should be further investigated as biomarkers for infection vs. +1O and
+2O oxylipins which may be more general stress biomarkers.

300
Figure 5. Oxylipin biomarkers of infection state from the DYEatom meta-lipidome. A) Heatmap showing the
relative abundance of 175 compounds identified as free fatty acids and oxylipins in eleven dissolved lipidome

samples from the DYEatom cruise. Samples on the X-axis (lipid compounds) and Y-axis (station and depth) are
organized by similarity depicted by the Ward clustering in the dendrogram. **B)** Loading of the dissolved meta-
lipidome samples on Components 1 and 2 derived from sPLS-DA for samples collected at Point Reyes (coral
triangles), Monterey Bay (cyan circles), and non-bloom sites (grey x). The 95% confidence intervals (colored
ellipses) for the three bloom states do not overlap. Component 1 (55.1% of the variability in the data) distinguishes
the post-lytic samples from the other two categories; Component 2 (19.7% of the variability in the data) describes
active infection. **C- D)** The ten compounds from the dissolved meta-lipidome used to derive **C)** Component 1 and **D)**
Component 2 in the sPLS-DA. Compounds are ordered by their loading as noted by the dot-plot. Heatmaps show the
average relative abundance of each compound in the dissolved meta-lipidomes from each bloom state. Compound
annotations in red text are biomarkers of viral infection in *Chaetoceros* cultures (Fig. 3B), while those in bolded
black text are more general cell death biomarkers (Fig. 3C).
314

Oxylipins function as chemical signaling across trophic levels in the California Current
Similar to the compounds pulled out of the laboratory experiment, the NVOs associated with Post-lytic
stage of infection at Point Reyes are also known to have allelopathic effects on microzooplankton grazers
in culture and copepods at sea[8,10,33] (Table S5; Fig. 5C). An isomer of the NVO 20:4 +3O oxylipin,
annotated as a 5-epoxy-8,14-dihydroxy eicosatrienoic acid, was associated with dinoflagellate grazing
deterrence and chronic stress in *P. tricornutum* experiments. Furthermore, the C16:1, C16:2, C20:4 and
C20:5 oxylipins from the Post-lytic site have been observed in diatom blooms in the Adriatic Sea that are
detrimental to copepods[8,33].
323

Unlike grazers, the effects of oxylipins on eukaryotic phytoplankton and bacteria have only been widely
studied using commercially available PUAs not NVOs[23,27,28,30,34,35]. NVOs are upstream in the oxylipin
biosynthetic pathway from PUAs (Fig. 1A) and tend to be toxic at lower doses than PUAs when tested
against microzooplankton grazers in the laboratory[33] and copepods at sea[8,10,13]. To assess the role of
NVOs and PUAs in signaling across trophic levels, we conducted deck-board incubations of the surface
assemblages and sinking particle-associated microbes treated with exogenous PUAs and NVOs (Fig. 6)
and assessed trends in oxylipin concentration compared to metrics of microbial metabolism across the
cruise.
332

Negative feedback loop between oxylipins and grazing
A mixture of PUAs—heptadienal, octadienal, and decadienal—had differential effects on natural
microbial communities from the California Current. PUAs inhibited microzooplankton grazing and
phytoplankton growth (Fig. 6A-C) but stimulated the activity of different hydrolytic enzymes (peptidase
and phosphatases) in surface communities and sinking particles (Fig 6D-H). A map of grazing, growth,
and a proxy for oxylipin concentration (total peak area of oxylipins and FFA in the dissolved lipidome)
across the study site showed the highest oxylipin concentrations co-occurred with the lowest grazing rates
and the lowest phytoplankton growth rates (Fig. S4), consistent with laboratory-based interactions[33].
341

The relationships between oxylipin concentration, grazing and growth across the cruise sites reflect the
dose dependence of the allelopathic effects of oxylipins and the community's threshold of response to
these chemical cues. Excluding the highest oxylipin concentration, a positive relationship between the
oxylipin concentration proxy and phytoplankton growth was observed (Fig. S5 C). The highest oxylipin
concentrations occurred at the lytic Point Reyes site where the oxylipin threshold for inhibiting
phytoplankton had been crossed, resulting in negative growth rates (Fig. S4 A, D).
348

A negative feedback loop between grazing and oxylipin production was observed, whereby grazing
induces oxylipin production and oxylipins inhibit further grazing (Fig. S5 D). The lowest oxylipin
concentrations were observed at stations with low grazing rates. Subsequently, a negative correlation
between grazing and oxylipin concentrations was observed for the four stations with the highest oxylipin
concentrations once the threshold concentration for allelopathy had been surpassed (Fig. S5 D). Several
studies have explored the synergy amongst various oxylipins whereby having multiple oxylipins present
leads to a lower effective concentration[11,13,65]. Thus, the higher diversity of lipids at the lytic sites (Table
S6 and S7) may also contribute to the lower growth and grazing rates compared to the other sites (Fig.
S4).
358
Dose-dependent PUA signaling in surface and sinking particle-associated microbial communities
We tested the sensitivity of surface and sinking particle-associated microbial communities to oxylipins by
exposing them to a range of concentrations based on the threshold responses observed in the literature
(Fig. S1; Materials and Methods). Depending on the bloom state and community, the response to
exogenous oxylipins varied with shifting effective concentration thresholds and sign of response (negative
vs. positive). At the Active Infection and Post-lytic sites, where background oxylipin concentrations were
relatively high, addition of 100 μM PUAs to sinking particle-associated communities increased
phosphatase activity and inhibited peptidase activity (Fig. 6F,H). The communities on particles collected
from the lower-oxylipin dinoflagellate-dominated, non-bloom site (St 6) were inhibited for both enzymes
after exposure to 100 μM PUA but stimulated by a 10 μM addition (Fig 6G). Thus the microbial
communities that had prior exposure to oxylipins were more tolerant to exogenous oxylipin additions.
Our findings are in agreement with experiments conducted across the Pacific in the South China Sea
showing stimulation of particle-associated communities in response to 10-100 μM PUAs[34]. The threshold
of inhibit was lower in particle-associated communities from the Atlantic, where inhibition of enzyme
activity, respiration, and bacteria growth was observed at 100 μM[35] and stimulation between 1 and 10 μM
PUA addition. The community dependent response to PUAs—inhibition of both peptidase and
phosphatase activity on particles collected from non-blooming sites at 100 μM (Fig 6G) —is consistent
with a screening of 33 bacterial isolates showing stimulation only amongst strains known to be epibionts
of diatoms[22]
378
In surface microbial communities from the Active Infection site, peptidase activity was stimulated and
phosphatase activity was inhibited at a 10 μM PUA dosage. Unfortunately, we did not replicate the 10 μM
PUA treatment in the other experiments to explore whether non-blooming communities are more
sensitive to oxylipins and if post-lytic surface communities are equally stimulated. This is the first report
of oxylipins stimulating natural assemblages of surface ocean microbes. A study conducted with natural
bacterial communities from the NW Mediterranean found a significant decreases in the single cell activity
for a subset of the community in response to nM concentrations of PUAs[66.] Another study testing daily 1-
2 μM PUA additions in 21-day mesocosms found no effect on microbial composition in response to
PUAs alone but did see a difference in bacterial community structure with time and in the two mesocosms
that ended up with different strains of *Skeletonema* blooming after nutrient addition[67]. We previously
reported enhanced peptidase activity at the Post-lytic site likely reflecting the stimulatory effect of
oxylipins on enzyme activity. Culture studies show enhanced peptidase activity in infected *Chaetoceros*
cells compared to control diatom cells and more peptidase activity but less growth in bacteria exposed to

CtenRNAV lysate compared to CtenDNAV lysate[20], introducing the caveat that stimulation may not
translate to growth or secondary production by the bacterial community.

Figure 6. Microzooplankton grazers, phytoplankton, and both the surface and the particle associated
heterotrophic microbial communities exhibit dose- and community-dependent responses to specific PUAs and
non-volatile oxylipins. A-C, I) Fold-change in growth rate of phytoplankton (white) and grazing rate of
microzooplankton (black) for control (untreated) and oxylipin amendments at seven stations. The mean and standard
deviation are shown (n=3). Asterisks denote a statistical difference from the control ($p < 0.05$ Mann-Whitney U).
Shaded panels denote statistically insignificant results. **D-E, J)** Enzymatic activity of the surface and **F-H, K)**
sinking particle-associated microbial communities in response to varying concentrations of a PUA mixture **(D-H)**
and the oxylipin 15-hydroxyeicosapentaenoic acid (15-HpETE; **I-K**). Alkaline phosphatase activity is in grey and

white stripe and peptidase activity is in solid grey. Values are normalized to the triplicate controls. Asterisks denote
a statistically significant difference from the control ($p < 0.1$ Krushal-Wallis with a $p < 0.05$ Dunn's post-hoc).
405
Exploring the bioactivity of NVO vs PUAs
As mentioned above, NVOs are thought to be more deleterious to grazers than PUAs. For example,
dinoflagellate grazing on diatom cultures is inhibited by 50% at 1 nM NVO but exposure to 10 nM PUA
only decreased grazing by 33%; PUA concentrations in the 1000 nM range were needed for a 50%
reduction in grazing[33]. To better understand the response to NVOs vs. PUA across multiple trophic levels,
a set of grazing-growth experiments (Fig. 6I), two small sets of surface microbial enzymatic activity
experiments (Fig. 6J), and one set of particle-associated microbial enzymatic activity experiments (Fig.
6K) were conducted with the NVO, 15-HpETE replicating the PUA experiment present in the previous
section. The grazing experiment with 15-HpETE (Fig. 6I) displayed large variability amongst the
triplicates and a negative growth rate in the control and was inconclusive (Fig. S4D). Enzymatic activity
was stimulated by NVOs in both surface and sinking particle-associated microbial communities (Fig. 6J-
K). The stimulation of the surface community was more significant for NVOs than PUAs at 1 µM (Fig. 6J
vs. 6E; St. 7). Similarly, particle-associated community responses to 10 µM PUAs and NVOs additions
were observed at the lytic Point Reyes sites (St. 8 and St.10, respectively; Fig. 6H and 6K). We had a
limited amount of 15-HpETE stock with us so we were restricted to a smaller number of experiments and
a limited range of concentrations tested on this cruise. Additional parallel experiments with NVOs and
PUAs are needed to fully understand dose-dependent impacts of oxylipins on the same population.
Nonetheless, our results are congruent with studies exploring the effect of various oxylipins on copepods,
sea urchins, and dinoflagellate grazing, which suggest that the range of effective concentrations varies by
oxylipin[8,10,12,33,65,66].
426
Notably, 15-HpETE, the compound tested in deck-board incubations, was observed in the DYEatom
cruise meta-lipidome (Fig. S6A). HpETE analogs (*i.e.* compounds annotated as NVO 20:4 +2O at all
retention times) were present in significantly higher concentrations at the lytic Point Reyes sites compared
to mid infection or non-bloom states (Fig. S6B). HpETE analogs represented less than 1% of the average
annotated peak area (Fig. S7). C20:5 and C16:n oxylipins and their free fatty acid precursors dominated
the dissolved lipidome representing 58% of the peak area (Fig. S6). Given the dose dependence of
oxylipin bioactivity, oxylipin quantification is critical to understanding and contextualizing their
bioactivity *in situ*. Our general untargeted approach better captured the wide diversity of compounds
present but it only allows for presentation of relative abundances. Future work should take advantage of
tractable absolute quantification with an adequate suite of representative authentic standards.
437
Biogeochemical impacts of oxylipin chemical signaling during viral infection
We previously reported enhanced ectoprotease activity at the lytic Point Reyes site, which we interpreted
as stimulation of remineralization and the 'viral shunt'[17]. Deck-board experiments in this study revealed
oxylipin-inducible enzyme activity in surface and sinking particle-associated microbial communities
further connecting virus infection (and the induced chemical signaling) with enhanced organic matter
turnover. Silicon dissolution is critically controlled by proteolytic degradation of the organic matrix
protecting diatom frustules, denuding and exposing the opal (SiO_2) to the undersaturated seawater[68]. In
turn, this dissolved silicic acid ($Si(OH)_4$) can support regenerative biogenic silica production[69]. Enhanced
silica dissolution can also negatively impact mineral ballasting of exported carbon and diminish the

oceanic silica pump[70]. We conducted experiments to test whether oxylipin enhanced enzymatic activity
impacted biogenic silica dissolution, but it was inconclusive (data not presented). The stimulation of
ectopeptidase in surface microbial communities has been observed in diatom blooms in this region[70]. The
mechanisms underlying peptidase stimulation during viral infection of *Chaetoceros*[20] are not well
understood but thought to be linked to the production of nitrogen-rich, Coomassie Stainable Particles
(CSP), which have been observed in virus-infected *Chaetoceros* culture studies[71]. Stimulation of
peptidase over phosphatase in the surface ocean would serve to preferentially remineralize nitrogen from
dissolved organic matter and is consistent with nitrogen being the colimiting nutrient in the sunlit CCE
along with Fe rather than phosphorus[72]. A preference for phosphatase stimulation on particles reflects a
shift in community activity towards accessing organically bound phosphate in response to oxylipin
chemical signaling below the euphotic zone.

At the same time, our geochemical data (Fig. 4A-C) strongly suggest that virus infection of this natural
diatom community induced the viral shuttle and enhanced carbon export even with enhanced enzymatic
turnover of organic material in surface communites[17]. Export was positively correlated with the total
peak intensity of oxylipins in the dissolved lipidome (Fig. S5A), suggesting that oxylipins may play a role
in enhanced export during the viral shuttle. The relative balance between shunt versus shuttle outcomes
will ultimately be determined by the remineralization length scales relative to the mixed layer depth (see
Figure 5 in ref[20]). Here, virus infection stimulated export flux and high e-ratios to 50 m; the observed
MLD at this station was 10 m, indicative of a short remineralization length scale driving net outcome
towards the shuttle.

Evidence of a strong viral shuttle was previously observed in coccolithophores populations that were
actively infected with giant, double-stranded DNA-containing Coccolithoviruses in the North Atlantic[59];
diagnostic lipid biomarkers[48,73,74] showed that sinking aggregates were enriched with infected cells and
resulted in very high POC and PIC fluxes. Similarly, eukaryotic viral marker genes in a meta'omic study
of the TARA Oceans Expedition global dataset explained 67% of the variability in carbon export
efficiency with 58 viruses positively correlated and 25 negatively correlated with export efficiency[19].
Many of the positively correlated viruses were linked to silicifying eukaryotes, like diatoms.
Mechanistically, culture studies with eukaryotic phytoplankton, including diatoms and coccolithophores,
have shown that virus infection enhances spore formation[75], TEP and CSP excretion, and particle
aggregation [71,76]. Oxylipins are known to increase aggregation in the benthic diatom *Fistulifera*
saprophila[77] but these effects are strain dependent.

In addition to remineralization length scales of sinking particles, population dynamics and have been
proposed as controls on whether viral infection leads to enhanced export or enhanced recycling. Large
scale mesocosm experiments show a 2-4x increase in TEP and PIC production per cell for *G. huxleyi*
during infection feeding into the viral shuttle and resulting in adsorption of viruses from the water column
on to aggregated particles. Taken with the data presented here and the conceptualization of larger
phytoplankton being associated with more zooplankton degradation and higher overall fluxes, we
hypothesize that eukaryotic algae are more subject to the viral shuttle compared to prokaryotic hosts. The
length scale plays into this as eukaryotic phytoplankton sink faster than prokaryotes as a function of size
and added density due to their associated biominerals, thus the length scale over which their biomass can
be remineralized is shorter leading to more efficient carbon export. Our analysis adds a layer of

complexity to the population dynamics control on viral carbon cycling with oxylipin-producing diatom
populations altering organic matter recycling, grazing, growth, and export. The dose dependent nature of
oxylipin signaling may contribute to episodic export once inhibitory thresholds for the community are
surpassed, resulting in a toggle between shuttle and shunt (Fig. 7).

495
Figure 7. Oxylipin chemical signaling during virus infection and bloom demise toggles the fate of organic
matter between export and remineralization. Conceptual link between the state of virus infection, oxylipin
production, export flux, remineralization and remineralization length scale in a natural diatom population. **A)**
Communities undergoing active infection (such as those surveyed near Monterey Bay on June 28[th]) were at the peak
bloom NPP, as per satellite measurements and *in situ* chlorophyll accumulation. Some oxylipins are detected in the
environment but at lower levels and were characterized by a different suite of compounds. **B)** Lytic populations
(such as those surveyed at Point Reyes) had higher dissolved oxylipin concentrations, including those specifically
viral biomarkers from culture infection studies, along with decreases in NPP over the proceeding 8 d, higher export
flux and e-ratios of carbon to 50 m (compared to Monterey Bay), and higher ectoenzymatic activities. Taken
together, these data suggest that virus infection and oxylipins induced the viral shuttle and enhanced carbon export
even with enhanced turnover of organic matter remineralization indicative of the viral shunt[17].

Conclusions
Several, specific oxylipins were identified as biomarkers for viral infection and cell mortality in cultured
Chaetoceros sp. and were subsequently found in the meta-lipidomes of virus-infected natural diatom
populations infected California Current Ecosystem. Deck-board incubations agreed with microbial
responses to oxylipin observed in previous studies, namely dose-, compound-, and community-dependent
inhibition of grazing and phytoplankton growth and stimulation of particle-associated microbial
communities. Oxylipins stimulated phosphatase activity on sinking particles and peptidase activity in
surface communities, while inhibiting microzooplankton grazing and phytoplankton growth.
Biogeochemical measurements suggest that virus-mediated diatom bloom decline led to higher and more
efficient carbon export compared to earlier phases of infection, despite stimulatory effects on enzymatic
remineralization of organic matter. Taken together, these findings highlight how viruses and chemical

signaling via oxylipins mediate trophic interactions and biogeochemical fate of organic matter in
productive upwelling regimes.
521
522   **Materials and Methods**
523
Viral Infection Experiments with Cultured Chaetoceros sp.
Cultures of the diatom hosts *C. socialis* strain NIES-3713 (formerly L-4) and *C. tenuissimus* strains
NIES-3714 (formerly 2-6) and NIES-3715 (formerly 2-10) were grown in SWM-3 media at 15°C under
150 µmols photons m^{-2} s^{-1} irradiance on a 12:12 L:D photoperiod. Triplicate cultures were infected during
exponential growth with CsfrRNAV (*C. socialis*), CtenRNAV (*C. tenuissimus* NIES-3715) or
CtenDNAV (*C. tenuissimus* NIES-3715). Samples were collected for cell counts via microscopy or flow
cytometry. For experiments with CsfrRNAV and CtenRNAV, samples for dissolved lipidome analysis
were collected at 0, 1, and 3 dpi, representing early, mid, and late infection, respectively (Fig. S2). For
experiments with CtenDNAV, samples for dissolved lipidome analysis were collected 0, 1, 3, and 4 dpi.
Triplicate uninfected control cultures were sampled concurrently to exclude changes during normal
growth. Follow-up experiments grew the two hosts, along with *C. tenuissimus* NIES-3714 into decline
phase to assess changes associated with death (likely induced by nutrient limitation). Samples were pre-
filtered through a 0.2 µM Durapore filter to remove particles. The deuterated internal standard, 15(S)-
Hydroxy eicosatetraenoic acid-d8 (Cayman Chemical), was added to each filtered sample to a final
concentration of 10 µM. The dissolved lipids were then extracted onto a Waters HLB SPE cartridge.
539
Lipidomic Analysis
The dissolved lipids were eluted from the SPE cartridge with 2 ml of 70:30 acetonitrile: isopropanol into
pre-combusted collection vials that were treated with the antioxidant BHT to prevent autoxidation of the
samples. The samples were immediately transferred from the collection vials into HPLC vials and capped
under argon. The samples were stored at -80°C until mass spectrometric analysis. Within a week of
extraction the dissolved lipidome samples were analyzed using reverse phase HPLC (Agilent 1200
system; Agilent, Santa Clara, CA, USA) paired with high resolution, accurate mass (HRAM) data from a
Thermo Exactive Plus Orbitrap mass spectrometer (ThermoFisher Scientific, Waltham, MA, USA). The
chromatographic method used a Xbridge C8 column (Waters, Milford, MA, USA) as the stationary phase,
18 MΩ water as eluent A, 70:30 acetonitrile: isopropanol as eluent B, and ammonium acetate and acetic
acid as the adduct forming additives[70]. The gradient began with a 1-minute isocratic hold of 45% B and
shifted to 99% B over 25 minutes with a flow rate of 0.4 mL min^{-1}. Full scan data (m/z 100-1500) was
collected in negative mode at a mass resolution of 170,000. For the follow-up experiments and to annotate
the oxylipins in the original Chaetoceros and DYEatom cruise samples, the dissolved lipid extracts were
analyzing using reverse phase chromatography (Vanquish UHPLC, ThermoFisher Scientific, San Jose,
CA, USA) paired with HRAM data-dependent acquisition on a Orbitrap ID-X mass spectrometry
(ThermoFisher Scientific, San Jose, CA, USA). A mass inclusion list with the exact masses of [M-H]$^-$
adducts of important oxylipins was used to acquire ms^2 for further annotation (*Chaetoceros* experiment
oxylipin annotation described in Edwards et al. 2024; DYEatom annotation in Table S5). For further
details, please refer to the manuscript detailing the intercomparison of diatom host-virus pairs[16].
560
Chemoinformatic Pipeline

The raw mass spectrometric data was converted to mzXML format with mscovert and analyzed with the
R packages xcms[78–80] and CAMERA[81], which aligned the chromatograms, picked and integrated peaks,
and identified and removed secondary isotopic peaks (specific details in Edwards *et al.*, 2024). The
lipidomic features were further annotated using the LOBSTAHS package which included m/z information
of the various acyl chain-lengths, degrees of unsaturation, and oxidations of triacylglycerols, free fatty
acids (FFA), polyunsaturated aldehydes (PUA), and eight intact polar diacylglycerol lipids and their most
common adducts[82]. The features annotated as oxylipins, and free fatty acids were retained and manually
verified as high-quality peaks in El-MAVEN[83–85]. The fragmentation of the compounds was assessed
using the rerun of the 2016 samples on the UPLC-Orbitrap ID-X and in silico databases created by the
Edwards Lab in CFM-ID[86], and iterative database of mammalian oxylipins from Watrouse[87], and the *in*
silico database of algae, plant, and animal lipids available through the LipidBlast pathway in MS-DIAL[88].
Peak areas were normalized to the recovery of the internal standard and volume filtered. Statistical
analyses were performed using the R package MetaboAnalyst through its web interface[54].
575
DYEatom cruise
The DYEatom cruise aboard the R/V Point Sur (PS1312) disembarked from Moss Landing, CA on June
27, 2013. The 9-day cruise track transected the California coastal upwelling region (Fig 4 and Table S4).
Nine stations were occupied. The details of the metatranscriptomic analysis of cell associated viruses and
qPCR of free-living viruses used for assessing viral infection by diatom RNA viruses can be found in
Kranzler *et al.* 2019, where the details of the enzyme activity protocols can also be found[17].
582
Dissolved Meta-lipidome
The dissolved lipidome was sampled at two depths in the water column along the cruise track, the 55% Io
depth and the chlorophyll maximum. One liter of seawater from each depth was filtered through a bottle
top 0.2 μm filter (Corning) to remove particulate material. The internal standard benzaldehyde (Sigma-
Aldrich) was added to a final concentration of 10 μM in one liter of seawater. Using a vacuum manifold,
the samples were filtered onto a Water HLB solid phase extraction (SPE) cartridge for approximately one
hour and the final volume filtered was recorded. The SPE cartridges were acidified with 0.1%
hydrochloric acid, placed in a whirl-pak bag, and stored at -80°C until elution and analysis at Woods Hole
Oceanographic Institution. Samples were analyzed using the chromatographic, mass spec, and lipidomic
pipeline outlined above.
593
Sediment Trap Collection of Sinking Particles
At Stations 1, 6, 8, and 10, sinking particles were collected using a surface tethered net trap deployed at a
depth of 50m for ~8 hrs. Trap material was screened through a 350 μm mesh to exclude larger
mesozooplankton and split into 8 equal 500 mL fractions using a rotating electric splitter[89,90]. One 500-
mL split of the trap material was filtered onto three pre-combusted 47mm glass fiber filters (nominal pore
size- 0.7-μm) for particulate organic carbon and nitrogen (POC/PON) analysis. The samples were kept at
−80 °C until analysis at Woods Hole Oceanographic Institution (Woods Hole, MA) on a Finnigan- MAT
DeltaPlus stable light isotope ratio mass spectrometer[91]. The POC and PON values of the three filters
were summed together for each trap and deployment time was used to calculate flux to 50 m over a 24-
hour period.
604
605 *Net Primary Productivity Estimates*

606   High-resolution, 8-day integrated satellite-based estimates of net primary productivity in the CCE for the
607   month surrounding our cruise were downloaded from the Oregon State University Ocean Productivity
608   website. The NPP estimates come from the CAFE model, which is depth, time, and spectrally resolved[55].
609   Clara Douglas wrote the MatLab script for this analysis.
610
Deck-board Incubations
Three types of amendment experiments were conducted at sea: 1) three deck-board incubation
experiments with surface seawater communities and varying concentrations, of either a mix of PUAs (0,
0.1, and 1μM) or the NVO, 15-HpETE (0 , 0.1, and 1 μM; Fig. 6B), 2) four dark incubation experiment
with sinking particles from 50 m amended with 0 , 1, and 10 μM HpETE or 0, 1, 10, and 100 μM PUA
(Fig. 6C); 3) seven deck-board paired dilution experiments to compare microzooplankton grazing rates
and phytoplankton growth rate in no amendment controls and 1 μM PUA or 0.8 μM HpETE treatments
(Fig. 6A). These concentrations were chosen based on the bioactive PUA concentrations for marine
bacterial isolates[28], PUA concentrations that enhanced the metabolism of marine microbes associated with
sinking particles in the North Atlantic[35], and PUA concentrations that decreased microzooplankton
growth rate in the Chesapeake Bay[32].
622
Stock solutions
The NVO 15-HpETE stock solution was biosynthesized in the lab by adding the enzyme 15-lipoxygenase
(Sigma-Aldrich) to arachidonic acid (Sigma-Aldrich) using the protocol outlined in Iacazio *et al.* (1990)[92].
Once the reaction reached completion, the oxylipins were extracted onto an SPE cartridge (HBL Waters)
and immediately eluted with methanol. Mass spectrometric analysis determined the composition of the
oxylipin stock was 825 mmol L^{-1} 15-HpETE and a background of its precursor and positional isomers.
The oxylipin stock solution was capped under N_2 and stored at -20 °C. In our LOBSTAHs notation
HpETE is FFA 20:4 +2O.
631
632   A stock solution containing a mix of polyunsaturated aldehydes was prepared at sea and contained 100
633   mM of 2,4-heptdienal, 2.4-octadienal, and 2,4- decadienal (Sigma-Aldrich) dissolved in methanol. The
634   solution was kept at 4C and was remade every two days to prevent degradation. In our LOBSTAHS
635   notation these compounds would be PUA 7:2, PUA 8:2 and PUA 10:2.
636
Surface Ocean Community
At Stations 4, 7, and 10, seawater was collected from the 55% I_0 depth into 2 L Nalgene bottles. At St. 4,
0, 0.1, 1 and 10 μM PUAs were added to triplicate bottles using methanol as a carrier liquid. Care was
taken such that the same volume of methanol was added to each treatment including the 0 μM control. At
St. 7, 0, 0.1, and1 μM PUA and 1 μM HpETE were added to bottles in triplicate. At St. 10, we began to
run out of oxylipin stocks so only three conditions were made up in triplicate: 0, 0.1, and 1 μM HpETE.
The 2 L microcosms were placed in on-deck incubators screened to mimic the light field at 55% I_0 and
temperature controlled by a flow through system for ~24 hours. Enzymatic activity was measured at the
end of the experiment (Fig. 6B) using fluorogenic substrates as described in Kranzler *et al.* 2019[17].
646
Particle Associated Community
At the stations where sinking particles were collected, a 500 mL split was used to set up dark incubations
using the experimental design laid out in Edwards et al 2015. In a dark on-deck incubator that was

temperature controlled by a flow-through system, the trap material was incubated in opaque 150 mL
Nalgene bottles in triplicate for 24 hours. At St 1, 6, and 8, the treatments were 0, 1, 10, and 100 μM
PUA. At St. 10, the treatment was 0, 1, or 10 μM HpETE Enzymatic activity was measured at the end of
each experiment (Fig. 6C). Samples were also taken for biogenic silica quantification by filtering 25 mL
onto a 1.2 μm nucleopore filter for analysis at Rutgers (Fig. S8A). The same volume was filtered through
a 0.7 μm nucleopore and frozen for dissolved Si analysis at UCSB (Fig. S8B).

Grazing and Growth Rate- Dilution Experiments
Paired dilution experiments were conducted to measure microzooplankton grazing rates and
phytoplankton growth rates with and without exposure to exogenous oxylipins, at seven stations along the
cruise track (Fig. 6A and S4). At each station, 40 L of water was collected from the depth of 50% surface
photosynthetically active radiation (PAR). Half of this volume was then gravity filtered through a 200 μm
Nitex mesh, to be used as whole seawater (WSW), and stored in an on-deck incubator at *in situ* surface
temperature until the experimental set-up. The other 20 L was gravity filtered through a 20 μm Nitex
mesh, to gently remove large phytoplankton (i.e. diatoms), and then gravity filtered through a Pall 0.2 μm
vented sterile filter capsule, in order to make dilution water. Once the dilution water was prepared, we
used a modified dilution method[93] with three points, using 5, 20, and 100% WSW, in triplicate 1L bottles.
Nutrients were not added to bottles. For treated bottles, a final concentration of 0.8 μM oxylipins were
added using a stock solution prepared in the lab. All dilution bottles were sampled at time 0 and after 20-
24h for chlorophyll, cell counts (1% gluteraldehyde), and FIRe analysis, while only 100% WSW bottles
were sampled at both time points for microzooplankton abundance (10% Lugol's).

Statistical analyses for incubation experiments
For the oxylipin amendment experiments, Shapiro-Wilk tests showed that the data was not normally
distributed and non-parametric rank-order tests were used to determine significant differences between
treatments (Sigma-Plot). A Mann-Whitney U was utilized to test whether microzooplankton grazing rate
and the phytoplankton growth rates from the paired dilution experiments were significantly different in
the no amendment controls compared to the 0.8 μM HpETE treatments. A Kruskal-Wallis analysis of
variance on ranks was used to determine if the enzymatic activity of peptidase and phosphatase was
significantly different in the controls compared to the oxylipin treatments in amendment experiments, as
well as, to determine if the total peak area of annotated free fatty acids and oxylipins was significantly
different between the significant groups of samples revealed by similarity profile analysis ($p < 0.1$).
Dunn's method was used as a post-hoc to determine which groups were different from one another ($p <$
0.05).

Data Accessibility
The dissolved lipidomes are available on the MassIVE repository. Biogeochemical data from the
DYEatom cruise can be found on BCO-DMO (PS1312, http://www.bco-dmo.org/project/550825).

Acknowledgements
The captain and crew of the *R/V Pt. Sur* and the Brzezinski lab group who also sailed with us. We thank
Yuji Tomaru (Fisheries Technology Institute, Japan Fisheries Research and Education Agency) for
providing the *Chaetoceros* diatom host-virus systems. We thank the funding sources that supported this

work: NSF OCE-1155663 (to JWK), Gordon and Betty Moore Foundation (BVM, KDB, MJ), McMinn
Endowed Fund (BRE), NSF OCE- (BRE),

1. Leblanc, K. *et al. Distribution of Calcifying and Silicifying Phytoplankton in Relation to*
Environmental and Biogeochemical Parameters during the Late Stages of the 2005 North
East Atlantic Spring Bloom. Biogeosciences vol. 6 www.biogeosciences.net/6/2155/2009/
(2009).
2. Behrenfeld, M. J., Boss, E. S. & Halsey, K. H. Phytoplankton community structuring and
succession in a competition-neutral resource landscape. *ISME Communications* **1**,
(2021).
3. Jin., X., Gruber, N., Dune, J. P., Sarmiento, J. L. & Armstrong, R. A. Diagnosing the
contributions of phytoplankton functional groups to the production and export of
particulate organic carbon, CaCO3, and opal from global nutrient and alkalinity
distributions. *Global Biogeochem Cycles* **20**, (2006).
4. Benoiston, A. S. *et al.* The evolution of diatoms and their biogeochemical functions.
Philosophical Transactions of the Royal Society B: Biological Sciences vol. 372 Preprint
at https://doi.org/10.1098/rstb.2016.0397 (2017).
5. Bianchi, T. S. *et al.* Centers of organic carbon burial and oxidation at the land-ocean
interface. *Organic Geochemistry* vol. 115 138–155 Preprint at
https://doi.org/10.1016/j.orggeochem.2017.09.008 (2018).
6. Miralto, A. *et al.* The insidious effect of diatoms on copepod reproduction. *Nature* **402**,
173–176 (1999).
7. Ianora, A. *et al.* Aldehyde suppression of copepod recruitment in blooms of a ubiquitous
planktonic diatom. *Nature* **429**, 403–407 (2004).
8. Ianora, A. *et al.* Non-volatile oxylipins can render some diatom blooms more toxic for
copepod reproduction. *Harmful Algae* **44**, 1–7 (2015).
9. Brugnano, C. *et al.* The deleterious effect of diatoms on the biomass and growth of early
stages of their copepod grazers. *J Exp Mar Biol Ecol* **476**, 41–49 (2016).
10. Lauritano, C. *et al.* New oxylipins produced at the end of a diatom bloom and their effects
on copepod reproductive success and gene expression levels. *Harmful Algae* **55**, 221–
229 (2016).
11. Turner, J. T., Ianora, A., Miralto, A., Laabir, M. & Esposito, F. Decoupling of copepod
grazing rates, fecundity and egg-hatching success on mixed and alternating diatom and
dinoflagellate diets. *Mar Ecol Prog Ser* **220**, 187–199 (2001).
12. Ianora, A. *et al.* Low reproductive success for copepods during a bloom of the non-
aldehyde-producing diatom Cerataulina pelagica in the North Adriatic Sea. *Marine*
Ecology **29**, 399–410 (2008).
13. Barreiro Felpeto, A. *et al.* Diatom induction of reproductive failure in copepod: The effect
of PUAs versus non volatile oxylipins. *J Exp Mar Biol Ecol* **401**, 13–19 (2011).
14. Ribalet, F. *et al.* Age and nutrient limitation enhance polyunsaturated aldehyde
production in marine diatoms. *Phytochemistry* **68**, 2059–2067 (2007).
15. Ribalet, F. *et al.* High Plasticity in the Production of Diatom-derived Polyunsaturated
Aldehydes under Nutrient Limitation: Physiological and Ecological Implications. *Protist*
160, 444–451 (2009).

16.   Edwards, B. R., Thamatrakoln, K., Fredricks, H. F., Bidle, K. D. & Van Mooy, B. A. S.
738   Viral Infection Leads to a Unique Suite of Allelopathic Chemical Signals in Three Diatom
739   Host–Virus Pairs. *Mar Drugs* **22**, (2024).
740   17.   Kranzler, C. F. *et al.* Silicon limitation facilitates virus infection and mortality of marine
741   diatoms. *Nat Microbiol* (2019) doi:10.1038/s41564-019-0502-x.
742   18.   Vincent, F. *et al.* Viral infection switches the balance between bacterial and eukaryotic
743   recyclers of organic matter during coccolithophore blooms. *Nat Commun* **14**, (2023).
744   19.   Kaneko, H. *et al.* Eukaryotic virus composition can predict the efficiency of carbon export
745   in the global ocean. *iScience* **24**, (2021).
746   20.   Kranzler, C. F. *et al. Taxonomically Distinct Diatom Viruses Differentially Impact Microbial
747   Processing of Organic Matter. Sci. Adv* vol. 11 https://www.science.org (2025).
748   21.   Jagusch, H., Baumeister, T. U. H. & Pohnert, G. Mammalian-Like Inflammatory and Pro-
749   Resolving Oxylipins in Marine Algae. *ChemBioChem* **21**, 2419–2424 (2020).
750   22.   Wichard, T. *et al.* Survey of the Chemical Defence Potential of Diatoms: Screening of
751   Fifty Species for α,β,γ,δ-unsaturated aldehydes. *J Chem Ecol* **31**, 949–958 (2005).
752   23.   Ribalet, F. *et al.* Phytoplankton cell lysis associated with polyunsaturated aldehyde
753   release in the Northern Adriatic Sea. *PLoS One* **9**, e85947–e85947 (2014).
754   24.   Gerecht, A. *et al.* Oxylipin production during a mesocosm bloom of Skeletonema marinoi.
755   *J Exp Mar Biol Ecol* **446**, 159–165 (2013).
756   25.   Russo, E. *et al.* Density-dependent oxylipin production in natural diatom communities:
757   possible implications for plankton dynamics. *ISME J* **14**, 164–177 (2020).
758   26.   Barbarinaldi, R. *et al.* Prostaglandin pathway activation in the diatom Skeletonema
759   marinoi under grazer pressure. *Mar Environ Res* **196**, (2024).
760   27.   Ribalet, F., Berges, J. A., Ianora, A. & Casotti, R. Growth inhibition of cultured marine
761   phytoplankton by toxic algal-derived polyunsaturated aldehydes. *Aquatic Toxicology* **85**,
762   219–227 (2007).
763   28.   Ribalet, F., Intertaglia, L., Lebaron, P. & Casotti, R. Differential effect of three
764   polyunsaturated aldehydes on marine bacterial isolates. *Aquat Toxicol* **86**, 249–255
765   (2008).
766   29.   Vardi, A. *et al.* A Stress Surveillance System Based on Calcium and Nitric Oxide in
767   Marine Diatoms. *PLoS Biol* **4**, e60- (2006).
768   30.   Pepi, M. *et al.* Toxicity of diatom polyunsaturated aldehydes to marine bacterial isolates
769   reveals their mode of action. *Chemosphere* **177**, 258–265 (2017).
770   31.   Franzè, G., Pierson, J. J., Stoecker, D. K. & Lavrentyev, P. J. Diatom-produced
771   allelochemicals trigger trophic cascades in the planktonic food web. *Limnol Oceanogr* **63**,
772   1093–1108 (2018).
773   32.   Lavrentyev, P. J., Franzè, G., Pierson, J. J. & Stoecker, D. K. The effect of dissolved
774   polyunsaturated aldehydes on microzooplankton growth rates in the Chesapeake Bay
775   and Atlantic coastal waters. *Mar Drugs* **13**, 2834–2856 (2015).
776   33.   Johnson, M. D., Edwards, B. R., Beaudoin, D. J., Van Mooy, B. A. S. & Vardi, A. Nitric
777   oxide mediates oxylipin production and grazing defense in diatoms. *Environ Microbiol* **22**,
778   629–645 (2020).

34. Wu, Z., Li, Q. P., Ge, Z., Huang, B. & Dong, C. Impacts of biogenic polyunsaturated
aldehydes on metabolism and community composition of particle-attached bacteria in
coastal hypoxia. *Biogeosciences* **18**, 1049–1065 (2021).
35. Edwards, B. R., Bidle, K. D. & Van Mooy, B. A. S. Dose-dependent regulation of microbial
activity on sinking particles by polyunsaturated aldehydes: Implications for the carbon
cycle. *Proc Natl Acad Sci U S A* **112**, 5909–5914 (2015).
36. Ianora, A. & Miralto, A. Toxigenic effects of diatoms on grazers, phytoplankton and other
microbes: A review. *Ecotoxicology* **19**, 493–511 (2010).
37. Stoecker, D. K. *et al.* Underestimation of microzooplankton grazing in dilution
experiments due to inhibition of phytoplankton growth. *Limnol Oceanogr* **60**, 1426–1438
(2015).
38. Vidoudez, C., Casotti, R., Bastianini, M. & Pohnert, G. Quantification of dissolved and
particulate polyunsaturated aldehydes in the Adriatic Sea. *Mar Drugs* **9**, 500–503 (2011).
39. Wu, Z., Li, Q. P., Dong, Y., Xu, J. & Luo, L. High-Resolution Surveys of Phytoplankton-
Derived Polyunsaturated Aldehydes at Frontal Zones Outside a Eutrophic Estuary. *J
Geophys Res Biogeosci* **126**, e2020JG005808 (2021).
40. Wu, Z., Li, Q. P., Ge, Z., Huang, B. & Dong, C. Impacts of biogenic polyunsaturated
aldehydes on metabolism and community composition of particle-Attached bacteria in
coastal hypoxia. *Biogeosciences* **18**, 1049–1065 (2021).
41. Stoecker, D. K. *et al.* Underestimation of microzooplankton grazing in dilution
experiments due to inhibition of phytoplankton growth. *Limnol Oceanogr* **60**, 1426–1438
(2015).
42. Kolody, B. C. *et al.* Diel transcriptional response of a California Current plankton
microbiome to light, low iron, and enduring viral infection. *ISME J* **13**, 2817–2833 (2019).
43. Krause, J. W. *et al.* The interaction of physical and biological factors drives phytoplankton
spatial distribution in the northern California Current. *Limnol Oceanogr* **65**, 1974–1989
(2020).
44. Bolaños, L. M. *et al.* Small phytoplankton dominate western North Atlantic biomass. *ISME
Journal* **14**, 1663–1674 (2020).
45. de Vargas, C. *et al. Eukaryotic Plankton Diversity in the Sunlit Ocean. Stefanie Kandels-
Lewis* vol. 15 https://www.science.org.
46. Mojica, K. D. A., Huisman, J., Wilhelm, S. W. & Brussaard, C. P. D. Latitudinal variation in
virus-induced mortality of phytoplankton across the North Atlantic Ocean. *ISME J* **10**,
500–513 (2016).
47. Eich, C. *et al.* Ecological Importance of Viral Lysis as a Loss Factor of Phytoplankton in
the Amundsen Sea. *Microorganisms* **10**, (2022).
48. Vardi, A. *et al.* Viral glycosphingolipids induce lytic infection and cell death in marine
phytoplankton. *Science (1979)* **326**, 861–865 (2009).
49. Diaz, B. P. *et al.* Seasonal mixed layer depth shapes phytoplankton physiology, viral
production, and accumulation in the North Atlantic. *Nat Commun* **12**, 6634 (2021).
50. Kimura, K. & Tomaru, Y. Isolation and Characterization of a Single-Stranded DNA Virus
Infecting the Marine Diatom Chaetoceros sp. Strain SS628-11 Isolated from Western
JAPAN. *PLoS One* **8**, e82013- (2013).

51.   Tomaru, Y., Takao, Y., Suzuki, H., Nagumo, T. & Nagasaki, K. Isolation and
Characterization of a Single-Stranded RNA Virus Infecting the Bloom-Forming Diatom
Chaetoceros socialis *Appl Environ Microbiol* **75**, 2375 (2009).
52.   Tomaru, Y. *et al.* Isolation and characterization of a single-stranded RNA virus that infects
the marine planktonic diatom Chaetoceros sp. (SS08-C03). *Phycological Res* **61**, 27–36
(2013).
53.   Pang, Z. *et al.* Using MetaboAnalyst 5.0 for LC–HRMS spectra processing, multi-omics
integration and covariate adjustment of global metabolomics data. *Nat Protoc* **17**, 1735–
1761 (2022).
54.   Chong, J., Wishart, D. S. & Xia, J. Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. *Curr Protoc Bioinformatics* **68**, e86 (2019).
55.   Silsbe, G. M., Behrenfeld, M. J., Halsey, K. H., Milligan, A. J. & Westberry, T. K. The
CAFE model: A net production model for global ocean phytoplankton. *Global
Biogeochem Cycles* **30**, 1756–1777 (2016).
56.   Boeuf, D. *et al.* Biological composition and microbial dynamics of sinking particulate
organic matter at abyssal depths in the oligotrophic open ocean. *Proc Natl Acad Sci U S
A* **116**, 11824–11832 (2019).
57.   Luo, E., Leu, A. O., Eppley, J. M., Karl, D. M. & DeLong, E. F. Diversity and origins of
bacterial and archaeal viruses on sinking particles reaching the abyssal ocean. *ISME
Journal* **16**, 1627–1635 (2022).
58.   Karl, D. M., Church, M. J., Dore, J. E., Letelier, R. M. & Mahaffey, C. Predictable and
efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen
fixation. *Proc Natl Acad Sci U S A* **109**, 1842–1849 (2012).
59.   Laber, C. P. *et al.* Coccolithovirus facilitation of carbon export in the North Atlantic. *Nat
Microbiol* **3**, 537–547 (2018).
60.   Mojica, K. D. A. & Brussaard, C. P. D. Factors affecting virus dynamics and microbial
host–virus interactions in marine environments. *FEMS Microbiol Ecol* **89**, 495–515
(2014).
61.   Buesseler, K. O. *et al.* An assessment of the use of sediment traps for estimating upper
ocean particle fluxes. *Journal of Marine Research* vol. 65 345–416 Preprint at
https://doi.org/10.1357/002224007781567621 (2007).
62.   Siegel, D. A., Devries, T., Cetini´c, I. C. & Bisson, K. M. Quantifying the Ocean's
Biological Pump and Its Carbon Cycle Impacts on Global Scales. *Downloaded from
www.annualreviews.org. Guest* (2023) doi:10.1146/annurev-marine-040722.
63.   Henson, S. A. *et al.* A reduced estimate of the strength of the ocean's biological carbon
pump. *Geophys Res Lett* **38**, (2011).
64.   Nanjappa, D., d'Ippolito, G., Gallo, C., Zingone, A. & Fontana, A. Oxylipin diversity in the
diatom family Leptocylindraceae reveals DHA derivatives in marine diatoms. *Mar Drugs*
12, 368–384 (2014).
65.   Esposito, R. *et al.* Combined Effects of Diatom-Derived Oxylipins on the Sea Urchin
Paracentrotus lividus. *Int J Mol Sci* **21**, 719 (2020).
66.   Balestra, C., Alonso-Sáez, L., Gasol, J. M. & Casotti, R. Group-specific effects on coastal
bacterioplankton of polyunsaturated aldehydes produced by diatoms. *Aquatic Microbial
Ecology* **63**, 123–131 (2011).

67.   Paul, C. *et al.* Diatom derived polyunsaturated aldehydes do not structure the planktonic
867   microbial community in a mesocosm study. *Mar Drugs* **10**, 775–792 (2012).
868   68.   Bidle, K. D. & Azam, F. Accelerated dissolution of diatom silica by marine bacterial
869   assemblages. *Nature* **397**, 508–512 (1999).
870   69.   Brzezinski, M. A., Jones, J. L., Bidle, K. D. & Azam, F. The balance between silica
871   production and silica dissolution in the sea: Insights from Monterey Bay, California,
872   applied to the global data set. *Limnol Oceanogr* **48**, 1846–1854 (2003).
873   70.   Bidle, K. D., Brzezinski, M. A., Long, R. A., Jones, J. L. & Azam, F. Diminished efficiency
874   in the oceanic silica pump caused by bacteria-mediated silica dissolution. *Limnol
875   Oceanogr* **48**, 1855–1868 (2003).
876   71.   Yamada, Y., Tomaru, Y., Fukuda, H. & Nagata, T. Aggregate formation during the viral
877   lysis of a marine diatom. *Front Mar Sci* **5**, (2018).
878   72.   Deutsch, C. *et al.* Biogeochemical variability in the California Current System. *Prog
879   Oceanogr* **196**, (2021).
880   73.   Hunter, J. E., Frada, M. J., Fredricks, H. F., Vardi, A. & Van Mooy, B. A. S. Targeted and
881   untargeted lipidomics of Emiliania huxleyi viral infection and life cycle phases highlights
882   molecular biomarkers of infection, susceptibility, and ploidy. *Front Mar Sci* **2**, (2015).
883   74.   Vardi, A. *et al.* Host–virus dynamics and subcellular controls of cell fate in a natural
884   coccolithophore population. *Proceedings of the National Academy of Sciences* **109**,
885   19327–19332 (2012).
886   75.   Pelusi, A. *et al.* Virus-induced spore formation as a defense mechanism in marine
887   diatoms. *New Phytologist* **229**, 2251–2259 (2021).
888   76.   Nissimov, J. I. *et al.* Dynamics of transparent exopolymer particle production and
889   aggregation during viral infection of the coccolithophore, Emiliania huxleyi. *Environ
890   Microbiol* **20**, 2880–2897 (2018).
891   77.   Leflaive, J. & Ten-Hage, L. Effects of 2E,4E-Decadienal on Motility and Aggregation of
892   Diatoms and on Biofilm Formation. *Microb Ecol* **61**, 363–373 (2011).
893   78.   Smith, C. A., Want, E. J., O'Maille, G., Abagyan, R. & Siuzdak, G. XCMS: Processing
894   mass spectrometry data for metabolite profiling using nonlinear peak alignment,
895   matching, and identification. *Anal Chem* **78**, 779–787 (2006).
896   79.   Benton, H. P., Want, E. J. & Ebbels, T. M. D. Correction of mass calibration gaps in liquid
897   chromatography-mass spectrometry metabolomics data. *Bioinformatics* **26**, 2488–2489
898   (2010).
899   80.   Tautenhahn, R., Böttcher, C. & Neumann, S. Highly sensitive feature detection for high
900   resolution LC/MS. *BMC Bioinformatics* **9**, 504 (2008).
901   81.   Kuhl, C., Tautenhahn, R., Böttcher, C., Larson, T. R. & Neumann, S. CAMERA: An
902   Integrated Strategy for Compound Spectra Extraction and Annotation of Liquid
903   Chromatography/Mass Spectrometry Data Sets. *Anal Chem* **84**, 283–289 (2012).
904   82.   Collins, J. R., Edwards, B. R., Fredricks, H. F. & Van Mooy, B. A. S. LOBSTAHS: An
905   Adduct-Based Lipidomics Strategy for Discovery and Identification of Oxidative Stress
906   Biomarkers. *Anal Chem* **88**, 7154–7162 (2016).
907   83.   Melamud, E., Vastag, L. & Rabinowitz, J. D. Metabolomic Analysis and Visualization
908   Engine for LC−MS Data. *Anal Chem* **82**, 9818–9826 (2010).

84.   Clasquin, M. F., Melamud, E. & Rabinowitz, J. D. LC-MS Data Processing with MAVEN:
A Metabolomic Analysis and Visualization Engine. *Curr Protoc Bioinformatics* **37**,
14.11.1-14.11.23 (2012).
85.   Wang, M. *et al.* Sharing and community curation of mass spectrometry data with Global
Natural Products Social Molecular Networking. *Nat Biotechnol* **34**, 828–837 (2016).
86.   Wang, F. *et al.* CFM-ID 4.0: More Accurate ESI-MS/MS Spectral Prediction and
Compound Identification. *Anal Chem* **93**, 11692–11700 (2021).
87.   Watrous, J. D. *et al.* Directed Non-targeted Mass Spectrometry and Chemical Networking
for Discovery of Eicosanoids and Related Oxylipins. *Cell Chem Biol* **26**, 433-442.e4
(2019).
88.   Tsugawa, H. *et al.* A lipidome atlas in MS-DIAL 4. *Nat Biotechnol* **38**, 1159–1163 (2020).
89.   Lamborg, C. H. *et al.* The flux of bio- and lithogenic material associated with sinking
particles in the mesopelagic 'twilight zone' of the northwest and North Central Pacific
Ocean. *Deep Sea Res 2 Top Stud Oceanogr* **55**, 1540–1563 (2008).
90.   Collins, J. R. *et al.* The multiple fates of sinking particles in the North Atlantic Ocean.
Global Biogeochem Cycles **29**, 1471–1494 (2015).
91.   Whiteside, J. H. *et al.* Pangean great lake paleoecology on the cusp of the end-Triassic
extinction. *Palaeogeogr Palaeoclimatol Palaeoecol* **301**, 1–17 (2011).
92.   Iacazio, G., Langrand, G., Baratti, J., Buono, G. & Triantaphylides, C. Preparative,
Enzymatic Synthesis of Linoleic Acid (13S)-Hydroperoxide Using Soybean Lipoxygenase-
1. *J. Organic Chem* **55**, 1690–1691 (1990).
93.   Landry, M. R. & Hassett, R. P. *MARINE BIOLOGY Estimating the Grazing Impact of
Marine Micro-Zooplankton *. *Marine Biology* vol. 67 (1982).
94.   Bartual, A., Morillo-García, S., Ortega, M. J. & Cózar, A. First report on vertical
distribution of dissolved polyunsaturated aldehydes in marine coastal waters. *Mar Chem*
204, 1–10 (2018).
95.   Bartual, A. *et al.* Polyunsaturated aldehydes from large phytoplankton of the Atlantic
Ocean surface (42°N to 33°S). *Mar Drugs* **12**, 682–699 (2014).
96.   Fontana, A. *et al.* Chemistry of oxylipin pathways in marine diatoms. *Pure and Applied
Chemistry* **79**, 481–490 (2007).
97.   d'Ippolito, G. *et al.* New C16 fatty-acid-based oxylipin pathway in the marine diatom
Thalassiosira rotula. *Org Biomol Chem* **3**, 4065–4070 (2005).
98.   D'Ippolito, G. *et al.* 15S-Lipoxygenase metabolism in the marine diatom Pseudo-nitzschia
delicatissima. *New Phytologist* **183**, 1064–1071 (2009).
99.   Lauritano, C. *et al.* Effects of the oxylipin-producing diatom Skeletonema marinoi on gene
expression levels of the calanoid copepod Calanus sinicus. *Mar Genomics* **24**, 89–94
(2015).
946
947
948

950
951 **Figure S1. A summary of the range of concentrations known to be bioactive in amendment experiments and**
952 **the maximum concentrations observed across the global ocean.** The asterisk is the highest dissolved PUA
953 concentration observed *in situ.*[94]. The caret is the highest particulate PUA concentration observed *in situ*[94]. The 'x' is
954 the highest PUA concentration observed within sinking particles[35].

955
Figure S2. Time course of virus infection of *C. tenuissimus* and *C. socialis*. Growth curves for uninfected host-
only control triplicates (green circles), *C. tenuissimus* infected with CtenRNAV (purple triangles) and CtenDNAV
(pink triangle) viruses, and C. socialis infected with CsfrRNAV (blue triangles). Initial experiments at Rutgers in
2016 followed the diatom hosts, *Chaetoceros socialis* L4 (grass green circles) and *Chaetoceros tenuissimus* 2-10
(darkest green circle), to late exponential/ stationary phase and the dissolved lipid extracts were processed at WHOI.
Subsequent experiments at UC-Berkely in 2021 followed cells well into stationary and decline phase (mint green,
light green, and grey circles, denoted by an asterisk in the legend) were analyzed at UC-Berkeley.
963

964
Figure S3. Outline of dissolved metabolome sampling at sea as well as sample extraction and analysis in the
lab. One liter of water was collected from the surface ocean (55% I0 depth approx. 5 m) and the chlorophyll
maximum (4-35 m) at each station. The 1L sample was filtered through a 0.2 um filter and benzaldehyde was added
to the filtrate as an internal standard. The dissolved organics were extracted from the filtrate onto a Water HLB
Solid Phase Extraction cartridge using a vacuum manifold. SPE Samples were flash frozen, and the dissolved
metabolome was eluted from the SPE back in the lab. Dissolved metabolome samples were run for lipidomic
analysis using HPLC paired with Orbitrap Exactive mass spectrometer in early 2014. Three samples were excluded
from further analysis due to missing internal standard. The eleven remaining raw lipidomes were processed through
the xcms-CAMERA-LOBSTAHS-MetaboAnalyst Chemoinformatic pipeline in the black box. Representative

samples were rerun in 2021 using an Orbitrap ID-X mass spectrometer which allowed for more targeted ms2
fragmentation for higher confidence oxylipin annotation of important features from the comparative statistical
analysis provided by MetaboAnalyst; these annotations can be found in Table S5.
977
978

979
Fig S4. Grazing and growth were lowest at the station with the highest oxylipin concentration and deckboard
amendment experiments with exogenous oxylipins show inhibition of growth and grazing. Contour maps of **A)**
the maximum total oxylipin and free fatty acid peak area, **B)** microzooplankton grazing rate at the 55% I0 depth, and
C) phytoplankton growth rate at the 55% I0 depth for each station of the DYEatom cruise. Plotted with Ocean Data
View using the DIVA contouring model which accounts for general ocean current patterns. D) Grazing rate
presented as a bar chart with the parallel oxylipin amendment rate. Asterisk denotes statistical difference in +
oxylipin amendment from the control treatment. E) Growth rate presented as a bar chart with the parallel oxylipin
amendment rate. Asterisks denote statistical difference from the control treatment.

988
Fig S5. Relationship between the total peak area and **A)** export, **B)** Si-stress of the surface ocean community **C)**
Grazing rate, and **D)** Phytoplankton growth rate across the DYEatom cruise (PS1312) reflect allelopathy of
oxylipins on the community, oxylipin production under stress, and evidence of a link between oxylipin production
and the viral shuttle. The silica stress metric is described in ref[17]. Briefly, the Y-axis is the percentage of the Si
uptake rate (B) that happens in the ambient Si concentration vs. a deck-board amendment with exogenous Si. A
lower value indicates more silicon stress in that environment. For the phytoplankton growth regression (D) all but
the negative growth rate was included. And for the microzooplankton grazing rate regression (C) the low peak area
samples were excluded.
997
998

999
Figure S6. Analogs to the exogenous oxylipins found to be bioactive in Fig 6I-K were found *in situ* in addition to
the various similar oxylipins of different carbon chain lengths and degrees of unsaturation highlighted in Fig 5. **A)**
Extracted ion chromatogram of HpETE analogs (i.e. compounds with a [M-H]⁻ adduct m/z of 335.2232) observed in
the DYEatom dissolved meta-lipidome and the integrated peak height displayed in the inset bar charts. **B)** The
normalized peak area for HpETE analogs averaged across each bloom state. Box plots show the 1st and 3rd quartile
range as the upper and lower bounds of the box and the mean is denoted as a line through the box. The error bars
denote the standard deviation with the dots representing outliers.

1007
1008 **Figure S7.** Features representing greater than 1% of the peak area in the annotated dissolved meta-lipidome on
1009 average. The heat map displays the percentage of the total annotated peak area attributable to each molecular species
1010 across the samples. Samples are grouped by similarity profile analysis of the structure of the dissolved meta-
1011 lipidome across the cruise track as presented in Fig. 5A.

Table S1. Compilation of maximum PUA and NVO concentrations observed in situ. Each oxylipin subclass is broken down into particulate (p-prefix) and dissolved (d-prefix). Note that no dNVO quantifications have been published. The pPUA values are derived from sonicated or freeze-thawed cells that were collected on a filter presenting a potential oxylipin concentration. The dPUA values were collected by extracting the organics on to a solid phase extraction cartridge. The pNVO values in Mar 2005 and April 2011 were estimated from cell abundances and fg of oxylipin per cell.

Oceanic Region	Date	pPUA	dPUA	pNVO	dNVO	PUA$_{sinking/suspended}$	Citation
Adriatic	Mar 2005			1.76 nM			Ianora *et al.*, 2015[8]
	Mar 2006	9 nM	2.3 nM				Vidoudez *et al.*, 2011[38]
	Feb 2008	20 nM	0.1 nM				Ribalet *et al.*, 2014[23]
	April 2011			48 pM			Lauritano *et al.*, 2016[10]
	June 2017			1.75 nM			Russo *et al*, 2020[25]
	Avg 2017			0.4 nM			
Pearl River Estuary	June 2015	20 nM	0.37 nM				Wu…Luo, 2021a[39]
	June 2019	0.2 nM	0.1 nM			10 µM	Wu...Dong, 2021b [34]
Western Mediterranean	Aug 2011	1.65 nM	1.87 nM				Bartual *et al.*, 2018[94]
	Mar 2012	0.2 nM	0.54 nM				
	Oct 2015	5.3 pM	34.9 pM				
North Atlantic	July 2011	1 pM					Bartual *et al.*, 2014[95]
	Jun 2012					37 µM	Edwards *et al.*, 2015[35]

Table S2. Authentic fatty acid, polyunsaturated aldehyde, non-volatile oxylipin standards with the common name or abbreviation, the structural level annotation assigned by the lipidomics pipeline presented here, and the retention time (RT) of each compound. Comparisons to the RT of the authentic standards were used to assign putative annotations to features in the dissolved lipidome.

Authentic Standard	Common Abbreviation/ Name	Structural Level Annotation	RT
Octanoic acid	--	FFA 8:0	2.65
Arachidonic acid	AA	FFA 20:4	8.09
Docosahexaenoic acid	DHA	FFA 22:6	7.71
9-Hydroperoxy octadecadienoic acid	9-HpODE	NVO 18:2 2O	5.44
13-Hydroperoxy octadecadienoic acid	13-HpODE	NVO 18:2 2O	5.44
9-Hydroxy octadecatrienoic acid	9-HOTrE	NVO 18:3 1O	4.02
13-Hydroperoxy octadecatrienoic acid	13-HpOTrE	NVO 18:3 2O	4.65
9-Hydroperoxy octadecatrienoic acid	9-HpOTrE	NVO 18:3 2O	4.28
15-Hydroxy eicosatrienoic acid	15-HETrE	NVO 20:3 1O	6.11
15-Hydroxy eicosatetraenoic acid	15-HETE	NVO 20:4 1O	5.43
5-Hydroperoxy eicosatetraenoic acid	5-HpETE	NVO 20:4 2O	5.01
12-Hydroperoxy eicosatetraenoic acid	12-HpETE	NVO 20:4 2O	5.93
15-Hydroperoxy eicosatetraenoic acid	15-HpETE	NVO 20:4 2O	5.73
5,6-Dihydroxy eicosatetraenoic acid	5,6-diHETE	NVO 20:4 2O	3.48
14,15- Dihydroxy eicosatetraenoic acid	14,15-diHETE	NVO 20:4 2O	3.00
5,12- Dihydroxy eicosatetraenoic acid	Leukotriene B4	NVO 20:4 2O	2.89
15-Hydroxy eicosapentaenoic acid	15-HEPE	NVO 20:5 1O	4.56
5,6,15- Trihydroxy eicosapentaenoic acid	Lipoxin A5	NVO 20:5 3O	1.72
Heptadienal	--	PUA 7:2	1.90
Octadienal	--	PUA 8:2	2.28
Nonadienal	--	PUA 9:2	2.69
Decadienal	DD	PUA 10:2	3.65

Table S3. Viral Infection and general death oxylipin-biomarkers from our *Chaetoceros*-virus experiments that were observed in culture, field studies, and their known allelopathy. The diatom species that each compound has been observed in is noted: *Skeletonema marinoi* (Sm), *Chaetoceros socialis* (Cs), *Chaetoceros affinis* (Ca)[96], *Thalassiosira rotul*[97] (Tr), *Pseudo-nitzchia delicatissima* (Pd)[98], *Skeletonema pseudocostatum* (Sp)[13], and *Phaeodactylum tricornutum* – microzooplankton grazing experiments (Pt-MZP)[33]. The lipidomic pipeline in this study automatically annotates to the structural level. Structural isomers of these compounds were observed in the DYEatom cruise dissolved lipidome (denoted as this study) and studies in the Mediterranean[8,10,25,99]. Compounds with known allelopathic effects are [10,25]. Many of these compounds were annotated to the function group level using a retention time model or *in silico* fragmentation model as outlined in Edwards *et al.* (2024); compound abbreviations in Fig 3B-C.

Viral Biomarkers	Observed in culture	Observed in the Field	Known allelopathy	RT-model	CFMID model
FFA 14:2 +1O_mz_239.165_RT_1.71	isomer Pt-MZP		*MZP grazing*		
FFA 20:5 +2O_mz_333.2073_RT_2.08	Sm, Ca, isomer Pt-MZP		Copepods; 20:4 +2O dihydro and hydroperoxy decrease MZP grazing; HpETE modulate particle associated microbial activity (this study)	10,11-diHEPE	
FFA 16:3 +3O_mz_297.1708_RT_2.13	16:3 oxylipins general Sm, Tr				
FFA 14:2 +1O_mz_239.165_RT_2.56	isomer Pt -MZP		*MZP grazing*		
FFA 16:3_mz_249.1857_RT_7.36	16:3 oxylipins general Sm, Tr	Lauritano *et al.,* 2016			
FFA 22:5 +1O_mz_345.2436_RT_7.67	*Lepidocylindrus*	Lauritano *et al.,* 2016; Russo *et al.,* 2020; this study	*Copepods*		
FFA 14:2 +1O_mz_239.165_RT_2.25	isomer Pt -MZP		*MZP grazing*		
FFA 22:4 +1O_mz_347.2593_RT_7.77	*Lepidocylindrus*				
FFA 18:3 +2O_mz_309.2074_RT_5.01	cyanobacteria, green algae, moss				
FFA 20:4 +1O_mz_319.2279_RT_5.95	Tr, *Stephanopyxis turris*, red algae, fungi	Lauritano *et al.* 2016; Russo *et al.* 2020; this study	Copepods; analog impacts sea urchins		

Death Biomarkers	Observed in culture	Observed in the Field	Known allelopathy	RT-model	CFMID model
FFA 16:1 +1O_mz_269.2123_RT_4.37	Sm, Tr	Lauritano *et al.* 2016; this study	Copepods		
FFA 18:3 +2O_mz_309.2073_RT_1.98	cyanobacteria, green algae, moss				
FFA 16:2 +1O_mz_267.1966_RT_4.78	Sm, Tr	this study	Copepods	9-KHME	9-epHME
FFA 16:3 +3O_mz_297.1707_RT_1.52	16:3 oxylipins general Sm, Tr				
FFA 16:2 +1O_mz_267.1966_RT_4.28	Sm, Tr	this study	Copepods	9-KHME	5-epHME
FFA 16:3 +1O_mz_265.1809_RT_2.99	Sm, Tr	Lauritano *et al.* 2016; Russo *et al.* 2020; this study	Copepods		9-HHTrE
FFA 16:2 +1O_mz_267.1966_RT_3.46	Sm, Tr, Flowering plants	this study	Copepods	9-KHME	
FFA 18:4 +1O_mz_291.1967_RT_2.02					
FFA 18:3 +2O_mz_309.2073_RT_7.32	cyanobacteria, green algae, moss				
FFA 16:4 +2O_mz_279.1602_RT_1.85	Sm	this study	Copepods		9-HpHTE

1060
1061
1062

Table S4. Location and data of collection for the dissolved meta-lipidomic samples across the DYEatom cruise track

Station	Depth (m)	Latitude N	Longitude W	Date	MLD
1	10	36°46.864 '	121°59.1257 '	June 27, 2013	7
4	2	36°27.31 '	122°20.32 '	June 29, 2013	16
4	4	36°27.31 '	122°20.32 '	June 29, 2013	16
6	4	36°42.61 '	123°29.62 '	June 30, 2013	26
6	35	36°42.61 '	123°29.62 '	June 30, 2013	26
7	8	37°33.42 '	123°37.96 '	July 1, 2013	18
8	1	38°8.96 '	123°5.95 '	July 2, 2013	7
8	4	38°8.96 '	123°5.95 '	July 2, 2013	7
9	7	38°15.89 '	123°58.12 '	July 3, 2013	27
10	10	37°50.24 '	123°6.6 '	July 4, 2013	10
11	2	36°46.67 '	121°58.83 '	July 5, 2013	11

Table S5. Further annotation of significant compounds identified by sPLS-DA analysis of either lysis or mid infection within the DYEatom cruise dissolved lipidome. LOBSTAHS annotation, peak retention time, the mass to charge ration (m/z) of the feature, and the annotation from considering the ms^2 fragmentation of samples analyzed using HRAM-MS^n analysis outlined in Fig. S3. It is also noted whether each compound was a biomarker observed in the *Chaetoceros* culture experiments in Fig. 3B-C and whether the compound has known allelopathy (based on the studies cited in Table S3).

Lysis	LOBSTAHS annotation	Retention Time (min)	m/z	Further Annotation	Annotation Method	Biomarker/ Allelopathic
	PUA 12:5 1O	2.5	189.0921	$C_{12}H_{14}O_2$	-	No/No
	FFA 20:4 +3O	4.1	351.2177	5-epoxy 8,14 dihydroxy eicosatrienoic acid	hypothesized from Edwards Lab DB	No/No
	FFA 22:5 +1O	6	345.2435	signal too low	-	No/No
	FFA 16:1 +1O	4.2	269.2122	9-HHME	Edwards Lab DB	Death/Yes
	FFA 20:4 +4O	4.5	367.2126	signal too low	-	No/No
	FFA 20:4 +1O	5.4	319.2279	signal too low	-	Virus/Yes
	FFA 16:2 +1O	4	267.1966	9-oxoHME	Edwards Lab DB	Death/Yes
	FFA 20:5 +1O	4.5	317.2122	14-HEPE	Edwards Lab DB	No/Yes
	FFA 16:1 +2O	4.5	285.2071	9-HpHME	Edwards Lab DB	No/No
	FFA 16:2 +1O	4.5	267.1966	9-HHDE	Edwards Lab DB	Death/Yes

Mid Infection	LOBSTAHS annotation	Retention Time (min)	m/z	Further Annotation	Annotation Method	
	FFA 22:5 +4O	2.6	393.2283	signal too low	-	No/No
	FFA 22:4 +4O	3.2	395.2439	signal too low	-	No/No
	FFA 22:4 +4O	3.7	395.2439	signal too low	-	No/No
	FFA 22:5 +4O	4.2	393.2283	signal too low	-	No/No
	FFA 22:4 +4O	4.2	395.2439	signal too low	-	No/No
	FFA 22:4 +4O	2.7	395.2439	signal too low	-	No/No
	FFA 22:5 +4O	3.7	393.2283	signal too low	-	No/No
	FFA 22:5 +4O	3.3	393.2283	signal too low	-	No/No
	FFA 16:4	5.4	247.1704	hexadecatetraenoic acid	MS-Dial Lipid Blast	No/No
	FFA 18:4 +2O	5.4	307.1915	signal too low	-	No/No

1066
1067
1068
1069

Table S6. Average richness and diversity of the oxylipins and free fatty acids in the dissolved lipidome collected from the surface ocean along the DYEatom cruise. Samples subdivided by the state of diatom infection.

Non-bloom			
Samples	Richness	Diversity	Sum
St6_4m	165	-4.03134	2×10^8
St6_35m	164	-3.97726	1.28×10^8
St9_7m	156	-3.92887	3.25×10^8
Avg	161.6667	-3.97915	2.17×10^8
stdev	4.932883	0.051261	99692081
Post-Lytic			
Samples	Richness	Diversity	Sum
St7_8m	175	-3.60247	5.27×10^9
St8_1m	175	-3.47831	2.31×10^9
St8_4m	174	-3.74992	3.74×10^9
St10_10 m	174	-3.54632	1.4×10^{10}
Avg	174.5	-3.59426	5.48×10^9
stdev	0.57735	0.115527	6×10^9
Active Infection			
Samples	Richness	Diversity	Sum
St11_2m	174	-2.94217	2.62×10^9
St4_2m	174	-3.28648	1.37×10^9
St1_5m	175	-2.94248	2.17×10^9
St4_9m	172	-2.71614	1.96×10^9
Avg	173.75	-2.97182	2.03×10^9
stdev	1.258306	0.235316	5.18×10^8

Table S7. T-test results showing a significant difference between the average diversity of the dissolved lipidome collected during three different states of viral infection. Bold italics values are significant <0.01, bold values are significant <0.05 with a Bonferroni correction.

	Richness	Diversity	Sum
Post-lytic *vs.* Non-bloom	0.044609	***0.003134***	0.177903
Active Infection *vs.* Post-lytic	0.336716	**0.007219**	0.334421
Active Infection *vs.* Non-bloom	0.045554	***0.002336***	**0.004728**

1073
1074
1075
1076 Supplemental File 1: Dissolved Lipidome Rutgers *Chaetoceros* Viral Infection Experiments
1077 Supplemental File 2: Dissolved Meta-Lipidome DYEatom Cruise.
1078 Supplemental File 3: DYEatome Cruise Lipidome Retention Time Model
1079