An aphid-resistant plant metabolite as a candidate aphicide: Insight into the bioactivity and action mode of betulin against aphids

Junxiu Wang [a†], Matthana Klakong [b†], Qiuyu Zhu [b], Jinting Pan [a], Yudie Duan [a], Lirong Wang [d], Yong Li [d], Jiangbo Dang [a], Danlong Jing [a], Hong Zhou [b, c*]

[a] Key Laboratory of Agricultural Biosafety and Green Production of Upper Yangtze River (Ministry of Education), Chongqing Key Laboratory of Forest Ecological Restoration and Utilization in the Three Gorges Reservoir Area, College of Horticulture and Landscape Architecture, Southwest University, Beibei, Chongqing 400715, China.

[b] College of Plant Protection, Southwest University, Chongqing, China

[c] Yibin Academy of Southwest University, Southwest University, Chongqing, China

[d] Zhengzhou Fruit Research Institute, Chinese Academy of Agricultural Sciences, Zhengzhou, 451100, China

[†] These authors contributed equally.

[*] Correspondence to Dr. Hong Zhou, College of Plant Protection, Southwest University, Chongqing 400715, China. E-mail: Hong Zhou: zhouhong1990@swu.edu.cn.

ABSTRACT

Pest-resistant plants usually utilize secondary metabolites to cope with insect infestation. Betulin, a key bioactive compound in aphid-resistant wild peach, possesses promising applications in crop protection. Here, betulin, in both greenhouse and field experiments, displayed excellent control efficacy against *Myzus persicae*. RNA-seq, qRT–PCR, and western blotting revealed that betulin significantly inhibited the expression of *MpGABR* (encoding a GABA$_A$ receptor). Besides, RNAi-mediated silencing of *MpGABR* markedly increased aphid sensitivity to betulin. Furthermore, MST (microscale thermophoresis) and voltage-clamp assays indicated that betulin bound to MpGABR (K_d = 2.24 μM) and acted as an inhibitor of MpGABR. Molecular docking, mutagenesis and genome editing suggested that THR228 is a critical and highly conserved site in MpGABR that betulin binds to specifically, causing aphid death. Overall, the activity of betulin depends on specific targeting and inhibition of MpGABR. Elucidating the mechanism of action of this peach-derived insecticide may offer a sustainable green strategy for aphid control.

KEYWORDS: betulin; aphid; GABA$_A$ receptor; insecticide; insecticidal mechanism

1. INTRODUCTION

Over more than 400 million years of coevolution with herbivores, plants have developed a variety of morphological (Soujanya et al. 2023), biochemical (Zhou et al. 2024b), and molecular (Wang et al. 2024) defense strategies against insect herbivore attack. Plant secondary metabolites are highly valuable natural compounds, offering abundant and potent biochemical defense driven by plant-insect coevolution (Jahan et al. 2025a). Interestingly, wild relatives of crops with insect resistance usually accumulate large amounts of biologically active compounds (Alonso-Salces et al. 2022; Wang et al. 2022a). Accordingly, investigating the bioactivities and potential mechanisms of action of compounds derived from wild resistant germplasms may provide effective and sustainable strategies for pest control in crop protection.

Myzus persicae (Sülzer) is one of the most destructive sap-feeding pests worldwide. It is able to settle on more than 400 plant species globally, displaying extraordinary polyphagia and a wide range of hazards (Bass et al. 2014). In addition to ingesting phloem saps and secreting honeydew, *M. persicae* can transmit various plant viruses (Guo et al. 2022), resulting in severe losses in crop yields. To date, the principal control strategy for *M. persicae* has relied on synthetic insecticides, including pyrethroids, organophosphorus compounds, carbamates and neonicotinoids (Jiménez-Jiménez et al. 2019; Martins et al. 2021; Zhang et al. 2023). However, long-term irrational application of these chemical insecticides has led to environmental pollution and promoted the resistance of aphids to insecticides (Troczka et al. 2021; Zhang et al. 2023; Stará et al. 2024). Therefore, it is necessary to develop novel green insecticides as potential alternatives to synthetic insecticides. Fortunately, plant secondary metabolites are a "treasure trove of active compounds" that serve as a rich resource for the development of promising green insecticides.

Our previous study revealed that *Prunus davidiana*, a close wild relative of cultivated peach,

displays strong resistance to *M. persicae* through the accumulation of high levels of betulin

(Wang et al. 2022a; Wang et al. 2024). In addition, betulin, a lupane-type triterpene, possesses

strong insecticidal activity and is a promising substance for the development of novel aphid-

control insecticides. A number of studies have reported that betulin and its derivatives exhibit

a wide range of pharmacological activities (Amiri et al. 2020). An immune stimulant, Ir-Bet,

was prepared using iridium complex and betulin, which evoked ferritinophagy-enhanced

ferroptosis, thereby activating anti-tumor immunity (Lv 2023). The anti-inflammatory effect of

betulin has been reported in macrophages at lymphoma site in mice (Szlasa et al. 2023). Betulin

has been found to improve hyperlipidemia and insulin resistance and decrease atherosclerotic

plaques by inhibiting the maturation of sterol regulatory element-binding protein (Tang et al.

2011). Additionally, betulin and its derivatives have been found to exhibit insecticidal activity

against *Plutella xylostella* L. (Huang et al. 2025), *Aedes aegypti* (de Almeida Teles et al. 2024), and

Drosophila melanogaster (Lee and Min 2024). However, the insecticidal mechanism of betulin

against aphids remains unclear.

Gamma-aminobutyric acid (GABA) receptors have been confirmed to be targets of

terpenoids that impair neuronal function in insect herbivores (Guo et al. 2023). There are two

types of GABA receptors: ionotropic (GABA$_A$) and metabotropic (GABA$_B$) receptors. The first

GABA receptor subunit identified in insects is encoded by *Rdl*, which confers resistance to

dieldrin in *Drosophila melanogaster* (Ffrench-Constant et al. 1991). Although a variety of

terpenoids play roles as positive allosteric modulators or noncompetitive antagonists of

GABA$_A$ receptors (GABRs) (Guo et al. 2023), whether betulin targets the GABR in *M. persicae*

needs to be further investigated. GABRs are heteropentameric ligand-gated ion channels in the

central nervous system that conduct chloride and bicarbonate ions and are the target of

numerous drugs for the treatment of neuropsychiatric disorders (Thompson 2024). GABRs are

composed of 5 different types of subunits, each with 4 helical transmembrane domains (TM1-

TM4), of which TM2 is located at the center of the pentamer and forms an ion channel (Fan et

al. 2024). Previous study has proposed that vertebrate and human GABR genes maintain a

broad and conservative gene clustering pattern, while in invertebrates, this pattern is missing,

indicating that these gene clusters were formed early in vertebrate evolution and were

established after invertebrates diverged. Notably, invertebrates each possess a unique GABR

gene pair, which are homologous with human GABR α and β subunits, suggesting that the

existing GABR gene cluster evolved from an ancestral α-β subunit gene pair (Tsang et al. 2006).

During the coevolution of plants and insects, the duplications and amino acid substitutions in

GABR may have been beneficial for adaptation to insecticides and terpenoid compounds (Guo

et al. 2023).

In this study, betulin was confirmed to have excellent control efficacy against *M. persicae*

in both greenhouse and field experiments. RNA-seq, qRT–PCR, and western blotting assays

revealed that betulin significantly inhibited the expression of *MpGABR* in aphids. In addition,

RNAi-mediated silencing of *MpGABR* markedly increased aphid sensitivity to betulin.

Furthermore, microscale thermophoresis (MST) and voltage-clamp assays indicated that

betulin can bind to MpGABR (K_d = 2.24 μM) and act as an inhibitor (EC_{50} = 20.66 μM) of

MpGABR. Molecular docking analysis suggested that betulin bound to MpGABR via the

THR228 amino acid residue. This site is highly conserved across 11 species in the Aphididae

family of Hemiptera and may be a critical specific binding site for betulin in MpGABR in aphids.

Mutagenesis and genome editing experiments revealed that betulin bound specifically to this

amino acid residue in aphids but not in *Drosophila*, resulting in aphid death. This study

elucidated the insecticidal mechanism of betulin involving the targeting of MpGABR,

providing a sustainable green strategy for aphid control. However, given that betulin may

affect a wider range of organisms, it should be used with caution.

2. MATERIALS AND METHODS

2.1 Test insects

M. persicae populations were collected from tobacco (*Nicotiana tabacum* L.) leaves. Wild-

type (WT) flies (*D. melanogaster*) were purchased from Fungene Biotechnology Co., Ltd.,

Jiangsu, China. The test insects were cultured in a greenhouse (25 °C, 70% relative humidity

and 14 h light/10 h dark cycle) in Beibei, Chongqing, China.

2.2 Leaf bioassay

The toxicity of betulin against *M. persicae* was measured using a slip-dip bioassay as

previously described (Wang et al. 2022a). Thirty apterous adult *M. persicae* individuals were

gently transferred on tobacco (*N. tabacum*) leaves using a fine brush. The tobacco leaves were

subsequently immersed in 0.1% (v/v) Tween-80 and 3% (v/v) acetone solution containing

betulin or pymetrozine (as a positive control) at 7 doses (0, 0.0625, 0.125, 0.25, 0.5, 1, and 2

mg· mL^{-1}) for 5 seconds. After drying, the leaves with *M. persicae* were placed on wet filter paper

and cultured on a water-soaked sponge for 48 h. Three replicates were tested for each treatment.

Furthermore, the LC_{30}, LC_{50} and LC_{70} values of betulin were calculated using log-probit analysis

via IBM SPSS Statistics (v.22.0, Chicago, IL, USA) (Zhou et al. 2024b).

2.3 Greenhouse and field bioassays

The control effects of betulin against *M. persicae* in the greenhouse and field were evaluated

by a spraying-based method as described previously (Zhou et al. 2023). The commercial

insecticide pymetrozine and an aqueous solution containing 0.1% (v/v) Tween-80 and 3% (v/v)

acetone were employed as positive and negative controls, respectively. Sprays of 0.1641

mg·mL^{-1} (LC$_{50}$) betulin and 1.0612 mg·mL^{-1} (LC$_{50}$) pymetrozine (Table S1) were prepared in

aqueous solutions including 3% (v/v) acetone and 0.1% (v/v) Tween-80, respectively. Before the

control effect assays, each tobacco (*N. tabacum* L.) seedling was pre-inoculated with 60 apterous

adults of *M. persicae*. In the greenhouse trial, each treatment included 8 replications, and each

replication consisted of 3 tobacco plants growing under controlled conditions as described

above. The field trial (Beibei District, Chongqing, China.) was conducted under natural

conditions with an average temperature of 20 °C. Each treatment had 8 replications, covering

an area of 20 m^2, with 3 protected rows per replication. Furthermore, each tobacco plant was

sprayed with 20 mL of the corresponding solution with an electric sprayer. Mortality was

recorded at 1, 5, 9, and 14 days after treatment. The control efficacy of betulin against *M. persicae*

was calculated according to the methods of Zhou (Zhou et al. 2023).

2.4 RNA-seq assay

After exposure to 0.1641 mg· mL^{-1} (48 h LC$_{50}$) betulin for 48 h, 100 apterous *M. persicae*

adults were sampled for each of 3 replicates. and total RNA was extracted via TRIzol reagent

(Tiangen, Beijing, China) to construct mRNA sequencing libraries. RNA-seq analysis was

performed on the Ion Proton BGISEQ-500 platform (Beijing Genomic Institute, BGI, China).

After the raw reads were filtered, the clean reads were aligned to the *M. persicae* reference

genome, which was downloaded from the National Center for Biotechnology Information

database, using HISAT2 (v2.1.0) (Kim et al. 2015). The expression levels of genes in each sample

were normalized to the fragments per kilobase million mapped reads (FPKM) value. The

differentially expressed genes (DEGs) between groups were identified using the fold change

factor ($|Log_2(FC)| > 1$). Functional enrichment analysis of DEGs was performed using the Gene

Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases.

2.5 Validation by quantitative real-time PCR (qRT–PCR)

The gene-specific primers used for qRT–PCR, which were designed with Primer Premier

5.0, are listed in Table S2. According to the manufacturer's specifications for the Bio-Rad iQ™

SYBR® Green Supermix Kit, qRT–PCR was performed with the following conditions: 60 °C for

15 s, followed by 48 cycles of 60 °C for 30 s, and 72 °C for 30 s. The *ribosomal protein S18* gene

(*RPS18*) was used as the reference gene to normalize the expression levels. The relative

expression levels of the target genes were calculated using the $2^{-\Delta\Delta Ct}$ approach (Zhou et al.

2024b). Three biological replicates were used for each assay.

2.6 Cloning, bioinformatics and phylogenetic analysis

The cDNA of *M. persicae* was synthesized from the extracted total RNA using the

PrimeScript® 1st Strand cDNA Synthesis Kit (TaKaRa, Japan). The coding sequences of *MpGABR*

(encoding GABA$_A$ receptor), *MpGABRAP* (encoding GABA$_A$ receptor-associated protein) and

MpGABRB (encoding GABA$_A$ receptor β subunit) were cloned from *M. persicae* cDNA using

the corresponding specific primers (Table S2) and then inserted separately into the pGEM-T

Easy vector. The recombinant plasmids were subsequently transferred into *Escherichia coli*

DH5α competent cells for sequencing. BioXM (v2.7.1) (http://202.195.246.60/BioXM/) was used

to analyze the sequence information of these three genes, including the length, molecular

weight, and isoelectric point of the deduced amino acid sequence. MEGA7 was used for

multiple amino acid sequence alignment (ClustalW) and phylogenetic analysis (neighbor-

joining method with 1000 replicates) of MpGABR.

2.7 RNAi assay

To knock down the target genes (*MpGABR*, *MpGABRAP*, and *MpGABRB*), *MpGABR-*,

MpGABRAP-, and *MpGABRB*-dsRNA were artificially synthesized using the dsRNA Synthesis

Kit (Thermo Scientific, Vilnius, Lithuania, EU). In accordance with a previously described

method (Zhou et al. 2024a), fresh tobacco leaves were cut into 3.0 cm-diameter discs and placed

in an oven at 50 °C for 3 min. Subsequently, the dried leaf discs were exposed to 10 µL of diethyl

pyrocarbonate-treated nuclease-free water (DEPC-water), dsGFP (green fluorescent protein,

1000 ng· µL⁻¹, negative control), and *MpGABR-*, *MpGABRAP-*, or *MpGABRB*-dsRNA (1000

ng/·µL⁻¹) at 25 °C for 5 h. After the leaf discs had absorbed the solution, they were placed on

wet filter paper on a water-soaked sponge. Subsequently, 30 apterous adult *M. persicae*, starved

for 24 h, were fed on the treated leaf discs for 48 h. The surviving aphids were collected after

exposure for qRT–PCR. Three biological replicates were included for each treatment.

2.8 Prokaryotic expression of *MpGABR* and western blotting

To produce the MpGABR protein *in vitro*, an *Escherichia coli* expression system was

constructed (Zhou et al. 2021). Briefly, the coding sequence of *MpGABR* was inserted into the

expression vector PET-B2M, obtained from BGI (Shenzhen, China), and transformed into BL21

(*DE3*) competent cells. To induce the expression of the MpGABR protein, 2 mM isopropyl *β*-D-

1-thiogalactopyranoside (IPTG) was used, and then, the transformed BL21 cells were incubated

at 180 rpm and 30 °C for 24 h. Finally, the protein was extracted and purified using Solarbio

GST-tag Purification Resin for subsequent assays.

The purified MpGABR protein resolved on a 15% SDS–PAGE (sodium dodecyl sulfate–

polyacrylamide gel electrophoresis) gel and then transferred to a polyvinylidene fluoride

(PVDF) membrane at 20 V for 1 h. Moreover, 1 × TBST (Tris-Borate-Sodium Tween-20)

containing 5% fat-free milk powder was used to block the PVDF membrane. Then, the PVDF

membrane was incubated with an antibody against MpGABR (polyclonal rabbit antibody,

provided by GeneCreate Biotechnology) at 25 °C for 1 h. Subsequently, the membrane was

further incubated with the corresponding horseradish peroxidase (HRP)-conjugated goat anti-

rabbit immunoglobulin G (IgG) antibody at 25 °C for 1 h and then washed three times with

TBST. Finally, the bands on the membrane were visualized using a Chemi Doc MP Imaging

System (Bio-Rad) after incubation with a chemiluminescence reagent (ECL, Bio-Rad). GAPDH

was used as a reference protein.

2.9 Molecular docking

In accordance with a previously reported method (Zhou et al. 2023), the protein structure

and function of MpGABR were predicted via the I-TASSER server (Iterative Threading

ASSEmbly Refinement). AutoDockTools software (v1.5.7) was used to conduct molecular

docking. Furthermore, PyMOL (v3.1) and Discovery Studio (v4.5) were used to evaluate the

molecular docking model.

2.10 Site-directed mutagenesis

The coding sequence of *MpGABR* was inserted into the PET-B2M vector and used as the

template for site-directed mutagenesis. In accordance with the manufacturer's specifications,

the Fast MultiSite Mutagenesis Kit (TransGen, Beijing, China) was used to generate the

multisite mutations in MpGABR (Ma et al. 2020). Then, the MpGABR mutants, with the

mutations of R224A (ARG224 replaced by ALA), A226T (ALA226 replaced by THR), F227Y

(PHE227 replaced by TYR), or T228R (THR228 replaced by ARG), were amplified, and the

amplicons were used to replace the *Bam*HI-*Sma*I fragment in the recombinant MpGABR

plasmid.

2.11 Method for *Drosophila* genome editing

To generate *Drosophila* mutants bearing the DmGABR[R122T] mutation, homologous to

THR228 in MpGABR, the Clustered Regularly Interspaced Short Palindromic Repeats-Cas9

(CRISPR/Cas9) genomic editing strategy was used. First, CRISPR Optimal Target Finder

(http://targetfinder.flycrispr.neuro.brown.edu/) was used to generate short guide RNA

(sgRNA) targets without off-target hits. Then, the Precision gRNA Synthesis Kit (Thermo

Scientific) was used to synthesize the sgRNA. The 228 bp homologous arm in R122T, for

homology-directed repair, was subsequently employed to synthetize single-stranded DNA

(ssDNA) bearing the desired mutation (DmGABR[R122T]). Subsequently, 1 nL of a mixture

containing 160 ng· μL^{-1} ssDNA, 300 ng· μL^{-1} both sgRNAs, and 90 ng· μL^{-1} Cas9 protein was

injected into *Drosophila* embryos via the Drummond Nanoject III (Broomall, PA, USA) system.

CRISPR/Cas9-induced mutations were verified by amplifying and sequencing genomic DNA

flanking the genome editing target site. Finally, according to the traditional topical application

and bioassay approach described previously (Zhou et al. 2023), the acute insect toxicity of

betulin in genome-modified or nonmodified (WT) 3-day-old adult flies was determined. The

oligonucleotide sequences used for PCR, the sgRNA synthesis template and the fragment of

donor DNA used for homology-directed repair (HDR) are listed in Table S3.

2.12 Microscale thermophoresis (MST) assay

The binding affinity of MpGABR for betulin was measured via a NanoTemper Technologies Monolith Kit (MO, Munich, Germany) according to the manufacturer's specifications. Briefly, MpGABR was fluorescently labeled via a protein labeling kit (RED-NHS 2nd generation, Monolith™) and then mixed with sixteen concentrations (from 5 nM to 60 mM) of betulin. The mixtures were subsequently loaded into NanoTemper capillaries for MST measurements with 40% LED and 60% MST power at 25 °C. MO Control NanoTemper Technologies GmbH software (v2.3) and MO Affinity Analysis software (v2.3, NanoTemper Technologies GmbH) were used to acquire and analyze the recorded data, respectively. Three replicates were set for each concentration. The dissociation constant (K_d), indicating the binding affinity, was calculated using data from three replicates for each experiment.

2.13 Electrophysiological recording

In accordance with previously described methods (Chen et al. 2019), oocytes isolated from mature healthy *Xenopus* females were digested at 25 °C for 1 h to remove the follicular layer. Individual oocytes were selected to express *MpGABR* and incubated at 18 °C with 1× Ringer's solution (96 mM NaCl, 5 mM $MgCl_2$, 5 mM HEPES, 2 mM KCl, and 0.8 mM $CaCl_2$) supplemented with 550 mg·mL^{-1} sodium pyruvate, 100 mg·mL^{-1} streptomycin, 50 mg·mL^{-1} tetracycline, and 5% dialyzed horse serum. After 3 days of incubation, the oocytes were used for electrophysiological recording. GABA was dissolved in 1 × Ringer's solution to prepare 100 μM GABA solution. Subsequently, the 100 μM GABA solutions containing different concentrations of betulin (0, 5, 10, 20, 40, 80, 160, 320 μM) were used to perfuse the oocytes. The whole-cell currents were recorded via a two-electrode voltage-clamp. The data were collected

and analyzed using Digidata 1440A and pCLAMP 10.2 software (Axon Instruments Inc., Union

City, CA, USA), respectively (Xu et al. 2020).

2.14 Statistical analysis

Statistical analysis was performed using SPSS (v22). Statistical significance was

determined using one-way analysis of variance (ANOVA) for differences among groups. The

replicate numbers are indicated by the black dots in the figures.

3. RESULTS

3.1 Control efficacy of betulin against aphids

Bioassays revealed that the LC_{50} values of betulin and pymetrozine against *M. persicae* at

48 h were 0.1641 and 1.0612 mg·mL^{-1}, respectively (Table S1). To assess the control efficacy of

betulin, tobacco plants infected with *M. persicae* were exposed to betulin (0.1641 mg·mL^{-1}) and

pymetrozine (1.0612 mg·mL^{-1}, as a positive control) in both the greenhouse and field. In the

greenhouse test (Fig. 1A-D), at 14 days after treatment, *M. persicae* in the CK group (negative

control) reproduced rapidly, and tobacco leaves turned yellow and wrinkled, whereas tobacco

leaves in the betulin and pymetrozine treatment groups were barely affected by the aphids (Fig.

1A-B). The control efficacy of betulin against *M. persicae* was significantly greater than that of

pymetrozine at 1 day after treatment ($P < 0.01$), while there were no significant differences

between the control efficacies of betulin and pymetrozine at 5, 9, or 14 days after treatment (Fig.

1E). After 14 days of treatment, the control efficacies of betulin and pymetrozine reached 91.54%

and 95.15%, respectively. Throughout the field test (Fig. 1F), no difference in the control efficacy

was observed between betulin and pymetrozine, with values of 87.79% and 90.14%,

respectively, after 14 days of treatment (Fig. 1G). These results suggest that betulin has

immense potential for development as a commercial aphid insecticide like pymetrozine.

3.2 Expression patterns of genes in aphids after treatment with betulin

To investigate the expression patterns of genes in *M. persicae* after exposure to betulin,

RNA-seq was performed on *M. persicae* with 0.1641 mg·mL^{-1} betulin treatment for 48 h and

without (CK) (Fig. 2A). Principal coordinate analysis (PCA) revealed that the expressed genes

were strongly clustered in the CK group and betulin group (Fig. 2B). Compared with those in

the CK group, there were 130 up- and 41 downregulated significant differentially expressed

genes (DEGs) in *M. persicae* after betulin treatment (Fig. 2C and D, Table S4). To validate the

RNA–seq results, fifteen DEGs were randomly selected with different expression levels. The

relative expression trends of the fifteen DEGs determined via qRT–PCR were similar to those

detected via RNA–seq, supporting the reliability of the RNA–seq data (Fig. 2E). To further

analyze the functions of the DEGs, GO (Gene Ontology) and KEGG (Kyoto Encyclopedia of

Genes and Genomes) enrichment analyses were performed. GO term enrichment

demonstrated that the DEGs were enriched mainly in the GABA$_A$ receptor activity term (Fig.

2F). Interestingly, KEGG enrichment also found that the DEGs were enriched in the GABAergic

signaling related pathway and GABAergic synapse terms (Fig. 2G). Additionally, the

expression of DEGs related to GABR, including *MpGABR* (encoding GABA$_A$ receptor),

MpGABRAP (encoding GABA$_A$ receptor-associated protein) and *MpGABRB* (encoding GABA$_A$

receptor β subunit), in the betulin group was significantly lower than that in the CK group (Fig.

2H). In particular, the log$_2$(fold change) value of *MpGABR* was the lowest, at -3.8 (Table S4).

These results suggested that *MpGABR* may be a candidate target for betulin against *M. persicae*.

3.3 *MpGABR* expression was inhibited by betulin

Analysis of the sequence information revealed that the MpGABR, MpGABRAP and

MpGABRB proteins have 694, 118, and 250 amino acids, respectively, with calculated molecular

weights of 77.20, 14.08, and 28.01 kDa and isoelectric points of 9.85, 9.63, and 7.20, respectively

(Table S5). Additionally, structure prediction indicated that MpGABR, MpGABRAP and

MpGABRB contain 4, 0, and 1 transmembrane helical domain, respectively (Fig. 3A), implying

that only MpGABR has a complete transmembrane structure.

To further verify the inhibitory effect of betulin on GABR-related gene expression, the

relative expression of the three GABR-related genes was detected in *M. persicae* exposed to the

LC_{30}, LC_{50}, and LC_{70} of betulin for 48 h. The relative expression of all three genes decreased

gradually as the concentration of betulin increased (Fig. 3B-D). After exposure to the LC_{30} of

betulin for 48 h, the relative expression of *MpGABR, MpGABRAP* and *MpGABRB* decreased by

82.91%, 10.53%, and 11.49%, respectively. These results indicated that *MpGABR* was the most

sensitive to betulin. To further investigate the effect of betulin on the MpGABR protein, *M.*

persicae was exposed to betulin concentrations of LC_{30}, LC_{50}, and LC_{70} for 48 h. As the

concentration of betulin increased, the MpGABR protein content in *M. persicae* gradually

decreased (Fig. 3E and F). Compared with that in the CK group, the MpGABR protein content

in *M. persicae* exposed to betulin concentrations of LC_{30}, LC_{50}, and LC_{70} decreased by 33.68%,

44.89%, and 76.89%, respectively.

Furthermore, after these three genes were silenced via RNAi (Fig. 4D), the expression

levels of *MpGABR, MpGABRAP* and *MpGABRB* in *M. persicae* significantly decreased by 64.57%,

63.28%, and 67.33%, respectively, compared with those in the control groups (DEPC-treated

water and dsGFP) ($P < 0.0001$, Fig. 4A-C). After exposure to the LC_{50} of betulin for 48 h, the

mortality of *M. persicae* with *MpGABR* silenced markedly increased ($P < 0.001$) by 30.44%,

compared with that in the control groups, whereas the mortalities of *M. persicae* in the

MpGABRAP- and *MpGABRB*-silenced groups did not significantly differ from that in the

control groups (Fig. 4E). Additionally, western blotting analysis revealed that MpGABR

protein expression significantly decreased by 69.91% after RNAi ($P < 0.0001$, Fig. S1A, B). These

results further implied that *MpGABR* might be a target for betulin against *M. persicae*.

3.4 Phylogenetic analysis of GABRs in insects

Furthermore, phylogenetic analysis of GABRs in insects was performed using the amino

acid sequences of 70 GARB proteins from Hemiptera, Diptera, Lepidoptera, Thysanoptera,

Hymenoptera, and Coleoptera. MpGABR (XP_022173711.1) was found to be genetically closely

related to CAI6365831.1 from *Macrosiphum euphorbiae*, XP 008183008.2 from *Acyrthosiphon pisum*,

and XP 060864885.1 from *Metopolophium dirhodum* (Fig. 5 and Table S6), indicating that

MpGABR may be functionally analogous to CAI6365831.1, XP 008183008.2, and XP 060864885.1.

3.5 Interaction between MpGABR and betulin

The WT (wild type) MpGABR was expressed in *Escherichia coli* (Fig. 6A). The binding

affinity of betulin with WT MpGABR was further measured using MST. The K_d value was 2.24

μM (Fig. 6B), suggesting that betulin could bind to MpGABR. Additionally, voltage-clamp-

based electrophysiological recording demonstrated that MpGABR strongly responded to

betulin (Fig. 6C). In the presence of only GABA, a fast inward current was generated. As the

concentration of added betulin increased, the current and channel activity of MpGABR

decreased correspondingly, with an EC_{50} value of 20.66 μM (Fig. 6C and D). These results

indicated that betulin acted as an inhibitor of MpGABR.

3.6 Key sites for the binding of betulin to MpGABR

To further study the binding mode of betulin in the active pocket of MpGABR, molecular

docking was performed to explore the structure–function relationships between betulin and

MpGABR. The docking results revealed that the predicted binding energy between betulin and

MpGABR was −6.38 kcal·mol^{-1} (Table S7), verifying that betulin could act as a specific ligand

for MpGABR. The three-dimensional binding pattern of betulin with MpGABR indicated that

the four key amino acid residues (ARG224, ALA226, PHE227 and THR228) interacted with

betulin in the MpGABR binding pocket (Fig. 7A). Among the four key amino acid residues,

ALA226 and THR228 interacted with betulin via hydrogen bonding in the MpGABR binding

pocket (Table S7). Betulin generated nonconventional H-bonds (C…H) with ALA226 (3.31 Å)

and conventional H-bonds (C-O H) with THR228 (2.16 Å). Furthermore, sequence alignment

of the key amino acid residues of GABR in different species from Hemiptera, Diptera,

Lepidoptera, Hymenoptera, Thysanoptera, and Coleoptera indicated that, among the four

residues, only THR228 was conserved across 11 species in the Aphididae family of Hemiptera

(Fig. 7B), implying that THR228 may be an essential specific site for the action of betulin on

MpGABR in aphids. Mutants of aphid MpGABR were subsequently constructed, including

R224A (ARG224 replaced by ALA), A226T (ALA226 replaced by THR), F227Y (PHE227

replaced by TYR, similar to the site in *Drosophila*), or T228R (THR228 replaced by ARG, similar

to the site in *Drosophila*) (Fig. 7C). The binding affinity of betulin with these mutants was

measured using MST. The results revealed that the K_d values of R224A, A226T, F227Y, T228R,

and WT were 2.31, 2.27, 2.25, 5321, and 2.24 μM, respectively (Fig. 7D and E, Table S8). This

result suggested that the mutation at the THR228 site, with the highest K_d value, markedly

reduced the binding ability of betulin to MpGABR, implying that the THR228 site may be an

essential specific site for the binding of betulin to MpGABR.

3.7 Betulin binds specifically to MpGABR via THR228

To further prove that THR228 is the specific binding site for betulin in MpGABR, the WT

and mutant (R122T, equivalent to THR228 in MpGABR) *Drosophila* DmGABR proteins were

expressed in *E. coli* and extracted (Fig. 8A and D). The binding affinities of betulin and

pymetrozine with DmGABR^{WT} and DmGABR^{R122T} were evaluated using MST. As shown in Fig.

8B, betulin was able to bind to DmGABR^{R122T} (K_d = 342.4 μM) but not to DmGABR^{WT}. However,

pymetrozine exhibited no binding affinity for DmGABR^{R122T} or DmGABR^{WT} (Fig. 8C).

Moreover, to generate *Drosophila* mutant individuals carrying DmGABR^{R122T}, the CRISPR/Cas9

genomic editing strategy was used. After exposure to different concentrations of betulin, the

mortality rate of DmGABR^{R122T} *Drosophila* was significantly greater than that of DmGABR^{WT}

Drosophila (Fig. 8E). In addition, the LD$_{50}$ value of betulin against DmGABR^{R122T} *Drosophila* at 72

h was 27.19 μg·fly^{-1}, whereas the LD$_{50}$ value of betulin against DmGABR^{WT} *Drosophila* could not

be calculated because the lethal concentration exceeded 1000 μg·fly^{-1} (Table S9). Additionally,

there was no significant difference in mortality between DmGABR^{WT} and DmGABR^{R122T}

Drosophila after exposure to pymetrozine (Fig. 8F). The LD$_{50}$ values of pymetrozine against

DmGABR^{WT} and DmGABR^{R122T} *Drosophila* at 72 h were 0.23 and 0.24 μg·fly^{-1}, respectively (Table

S9). Moreover, THR228 is located at the neurotransmitter-gated ion-channel ligand-binding

domain in MpGABR (Fig. S2), implying that betulin may play a role as a competitive antagonist

of MpGABR. Taken together, these findings suggested that betulin bound specifically to

MpGABR via THR228, resulting in the death of *M. persicae*.

4. DISCUSSION

Although chemical pesticides are effective in pest control, the long-term unreasonable application of these substances has led the emergence of resistant pests, environmental deterioration, and deleterious effects on nontarget organisms, including humans, raising widespread and intense concerns (Devi et al. 2022). These challenges have made it urgent for the development of alternative strategies for pest control. Recently, the utilization of plant secondary metabolites as insecticides has become increasingly popular as an eco-friendly and biocontrol approach (Ling et al. 2022; Ayilara et al. 2023; Jahan et al. 2025b). Our previous study revealed that *P. davidiana*, a wild relative of cultivated peach, strongly resists *M. persicae* by accumulating high contents of betulin (Wang et al. 2022a; Wang et al. 2024). In addition, betulin, a lupane-type triterpene, possesses potent insecticidal activity and is a promising substance for the development of novel insecticides for aphid control. In this study, the insecticidal effect of betulin was further evaluated by comparing the control efficacy of betulin with that of pymetrozine against aphids in greenhouses and fields (Fig. 1A-G). These results indicate that betulin has a similar control effect to pymetrozine and has immense potential for development as a plant-derived insecticide.

Terpenes are a diverse group of plant secondary metabolites that can increase the resistance of plants to insect herbivores through direct (Wang et al. 2025b) and indirect (Wang et al. 2025a) defense mechanisms. In direct defense against herbivores, triterpenes play important roles in diverse biological activities, including antiparasitic, insecticidal, and antifeedant activities (Tian et al. 2021; Ivana et al. 2023). Azadirachtin, a tetracyclic triterpenoid compound isolated from the Indian neem tree (*Azadirachta indica*), is one of the most prominent

commercial biopesticides, exhibiting strong insect antifeedant properties as well as growth-

and reproduction-regulating effects (Dawkar et al. 2019; Bae et al. 2022). Besides, triterpene

glycoside compounds play crucial roles in the defense of tobacco (*N. attenuata*) against tobacco

hornworm (*Manduca sexta*) larvae (Yang et al. 2023). Although betulin exhibits various

pharmacological activities (Amiri et al. 2020; Li et al. 2022; Yan 2022; Lv 2023), reports on its

insecticidal activity are limited. Encouragingly, the effects of betulin and its derivatives on pests

have attracted increasing attention. Betulinic acid and its derivatives showed larvicidal activity

against *Aedes aegypti* larvae (da Silva et al. 2016). Betulin-cinnamic acid-related hybrid

compound 5b exhibited strong aphicidal activity, and compound 2l could destroy the

ultrastructure of midgut cells and significantly inhibit the activity of α-amylase in

diamondback moth (*Plutella xylostella* L.) larvae (Huang et al.). Our previous studies also

indicated that betulin possesses potent insecticidal activity and is a key endogenous secondary

metabolite related to the defense of peach against *M. persicae* (Wang et al. 2022a). Elucidating

the insecticidal mechanism of betulin against aphids will provide a basis for the development

of novel aphicides and sustainable strategies for aphid control.

GABA receptors have been confirmed to be targets of terpenoids that impair insect

neuronal function in herbivores (Guo et al. 2023). GABRs are heteropentameric ligand-gated

ion channels in the central nervous system that conduct chloride and bicarbonate ions. These

receptors are targets of numerous drugs for the treatment of neuropsychiatric disorders

(Thompson 2024). A variety of terpenoids act as positive allosteric modulators or

noncompetitive antagonists of GABRs, such as diterpenoids (isopimaric acid and miltirone),

sesquiterpenoids (picrotoxin, bilobalide and ginkgolides) and monoterpenoids (α-thujone and

thymol) (Guo et al. 2023). In this study, both GO and KEGG enrichment analyses revealed that

the DEGs identified by RNA-seq were enriched in GABAergic signaling-related pathways (Fig.

2F and G). Additionally, the expression of the DEGs related to GABRs, particularly *MpGABR*

(Table S4), in the betulin group was significantly lower than that in the CK group (Fig. 2H).

Besides, the relative expression of *MpGABR*, *MpGABRAP* and *MpGABRB* decreased gradually

after *M. persicae* was exposed to the LC_{30}, LC_{50}, and LC_{70} of betulin for 48 h (Fig. 3B-D). Among

them, *MpGABR* was the most sensitive to betulin, and its expression was reduced by 82.91%

after exposure to the LC_{30} of betulin for 48 h. Furthermore, compared with the control group,

the *M. persicae* group with *MpGABR* silenced by RNAi presented a significant increase in

mortality ($P < 0.001$), by 30.44%, after 48 h of exposure to the LC_{50} of betulin (Fig. 4E).

Collectively, these results suggest that betulin may have insecticidal effects on aphids by

inhibiting *MpGABR* expression. The regulation of gene expression is sophisticated and delicate

(Pope and Medzhitov 2018). The regulatory network controlling *GABR* expression remains

unclear. In adult rats, epileptic seizures have been reported to increase the levels of brain-

derived neurotrophic factor (BDNF), which in turn prompted the transcription factors CREB

and ICER to reduce the gene expression of the GABR α1 subunit (Lund et al. 2008). In *Drosophila*,

it has been demonstrated that WIDE AWAKE, which regulated the onset of sleep, interacted

with the GABR and upregulated its expression level (Liu et al. 2014). In *Drosophila* brain,

circular RNA circ_sxc was found to inhibit the expression of miR-87-3p in the brain through

sponge adsorption, thereby regulating the expression of neurotransmitter receptor ligand

proteins, including GABR, and ensuring the normal function of synaptic signal transmission in

brain neurons (Li et al. 2024). However, it remains unclear how betulin reduces *MpGABR*

expression, and further research is needed.

Additionally, betulin has been reported to be able to bind to GABRs (Manayi et al. 2016).

We further investigated the interaction of betulin with the MpGABR protein. The MST assay

revealed that betulin was able to bind to MpGABR (K_d = 2.24 μM) (Fig. 6B), which is consistent

with previous findings showing that betulin binds to GABA receptors in mouse brains *in vitro*

(Muceniece et al. 2008). Voltage-clamp-based electrophysiological recordings indicated that

betulin acted as an inhibitor (EC_{50} = 20.66 μM) for MpGABR (Fig. 6C and D). Subsequent

molecular docking analysis suggested that four key amino acid residues (ARG224, ALA226,

PHE227 and THR228) interact with betulin in the MpGABR binding pocket (Fig. 7A), among

which merely ALA226 and THR228 interact with betulin via hydrogen bonding (Table S7). The

results of the sequence alignment revealed that only THR228 was conserved across 11 species

in the Aphididae family of Hemiptera (Fig. 7B). Furthermore, evaluation of the ability of betulin

to bind to MpGABR mutants with mutations at those four sites revealed that the ability of

betulin to bind to T228R was significantly weaker than its ability to bind to the WT (Fig. 7D and

E, Table S8), indicating that THR228 is an essential specific site for the binding of betulin to

MpGABR. Moreover, to further prove that THR228 is the specific binding site for betulin in

MpGABR, the binding affinities of betulin with the WT and mutant (R122T, equivalent to

THR228 in MpGABR) *Drosophila* DmGABR proteins were assessed using MST. The results

showed that betulin was able to bind to DmGABR[R122T] (K_d = 342.4 μM) but not DmGABR[WT] (Fig.

8B). Additionally, after exposure to different concentrations of betulin, the mortality rate of

DmGABR[R122T] *Drosophila* was significantly greater than that of DmGABR[WT] *Drosophila* (Fig. 8E

and Table S9). Similarly, a previous study indicated that the R122G amino acid site substitution,

generated via RNA editing, affects the sensitivity of *Drosophila* to fipronil (Es-Salah et al. 2008).

Together, these findings suggest that betulin binds specifically to MpGABR via THR228, acting

as an inhibitor of MpGABR and causing aphid death. Studies on key amino acids that are

crucial for GABR function has primarily focused on transmembrane regions. For instance,

based on the mutational research and *Drosophila* GABR modeling approach, multiple key

amino acids have been identified as insecticide targets in the transmembrane domain (Nakao

and Banba 2021). Guo et al. proposed that amino acid substitutions in transmembrane domain

2 contribute to terpenoid insensitivity during plant-insect coevolution (Guo et al. 2023).

However, these studies have neglected the extracellular domain. Our study signified that

betulin targets the THR228 site in the extracellular domain of MpGABR, which is conserved

only in the Aphididae family. Therefore, betulin is speculated to be a specific insecticidal

substance evolved by plants in response to aphid infestation. Besides, further verification is

needed to determine whether betulin is toxic to other insect species.

GABRs belong to the cysteine loop (Cys)-loop superfamily of neurotransmitter receptors

and are essential heteropentameric ligand-gated ion channels in the central nervous system.

After GABA binds to the extracellular Cys-loop of a GABR (Ashby et al. 2012), the GABR is

activated and opens up its central pore to allow chloride ions to pass through, thereby

hyperpolarizing neurons and attenuating excitatory neurotransmission (Tremblay et al. 2016).

In insects, GABRs play important roles in circadian rhythms (Schellinger et al. 2022), sleep

(Chaturvedi et al. 2022), movement (Eick et al. 2022), and olfactory memory (Yamagata et al.

2021). Additionally, GABR is a critical target for a variety of insecticides and ectoparasiticides.

Insecticides targeting GABRs are categorized into noncompetitive antagonists (NCAs) and

competitive antagonists (CAs) according to their different binding sites. NCAs block chloride

channels in nerve cells by interacting with amino acid residues of the GABA-gated chloride

channel, causing a conformational change in the receptor, which interferes with the normal

function of the central nervous system and ultimately leads to insect death (Nakao and Banba

2021; Guo et al. 2023). A series of first- and second-generation NCA insecticides have been

successfully developed. Picrotoxinin, isolated from *Anamirta cocculus* fruit, is the oldest natural

NCA and is toxic to houseflies (Tong et al. 2023). Among the first-generation NCAs,

polychlorocycloalkanes, including dieldrin and lindane, were commonly used as pesticides in

the middle and late 20th centuries (Hainzl et al. 1998; Tanaka 2019). As second-generation

NCAs, fipronil and its derivatives are commercially available phenylpyrazole insecticides that

have been widely used for agricultural pest control (Sheng et al. 2018; Li et al. 2021).

Additionally, CAs bind to orthogonal binding sites, such as the GABA recognition site in the

extracellular region, competitively inhibiting the binding of GABA to its receptor, leading to

toxic effects in insects. Given the lack of interference from existing insect resistance mechanisms,

CAs hold promise for the development of efficient new insecticides. Novel 1, 6-dihydro-6-

iminopyridazine-derived insecticides, as CAs, have insecticidal properties against common

cutworms and houseflies (Liu et al. 2022). Besides, nootkatone, a sesquiterpenoid, also acts as

a CA to induce insect mortality (Norris et al. 2022). Fluralaner, an isooxazoline ectoparasiticide,

inhibits both parasites and *Aedes aegypti* by acting as a CA (Wang et al. 2022b; Asahi et al. 2023).

Our results revealed that the binding site (THR228) for betulin in MpGABR was located in the

extracellular neurotransmitter-gated ion-channel ligand-binding domain (Fig. S2), implying

that betulin acts as a CA of MpGABR. Although the mechanism by which betulin competes

with GABA for binding to MpGABR requires further experimental validation, our work may

have provided a novel target for developing insecticides. In this study, betulin, on the one hand,

inhibited the expression of *MpGABR* and, on the other hand, specifically bound to MpGABR

through THR228, acting as an inhibitor of MpGABR and causing aphid death (Fig. 9).

The development of bioinsecticides should not only focus on the toxic effects of active

substance on target organisms, but also on their influence on the ecosystem (Haddi et al. 2020).

Although our results indicate that betulin has specific toxicity to aphids, previous studies have

reported that betulin and its derivatives had effects on *Plutella xylostella* L. (Huang et al. 2025),

Aedes aegypti (de Almeida Teles et al. 2024), and *Drosophila melanogaster* (Lee and Min 2024).

Therefore, further research is needed to determine whether there are other insecticidal

mechanisms or off target effects of betulin. Additionally, betulin exhibits a wide range of

pharmacological activities (Amiri et al. 2020), which have been used to treat various diseases,

such as cancer (Lv 2023), glioblastoma (Li et al. 2022), inflammation (Szlasa et al. 2023) and

hyperlipidemia (Tang et al. 2011). Before applying betulin in the field, it is necessary to fully

verify and consider whether betulin has any impact on farmers' health. Furthermore, will

betulin cause residue or diffusion during the field application process? Will long-term

application promote the evolution of resistance to aphids or other insects? These issues also

need further experimental verification. In summary, before any field application, further

research on the environmental behavior, degradation process, and safety of betulin is needed.

5. CONCLUSION

Betulin, a key metabolite in the aphid-resistant wild peach *P. davidiana*, possesses potent

aphicidal effects on *M. persicae.* This study confirmed that betulin exhibited excellent control

efficacy against *M. persicae* in both greenhouse and field experiments. RNA-seq, qRT–PCR and

western blotting assays revealed that betulin significantly inhibited the expression of *MpGABR*

in aphids. In addition, RNAi-mediated silencing of *MpGABR* markedly increased the

sensitivity of aphids to betulin. Moreover, MST and voltage-clamp-based electrophysiological

recording assays indicated that betulin was able to bind to MpGABR (K_d = 2.24 μM) and acted

as an inhibitor (EC$_{50}$ = 20.66 μM) of MpGABR. Molecular docking analyses suggested that the

amino acid residue THR228, which is highly conserved across 11 species in the Aphididae

family of Hemiptera, might be a critical specific binding site for betulin in MpGABR.

Mutagenesis and genome editing assays revealed that betulin bound specifically to this amino

acid residue in aphids but not in *Drosophila*, resulting in aphid death. Collectively, the results

suggest that the aphicidal effects of betulin on aphids occur in a two-pronged manner: on the

one hand, betulin inhibits *MpGABR* expression; on the other hand, it specifically binds to

MpGABR via THR228 and acts as an inhibitor of MpGABR. Elucidating the insecticidal

mechanism of betulin against aphids will provide a basis for the development of novel

insecticides and sustainable strategies for aphid control.

ASSOCIATED CONTENT

Supporting Information

Protein expression of MpGABR after RNAi (Figure S1); Betulin binding site on the domain of

MpGABR (Figure S2); LC$_{50}$ values of betulin and pymetrozine against *M. persicae* at 48 h (Table

S1); Primers used for cloning, RNAi, heterologous expression, and qRT-PCR (Table S2);

Oligonucleotide sequences for PCR, the sgRNA synthesis template and the fragment of donor

DNA for homology-directed repair (HDR) (Table S3); Screening of genes significantly

differentially expressed in *M. persicae* in response to betulin at 48 h post treatment (Table S4);

Complete sequence information for the GABA$_A$ receptor gene of *M. persicae* (Table S5);

Sequences and relevant information for phylogenetic analysis of GABA$_A$ receptor (Table S6);

Binding energy and nonbonding interactions between betulin and GABA$_A$ receptor (Table S7);

Parameters of dose-response curves from microscale thermophoresis experiments (Table S8);

LD$_{50}$ values of betulin and pymetrozine against *D. melanogaster* at 72 h (Table S9).

AUTHOR INFORMATION

Corresponding author:

Hong Zhou – College of Plant Protection, Southwest University, Chongqing, 400715, China. E-

mail: Hong Zhou: zhouhong1990@swu.edu.cn

Authors

Junxiu Wang – Key Laboratory of Agricultural Biosafety and Green Production of Upper

Yangtze River (Ministry of Education), Chongqing Key Laboratory of Forest Ecological

Restoration and Utilization in the Three Gorges Reservoir Area, College of Horticulture and

Landscape Architecture, Southwest University, Beibei, Chongqing 400715, China

Matthana Klakong – College of Plant Protection, Southwest University, Chongqing, 400715,

China

Qiuyu Zhu – College of Plant Protection, Southwest University, Chongqing, 400715, China

Jinting Pan – Key Laboratory of Agricultural Biosafety and Green Production of Upper Yangtze

River (Ministry of Education), Chongqing Key Laboratory of Forest Ecological Restoration and

Utilization in the Three Gorges Reservoir Area, College of Horticulture and Landscape

Architecture, Southwest University, Beibei, Chongqing 400715, China

Yudie Duan – Key Laboratory of Agricultural Biosafety and Green Production of Upper

Yangtze River (Ministry of Education), Chongqing Key Laboratory of Forest Ecological

Restoration and Utilization in the Three Gorges Reservoir Area, College of Horticulture and

Landscape Architecture, Southwest University, Beibei, Chongqing 400715, China

Lirong Wang – Zhengzhou Fruit Research Institute, Chinese Academy of Agricultural Sciences,

Zhengzhou, 451100, China

Yong Li – Zhengzhou Fruit Research Institute, Chinese Academy of Agricultural Sciences,

Zhengzhou, 451100, China

Jiangbo Dang –Key Laboratory of Agricultural Biosafety and Green Production of Upper

Yangtze River (Ministry of Education), Chongqing Key Laboratory of Forest Ecological

Restoration and Utilization in the Three Gorges Reservoir Area, College of Horticulture and

Landscape Architecture, Southwest University, Beibei, Chongqing 400715, China

Danlong Jing – Key Laboratory of Agricultural Biosafety and Green Production of Upper

Yangtze River (Ministry of Education), Chongqing Key Laboratory of Forest Ecological

Restoration and Utilization in the Three Gorges Reservoir Area, College of Horticulture and

Landscape Architecture, Southwest University, Beibei, Chongqing 400715, China

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China

(32402492, 32302394), Special Funding for Chongqing Postdoctoral Research Project

(2312013543265057), Sichuan Science and Technology Program (2025ZNSFSC1120), and

Fundamental Research Fund for the Central Universities of China (SWU-KR25018).

ABBREVIATIONS

GABA, Gamma-aminobutyric acid; GABR, GABA$_A$ receptor; GABRB, GABA$_A$ receptor β subunit; GABRAP, GABA$_A$ receptor associated protein; MST, microscale thermophoresis; PCA, principal coordinates analysis; DEGs, differentially expressed genes; GO, Gene Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; K_d, dissociation constant; RDL, resistance to dieldrin; NCA, noncompetitive antagonists; CA, competitive antagonists.

REFERENCES

Alonso Salces RM, Berrueta LA, Abad García B, Sasía Arriba A, Asensio Regalado C, Dapena E, Gallo B. 2022. Untargeted metabolomic liquid chromatography high-resolution mass spectrometry fingerprinting of apple cultivars for the identification of biomarkers related to resistance to rosy apple aphid. *Journal of Agricultural and Food Chemistry* **70**: 13071-13081.

Amiri S, Dastghaib S, Ahmadi M, Mehrbod P, Khadem F, Behrouj H, Aghanoori M, Machaj F, Ghamsari M, Rosik J, Hudecki A, Afkhami A, Hashemi M, Los M, Mokarram P, Madrakian T, Ghavami S. 2020. Betulin and its derivatives as novel compounds with different pharmacological effects. *Biotechnology Advances* **38**: 107409.

Asahi M, Yamato K, Ozoe F, Ozoe Y. 2023. External amino acid residues of insect GABA receptor channels dictate the action of the isoxazoline ectoparasiticide fluralaner. *Pest Management Science* **79**: 4078-4082.

Ashby Jamie A, McGonigle Ian V, Price Kerry L, Cohen N, Comitani F, Dougherty Dennis A, Molteni C, Lummis Sarah CR. 2012. GABA binding to an insect GABA receptor: a molecular dynamics and mutagenesis study. *Biophysical Journal* **103**: 2071-2081.

Ayilara MS, Adeleke BS, Akinola SA, Fayose CA, Adeyemi UT, Gbadegesin LA, Omole RK,

Johnson RM, Uthman QO, Babalola OO. 2023. Biopesticides as a promising alternative to

synthetic pesticides: A case for microbial pesticides, phytopesticides, and nanobiopesticides.

Frontiers in Microbiology **14**: 1040901.

Bae M, Lewis A, Liu S, Arcot Y, Lin YT, Bernal JS, Cisneros Zevallos L, Akbulut M. 2022.

Novel biopesticides based on nanoencapsulation of azadirachtin with whey protein to control

fall armyworm. *Journal of Agricultural and Food Chemistry* **70**: 7900-7910.

Bass C, Puinean AM, Zimmer CT, Denholm I, Field LM, Foster SP, Gutbrod O, Nauen R,

Slater R, Williamson MS. 2014. The evolution of insecticide resistance in the peach potato

aphid, Myzus persicae. *Insect Biochemistry and Molecular Biology* **51**: 41-51.

Chaturvedi R, Stork T, Yuan C, Freeman MR, Emery P. 2022. Astrocytic GABA transporter

controls sleep by modulating GABAergic signaling in *Drosophila* circadian neurons. *Current*

Biology **32**: 1895-1908.

Chen Z, Yao X, Dong F, Duan H, Shao X, Chen X, Yang T, Wang G, Zheng Y. 2019. Ecological

toxicity reduction of dinotefuran to honeybee: New perspective from an enantiomeric level.

Environment International **130**: 104854.

da Silva GN, Trindade FT, dos Santos F, Gosmann G, e Silva AA, Gnoatto SC. 2016.

Larvicidal activity of natural and modified triterpenoids against *Aedes aegypti* (Diptera:

Culicidae). *Pest Management Science* **72**: 1883-1887.

Dawkar VV, Barage SH, Barbole RS, Fatangare A, Grimalt S, Haldar S, Heckel DG, Gupta

VS, Thulasiram HV, Svatoš A, Giri AP. 2019. Azadirachtin-a from *Azadirachta indica* impacts

multiple biological targets in cotton bollworm *Helicoverpa armigera*. *ACS Omega* **4**: 9531-9541.

de Almeida Teles AC, dos Santos BO, Santana EC, Durço AO, Conceição LSR, Roman

Campos D, de Holanda Cavalcanti SC, de Souza Araujo AA, dos Santos MRV. 2024.

Larvicidal activity of terpenes and their derivatives against *Aedes aegypti*: a systematic review

and meta-analysis. *Environmental Science and Pollution Research* **31**: 64703-64718.

Devi PI, Manjula M, Bhavani RV. 2022. Agrochemicals, environment, and human health.

Annual Review of Environment and Resources **47**: 399-421.

Eick AK, Ogueta M, Buhl E, Hodge JJL, Stanewsky R. 2022. The opposing chloride

cotransporters KCC and NKCC control locomotor activity in constant light and during long

days. *Current Biology* **32**: 1420-1428

Es Salah Z, Lapied B, Le Goff G, Hamon A. 2008. RNA editing regulates insect gamma-

aminobutyric acid receptor function and insecticide sensitivity. *NeuroReport* **19**: 939-943.

Fan C, Cowgill J, Howard RJ, Lindahl E. 2024. Divergent mechanisms of steroid inhibition in

the human ϱ1 GABAA receptor. *Nature Communications* **15**: 7795.

Ffrench Constant RH, Mortlock DP, Shaffer CD, MacIntyre RJ, Roush RT. 1991. Molecular

cloning and transformation of cyclodiene resistance in *Drosophila*: an invertebrate gamma-

aminobutyric acid subtype A receptor locus. *Proceedings of the National Academy of Sciences* **88**:

7209-7213.

Guo L, Qiao X, Haji D, Zhou T, Liu Z, Whiteman NK, Huang J. 2023. Convergent resistance

to GABA receptor neurotoxins through plant–insect coevolution. *Nature Ecology & Evolution* **7**:

1444-1456.

Guo Y, Ji N, Bai L, Ma J, Li Z. 2022. Aphid viruses: A brief view of a long history. *Frontiers in*

Insect Science **2**: 846716.

Haddi K, Turchen LM, Viteri Jumbo LO, Guedes RN, Pereira EJ, Aguiar RW, Oliveira EE.

2020. Rethinking biorational insecticides for pest management: unintended effects and

consequences. *Pest Management Science* **76**: 2286-2293.

Hainzl D, Cole LM, Casida JE. 1998. Mechanisms for selective toxicity of fipronil insecticide

and it sulfone metabolite and desulfinyl photoproduct. *Chemical Research in Toxicology* **11**: 1529-

1535.

Huang X, Hao N, Shu L, Wei Z, Shi J, Tian Y, Chen G, Yang X, Che Z. 2025. Preparation and

insecticidal activities of betulin-cinnamic acid-related hybrid compounds and insights into the

stress response of *Plutella xylostella* L. *Pest Management Science* **81**: 4243-4255.

Ivana ZK, Marina PS, Jovana JA, Andrea RN. 2023. Steroid and triterpenoid compounds with

antiparasitic properties. *Current Topics in Medicinal Chemistry* **23**: 791-815.

Jahan T, Huda M, Zhang K, He Y, Lai D, Dhami N, Quinet M, Ali M, Kreft I, Woo S, Georgiev

M, Fernie A, Zhou M. 2025a. Plant secondary metabolites against biotic stresses for sustainable

crop protection. *Biotechnology Advances* **79**: 108520.

Jahan T, Huda MN, Zhang K, He Y, Lai D, Dhami N, Quinet M, Ali MA, Kreft I, Woo SH,

Georgiev MI, Fernie AR, Zhou M. 2025b. Plant secondary metabolites against biotic stresses

for sustainable crop protection. *Biotechnology Advances* **79**: 108520.

Jiménez Jiménez S, Casado N, García MÁ, Marina ML. 2019. Enantiomeric analysis of

pyrethroids and organophosphorus insecticides. *Journal of Chromatography A* **1605**: 360345.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory

requirements. *Nature Methods* **12**: 357-360.

Lee HY, Min KJ. 2024. Betulinic acid increases the lifespan of *Drosophila melanogaster* via Sir2

and FoxO activation. *Nutrients* **16**: 441.

Li BJ, Wang KK, Chen DP, Yan Y, Cai XL, Chen HM, Dong K, Lin F, Xu HH. 2021. Distinct roles of two RDL GABA receptors in fipronil action in the diamondback moth (*Plutella xylostella*). *Insect Science* **28**: 1721-1733.

Li Q, Wang L, Tang C, Wang X, Yu Z, Ping X, Ding M, Zheng L. 2024. Adipose tissue exosome circ_sxc mediates the modulatory of adiposomes on brain aging by inhibiting brain dme-miR-87-3p. *Molecular Neurobiology* **61**: 224-238.

Li Y, Wang Y, Gao L, Tan Y, Cai J, Ye Z, Chen A, Xu Y, Zhao L, Tong S, Sun Q, Liu B, Zhang S, Tian D, Deng G, Zhou J, Chen Q. 2022. Betulinic acid self-assembled nanoparticles for effective treatment of glioblastoma. *Journal of Nanobiotechnology* **20**: 39.

Ling W, Kaliaperumal K, Huang M, Liang Y, Ouyang Z, Zhou Z, Jiang Y, Zhang J. 2022. Pomelo seed oil: Natural insecticide against cowpea aphid. *Frontiers in Plant Science* **13**: 1048814.

Liu G, Zhou C, Zhang Z, Wang C, Luo X, Ju X, Zhao C, Ozoe Y. 2022. Novel insecticidal 1,6-dihydro-6-iminopyridazine derivatives as competitive antagonists of insect RDL GABA receptors. *Pest Management Science* **78**: 2872-2882.

Liu S, Lamaze A, Liu Q, Tabuchi M, Yang Y, Fowler M, Bharadwaj R, Zhang J, Bedont J, Blackshaw S, Lloyd Thomas E, Montell C, Sehgal A, Koh K, Wu Mark N. 2014. WIDE AWAKE mediates the circadian timing of sleep onset. *Neuron* **82**: 151-166.

Lund IV, Hu Y, Raol YH, Benham RS, Faris R, Russek SJ, Brooks Kayal AR. 2008. BDNF selectively regulates GABA$_A$ receptor transcription by activation of the JAK/STAT pathway. *Science Signaling* **1**: ra9.

Lv M, Zheng Y, Wu J, Shen Z, Guo B, Hu G, Huang Y, Zhao J, Qian Y, Su Z, Wu C, Xue X,

Liu H, Mao Z. 2023. Evoking ferroptosis by synergistic enhancement of a cyclopentadienyl iridium-betulin immune agonist. *Angewandte Chemie International Edition* **62**: e202312897.

Ma R, Haji Ghassemi O, Ma D, Jiang H, Lin L, Yao L, Samurkas A, Li Y, Wang Y, Cao P, Wu S, Zhang Y, Murayama T, Moussian B, Van Petegem F, Yuchi Z. 2020. Structural basis for diamide modulation of ryanodine receptor. *Nature Chemical Biology* **16**: 1246-1254.

Manayi A, Nabavi SM, Daglia M, Jafari S. 2016. Natural terpenoids as a promising source for modulation of GABAergic system and treatment of neurological diseases. *Pharmacological Reports* **68**: 671-679.

Martins FCOL, Batista AD, Melchert WR. 2021. Current overview and perspectives in environmentally friendly microextractions of carbamates and dithiocarbamates. *Comprehensive Reviews in Food Science and Food Safety* **20**: 6116-6145.

Muceniece R, Saleniece K, Rumaks J, Krigere L, Dzirkale Z, Mezhapuke R, Zharkova O, Klusa V. 2008. Betulin binds to γ-aminobutyric acid receptors and exerts anticonvulsant action in mice. *Pharmacology Biochemistry and Behavior* **90**: 712-716.

Nakao T, Banba S. 2021. Important amino acids for function of the insect Rdl GABA receptor. *Pest Management Science* **77**: 3753-3762.

Norris EJ, Chen R, Li Z, Geldenhuys W, Bloomquist JR, Swale DR. 2022. Mode of action and toxicological effects of the sesquiterpenoid, nootkatone, in insects. *Pesticide Biochemistry and Physiology* **183**: 105085.

Pope SD, Medzhitov R. 2018. Emerging principles of gene expression programs and their regulation. *Molecular Cell* **71**: 389-397.

**Schellinger JN, Sun Q, Pleinis JM, An SW, Hu J, Mercenne G, Titos I, Huang CL, Rothenfluh

A, Rodan AR. 2022. Chloride oscillation in pacemaker neurons regulates circadian rhythms

through a chloride-sensing WNK kinase signaling cascade. *Current Biology* **32**: 1429-1438.

Sheng CW, Casida JE, Durkin KA, Chen F, Han ZJ, Zhao CQ. 2018. Fiprole insecticide

resistance of *Laodelphax striatellus*: electrophysiological and molecular docking characterization

of A2′N RDL GABA receptors. *Pest Management Science* **74**: 2645-2651.

Soujanya PL, Sekhar JC, Karjagi CG, Ratnavathi CV, Venkateswarlu R, Yathish KR, Suby SB,

Sunil N, Rakshit S. 2023. Role of morphological traits and cell wall components in imparting

resistance to pink stem borer, *Sesamia inferens* Walker in maize. *Frontiers in Plant Science* **14**:

1167248.

Stará J, Hovorka T, Horská T, Zusková E, Kocourek F. 2024. Pyrethroid and carbamate

resistance in Czech populations of *Myzus persicae* (Sulzer) from oilseed rape. *Pest Management*

Science **80**: 2342-2352.

Szlasa W, Ślusarczyk S, Nawrot Hadzik I, Abel R, Zalesińska A, Szewczyk A, Sauer N,

Preissner R, Saczko J, Drąg M, Poręba M, Daczewska M, Kulbacka J, Drąg Zalesińska M.

2023. Betulin and its derivatives reduce inflammation and COX-2 cctivity in macrophages.

Inflammation **46**: 573-583.

Tanaka K. 2019. Studies on the metabolism, mode of action, and development of insecticides

acting on the GABA receptor. *Journal of Pesticide Science* **44**: 71-86.

Tang JJ, Li JG, Qi W, Qiu WW, Li PS, Li BL, Song BL. 2011. Inhibition of SREBP by a small

molecule, betulin, improves hyperlipidemia and insulin resistance and reduces atherosclerotic

plaques. *Cell Metabolism* **13**: 44-56.

Thompson SM. 2024. Modulators of GABAA receptor-mediated inhibition in the treatment of

neuropsychiatric disorders: past, present, and future. *Neuropsychopharmacology* **49**: 83-95.

Tian X, Li Y, Hao N, Su X, Du J, Hu J, Tian X. 2021. The antifeedant, insecticidal and insect

growth inhibitory activities of triterpenoid saponins from *Clematis aethusifolia* Turcz against

Plutella xylostella (L.). *Pest Management Science* **77**: 455-463.

Tong G, Griffin S, Sader A, Crowell AB, Beavers K, Watson J, Buchan Z, Chen S, Shenvi RA.

2023. C5 methylation confers accessibility, stability and selectivity to picrotoxinin. *Nature*

Communications **14**: 8308.

Tremblay R, Lee S, Rudy B. 2016. GABAergic interneurons in the neocortex: From cellular

properties to circuits. *Neuron* **91**: 260-292.

Troczka BJ, Singh KS, Zimmer CT, Vontas J, Nauen R, Hayward A, Bass C. 2021. Molecular

innovations underlying resistance to nicotine and neonicotinoids in the aphid *Myzus persicae*.

Pest Management Science **77**: 5311-5320.

Tsang SY, Ng SK, Xu Z, Xue H. 2006. The evolution of GABA$_A$ receptor–like genes. *Molecular*

Biology and Evolution **24**: 599-610.

Wang B, Jacquin Joly E, Wang G. 2025a. The role of (*E*)-β-farnesene in tritrophic interactions:

biosynthesis, chemoreception, and evolution. *Annual Review of Entomology* **70**: 313-335.

Wang JX, Li Y, Wang XW, Cao K, Chen CW, Wu JL, Fang WC, Zhu GR, Chen XJ, Guo DD,

Wang J, Zhao YL, Fan JQ, Liu SN, Li WQ, Bie HL, Xu Q, Wang LR. 2024. Haplotype-resolved

genome of a heterozygous wild peach reveals the *PdaWRKY4-PdaCYP716A1* module mediates

resistance to aphids by regulating betulin biosynthesis. *Journal of Integrative Plant Biology* **66**:

2716-2735.

Wang J, Li Y, Wang X, Cao K, Zhu G, Fang W, Chen C, Wu J, Guo J, Xu Q, Wang L. 2022a.

Betulin, synthesized by *PpCYP716A1*, is a key endogenous defensive metabolite of peach against aphids. *Journal of Agricultural and Food Chemistry* **70**: 12865-12877.

Wang Q, Wang H, Zhang Y, Chen J, Upadhyay A, Bhowmick B, Hang J, Wu S, Liao C, Han Q. 2022b. Functional analysis reveals ionotropic GABA receptor subunit RDL is a target site of ivermectin and fluralaner in the yellow fever mosquito, *Aedes aegypti*. *Pest Management Science* **78**: 4173-4182.

Wang Y, Gao R, Gu T, Li X, Wang M, Wang A, Qiu Y. 2025b. Metabolomics and transcriptomics reveal the role of the terpene biosynthetic pathway in the mechanism of insect resistance in *Solanum habrochaites*. *Journal of Agricultural and Food Chemistry* **73**: 6253-6269.

Xu Z, Hu Y, Hu J, Qi C, Zhang M, Xu Q, He L. 2020. The interaction between abamectin and RDL in the carmine spider mite: a target site and resistant mechanism study. *Pesticide Biochemistry and Physiology* **164**: 191-195.

Yamagata N, Ezaki T, Takahashi T, Wu H, Tanimoto H. 2021. Presynaptic inhibition of dopamine neurons controls optimistic bias. *ELife* **10**: e64907.

Yan X, Yang C, Yang M, Ma Y, Zhang Y, Zhang Y, Liu C, Xu Q, Tu K, Zhang M. 2022. All-in-one theranostic nano-platform based on polymer nanoparticles for BRET/FRET-initiated bioluminescence imaging and synergistically anti-inflammatory therapy for ulcerative colitis. *Journal of Nanobiotechnology* **20**: 99.

Yang C, Halitschke R, O'Connor SE. 2023. OXIDOSQUALENE CYCLASE 1 and 2 influence triterpene biosynthesis and defense in *Nicotiana attenuata*. *Plant Physiology* **194**: 2580-2599.

Zhang X, Huang Y, Chen WJ, Wu S, Lei Q, Zhou Z, Zhang W, Mishra S, Bhatt P, Chen S. 2023. Environmental occurrence, toxicity concerns, and biodegradation of neonicotinoid insecticides.

Environmental Research **218**: 114953.

Zhou H, Guo F, Luo J, Zhang Y, Liu J, Zhang Y, Zheng X, Wan F, Ding W. 2021. Functional

analysis of an upregulated calmodulin gene related to the acaricidal activity of curcumin

against *Tetranychus cinnabarinus* (Boisduval). *Pest Management Science* **77**: 719-730.

Zhou H, Jian Y, Shao Q, Guo F, Zhang M, Wan F, Yang L, Liu Y, Yang L, Li Y, Yang P, Li Z, Li

S, Ding W. 2023. Development of sustainable insecticide candidates for protecting pollinators:

insight into the bioactivities, selective mechanism of action and QSAR of natural coumarin

derivatives against aphids. *Journal of Agricultural and Food Chemistry* **71**: 18359-18374.

Zhou H, Ning Y, Jian Y, Zhang M, Klakong M, Guo F, Shao Q, Li Y, Yang P, Li Z, Yang L, Li

S, Ding W. 2024a. Functional analysis of a down-regulated transcription factor-SoxNeuroA

gene involved in the acaricidal mechanism of scopoletin against spider mites. *Pest Management*

Science **80**: 1593-1606.

Zhou H, Wan F, Lai X, Yan F, Zhang M, Ni Y, Guo Y, Zhang P, Guo F, Klakong M, Peng G,

Guo W, Zeng X, Zhang Z, Pan X, Liu Y, Yang L, Li S, Ding W. 2024b. Synergistic action and

mechanism of scoparone, a key bioactive component of *Artemisia capillaris,* and spirodiclofen

against spider mites. *Pest Management Science* **80**: 5035-5049.

Figure 1. The aphicidal activity of plant-derived derivative betulin against *M. persicae*. (A-G)

Representative images of the control effects of betulin (A, 0.1641 mg·mL^{-1}, 48 h LC$_{50}$ value of betulin),

pymetrozine (B, 1.0612 mg·mL^{-1}, a positive control), and CK (C, a negative control) against *M. persicae* in

greenhouse (A-E) and field (F, G) tests after treatment for 14 d. The data are shown as the mean ± SD from

twelve independent experiments. ** $P < 0.01$. ns, not significant.

Figure 2. RNA-Seq analysis revealed candidate targets for betulin against *M. persicae.* (A) Diagram of the

candidate targets for betulin against aphids as determined by RNA-Seq. Water containing 0.1% (v/v)

Tween-80 and 3% (v/v) acetone was used as the control treatment (CK). (B) Principal coordinates analysis

(PCA) analysis differentially expressed genes in RNA-Seq. (C-D) Distribution (C) and heatmap (D) of the

significantly differentially expressed genes (DEGs) in *M. persicae.* (E) qPCR validation of 15 DEGs

identified by RNA-Seq. (F-G) Top 10 enriched GO (F) and KEGG (G) pathways of the DEGs. The "rich

ratio" was defined as the ratio of the number of DEGs enriched in the pathway to the total number of

genes enriched in the same pathway. (H) Heatmap of the expression level of the GABR genes identified

by RNA-Seq. *MpGABR*, encoding GABA$_A$ receptor; *MpGABRAP*, encoding GABA$_A$ receptor associated

protein; *MpGABRB*, encoding GABA$_A$ receptor β subunit.

Figure 3. *MpGABR* expression in aphids was significantly inhibited by treatment with betulin. (A)

Schematic drawing of MpGABR, MpGABRAP and MpGABRB. TM, transmembrane helices. (B-D) qPCR

expression analysis of *MpGABR* (B), *MpGABRAP* (C) and *MpGABRB* (D) transcripts in *M. persicae* exposed

to LC_{30}, LC_{50}, and LC_{70} of betulin for 48 h. The bars represent the average (± SD). Different letters above

the error bars indicate significant difference (ANOVA, Tukey's test, $P < 0.05$). (E, F) Western blotting

analysis of MpGABR protein after betulin treatment for 48 h at three different concentrations (LC_{30}, LC_{50}

and LC_{70}). The bars represent the average (± SD). Different letters above the error bars indicate significant

difference (ANOVA, Tukey's test, $P < 0.05$).

Figure 4. Silencing the expression of *MpGABR*, *MpGABRAP* and *MpGABRB* via RNAi. (A-C) qPCR

expression analysis of *MpGABR* (A), *MpGABRAP* (B) and *MpGABRB* (C) after RNAi at 48 h post-dsRNA

feeding relative to the expression levels after DEPC-water treatment. (D) Schematic drawing of the RNAi

assay in *M. persicae*. (E) Mortality of aphids exposed to the LC$_{50}$ of betulin for 48 h after RNAi. An asterisk

(*) on the error bar indicates a significant difference between the treatment and group CK according to *t*

tests, ***$P < 0.001$, ****$P < 0.0001$. ns, not significant.

Figure 5. Phylogenetic analysis of GABR in insects. Neighbor-joining method with 1000 replicates based phylogenetic tree of GABR from six orders: Hemiptera, Diptera, Lepidoptera, Thysanoptera, Hymenoptera, and Coleoptera. Protein sequence alignment was performed using ClustalW in MEGA 7. The sequences used for constructing the tree were listed in Table S6.

Figure 6. The interaction of betulin with MpGABR protein. (A) Expression of recombinant wild type-aphid MpGABR (1, WT) protein by *Escherichia coli*. (B) Quantification of the binding affinity of betulin with wild type-aphid MpGABR using MST. Current responses (C) and inhibition activity (D) of MpGABR induced by different concentration of betulin in the presence of GABA. 0 µM MpGABR indicates the presence of only GABA.

Figure 7. Molecular docking, binding site and inhibitory effect of betulin on MpGABR. (A) Best conformations of betulin docked to the binding pocket of MpGABR in aphids. An enlarged view of the betulin binding sites in MpGABR is indicated by a dashed frame. Potential Pi–alkyl (green), Alkyl (yellow), and hydrogen bond (red) interactions are indicated by dashed lines. (B) Sequence alignment of the key amino acids bound to betulin. The conserved residues among the different species in GABR are shown in orange, cyan, and light green. The numbers next to the amino acid indicates the site of the last residue of the key amino acids. (C) Expression of the recombinant mutation type-aphid MpGABR by *Escherichia coli*. Lane 1: R224A, Lane 2: A226T, Lane 3: F227Y, Lane 4: T228R. (D, E) Quantification of the binding affinity of betulin with wild-type and mutant type-aphid MpGABR using MST. The bars represent the average (± SD) values. * indicates a significant difference (Student's *t* test, ****P < 0.0001).

Figure 8. Gene editing in *Drosophila* validated the species-specific binding site of betulin. (A) Expression

of the recombinant wild type (WT) and mutation type (R122T) of DmGABR (*Drosophila melanogaster*) by

E. coli. Lane 1: wild type, Lane 2: R122T. (B, C) Quantification of the binding affinity of betulin (B) and

pymetrozine (C) with WT and R122T of DmGABR using MST. (D) Sanger sequencing of the *DmGABR*

gene in flies. Direct sequencing chromatograms of PCR products amplified from a fragment of gDNA

flanking the WT (a), and introduced R122T (b) mutant flies. (E-F) Toxicity curves of betulin (E), and

pymetrozine (F) against WT, and the DmGABR[R122T] of the Cas9 *Drosophila*.

Figure 9. Proposed model for the mechanism of action against *Myzus persicae* by targeting GABA$_A$

receptors (GABR). After exposure to betulin, the expression of *MpGABR* was inhibited the level of

MpGABR protein is decreased, resulting in a decrease in the channel of chloride ion influx. Besides,

betulin directly and specifically binds to the amino acid residue THR228 of MpGABR, thereby disabling

it.