Developmental constraints mediate the summer solstice reversal of climate effects on the autumn phenology of European beech

Dominic Rebindaine[1,2]*, Thomas W. Crowther[1], Susanne S. Renner[3], Zhaofei Wu[4,2], Yibiao Zou[1,2], Lidong Mo[5], Haozhi Ma[2], Raymo Bucher[1], Constantin M. Zohner[1]

[1]Department of Environmental Systems Science, ETH Zurich (Swiss Federal Institute of Technology), Universitätsstrasse 16, 8092 Zurich, Switzerland; [2]WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland; [3]Department of Biology, Washington University, Saint Louis, United States of America; [4]College of Water Sciences, Beijing Normal University, Beijing, China; [5]Department of Plant Biology and Ecology, College of Life Science, Nankai University, 300071 Tianjin, PR China.

*Author for correspondence: Dominic Rebindaine, dominic.rebindaine@usys.ethz.ch

Abstract

Accurate projections of temperate tree growing seasons under climate change require representing developmental constraints that determine tree resource allocation. Recent work has identified the summer solstice (21 June) as a phenological "switch point" with pre-solstice warming advancing autumn phenology and post-solstice warming delaying it. However, the flexibility of this switch point remains poorly understood. We performed trans-solstice climate manipulation experiments on potted European beech (*Fagus sylvatica*) saplings to test (i) how spring leaf-out timing and June-August temperatures influence end-of-season timing (bud set and leaf senescence [50% loss of leaf chlorophyll content]), and (ii) whether daytime and nighttime temperatures before and after the solstice have different effects, given that trees primarily grow at night. Bud set and leaf senescence responses were tightly coupled (R^2 = 0.49), with bud responses being generally stronger. Each day delay in spring leaf-out delayed bud set by 0.24 ± 0.06 days and senescence by 0.22 ± 0.08 days. Post-solstice full-day cooling in July delayed autumn phenology in late-leafing individuals (bud set +4.9 ± 2.6 days; senescence +3.1 ± 2.8 days) but had negligible impact on early-leafing trees (bud set +1.4 ± 2.6 days; leaf senescence +2.2 ± 2.8 days). Conversely, August full-day cooling advanced both

stages. Daytime cooling before the solstice had no effect, while after the solstice it advanced

autumn phenology. Nighttime cooling always delayed bud set. These findings support the

Solstice-as-Phenology-Switch model and highlight the central role of developmental

progression in constraining growing seasons. Faster early-season development –especially

under nighttime warming– moves trees past the switch earlier, increasing sensitivity to late-

season cooling and thereby triggering earlier autumn phenology. To improve growing season

length projections, phenology models should account for these developmentally-mediated

and diel-specific temperature responses.

Keywords: Carbon cycle | Climate change | Global warming | Plant-climate interactions | Plant

development| Plant growth | Phenology

Introduction

Climate shifts are leading to rapid, species-specific changes in phenology and ecosystem

productivity (Boisvenue & Running, 2006; Menzel *et al.*, 2006; Yang & Rudolf, 2010; Thackeray

et al., 2016). In temperate forests, changes in the timing of spring leaf-out, autumn leaf-

senescence and bud set are modifying water, energy and carbon cycles (Peñuelas *et al.*, 2009;

Richardson *et al.*, 2013), with extended growing seasons increasing net ecosystem carbon

uptake by up to 9.8 gC m^{-2} day^{-1} (Keenan *et al.*, 2014). Therefore, understanding the

interaction between climate change and temperate forest phenology is pivotal to improving

forecasts of community dynamics and carbon sequestration.

The past few decades have seen delays in the onset of temperate autumn phenology, but

these changes are much smaller in magnitude compared to the advances in spring leaf-out

observed during the same period (Gill *et al.*, 2015; Piao *et al.*, 2019). This is unexpected, given

that experiments have demonstrated a high sensitivity of leaf senescence to autumn warming,

with phenological responses even surpassing the temperature sensitivity (days per °C) of

spring leaf-out (Fu *et al.*, 2018). One possible explanation for this discrepancy is that other

factors may counterbalance the effects of autumn warming, with some studies finding that

earlier leaf-out leads to earlier leaf senescence (Fu *et al.*, 2014; Keenan & Richardson, 2015;

Zani *et al.*, 2020). This connection between spring and autumn phenophases could be due to

developmental and nutrient constraints that affect carbon source-sink dynamics in temperate

trees (Paul & Foyer, 2001; Zani *et al.*, 2020; Zohner *et al.*, 2023; Gessler & Zweifel, 2024).

The Solstice-as-Phenology-Switch hypothesis, supported by experiments and monitoring data,

posits that warmer air temperatures before the summer solstice accelerate tree growth and

development, leading to an earlier onset of leaf senescence, whereas warmer temperatures

after the solstice slow the progression of leaf senescence (Zohner *et al.,* 2023). The opposite

is equally true in colder years, with pre-solstice cooling delaying autumn phenology and post-

solstice cooling advancing it. This reversal in temperature effects provides an explanation for

the less pronounced shifts in autumn phenology compared to spring (Piao *et al.*, 2019). The

effect reversal after the solstice points to the role of photoperiod in regulating plant

physiology (Bauerle *et al.*, 2012; Petterle *et al.*, 2013; Singh *et al.*, 2017), with trees becoming

increasingly responsive to cooling temperatures as days continue to shorten. The precise

timing of this reversal appears to be flexible and has been advancing in recent decades,

potentially due to faster development driven by climate warming (Zohner *et al.*, 2023). For

temperate trees, the key outcomes of development are production of viable seeds, cessation

of primary and secondary growth, and maturation of perennating tissues including leaf or

flower buds (completion of bud set) before the onset of frost (Rohde & Bhalerao, 2007; Tanino

et al., 2010; Cooke *et al.*, 2012). Because opportunities for development will vary among

years, the timing of the temperature effect reversal should be flexible. However, this flexibility

has yet to be demonstrated in processes directly related to development such as autumn bud

set.

In this study, we investigate how air temperature changes around the summer solstice affect

end-of-season (EOS) timing in European beech (*Fagus sylvatica* L.), using bud set and leaf

senescence as key physiological markers of autumn phenology (Singh *et al.*, 2017; Zohner &

Renner, 2019; Mariën *et al.*, 2021). We ask three main questions: i) How does cooling before

vs. after the solstice affect EOS timing? ii) How do diel (daytime vs. nighttime) temperature

changes differ in their effects? iii) Does the timing of leaf-out modulate the reversal of

temperature effects on autumn phenology? Diel temperature variations are relevant because

trees typically grow more at night when temperatures and water deficit are lower (Steppe *et*

al., 2015; Mencuccini *et al.*, 2017; Zweifel *et al.*, 2021). Thus, developmental effects mediated

by temperature may be especially pronounced during nights. Moreover, long-term trends in

climate show that nighttime temperatures rose faster than daytime temperatures from 1950

to 2004 (Vose et al., 2005), though this trend may have shifted since 1991 (Zhong et al., 2023).

To address the third question, we experimentally manipulated spring conditions to create

early-leafing and late-leafing individuals within the same growing season, allowing us to assess

whether slowed early-season development postpones the reversal date of temperature

effects on autumn phenology (Fig. 1). As depicted in our conceptual model (Fig. 1), trees are

dynamic systems that continuously integrate and adapt their physiological responses to

environmental conditions over time. We therefore expected their temperature responses to

differ both across months (June-August) and between day and night.

Materials and Methods

Selected species

Fagus sylvatica L. has a large temperate European distribution, is highly economically and

ecologically important, and is likely threatened by climate change (Gessler *et al.*, 2006).

Therefore, improving our understanding of how *F. sylvatica*'s growing season is modulated is

vital to the management of European forests. Moreover, *F. sylvatica* has frequently been used

as a key species to study temperate tree phenology (Dittmar & Elling, 2006; Vitasse *et al.*,

2011; Zohner & Renner, 2019; Zani *et al.*, 2020; Zohner *et al.*, 2023; Mariën *et al.*, 2025; Švik

et al., 2025).

Experiment 1

To test the antagonistic effects of early-season development and late-season air temperature

on autumn phenology, we set up an experimental population (n = 267) of 40-60 cm tall

European beech (*F. sylvatica*) trees in Zurich, Switzerland in 2023. The trees were sourced from

a local nursery, and each tree was placed individually in a 20 L plastic pot containing a 1:1:1

sand / peat / organic soil mixture with a Nitrogen (N) concentration of ~65 g m^{-3}, a Phosphate

(P_2O_5) concentration of ~140 g m^{-3}, and a Potassium (K_2O) concentration of ~400 g m^{-3}.

Individual trees were assigned randomly to one of 10 treatment groups (26 ≤ n ≤ 27).

Treatment groups were cooled at different times of the year, using different cooling levels (Fig.

2, Table 1). To arrest spring development and thereby generate early-leafing and late-leafing

individuals, half of the experimental population was placed outside under ambient conditions,

while the other half was cooled in climate chambers from 4 April to 24 May. The chambers

were set to a low of 2°C at night and a high of 7°C during the day, following a simulated day-

night cycle of temperature and light availability (13 h photoperiod at ~4,300 lux). These cold

conditions were not intended to mimic natural European conditions, but rather to slow spring

development as much as possible without damaging the trees. Temperatures around 5°C are

associated with low rates of growth and development processes such as cell division,

expansion and maturation (Körner, 2003; Tumajer *et al.*, 2021). From 24 May to 21 June, all

trees were kept outside under ambient conditions in a randomised block design. All trees were

monitored to observe their individual leaf-out dates, which was defined as the date when >

50% of their leaves had unfolded, corresponding to BBCH15 (Capdevielle-Vargas *et al.*, 2015).

During the summer treatments, control trees were placed outside under natural ambient

conditions. We did not use climate chambers for the control groups because warm conditions

in these chambers can introduce unique physiological stressors—such as aphid infestations—

not present in cold chambers (Bezemer *et al.*, 1998). In experiment 2, however, we included

an additional chamber control to test for potential chamber effects. Although we observed

higher aphid abundances in these warm control chambers, chamber exposure itself had no

detectable effect on autumn phenology. This strengthens the inference that the phenological

shifts observed in our treatments reflect the intended temperature manipulations.

Nevertheless, for experiment 1, where chamber controls were not included, we cannot

entirely rule out the possibility that chamber conditions exerted some unintended influence

on treated trees compared with ambient controls.

The July treatments took place directly after the summer solstice from 22 June to 23 July.

Treated trees were placed in climate chambers set to a low of 2°C at night and a high of 7°C

during the day or a low of 8°C at night and a high of 13°C during the day depending on their

cooling level (extreme and moderate, respectively). Trees in the chambers experienced a

photoperiod of 16 h at ~7,300 lux. The extreme cooling was designed to severely impair cell

division, expansion, and maturation as described above. Additionally, under this temperature

regime, photosynthesis should be reduced by > 40% (Körner, 2006). The moderate cooling

regime, which was still much colder than the ambient conditions, should have also impaired

growth and development as well as reduced photosynthesis by at least 30% (Körner, 2003,

2006; Lenz *et al.*, 2014). These settings aimed to generate large temperature differences

between treatments and maximise our ability to detect the mechanism underlying the solstice

switch, rather than to represent naturally occurring conditions. The August treatments took

place from 24 July to 25 August under the same conditions as the July treatments except with

a 15 h photoperiod. For the remainder of the experiment, all trees were kept outside in a

randomised block design. Throughout the experiment, all trees were watered frequently to

ensure constant water supply (Fig. S1).

To observe the effects of our treatments on the development of overwintering buds, we

monitored bud growth to derive bud set dates, our marker for the cessation of primary

aboveground development and the beginning of autumn phenology, for all trees. On each

tree, the terminal bud of the primary shoot and the terminal bud on a random lateral stem

were selected and tagged for measurement. Each selected bud was measured to 0.01 mm

precision using a digital calliper (CD-P8"M, Mitutoyo Corp, Japan). We measured all buds on

a regular basis from 4 July to 2 November. Bud set was defined as the date when each bud

reached 90% of its own maximum length, which is considered to be an indicative stage of

aboveground primary growth cessation (Signarbieux *et al.*, 2017; Zohner & Renner, 2019).

Bud lengths were linearly interpolated between measurement dates to derive the date at

which they reached the 90% threshold.

To obtain a more holistic perspective of autumn phenology, we also derived leaf senescence

dates from leaf spectral index measurements taken using a SPAD-502 Plus (Soil Plant Analysis

Development, Minolta Camera Co., Ltd, Tokyo, Japan). We measured nine leaves per individual

(three each from the top, middle and bottom of the crown) on a monthly basis in summer,

every other week in September, and on a weekly basis from October to mid-December. We

removed measurements taken on 7 December from the analysis as they were unreasonably

high, likely due to false readings caused by extensive leaf browning. SPAD readings were then

converted to total chlorophyll content (*Chl* in µg/g fresh weight) using an empirically

established equation for *Fagus sylvatica* leaves (Percival *et al.*, 2008):

$$Chl = -0.0029 \times SPAD^2 + 1.175 \times SPAD + 3.8506.$$

The chlorophyll content between two consecutive measurement dates was estimated using

linear interpolation. For each treatment and measurement date, we removed individual

chlorophyll measurements that were more than 1.5 × the interquartile range below the lower

quartile or above the upper quartile. As an additional cleaning step, we completely removed

any trees that had more than one data point removed in the last step. Finally, we calculated

individual leaf senescence dates as the day-of-year when chlorophyll content last fell below

50% of the observed peak chlorophyll content.

Experiment 2

To observe the effects of pre- and post-solstice daytime and nighttime temperature on autumn

phenology, we set up an experimental population (n = 180) of four-year-old *Fagus sylvatica*

trees in Zurich, Switzerland in 2022. The trees were sourced from a local nursery and each

tree was placed individually in a 20 L plastic pot containing a 1:1:1 sand / peat / organic soil

mixture with a Nitrogen (N) concentration of ~65 g m^{-3}, a Phosphate (P_2O_5) concentration of

~140 g m^{-3}, and a Potassium (K_2O) concentration of ~400 g m^{-3}.

The ambient control treatment consisted of 36 trees exposed to natural ambient conditions.

The remaining eight treatments were each applied to 18 trees and included cooling in climate

chambers with simulated ambient day length (16 h) and light intensity (~6,900 lux). The pre-

solstice cooling treatments were applied between 22 May and 21 June. The post-solstice

cooling treatments were applied between 22 June and 21 July. The pre- and post-solstice

treatments included four levels each: Chamber control, where the trees were continuously

subjected to 20°C; Day cooling, where trees were subjected to 8°C in the daytime and 20°C at

night; Night cooling, where trees were subjected to 20°C in the day and 8°C at night; Full-day

cooling, where trees were continuously subjected to 8°C (Fig. 3, Table 2). Due to warm

temperatures, chamber control trees were subject to an increase in aphid abundance,

however this did not alter their bud set timing (see data analyses). After treatment, all trees

were placed in a randomised block design outside under ambient conditions. Soil moisture

content was regulated by frequent watering. We measured all buds on a weekly basis from 25

August to 3 November following the same methodology as in experiment 1. We also measured

leaf-level CO_2 assimilation rates during the pre-solstice treatment window (see Zohner *et al.,*

2023 for methodology), and derived leaf senescence dates following experiment 1.

Data analyses

For experiment 2, we performed one-way ANOVAs that showed no significant differences between the end-of-season dates of the ambient, pre-chamber and post-chamber control groups (for bud set: $F_{2,136}$ = 0.346, p = 0.708, see Fig. S7; for leaf senescence: $F_{2,63}$ = 0.931, p = 0.399). Therefore, for the following analyses we treated all three control treatments as one control treatment with 72 trees. For both experiments we ran linear models using treatment and bud type (apical vs. lateral) as predictors and bud set day-of-year, absolute bud growth or relative bud growth as the response variable. When modelling leaf senescence day-of-year, we used treatment as the sole predictor. For experiment 1, effect sizes were calculated in comparison to the corresponding control treatment, i.e., all early-leafing treatments were compared to the early-leafing control treatment, and all late-leafing treatments were compared to the late-leafing control. Absolute and relative bud growth were calculated as:

$$Absolute = max\ length - min\ length$$

$$Relative = \left(1 - \frac{min\ length}{max\ length}\right) * 100$$

Where:

min length = the first measured bud length,

max length = the measured bud length when the bud first surpassed 90% of its final length.

To determine the sensitivity of autumn bud set timing to spring leaf-out timing, we ran a linear mixed-effects model using leaf-out day-of-year and bud type as the fixed predictor variables, summer temperature treatment as random effect, and bud set day-of-year as the response variable. We ran the same analysis for leaf senescence timing without the bud type effect.

To quantify the relative contributions of early-season development and late-season

temperature to variation in bud set timing in experiment 1, we ran a variance partitioning

analysis. We fit a linear model with leaf-out day-of-year, summer cooling treatment

(combination of cooling timing and level, e.g. July_Moderate), and bud type as explanatory

variables. Variance partitioning was conducted using the *varpart* function in the *vegan* R

package (Oksanen *et al.*, 2025), which decomposes explained variance into unique and shared

components. All statistical analyses were performed in R version 4.5.1 (R Core Team, 2025).

Results

Experiment 1

Linear modelling showed that across all treatments, late-leafing trees cooled during spring

delayed their bud set by 4.6 ± 1.3 days (mean ± 2 SE, $p < 0.01$). Across all treatment pairings,

the mean bud set date for the late-leafing group always occurred later than for the early-

leafing group (Fig. S2). A linear mixed effects model including leaf-out day-of-year and bud-

type as fixed effects and treatment as a random effect revealed that, on average, each day

delay in spring leaf-out was associated with a delay of 0.24 ± 0.06 days in bud set timing.

Across all treatments, bud type had a small effect, lateral buds set on average 1.1 ± 1.2-days

earlier than apical buds ($p = 0.06$). When modelling only the early-leafing groups, bud type

had no discernible effect (0.04 ± 1.64 days earlier for lateral buds, $p = 0.96$). However, when

modelling only the late-leafing groups, lateral buds set earlier than apical buds (2.17 ± 1.64

days advancement, $p = 0.01$).

The effect of July cooling differed strongly between the early- and late-leafing trees: Within

the early-leafing group, moderate cooling in July lead to a small, non-significant delay in bud

set compared to the ambient July treatment (1.4 ± 2.6 days, p = 0.29). July cooling had a much

greater impact on late-leafing trees, leading to a 4.8 ± 2.6-day delay in bud set compared to

the ambient July treatment in the late-leafing group (p < 0.01).

In contrast to July cooling, the effects of August cooling did not differ between early- and late-

leafing trees, advancing bud set in all treatment groups (Fig. 4b). Cooling in August lead to a

4.5 ± 2.6-day (p < 0.01) and a 4.4 ± 2.6-day (p < 0.01) advancement in bud set for early- and

late-leafing trees, respectively. The extreme cooling treatments showed similar patterns,

although slightly less clearly (Fig. S3). Variance partitioning revealed that the early-season

development effect (leaf-out day-of-year) explained 8.8% of the total variance in bud set

timing, the late-season temperature effect (summer cooling treatment) explained 14.6%, and

bud type explained 0.3%. Shared variance values were ≈ 0%, indicating no meaningful overlap.

Effects of absolute and relative growth on bud set date can be found in the supporting

information (Figs. S4-S5). Bud set timing had no effect on final bud length (Fig. S6).

Experiment 2

Linear modelling showed that pre-solstice full-day (day and night) cooling delayed autumn

bud set by 4.1 ± 3.6 days (mean ± 2SE, p = 0.02) (Fig. 5). Pre-solstice nighttime cooling had a

similar effect, delaying bud set by 4.2 ± 3.6 days (p = 0.02). By contrast, pre-solstice daytime

cooling had no significant effect on bud set (0.1 ± 3.7 days, p = 0.94). Post-solstice full-day

cooling advanced autumn bud set by 5.2 ± 3.6 days (p < 0.01). Similarly, post-solstice daytime

cooling advanced bud set by 5.3 ± 4.3 days (p = 0.01). Conversely, post-solstice nighttime

cooling delayed bud set by 3.8 ± 3.5 days (p = 0.03). Across all treatments, lateral buds set

considerably earlier than apical buds (9.4 ± 2.08 days, p < 0.01). Effects of relative growth on

bud set date can be found in the supporting information (Figs. S8).

Comparison of phenological metrics

Analyses using leaf senescence (50% drop in chlorophyll content) as the phenological marker

produced results that were overall consistent with the bud set analyses. For example, in

experiment 1, senescence was delayed by 4.22 ± 1.61 days in late-leafing trees and each day

delay in spring leaf-out led to a 0.22 ± 0.08-day delay in leaf senescence, closely matching with

bud set. Across both experiments the observed patterns were largely the same, although the

strengths of each effect differed slightly, and overall, the effects on leaf senescence tended to

be weaker (Figs. S9-10). Out of all treatments, opposing responses from bud set and leaf

senescence timing were only observed in the early-leafing extreme August cooling group

(Treatment 9, bud effect = 1.9 ± 2.59 days advancement, p = 0.14, leaf effect = 1.77 ± 3.53 days

delay, p = 0.32) from experiment 1, and the post-full-day cooling group from experiment 2

(bud effect = 5.2 ± 1.81 days advancement, p < 0.01, leaf effect = 1.66 ± 3.34 days delay, p =

0.32). Out of 66 pairwise comparisons of estimated marginal means (every treatment

compared against every other treatment in the same experiment), 55 (83%) showed

directional agreement between bud set and leaf senescence effects (Fig. S11, adjusted R^2 =

0.49).

Discussion

In this study, we conducted experiments on European beech to test how bud set—a marker

of primary growth cessation—and leaf senescence respond to monthly and diel temperature

changes around the summer solstice. We found large differences in the responses between

trees subjected to daytime versus nighttime cooling before and after the solstice (Fig. 5) and

between early- and late-leafing trees (Fig. 4). These differences suggest that early-season

developmental timing not only influences the timing of autumn phenology *per se* but also

plays a critical role in determining the timing of the summer solstice effect reversal. Therefore,

our findings support our conceptual model (Fig. 1), highlighting development as a key factor

underpinning the Solstice-as-Phenology-Switch hypothesis. In the following sections, we

discuss the effects of pre-solstice and post-solstice air temperature on development, along

with the distinct impacts of daytime versus nighttime temperature variations.

Early-season development rates alter responses to late-season temperature

Our trans-solstice climate manipulation experiments showed that trees delayed the end of

their growing season in response to slow/late early-season development (delayed spring leaf-

out). Regardless of the summer cooling treatments, late-leafing trees consistently set buds

and senesced leaves after their early-leafing counterparts (Fig. S2), with each day delay in

spring leaf-out delaying bud set by an average of 0.24 days and senescence by 0.22 days. This

is in line with previous studies demonstrating a tight linkage between within-year variations

in spring and autumn phenology (Fu *et al.*, 2014; Keenan & Richardson, 2015; Signarbieux *et*

al., 2017), likely governed by developmental constraints (Zohner *et al.*, 2023), buildup of

water and nutrient stress (Paul & Foyer, 2001; Buermann *et al.*, 2018; Bigler & Vitasse, 2021),

or leaf aging (Lim *et al.*, 2007).

Despite the importance of early-season development, summer cooling treatment uniquely

explained more variance in bud set timing than leaf-out day-of-year (Adjusted R^2 = 0.15 and

0.09, respectively). July cooling induced a delay in bud set dates 3.4 times greater in late-

leafing trees compared to early-leafing ones (4.8 versus 1.4 days delay), which agrees with the

expectations derived from our conceptual model (Fig. 1). As we had expected, the natural

interannual variation in early-season development causes year-to-year differences in the point

at which cool temperatures switch from delaying autumn phenology to advancing it. This

offers a physiological explanation for the advancement in the effect reversal timing observed

between 1966 and 2015 in *Fagus sylvatica, Aesculus hippocastanum, Quercus robur, and*

Betula pendula (Zohner *et al.*, 2023; because control trees did not experience chamber time,

we cannot entirely rule out some influence of this factor on our comparisons).

August cooling induced comparable advances in bud set timing in both early- and late-leafing

trees (4.4-4.5 days). The diminishing difference in cooling responses between the two groups

from July to August suggests that the influence of early-season development weakens as

individuals approach completion of their developmental requirements. By August, phenology

appears primarily governed by the delaying effect of warm air temperature (see direct

warming effect in Fig. 1b).

Two mechanisms could underlie this shift: shortening daylengths and declining photosynthetic

assimilation, the latter of which has been linked to accelerated senescence (Krieger-Liszkay *et*

al., 2019). However, an experiment that reduced August photosynthetic rates in beech by 52-

72% – either through cooling or shading – showed that photosynthetic assimilation alone

cannot account for autumn phenology (Zohner *et al.*, 2023). Despite similar reductions in

photosynthesis, August cooling advanced senescence, while August shading delayed it,

revealing contrasting outcomes. This suggests that photosynthetic assimilation cannot have

been the driver of the observed phenological responses in beech. Instead, the observed

change in temperature responses is more likely regulated by daylength (Kramer, 1936; Petterle

et al., 2013; Singh *et al.*, 2021). As days shorten, trees become increasingly responsive to

cooling (Delpierre *et al.*, 2009; Körner *et al.*, 2016). Together with the waning influence of

developmental carry-over, this can explain why by August trees responded similarly to cooling

regardless of their earlier developmental trajectories (Fig. 1). These findings highlight the

critical interaction between photoperiod and temperature in shaping the autumn phenology

of *Fagus sylvatica.*

Effects of daytime vs. nighttime temperature

Our experiments showed that daytime and nighttime temperatures had different effects on

autumn phenology before and after the summer solstice: Before the solstice, daytime cooling

had no impact on the timing of bud set, while nighttime cooling and full-day cooling delayed

bud set by 4.2 and 4.1 days, respectively (Fig. 5). This pattern was similar when using leaf

senescence as the end-of-season marker. Because experimental cooling reduced

photosynthesis by 69-83% across all treatments (Fig. S12), these effects are unlikely to be

explained by changes in carbon assimilation. Instead, they highlight the importance of

developmental processes—such as cell division and expansion—which mostly occur at night

(Steppe *et al.*, 2015; Mencuccini *et al.*, 2017; Zweifel et al., 2021; Fig. 6a). The developmental

arrest induced by night cooling—and therefore also by full-day cooling—likely slowed growth

and maturation, effectively extending the growing season.

After the solstice, daytime and nighttime cooling of 8°C elicited opposite responses in both

phenological markers, with weaker effects on senescence. Trees subjected to post-solstice

daytime cooling and full-day cooling set their buds earliest, on average more than five days

earlier than controls. As days shorten, the temperature threshold inducing this growth

cessation must be lower at night, otherwise trees would risk prematurely ending their growing

season due to the daily minimum temperatures occurring at night (Fig. 6b). This may explain

why post-solstice nighttime cooling delayed, rather than advanced, autumn phenology:

Cooling of 8°C was sufficient to trigger overwintering responses—growth cessation and

maturation of perennating tissues—when applied during the day but insufficient when

applied at night (see Fig. S13 for a conceptual schematic).

An alternative explanation is that daytime cooling indirectly triggered dormancy induction by

suppressing photosynthesis (Fig 6a), under the assumption that trees prioritise dormancy

once photosynthesis is no longer possible (Körner *et al.*, 2016). Though, as discussed above,

late-season photosynthesis appears to assert little control over the autumn phenology of *F.*

sylvatica. Taken together, these results support a Diel Cooling Hypothesis, in which the impact

of cooling depends on whether it occurs during the day or at night, because the temperature

thresholds that trigger growth cessation differ between daytime and nighttime conditions.

Nighttime cooling always led to a delay in bud set of approximately four days (Fig. 5). As trees

mainly grow at night (Steppe *et al.*, 2015; Mencuccini *et al.*, 2017; Zweifel *et al.*, 2021), colder

nighttime temperatures likely slowed down key developmental processes such as

meristematic activity, tissue expansion and maturation, which in turn delayed primary growth

cessation (Fig. 6). Similar responses may occur in other temperate tree species: Cold autumn

nights have been shown to delay growth cessation and slow bud development in *Populus,*

Pinus and *Picea* species (Kramer, 1956; Malcolm & Pymar, 1975; Kalcsits *et al.*, 2009).

Moreover, the summer solstice effect reversal has been observed consistently across Northern

Hemisphere temperate forests (Zohner *et al.*, 2023). Further experiments should explicitly

test these diel responses in other temperate tree species.

Support for the Solstice-as-Phenology Switch Hypothesis

The Solstice-as-Phenology-Switch hypothesis posits that changes in temperature before and

after the summer solstice have opposite effects on the autumn phenology of northern forest

trees (Zohner *et al.*, 2023). The hypothesis does not assume that the reversal is fixed precisely

on June 21, but rather that it occurs around the solstice, when photoperiod cues shift trees

from accelerating to decelerating their seasonal development. Our conceptual model (Fig. 1)

explicitly incorporates this flexibility by showing how the timing of the reversal depends on

developmental speed: individuals that develop more slowly (or leaf out later) cross the

compensatory point later in the summer, whereas fast-developing individuals reach it earlier.

Our experiments support this framework: pre-solstice full-day cooling delayed bud set,

whereas post-solstice full-day cooling advanced it, with differences between early- and late-

developing individuals consistent with the model. Moreover, the contrasting impacts of

daytime vs. nighttime cooling demonstrate how diel conditions can further shape when the

reversal is expressed. Thus, our findings reinforce the Solstice-as-Switch hypothesis and

extend it by showing how flexibility arises from interactions between developmental

progression, diel temperature responses, and photoperiod.

In conclusion, earlier primary growth cessation in temperate trees under climate change may

result from higher developmental rates under warmer temperatures. In European beech, we

show that this effect is predominantly driven by nighttime temperatures, reflecting the fact

that growth primarily occurs at night. To better predict future growing season lengths, models

must begin to account for the differential impacts of the seasonal and daily temperature cycles

that temperate zone trees are physiologically adapted to.

Acknowledgments
C.M.Z. was supported by the SNF Ambizione Fellowship programme (no. PZ00P3_193646) and
T.W.C. by DOB Ecology and the Bernina Foundation.
444

445   ## Competing interests
446   The authors declare no competing interests.
447

Author contributions
D.R. conceived and developed the study, conducted the experiments and analyses, and wrote
the manuscript. C.M.Z equally contributed to the conception and development of the study,
contributed to the experiments and analyses, and the text. L.M., H.M., Z.W. and Y.Z.
contributed to the conception and implementation of the first experiment. R.B. contributed
to the conception and implementation of the second experiment. T.W.C. and S.S.R.
contributed to the text. All authors provided comments and approved the final manuscript.
455

456   ## Data availability
457   The code and data for this study are available at Zenodo (Rebindaine *et al.*, 2025).
458

459   ## References
460

Bauerle WL, Oren R, Way DA, Qian SS, Stoy PC, Thornton PE, Bowden JD, Hoffman FM,
Reynolds RF. **2012**. Photoperiodic regulation of the seasonal pattern of photosynthetic
capacity and the implications for carbon cycling. *Proceedings of the National Academy of*
Sciences **109**: 8612–8617.

Bezemer TM, Jones TH, Knight KJ. **1998**. Long-term effects of elevated CO2 and temperature
on populations of the peach potato aphid Myzus persicae and its parasitoid Aphidius
matricariae. *Oecologia* **116**: 128–135.

Bigler C, Vitasse Y. **2021**. Premature leaf discoloration of European deciduous trees is caused
by drought and heat in late spring and cold spells in early fall. *Agricultural and Forest*
Meteorology **307**: 108492.

Boisvenue C, Running SW. **2006**. Impacts of climate change on natural forest productivity -
evidence since the middle of the 20th century: climate change impacts on forest vegetation.
Global Change Biology **12**: 862–882.

Buermann W, Forkel M, O'Sullivan M, Sitch S, Friedlingstein P, Haverd V, Jain AK, Kato E, Kautz M, Lienert S, *et al.* 2018. Widespread seasonal compensation effects of spring warming on northern plant productivity. *Nature* **562**: 110–114.

Capdevielle-Vargas R, Estrella N, Menzel A. 2015. Multiple-year assessment of phenological plasticity within a beech (Fagus sylvatica L.) stand in southern Germany. *Agricultural and Forest Meteorology* **211–212**: 13–22.

Cooke JEK, Eriksson ME, Junttila O. 2012. The dynamic nature of bud dormancy in trees: environmental control and molecular mechanisms. *Plant, Cell & Environment* **35**: 1707–1728.

Delpierre N, Dufrêne E, Soudani K, Ulrich E, Cecchini S, Boé J, François C. 2009. Modelling interannual and spatial variability of leaf senescence for three deciduous tree species in France. *Agricultural and Forest Meteorology* **149**: 938–948.

Dittmar C, Elling W. 2006. Phenological phases of common beech (Fagus sylvatica L.) and their dependence on region and altitude in Southern Germany. *European Journal of Forest Research* **125**: 181–188.

Fu YSH, Campioli M, Vitasse Y, De Boeck HJ, Van den Berge J, AbdElgawad H, Asard H, Piao S, Deckmyn G, Janssens IA. 2014. Variation in leaf flushing date influences autumnal senescence and next year's flushing date in two temperate tree species. *Proceedings of the National Academy of Sciences* **111**: 7355–7360.

Fu YH, Piao S, Delpierre N, Hao F, Hänninen H, Liu Y, Sun W, Janssens IA, Campioli M. 2018. Larger temperature response of autumn leaf senescence than spring leaf-out phenology. *Global Change Biology* **24**: 2159–2168.

Gessler A, Keitel C, Kreuzwieser J, Matyssek R, Seiler W, Rennenberg H. 2006. Potential risks for European beech (Fagus sylvatica L.) in a changing climate. *Trees* **21**: 1–11.

Gessler A, Zweifel R. 2024. Beyond source and sink control – toward an integrated approach to understand the carbon balance in plants. *New Phytologist* **242**: 858–869.

Gill AL, Gallinat AS, Sanders-DeMott R, Rigden AJ, Short Gianotti DJ, Mantooth JA, Templer PH. 2015. Changes in autumn senescence in northern hemisphere deciduous trees: a meta-analysis of autumn phenology studies. *Annals of Botany* **116**: 875–888.

Kalcsits L, Kendall E, Silim S, Tanino K. 2009. Magnetic resonance microimaging indicates water diffusion correlates with dormancy induction in cultured hybrid poplar (Populus spp.) buds. *Tree Physiology* **29**: 1269–1277.

Keenan TF, Gray JM, Friedl MA, Toomey M, Bohrer G, Hollinger DY, Munger JW, O'Keefe J, Schmid HP, Wing IS, *et al.* 2014. Net carbon uptake has increased through warming-induced changes in temperate forest phenology. *Nature Climate Change*.

Keenan TF, Richardson AD. 2015. The timing of autumn senescence is affected by the timing of spring phenology: implications for predictive models. *Global Change Biology* **21**: 2634–2641.

Körner C. **2003**. *Alpine Plant Life*. Berlin, Heidelberg: Springer Berlin Heidelberg.

Körner C. **2006**. Significance of Temperature in Plant Life. In: Morison JIL, Morecroft MD, eds. Plant Growth and Climate Change. Wiley, 48–69.

Körner C, Basler D, Hoch G, Kollas C, Lenz A, Randin CF, Vitasse Y, Zimmermann NE. **2016**. Where, why and how? Explaining the low-temperature range limits of temperate tree species (M Turnbull, Ed.). *Journal of Ecology* **104**: 1076–1088.

Kramer PJ. **1936**. Effect of variation in length of day on growth and dormancy of trees. *Plant Physiology* **11**: 127–137.

Kramer PJ. **1956**. Some effects of various combinations of day and night temperatures and photoperiod on the height growth of loblolly pine seedlings. *Forest Science* **3**: 45–55.

Krieger-Liszkay A, Krupinska K, Shimakawa G. **2019**. The impact of photosynthesis on initiation of leaf senescence. *Physiologia Plantarum* **166**: 148–164.

Lenz A, Vitasse Y, Hoch G, Körner C. **2014**. Growth and carbon relations of temperate deciduous tree species at their upper elevation range limit (M Turnbull, Ed.). *Journal of Ecology* **102**: 1537–1548.

Lim PO, Kim HJ, Gil Nam H. **2007**. Leaf Senescence. *Annual Review of Plant Biology* **58**: 115–136.

Malcolm DC, Pymar CF. **1975**. The influence of temperature on the cessation of height growth of Sitka spruce (Picea sitchensis Bong. Carr.). *Silvae Genet* **24**: 5–6.

Mariën B, Dox I, De Boeck HJ, Willems P, Leys S, Papadimitriou D, Campioli M. **2021**. Does drought advance the onset of autumn leaf senescence in temperate deciduous forest trees? *Biogeosciences* **18**: 3309–3330.

Mariën B, Robinson KM, Jurca M, Michelson IH, Takata N, Kozarewa I, Pin PA, Ingvarsson PK, Moritz T, Ibáñez C, *et al.* **2025**. Nature's Master of Ceremony: The Populus Circadian Clock as Orchestrator of Tree Growth and Phenology. *npj Biological Timing and Sleep* **2**: 1–19.

Mencuccini M, Salmon Y, Mitchell P, Hölttä T, Choat B, Meir P, O'Grady A, Tissue D, Zweifel R, Sevanto S, *et al.* **2017**. An empirical method that separates irreversible stem radial growth from bark water content changes in trees: theory and case studies. *Plant, Cell & Environment* **40**: 290–303.

Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kübler K, Bissolli P, Braslavská O, Briede A, *et al.* **2006**. European phenological response to climate change matches the warming pattern: european phenological response to climate change. *Global Change Biology* **12**: 1969–1976.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos P, Stevens MHH, Szoecs E, *et al.* **2025**. vegan: Community Ecology Package.

Paul MJ, Foyer CH. **2001**. Sink regulation of photosynthesis. *Journal of Experimental Botany*
52: 1383–1400.

Peñuelas J, Rutishauser T, Filella I. **2009**. Phenology Feedbacks on Climate Change. *Science*
324: 887–888.

Percival GC, Keary IP, Noviss K. **2008**. The Potential of a Chlorophyll Content SPAD Meter to
Quantify Nutrient Stress in Foliar Tissue of Sycamore (Acer pseudoplatanus), English Oak
(Quercus robur), and European Beech (Fagus sylvatica). *Arboriculture & Urban Forestry* **34**:
89–100.

Petterle A, Karlberg A, Bhalerao RP. **2013**. Daylength mediated control of seasonal growth
patterns in perennial trees. *Current Opinion in Plant Biology* **16**: 301–306.

Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, Liu L, Lian X, Shen M, Zhu X. **2019**. Plant
phenology and global climate change: Current progresses and challenges. *Global Change*
Biology **25**: 1922–1940.

R Core Team. **2025**. R: A Language and Environment for Statistical Computing.

Rebindaine D, Crowther TW, Susanne S. Renner, Zhaofei Wu, Yibiao Zou, Lidong Mo, Haozhi
Ma, Raymo Bucher, Constantin M. Zohner. **2025**. Developmental constraints mediate the
summer solstice reversal of climate effects on the autumn phenology of European beech.
Zenodo, zenodo.org/doi/10.5281/zenodo.17192402.

Richardson AD, Keenan TF, Migliavacca M, Ryu Y, Sonnentag O, Toomey M. **2013**. Climate
change, phenology, and phenological control of vegetation feedbacks to the climate system.
Agricultural and Forest Meteorology **169**: 156–173.

Rohde A, Bhalerao RP. **2007**. Plant dormancy in the perennial context. *Trends in Plant Science*
12: 217–223.

Signarbieux C, Toledano E, Sanginés de Carcer P, Fu YH, Schlaepfer R, Buttler A, Vitasse Y.
2017. Asymmetric effects of cooler and warmer winters on beech phenology last beyond
spring. *Global Change Biology* **23**: 4569–4580.

Singh RK, Bhalerao RP, Eriksson ME. **2021**. Growing in time: exploring the molecular
mechanisms of tree growth. *Tree Physiology* **41**: 657–678.

Singh RK, Svystun T, AlDahmash B, Jönsson AM, Bhalerao RP. **2017**. Photoperiod- and
temperature-mediated control of phenology in trees – a molecular perspective. *New*
Phytologist **213**: 511–524.

Steppe K, Sterck F, Deslauriers A. **2015**. Diel growth dynamics in tree stems: linking anatomy
and ecophysiology. *Trends in Plant Science* **20**: 335–343.

Švik M, Brovkina O, Veljanovski T, Čater M. **2025**. Phenological trends of European beech
stands along the Carpathian arc: a 20-year MOD13Q1/MYD13Q1 based analysis. *European*
Journal of Remote Sensing **58**: 2506576.

Tanino KK, Kalcsits L, Silim S, Kendall E, Gray GR. **2010**. Temperature-driven plasticity in
growth cessation and dormancy development in deciduous woody plants: a working
hypothesis suggesting how molecular and cellular function is affected by temperature during
dormancy induction. *Plant Molecular Biology* **73**: 49–65.

Thackeray SJ, Henrys PA, Hemming D, Bell JR, Botham MS, Burthe S, Helaouet P, Johns DG,
Jones ID, Leech DI, *et al.* 2016. Phenological sensitivity to climate across taxa and trophic
levels. *Nature* **535**: 241–245.

Tumajer J, Kašpar J, Kuželová H, Shishov VV, Tychkov II, Popkova MI, Vaganov EA, Treml V.
2021. Forward Modeling Reveals Multidecadal Trends in Cambial Kinetics and Phenology at
Treeline. *Frontiers in Plant Science* **12**: 613643.

Urban O, Klem K, Holišová P, Šigut L, Šprtová M, Teslová-Navrátilová P, Zitová M, Špunda V,
Marek MV, Grace J. **2014**. Impact of elevated CO2 concentration on dynamics of leaf
photosynthesis in Fagus sylvatica is modulated by sky conditions. *Environmental Pollution* **185**:
271–280.

Vitasse Y, François C, Delpierre N, Dufrêne E, Kremer A, Chuine I, Delzon S. **2011**. Assessing
the effects of climate change on the phenology of European temperate trees. *Agricultural and*
Forest Meteorology **151**: 969–980.

Yang LH, Rudolf VHW. **2010**. Phenology, ontogeny and the effects of climate change on the
timing of species interactions. *Ecology Letters* **13**: 1–10.

Zani D, Zani D, Crowther TW, Mo L, Renner SS, Zohner CM. **2020**. Increased growing-season
productivity drives earlier autumn leaf senescence in temperate trees. *Science*.

Zohner CM, Mirzagholi L, Renner SS, Mo L, Rebindaine D, Bucher R, Palouš D, Vitasse Y, Fu
YH, Stocker BD, *et al.* 2023. Effect of climate warming on the timing of autumn leaf senescence
reverses after the summer solstice. *Science* **381**: eadf5098.

Zohner CM, Renner SS. **2019**. Ongoing seasonally uneven climate warming leads to earlier
autumn growth cessation in deciduous trees. *Oecologia* **189**: 549–561.

Zweifel R, Sterck F, Braun S, Buchmann N, Eugster W, Gessler A, Häni M, Peters RL, Walthert
L, Wilhelm M, *et al.* 2021. Why trees grow at night. *New Phytologist* **231**: 2174–2185.

619 Figures

620
Figure 1. Conceptual model of autumn phenological responses of temperate trees to early-season
development and late-season temperature. Autumn phenology, represented in this study by the timing of
primary growth cessation (bud set) and leaf senescence (50% loss of leaf chlorophyll content), is influenced
by two opposing factors: early-season development and late-season temperature. **a)** In our model, early-
season development, which is driven by temperature, has an advancing effect on autumn phenology that
lasts until shortly after the summer solstice (green curve). Higher temperatures cause trees to complete
their annual life cycles faster, allowing them to set buds and senesce leaves. After the summer solstice, as
days shorten, trees become increasingly sensitive to cooling conditions, so late-season warming slows the
progression of bud set and senescence, delaying autumn phenology (red curve). The compensatory point,
where the advancing effect of early-season development is balanced by the delaying effect of late-season
warming, is represented by the green circle. **b)** According to the model, the timing of this effect reversal is
flexible and varies between years based on the speed of development. When development is slow or starts
late (blue curve), the effect reversal occurs later than under fast or early development (green curve).
Therefore, shortly after the solstice, the effect of temperature on autumn phenology differs between
fast/early and slow/late developing individuals. For example, point 1 (green circle) shows no net
temperature effect on phenology in fast/early trees. By contrast, point 2 (blue circle) shows that in slow/late
developing trees, warmer temperatures shortly after the solstice (in July) still advance autumn phenology.
However, as the growing season progresses and days shorten, trees become more responsive to cooling
regardless of prior developmental speed. By August, both fast/early and slow/late trees should therefore
exhibit similar phenological responses, with warming consistently delaying autumn phenology (points 3 and
4).
642

Spring
4 April - 24 May

June
24 May - 21 June

July
22 June - 23 July

August
24 July - 25 August

Figure 2. Depiction of the experimental timeline and settings for experiment 1. Each box corresponds to a specific treatment block at that point in time. The numbers inserted above the boxes refers to the treatment group (see Table 1 for details). Each box contains information on the location of the trees, the specific conditions they were under and the intended physiological effects of those conditions. The sapling graphics highlight differences in early-season developmental progression for the early-leafing and late-leafing groups. Following the August treatments all trees were placed outside under ambient conditions in a randomised block design.

Spring

Pre-solstice
22 May - 21 June

Post-solstice
22 June - 21 July

Autumn

Figure 3. Depiction of the experimental timeline and settings for experiment 2. Each box corresponds to a specific treatment block at that point in time. The text inserted above the boxes refers to the treatment group (see Table 2 for details). Each box contains information on the location of the trees and the specific conditions they were under. The sapling graphics indicate that all trees had equal opportunities for early-season development.

660
Figure 4. Effects of early-season development and late-season temperature on the timing of autumn bud set in *Fagus sylvatica* (experiment 1). a) Bud set dates for early- (green) and late- (blue) leafing trees including all treatments. Late-leafing trees were cooled (2-7°C) in climate chambers from 4 April to 24 May to arrest their development and delay leaf-out. **b)** Effects of July (22 June to 23 July) and August (24 July to 25 August) moderate cooling (8-13°C) on bud set date for early- (green) and late- (blue) leafing trees (see Fig. S3 for extreme cooling effects). Analyses show effect size means ± 95% confidence intervals from linear models, including treatment and bud-type (apical vs lateral) as predictors. Early-leafing effects are calculated against the early-leafing control and late-leafing effects are calculated against the late-leafing control. The bud type effect is not shown. Number labels (1-4) above each point are shown to aid comparison between points 1-4 in the conceptual model (Fig. 1b) and the observed effects. Positive values indicate advances in bud set and negative values indicate delays.

673
Figure 5. The pre- and post-solstice effects of night, full-day and day cooling on the timing of autumn
primary growth cessation in *Fagus sylvatica* (experiment 2). Effects of pre-solstice (22 May to 21 June) and
post-solstice (22 June to 21 July) cooling on bud set date. Full-day cooling trees were continuously cooled
to 8°C, day cooling trees were cooled to 8°C in the day and kept at 20°C at night, night cooling trees were
cooled to 8°C at night and kept at 20°C in the day. Analyses show effect size means ± 95% confidence
intervals from linear models, including treatment and bud-type (apical vs lateral) as predictors. The bud
type effect is not shown.
681
682

Figure 6. Diel patterns of relative growth, photosynthetic rate and theoretical thresholds for cold-induced bud set in *Fagus sylvatica*. Vertical red lines indicate the start of the day, and vertical blue lines indicate the start of the night, marking the boundaries between the 12-hour treatment windows used in experiment 2. **a)** The green curve shows relative carbon assimilation rate, the raw values were taken from the literature in a study that measured assimilation rates under controlled conditions (Urban *et al.*, 2014). Values were linearly interpolated between measurements then converted to a percentage of the peak value. Finally, the curve was smoothed by taking the running mean of the target value, the previous value and the following value. The black curve shows the relative probability for growth, the raw values were taken from the literature in a study that measured growth rates in the field (Zweifel *et al.*, 2021), then processed in the same way as the assimilation data. At night, low temperatures slow the trees' developmental processes, possibly leading to delayed tissue maturation, while low temperatures during the day reduce photosynthetic activity. **b) The Diel Cooling Hypothesis:** As days shorten after the solstice, autumn bud set becomes increasingly responsive to cooling. Temperatures below a certain threshold induce overwintering responses, advancing autumn phenology. Our results indicate that daytime cooling of 8°C is below this threshold. However, because daily temperatures reach their minimum during the night, trees' induction thresholds should be lower at night than during the day. Post-solstice night-time cooling of 8°C may thus have delayed bud set by slowing development rather than inducing overwintering responses.

Tables

Table 1. Description of the temperature treatments applied in experiment 1. July treatments were from
22 June to 23 July. August treatments were from 24 July to 25 August. Ambient means outside under natural
conditions. Temperature ranges indicate the daily minimum and maximum temperatures experienced by
trees inside climate chambers. The early-leafing and late-leafing trees were experimentally generated by
placing potted trees in climate chambers from 4 April to 24 May and cooling them to 2°C at night and 7°C
during the day to arrest spring development.

Treatment	1	2	3	4	5	6	7	8	9	10
Leaf-out timing	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late
Treatment timing	-	-	July	July	July	July	August	August	August	August
Cooling level	Control (ambient)	Control (ambient)	Moderate (8-13°C)	Moderate (8-13°C)	Extreme (2-7°C)	Extreme (2-7°C)	Moderate (8-13°C)	Moderate (8-13°C)	Extreme (2-7°C)	Extreme (2-7°C)

714 **Table 2. Description of the temperature treatments applied in experiment 2.** June treatments were from
715 22 May to 21 June. July treatments were from 22 June to 21 July. Ambient means outside under natural
716 conditions. The remaining temperature regimes are in the format day/night and refer to the temperatures
717 applied to trees in climate chambers.

Treatment	Control	Pre-Chamber control	Pre-Day	Pre-Night	Pre-Full-day	Post-Chamber control	Post-Day	Post-Night	Post-Full-day
Cooling time of year	-	June	June	June	June	July	July	July	July
Cooling time of day	-	-	Day	Night	Full-day	-	Day	Night	Full-day
Temperature regime	Ambient	20/20°C	8/20°C	20/8°C	8/8°C	20/20°C	8/20°C	20/8°C	8/8°C