1 Developmental constraints mediate the summer solstice reversal of climate effects
2 on European beech bud set
3

4 Dominic Rebindaine[1]*, Thomas W. Crowther[1], Susanne S. Renner[2], Zhaofei Wu[3], Yibiao Zou[1],
5 Lidong Mo[1], Haozhi Ma[4], Raymo Bucher[1], Constantin M. Zohner[1]
6
7 [1]Department of Environmental Systems Science, ETH Zurich (Swiss Federal Institute of
8 Technology), Universitätsstrasse 16, 8092 Zurich, Switzerland; [2]Department of Biology,
9 Washington University, Saint Louis, United States of America; [3]College of Water Sciences,
10 Beijing Normal University, Beijing, China; [4]WSL Swiss Federal Institute for Forest, Snow and
11 Landscape Research, Birmensdorf, Switzerland.
12
13 *Author for correspondence: Dominic Rebindaine, dominic.rebindaine@usys.ethz.ch

14
Abstract

Predicting the growing season of temperate zone trees under climate change requires

factoring in developmental constraints that determine tree resource allocation. Recent work

has identified the summer solstice (21 June) as a phenological "switch point," with pre-

solstice warming advancing autumn bud set and post-solstice warming delaying it. However,

the flexibility of this switch point remains poorly understood. We performed trans-solstice

climate manipulation experiments on potted European beech (*Fagus sylvatica*) saplings to

test (i) how spring leaf-out timing and June-August temperatures influence growth cessation

(bud set), and (ii) whether daytime and nighttime temperatures before and after the solstice

have different effects, given that trees primarily grow at night. Each day delay in spring leaf-

out delayed bud set by 0.24 ± 0.06 days on average. July cooling delayed bud set by 4.9 ± 2.6

days in late-leafing individuals but had no effect in early-leafing ones. Conversely, August

cooling advanced bud set by 4.4-4.5 ± 2.6 days in all trees. Daytime cooling before the

solstice had no effect, while after the solstice it advanced bud set. Nighttime cooling always

delayed bud set. Our findings support the Solstice-as-Phenology-Switch model and highlight

the central role of developmental progression in constraining the growing seasons of

temperate trees. Faster early-season development–especially under nighttime warming–advances the point at which trees become sensitive to late-season cooling, thereby triggering senescence earlier. To improve growing season length projections, phenological models must account for these developmentally-mediated and diel-specific responses to warming.

Keywords: Carbon cycle | Climate change | Global warming | Plant-climate interactions | Plant development| Plant growth | Phenology

Introduction

Climate shifts are leading to rapid, species-specific changes in phenology and ecosystem productivity (Boisvenue & Running, 2006; Menzel *et al.*, 2006; Yang & Rudolf, 2010; Thackeray *et al.*, 2016). In temperate forests, changes in the timing of spring leaf-out, autumn leaf-senescence and bud set are modifying water, energy and carbon cycles (Peñuelas *et al.*, 2009; Richardson *et al.*, 2013), with extended growing seasons increasing net ecosystem carbon uptake by up to 9.8 gC m^{-2} day^{-1} (Keenan *et al.*, 2014). Therefore, understanding the interaction between climate change and temperate forest phenology is pivotal to improving forecasts of community dynamics and carbon sequestration.

The past few decades have seen delays in the onset of temperate autumn phenology, but these changes are much smaller in magnitude compared to the advances in spring leaf-out observed during the same period (Gill *et al.*, 2015; Piao *et al.*, 2019). This is unexpected, given that experiments have demonstrated a high sensitivity of leaf senescence to autumn warming, with phenological responses even surpassing the temperature sensitivity (days per

°C) of spring leaf-out (Fu *et al.*, 2018). One possible explanation for this discrepancy is that other factors may counterbalance the effects of autumn warming, with some studies finding that earlier leaf-out leads to earlier leaf senescence (Fu *et al.*, 2014; Keenan & Richardson, 2015; Zani *et al.*, 2020). This connection between spring and autumn phenophases could be due to developmental and nutrient constraints that affect carbon source-sink dynamics in temperate trees (Paul & Foyer, 2001; Zani *et al.*, 2020; Zohner *et al.*, 2023; Gessler & Zweifel, 2024).

The Solstice-as-Phenology-Switch hypothesis, supported by experiments and monitoring data, posits that warmer air temperatures before the summer solstice accelerate tree growth and development, leading to an earlier onset of leaf senescence, whereas warmer temperatures after the solstice slow the progression of leaf senescence (Zohner *et al.,* 2023). This reversal in temperature effects provides an explanation for the less pronounced shifts in autumn phenology compared to spring (Piao *et al.*, 2019). The effect reversal after the solstice points to the role of photoperiod in regulating plant physiology (Bauerle *et al.*, 2012), with trees becoming increasingly responsive to cooling temperatures as days continue to shorten. The precise timing of this reversal appears to be flexible and has been advancing in recent decades, potentially due to faster development driven by climate warming (Zohner *et al.*, 2023). For temperate trees, the key outcomes of development are production of viable seeds, cessation of primary and secondary growth, and maturation of tissues including leaf or flower buds (bud set) ready for overwintering before the onset of frost (Rohde & Bhalerao, 2007; Tanino *et al.*, 2010; Cooke *et al.*, 2012). Because opportunities for development will vary among years, the timing of the temperature effect

reversal should be flexible. However, this flexibility has yet to be demonstrated in processes

directly related to development such as autumn bud set.

In this study, we investigate how changes in air temperature around the summer solstice

affect primary growth cessation, using bud set as a key physiological marker for the end of

primary growth and the beginning of autumn phenology (Zohner & Renner, 2019).

Specifically, we explore the effects of pre- and post-solstice cooling, and of daytime versus

nighttime cooling, on bud set timing. Diel temperature variations are relevant because trees

mainly grow at night when temperatures and water deficit are lower than during the day

(Zweifel *et al.*, 2021). Between 1950 and 2004, nighttime temperatures rose faster than

daytime temperatures (Vose et al., 2005), but from 1991 to 2020 this appears to have

reversed (Zhong et al., 2023). We also experimentally created early-leafing vs. late-leafing

trees to assess whether slowed early-season development postpones the reversal date of

temperature effects on autumn phenology (Fig. 1). As depicted in our conceptual model (Fig.

1), tree individuals are dynamic systems that continuously integrate and adapt their

physiological responses to environmental conditions over time. Their responses to

temperature between June and August and between day and night are therefore expected

to differ, an expectation tested in this study.

Results

Experiment 1

Linear mixed-effects modelling showed that across all treatments, late-leafing trees cooled

during spring experienced a delay to their bud set of 4.6 ± 1.3 days (mean ± 2 SE). Across all

treatment pairings, the mean bud set date for the late-leafing group always occurred later

than for the early-leafing group (Fig. S2). On average, each day delay in spring leaf-out was

associated with a delay of 0.24 ± 0.06 days in bud set timing.

The effect of July cooling differed strongly between the early- and late-leafing trees: Within

the early-leafing group, moderate cooling in July lead to a small, non-significant delay in bud

set compared to the ambient July treatment (1.4 ± 2.6 days). July cooling had a much greater

impact on late-leafing trees, leading to a 4.9 ± 2.6-day delay in bud set compared to the

ambient July treatment in the late-leafing group.

In contrast to July cooling, the effects of August cooling did not differ between early- and

late-leafing trees, advancing bud set in all treatment groups (Fig. 2b). Cooling in August lead

to a 4.5 ± 2.6-day and 4.4 ± 2.6-day advancement in bud set for early- and late-leafing trees,

respectively. The extreme cooling treatments showed similar patterns, although slightly less

clearly (Fig. S3). Effects of absolute and relative growth on bud set date can be found in the

supporting information (Figs. S4-S5). Bud set timing had no effect on final bud length (Fig.

S6).

Experiment 2

Linear mixed-effects modelling showed that pre-solstice full-day (day and night) cooling

delayed autumn bud set by 4.1 ± 3.6 days (mean ± 2SE) (Fig. 3). Pre-solstice nighttime

cooling had a similar effect, delaying bud set by 4.2 ± 3.6 days. By contrast, pre-solstice

daytime cooling had no significant effect on bud set (0.2 ± 3.7 days). Post-solstice full-day

cooling advanced autumn bud set by 5.2 ± 3.6 days. Similarly, post-solstice daytime cooling

advanced bud set by 5.3 ± 4.3 days. Conversely, post-solstice nighttime cooling delayed bud

set by 3.8 ± 3.5 days.

Discussion

In this study, we conducted experiments on European beech to test how primary growth

cessation and bud set respond to monthly and diel temperature changes around the

summer solstice. We found large differences in the responses between trees subjected to

daytime versus nighttime cooling before and after the solstice (Fig. 3) and between early-

and late-leafing trees (Fig. 2). These differences suggest that early-season developmental

timing not only influences the timing of autumn bud set *per se*, but also plays a critical role

in determining the timing of the summer solstice effect reversal. Therefore, our findings

support our conceptual model (Fig. 1), highlighting development as a key factor

underpinning the Solstice-as-Phenology-Switch hypothesis. In the following sections, we

discuss the effects of pre-solstice and post-solstice air temperature on development, along

with the distinct impacts of daytime versus nighttime temperature variations.

Early-season development rates alter responses to late-season temperature

Our trans-solstice climate manipulation experiments showed that trees delayed their

aboveground primary growth cessation (bud set) in response to slow/late early-season

development (delayed spring leaf-out). Regardless of the summer cooling treatments, late-

leafing trees consistently set buds after their early-leafing counterparts (Fig. S2), with each

day delay in spring leaf-out delaying bud set by an average of 0.24 days. This is in line with

previous studies demonstrating a tight linkage between within-year variations in spring and

autumn phenology (Fu *et al.*, 2014; Keenan & Richardson, 2015; Signarbieux *et al.*, 2017),

likely governed by developmental constraints (Zohner *et al.*, 2023), buildup of water and

nutrient stress (Paul & Foyer, 2001; Buermann *et al.*, 2018; Bigler & Vitasse, 2021), or leaf

aging (Lim *et al.*, 2007).

July cooling induced a delay in bud set dates 3.5 times greater in late-leafing trees compared

to early-leafing ones (4.9 versus 1.4 days delay), which agrees with the expectations derived

from our conceptual model (Fig. 1). As we had expected, the natural interannual variation in

early-season development causes year-to-year differences in the point at which cool

temperatures switch from delaying bud set to advancing it. This offers a physiological

explanation for the advancement in the effect reversal timing observed between 1966 and

2015 in *Fagus sylvatica, Aesculus hippocastanum, Quercus robur, and Betula pendula*

(Zohner *et al.*, 2023).

August cooling induced comparable advances in bud set timing in both early- and late-

leafing trees (4.4-4.5 days). As trees become increasingly responsive to cooling with

shortening days (Delpierre *et al.*, 2009; Körner *et al.*, 2016), they exhibit increasingly similar

responses to late-season (August) cooling, regardless of their prior developmental speeds

(Fig. 1). That the differences in the responses of early-leafing and late-leafing trees to cooling

diminishes between July and August suggests tree individuals are dynamic systems that

continuously integrate and adapt their physiological responses to environmental conditions

over time.

Effects of daytime vs. nighttime temperature

Our experiments showed that daytime and nighttime temperatures had different effects on

primary growth cessation before and after the summer solstice: Before the solstice, daytime

cooling had no impact on the timing of bud set, while nighttime cooling and full-day cooling

delayed bud set by 4.2 and 4.1 days, respectively (Fig. 3). Given that experimental cooling

reduced photosynthesis by 69-83% across all treatments (Fig. S9), these findings underscore

the role of developmental processes—such as cell division and expansion—which mostly

occur at night (Zweifel et al., 2021; Fig. 4a). The developmental arrest induced by night

cooling—and therefore also by full-day cooling—likely slowed bud growth and maturation,

effectively extending the growing season.

After the solstice, daytime and nighttime cooling of 8°C elicited opposite responses. Trees

subjected to post-solstice daytime cooling and full-day cooling set their buds earliest, more

than five days earlier on average than the control group. As days shorten, the temperature

threshold inducing this growth cessation must be lower at night, otherwise trees would risk

prematurely ending their growing season due to the daily minimum temperatures occurring

at night (Fig. 4b). This may explain why post-solstice nighttime cooling delayed, rather than

advanced, bud set: Cooling of 8°C was sufficient to trigger overwintering responses when

applied during the day but not during the night (see Fig. S10 for a conceptual schematic). It

is also possible that daytime cooling indirectly triggered bud set by reducing photosynthesis

(Fig 4a). Such a mechanism is based on the assumption that trees will prioritise dormancy

induction over growth once photosynthesis is not possible anymore (Körner *et al.*, 2016).

Indeed, downregulation of photosynthesis has previously been reported to accelerate

senescence (Krieger-Liszkay *et al.*, 2019), although this may simply follow a reduction in

growth activity (Paul & Foyer, 2001). However, reduced photosynthesis has also been linked

to delayed senescence in both observational (Meng *et al.*, 2023) and experimental studies

(Vitasse *et al.*, 2021; Zohner *et al.*, 2023). Therefore, the direct effects of post-solstice

photosynthesis on autumn phenology still need to be clarified.

Nighttime cooling always led to a delay in bud set of approximately four days (Fig. 3). As

trees mainly grow at night (Zweifel *et al.*, 2021), colder nighttime temperatures slow down

key developmental processes such as meristematic activity, tissue expansion and

maturation, which in turn likely caused the delays in primary growth cessation (Fig. 4). Cold

autumn nights also delay growth cessation and slow bud development in *Populus, Pinus* and

Picea species (Kramer, 1956; Malcolm & Pymar, 1975; Kalcsits *et al.*, 2009). Given the

consistency of the effect reversal across Northern Hemisphere temperate forests (Zohner *et*

al., 2023), and the patterns seen in previous experiments (Kramer, 1956; Malcolm & Pymar,

1975; Kalcsits *et al.*, 2009; Zohner *et al.*, 2023), our results likely are generalisable across

temperate tree species.

In conclusion, earlier primary growth cessation in temperate trees under climate change can

be explained by higher developmental rates under warmer temperatures, an effect

predominantly driven by nighttime temperatures (since trees primarily grow at night).

Ultimately, to better predict future growing season lengths, models must begin to account

for the differential impacts of the annual and daily temperature cycles that temperate zone

trees are physiologically adapted to.

Materials and Methods

Experiment 1

To test the antagonistic effects of early-season development and late-season air

temperature on autumn primary growth phenology, we set up an experimental population

(n = 267) of 40-60 cm tall European beech (*Fagus sylvatica* L.) trees in Zurich, Switzerland in

2023. The trees were sourced from a local nursery, and each tree was placed individually in a

20 L plastic pot containing a 1:1:1 sand / peat / organic soil mixture with a Nitrogen (N)

concentration of ~65 g m^{-3}, a Phosphate (P_2O_5) concentration of ~140 g m^{-3}, and a Potassium

(K_2O) concentration of ~400 g m^{-3}. Individual trees were assigned randomly to one of 10

treatment groups (26 ≤ n ≤ 27). Treatment groups were cooled at different times of the year,

using different cooling levels (Table 1). To arrest spring development and thereby generate

early-leafing and late-leafing individuals, some trees were placed outside under ambient

conditions, while others were cooled in climate chambers from 4 April to 24 May. The

chambers were set to a low of 2°C at night and a high of 7°C during the day, following a

simulated day-night cycle of temperature and light availability (13 h photoperiod at ~4,300

lux). Temperatures around 5°C are associated with low rates of growth and development

processes such as cell division, expansion and maturation (Körner, 2003; Tumajer *et al.*,

2021). From 24 May to 21 June, all trees were kept outside under ambient conditions in a

randomised block design. All trees were monitored to observe their individual leaf-out

dates, which was defined as the date when >50% of their leaves had unfolded,

corresponding to BBCH15 (Capdevielle-Vargas *et al.*, 2015).

During the summer treatments, control trees were placed outside under natural ambient

conditions. We did not use climate chambers for the control groups because warm

conditions in these chambers can introduce unique physiological stressors—such as aphid

infestations—not present in cold chambers (Bezemer *et al.*, 1998). We noticed increased

aphid abundance in control chambers during experiment 2, however these stressors did not

appear to affect autumn phenology (see Experiment 2). The July treatments took place

directly after the summer solstice from 22 June to 23 July. Treated trees were placed in

climate chambers set to a low of 2°C at night and a high of 7°C during the day or a low of 8°C

at night and a high of 13°C during the day depending on their cooling level (extreme and

moderate, respectively). Trees in the chambers experienced a photoperiod of 16 h at ~7,300

lux. The extreme cooling was designed to severely impair cell division, expansion, and

maturation as described above. Additionally, under this temperature regime, photosynthesis

should be reduced by >40% (Körner, 2006). The moderate cooling regime should still impair

growth and development in temperate deciduous trees as well as reducing photosynthesis

by at least 30% (Körner, 2003, 2006; Lenz *et al.*, 2014). The August treatments took place

from 24 July to 25 August under the same conditions as the July treatments except with a 15

h photoperiod. For the remainder of the experiment, all trees were kept outside in a

randomised block design. Throughout the experiment, all trees were watered frequently to

ensure constant water supply (Fig. S1).

To observe the effects of our treatments on the development of overwintering buds, we

monitored bud growth to derive bud set dates, our marker for the cessation of primary

aboveground development and the beginning of autumn phenology, for all trees. On each

tree, the terminal bud of the primary shoot and the terminal bud on a random lateral stem

were selected and tagged for measurement. Each selected bud was measured to 0.01 mm

precision using a digital calliper (CD-P8"M, Mitutoyo Corp, Japan). We measured all buds on

a regular basis from 4 July to 2 November. Bud set was defined as the date when each bud

reached 90% of its own maximum length, which is considered to be an indicative stage of

aboveground primary growth cessation (Signarbieux *et al.*, 2017; Zohner & Renner, 2019).

Bud lengths were linearly interpolated between measurement dates to derive the date at

which they reached the 90% threshold.

Experiment 2

To observe the effects of pre- and post-solstice daytime and nighttime temperature on bud

set phenology, we set up an experimental population (n = 180) of four-year-old *Fagus*

sylvatica trees in Zurich, Switzerland in 2022. The trees were sourced from a local nursery

and each tree was placed individually in a 20 L plastic pot containing a 1:1:1 sand / peat /

organic soil mixture with a Nitrogen (N) concentration of ~65 g m^{-3}, a Phosphate (P$_2$O$_5$)

concentration of ~140 g m^{-3}, and a Potassium (K$_2$O) concentration of ~400 g m^{-3}.

The ambient control treatment consisted of 36 trees exposed to natural ambient conditions.

The remaining eight treatments were each applied to 18 trees and included cooling in

climate chambers with simulated ambient day length (16 h) and light intensity (~6,900 lux).

The pre-solstice cooling treatments were applied between 22 May and 21 June. The post-

solstice cooling treatments were applied between 22 June and 21 July. The pre- and post-

solstice treatments included four levels each: Chamber control, where the trees were

continuously subjected to 20°C; Day cooling, where trees were subjected to 8°C in the

daytime and 20°C at night; Night cooling, where trees were subjected to 20°C in the day and

8°C at night; Full-day cooling, where trees were continuously subjected to 8°C (Table 2). Due

to warm temperatures, chamber control trees were subject to an increase in aphid

abundance, however this did not alter their bud set timing (see data analyses). After

treatment, all trees were placed in a randomised block design outside under ambient

conditions. Soil moisture content was regulated by frequent watering. We measured all buds

on a weekly basis from 25 August to 3 November following the same methodology as in

experiment 1. We also measured leaf-level CO_2 assimilation rates (see Zohner *et al.,* 2023 for

methodology).

Data analyses

For experiment 2, we performed a one-way ANOVA that showed no significant differences

between the bud set dates of the ambient, pre-chamber and post-chamber control groups

($F_{2,136}$ = 0.346, p =0.708, see Fig. S7). Therefore, for the following analyses we treated all three

control treatments as one control treatment with 72 trees. For both experiments we ran

linear mixed-effects models using treatment as a categorical fixed effect, bud type (apical vs.

lateral) as a random effect and bud set day-of-year, absolute bud growth or relative bud

growth as the response variable. Effect sizes were calculated in comparison to the

corresponding control treatment, i.e., all early-leafing treatments were compared to the

early-leafing control treatment and all late-leafing treatments were compared to the late-

leafing control. Absolute and relative bud growth were calculated as:

$$Absolute = max\ length - min\ length$$

$$Relative = \left(1 - \frac{min\ length}{max\ length}\right) * 100$$

311 Where:

312 *min length* = the first measured bud length

313 *max length* = the measured bud length when the bud first surpassed 90% of its final length

To determine the sensitivity of autumn bud set timing to spring leaf-out timing, we ran a

linear mixed-effects model using leaf-out day-of-year as the fixed predictor variable, bud

type and summer temperature treatment as random effects, and bud set day-of-year as the

response variable.

Acknowledgments

C.M.Z. was supported by the SNF Ambizione Fellowship programme (no. PZ00P3_193646) and T.W.C. by DOB Ecology and the Bernina Foundation.

Competing interests

The authors declare no competing interests.

Author contributions

D.R. conceived and developed the study, conducted the experiments and analyses, and wrote the manuscript. C.M.Z equally contributed to the conception and development of the study, contributed to the experiments and analyses, and the text. L.M., H.M., Z.W. and Y.Z. contributed to the conception and implementation of the first experiment. R.B. contributed to the conception and implementation of the second experiment. T.W.C. and S.S.R. contributed to the text. All authors provided comments and approved the final manuscript.

Data availability

The code and data for this study will be made available upon request or at Zenodo on publication of the manuscript.

References

Bauerle WL, Oren R, Way DA, Qian SS, Stoy PC, Thornton PE, Bowden JD, Hoffman FM, Reynolds RF. **2012**. Photoperiodic regulation of the seasonal pattern of photosynthetic capacity and the implications for carbon cycling. *Proceedings of the National Academy of Sciences* **109**: 8612–8617.

Bezemer TM, Jones TH, Knight KJ. **1998**. Long-term effects of elevated CO_2 and temperature on populations of the peach potato aphid Myzus persicae and its parasitoid Aphidius matricariae. *Oecologia* **116**: 128–135.

Bigler C, Vitasse Y. **2021**. Premature leaf discoloration of European deciduous trees is caused by drought and heat in late spring and cold spells in early fall. *Agricultural and Forest Meteorology* **307**: 108492.

Boisvenue C, Running SW. 2006. Impacts of climate change on natural forest productivity - evidence since the middle of the 20th century: climate change impacts on forest vegetation. *Global Change Biology* **12**: 862–882.

Buermann W, Forkel M, O'Sullivan M, Sitch S, Friedlingstein P, Haverd V, Jain AK, Kato E, Kautz M,
Lienert S, *et al.* **2018**. Widespread seasonal compensation effects of spring warming on northern plant
productivity. *Nature* **562**: 110–114.

Capdevielle-Vargas R, Estrella N, Menzel A. **2015**. Multiple-year assessment of phenological plasticity
within a beech (Fagus sylvatica L.) stand in southern Germany. *Agricultural and Forest Meteorology* **211–**
212: 13–22.

Cooke JEK, Eriksson ME, Junttila O. **2012**. The dynamic nature of bud dormancy in trees: environmental
control and molecular mechanisms. *Plant, Cell & Environment* **35**: 1707–1728.

Delpierre N, Dufrêne E, Soudani K, Ulrich E, Cecchini S, Boé J, François C. **2009**. Modelling interannual
and spatial variability of leaf senescence for three deciduous tree species in France. *Agricultural and*
Forest Meteorology **149**: 938–948.

Fu YSH, Campioli M, Vitasse Y, De Boeck HJ, Van den Berge J, AbdElgawad H, Asard H, Piao S, Deckmyn
G, Janssens IA. **2014**. Variation in leaf flushing date influences autumnal senescence and next year's
flushing date in two temperate tree species. *Proceedings of the National Academy of Sciences* **111**: 7355–
7360.

Fu YH, Piao S, Delpierre N, Hao F, Hänninen H, Liu Y, Sun W, Janssens IA, Campioli M. **2018**. Larger
temperature response of autumn leaf senescence than spring leaf-out phenology. *Global Change Biology*
24: 2159–2168.

Gessler A, Zweifel R. **2024**. Beyond source and sink control – toward an integrated approach to
understand the carbon balance in plants. *New Phytologist* **242**: 858–869.

Gill AL, Gallinat AS, Sanders-DeMott R, Rigden AJ, Short Gianotti DJ, Mantooth JA, Templer PH. **2015**.
Changes in autumn senescence in northern hemisphere deciduous trees: a meta-analysis of autumn
phenology studies. *Annals of Botany* **116**: 875–888.

Kalcsits L, Kendall E, Silim S, Tanino K. **2009**. Magnetic resonance microimaging indicates water diffusion
correlates with dormancy induction in cultured hybrid poplar (Populus spp.) buds. *Tree Physiology* **29**:
1269–1277.

Keenan TF, Gray JM, Friedl MA, Toomey M, Bohrer G, Hollinger DY, Munger JW, O'Keefe J, Schmid HP,
Wing IS, *et al.* **2014**. Net carbon uptake has increased through warming-induced changes in temperate
forest phenology. *Nature Climate Change*.

Keenan TF, Richardson AD. **2015**. The timing of autumn senescence is affected by the timing of spring
phenology: implications for predictive models. *Global Change Biology* **21**: 2634–2641.

Körner C. **2003**. *Alpine Plant Life*. Berlin, Heidelberg: Springer Berlin Heidelberg.

Körner C. **2006**. Significance of Temperature in Plant Life. In: Morison JIL, Morecroft MD, eds. Plant
Growth and Climate Change. Wiley, 48–69.

Körner C, Basler D, Hoch G, Kollas C, Lenz A, Randin CF, Vitasse Y, Zimmermann NE. **2016**. Where, why
and how? Explaining the low-temperature range limits of temperate tree species (M Turnbull, Ed.).
Journal of Ecology **104**: 1076–1088.

Kramer PJ. **1936**. Effect of variation in length of day on growth and dormancy of trees. *Plant Physiology*
11: 127–137.

Kramer PJ. **1956**. Some effects of various combinations of day and night temperatures and photoperiod
on the height growth of loblolly pine seedlings. *Forest Science* **3**: 45–55.

Krieger-Liszkay A, Krupinska K, Shimakawa G. **2019**. The impact of photosynthesis on initiation of leaf
senescence. *Physiologia Plantarum* **166**: 148–164.

Lenz A, Vitasse Y, Hoch G, Körner C. **2014**. Growth and carbon relations of temperate deciduous tree
species at their upper elevation range limit (M Turnbull, Ed.). *Journal of Ecology* **102**: 1537–1548.

Lim PO, Kim HJ, Gil Nam H. **2007**. Leaf Senescence. *Annual Review of Plant Biology* **58**: 115–136.

Malcolm DC, Pymar CF. **1975**. The influence of temperature on the cessation of height growth of Sitka
spruce (Picea sitchensis Bong. Carr.). *Silvae Genet* **24**: 5–6.

Meng F, Liu D, Wang Y, Wang S, Wang T. **2023**. Negative relationship between photosynthesis and late-
stage canopy development and senescence over Tibetan Plateau. *Global Change Biology* **29**: 3147–3158.

Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kübler K, Bissolli P, Braslavská O, Briede A,
et al. **2006**. European phenological response to climate change matches the warming pattern: european
phenological response to climate change. *Global Change Biology* **12**: 1969–1976.

Paul MJ, Foyer CH. **2001**. Sink regulation of photosynthesis. *Journal of Experimental Botany* **52**: 1383–
1400.

Peñuelas J, Rutishauser T, Filella I. **2009**. Phenology Feedbacks on Climate Change. *Science* **324**: 887–888.

Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, Liu L, Lian X, Shen M, Zhu X. **2019**. Plant phenology and
global climate change: Current progresses and challenges. *Global Change Biology* **25**: 1922–1940.

Richardson AD, Keenan TF, Migliavacca M, Ryu Y, Sonnentag O, Toomey M. **2013**. Climate change,
phenology, and phenological control of vegetation feedbacks to the climate system. *Agricultural and*
Forest Meteorology **169**: 156–173.

Rohde A, Bhalerao RP. **2007**. Plant dormancy in the perennial context. *Trends in Plant Science* **12**: 217–
223.

Signarbieux C, Toledano E, Sanginés de Carcer P, Fu YH, Schlaepfer R, Buttler A, Vitasse Y. **2017**.
Asymmetric effects of cooler and warmer winters on beech phenology last beyond spring. *Global Change*
Biology **23**: 4569–4580.

Tanino KK, Kalcsits L, Silim S, Kendall E, Gray GR. **2010**. Temperature-driven plasticity in growth cessation
and dormancy development in deciduous woody plants: a working hypothesis suggesting how molecular
and cellular function is affected by temperature during dormancy induction. *Plant Molecular Biology* **73**:
49–65.

Thackeray SJ, Henrys PA, Hemming D, Bell JR, Botham MS, Burthe S, Helaouet P, Johns DG, Jones ID,
Leech DI, ***et al.*** **2016**. Phenological sensitivity to climate across taxa and trophic levels. *Nature* **535**: 241–
245.

Tumajer J, Kašpar J, Kuželová H, Shishov VV, Tychkov II, Popkova MI, Vaganov EA, Treml V. **2021**.
Forward Modeling Reveals Multidecadal Trends in Cambial Kinetics and Phenology at Treeline. *Frontiers in*
Plant Science **12**: 613643.

Urban O, Klem K, Holišová P, Šigut L, Šprtová M, Teslová-Navrátilová P, Zitová M, Špunda V, Marek MV,
Grace J. **2014**. Impact of elevated CO2 concentration on dynamics of leaf photosynthesis in Fagus sylvatica
is modulated by sky conditions. *Environmental Pollution* **185**: 271–280.

Vitasse Y, Baumgarten F, Zohner CM, Kaewthongrach R, Fu YH, Walde MG, Moser B. **2021**. Impact of
microclimatic conditions and resource availability on spring and autumn phenology of temperate tree
seedlings. *New Phytologist* **232**: 537–550.

Yang LH, Rudolf VHW. **2010**. Phenology, ontogeny and the effects of climate change on the timing of
species interactions. *Ecology Letters* **13**: 1–10.

Zani D, Zani D, Crowther TW, Mo L, Renner SS, Zohner CM. **2020**. Increased growing-season productivity
drives earlier autumn leaf senescence in temperate trees. *Science*.

Zohner CM, Mirzagholi L, Renner SS, Mo L, Rebindaine D, Bucher R, Palouš D, Vitasse Y, Fu YH, Stocker
BD, *et al.* **2023**. Effect of climate warming on the timing of autumn leaf senescence reverses after the
summer solstice. *Science* **381**: eadf5098.

Zohner CM, Renner SS. **2019**. Ongoing seasonally uneven climate warming leads to earlier autumn
growth cessation in deciduous trees. *Oecologia* **189**: 549–561.

Zweifel R, Sterck F, Braun S, Buchmann N, Eugster W, Gessler A, Häni M, Peters RL, Walthert L, Wilhelm
M, *et al.* **2021**. Why trees grow at night. *New Phytologist* **231**: 2174–2185.

Figures

446
Figure 1. Conceptual model of autumn phenological responses of temperate trees to early-season
development and late-season temperature. Autumn phenology, represented in this study by the timing
of primary growth cessation (bud set), is influenced by two opposing factors: early-season development
and late-season temperature. **a)** In our model, early-season development, which is driven by
temperature, has an advancing effect on autumn phenology that lasts until shortly after the summer
solstice (green curve). Higher temperatures cause trees to complete their annual life cycles faster,
allowing them to set buds. After the summer solstice, as days shorten, trees become increasingly sensitive
to cooling conditions, so late-season warming slows the progression of bud set and senescence, delaying
autumn phenology (red curve). The compensatory point, where the advancing effect of early-season
development is balanced by the delaying effect of late-season warming, is represented by the green
circle. **b)** According to the model, the timing of this effect reversal is flexible and varies between years
based on the speed of development. When development is slow or starts late (blue curve), the effect
reversal occurs later than under fast or early development (green curve). Therefore, shortly after the
solstice, the effect of temperature on autumn phenology differs between fast/early and slow/late
developing individuals. For example, point 1 (green circle) shows no net temperature effect on phenology
in fast/early trees. By contrast, point 2 (blue circle) shows that in slow/late developing trees, warmer
temperatures shortly after the solstice (in July) still advance autumn phenology. However, as the growing
season progresses and days shorten, trees become more responsive to cooling regardless of prior
developmental speed. By August, both fast/early and slow/late trees should therefore exhibit similar
phenological responses, with warming consistently delaying autumn phenology (points 3 and 4).
467

Figure 2. Effects of early-season development and late-season temperature on the timing of autumn bud set in *Fagus sylvatica* (experiment 1). a) Bud set dates for early- (green) and late- (blue) leafing trees including all treatments. Late-leafing trees were cooled (2-7°C) in climate chambers from 4 April to 24 May to arrest their development and delay leaf-out. **b)** Effects of July (22 June to 23 July) and August (24 July to 25 August) moderate cooling (8-13°C) on bud set date for early- (green) and late- (blue) leafing trees (see Fig. S3 for extreme cooling effects). Analyses show effect size means ± 95% confidence intervals from linear mixed models, including treatment as fixed effect and bud-type (apical vs lateral) as random effect. Early-leafing effects are calculated against the early-leafing control and late-leafing effects are calculated against the late-leafing control. Number labels (1-4) above each point are shown to aid comparison between points 1-4 in the conceptual model (Fig. 1b) and the observed effects. Positive values indicate advances in bud set and negative values indicate delays.

481
Figure 3. The pre- and post-solstice effects of night, full-day and day cooling on the timing of autumn
primary growth cessation in _Fagus sylvatica_ (experiment 2). Effects of pre-solstice (22 May to 21 June)
and post-solstice (22 June to 21 July) cooling on bud set date. Full-day cooling trees were continuously
cooled to 8°C, day cooling trees were cooled to 8°C in the day and kept at 20°C at night, night cooling
trees were cooled to 8°C at night and kept at 20°C in the day. Analyses show effect size means ± 95%
confidence intervals from linear mixed models, including treatment as fixed effect and bud-type (apical vs
lateral) as random effect. Positive values indicate advances to bud set and negative values indicate delays.
489

Figure 4. Diel patterns of relative growth, photosynthesis and theoretical thresholds for cold-induced bud set in *Fagus sylvatica*. Vertical red lines indicate the start of the day, and vertical blue lines indicate the start of the night, marking the boundaries between the 12-hour treatment windows used in experiment 2. **a)** The green curve shows relative carbon assimilation rate measured under controlled conditions (Urban *et al.*, 2014), with values linearly interpolated between measurements. The black curve shows the relative probability for growth (Zweifel *et al.*, 2021). Both curves were smoothed by taking the running mean of the target value, the previous value and the following value. At night, low temperatures slow the trees' developmental processes, possibly leading to delayed primary growth cessation, while low temperatures during the day reduce photosynthetic activity. **b)** As days shorten after the solstice, autumn bud set becomes increasingly responsive to cooling. Temperatures below a certain threshold induce overwintering responses, advancing autumn phenology. Our results indicate that daytime cooling of 8°C is below this threshold. However, because daily temperatures reach their minimum during the night, trees' induction thresholds should be lower at night than during the day. Post-solstice night-time cooling of 8°C may thus have delayed senescence by slowing development rather than inducing it.

Tables

Table 1. Description of the temperature treatments applied in experiment 1. July treatments were from
22 June to 23 July. August treatments were from 24 July to 25 August. Ambient means outside under
natural conditions. Temperature ranges indicate the daily minimum and maximum temperatures
experienced by trees inside climate chambers. The early-leafing and late-leafing trees were
experimentally generated by placing potted trees in climate chambers from 4 April to 24 May and cooling
them to 2°C at night and 7°C during the day to arrest spring development.

Treatment	1	2	3	4	5	6	7	8	9	10
Leaf-out timing	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late
Treatment timing	-	-	July	July	July	July	August	August	August	August
Cooling level	Control (ambient)	Control (ambient)	Moderate (8-13°C)	Moderate (8-13°C)	Extreme (2-7°C)	Extreme (2-7°C)	Moderate (8-13°C)	Moderate (8-13°C)	Extreme (2-7°C)	Extreme (2-7°C)

523 **Table 2. Description of the temperature treatments applied in experiment 2.** June treatments were from
524 22 May to 21 June. July treatments were from 22 June to 21 July. Ambient means outside under natural
525 conditions. The remaining temperature regimes are in the format day/night and refer to the temperatures
526 applied to trees in climate chambers.

Treatment	Control	Pre-Chamber control	Pre-Day	Pre-Night	Pre-Full-day	Post-Chamber control	Post-Day	Post-Night	Post-Full-day
Cooling time of year	-	June	June	June	June	July	July	July	July
Cooling time of day	-	-	Day	Night	Full-day	-	Day	Night	Full-day
Temperature regime	Ambient	20/20°C	8/20°C	20/8°C	8/8°C	20/20°C	8/20°C	20/8°C	8/8°C

Supporting Information

Figure S1. Seasonal soil water content (%) for each treatment (experiment 1). Five individuals per treatment were measured at each time step.

Figure S2. Effects of slow spring development on the timing of autumn bud set in *Fagus sylvatica* (experiment 1). Spring development was slowed by cooling the late-leafing trees from 4 April to 24 May in climate chambers. The late-leafing trees experienced a simulated day-night cycle of temperature and light availability, with minimum (2°C) and maximum (7°C) temperatures reached during the night- and daytime, respectively and otherwise simulated ambient conditions. Analyses show effect size means ± 95% confidence intervals from linear mixed models, including treatment as fixed effect and bud-type (apical vs lateral) as random effect. For each late-leafing treatment, the effect is calculated against the

548 equivalent early-leafing treatment. Positive values indicate advances to bud set and negative values
549 indicate delays.

550
Figure S3. Effects of early-season development and extreme late-season cooling on the timing of
autumn primary growth cessation in *Fagus sylvatica* (experiment 1). Effects of July (22 June to 23 July)
and August (24 July to 25 August) extreme cooling (2-7°C) on bud set date for early- (grey) and late- (blue)
leafing trees. Analyses show effect size means ± 95% confidence intervals from linear mixed models,
including treatment as fixed effect and bud-type (apical vs lateral) as random effect. Early-leafing effects
are calculated against the early-leafing control and late-leafing effects are calculated against the late-
leafing control. Positive values indicate advances to bud set and negative values indicate delays.

564
Figure S4. Effects of early-season development and late-season cooling on total bud growth in *Fagus*
***sylvatica* (experiment 1).** Effects of July (22 June to 23 July) and August (24 July to 25 August) moderate
and extreme cooling (8-13°C and 2-7°C, respectively) on total seasonal (4 July to bud set date) bud growth
for early- (grey) and late- (blue) leafing trees. Analyses show effect size means $\pm$ 95% confidence intervals
from linear mixed models, including treatment as fixed effect and bud-type (apical vs lateral) as random
effect. Early-leafing effects are calculated against the early-leafing control and late-leafing effects are
calculated against the late-leafing control.

572
573

574
Figure S5. Effects of early-season development and late-season cooling on relative bud growth in *Fagus*
***sylvatica* (experiment 1).** Effects of July (22 June to 23 July) and August (24 July to 25 August) moderate
and extreme cooling (8-13°C and 2-7°C, respectively) on relative seasonal (4 July to bud set date) bud
growth for early- (grey) and late- (blue) leafing trees. Analyses show effect size means ± 95% confidence
intervals from linear mixed models, including treatment as fixed effect and bud-type (apical vs lateral) as
random effect. Early-leafing effects are calculated against the early-leafing control and late-leafing effects
are calculated against the late-leafing control.
582

$$y = 16.4 + 0.00129\,x$$

$$R^2 < 0.01$$

583
584 **Figure S6. Relationship between end of season bud length and the timing of primary growth cessation**
585 **in *Fagus sylvatica* (experiment 1).** Effect of bud set DOY (day-of-year) on the final bud length at the end
586 of the growing season from a linear model with 95% confidence interval.

592 **Table S1. Mean and 95% confidence intervals of 50% leaf-out (day-of-year), bud set (day-of-year) and**
593 **absolute bud growth (cm) for each treatment in experiment 1.**

TREATMENT	LEAF-OUT DOY (95% CI)	BUD SET DOY (95% CI)	ABSOLUTE BUD GROWTH CM (95% CI)
1	**131.6 (130.2-133.)**	**260.5** (259.1-261.9)	**12.88** (11.96-13.80)
2	**148.1 (147.8-148.4)**	**264.1** (262.4-265.8)	**14.21** (13.53-14.89)
3	**131.2 (129.7-132.7)**	**261.9** (260.0-263.8)	**10.95** (10.29-11.61)
4	**148.6 (148.2-149.0)**	**268.9** (267.2-270.6)	**12.72** (11.99-13.45)
5	**130.1 (128.9-131.3)**	**263.1** (261.1-265.1)	**10.89** (10.41-11.37)
6	**148.7 (148.4-149.0)**	**268.1** (266.2-270.0)	**10.97** (10.35-11.59)
7	**130.2 (128.6-131.8)**	**256.0** (254.6-257.4)	**12.85** (12.17-13.53)
8	**148.9 (148.6-149.2)**	**259.7** (257.7-261.7)	**11.28** (10.63-11.93)
9	**132.3 (130.8-133.8)**	**258.6** (256.5-260.7)	**9.71** (9.22-10.20)
10	**148.0 (147.6-148.4)**	**262.2** (260.5-263.9)	**10.46** (9.81-11.11)

594
595
596
597

598 **Table S2. Mean and 95% confidence intervals of bud set (day-of-year) and absolute bud growth (cm) for**
599 **each treatment in experiment 2.**

TREATMENT	BUD SET DOY (95% CI)	ABSOLUTE BUD GROWTH CM (95% CI)
AMBIENT CONTROL	**257.2** (254.7-259.7)	**4.47** (3.80-5.14)
PRE-CHAMBER-CONTROL	**257.1** (254.1-260.1)	**4.73** (3.82-5.54)
PRE-FULL-DAY	**260.6** (257.3-263.9)	**5.18** (4.26-6.10)
PRE-DAY	**257.2** (252.7-261.7)	**4.95** (3.96-5.94)
PRE-NIGHT	**261.0** (257.9-264.1)	**5.01** (4.34-5.68)
POST-CHAMBER-CONTROL	**255.4 (251.8-259.0)**	**4.24 (3.45-5.03)**
POST-FULL-DAY	**251.6** (248.3-254.9)	**2.58** (2.08-3.08)
POST-DAY	**251.5** (248.0-255.0)	**2.92** (2.32-3.52)
POST-NIGHT	**260.6** (256.7-264.5)	**5.58** (4.70-6.46)

604
Figure S7. Bud set timing (day-of-year) for each control treatment in experiment 2. One-way ANOVA
revealed no significant differences in bud set timing between the three control groups ($F_{2,136}$ = 0.346, *p*
=0.708). Ambient control trees were placed outside throughout the experiment, Pre-Chamber control
trees were in a climate chamber under simulated ambient conditions between 22 May and 21 June, Post-
Chamber control trees were placed in a climate chamber under simulated ambient conditions between 22
June and 21 July.

613
Figure S8. Effects of pre- and post-solstice daytime, nighttime and full-day cooling on relative bud
growth in *Fagus sylvatica* (experiment 2). Pre-solstice (22 May to 21 June) and post-solstice (22 June to
21 July) cooling treatments were applied. Full-day cooling trees were continuously cooled to 8°C, day
cooling trees were cooled to 8°C in the day and kept at 20°C at night, night cooling trees were cooled to
8°C at night and kept at 20°C in the day. Analyses show effect size means ± 95% confidence intervals
from linear mixed models, including treatment as fixed effect and bud-type (apical vs lateral) as random
effect.

a

Pre-solstice

Legend:
- Normal plant function
- Reduced photosynthesis
- Arrested development
- Induced senescence

b

Post-solstice

Figure S10. Conceptual model of pre- and post-solstice effects of day and nighttime temperatures on
the autumn phenology of *Fagus sylvatica*. The colours of the shaded areas represent the effect of cooling
to different temperatures at specific times of the day. Above a certain threshold, plants function normally,
and cooling has little effect on the timing of primary growth cessation (green; here depicted as 10°C, n.b.
this is an intended simplification and does not necessarily accurately represent the reality). Light blue
areas show combinations that lead to arrested development, which delays primary growth cessation. Pale
green areas depict combinations that reduce photosynthesis and potentially growth but do not affect
phenology. Dark purple areas depict combinations that lead to the induction of senescence and
overwintering processes, therefore advancing primary growth cessation. **a)** Before the solstice,
lengthening days give temperate trees a reliable cue of the favourable conditions to come, so there is no
temperature that can induce senescence at this time of year (Kramer, 1936). Nighttime cooling (< ~10°C)
slows down beech's developmental processes such as meristematic activity, tissue expansion and
maturation, which leads to a growth and development debt (Zweifel *et al.*, 2021). As the days are
sufficiently warm, photosynthesis is unaffected. This means that the trees accumulate excess carbon that
can only be used later in the growing season to make up for this developmental debt, thus delaying

primary growth cessation. Daytime cooling (< ~10°C) slows development to a lesser extent, however, it also reduces carbon gain by inhibiting photosynthesis but not respiration. Due to carbon limitation, affected trees may prioritise development over growth, so buds may be shorter but there is no resulting delay to primary growth cessation. **b)** After the solstice and due to shortening days, trees become increasingly sensitive to cooling, especially during the day. Therefore, daytime temperatures < ~10°C are likely to induce overwintering responses and therefore advance autumn phenology. However, due to typical daily cycles of temperature always reaching their minimum during the night, trees are less susceptible to cooling in the night than in the day. At night, temperatures between ~4°C and ~10°C slow the trees' developmental processes, leading to delayed primary growth cessation.