Revisiting the habitat selection of a specialist forest bird,
Scolopax rusticola, during its wintering in the Mediterranean
region

Camille Beaumelle[a,*], Jessica Barbet[b], Aurélie Cuby[a], Marc Chautan[a], Fabrice

Etienne[c], Michel Martel[d], Alix Du Roure[a], Remy Chabanne[e], Estelle Lauer[f], Kévin Le

Rest[b]

[a] *Fédération régionale des chasseurs Auvergne-Rhône-Alpes, 42163 Andrézieux-*
Bouthéon, France
[b] *Office Français de la Biodiversité, 44300 Nantes, France*
[c] *Fédération départementale des chasseurs de l'Ardèche, 07200 Saint Etienne de*
Boulogne, France
[d] *Office national des forêts, 34172 Montpellier, France*
[e] *Office Français de la Biodiversité, 26400 Eurre, France*
[f] *Fédération départementale des chasseurs de l'Isère, 38610 Gières, France*
* Corresponding author : beaumelle.camille @gmail.com
17

Abstract

Habitat selection is a key mechanism that enables animals to optimize their fitness in
response to varying environmental conditions. Differences in habitat selection
between geographical areas may imply behavioral adaptations to local environmental
conditions. Understanding the potential of adaptation of species along broad
geographical ranges is of primary interest to anticipate potential changes in species
behavior or distribution range in the context of climate changes.

In this study, we investigated the habitat selection and the daily movement patterns of
the Eurasian Woodcock, *Scolopax rusticola*, a bird species that winters across widely
varying climatic zones. We tracked 89 individuals wintering in the Mediterranean
regions with GPS-VHF transmitters, where climate and habitats conditions differ
significantly from the oceanic regions. To assess how they responded to varying
habitat and environmental conditions, we collected data across four geographical
regions spanning a gradient of Mediterranean climatic influence — ranging from

northern subareas with denser forest and deeper soil to southern subareas characterized by less productive forests, garrigues, and rocky soil.

In wetter regions, woodcocks generally commuted between diurnal resting in forested areas and nocturnal feeding in open habitats, though less frequently than described in studies from other climatic regions. Under drier conditions, they often remained within forest habitats, displaying higher daytime activity and reduced nocturnal movement. Across all subareas, denser forests were selected during the day. The selection for nocturnal forest microhabitats varied by subarea, e.g., denser low vegetation was more strongly selected in southern subareas, while forest habitats highly covered by forest herbs were mostly avoided in northern subareas.

Our findings underscore the remarkable behavioral flexibility of woodcocks, highlighting their potential to adapt to global change. However, the occurrence of escape movements under the driest conditions suggest that this change in behavior and habitat selection may be an early warning sign of the effects of climate change on the wintering areas. Overall, our study emphasizes the need to study the ecology of species across diverse environmental conditions to better understand their habitat requirements and adaptive capacity.

Introduction

Global change is driving species to respond rapidly to new environmental conditions and altered biotic interactions (IPBES, 2019). While empirical niche models predicting the future species distribution and survival underscore the urgency of planning conservation and management strategies to protect biodiversity, they often fall short in capturing species' vulnerability or their potential for adaptation in response to rapid changes. Yet, understanding these processes is essential for developing effective conservation strategies (Bellard et al., 2012; Dawson et al., 2011).

Habitat selection — the disproportionate use of resources or conditions by an animal (Mayor et al., 2009) — is one of the behavioral mechanisms enabling species to cope with global changes and persist in their environment (e.g., Bailey et al., 2019). The study of habitat selection is crucial for understanding the distribution and abundance of species. It also helps to identify priority habitats that should be preserved or

managed to support the persistence of species in a region, providing practical and
effective devices for conservation strategies.

Habitat selection occurs at multiple scales, both temporally and spatially (Mayor et
al., 2009), sometimes across large and heterogeneous distribution areas that do not
necessarily provide optimal conditions for the species. These areas may also be
remote or difficult to access, further complicating data collection. Consequently,
habitat selection studies often rely on data from a small part of an animal's range and
cover only a limited period of the animal's lifespan. The insights gained from such
studies and the management recommendations derived from them are therefore
rarely transferable to other contexts within the species' range (Beyer et al., 2010) or
to other taxa (Yates et al., 2018). This underlines the need for broader, integrative
studies that take into account the spatiotemporal variability of habitat selection (Yates
et al., 2018).

Habitat selection is a complex process shaped by the interaction between habitat
availability and the behavioral motivations of individuals to meet their biological needs
(Beyer et al., 2010; Roever et al., 2014). For example, habitat selection may be
influenced by factors such as the physiology of the individual (e.g., age, Skórka et al.,
2016), access to specific resources (e.g., landfill availability for *Larus fuscus*, Langley
et al., 2021) and predation risk (Wasserlauf et al., 2023). This flexibility in habitat
selection is particularly critical for species occupying habitats that are spatially
heterogeneous or/and subject to temporal changes (e.g., climate and land use
changes), as optimal space occupancy decisions may vary according to these
variable conditions (Mabille et al., 2012; Merkle et al., 2022; Pearse et al., 2024;
Shahsavarzadeh et al., 2023; Verzuh et al., 2021).

Behavioral flexibility is one of the key mechanisms by which animal species can
swiftly adapt to a changing environment. However, it remains underexplored and
exhibits considerable taxonomic and geographic disparities (Beever et al., 2017;
Thurman et al., 2020). Research on behavioral adaptation has often been restricted
to limited areas and short time periods or focused on the responses to chronic stimuli
(e.g., interannual fluctuation in average temperature, Mazerolle et al., 2011), limiting
insights into species' capacity for flexible responses to more extreme environmental
changes, such as those involving climate variability (Beever et al., 2017).

Migratory birds often occupy a remarkable diversity of environmental niches
throughout their annual cycle, suggesting their behavioral flexibility in optimizing the

use of available resources. Despite this potential adaptability, they often exhibit
strong fidelity to their breeding, wintering and stopover sites, likely because such
consistency offers advantages such as more efficiency of movement and foraging, as
well as better predator avoidance (Piper, 2011). However, when these sites are
degraded, migratory birds may prioritize fidelity over relocation to alternative areas, a
behavior that can sometimes result in maladaptation (Merkle et al., 2022). This trade-
off suggests that species with the highest site fidelity may be particularly vulnerable
to the challenges posed by rapid global change (Chan et al., 2023; but see Blackburn
& Cresswell, 2016; Wellbrock et al., 2017). To cope with these changes, behavioral
compensation within existing habitats could be an important mechanism that enables
highly site-faithful species to adapt to local habitat alterations (Merkle et al., 2022).
For example, one potential form of behavioral compensation is the adjustment of
habitat selection within the occupied site, but few studies have investigated the ability
of species to employ these strategies in response to environmental change (Mallet et
al., 2023).
The aim of this study is to determine whether a migratory bird occupying a broad
geographical wintering range can adjust its habitat selection and behavior along a
latitudinal gradient. Specifically, we evaluate the variability in habitat selections of
Eurasian Woodcocks wintering in southern France. The ecology of the Eurasian
Woodcock has been extensively documented in oceanic and continental regions of
Europe (Arizaga et al., 2015; Braña et al., 2010, 2013; Duriez, Ferrand, et al., 2005;
Duriez, Fritz, Said, et al., 2005; Guzmán et al., 2017; Hirons & Johnson, 1987;
Hoodless & Hirons, 2007; Sládeček et al., 2023). Typically, Eurasian woodcocks
commute between open nocturnal foraging sites, where they feed mainly on
earthworms, and diurnal forest sectors, where they rest in shelter. This species also
winters in the Mediterranean region, which constitutes a significant part of its
wintering range (Gonçalves et al., 2019). However, in these regions, observations of
woodcocks in open areas at night are less frequent, suggesting a different behavior.
To investigate this, we used GPS-VHF devices to track the movements of 89
wintering woodcocks in four subareas with varying degrees of Mediterranean climate
influence and winter dryness.
First, we analyzed general movement patterns during wintering to assess woodcock
behavior in the different Mediterranean subareas. At the landscape scale (as defined
by Mayor et al., 2009), we examined whether woodcocks continued to select forest

habitats during the day and more open habitats at night. At the local scale (as defined
by Mayor et al., 2009), we analyzed the habitat structures within the forests selected
by woodcocks. Finally, we compared our findings with studies from other climatic
regions and discussed their broader implications for the conservation of other taxa in
the context of global change.

Material and methods

Study area

Four subareas were selected in distinct sectors of southern France (Figure 1), each
of which is characterized by vegetation adapted to dry summers and strong winds
(called "le mistral" and "la tramontane") but is influenced to varying degrees by the
Mediterranean climate (Table 1).

Figure 1: Location of the four subareas and the 14 298 recorded locations of 89
woodcocks in the study area located southern France. Subareas: S1: Central Drôme,
S2: Southern Drôme, S3: Ardèche gorges, S4: Hérault gorges. The background of
the map is the tree cover density, dark grey corresponding to high level of tree cover
density (land.copernicus.eu).

The two Mediterranean subareas of the central Drôme (S1 and S2) belong to the
ecoregion of the Plains and Hills of the Rhône Valley and Languedoc Region (see
Figure 1). These two subareas are relatively humid due to the influence of the

continental and mountain climate, have moderately deep soils, and the forests are
dominated by *Quercus pubescens* (a deciduous tree species). The Ardèche (S3) and
Hérault (S4) gorges, located in the southern part of the study area, belong to the
same forest ecoregion, the *garrigues* (or shrublands), as defined by the French
National Geographic Institute (Figure 1). Their soils are shallower and much rockier
than further north, so they drain better. The *garrigues* are characterized by low, slow-
growing vegetation. *Quercus ilex* (a sempervirens tree species) is the main tree
species. Others are *Pinus halepensis*, *Pinus pinea, Quercus coccifera*, *Erica* spp.,
Cistus spp. and *Arbutus unedo*.
The differences in soil characteristics and climates among subareas lead to different
agricultural activities, with orchards and crops predominating in the Drôme subareas,
while vineyards predominate in the southern subareas.
166

Capture and tracking

The study was conducted from winters 2019-20 to 2022-23. For each subarea,
Woodcocks were caught over two or three consecutive winters, from late December
to early January (Table 1), i.e., a period where the individuals showed high site fidelity
(Ferrand et al., 2008). The captures took place mostly at night in fields, vineyards or
orchards, where the birds were detected and immobilized using a spotlight (glared)
and then caught with a landing net (Gossmann et al., 1988). In the Hérault gorges,
half of the birds were caught at night near ponds using vertical nets (misnets) set up
before dusk, as woodcocks are sometimes observed visiting these ponds at the
beginning of the night to groom themselves. The woodcocks were ringed and aged
by plumage characteristics (Clausager, 1973). To determine sex, a feather was
collected and analyzed by the CIBIO/CTM laboratory in Portugal (DAN analyses,
Griffiths et al., 1998). If the individual weighed more than 300 g and was in good
physical and health condition, it was equipped with a VHF-GPS tag (PinPoint VHF
240, Lotek Ltd, ~8 g) using a leg-loop harness made with an elastic rubber cord (1.5
mm EPDM) lined within a silicone sheaths (IN 2.0; OUT 2.6 mm) to limit friction and
weighting in total ~ 2 g. Tag plus harness thus weight less than 3% of bird's body
mass. The elastic rubber cord sometimes broke too early, few weeks after bird's
release, so we use only the silicone tube in 2022-23 in S4 (harness being event

lighter, ~1g). Many of these birds kept the tag for more than one year (half of them
being detected with VHF next winter).

Each bird was first located by triangulation and the data (GPS positions) were
downloaded every two weeks, from approximately 50 meters to avoid any
disturbance (depending on vegetation and topography). The tags were programmed
to record four locations every 6 hours, comprising two nocturnal locations (9 pm and
3 am) and two diurnal locations (9 am and 3 pm). In addition to the GPS data,
temperature, elevation, and GPS quality metrics (HDPO, eRes, number of satellites
used for the location) were recorded. Some tags also recorded activity using
accelerometer values (ODBA, Wilson et al., 2006) but this data is not available for all
individuals and are thus not considered in this paper. The activity levels and daily
distances calculated below (see section Statistical analysis of movement patterns)
are however highly correlated.

During the study period, we tagged 97 woodcocks. However, one bird was killed by a
hunter a few days after being equipped, another was predated, and for six of them
we could not successfully download the GPS data (the birds may have flown out of
the study area or been predated, or the tags were defective). Therefore, in this study
we analyzed the movements of 89 woodcocks (Table 1).

Table 1: Descriptive data on the four subareas and sample size of woodcocks. N:
number of woodcocks fitted with GPS, n: number of GPS locations per woodcock.
Temperature and soil wetness were computed based on the daily Safran model
(Météo-France) data during the tracking period (Begin date within the subarea to end
February). The area of subareas was calculated using the 100% minimum convex
hull of woodcock locations.

Subarea	Area (km²)	Year	N	Begin date	N Mean [min-max]	Temperature Mean [min-max]	Soil wetness Mean % [min-max]
(S1)	29	2019-2020	8	21st Dec.	187 [61; 261]	7 [-2; 19]	80 [67; 97]
		2020-2021	6	9th Jan.	126 [59; 170]	7 [-5; 18]	80 [72; 83]
		2021-2022	5	25th Dec.	162 [115; 234]	6 [-5; 18]	76 [66; 85]
(S2)	75	2020-2021	4	8th Jan.	172 [161; 187]	6 [-4; 17]	87 [78; 93]
		2021-2022	1	29th Dec.	175	5 [-3; 17]	75 [65; 84]

(S3)	150	2019-2020	7	21st Dec.	168 [62; 273]	7 [-3; 21]	92 [76; 116]
		2020-2021	10	7th Jan.	176 [144; 183]	6 [-5; 17]	91 [84; 97]
		2021-2022	10	29th Dec.	143 [40; 224]	6 [-5; 18]	75 [63; 90]
(S4)	118	2021-2022	20	16th Dec.	141 [42; 207]	8 [-3; 18]	59 [50; 67]
		2022-2023	18	21st Dec.	182 [62; 257]	8 [-4; 20]	45 [40; 51]

Subareas: S1: Central Drôme, S2: Southern Drôme, S3: Ardèche gorges, S4: Hérault gorges

GPS data filtering

Following the manufacturer's instructions, we excluded locations recorded with fewer

than four satellites, an HDPO≥10 or an eRes≤0 or ≥0. For isolated locations (i.e.,

distance >120 meters from other locations, which corresponds to the 75% quartile of

nocturnal distance covered by woodcocks with a unique nocturnal feeding place in

Duriez, Fritz, Said, et al., (2005), we increased the threshold for satellite to six

because woodcock is unexpected to move significant distances during daytime or

nighttime without being disturbed. This removed 28 isolated points with doubts about

their accuracy, which represented less than 0.2% of the data). Additionally, locations

with a significant difference between the recorded elevation and the ground elevation

were removed because this may indicate a poor GPS reading (Sládeček et al., 2023).

The ground elevation data were estimated using the *elevatr* R package with a

resolution of 54 meters (Hollister et al., 2023). We set the threshold at 250 m to

account for the large elevation differences in our study area.

Some tags fall off the bird due to harness breakage (those with elastic EPDM 1.5

mm), and some birds have been predated (23, but it is not always possible to tell

whether it is predation or harness breakage). To account for cases of detachment or

predation, the temperature recorded by the tag dropped suddenly, allowing us to

remove all non-informative data. Similarly, we excluded the locations for the birds

having been killed by hunters (one bird). Due to these exclusions, two individuals had

less than 10 days of tracking data, so we decided to exclude them.

To minimize the influence of capture stress and transmitter attachment on the results,

we systematically removed locations recorded within 48 hours of capture from all

individuals. However, data collected during the first days of monitoring showed that

individuals appeared to recover their normal behavior at varying rates. As a

compromise between using a maximum number of data and excluding data that were

influenced by capture, we examined the behavior of each bird during the first week

after capture. We removed locations until each bird had stabilized its wintering

behavior, i.e., found its first diurnal sector and established a routine commuting flight

pattern or not (Duriez, Fritz, Said, et al., 2005). Thus, the final dataset included data

from two days after capture for 59 woodcocks, three days for 16 birds, and four to

eight days for 14 individuals for which it was more difficult to detect a routine in

behavior. All filtering steps and the corresponding sample size are provided in

Appendix 1, Figure A1.

In addition, eight woodcocks in the Hérault gorges escaped during the driest period

of the study and later returned to their primary wintering area. Data from these

escape locations (18 to 74 km away) and points between the primary wintering area

and the escape area were excluded from the analysis, resulting in the removal of 413

locations (24% of the total from these eight individuals).

Our study focused on the wintering habitat, so we excluded locations recorded in

March and April because woodcocks may change their habits to prepare for migration

(Duriez, Fritz, Said, et al., 2005; Tedeschi et al., 2020).

Statistical analysis of movement patterns

To analyze the global movement of woodcock, we computed diurnal and nocturnal

distances as well as the distances between diurnal and nocturnal sites (commuting

flights). All distances were calculated between pairs of consecutive locations using

the "st_distance" function of the *sf* package (Pebesma et al., 2024). Commuting

distances were estimated by measuring the distances between consecutive nocturnal

and diurnal locations (see Duriez, Fritz, Said, et al., 2005). These include the

distances of evening commute flights (occurring between 3 pm and 9 pm) and

morning commute flights (occurring between 3 am and 9 am). Nocturnal and diurnal

movements were estimated by measuring the distances between consecutive

nocturnal locations (9 pm and 3 am) and between consecutive diurnal locations (9

am and 3 pm), respectively. We tested whether distances differed between

woodcocks of different age and sex, and whether they varied by subareas and

dryness of winter. Winter dryness was assessed using the mean soil wetness index

(from the daily Safran-Isba model, Météo-France) as an indicator of food availability,

as soil that is too dry may prevent woodcocks from probing the soil to feed on

earthworms (Duriez, Ferrand, et al., 2005; Péron et al., 2011). We computed the

mean soil wetness during the tracking period in each subarea for each year and
assigned the same value to all woodcocks in the same winter and subarea (Table 1).
To identify the main drivers of each pairwise distance, the distances were log-
transformed, and since they were normally distributed, we fitted a linear mixed model
with the four explanatory variables and the interaction between the subareas and the
soil wetness using the *lmerTest* package (Kuznetsova et al., 2017). Confidence
intervals were calculated using the *profile* method of the *confint* function of the *lme4*
package (Bates et al., 2015). The p-values were produced by the Satterthwaite
approximation. The soil wetness variable was scaled. We included individuals as a
random intercept to account for repeated measures. The three models were fitted
using the following formula, with subarea being a factor with four levels (S1, S2, S3,
S4) and *SWI* being the mean soil witness index.
285

$$\log(distance) \sim age + sex + subarea + SWI + subarea\!:\!SWI + 1|individual$$

Statistical analysis of habitat selection at landscape scale
To analyze the habitat selection of woodcock at the landscape scale, we determined
whether woodcock locations were situated in forested or open habitats, using a high-
resolution tree cover density (TCD) layer (10m raster from copernicus.eu). The TCD
for each location was calculated as the mean value of raster cells within a 30 m
radius buffer to account for the accuracy of GPS locations (estimated from data
collected by transmitters left in the forest).
To measure the degree of TCD selection by woodcock, we fitted a resource selection
function using logistic regression to model use-availability data (Boyce et al., 2002;
Fieberg et al., 2021; Northrup et al., 2022). Habitats used by woodcock that were
confirmed by GPS locations, were classified as 1, while randomly sampled available
habitats were classified as 0 (hereafter named pseudo-absence). Due to the large
variation in the number of locations available per woodcock (min= 40; max=273),
there was a risk that the model would be more influenced by woodcocks with a higher
number of locations. To limit this bias, we added individuals as random intercepts
(Gillies et al., 2006). In our study, several woodcocks frequently commuted between
areas more than 1 km apart, a behavior that was also observed in other wintering
areas (e.g., in Brittany: Duriez, Fritz, Said, et al., 2005; in Spain: Guzmán et al.,

2017) and some reach 5 km. Therefore, for each woodcock, the available habitat was
defined as the area within a buffer of 1 km, 3 km or 5 km around all locations and we
compared the results according to the size of the buffer considered. For each true
location, 100 pseudo-absences were sampled, excluding areas never used by
woodcocks, i.e., water bodies or artificial surfaces determined by the Corine Land
Cover layer (100m raster from copernicus.eu). Predictor variables included the
scaled TCD, a DayNight factor, the four subareas, the scaled soil wetness (Table 1),
the sex and the age. The day/night factor for the pseudo-absence data was defined
from the nearest available GPS location from each woodcock. The model was fitted
using the following formula:

$$used \sim TCD{:}DayNight + TCD{:}DayNight{:}(age + sex + subarea + SWI)$$
$$+ TCD{:}DayNight{:}subarea{:}SWI + 1|individual$$

The response variable *used* take on a value of 0 or 1, DayNight factor have two
levels, subarea is a factor of four levels (S1, S2, S3, S4) and *SWI* is the mean soil
witness index. We repeated the sampling of the 100 random pseudo-absence
locations 10 times for each buffered area (1, 3 and 5 km) to assess the sensitivity of
the results to the sampling of these random locations which represented the available
habitat. The 10 models were averaged using the default *model.avg* function of the
MuMIn R package (Bartoń, 2024).

Statistical analysis of forest habitat selection at local scale
To determine the environmental characteristics of habitat selected by woodcock at
the local scale, we randomly visited six plots of 25 m x 25 m per woodcock fitted with
GPS. Of the six plots, three contained nocturnal locations and three contained diurnal
locations. To capture a gradient of selection, for both nocturnal and diurnal plots, we
selected one plot containing, before data filtering, less than 3% of raw data locations
(low use), one plot containing between 3% and 6% of raw locations (moderate use),
and one plot with the maximum number of raw data locations (high use). To avoid
disturbing the tracked woodcocks, plots were visited in the winter of the following
year, assuming that vegetation would not change significantly within a year, given the
slow growth of Mediterranean vegetation. The plot having undergone an obvious

change in vegetation (e.g., local fire or human clearing) were not monitored. In total,
we assessed the environmental characteristics of 403 plots in forest habitats.

In the forests, we described the main tree species, the dominant tree height, the
proportion of vegetation cover >0.5 meters, the accessibility of the soil (which could
be completely covered by rocks), the proportion and distribution of grass and rocks,
and the vegetation density at 10 cm, 50 cm, 100 cm and 150 cm (Appendix 1, Figure
A2). Vegetation density was determined as the probability of not seeing a mark (a 5
cm long piece of orange tape) at the corresponding height on a 5 cm diameter stick
placed approximately in the center of the plot. Observations were made from a
distance of 10 meters in eight cardinal directions, with the observer's eyes positioned
at the same height as the mark (Appendix 1, Figure A3).

To analyze forest habitat selection, we used a model selection approach to test the
selection of environmental variables (Table 2) and their interaction with age, sex, soil
wetness, day/night, and the subareas. To limit the number of parameters, we
transformed the subarea factor variable into a continuous variable coded 1 (S1), 2
(S2), 3 (S3) and 4 (S4), reflecting the degree of Mediterranean climate influence. We
quantified the intensity of plot selection (where environmental variables were
described) by summing the number of nocturnal and diurnal locations, respectively, of
each woodcock in the corresponding sampled plots and during the studied period
(Table 1). An offset in the model controlled the different number of locations per
individual (see below). After controlling environmental variables and correlations, we
retained six environmental variables (Table 2) as explanatory variables for the model.
Due to the strong structuring of the proportion and distribution of forest herbs among
the four subareas (Appendix 1, Figure A2), we choose to group these two variables
into three classes: forest herbs with scattered distribution, connected patches of
forest herbs covering ≥50% or <50% of the plot. Concerning rocks, we choose to
retain only the proportion of rocks to describe this environmental variable because
there was a redundance of information between the proportion and distribution of
rocks, but no evident pattern (Appendix 1, Figure A2). Due to high correlation (r>0.7,
Spearman's test) between vegetation density of different height and soil wetness, we
kept only the vegetation density of 50 cm and 150 cm height (Table 2). After this initial
variable selection, only soil wetness and the degree of Mediterranean climate
influence were highly correlated (r=-0.8, Spearman's test, other variables correlations
were below 0.47), so these variables were not used in the same models.

Table 2: Description of the six retained environmental variables measured in plots
located in forest.

Variables	Description
Dominant height of tree	Height <4 meters or >4 meters
Cover	Proportion of the ground covered by vegetation of >0.5 meters
Vegetation density 150 cm	Probability of not seeing the mark at 150cm
Vegetation density 50 cm	Probability of not seeing the mark at 50cm
Forest herb distribution	Scattered, connected covering <50% of the ground, connected covering ≥50% of the ground
Proportion of rocks	Proportion of the ground covered by rocks outcrops and loose rocks

Given the number of variables and sample size, we limited the number of competing
models to seven in the model selection procedure to avoid overfitting. These included
the null model, the model with environmental variables only, the model with
environmental variables interacted with the DayNight factor, and four models in which
the environmental variables also interacted with the DayNight factor, age, sex, soil
wetness or the influence of the Mediterranean gradient. To account for overdispersion
in the count data, we used a negative binomial family. Due to the varying numbers of
locations among woodcocks, we included an offset in the models corresponding to
the total number of forest locations for each woodcock (log-transformed). During the
day, woodcocks were always in forest; therefore, the offset for diurnal locations
corresponded to the total number of diurnal locations. At night, woodcocks could use
either forest or open areas. Nocturnal locations where the mean TCD exceeded 5%
within a 12.5 m radius buffer (to match the 25 m x 25 m plot resolution) were
considered forest habitat, which includes clear forests typical of Mediterranean
landscapes. Models were ranked based on their AICc values, and we selected the
model with the lowest AICc value. Models with ΔAICc <2 were considered equivalent.
In this case, we selected the most parsimonious model, i.e., the model with the
fewest degrees of freedom (Burnham & Anderson, 2004).

We used R 4.4.1 (R Core Team, 2024) for data processing and statistics, and QGIS
3.28.1 (QGIS.org, 2024) and Inscape 1.3.2 (Inkscape Project, 2023) for data
visualization.

Results

In this study we analyzed the movements of 89 woodcocks, including 23 1[st] winter
male, 24 1[st] winter female, 25 adult males and 17 adult females. From the start of the
tracking record (Table 1) to the end of February, there were a total of 14 344 locations
with an average of 161 locations per woodcock (min= 40; max=273).

Movement patterns

The movement patterns of woodcocks varied greatly depending on the individual and
subarea (Table 3, Appendix 1, Figure A4). The sex and age of individuals were not
significant factors in explaining the distances between locations (Table 4).

<u>Table 3: Summary of commute, diurnal and nocturnal distances (meters) in each</u>
<u>subarea</u>. The mean, the standard deviation (sd) and the maximum (max) of each
type of distance were reported. N: number of individuals. n: number of location pairs
to calculate the distances

	S1 (N=19)			S2 (N=5)			S3 (N=27)			S4 (N=38)		
	mean	sd	max	**mean**	sd	max	**mean**	Sd	max	**Mean**	sd	max
Commute (n=6223)	**1087**	1329	5345	**909**	1553	6264	**919**	1148	12230	**358**	578	13796
Diurnal (n=2695)	**32**	43	645	**32**	34	258	**40**	33	327	**74**	57	418
Nocturnal (n=3709)	**182**	592	5284	**251**	832	5879	**178**	633	12237	**76**	206	3511

Subareas: S1: Central Drôme, S2: Southern Drôme, S3: Ardèche gorges, S4: Hérault gorges

woodcocks in S4 commuted less frequently between distinct diurnal and nocturnal
sectors than woodcocks in other subareas, and commuting distances tended to be
shorter (Table 3, Table 4). For instance, the 75% quartile of commuting distances in
S4 was <450m, whereas commuting distances in the other subareas almost always
exceeded this value (75% quartile of commuting distances in S1 = 1 608 m, S2 = 763
m, S3 = 1383 m). They were also more active during the day, as shown by the diurnal

distances, which were twice as high as those of woodcocks in the other three

subareas (Table 3, Table 4).

Woodcocks in S4 were less active at night than woodcocks in S1 (Table 3, Table 4).

In S3, woodcocks were also significantly less active than woodcocks in S1 (Table 4),

albeit to a much lesser extent (Table 3) and especially during the dry winter

(Appendix 1, Figure A4). This pattern was indeed slightly more pronounced with

decreasing soil wetness, although it was not significant as *p-values* were below 0.1

but exceeded the 0.05 threshold (Table 4). For example, woodcocks in S3 covered

137 m [94, 180]$_{CI95\%}$ between consecutive locations at night in the drier winter (2021-

2022, Table 1), whereas they covered 169 m [118; 219]$_{CI95\%}$ and 215 m [141;

290]$_{CI95\%}$ in the wetter winters of 2019-2020 and 2020-2021 respectively.

Table 4: Estimates and 95% confidence intervals of the linear mixt models explaining

commute, diurnal and nocturnal distances. Females and adults were the references

levels for the sex and age factors. The subarea S1 was the reference factor level.

SWI: soil wetness index.

	Commute			Diurnal			Nocturnal		
	Estimate	CI95%	P	Estimate	CI95%	p	Estimate	CI95%	p
Sex (Male)	0.15	[-0.20; 0.50]	0.420	0.03	[-0.10; 0.16]	0.675	0.00	[-0.18; 0.19]	0.980
Age (1st winter)	-0.17	[-0.54; 0.19]	0.382	0.05	[-0.08; 0.18]	0.498	-0.11	[-0.30; 0.08]	0.280
Intercept (S1)	6.67	[-4.69; 8.66]	<0.001	3.04	[2.26; 3.81]	<0.001	4.11	[3.19; 5.03]	<0.001
Subarea S2	-1.12	[-4.21; 1.96]	0.499	-0.48	[-1.64; 0.68]	0.448	-0.27	[-1.72; 1.18]	0.732
Subarea S3	-1.23	[-3.31; 0.86]	0.275	0.55	[-0.26; 1.37]	0.209	-1.19	[-2.16; -0.21]	0.025
Subarea S4	-1.87	[-3.95; 0.21]	0.097	1.07	[0.26; 1.87]	0.016	-1.16	[-2.16; -0.18]	0.030
SWI (S1)	-1.48	[-5.05; 2.10]	0.443	0.09	[-1.21; 1.37]	0.906	-1.62	[-3.46; 0.23]	0.106
Subarea S2:SWI	1.74	[-2.63; 6.10]	0.461	0.39	[-1.17; 1.95]	0.648	1.27	[-0.98; 3.51]	0.295
Subarea S3:SWI	1.89	[-1.74; 5.52]	0.333	-0.34	[-1.65; 0.97]	0.628	1.73	[-0.16; 3.60]	0.090
Subarea S4:SWI	1.11	[-2.52; 4.74]	0.570	0.05	[-1.26; 1.36]	0.942	1.80	[-0.08; 3.68]	0.077

434 Subareas: S1: Central Drôme, S2: Southern Drôme, S3: Ardèche gorges, S4: Hérault gorges

Habitat selection at the landscape scale

During the day, woodcocks almost always stayed in tree-covered area (Figure 2).

Only 1% of diurnal locations were recorded in open areas (TCD<5%), which included

garrigues with sparse vegetation, hedgerows, and orchards. The selection of

nocturnal habitat at the landscape scale varied between individuals. There were 20

individuals that frequently used open habitats at night, with more than 50% of their

nocturnal locations in non-tree-covered areas. In contrast, 69 individuals used less

than 50% open areas, including 27 woodcocks that had less than 5% of their

locations in non-tree-covered areas. Nevertheless, woodcocks that remained in the

forest at night generally stayed in areas with lower TCD at night than during the day
(Appendix 1, Figure A5).

448
<u>Figure 2: Frequency of open areas and forest in used habitats in comparison with the</u>
<u>available habitats, computed with a buffer size of 1 km,3 km and 5 km around used</u>
<u>bird's locations</u>. The area was considered as forest if the tree cover density (TCD)
was ≥ 5%. Each bar displays the mean habitat composition observed for all
individuals pooled within the same subarea during nighttime and daytime. Subareas:
S1: Central Drôme, S2: Southern Drôme, S3: Ardèche gorges, S4: Hérault gorges.

Parameter estimates were generally consistent regardless of the size of the buffer
used to sample the pseudo-absence around the bird's location (Figure 3, Appendix 2
Table A1).
During the day, woodcocks significantly selected habitats with a higher tree cover
density (TCD) than available in the surrounding area. This selection was slightly more
pronounced in males and in wetter conditions in S1 and S2. In contrast, in S3 and
S4, woodcocks showed a stronger preference for higher TCD under drier conditions
than under wetter conditions (Figure 3). During the drier winter in S3 (2021-2022), the
mean TCD of pseudo-absence during daytime within the 1-km, 3-km and 5-km
buffers was 57%, 55% and 50%, respectively, while the mean TCD of woodcock
locations during daytime was 66% (i.e., 16 to 32% higher). In contrast, during the
wetter winter in S3 (2019-2020), the mean TCD of pseudo-absence during the day
within 1-km, 3-km and 5-km buffers was 58%, 55% and 49%, respectively, and
woodcocks selected diurnal habitats with a mean TCD of 57% (i.e., variation of -2%
to +16%).
At night, woodcocks selected habitats with a lower TCD (considering available
habitats within the 1-km, 3-km and 5-km buffers), except in S2 and in drier winters.
Under drier soil conditions, selection for higher TCD was significant in S3 and S4
(Figure 3). 1[st] winter birds showed slightly stronger selection for lower TCD than

adults (Figure 3), although 1st winter birds were not more likely to commute in open
areas at night than adults, i.e., the proportion of nocturnal location in open areas was
similar for 1st winter birds and adults (1st winter birds = 28%, adults = 26%). Indeed,
adult and 1st winter birds used similar TCD, but TCD in available habitats was lower
for adults than for 1st winter birds. For example, in S1, 1st winters birds and adults
used habitats at night with a mean TCD of 29% and 26%, respectively, while the
mean TCD of pseudo-absence within the 1-km, 3-km and 5-km buffers was 41%,
43% and 43%, and 30%, 36% and 39%, respectively.

<u>Figure 3: Odd ratios of the tree cover density (TCD) at the landscape scale obtained</u>
<u>from the model averaging. The available habitat was computed with a buffer size of 1</u>
<u>km, 3 km and 5 km.</u> The abscise axis was set as square root scale. Females and
adults were the references levels for the sex and age factors. TCD: tree cover
density, Subareas: S1 (reference factor): Central Drôme, S2: Southern Drôme, S3:
Ardèche gorges, S4: Hérault gorges.

Forest habitat selection at the local scale
Selection for specific environmental characteristics within the forest habitat was low,
but significant in the selected model (Appendix 2, Table A2) for several environmental
characteristics, especially during the night (Figure 4, Figure 5, Appendix 2, Table A3).
Within the forest, woodcocks preferred more covered plots during the day, while at
night they selected less covered areas and a higher proportion of rocks (Figure 4,

Figure 5). Visited plots were mainly covered with dispersed forest herbs in all
subareas, and plots with ≥ 50% forest herb cover were avoided at night, especially in
subareas with low degree of Mediterranean climate influence (Figure 4, Figure 5).

The dominant height of trees differed significantly between S1/S2 and S3/S4, as
trees in the south are rarely >4m, while they are well above 4 m in the north (Figure
5). The selection for a dominant height of trees >4m increased with the degree of
Mediterranean climate influence (Figure 4). This pattern could be explained by
specific forest structures or an atypical behavior of a few individuals. Indeed, forests
<4m are represented on <10% of the plots in the northern subarea (S1), but the two
plots frequently used during the night with a dominant height of trees <4m are highly
used, i.e., >20% of the nocturnal locations in the forest, by these two different
woodcocks. In the southern subarea (S4), the forests rarely reached a height of 4 m
(80% of the plots were <4m), yet three woodcocks very frequently used plots with a
dominant height of trees >4m, i.e., >20% of the nocturnal locations in the forest.

Woodcocks mainly visited plots that were moderately occupied by shrubs and low
branches (mean vegetation density at 50 cm: 0.41 $[0.12; 0.62]_{75\%IQR}$, mean
vegetation density at 150 cm: 0.37 $[0.12; 0.50]_{75\%IQR}$, Figure 3). With increasing
degree of Mediterranean climate influence, woodcocks preferred significantly higher
vegetation densities at 50 cm and 150 cm (Figure 4).

<u>Figure 4: Odd ratios of the environmental characteristics at the local scale obtained</u>
<u>from the selected model.</u>

<u>Figure 5: Graphical interpretation of local habitats rarely or frequently used by</u>
<u>woodcocks</u>. "Rarely used" habitats refer to plots used ≤2 times. The habitat
frequently used are plots with at least 5% of diurnal or nocturnal locations of each
woodcock. The values used for graphical interpretation are available in Appendix 2,
Table A4. Subareas: S1: Central Drôme, S2: Southern Drôme, S3: Ardèche gorges,
S4: Hérault gorges.

Discussion

Tracking of wintering woodcocks in the southern part of their geographic range
revealed different behaviors and diverse habitat selection strategies that are in
marked contrast to observations from other parts of their wintering range. Our results
suggest that woodcocks have adjusted their behavior to varying environmental
conditions along the Mediterranean climate gradient. This emphasizes the
importance of studying species across diverse habitats and under variable climatic
conditions to get a complete overview of species-environment relationships.

In the northern parts of the study area, woodcocks frequently commuted between
open and forested areas, which is consistent with the results of other studies (along
the Atlantic coast : Duriez et al., 2005; Hoodless & Hirons, 2007; northern Spain :
Braña et al., 2010), resulting in large differences in the type of habitat selected
between day and night. In contrast, woodcocks in southern regions tended to remain
in forested areas both day and night. However, when they were in forested areas at
night (77% of the nocturnal locations in S3 and S4), they selected less covered
environments, similar to woodcocks in Central European forests tracked during the
breeding period (Sládeček et al., 2023), and with low grass cover. It is unlikely that
this difference in forest habitat selection between day and night was caused by prey
availability (Duriez, Ferrand, et al., 2005). Indeed, there is no evidence that habitats
with lower forest cover provided more prey than those with high forest cover.
Moreover, woodcocks selected higher rock cover at night where stony habitats are
available (mainly in S3 and S4), which restricts woodcocks from probing for
earthworms, their principal prey (Granval, 1986), or alternative prey (Aradis et al.,
2019). Instead, nocturnal forest habitats were probably selected for protection from
predators (Braña et al., 2010). Adequate overhead tree and shrub cover may shield

woodcocks from nocturnal avian predators, e.g., the Eurasian eagle-owl, *Bubo bubo*

(Obuch, 2024), while relatively low tree cover allows for easier takeoff in response to

ground predators. Although the selection of specific microhabitats is too small to

provide precise recommendations for the management of the forest

microenvironment, our study reveals some patterns. Avoidance of microhabitats that

are heavily covered with continuous patches of forest herb at night could be due to

the lower visibility, which makes it difficult for birds to detect approaching ground

predators. The selection of habitats with higher vegetation density in the nocturnal

areas of the southern subareas could indicate that the birds prioritize shelter over

terrain traversability at night, which is associated with reduced feeding activity as

shown by movement patterns.

Habitat selection and movement patterns were generally consistent between males

and females, as well as between adults and 1[st] winter birds, except for the selection

of tree cover density at the landscape scale. During the day, males showed a

significantly stronger preference for areas with denser tree cover compared to

females. At night, 1[st] winter birds selected areas with lower tree cover density than

adults. 1[st] winter birds might be compelled to use less favorable wintering areas (e.g.,

areas with higher predation risk) due to competition or inexperience (Skórka et al.,

2016), while the difference between the sexes might be due to small differences in

the diet (Aradis et al., 2019). While wintering, there are few dimorphisms or

differences in energy requirements between sexes and age classes (Duriez et al.,

2004), so few differences in physiological requirements are expected. The sample

size for the habitat selection model at the local scale was probably insufficient to

detect such small differences.

Although open areas were available within their home range, woodcocks rarely

utilized them at night, especially in the three southernmost subareas and in drier

winters. The proportion of nocturnal locations in open areas was markedly lower (S1=

53%, S2= 29%, S3 = 40%, S4= 11%) than in observations in oceanic climates in

winter, where woodcocks are more frequently found in open areas at night (e.g., in

England: 94% (Hoodless & Hirons, 2007), and in Brittany, France: 85% (Duriez, Fritz,

Said, et al., 2005). Additionally, the analysis of woodcock movements suggested

higher diurnal activity, especially in the warmest subarea, the Hérault gorges. This

aligns with the hypothesis proposed by Duriez, Fritz, Binet, et al. (2005) and with the

observations of Hoodless & Hirons (2007), which suggest that woodcocks shift to

diurnal activity under milder climatic conditions. First, in milder climates, the costs of thermoregulation are lower, potentially reducing the birds' overall food requirements and decreasing the need to forage in riskier open habitats at night (Duriez, Fritz, Binet, et al., 2005). Measuring the activity rate of woodcocks using accelerometry, together with recording their location, could provide valuable data to analyze both their energy expenditure and the timing and locations of their feeding under varying climatic conditions (Patterson et al., 2019; Wilson et al., 2006). The activity rate of some woodcocks was recorded during this study and confirmed that the birds in S4 were active during the day. However, we chose not to present this data as it only concerns a subset of the total number of woodcocks tagged and would require additional methods and results. Furthermore, forests may provide shelter from strong winds, which are common in Mediterranean regions and can impose significant thermoregulation costs (Duriez et al., 2004). Second, prey species in southern subareas differ from those in northern areas (Aradis et al., 2019; Granval, 1986), and food resources are not necessarily more abundant in open area, reducing the incentive for woodcocks to leave the forest. To confirm the effects of prey diversity and distribution on woodcock behavior, it would be interesting to study both the soil macrofauna in the habitats used and to refine our knowledge of woodcock diet in Mediterranean regions (Aradis et al., 2019; Granval, 1986). Assessing diet and prey availability for woodcocks is labor-intensive and requires specialists who can identify taxa at taxonomic level with sufficient resolution from different species groups. DNA-based methods could help with species identification, but are not able to provide quantitative information (Gongalsky, 2021; Hoenig et al., 2022; Smith et al., 2008). We collected data to compare food availability based on earthworm abundance (Duriez, Ferrand, et al., 2005; Granval, 1986) at random sites and at sites used by some of the GPS transmitter equipped woodcocks. Variability in earthworm abundance was high at both the utilized and randomly selected sites (results not presented in this article), so it was difficult to detect a significant difference with the protocol used. Future protocols in Mediterranean regions should include the identification of additional prey species (e.g., Coleoptera and Diplopoda, Aradis et al., 2019) to achieve greater concordance with woodcock prey in Mediterranean regions. In addition, a significant number of habitats should be included in the food resource study to account for differences in the habitat selection strategy of woodcocks wintering in different climatic regions.

The winter of 2021-2022 was particularly dry, which affected soil moisture and
possibly prey availability, which could explain the observed differences in habitat
selection across winters. Specifically, woodcocks in the two southern subareas
selected areas with higher than usual cover density both during the day and at night
when soil wetness decreased. They also appeared to change their diel activity
pattern, although additional analyses of activity rates and associated behavior are
needed to confirm this change. Forest habitats can help to maintain soil moisture
during drier periods in Mediterranean regions (Sardans & Peñuelas, 2013), ensuring
an adequate food supply in forests. Furthermore, we recorded eight woodcocks
relocating from the southern subarea during the winter, while woodcocks are known
to be very faithful to their wintering areas throughout the winter (Ferrand et al., 2008),
suggesting that this wintering area did not provide anymore the necessary conditions
to meet the bird's needs, probably due to a lack of food availability. In addition, we
observed that the proportion of 1[st] winter birds among the birds hunted or trapped
was much lower in dry winters in the Southern subarea (*Réseau Bécasse - Lettre*
d'information n° 27, 2018), suggesting that inexperienced 1[st] winter birds tend to
avoid these areas when dry, while adult birds are more likely to return and remain
even in such less suitable environmental conditions. One of the eight individuals
relocated at the end of winter was recaptured and was in poor body condition (260 g
compared to 320 g when first captured). Also, the fact that most of the tagged birds in
this southern subarea (30/38 woodcocks) preferred to stay in a known location with
poor food availability rather than relocate suggests that relocation comes at a
significant cost compared to maintaining site fidelity. This unusual behavior of
relocating during winter could serve as an early warning signal of declining habitat
suitability for wintering woodcocks in Mediterranean regions. Further research is
needed to identify the factors that trigger these relocations. We hypothesize that food
availability is the most important factor (Duriez, Ferrand, et al., 2005; Martín et al.,
2023). In addition, the relocation of wintering woodcocks may lead to a change in the
local population density, which will imply a revision of population management, e.g.,
identification of relocation areas and local reduction of the number of birds that can
be legally harvested during hunting.

The differences in movement and habitat selection strategies between the four
Mediterranean subareas probably reflect differences in climate influence, structure
and distribution of forests and open habitats, all of which influence the diversity and

density of prey and predators. Forests in the southernmost subareas were used for
both shelter and foraging, whereas woodcocks in the northern subareas segregated
feeding in open areas and resting in forests. Consequently, habitat management
recommendations should be tailored to each subarea. For example, in the central
and south Drôme (S1 and S2), maintaining open habitats with high earthworm
densities could be beneficial, whereas in the Ardèche and the Hérault gorges, a
mosaic of dense and sparse forests could be more advantageous than maintaining
open areas. Overall, our results reaffirm the importance of preserving heterogeneous
habitats that provide multiple options for species to persist locally under varying
conditions (Oliver et al., 2010; Suggitt et al., 2018). Even within a single climatic
region (<200 km apart), woodcock ecology varies considerably, highlighting the
complexity of defining suitable habitat for a population. Although woodcocks are
widespread and a well-studied species, our study reveals important knowledge gaps
in their ecology. This highlights the misunderstanding we likely have about a variety
of species because they are only studied in a small portion of their temporal and
geographic range, even though animal responses to available habitat are often
context-dependent (Broekman et al., 2022; Constible et al., 2010). Our results
therefore highlight a well-known but often overlooked principle in species-habitat
relationships: extrapolating of results beyond the geographic or temporal range of the
studied population is precarious. To refine the assessment of habitat suitability, it is
essential to study a sufficiently representative variety of environments within the
distribution and behavior-specific habitat of a species (Roever et al., 2014).

Conclusion

The study of woodcocks wintering in the Mediterranean region shows the remarkable
behavioral flexibility of this species at the climatic edges of its range. However, this
flexibility makes it difficult to predict the dynamics of the woodcocks' geographic
range under climate change (Oldfather et al., 2020). While behavioral adaptations
appear to mitigate some challenges, their long-term effectiveness remains unclear.
Our results suggest that behavioral adaptations, such as habitat use modification, are
often preferred over relocation to more suitable wintering areas. Nevertheless, we
lack data to determine whether these behavioral changes maintain or worsen fitness

benefits. Measuring fitness is crucial to assess whether the persistence of woodcocks
in the Mediterranean regions during wintering is at risk in the near future.

The Mediterranean region is probably one of the most vulnerable areas to exceed the
climatic conditions tolerable by woodcocks due to ongoing climate change. The
effects of climate change in the Mediterranean region, including decreasing
precipitation combined with an increase in the frequency and severity of droughts,
could have a significant impact on local fauna, but are not yet well understood
(Cramer et al., 2018; Ferreira et al., 2022) and present a pressing concern (Moatti &
Thiébault, 2016). For instance, drought conditions can induce physiological stress
(Dezetter et al., 2022), alter prey communities (Martín et al., 2023) and influence
habitat use (Pearse et al., 2024). This underlines the urgent need to study species-
specific ecological requirements in Mediterranean ecosystems to predict how climate
change might affect species distribution and survival and to develop effective
conservation strategies.

Appendices

Supplementary data to this article can be found online in Zenodo
(https://doi.org/10.5281/zenodo.15056074).

Acknowledgement

We warmly thank the professionals, students and volunteers for their assistance in
the data collection: Luc Aurousseau, Christophe Mathez, Guillaume Nard, Damien
Dubosc, Ludovic Fallais, Sébastien Blanchard, Jean-Claude Etienne, Maxime
Passerault and Damien Coreau. We are also grateful to Elodie Coudour for her
valuable support with administrative tasks. This project was funded by the
ecocontribution (Office Français de la Biodiversité, grant no. 1251).

References

Aradis, A., Lo Verde, G., & Massa, B. (2019). Importance of millipedes (Diplopoda) in the

autumn-winter diet of *Scolopax rusticola*. *The European Zoological Journal*, *86*(1),

452–457. https://doi.org/10.1080/24750263.2019.1611955

Arizaga, J., Crespo-Díaz, A., Ansorregi, F., Galdós, A., Urruzola, A., & Iriarte, E. (2015). The

impact of several environmental factors on density of woodcocks (*Scolopax rusticola*)

wintering in a southern European region. *European Journal of Wildlife Research*,

61(3), 407–413. https://doi.org/10.1007/s10344-015-0908-6

Bailey, L. D., Ens, B. J., Both, C., Heg, D., Oosterbeek, K., & van de Pol, M. (2019). Habitat

selection can reduce effects of extreme climatic events in a long-lived shorebird.

Journal of Animal Ecology, *88*(10), 1474–1485. https://doi.org/10.1111/1365-

2656.13041

Bartoń, K. (2024). *MuMIn: Multi-Model Inference* (Version R package version 1.48.4)

[Computer software]. https://CRAN.R-project.org/package=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Ournal of Statistical Software*, *67*(1), 1–4.

https://doi.org/10.18637/jss.v067.i01

Beever, E. A., Hall, L. E., Varner, J., Loosen, A. E., Dunham, J. B., Gahl, M. K., Smith, F. A., &

Lawler, J. J. (2017). Behavioral flexibility as a mechanism for coping with climate

change. *Frontiers in Ecology and the Environment*, *15*(6), 299–308.

https://doi.org/10.1002/fee.1502

Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of

climate change on the future of biodiversity. *Ecology Letters*, *15*(4), 365.

https://doi.org/10.1111/j.1461-0248.2011.01736.x

Beyer, H. L., Haydon, D. T., Morales, J. M., Frair, J. L., Hebblewhite, M., Mitchell, M., &

Matthiopoulos, J. (2010). The interpretation of habitat preference metrics under use–

availability designs. *Philosophical Transactions of the Royal Society B: Biological

Sciences*, *365*(1550), 2245–2254. https://doi.org/10.1098/rstb.2010.0083

Blackburn, E., & Cresswell, W. (2016). High winter site fidelity in a long-distance migrant:

Implications for wintering ecology and survival estimates. *Journal of Ornithology*,

157(1), 93–108. https://doi.org/10.1007/s10336-015-1252-z

Boyce, M. S., Vernier, P. R., Nielsen, S. E., & Schmiegelow, F. K. A. (2002). Evaluating

resource selection functions. *Ecological Modelling*, *157*(2), 281–300.

https://doi.org/10.1016/S0304-3800(02)00200-4

Braña, F., González-Quirós, P., Prieto, L., & González, F. (2013). Spatial distribution and

scale-dependent habitat selection by Eurasian Woodcocks *Scolopax rusticola* at the

south-western limit of its continental breeding range in northern Spain. *Acta*

Ornithologica, *48*(1), 27–37. https://doi.org/10.3161/000164513X669973

Braña, F., Prieto, L., & González-Quirós, P. (2010). Habitat change and timing of dusk flight in

the Eurasian Woodcock: A trade-off between feeding and predator avoidance?

Annales Zoologici Fennici, *47*(3), 206–214. https://doi.org/10.5735/086.047.0305

Broekman, M. J. E., Hilbers, J. P., Huijbregts, M. A. J., Mueller, T., Ali, A. H., Andrén, H.,

Altmann, J., Aronsson, M., Attias, N., Bartlam-Brooks, H. L. A., van Beest, F. M.,

Belant, J. L., Beyer, D. E., Bidner, L., Blaum, N., Boone, R. B., Boyce, M. S., Brown,

M. B., Cagnacci, F., … Tucker, M. A. (2022). Evaluating expert-based habitat

suitability information of terrestrial mammals with GPS-tracking data. *Global Ecology*

and Biogeography, *31*(8), 1526–1541. https://doi.org/10.1111/geb.13523

Burnham, K. P., & Anderson, D. R. (Eds.). (2004). *Model Selection and Multimodel Inference*.

Springer. https://doi.org/10.1007/b97636

Chan, Y.-C., Chan, D. T. C., Tibbitts, T. L., Hassell, C. J., & Piersma, T. (2023). Site fidelity of

migratory shorebirds facing habitat deterioration: Insights from satellite tracking and

mark-resighting. *Movement Ecology*, *11*(1), 79. https://doi.org/10.1186/s40462-023-

00443-9

Clausager, I. (1973). Age and sex determination of the Woodcock (*Scolopax rusticola*).

Danish Review of Game Biology, *8*(1), 18.

Constible, J. M., Gregory, P. T., & Larsen, K. W. (2010). The pitfalls of extrapolation in

779   conservation: Movements and habitat use of a threatened toad are different in the

780   boreal forest. *Animal Conservation*, *13*(1), 43–52. https://doi.org/10.1111/j.1469-

781   1795.2009.00291.x

782   Cramer, W., Guiot, J., Fader, M., Garrabou, J., Gattuso, J.-P., Iglesias, A., Lange, M. A.,

783   Lionello, P., Llasat, M. C., Paz, S., Peñuelas, J., Snoussi, M., Toreti, A., Tsimplis, M.

784   N., & Xoplaki, E. (2018). Climate change and interconnected risks to sustainable

785   development in the Mediterranean. *Nature Climate Change*, *8*(11), 972–980.

786   https://doi.org/10.1038/s41558-018-0299-2

787   Dawson, T. P., Jackson, S. T., House, J. I., Prentice, I. C., & Mace, G. M. (2011). Beyond

788   predictions: Biodiversity conservation in a changing climate. *Science*, *332*(6025), 53–

789   58. https://doi.org/10.1126/science.1200303

790   Dezetter, M., Le Galliard, J.-F., Leroux-Coyau, M., Brischoux, F., Angelier, F., & Lourdais, O.

791   (2022). Two stressors are worse than one: Combined heatwave and drought affect

792   hydration state and glucocorticoid levels in a temperate ectotherm. *Journal of*

793   *Experimental Biology*, *225*(7), jeb243777. https://doi.org/10.1242/jeb.243777

794   Duriez, O., Ferrand, Y., Binet, F., Corda, E., Gossmann, F., & Fritz, H. (2005). Habitat

795   selection of the Eurasian Woodcock in winter in relation to earthworms availability.

796   *Biological Conservation*, *122*(3), 479–490.

797   https://doi.org/10.1016/j.biocon.2004.08.011

798   Duriez, O., Fritz, H., Binet, F., Tremblay, Y., & Ferrand, Y. (2005). Individual activity rates in

799   wintering Eurasian Woodcocks: Starvation versus predation risk trade-off? *Animal*

800   *Behaviour*, *69*(1), 39–49. https://doi.org/10.1016/j.anbehav.2004.04.009

801   Duriez, O., Fritz, H., Said, S., & Ferrand, Y. (2005). Wintering behaviour and spatial ecology

802   of Eurasian Woodcock *Scolopax rusticola* in western France. *Ibis*, *147*(3), 519–532.

803   https://doi.org/10.1111/j.1474-919x.2005.00423.x

Duriez, O., Pastout-Lucchini, L., Boos, M., Chastel, O., Fritz, H., Ferrand, Y., & Clobert, J.

(2004). Low levels of energy expenditure in a nocturnal, forest-dwelling wader, the

Eurasian Woodcock *Scolopax Rusticola*. *Ardea*, *92*(1), 31.

Ferrand, Y., Gossmann, F., Bastat, C., & Guénézan, M. (2008). Monitoring of the wintering

and breeding Woodcock populations in France. *Revista Catalana d'Ornitologia*, *24*,

44–52.

Ferreira, C. S. S., Seifollahi-Aghmiuni, S., Destouni, G., Ghajarnia, N., & Kalantari, Z. (2022).

Soil degradation in the European Mediterranean region: Processes, status and

consequences. *Science of The Total Environment*, *805*, 150106.

https://doi.org/10.1016/j.scitotenv.2021.150106

Fieberg, J., Signer, J., Smith, B., & Avgar, T. (2021). A 'How to' guide for interpreting

parameters in habitat-selection analyses. *Journal of Animal Ecology*, *90*(5), 1027–

1043. https://doi.org/10.1111/1365-2656.13441

Gillies, C. S., Hebblewhite, M., Nielsen, S. E., Krawchuk, M. A., Aldridge, C. L., Frair, J. L.,

Saher, D. J., Stevens, C. E., & Jerde, C. L. (2006). Application of random effects to

the study of resource selection by animals. *Journal of Animal Ecology*, *75*(4), 887–

898. https://doi.org/10.1111/j.1365-2656.2006.01106.x

Gonçalves, D., Rodrigues, T. M., Pennacchini, P., Lepetit, J.-P., Taaffe, L., Tuti, M., Meunier,

B., Campana, J.-P., Gregori, G., Pellegrini, A., Raho, G., Duchein, P., Trotman, C.,

Minondo, M., Fitzgerald, D., Verde, A., Aurousseau, G., Díez, F., Le-Rest, K., &

Ferrand, Y. (2019). Survey of wintering Eurasian Woodcock in Western Europe.

American Woodcock Symposium. https://doi.org/10.24926/AWS.0131

Gongalsky, K. B. (2021). Soil macrofauna: Study problems and perspectives. *Soil Biology*

and Biochemistry, *159*, 108281. https://doi.org/10.1016/j.soilbio.2021.108281

Gossmann, F., Ferrand, Y., Loidon, Y., & Sardet, G. (1988). *Méthodes et résultats de*

baguages des bécasses des bois (Scolopax rusticola) en Bretagne (P. Havet & G.

Hirons, Eds.; pp. 34–41).

Granval, P. (1986). Variations du régime alimentaire diurne de la Bécasse des bois (*Scolopax*

rusticola L.). *3ème Symposium Européen Sur La Bécasse et La Bécassine*, 67–77.

Griffiths, R., Double, M. C., Orr, K., & Dawson, R. J. G. (1998). A DNA test to sex most birds.

Molecular Ecology, *7*(8), 1071–1075. https://doi.org/10.1046/j.1365-

294x.1998.00389.x

Guzmán, J. L., Caro, J., & Arroyo, B. (2017). Factors influencing mobility and survival of

Eurasian Woodcock wintering in Spain. *Avian Conservation and Ecology*, *12*(2).

https://doi.org/10.5751/ACE-01096-120221

Hirons, G., & Johnson, T. H. (1987). A quantitative analysis of habitat preferences of

Woodcock *Scolopax rusticola* in the breeding season. *Ibis*, *129*(2), 371–381.

https://doi.org/10.1111/j.1474-919X.1987.tb03181.x

Hoenig, B. D., Snider, A. M., Forsman, A. M., Hobson, K. A., Latta, S. C., Miller, E. T., Polito,

M. J., Powell, L. L., Rogers, S. L., Sherry, T. W., Toews, D. P. L., Welch, A. J., Taylor,

S. S., & Porter, B. A. (2022). Current methods and future directions in avian diet

analysis. *Ornithology*, *139*(1), ukab077. https://doi.org/10.1093/ornithology/ukab077

Hollister, J., Shah, T., Nowosad, J., Robitaille, A. L., Beck, M. W., & Johnson, M. (2023).

elevatr: Access Elevation Data from Various APIs (Version 0.99) [Computer software].

https://rdrr.io/cran/elevatr/

Hoodless, A. N., & Hirons, G. J. M. (2007). Habitat selection and foraging behaviour of

breeding Eurasian Woodcock *Scolopax rusticola*: A comparison between contrasting

landscapes. *Ibis*, *149*(s2), 234–249. https://doi.org/10.1111/j.1474-919X.2007.00725.x

Inkscape Project. (2023). *Inkscape* (Version 1.3.2) [Computer software]. https://inkscape.org

IPBES. (2019). *Global assessment report on biodiversity and ecosystem services of the*

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services

(E. S. Brondízio, J. Settele, S. Díaz, & H. T. Ngo, Eds.; IPBES secretariat).

https://zenodo.org/records/6417333

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest package: Tests in
linear mixed effects models. *Journal of Statistical Software*, *82*, 1–26.
https://doi.org/10.18637/jss.v082.i13

Langley, L. P., Bearhop, S., Burton, N. H. K., Banks, A. N., Frayling, T., Thaxter, C. B.,
Clewley, G. D., Scragg, E., & Votier, S. C. (2021). GPS tracking reveals landfill
closures induce higher foraging effort and habitat switching in gulls. *Movement*
Ecology, *9*(1), 56. https://doi.org/10.1186/s40462-021-00278-2

Mabille, G., Dussault, C., Ouellet, J.-P., & Laurian, C. (2012). Linking trade-offs in habitat
selection with the occurrence of functional responses for moose living in two nearby
study areas. *Oecologia*, *170*(4), 965–977. https://doi.org/10.1007/s00442-012-2382-0

Mallet, P., Bechet, A., Sirami, C., Mesleard, F., Blanchon, T., Calatayud, F., Dagonet, T.,
Gaget, E., Leray, C., & Galewski, T. (2023). Field margins as substitute habitat for the
conservation of birds in agricultural wetlands. *Peer Community Journal*, *3*.
https://doi.org/10.24072/pcjournal.299

Martín, J., Ortega, J., García-Roa, R., Rodríguez-Ruiz, G., Pérez-Cembranos, A., & Pérez-
Mellado, V. (2023). Coping with drought? Effects of extended drought conditions on
soil invertebrate prey and diet selection by a fossorial amphisbaenian reptile. *Current*
Zoology, *69*(4), 367–376. https://doi.org/10.1093/cz/zoac056

Mayor, S. J., Schneider, D. C., Schaefer, J. A., & Mahoney, S. P. (2009). Habitat selection at
multiple scales. *Écoscience*, *16*(2), 238–247. https://doi.org/10.2980/16-2-3238

Mazerolle, D. F., Sealy ,Spencer G., & and Hobson, K. A. (2011). Interannual flexibility in
breeding phenology of a Neotropical migrant songbird in response to weather
conditions at breeding and wintering areas. *Écoscience*, *18*(1), 18–25.
https://doi.org/10.2980/18-1-3345

Merkle, J. A., Abrahms, B., Armstrong, J. B., Sawyer, H., Costa, D. P., & Chalfoun, A. D.
(2022). Site fidelity as a maladaptive behavior in the Anthropocene. *Frontiers in*
Ecology and the Environment, *20*(3), 187–194. https://doi.org/10.1002/fee.2456

Moatti, J.-P., & Thiébault, S. (2016). The Mediterranean region under climate change: A

scientific update. In *The Mediterranean region under climate change: A scientific*

update. IRD Éditions. https://books.openedition.org/irdeditions/22908

Northrup, J. M., Vander Wal, E., Bonar, M., Fieberg, J., Laforge, M. P., Leclerc, M.,

Prokopenko, C. M., & Gerber, B. D. (2022). Conceptual and methodological advances

in habitat-selection modeling: Guidelines for ecology and evolution. *Ecological*

Applications, *32*(1), e02470. https://doi.org/10.1002/eap.2470

Obuch, J. (2024). The diet of the Eurasian eagle-owl (*Bubo bubo*) in various natural

environments across Eurasia. *Raptor Journal*, *18*(1), 25–46.

https://doi.org/10.2478/srj-2024-0004

Oldfather, M. F., Kling, M. M., Sheth, S. N., Emery, N. C., & Ackerly, D. D. (2020). Range

edges in heterogeneous landscapes: Integrating geographic scale and climate

complexity into range dynamics. *Global Change Biology*, *26*(3), 1055–1067.

https://doi.org/10.1111/gcb.14897

Patterson, A., Gilchrist, H. G., Chivers, L., Hatch, S., & Elliott, K. (2019). A comparison of

techniques for classifying behavior from accelerometers for two species of seabird.

Ecology and Evolution, *9*(6), 3030–3045. https://doi.org/10.1002/ece3.4740

Pearse, A. T., Caven, A. J., Baasch, D. M., Bidwell, M. T., Conkin, J. A., & Brandt, D. A.

(2024). Flexible migration and habitat use strategies of an endangered waterbird

during hydrological drought. *Conservation Science and Practice*, *6*(5), e13120.

https://doi.org/10.1111/csp2.13120

Pebesma, E., Bivand, R., Racine, E., Sumner, M., Cook, I., Keitt, T., Lovelace, R., Wickham,

H., Ooms, J., Müller, K., Pedersen, T. L., Baston, D., & Dunnington, D. (2024). *sf:*

Simple features for R (Version 1.0-16) [Computer software]. https://cran.r-

project.org/web/packages/sf/index.html

Péron, G., Ferrand, Y., Gossmann, F., Bastat, C., Guénézan, M., & Gimenez, O. (2011).

Escape migration decisions in Eurasian Woodcocks: Insights from survival analyses

using large-scale recovery data. *Behavioral Ecology and Sociobiology*, *65*(10), 1949–

1955. https://doi.org/10.1007/s00265-011-1204-4

Piper, W. H. (2011). Making habitat selection more "familiar": A review. *Behavioral Ecology*

and Sociobiology, *65*(7), 1329–1351. https://doi.org/10.1007/s00265-011-1195-1

QGIS.org. (2024). *QGIS Geographic Information System. Open source geospatial foundation*

project. (Version 3.28.1) [Computer software]. https://www.qgis.org/

R Core Team. (2024). *R: A language and environment for statistical computing.* (Version

4.4.1) [Computer software]. R Foundation for Statistical Computing.

Réseau Bécasse—Lettre d'information n° 27. (2018).

Roever, C. L., Beyer, H. L., Chase, M. J., & van Aarde, R. J. (2014). The pitfalls of ignoring

behaviour when quantifying habitat selection. *Diversity and Distributions*, *20*(3), 322–

333. https://doi.org/10.1111/ddi.12164

Sardans, J., & Peñuelas, J. (2013). Plant-soil interactions in Mediterranean forest and

shrublands: Impacts of climatic change. *Plant and Soil*, *365*(1), 1–33.

https://doi.org/10.1007/s11104-013-1591-6

Shahsavarzadeh, R., Hemami, M.-R., Farhadinia, M. S., Fakheran, S., & Ahmadi, M. (2023).

Spatially heterogeneous habitat use across distinct biogeographic regions in a wide-

ranging predator, the Persian leopard. *Biodiversity and Conservation*, *32*(6), 2037–

2053. https://doi.org/10.1007/s10531-023-02590-2

Skórka, P., Lenda, M., & Sutherland, W. J. (2016). Response of young and adult birds to the

same environmental variables and different spatial scales during post breeding

period. *Landscape Ecology*, *31*(9), 2063–2078. https://doi.org/10.1007/s10980-016-

0382-x

Sládeček, M., Pešková, L., Chajma, P., Brynychová, K., Koloušková, K., Trejbalová, K.,

Kolešková, V., Petrusová Vozabulová, E., & Šálek, M. E. (2023). Eurasian woodcock

(Scolopax rusticola) in intensively managed Central European forests use large home

ranges with diverse habitats. *Forest Ecology and Management*, *550*, 121489.

https://doi.org/10.1016/j.foreco.2023.121489

Smith, J., Potts, S., & Eggleton, P. (2008). Evaluating the efficiency of sampling methods in

assessing soil macrofauna communities in arable systems. *European Journal of Soil*

Biology, *44*(3), 271–276. https://doi.org/10.1016/j.ejsobi.2008.02.002

Tedeschi, A., Sorrenti, M., Bottazzo, M., Spagnesi, M., Telletxea, I., Ibàñez, R., Tormen, N.,

De Pascalis, F., Guidolin, L., & Rubolini, D. (2020). Interindividual variation and

consistency of migratory behavior in the Eurasian Woodcock. *Current Zoology*, *66*(2),

155–163. https://doi.org/10.1093/cz/zoz038

Thurman, L. L., Stein, B. A., Beever, E. A., Foden, W., Geange, S. R., Green, N., Gross, J.

E., Lawrence, D. J., LeDee, O., Olden, J. D., Thompson, L. M., & Young, B. E. (2020).

Persist in place or shift in space? Evaluating the adaptive capacity of species to

climate change. *Frontiers in Ecology and the Environment*, *18*(9), 520–528.

https://doi.org/10.1002/fee.2253

Verzuh, T. L., Hall, L. E., Cufaude, T., Knox, L., Class, C., & Monteith, K. L. (2021).

Behavioural flexibility in a heat-sensitive endotherm: The role of bed sites as thermal

refuges. *Animal Behaviour*, *178*, 77–86.

https://doi.org/10.1016/j.anbehav.2021.05.020

Wasserlauf, Y., Gancz, A., Ben Dov, A., Efrat, R., Sapir, N., Dor, R., & Spiegel, O. (2023). A

telemetry study shows that an endangered nocturnal avian species roosts in

extremely dry habitats to avoid predation. *Scientific Reports*, *13*(1), 11888.

https://doi.org/10.1038/s41598-023-38981-2

Wellbrock, A. H. J., Bauch, C., Rozman, J., & Witte, K. (2017). 'Same procedure as last

year?' Repeatedly tracked swifts show individual consistency in migration pattern in

successive years. *Journal of Avian Biology*, *48*(6), 897–903.

https://doi.org/10.1111/jav.01251

Wilson, R. P., White, C. R., Quintana, F., Halsey, L. G., Liebsch, N., Martin, G. R., & Butler, P.

J. (2006). Moving towards acceleration for estimates of activity-specific metabolic rate

in free-living animals: The case of the cormorant. *Journal of Animal Ecology*, *75*(5),

1081–1090. https://doi.org/10.1111/j.1365-2656.2006.01127.x

Yates, K. L., Bouchet, P. J., Caley, M. J., Mengersen, K., Randin, C. F., Parnell, S., Fielding,

968   A. H., Bamford, A. J., Ban, S., Barbosa, A. M., Dormann, C. F., Elith, J., Embling, C.

969   B., Ervin, G. N., Fisher, R., Gould, S., Graf, R. F., Gregr, E. J., Halpin, P. N., …

970   Sequeira, A. M. M. (2018). Outstanding challenges in the transferability of ecological

971   models. *Trends in Ecology & Evolution*, *33*(10), 790–802.

972   https://doi.org/10.1016/j.tree.2018.08.001

973