TEITbase: a database for transposable element (TE)-initiated transcripts in human cancers

Yun Zhang[1,#], Jianqi She[2,3,#], Xueyan Hu[2], Yueqi Jin[1], Changyu Tao[4], Minghao Du[2,*], Ence Yang[1,2,*]

[1] Department of Medical Bioinformatics, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China

[2] Department of Microbiology & Infectious Disease Center, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China

[3] Department of Radiation Medicine, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China

[4] Department of Human Anatomy, Histology and Embryology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China

[#] Contributed equally

[*] Correspondence: yangence@pku.edu.cn (Ence Yang), duminghao@pku.edu.cn (Minghao Du)

Abstract

Transposable elements (TEs) are abundant and play a crucial regulatory role in the human genome. Functioning as alternative promoters, TEs can be reactivated to produce TE-initiated transcripts. In cancer, TE-initiated transcription may upregulate oncogene expression and generate novel tumor-specific antigens, which could serve as potential targets for immunotherapy. However, there remains a lack of comprehensive databases that systematically investigate TE-initiated transcription in cancer. To address this gap, we developed a deep learning-based method and identified 38,995 TE-initiated transcripts across 33 cancer types from The Cancer Genome Atlas. Among these, 6,203 were tumor-specific expressed and strongly associated with the tumorigenesis. Using these annotations, we created TEITbase (http://teitbase.medbioinfo.org/), a user-friendly database that provides researchers with tools to conduct various downstream analyses and investigations for TE-initiated transcripts, including 546 novel onco-exaptation events. The establishment of TEITbase provides new insights into transcriptional reprogramming in cancers and enables further investigation into the potential roles of TE-initiated transcripts in cancer diagnostics and therapeutic strategies.

Keywords: transposable element, database, cancer, deep learning

Introduction

Transposable elements (TEs) are mobile genetic elements that are widespread in eukaryotic genomes (1). Approximately 45% of the human genome is derived from various classes of TEs, such as LINEs (long interspersed nuclear elements), SINEs (short interspersed nuclear elements), LTRs (long terminal repeats), and DNA transposons (2,3). In humans, only a small proportion of TEs, including active human-specific L1 (L1HS) elements, remain capable of transposition and inducing genetic changes (4,5). The majority of TEs in the human genome are incomplete and no longer transposable (6). However, TE-derived sequences retain significant regulatory potential, acting as promoters, enhancers, splicing sites, and more, playing crucial roles in both physiological and pathological processes (6–9).

As potential threats to human genome, most TEs have been kept inactive in somatic cells by mutational events and epigenetic repression (6,10). However, accumulating evidence suggests that certain TEs are transcriptionally reactivated in human cancers as a result of epigenetic dysregulation (11,12). Acting as alternative promoters, TEs can enhance their own transcription or regulate the expression of neighboring genes, leading to the production of TE-initiated transcripts with TE-derived transcription start sites (TSSs) (13). In cancer, TE-initiated transcription can upregulate the expression of oncogenes, a phenomenon known as onco-exaptation (14–16). Notable examples include Alujb-*LIN28B*, L1-*FGGY* and HERVH-*CALB1* (15,17,18). Furthermore, TE-initiated transcription may produce novel protein isoforms and tumor-specific antigens, which could serve as potential targets for immunotherapy (19,20). For instance, the tumor-specific membrane protein GABRG2, transcribed from the L1PA2 promoter, can generate aberrant epitopes on the extracellular surface of cancer cells (12).

Moreover, endogenous retrovirus (ERV) envelope glycoproteins have been identified as a

dominant anti-tumor antibody target and ERV-reactive antibodies exhibit anti-tumor activity

that extends survival in mouse models (21).

Given the recognized role of TEs in tumor progression, several databases, including

ERVcancer and CancerHERVdb, have been developed to investigate TE expression in human

cancers (22,23). However, these databases only provide TE expression data at the TE family

level, losing locus information and failing to distinguish between TE-initiated transcription and

TE exonization. Recently, Gu et al. established TE-TSS, a database compiling 2,681 RNA

sequencing datasets that identifies 5,768 human TE-derived TSSs and 2,797 mouse TE-derived

TSSs (24). Nevertheless, due to the repetitive nature of TEs and the limitations of short-read

RNA-seq, TE-TSS cannot provide accurate annotations of TE-initiated transcripts. Deep

learning techniques, which can automatically extract features from DNA sequence information,

have been widely applied to predict TSSs (25,26). By integrating RNA-seq data with DNA

sequence information, convolutional neural networks could accurately predict TSSs for each

assembled transcript, thereby enabling the identification of TE-initiated transcripts. However,

there remains a lack of databases that specifically annotate TE-initiated transcripts in human

cancers.

In this work, we developed the TEITbase (http://teitbase.medbioinfo.org/), an integrated

database that systematically studies the TE-initiated transcription across 33 cancer types from

The Cancer Genome Atlas (TCGA), offering data at both the TE family and transcript levels.

To identify TE-initiated transcripts, we first developed a deep learning-based method to predict

transcription start sites (TSSs) for each assembled transcript, utilizing both DNA sequences and

short-read RNA sequencing data. Subsequently, we applied this method to 10,079 tumor

samples from TCGA, and identified 38,995 TE-initiated transcripts. These annotations enable

TEITbase to provide researchers with the tools to conduct various downstream analyses, such

as promoter activity of TE families, tumor-specific TE-initiated transcripts, differential

expression analysis, stage analysis, and survival analysis. We believe that TEITbase will serve

as a valuable resource for further understanding the role of TE-initiated transcription in

epigenetically dysregulated tumor tissues and will be instrumental in investigating novel tumor-

specific antigens that could potentially be therapeutically targeted.

Results

A deep learning-based method for the identification of TE-initiated transcript

Due to the repetitive nature of TEs and the limitations of short-read RNA-seq, identifying TE-derived TSSs based on RNA-seq coverage is not sufficiently accurate. To overcome the challenge, we developed a deep learning-based method that integrates both DNA sequence and RNA-seq coverage information. Using one-hot encoding and convolutional neural network, we predicted TSSs for each assembled transcript and overlapped them with TE annotations to identify TE-initiated transcripts (Fig. 1A).

To train our model, we downloaded CapTrap-seq datasets along with matched short-read RNA-seq datasets (27). Unlike CAGE and RAMPAGE, CapTrap-seq combines the cap-trapping strategy with long-read sequencing technologies, enabling the identification of both TSSs and full-length transcripts. Using these CapTrap-seq datasets for training, our model can predict TSSs at the transcript level. For example, *PRKACB* has alternative promoters in brain tissue. While RAMPAGE identifies only two TSS signals, it cannot assign them to specific transcripts. In contrast, both CapTrap-seq and our model can identify the TSS for each transcript (Fig. 1B).

Compared to StringTie, which relies solely on RNA-seq coverage to determine TSS, our model significantly enhances the accuracy of TSS prediction ($P = 4.6 \times 10^{-4}$, Fig. 1C, Table S1). In the validation set, our model accurately identified an average of approximately 26.7% of TSSs, with about 84.7% of TSSs exhibiting identification errors within 100 bp, a notable improvement over StringTie's performance of 2.4% and 70.6%, respectively. As expected, our model performed significantly better in predicting highly expressed transcripts (Fig. 1C, $P = $

1.2×10^{-5}). For transcripts with counts per million (CPM) greater than 2, an average of

approximately 71.0% of TSSs were accurately identified by our model, with around 99.2% of

TSSs having identification errors within 100 bp. This is substantially better than for low-

expressed transcripts (CPM < 0.1), which had 9.1% and 72.7% accuracy within 100 bp,

respectively. Furthermore, the prediction score of our model was significantly and negatively

correlated with TSS prediction errors (Fig. 1D, Spearman's $r = -0.50$, $P < 2.2 \times 10^{-16}$), enabling

us to improve TSS identification accuracy by setting an appropriate prediction score threshold.

Through the application of the deep learning model and strict criteria, our method

significantly outperforms DeeReCT-TSS in identifying TE-TSS (Table S2, $P = 0.029$) (26). In

the H1 cell line, our method identified 954 TE-TSS, of which 626 (65.6%) were supported by

CapTrap-seq data. In contrast, DeeReCT-TSS identified only 390 TE-TSS, with just 12 (3.1%)

supported by CapTrap-seq data (Fig. 1E). When compared to the TE-TSS database (24), our

method identified 64 previously annotated TE-TSS and 908 novel TE-TSS, of which 580

(63.9%) were supported by CapTrap-seq data (Fig. 1E). To exclude the potential impact of

model overfitting on TE-TSS identification, we downloaded short-read RNA-seq data from 18

testis tissues as an external validation dataset (28), with RAMPAGE data serving as the gold

standard. Our method identified 3,979 TE-TSS, of which 2,909 (73.1%) were supported by

RAMPAGE data, while DeeReCT-TSS identified only 286 TE-TSS, with just 107 (37.4%)

supported by RAMPAGE data (Fig. 1F). Compared to the TE-TSS database, our method

identified 157 previously annotated TE-TSS and 3,822 novel TE-TSS, of which 2,752 (72.0%)

were supported by RAMPAGE data (Fig. 1F). These results suggest that we have developed a

deep learning-based method that can accurately identify TE-initiated transcripts using multiple

biological replicates of short-read RNA-seq data, significantly outperforming existing methods.

Due to the relatively accurate annotation of TE-initiated transcripts, our method effectively

distinguishes between TE-initiated transcription and TE exonization, enabling the measurement

of promoter activity in TE families. To evaluate the quantitative performance at the family level,

we downloaded short-read RNA-seq data and corresponding CAGE data (used as the gold

standard for quantitative evaluation) from six cancer cell lines (A549, Hela, HepG2, K562,

MCF-7, and SK-N-SH) from the ENCODE database (29). Our method demonstrated

significantly better consistency with CAGE data compared to existing TE family expression

quantification methods, TEtranscripts (Fig. 1G, $P < 2.2 \times 10^{-16}$). Among all TE families, the

median Spearman correlation coefficient between the quantitative results from our method and

CAGE data across these six cell lines was 0.26, while TEtranscripts had a value of -0.37.

Furthermore, our method identified 105 TE families with high accuracy (Spearman's $r \geq 0.7$),

whereas TEtranscripts identified only 6. Notably, the age of these 105 high-accuracy TE

families (measured by the evolutionary distance to ancestral TEs: Kimura value) was

significantly lower than that of other TE families (Fig. 1G, $P = 9.2 \times 10^{-4}$), which may be related

to the higher retention of promoter activity in younger TEs.

Genome-wide identification of TE-initiated transcripts in pan-cancers

In 10,079 tumor samples across 33 cancer types from the TCGA, we identified 38,995 TE-

initiated transcripts (Table S3), of which 34,670 were not annotated in GENCODE.v46 (30).

The expression of these TE-initiated transcripts was associated with the activation of 39,703

TE loci, including 11,857 SINE elements, 11,758 LTR elements, 10,911 LINE elements, and

2,580 DNA transposons (Fig. 2A). These active TEs contributed to the expression of 14,453

neighboring genes, including 9,315 protein-coding genes and 4,771 lncRNA genes (Fig. 2B),

suggesting that TEs play a widespread role in regulating gene expression in tumor tissues.

Tumor-specific expressed transcripts play a critical role in tumorigenesis and represent

important sources of tumor-specific antigens, which may provide new targets for cancer

immunotherapy. To investigate this, we performed the expression analysis of TE-initiated

transcripts in 10,079 tumor samples, 740 tumor-adjacent samples and 10,620 non-testicular

normal tissue samples. Tumor-specific scores were calculated, identifying 6,203 tumor-specific

TE-initiated transcripts (scores ≥ 8, Table S4). These activated tumor-specific TE-initiated

transcripts were associated with the expression of 1,532 adjacent protein-coding genes. As

anticipated, enrichment analysis revealed significant overrepresentation in tumorigenesis-

related pathways, including regulation of *PTEN* gene transcription and TP53 activity (Fig. 2C,

Table S5). Among the 702 oncogenes annotated in the ONGene database (31), we found that

TEs upregulated the expression of 333 oncogenes (a process known as onco-exaptation),

generating 696 TE-initiated transcripts (Table S6), including previously reported onco-

exaptations such as AluJb-*LIN28B*, L1HS-*SYT1*, L1HS-*FGGY*, and L1HS-*MET* (15,18,32).

Compared to a recent similar study (15), we identified 150 reported and 546 novel onco-

exaptation events, with 263 (48.2%) supported by CAGE data from at least one cancer cell line.

Among them, AluJr-*AKT2* was expressed across all cancer types and detected in 619 tumor

samples, further highlighting the universality of onco-exaptation events. Furthermore, TE

family enrichment analysis revealed significant overrepresentation of young families, including

L1HS, L1P1, and HERVH (Fig. 2D, Table S7). These evolutionarily young TE families retain

regulatory sequences that are typically suppressed in somatic cells but are reactivated in the

epigenetically dysregulated tumor tissues, driving the TE-initiated transcription and potentially

promoting tumor progression.

To illustrate, we analyzed the 10 TE-initiated transcripts with the highest tumor-specific

scores and found that 8 were initiated by the L1HS family (Fig. 2E). These 10 TE-initiated

transcripts were expressed in 275 to 1,069 tumor samples but only in 1 or 2 normal tissue

samples, indicating their high tumor specificity. Moreover, these activated TEs were critical in

regulating the expression of adjacent genes, with L1HS-*CNBD1*, L1HS-*CFAP47*, and L1P2-

PATE2 serving as the primary transcripts for their respective genes. Furthermore, *EYA4* and

NLRP7 have been shown to promote cell proliferation, invasion, and migration in various

cancers (33,34), suggesting that the expression of L1HS-*EYA4* and AluSz-*NLRP7* may have

significant oncogenic effects. Survival analysis further indicated that the expression of L1HS-

EYA4, L1HS-*CRPPA*, L1HS-*CFAP47*, and L1P2-*PATE2* was associated with poor prognosis in

at least one cancer type (Fig. 2F). In conclusion, these highly tumor-specific TE-initiated

transcripts may play a role in tumorigenesis and hold potential as biomarkers for tumor

diagnosis and prognosis.

Content and usage of TEITbase

Based on the identification of TE-initiated transcripts across various cancer types, we have

developed a free, user-friendly website, TEITbase (http://teitbase.medbioinfo.org/), which

provides expression data for 1,027 TE families and 38,995 TE-initiated transcripts (Fig. 3A).

The website consists of several main pages: Home, Browse, Analyze, Statistics, Download,

Help, and Contact. TEITbase offers an intuitive interface and a range of core functionalities,

enabling researchers to explore and understand the role of TE-initiated transcription in cancers.

The Browse page is divided into two sections: TE Family and TE-initiated Transcripts.

Users can query TE family entries by TE family, TE superfamily, and species. The query results

display the TE family, Dfam database ID, TE superfamily, TE classification, species, consensus

sequence length, Kimura value, and tissue-specificity score (Tau index). Users can also query

TE-initiated transcripts using TEITbase database ID, TE family, and gene. The results for these

queries include the TEITbase database ID, TE family, TE superfamily, TE classification,

chimeric gene, transcript classification, coding types, tumor specificity score, and TSS location.

According to the GENCODE v46 annotation, TE-initiated transcripts are classified into three

categories: annotated, chimeric (with a novel splice site at the 5' end), and intergenic (without

known splice sites). Based on predicted open reading frames (ORFs), TE-initiated transcripts

are further categorized into six types: annotated, 5' truncated, 5' chimeric, chimeric truncated,

other coding types, and non-coding. The Analyses page allows users to query tumor-specific

TE-initiated transcripts, as well as survival-associated, stage-associated, and differentially

expressed TE-initiated transcripts (or TE families). Additionally, the Statistics page provides

the basic information of TEITbase, such as the number of TE-initiated transcripts. The Help

page includes a user guide for the database, while the Download page offers links to download

all annotation information, expression matrices, and analysis results.

Using the functions of TEITbase, we investigated the dysregulation of the L1HS family in

cancer. We found that lung squamous carcinoma (LUSC), esophageal carcinoma (ESCA), and

bladder urothelial carcinoma (BLCA) exhibit higher L1HS promoter activity, while normal

tissues show lower activity (Fig. 3B). Differential expression analysis revealed significant

activation of the L1HS promoter in 16 out of 21 cancer types, with the most pronounced

increase in LUSC (Fig. 3C, $P < 2.2 \times 10^{-16}$). In LUSC, L1HS promoter activity in stage II

patients was significantly higher than in stage I patients (Fig. 3D, $P = 0.038$). Survival analysis

indicated that in 5 out of 33 cancer types, including liver hepatocellular carcinoma (LIHC),

abnormal activation of the L1HS promoter was associated with poorer tumor prognosis.

Conversely, activation of the L1HS promoter is significantly correlated with better tumor

prognosis in LUSC (Fig. 3E), adrenocortical carcinoma (ACC), and mesothelioma (MESO).

To explore how L1HS promoter activation improves prognosis, we conducted a correlation

analysis between L1HS promoter activity and gene expression, followed by enrichment

analysis of genes positively correlated with its expression. We discovered that in LUSC,

activation of the L1HS promoter was linked to apoptosis-regulating signaling pathways (Fig.

3F), suggesting that active transposition of L1HS may disrupt tumor genome stability, leading

to cancer cell apoptosis (35).

By utilizing the 'Tumor-specific transcripts' function of TEITbase, we investigated the

role of TE_11880 (L1HS-*IRF1*) in cancers. TE_11880 is initiated by L1HS with a tumor-

specific score of 32.5 and is chimeric with *IRF1*, a transcriptional regulator and tumor

suppressor that activates genes involved in antitumor immunity (36). These findings suggest

that the activation of TE promoters in cancer tissues may also confer benefits to the host. Further

details about TE_11880, including its sequence and other relevant information, can be accessed

in the TEITbase database. TE_11880 is silent in most normal tissue samples but activated in

several cancer types (Fig. 3G, H). We selected specific cancer types to explore the differential

expression, stage, and survival associations of TE_11880. Our differential expression analysis

revealed that TE_11880 expression was primarily restricted to LUSC and stomach

adenocarcinoma (STAD), with no expression in adjacent normal tissues (Fig. 3I). In STAD,

TE_11880 expression in stage III patients was significantly higher than in both stage I and stage

II patients (Fig. 3J). Additionally, TE_11880 expression is associated with a favorable prognosis

in colon adenocarcinoma (COAD), despite not reaching statistical significance ($P = 0.069$) (Fig.

3K), suggesting its potential prognostic value.

Discussion

The activation of TE promoters can significantly impact the transcriptomic and proteomic profiles of cancer (20). Recent studies suggest that TE-initiated transcription may be linked to tumorigenesis and could generate tumor-specific antigens, offering potential targets for immunotherapies (12,15). However, the widespread use of short-read RNA sequencing technologies poses challenges due to the loss of TSS information. To address this, we developed a deep learning-based approach that predicts TSSs and accurately identifies TE-initiated transcripts, enabling the creation of a specialized database, TEITbase. TEITbase offers abundant data resources and various analytical tools, allowing users to explore the roles of TE-initiated transcripts in cancer progression. Importantly, our findings have identified over 6000 tumor-specific TE-initiated transcripts, with the L1HS family being notably enriched. As the youngest L1 family in humans, the L1HS family retains transposable ability and may play a significant role in tumorigenesis (35).

Due to its powerful model fitting capabilities, deep learning has become widely applied to predicting key features from DNA sequences such as gene expression, RNA editing, TSS, alternative splicing, and polyadenylation (25,37–40). Several CNN-based tools have been developed for the accurate prediction of promoters and TSS, including Puffin, TSSPlant, TransPrise, and DeeReCT-TSS (25,26,41,42). Among them, DeeReCT-TSS accurately identifies active TSS across the genome by integrating DNA sequences and RNA-seq coverage information. However, DeeReCT-TSS uses CAGE-seq data to train the model, which loses relevant transcript structure information and cannot assign identified active TSS to specific transcripts. Besides, reads derived from TE sequences often produce ambiguous alignments,

leading to errors in RNA-seq coverage estimation and reducing the accuracy of the

identification of TE-derived TSS (43). To address this, we downloaded CapTrap-seq data to

train the model (27), enabling the prediction of TSS for each transcript, and applied strict

criteria to identify TE-initiated transcript. Our method significantly outperforms DeeReCT-TSS

in identifying TE-derived TSS, providing a new tool to investigate TE-initiated transcription.

In summary, TEITbase is a user-friendly database that provides valuable insights and tools

for investigating TE-initiated transcription in tumorigenesis, as well as exploring the potential

roles of these transcripts in cancer diagnostics and therapeutic strategies. While it already offers

a comprehensive data collection, there are still opportunities for optimization and further

development. Future versions will incorporate TE exonization, which results from

noncanonical splice junctions between exons and TEs, serving as a source of tumor-specific

antigens (8). With ongoing advancements in full-length long-read RNA-seq technologies (16),

TEITbase will integrate publicly available long-read RNA-seq data to capture the full length of

TE-initiated and TE-exonization transcripts. These updates are expected to further enhance

TEITbase's utility and broaden its applicability across various research fields.

Methods

Data collection

We downloaded short-read RNA-seq data from 10,819 samples across 33 cancer types from the The Cancer Genome Atlas (TCGA, phs000178.v11.p8) (44). Additionally, we obtained RNA-seq data from 10,620 samples across 45 human body sites from the GTEx project (by dbGaP, study accession: phs000424.v8.p2) (45). Short-read RNA-seq and CapTrap-seq datasets were downloaded from the ENCODE project and the ArrayExpress repository (E-MTAB-13063) (27,29). Ten samples of these samples were used as the training set (H1 cell line and heart), and three samples were used as the validation set (endodermal cells and brain). Short-read RNA-seq and RAMPAGE datasets of testis were download from the Gene Expression Omnibus database (GSE135791) and the ENCODE project (28). Short-read RNA-seq and CAGE datasets of six cancer cell lines were download from the ENCODE project. GENCODE Version 46 was download from https://www.gencodegenes.org/human/ and used as the transcript reference (30). TE annotations and consensus sequences of TE family were downloaded from the Dfam database (https://www.dfam.org/releases/Dfam_3.8/) (46).

Deep learning model for TSS prediction

CapTrap-seq and matched short-read RNA-seq datasets were used to train the model. For each 3' end of the first exon, the 10 kb DNA sequence on either side, along with the corresponding RNA-seq coverage (including both unique-mapping and multi-mapping reads), was extracted and used as input to the model. The corresponding TSS signal from the CapTrap-seq data served as the model's label. Our model is based on the architecture of the Puffin-D model, a

315   convolutional neural network specifically designed for TSS prediction using DNA sequences

316   (25). To enable TSS prediction for each sample, we adjusted the model to extract information

317   from both the DNA sequence and RNA-seq coverage. To address the imbalance between

318   positive and negative samples during model training, we employed the KL divergence loss

319   function:

$$L = \sum_i target_i^* [ln(target_i^* + \epsilon) - ln(pred_i^* + \epsilon)]$$

The *target* represents the true TSS signal at position *i*, *pred* represents the TSS signal

predicted by the model, and ϵ is a constant set to e^{-10}.

The model utilized the Adam optimizer for parameter optimization, with a learning rate of

0.005, a batch size of 64, and a maximum of 10 training epochs. Model construction and

training were carried out using PyTorch. The distance between the predicted and actual TSS

was used to evaluate the model's performance. The example of an alternative promoter was

visualized on WashU Epigenome Browser (47).

Pipeline for the detection of TE initiated transcripts

First, RNA-seq data were aligned to the reference human genome (GRCh38) using STAR

v2.7.10b (48). Sambamba was employed to remove PCR duplicates, and deepTools was used

to extract RNA-seq coverage (49,50). StringTie v2.1.7 was used to assemble the reads into full-

length transcripts (stringtie –m 200 –c 1 -t) (51). The TSS for each assembled transcript was

then predicted using the trained deep learning model. These assembled transcripts across

different samples were subsequently merged with a custom script. Briefly, the first exons of

multi-exon transcripts were extracted. Following this, first exons overlapping with other

internal exons were excluded, as they may have resulted from RNA degradation. Then, for each

first exon, the TSS was corrected according to the score calculated as follows (52):

339
$$S_t = s_t + \sum_{k \in T} \frac{w - d_k}{w * s_k}$$

The S is the weight score for position t; s represents the number of samples supporting the

position as a TSS of the first exon; w is the window size used to calculate the weight score; T

is the set of all positions within a distance from t that is less than the window size; and d is the

distance between positions t and k.

Corrected TSSs overlapping with TEs were retained, and the full-length transcripts of TE-

initiated transcripts were merged using a reference-guided approach across samples. The count

of junction reads supporting the first exon and the expression of TE-initiated transcripts at the

transcript level were then calculated using StringTie. TE-initiated transcripts with at least one

junction read supporting the first exon were considered expressed in the sample. The count of

junction reads supporting the first exon of TE-initiated transcripts from the same TE family was

summed, and the total read count mapped to the GRCh38 genome was used for normalization

to obtain TE promoter activity at the family level.

Computational validation of TE-derived TSSs and TE promoter activity

To evaluate the precision of TE-derived TSSs identified by our pipeline, we used CapTrap-seq

or RAMPAGE data as the gold standard. A TE-derived TSS with supporting reads within 100

bp was considered positive. Plots of the CapTrap-seq and RAMPAGE signals surrounding the

TSSs were generated using deepTools. To assess the precision of TE promoter activity

calculated by our pipeline, the Spearman correlation with CAGE data was calculated.

Annotation and expression analysis of TE-initiated transcripts

Identified TE-initiated transcripts were annotated with Dfam database by using BEDTools

v2.31.1 (46,53). Based on the relationship with reference transcripts, TE-initiated transcripts

were classified into three categories: annotated, chimeric, and intergenic. Coding potential of

TE-initiated transcripts was calculated by CPC2 (54). For coding TE-initiated transcripts, we

subsequently compared predicted CDS with annotated CDS using BLAST 2.14.1 (55). Then

the coding TE-initiated transcripts were annotated into five classes: annotated, 5' chimeric

normal, 5' chimeric truncated, 5' truncated and novel. Tumor-specific scores for each TE-

initiated transcript were calculated as follows:

$$S = \frac{(T_e + 1)/T_t}{(N_e + 1)/N_t}$$

The S represents the tumor-specific score for the TE-initiated transcript. T_e represents the

number of tumor samples expressing the TE-initiated transcript, N_e represents the number of

tumor-adjacent or normal tissue samples expressing the TE-initiated transcript (excluding

testicular tissue), and T_t and N_t represent the total number of tumor and normal samples,

respectively. A TE-initiated transcript with a tumor-specific score greater than eight was

considered tumor-specific expressed.

The reactome pathway enrichment analysis of gene-chimeric TE-initiated transcripts was

conducted by the ReactomePA package (56). The TE family enrichment analysis was conducted

using a two-tailed Fisher's exact test. All P-values were corrected using the Benjamini-

Hochberg (BH) procedure.

Clinically associated TE families or TE-initiated transcripts

The Wilcoxon rank-sum test was used to assess statistical differences in the expression levels of TE families and TE-initiated transcripts between tumor and adjacent normal tissues. The Kruskal-Wallis test was applied to evaluate statistical differences in the expression levels of TE families and TE-initiated transcripts across patients with different tumor stages. The survdiff function from the survival package was used to perform the Log-Rank test, comparing differences between two survival curves, and calculating the hazard ratio (HR) and P-value. All P-values were adjusted using the Benjamini-Hochberg (BH) procedure.

Construction of the TEITbase website

The backend application and frontend interface of the TEITbase website were developed using the Django framework, with Gunicorn serving as the WSGI HTTP server to execute the Django code. NGINX was employed as a reverse proxy server to forward client requests to Gunicorn. To ensure system stability, Supervisor was used to monitor the operation of the Gunicorn and Django processes. The frontend interface was designed and implemented using HTML, JavaScript, and CSS. Additionally, annotations for TE families and TE initiated transcripts were securely stored in an SQLite3 database.

Declarations

Ethics approval and consent to participate

Not applicable.

Availability of data and materials

Details about data analyzed in this study were included in the Methods section. The modified pipeline, together with the deep learning model for TSS prediction developed in this study are available in (https://github.com/yunzhang1998/Deep-TEIRI).

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by grant from the Young Scientists Fund of the National Natural Science Foundation of China (32400534), the China Postdoctoral Science Foundation (2024M760136), and the Ministry of Science and Technology of China (2021ZD0203203).

Author contributions

E.Y. conceived and designed the study. Y.Z., J.Q.S., X.Y.H., Y.Q.J., C.Y.T., and M.H.D. analyzed the data. Y.Z., J.Q.S., M.H.D., and E.Y. interpreted the data and wrote the manuscript. All authors approved the final version of the manuscript.

Acknowledgments

We would like to express our gratitude to the TCGA and GTEx committees for their valuable contributions. We also acknowledge the ENCODE Consortium and the ENCODE production laboratories.

Additional files

Additional file 1: Table S1-7. Supplementary tables (Table S1-S7, XLSX).

References

1. Britten RJ, Kohne DE. Repeated sequences in DNA. Hundreds of thousands of copies of DNA sequences have been incorporated into the genomes of higher organisms. *Science*. 1968;161(3841):529-540. doi:10.1126/science.161.3841.529

2. Kapitonov VV, Jurka J. A universal classification of eukaryotic transposable elements implemented in Repbase. *Nat Rev Genet*. 2008;9(5):411-412; author reply 414. doi:10.1038/nrg2165-c1

3. Wells JN, Feschotte C. A Field Guide to Eukaryotic Transposable Elements. *Annu Rev Genet*. 2020;54:539. doi:10.1146/annurev-genet-040620-022145

4. Beck CR, Collier P, Macfarlane C, et al. LINE-1 retrotransposition activity in human genomes. *Cell*. 2010;141(7):1159-1170. doi:10.1016/j.cell.2010.05.021

5. Brouha B, Schustak J, Badge RM, et al. Hot L1s account for the bulk of retrotransposition in the human population. *Proc Natl Acad Sci U S A*. 2003;100(9):5280-5285. doi:10.1073/pnas.0831042100

6. Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements: from conflicts to benefits. *Nat Rev Genet*. 2016;18(2):71. doi:10.1038/nrg.2016.139

7. Bourque G, Leong B, Vega VB, et al. Evolution of the mammalian transcription factor binding repertoire via transposable elements. *Genome Res*. 2008;18(11):1752-1762. doi:10.1101/gr.080663.108

8. Merlotti A, Sadacca B, Arribas YA, et al. Noncanonical splicing junctions between exons and transposable elements represent a source of immunogenic recurrent neo-antigens in patients with lung cancer. *Sci Immunol*. 2023;8(80):eabm6359. doi:10.1126/sciimmunol.abm6359

9. Miao B, Fu S, Lyu C, Gontarz P, Wang T, Zhang B. Tissue-specific usage of transposable element-derived promoters in mouse development. *Genome Biol*. 2020;21(1):255. doi:10.1186/s13059-020-02164-3

10. Deniz Ö, Frost JM, Branco MR. Regulation of transposable elements by DNA modifications. *Nat Rev Genet*. 2019;20(7):417-431. doi:10.1038/s41576-019-0106-6

11. Kong Y, Rose CM, Cass AA, et al. Transposable element expression in tumors is
associated with immune infiltration and increased antigenicity. *Nat Commun*.
2019;10(1):5228. doi:10.1038/s41467-019-13035-2

12. Rodriguez-Martin B, Alvarez EG, Baez-Ortega A, et al. Pan-cancer analysis of whole
genomes identifies driver rearrangements promoted by LINE-1 retrotransposition. *Nat
Genet*. 2020;52(3):306-319. doi:10.1038/s41588-019-0562-0

13. Modzelewski AJ, Shao W, Chen J, et al. A mouse-specific retrotransposon drives a
conserved Cdk2ap1 isoform essential for development. *Cell*. 2021;184(22):5541-
5558.e22. doi:10.1016/j.cell.2021.09.021

14. Babaian A, Mager DL. Endogenous retroviral promoter exaptation in human cancer. *Mob
DNA*. 2016;7:24. doi:10.1186/s13100-016-0080-x

15. Jang HS, Shah NM, Du AY, et al. Transposable elements drive widespread expression of
oncogenes in human cancers. *Nat Genet*. 2019;51(4):611-617. doi:10.1038/s41588-019-
0373-3

16. Shi Q, Li X, Liu Y, Chen Z, He X. FLIBase: a comprehensive repository of full-length
isoforms across human cancers and tissues. *Nucleic Acids Res*. 2023;52(D1):D124-D133.
doi:10.1093/nar/gkad745

17. Attig J, Pape J, Doglio L, et al. Human endogenous retrovirus onco-exaptation counters
cancer cell senescence through calbindin. *J Clin Invest*. 2023;133(14):e164397.
doi:10.1172/JCI164397

18. Sun Z, Zhang R, Zhang X, et al. LINE-1 promotes tumorigenicity and exacerbates tumor
progression via stimulating metabolism reprogramming in non-small cell lung cancer.
Mol Cancer. 2022;21:147. doi:10.1186/s12943-022-01618-5

19. Laumont CM, Vincent K, Hesnard L, et al. Noncoding regions are the main source of
targetable tumor-specific antigens. *Sci Transl Med*. 2018;10(470):eaau5516.
doi:10.1126/scitranslmed.aau5516

20. Liang Y, Qu X, Shah NM, Wang T. Towards targeting transposable elements for cancer
therapy. *Nat Rev Cancer*. 2024;24(2):123-140. doi:10.1038/s41568-023-00653-8

21. Ng KW, Boumelha J, Enfield KSS, et al. Antibodies against endogenous retroviruses
promote lung cancer immunotherapy. *Nature*. 2023;616(7957):563-573.
doi:10.1038/s41586-023-05771-9

22. Lei X, Mao S, Li Y, et al. ERVcancer: a web resource designed for querying activation of
human endogenous retroviruses across major cancer types. *J Genet Genomics Yi Chuan
Xue Bao*. Published online September 10, 2024:S1673-8527(24)00241-8.
doi:10.1016/j.jgg.2024.09.004

23.   Stricker E, Peckham-Gregory EC, Scheurer ME. CancerHERVdb: Human Endogenous
Retrovirus (HERV) Expression Database for Human Cancer Accelerates Studies of the
Retrovirome and Predictions for HERV-Based Therapies. *J Virol*. 2023;97(6):e0005923.
doi:10.1128/jvi.00059-23

24.   Gu X, Wang M, Zhang XO. TE-TSS: an integrated data resource of human and mouse
transposable element (TE)-derived transcription start site (TSS). *Nucleic Acids Res*.
2023;52(D1):D322. doi:10.1093/nar/gkad1048

25.   Dudnyk K, Cai D, Shi C, Xu J, Zhou J. Sequence basis of transcription initiation in human
genome. *Science*. 2024;384(6694):eadj0116. doi:10.1126/science.adj0116

26.   Zhou J, Zhang B, Li H, et al. Annotating TSSs in Multiple Cell Types Based on DNA
Sequence and RNA-seq Data via DeeReCT-TSS. *Genomics Proteomics Bioinformatics*.
2022;20(5):959-973. doi:10.1016/j.gpb.2022.11.010

27.   Carbonell-Sala S, Perteghella T, Lagarde J, et al. CapTrap-seq: a platform-agnostic and
quantitative approach for high-fidelity full-length RNA sequencing. *Nat Commun*.
2024;15(1):5278. doi:10.1038/s41467-024-49523-3

28.   Özata DM, Yu T, Mou H, et al. Evolutionarily conserved pachytene piRNA loci are highly
divergent among modern humans. *Nat Ecol Evol*. 2020;4(1):156-168.
doi:10.1038/s41559-019-1065-1

29.   ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the
human genome. *Nature*. 2012;489(7414):57-74. doi:10.1038/nature11247

30.   Harrow J, Frankish A, Gonzalez JM, et al. GENCODE: the reference human genome
annotation for The ENCODE Project. *Genome Res*. 2012;22(9):1760-1774.
doi:10.1101/gr.135350.111

31.   Liu Y, Sun J, Zhao M. ONGene: A literature-based database for human oncogenes. *J
Genet Genomics*. 2017;44(2):119-121. doi:10.1016/j.jgg.2016.12.004

32.   Hur K, Cejas P, Feliu J, et al. Hypomethylation of long interspersed nuclear element-1
(LINE-1) leads to activation of proto-oncogenes in human colorectal cancer metastasis.
Gut. 2014;63(4):635-646. doi:10.1136/gutjnl-2012-304219

33.   de la Peña Avalos B, Tropée R, Duijf PHG, Dray E. EYA4 promotes breast cancer
progression and metastasis through its role in replication stress avoidance. *Mol Cancer*.
2023;22(1):158. doi:10.1186/s12943-023-01861-4

34.   Li B, Qi ZP, He DL, et al. NLRP7 deubiquitination by USP10 promotes tumor progression
and tumor-associated macrophage polarization in colorectal cancer. *J Exp Clin Cancer
Res CR*. 2021;40(1):126. doi:10.1186/s13046-021-01920-y

35.   Gu Z, Liu Y, Zhang Y, et al. Silencing of LINE-1 retrotransposons is a selective

dependency of myeloid leukemia. *Nat Genet*. 2021;53(5):672-682. doi:10.1038/s41588-
021-00829-8

36.   Ghislat G, Cheema AS, Baudoin E, et al. NF-κB-dependent IRF1 activation programs
530   cDC1 dendritic cells to drive antitumor immunity. *Sci Immunol*. 2021;6(61):eabg3570.
531   doi:10.1126/sciimmunol.abg3570

532   37.   Avsec Ž, Agarwal V, Visentin D, et al. Effective gene expression prediction from sequence
533   by integrating long-range interactions. *Nat Methods*. 2021;18(10):1196-1203.
534   doi:10.1038/s41592-021-01252-x

535   38.   Bogard N, Linder J, Rosenberg AB, Seelig G. A deep neural network for predicting and
536   engineering   alternative   polyadenylation.   *Cell*.   2019;178(1):91-106.
537   doi:10.1016/j.cell.2019.04.046

538   39.   Fu ZC, Gao BQ, Nan F, Ma XK, Yang L. DEMINING: A deep learning model embedded
539   framework to distinguish RNA editing from DNA mutations in RNA sequencing data.
540   *Genome Biol*. 2024;25(1):258. doi:10.1186/s13059-024-03397-2

541   40.   Jaganathan K, Kyriazopoulou Panagiotopoulou S, McRae JF, et al. Predicting splicing
542   from   primary   sequence   with   deep   learning.   *Cell*.   2019;176(3):535-548.
543   doi:10.1016/j.cell.2018.12.015

544   41.   Pachganov S, Murtazalieva K, Zarubin A, Sokolov D, Chartier DR, Tatarinova TV.
545   TransPrise: a novel machine learning approach for eukaryotic promoter prediction. *PeerJ*.
546   2019;7:e7990. doi:10.7717/peerj.7990

547   42.   Shahmuradov IA, Umarov RK, Solovyev VV. TSSPlant: a new tool for prediction of plant
548   Pol II promoters. *Nucleic Acids Res*. 2017;45(8):e65. doi:10.1093/nar/gkw1353

549   43.   Lanciano S, Cristofari G. Measuring and interpreting transposable element expression.
550   *Nat Rev Genet*. 2020;21(12):721-736. doi:10.1038/s41576-020-0251-y

551   44.   Cancer Genome Atlas Research Network. Comprehensive genomic characterization
552   defines human glioblastoma genes and core pathways. *Nature*. 2008;455(7216):1061-
553   1068. doi:10.1038/nature07385

554   45.   GTEx Consortium, Laboratory, Data Analysis &Coordinating Center (LDACC)—
555   Analysis Working Group, Statistical Methods groups—Analysis Working Group, et al.
556   Genetic effects on gene expression across human tissues. *Nature*. 2017;550(7675):204-
557   213. doi:10.1038/nature24277

558   46.   Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. The Dfam community resource of
559   transposable element families, sequence models, and genome annotations. *Mob DNA*.
560   2021;12(1):2. doi:10.1186/s13100-020-00230-y

47.   Zhou X, Wang T. Using the Wash U Epigenome Browser to examine genome-wide

sequencing data. *Curr Protoc Bioinforma*. 2012;Chapter 10:10.10.1-10.10.14.
doi:10.1002/0471250953.bi1010s40

48. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics. 2012;29(1):15. doi:10.1093/bioinformatics/bts635

49. Ramírez F, Ryan DP, Grüning B, et al. deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res*. 2016;44(W1):W160-165.
doi:10.1093/nar/gkw257

50. Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. Sambamba: fast processing of NGS
alignment formats. *Bioinformatics*. 2015;31(12):2032-2034.
doi:10.1093/bioinformatics/btv098

51. Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. Transcriptome
assembly from long-read RNA-seq alignments with StringTie2. *Genome Biol*.
2019;20(1):278. doi:10.1186/s13059-019-1910-1

52. Tang AD, Soulette CM, van Baren MJ, et al. Full-length transcript characterization of
SF3B1 mutation in chronic lymphocytic leukemia reveals downregulation of retained
introns. *Nat Commun*. 2020;11(1):1438. doi:10.1038/s41467-020-15171-6

53. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinforma Oxf Engl*. 2010;26(6):841-842. doi:10.1093/bioinformatics/btq033

54. Kang YJ, Yang DC, Kong L, et al. CPC2: a fast and accurate coding potential calculator
based on sequence intrinsic features. *Nucleic Acids Res*. 2017;45(W1):W12-W16.
doi:10.1093/nar/gkx428

55. Camacho C, Coulouris G, Avagyan V, et al. BLAST+: architecture and applications. *BMC*
Bioinformatics. 2009;10:421. doi:10.1186/1471-2105-10-421

56. Yu G, He QY. ReactomePA: an R/Bioconductor package for reactome pathway analysis
and visualization. *Mol Biosyst*. 2016;12(2):477-479. doi:10.1039/c5mb00663e

Figure legends

Fig. 1 Detection of TE-initiated transcripts

(**A**) Pipeline designed to identify TE-initiated transcripts by using a deep learning model. (**B**)

RAMPAGE and CapTrap-seq signals of the *PRKACB* gene, along with TSS predictions of our

method. (**C**) The distances to the true TSSs of predicted TSSs from our method (left) and

Stringtie (right). (**D**) Prediction score and distance of predicted TSSs to the true TSSs. (**E**)

CapTrap-seq and CAGE signals around the TE-derived TSSs identified by our method or

DeeReCT-TSS (left) and those identified in this work, TE-TSS database, or both (right) in H1

cell lines. (**F**) RAMPAGE signals around the TE-derived TSSs identified by our method or

DeeReCT-TSS (left) and those identified in this work, TE-TSS database, or both (right) in testis

tissues. (**G**) Spearman correlation coefficient between CAGE data and the quantitative results

from our method or TEtranscripts (left). The Kimura value of high-accuracy TE families and

others (right).

Fig. 2 Tumor-specific TE-initiated transcription promotes tumorigenesis

(**A**) The distribution of the activated TE loci across the TE classes. (**B**) The distribution of the

neighboring genes across the gene type. (**C**) The reactome pathway enrichment analysis of

neighboring genes associated with the tumor-specific TE-initiated transcripts. (**D**) The TE

family enrichment analysis of tumor-specific TE-initiated transcripts. (**E**) The 10 TE-initiated

transcripts with the highest tumor-specific scores are presented. The left-most panel shows the

TE-gene along with the tumor-specific score. The next two panels display the number of tumor

samples in which each TE-initiated transcript is expressed and the distribution across cancer

types. The final two panels show the expression levels of the TE-initiated transcript and the

percentage of total oncogene expression contributed by TE-initiated transcription. (**F**) Kaplan–

Meier plots demonstrating overall survival associated with the expression of TE-initiated

transcripts.

Fig. 3 Applications of TEITbase

(**A**) The homepage of TEITbase. (**B**) Box plot showing the promoter activity of L1HS in TCGA tumors (top) and GTEx normal tissues (bottom). (**C**) Box plot showing the promoter activity of L1HS in LUSC and adjacent normal tissues. (**D**) Box plot illustrating the promoter activity of L1HS across different stages of LUSC. (**E**) Kaplan–Meier plots showing overall survival associated with the promoter activity of L1HS in LUSC. (**F**) The reactome pathway enrichment analysis of genes positively correlated with the promoter activity of L1HS in LUSC. (**G**, **H**) Proportion of TCGA tumor samples (**G**) and GTEx normal tissue samples (**H**) expressing TE_11880. (**I**) Box plot showing the expression level of TE_11880 in LUSC, STAD and adjacent normal tissues. (**J**) Box plot illustrating the expression level of TE_11880 across different stages of STAD. (**K**) Kaplan–Meier plots showing overall survival associated with the expression of TE_11880 in COAD.