Comparative Analysis of Deep Learning Models for Predicting Causative Regulatory Variants

Gaetano Manzo[1], Kathryn Borkowski[1,2] and Ivan Ovcharenko[1,*]

[1]Computational Biology Branch, Division of Intramural Research, National Library of Medicine, National Institutes of Health, 8600 Rockville Pike, Bethesda, MD, USA.,

[2]Case Western Reserve University, 10900 Euclid Ave., Cleveland, OH, USA.

[*]Corresponding author

Email: ovcharen@nih.gov

Abstract

Motivation: Genome-wide association studies (GWAS) have identified numerous noncoding variants associated with complex human diseases, disorders, and traits. However, resolving the uncertainty between GWAS association and causality remains a significant challenge. The small subset of noncoding GWAS variants with causative effects on gene regulatory elements can only be detected through accurate methods that assess the impact of DNA sequence variation on gene regulatory activity. Deep learning models, such as those based on Convolutional Neural Networks (CNNs) and transformers, have gained prominence in predicting the regulatory effects of genetic variants, particularly in enhancers, by learning patterns from genomic and epigenomic data. Despite their potential, selecting the most suitable model is hindered by the lack of standardized benchmarks, consistent training conditions, and performance evaluation criteria in existing reviews.

Results: This study evaluates state-of-the-art deep learning models for predicting the effects of genetic variants on enhancer activity using nine datasets stemming from MPRA, raQTL, and eQTL experiments, profiling the regulatory impact of 54,859 SNPs across four human cell lines. The results reveal that CNN models, such as TREDNet and SEI, consistently outperform other architectures in predicting the regulatory impact of SNP. However, hybrid CNN-transformer models, such as Borzoi, display superior performance in identifying causal SNPs within a linkage disequilibrium block. While fine-tuning enhances the performance of transformer-based models, it remains insufficient to surpass CNN and hybrid models when evaluated under optimized conditions.

Introduction

Genome-wide association studies (GWAS) have revealed that around 95% of disease-associated genetic variants occur in non-coding regions of the human genome, with causative variants commonly affecting regulatory elements that modulate gene expression [1, 2, 3]. These regulatory variants can profoundly affect phenotypes and alter disease susceptibility via dysregulation of their target genes [4].

Deep learning models have emerged as powerful tools for predicting the regulatory effects of genetic variants, particularly in enhancers. These methods leverage high-throughput genomic and epigenomic data to learn DNA sequence patterns associating sequence features with regulatory activity. For instance, convolutional neural networks (CNNs) have been used to identify variants that disrupt transcription factor binding or chromatin accessibility, providing insights into their potential phenotypic impact (e.g., DeepSEA [5], SEI [6], and TREDNet [7]). Similarly, modern, large transformer-based models can very accurately predict cell-type-specific regulatory effects, such as DNA methylation changes and fine-mapping of disease-associated loci; these primarily include DNABERT series [8, 9], Nucleotide Transformer series [10, 11], and Enformer [12].

Selecting the most suitable model for detecting the regulatory effects of genetic variants remains a significant challenge despite several surveys offering detailed overviews of the deep learning ecosystem in this domain [13, 14, 15]. These surveys have highlighted the unique strengths and limitations of various models. However, they lack a unified

framework for assessment. Specifically, existing reviews often fail to benchmark

models on a standardized dataset, train or fine-tune them under consistent conditions,

and evaluate their performance using uniform criteria. Furthermore, there is a

fundamental difference between benchmarking models on regulatory regions versus

regulatory variants, which is rarely addressed. While regulatory region analyses focus

on identifying broader functional elements, regulatory variant assessments require

evaluating the impact of specific sequence alterations within these regions, presenting

distinct challenges and opportunities for model evaluation.

In this study, we evaluate state-of-the-art deep learning models to predict the effects of

genetic variants on enhancer activity in the human genome. Our approach involved

curating and integrating nine datasets derived from diverse experimental

methodologies, including massively parallel reporter assay (MPRA), reporter assay

quantitative trait loci (raQTL), and expression quantitative trait loci (eQTL) studies.

These datasets encompass 54,859 single-nucleotide polymorphisms (SNPs) in enhancer

regions across four human cell lines.

We tackle three tasks: predicting log2-fold changes in enhancer activity, classifying

SNPs by regulatory impact, and identifying causal SNPs within LD blocks. We also

assess how data variability affects model performance, emphasizing the importance of

data quality. Finally, we illustrate how enhancer detection models can be effectively

leveraged for variant effect assessment, highlighting their practical utility in

interpreting genomic data and prioritizing candidate regulatory variants.

To ensure a robust and thorough assessment, we fine-tuned 22 deep learning models

for each cell line, systematically exploring a broad spectrum of architectural designs

and parameter configurations. Our results indicate that CNN models outperform more

"advanced" architectures, such as transformers, on causative regulatory variant

detection. However, fine-tuning significantly boosts the performance of transformer-

based architectures, revealing their potential to surpass CNNs under optimized

conditions.

Results

Comparison of Enhancer Variant Prediction Models

We assessed several state-of-the-art deep learning models for their ability to predict the

effects of genetic variants on enhancer activity in humans. The selected models differ

significantly, not only in terms of architecture and parameters but also in the type of

data used for training or pre-training. More importantly, these models were originally

designed for different purposes: some, like Nucleotide Transformer [10] and

DNABERT2 [9], were developed for a broad range of tasks, while others, such as

Geneformer [13] and TREDNet [7], were tailored for more specific applications.

Crucially, this means we are not only comparing model architectures but also the

underlying training datasets, which introduces additional complexity when

benchmarking their performance. To further disentangle these effects, we also evaluate

how different datasets impact the predictions of the same model. Despite these

differences, both model categories have been adapted to predict the effects of enhancer

variants (see Materials and Methods). Our evaluation encompasses a range of

architectural and parameter configurations across model families, focusing on 54,859

single-nucleotide polymorphisms located in enhancer regions from nine datasets

originating from four cell lines: K562, HepG2, NPC, and HeLa.

We performed regression analyses between model predictions and experimental log2-

fold changes for variant effects in enhancers across the human genome sequence

(Pearson Correlation Fig. 1, Spearman Correlation Fig. S1, Materials and Methods).

CNN-based models, such as TREDNet and SEI, demonstrate the best alignment with

experimentally recorded variant effects, achieving Pearson correlation coefficients of

0.297 and 0.276, respectively, while the most accurate transformer-based model,

Nucleotide Transformer v2, scores only at 0.105, well below the level of statistical

uncertainty. These findings highlight the effectiveness of CNNs in recognizing spatial

and structural genomic patterns essential for predicting enhancer variant effects.

Convolutional layers identify genomic features, such as local motifs and epigenetic

markers, that influence genetic variant effects [16].

Fig 1: Pearson correlation coefficients between model predictions and experimental log2-fold difference for enhancer variant effects across the human genome. Bar colors denote model architectures (CNN: green, transformer: blue, and hybrid: orange). All correlations have p-values < 0.05, and error bars show variance.

This capability is crucial for capturing the complex genomic relationships underlying

gene regulation and enhancer activity.

In contrast, transformer-based models fine-tuned for specific cell lines show lower

correlation values (0.019–0.105). Despite their ability to model long-range

relationships and complex sequences, they struggle to detect the subtle genomic

patterns linked to regulatory variant effects.

This challenge likely stems from the complexity of genomic data, lower granularity in

feature detection compared to CNNs, and the extensive data required to fully exploit

transformer capabilities.

To bridge the gap between local pattern recognition and long-range dependency

modeling, hybrid models combine components such as CNNs, transformers, and

LSTMs. Models like the HyenaDNA series and Borzoi show moderate correlations,

with HyenaDNA medium-160k-seqlen-hf achieving the highest among hybrids at

0.124. The integration of diverse components enhances performance compared to

transformer-based models while optimizing computational costs relative to CNN-based

models. However, despite their architectural versatility, these hybrid models do not

surpass the performance of CNN-based models, highlighting a trade-off between

flexibility and focused efficiency (Fig. 1, Fig. S1).

We acknowledge that our classification groups together all models that are not purely

CNN- or transformer-based, despite significant architectural differences—particularly

in the case of HyenaDNA, which diverges substantially from attention-based models.

Nonetheless, our primary objective is to establish a broad guideline for enhancer variant

prediction models, rather than to define rigid model categories. The classification task

 provides a deeper understanding of model performance by evaluating true positive and

 true negative rates in predicting variants that upregulate or downregulate gene

 expression (Fig. 2). Among 54,859 experimental SNPs, approximately half are linked

 to upregulation and half to downregulation.

Fig 2: Heatmap of model variant predictions (Predicted) versus experimental values (Actual Results). The color intensity represents the fraction of values predicted as positive (negative) relative to the experimental positive (negative) values. Red indicates higher fractions (desired in the first two columns from the left), while blue indicates lower fractions (desired in the follow two columns). Accuracy and F1-score are indicated in the last two columns.

 CNN-based models, like TREDNet and SEI, excel in predicting variant effects.

 TREDNet achieves balanced performance with correctly predicted upregulation (0.60)

 and downregulation (0.60) rates, while SEI leads in upregulation detection (0.62) but

 struggles with downregulation (0.56). Transformer-based models show mixed results.

 Nucleotide Transformer v2-500m-multi-species matches SEI's upregulation prediction

 rate (0.62) but underperforms in downregulation (0.52). DNABERT2 performs steadily

 (0.54, 0.55), while Geneformer shows the weakest outcomes. Borzoi achieves a strong

155 upregulation prediction rate (0.61) but falters in downregulation (0.47). Enformer

156 shows moderate, balanced results (0.58, 0.55), while HyenaDNA remains consistent

157 but unremarkable.

Fig 3: Percentage of top-scoring causal SNPs predicted within the Linkage Disequilibrium (LD) block by various models. Models are grouped by architecture: CNN (green), transformer (blue), and hybrid (orange). Bars are sorted by top-1 performance, with decreasing transparency indicating lower ranks (top-2 and top-3 results are progressively faded compared to top-1).

A crucial aspect of each model is its ability to identify causal SNPs within their

respective Linkage Disequilibrium (LD) blocks. Essentially, this is the most important

function of variant classification models, which can be directly applied to the resolution

of causal variants within large blocks of associated variants identified in GWAS studies.

To investigate this, we curated a dataset of 10,098 causal SNPs specific to the HepG2

cell line. For each causal SNP, we identified associated LD-block SNPs with an r^2-

value above 0.8 within a 500 kbp window, resulting in 263,286 SNPs. With this set, we

can directly assess each method's ability to retrieve the causal SNP within this ~26:1

pool of associated SNPs. For each identified SNP, we generated two sequences of 1

kbp: one containing the SNP and the other containing the reference nucleotide. In both

cases, the SNP or reference nucleotide were positioned at the center of the sequence.

These sequences were then input into several models to assess their ability to predict

causal SNPs by calculating the log2 foldchange between the scores of the alternative

sequence (containing the SNP) and the reference sequence (Methods and Materials

sections). This procedure was repeated for all LD-block associated SNPs and the log2

fold-change score of the known causal SNP was compared to its associated variants.

Finally, we computed the percentage of causal SNPs within an LD-block correctly

predicted as causal based on their score being either the highest ("top-1" test) or within

the top two and three highest scores ("top-2" and "top-3" tests).

The Borzoi model achieved the highest accuracy, correctly identifying 42.5% of causal

SNPs, followed by the SEI, Enformer, and TREDNet models, with 40.7%, 39.4%, and

33.8% in the top1 test, respectively, (Fig. 3). The causal variant detection rate increased

substantially in the top-2 and top-3 test with Borzoi (44.9%, 48.9), SEI (43.6%, 47.0%),

Enformer (42.4%, 46.2%), and TREDNet (36.1%, 40.1%). Among the transformer-

based models, the Gena LM bert-based-t2t-multi demonstrated the best performance

(top-1 30.4%, top-2 32.7%, and top-3 37.5%), marginally surpassing other Nucleotide

Transformer and Gena LM variants, which achieved accuracies between 28.0% and

30.0% in top-1 test. The HyenaDNA models, categorized as hybrid architectures,

displayed moderate performance, with accuracies ranging from 24.3% to 26.5% in top-

1 test, and larger sequence lengths, such as those used in the HyenaDNA large model,

led to improved results. The Geneformer model, however, had the lowest accuracy,

identifying 24.3% of causal SNPs in top-1 test. Overall, hybrid and CNN architectures

outperformed transformer-based models in this application, underscoring their

suitability for tasks involving causal SNP detection.

Overall, models like TREDNet and SEI consistently outperform others in both

regression analysis and classification tasks, showcasing their strength in capturing local

genomic patterns critical for regulatory variant effect prediction. Hybrid models, such

as Borzoi and Enformer, offer balanced performance, bridging the gap between local

pattern recognition and long-range dependency modeling but fail to surpass CNNs in

overall accuracy. Yet, these models excel at accurate detection of causal variants within

LD-blocks of associated SNPs. Transformer-based models, despite their potential to

model complex sequences and long-range interactions, struggle to match the

granularity of CNNs and hybrids.

Impact of Certainty in Experimental Results

The results of experimental assays of regulatory variants depend on the selected degree

of certainty in separating significant and insignificant variant effects. To investigate the

impact of experimental data significance on correlation with modeling results, we

analyzed the top models for each architecture type: TREDNet (CNN), HyenaDNA

(hybrid), and Nucleotide Transformer (transformer). The experimental data was binned

using the following significance thresholds, $p < 0.5$, $p < 0.1$, $p < 5 \cdot 10^{-5}$, $p < 10^{-5}$.

The results demonstrate that models align better with experimental findings when

applied to variants with higher statistical significance. For instance, data points with

greater experimental confidence show increased clustering in the first and third

quadrants, representing cases where both experimental and predicted results are either

positive (upregulation) or negative (downregulation) (Fig. 4, panels A-C, Figures S2,

S3a, S3b, S3c, and S3d).

Fig 4: Model performance relative to experimental data significance, using top models from each architecture category. Panels A-C show scatter plots for TREDNet, HyenaDNA, and Nucleotide Transformer, with the correlation between model predictions and experimental results as significance improves. Panel D compares model predictions for highly significant variants. Panel E presents heatmaps of classification metrics at different p-value thresholds, highlighting performance variations in identifying positive/negative outcomes across

Among the evaluated architectures, HyenaDNA exhibited the greatest improvement in

predictive accuracy with the increase of the statistical significance of experimental

results (from all data 0.124 to 0.226 with $p < 10^{-5}$, Fig. 4, panel C). However, TREDNet

consistently displayed superior overall performance, with a higher density of points

aligning with experimental outcomes (from all data 0.298 to 0.421 with $p < 10^{-5}$, Fig.

4, panel B). This trend is further illustrated in the density plots, where TREDNet

exhibits increased clustering in the first and third quadrants, demonstrating stronger

correlations between model predictions and experimental values compared to both

hybrid and transformer models (Fig. 4, panel D). Moreover, the correlation coefficients

across varying significance thresholds highlight TREDNet's robustness, achieving the highest values under all conditions. True positives and true negatives increase with the significance of the experimental results (positive/positive 0.63, negative/negative 0.66, with $p < 0.05$; positive/positive 0.72, negative/negative 0.78, with $p < 10^{-5}$), with the hybrid model and the transformer model following in performance (Fig. 4, panel E). The rest of models' performance relative to experimental data significance can be found in Fig. S2, and Figures S3a, S3b, S3c, and S3d.

Given the strong cell-line specificity of enhancers (Wu and Huang, 2024), determining whether model performance depends on the type of cell line used is essential for understanding their predictive capabilities and limitations. For this, we evaluated whether a model fine-tuned for a specific cell line demonstrates superior performance compared to models from other architectures, such as CNNs, which often rely on generalization rather than cell-line-specific adaptation. Indeed, transformer-based models, pre-trained on extensive datasets and fine-tuned for targeted tasks, can offer both flexibility and accuracy, making them well-suited for capturing the unique regulatory landscape of cell lines. In contrast, CNN-based models, such as TREDNet and SEI, though highly effective for specific tasks, could require additional steps, such as retraining or the addition of specialized layers, to adapt their outputs for cell-line-specific predictions. However, the results indicate that TREDNet and SEI remain the most effective models for predicting variant effects across different cell lines, even after model fine-tuning (Pearson Correlation Table 1, Spearman Correlation Sup Table 1, Dataset-wise Sup Table 2 and Sup Table 3). For example, the Nucleotide Transformer v2-250m-multi-species model aligns with experimental results in K562 and HepG2 (0.1998 and 0.11) but performs less effectively in NPC and HeLa cells. This variability

253   could stem from differences in pre-training datasets—such as multi-species versus

254   human-specific data—and the fine-tuning approach.

255   **Table 1**: Pearson correlation coefficient for various deep learning models across four cell lines: K562 (19321 SNPs),
256   HepG2 (16255 SNPs), NPC (14042 SNPs), and HeLa (5241 SNPs). Standard Error is reported in brackets. Bold and
257   underline styles denote the top and second-highest correlations per cell line, respectively.

Models	Cell Line			
	K562 (19321 SNPs)	**HepG2** (16255 SNPs)	**NPC** (14042 SNPs)	**Hela** (5241 SNPs)
DNABERT2	0.086 (0.039)	0.096 (0.191)	-0.004 (0.007)	0.120 (0.140)
NT v2-50m-ms	0.147 (0.058)	0.103 (0.036)	0.020 (0.003)	0.074 (0.047)
NT v2-100m-ms	0.152 (0.065)	0.127 (0.050)	0.016 (0.003)	0.042 (0.027)
NT v2-250m-ms	0.166 (0.064)	0.110 (0.020)	0.023 (0.003)	0.030 (0.018)
NT v2 500m-ms	0.199 (0.036)	0.114 (0.007)	0.028 (0.003)	0.036 (0.011)
NT 500m1000g	0.123 (0.084)	0.116 (0.023)	0.001 (0.002)	0.022 (0.018)
NT 500m-h-ref	0.149 (0.139)	0.119 (0.038)	0.013 (0.001)	0.067 (0.043)
NT 2.5b-1000g	0.147 (0.052)	0.112 (0.021)	0.005 (0.003)	0.087 (0.105)
NT 2.5b-m-s	0.153 (0.055)	0.139 (0.027)	0.027 (0.001)	0.066 (0.027)
Geneformer	0.005 (0.198)	0.027 (0.160)	-0.002 (0.138)	-0.007 (0.057)
Gena LM-base	0.077 (0.042)	0.050 (0.033)	0.024 (0.002)	0.029 (0.227)
Gena LM large	0.117 (0.044)	0.088 (0.036)	0.030 (0.002)	0.069 (0.216)
Gena LM b-multi	0.077 (0.051)	0.062 (0.024)	0.010 (0.001)	0.026 (0.061)
Gena LM bigbird	0.136 (0.055)	0.099 (0.029)	0.021 (0.002)	0.045 (0.015)
Hyenadna 32k	0.084 (0.071)	0.098 (0.061)	0.002 (0.002)	0.031 (0.013)
Hyenadna 160k	0.148 (0.053)	0.142 (0.040)	-0.007 (0.003)	0.046 (0.018)
Hyenadna 450k	0.077 (0.074)	0.134 (0.050)	-0.007 (0.003)	0.055 (0.015)
Hyenadna 1mf	0.118 (0.056)	0.136 (0.059)	0.001 (0.003)	0.046 (0.013)
TREDNet	**0.315 (0.025)**	**0.314 (0.018)**	**0.075 (0.003)**	0.076 (0.063)
SEI	<u>0.297 (0.022)</u>	<u>0.298 (0.029)</u>	<u>0.073 (0.003)</u>	**0.280 (0.008)**
Enformer	0.059 (0.059)	0.141 (0.110)	0.008 (0.080)	<u>0.245 (0.027)</u>
Borzoi	0.033 (0.016)	0.125 (0.025)	0.023 (0.004)	-0.066 (0.005)

Notably, the hybrid model Enformer, which integrates features from transformers and

CNNs, performs particularly well in HeLa cells (0.245), ranking second only to SEI

(0.279). As observed in the previous test, CNN-based models demonstrate superior

performance, showcasing robustness across cell-line variants. We also assess the

impact of the certainty of experimental results, SEI, which does not require training or

fine-tuning for specific cell lines, outperforms fine-tuned models. TREDNet, likely

benefiting from its second-phase training tailored to the specific cell line, achieves the

best overall performance, further highlighting its ability to leverage cell-line-specific

information effectively.

Enhancer Detection Models for Variant Effect Assessment

Models designed to predict variant effects rely primarily on the change in confidence

when detecting the enhancer region after introducing the variant. Next, we investigated

whether improving the accuracy of enhancer sequence detection enhances predictions

of variant effects. While SEI, Borzoi, and Enformer do not require fine-tuning,

TREDNet, despite being a CNN-based model, incorporates a two-phase training

process. The second phase can be considered fine-tuning [7].

Transformer-based architectures, particularly Nucleotide Transformer models pre-

trained on multi-species datasets, emerge as strong performers in enhancer

classification (Fig. S4). Models such as Nucleotide Transformer v2-250m-multi-

species and Nucleotide Transformer 2.5b-multi-species consistently demonstrate high

accuracy, as indicated by their auROC and auPRC metrics (auROC: 0.967, 0.963;

auPRC: 0.958, 0.951). However, the correlation between the models' outputs and the

MPRA experimental results, which measure the fold-change effect between reference

and alternative sequences, remains relatively low (ranging from 0.112 to 0.15). SEI

stands out as the top performer in detecting enhancer activity, achieving the highest auROC (0.974) and auPRC (0.971) scores among all models. It is also the second best for the correlation between model outputs and experimental results (0.295). TREDNet demonstrates strong performance with a higher correlation (0.318), while balancing high auROC (0.872) and auPRC (0.814) scores. The HyenaDNA models show competitive classification metrics, with the Hyenadna medium-160k-seqlen-hf and Hyenadna large-1m-seqlen-hf versions attaining notable auROC (0.933, 0.904) and auPRC (0.921, 0.893) scores.

Fig 5: *Comparison of F1 score (top) and Pearson Correlation Coefficient (bottom) between fine-tuned (blue dots) and one-shot (orange dots) models across various architectures, evaluated on Dataset 3, HepG2 cell line, (14182 SNPs). Arrows indicate the changes in performance between fine-tuned and one-shot implementations for each model.*

However, their Pearson correlation coefficients are moderate (0.152, 0.151), indicating

that while these models classify enhancer activity effectively, further optimization is

needed to improve their predictive accuracy for variant-specific effects. Borzoi

performs the weakest in detecting enhancer regions (auROC: 0.672, auPRC: 0.65),

while Geneformer fares poorly in detecting variants (auROC: 0.027). Yet, this weak

enhancer detection performance doesn't translate in inaccurate detection of causal

variants, as Borzoi outperforms all other methods in this critical test.

Fine-tuned models consistently outperform their one-shot counterparts, achieving F1

scores between 0.85 and 0.92, compared to 0.70 to 0.80 for one-shot models (Fig. 5,

top). This improvement in performance is reflected in the correlation between model

predictions and experimental results, particularly in detecting enhancer variants (Fig. 5,

bottom). The performance enhancement is consistent across most architectures, with

CNN-based models showing remarkable results even without fine-tuning. However,

SEI, which is available only in its 'one-shot' implementation, achieves a high F1 score

and correlation, surpassing many fine-tuned transformer models. Additionally, some

transformer-based models, such as Nucleotide Transformer 2.5b-1000g, show no

improvement in detecting enhancer variants.

Discussion

Our evaluation of deep learning models highlights the strengths and limitations of

different architectures in predicting the effects of genetic variants on enhancer activity.

CNN-based models, such as TREDNet and SEI, outperform transformer-based and

hybrid architectures in both regression and classification tasks. Notably, Borzoi, a

hybrid model, excels in causal SNP prediction, leveraging both local and global

genomic features critical for regulatory variant effect prediction.

CNNs-, Transformers-, and Hybrid-based Models

TREDNet and SEI achieve superior performance in regression (Fig. 1) and

classification tasks (Fig. 2). TREDNet demonstrates balanced predictive capability for

both upregulated and downregulated variants, while SEI excels in detecting upregulated

variants. CNNs effectively capture local genomic patterns like motifs and epigenetic

markers, making them well-suited for enhancer activity analysis. Their simpler

architecture yields high Pearson correlation coefficients (0.297 for TREDNet, 0.276 for

SEI). However, their inability to model long-range dependencies limits performance in

tasks requiring broader sequence context.

Transformers, including the Nucleotide Transformer and DNABERT 2, capture long-

range dependencies but struggle with fine-grained regulatory signals. They

underperform in enhancer variant effect prediction, with accuracies between 28.0% and

30.0% in causal SNP identification (Fig. 3). These models require substantial training

data, and their performance could improve with expanded datasets or refined focus on

regulatory signals.

Hybrid models, such as HyenaDNA and Borzoi, integrate CNNs, transformers, and

LSTMs to balance local and long-range dependencies. While HyenaDNA shows

moderate correlation with experimental results, Borzoi outperforms other models in

causal SNP detection (Fig. 3), making it valuable for GWAS analysis. However, hybrid

models do not consistently surpass CNNs in overall predictive accuracy and should be

assessed based on task-specific requirements.

Impact of Experimental Data Quality and Model Generalization

The reliability of model predictions is closely tied to the quality of experimental data. Model alignment with experimental results improves as statistical confidence in variant effects increases (Fig. 4). TREDNet, for instance, maintains high predictive accuracy even under varying significance thresholds, emphasizing the robustness of CNN-based models. These findings suggest that enhancing experimental rigor and statistical refinement can reduce uncertainty in variant effect assessments, leading to more reliable predictions.

In addition to data quality, model generalization across different cell lines is a critical factor. Transformer models fine-tuned for specific cell lines can capture regulatory landscapes but often underperform compared to CNNs. For instance, the Nucleotide Transformer v2-250m-multi-species achieves reasonable results in K562 and HepG2 but struggles in NPC and HeLa, likely due to pre-training limitations. In contrast, CNN models generalize well across cell lines without requiring fine-tuning, further demonstrating their robustness in variant effect prediction.

Enhancer detection models, such as the Nucleotide Transformer v2-250m-multi-species, achieve high classification accuracy (auROC: 0.967, auPRC: 0.958), yet their predictions show weak correlation with experimental results (Fig. S4). CNN-based models, particularly SEI, perform well in both enhancer detection and variant effect prediction. While fine-tuning benefits transformer models, SEI, despite its 'one-shot' implementation, still outperforms many fine-tuned transformers (Fig. 5). These results highlight the advantages of CNN architectures in both enhancer activity analysis and

cross-cell-line generalization, reinforcing their suitability for regulatory variant effect

prediction.

In conclusion, our findings suggest that SEI and TREDNet are the most effective

models for predicting the effects of genetic variants on enhancer activity. Hybrid

models like Borzoi demonstrate strong potential in extracting causal SNPs from GWAS

data, showcasing their value in applications that require the integration of both local

and global sequence contexts. Transformer-based models, on the other hand, are

particularly effective at identifying enhancer regions due to their ability to capture long-

range dependencies, although they still face challenges with fine-grained genomic

features and often require substantial data for optimal performance. These results

highlight the strengths of different model architectures for various aspects of regulatory

variant prediction and underscore the need for further optimization to enhance their

performance. Future work should aim to improve feature detection in transformer

models and refine the flexibility of hybrid architectures to better capture both local and

long-range genomic dependencies.

Materials and Methods

Data Pre-Processing

The datasets from HepG2, K562, NPC, and HeLa cell lines underwent a standardized

preprocessing pipeline to ensure consistency and feature extraction for downstream

analysis. Two main pipelines were applied: one for training and testing enhancer

classifier models, such as Nucleotide Transformers and DNABERT, and another for

mutagenesis experiments. For the enhancer classification pipeline, 1kb sequences were

extracted for both enhancer (positive) and control regions.

To maintain consistency across models, all nucleotides were capitalized, as some

models are case-sensitive. Unknown nucleotides marked as "N" were removed from

the sequences. The datasets were then split into training, validation, and test sets, and

tokenized using the corresponding model's tokenizer with a maximum length of 512

tokens. Labels were generated to distinguish between positive and control sequences.

In the mutagenesis experiment pipeline, the reference and alternative nucleotide

locations were extracted for each row of the dataset. For each nucleotide of interest, a

1kb sequence was generated with the target nucleotide positioned in the center.

Chromosome and position information, along with reference and alternative alleles,

were also extracted. Log2 activity metrics, such as fold-change values or effect sizes,

were calculated to assess regulatory activity. Additionally, signed p-values were

derived from raw or adjusted p-values to capture both the statistical significance and

directionality of the regulatory effects.

The preprocessing step ensured that variations in dataset structure did not interfere with

downstream analysis. To further enhance consistency, hg19 genomic coordinates were

used for variant mapping, and enhancer activity metrics were uniformly applied across

datasets. Differences in expression or activity levels between experimental conditions,

such as mutant versus control or modern versus archaic sequences, were computed

when applicable. Ultimately, all datasets were converted into a standardized format that

included chromosome information, positions, reference and alternative alleles, log2

activity ratios, signed p-values, raw p-values, and other relevant regulatory features.
21

Model Adaptation and Fine-tuning

The objective of fine-tuning is to adapt a model originally designed as an enhancer classifier to assess the impact of SNPs on enhancer activity. By introducing variant sequences, we evaluate how nucleotide changes affect the model's score. To achieve this, we fine-tuned pre-trained DNA language models, such as Geneformer (initially built for scRNA-seq data), as enhancer classifiers. Sequences were preprocessed as described earlier, with a length of 1 kbp, balancing genomic context and model compatibility. This choice reflects the average enhancer length of 400 bp.

Most models were loaded from the Hugging Face Hub using AutoTokenizer and AutoModelForSequenceClassification. Exceptions, such as TREDNet, Enformer, SEI, and Borzoi, were sourced from GitHub and required custom pipelines for preprocessing and inference. Fine-tuning leveraged the Hugging Face transformers library for flexible training and evaluation. Models like HyenaDNA, Geneformer, Gena-LM, DNABERT2, and Nucleotide Transformers utilized the Trainer API. We fine-tuned the models on four biosample-specific datasets: HeLa (BioS2), neural progenitor cells (NPC, BioS45), HepG2 (BioS73), and K562 (BioS74). Sequences were cleaned to remove ambiguous bases (Ns) and split into training, validation, and test sets. Model-specific tokenizers processed sequences with a maximum length of 512 tokens. For non-fine-tuned models like SEI and Enformer, the same 1 kbp sequence length was used with padding as needed.

Key hyperparameters included a batch size of 8, a learning rate of 1e-5, and a maximum of 200 epochs. Early stopping with a patience of 5 epochs was applied to prevent

overfitting, with progress logged every 500 steps. Fine-tuned models were saved in

output directories named by model checkpoint and biosample ID and uploaded to the

Hugging Face Hub for reproducibility. GPU memory was cleared after each session for

resource efficiency. This workflow allowed us to adapt general-purpose DNA language

models for cell-type-specific regulatory prediction while maintaining consistency with

non-fine-tuned models. Training on NIH BioWulf node clusters with A100 GPUs

enabled efficient handling of large datasets, such as the largest set (~130,000 sequences

for positive and control enhancer regions), which completed in hours using 8 A100

GPUs.

Validation and Post-Processing

During the fine-tuning process, the models were optimized to distinguish enhancer

sequences from control sequences. In contrast, the validation process focused on

evaluating the difference between the reference enhancer sequence and the alternative

sequence, where the middle nucleotide was mutated. The difference in output scores

between these two sequences, obtained from the fine-tuned model, allowed for the

estimation of the variant's effect on the enhancer. If the output scores for both sequences

were identical, the variant had no effect. Conversely, a change in output scores

indicated either up- or down-regulation.

The prediction process involved generating regulatory effect predictions for each

dataset by calculating the log2 ratio of alternative to reference allele probabilities for

each variant. This metric provided a standardized approach to assess the relative

regulatory effects of alternative alleles compared to reference alleles. Since each fine-

tuned model produced logits for each class (enhancer or not), predictions for both

reference and alternative alleles were processed by first calculating margin logits,

which represent the difference in predicted class probabilities. These logits were then

transformed into probabilities using the sigmoid function. The log2 ratio of these

probabilities was computed to quantify the regulatory differences between the two

alleles. Additionally, precomputed predictions stored in pickle files were loaded for

specific models and datasets. The predictions for reference and alternative alleles were

used to compute a ratio, representing the alternative-to-reference prediction ratio. This

ratio was subsequently log2-transformed to ensure consistency in the output format.

Additional preprocessing steps were applied to specific datasets as needed, such as

loading specialized prediction files.

The prediction pipeline was applied across all datasets by iterating through experiments

defined in a metadata table. For each experiment, data were preprocessed, and

predictions were either computed or loaded accordingly. The results were compiled into

a unified dictionary containing log2 prediction ratios for all variants across

experiments. As in the fine-tuning process, the validation of variant/reference

sequences was performed using 1 kb sequences, ensuring consistency in genomic

context and input length across models. Several metrics were employed to evaluate

model performance, including the Area Under the Curve (AUC), Precision-Recall

Curve (PRC), F1-score, and correlation coefficients such as Pearson and Spearman.

These metrics provided a comprehensive assessment of model performance,

encompassing both classification accuracy and the correlation between predicted and

observed values.

Deep Learning Models

This study utilizes several deep learning models designed for genomic sequence

analysis, with a particular focus on applications to the Homo sapiens genome. The

models represent the forefront of innovation in deep learning for genomics, leveraging

diverse architectures and pre-training paradigms to handle the complexity of human

genomic data. To maintain focus on models explicitly designed for the human genome,

this study excluded models like EVO and GPN, which are primarily aimed at other

species or biological domains.

Model	Architecture	Training Details	Applications
DNABERT 2 [9]	Transformer-based with ALiBi and Flash Attention	Pre-trained using Byte Pair Encoding (BPE) tokenization; compact representation of sequences; optimized for computational efficiency and scalability.	Sequence classification, motif discovery, variant effect prediction across species.
Nucleotide Transformer [10, 11]	Transformer-based (50M–2.5B parameters)	Pretrained on multi-species genomes with 6-mer tokenization; trained on 300B tokens using Adam optimizer with warmup and decay schedules; supports long-range genomic dependencies.	Comparative genomics, functional annotation, variant effect prediction.
Geneformer [13]	Transformer-based with rank-based analysis	Pretrained on single-cell transcriptome datasets using masked gene prediction; optimized for noise resilience and stability in single-cell data.	Gene network dynamics analysis, single-cell state classification.
GENA-LM [17]	BERT-base, BERT-large, BigBird variants	Pretrained on genomic sequences using sparse attention mechanisms (BigBird); supports long-range sequence modeling and local feature detection; memory augmentation techniques applied for scalability.	Local feature detection, long-range genomic dependency modeling.
Enformer [12]	Convolution + Transformer hybrid	Pretrained on DNA sequences up to 200,000bp using convolutional blocks for spatial reduction and transformer layers for long-range interactions; predicts over 5,000 epigenetic features.	Gene expression prediction, regulatory element analysis.
HyenaDNA [18]	Runtime-scalable models	Pretrained on human reference genome with extended sequence lengths (32,000–1,000,000bp); optimized for runtime scalability using AWS HealthOmics and SageMaker infrastructure.	Long-range genomic interaction analysis.
Borzoi [19]	Multi-layer regulatory model	Trained to predict RNA-seq coverage directly from DNA sequences; integrates multiple layers of regulatory predictions; fine-tuned for tissue-specific applications.	Cell- and tissue-specific RNA-seq coverage prediction, cis-regulatory pattern detection.
SEI [6]	Residual dilated convolutional layers	Trained on one-hot encoded sequences (4,096bp) using residual paths for multi-scale pattern detection; incorporates global integration layers for long-range dependencies; balances computational efficiency with expressiveness.	Variant effect prediction on cis-regulatory activity across chromatin profiles.

TREDNet [7]	Two-phase CNN architecture	Trained on ENCODE and NIH Roadmap Epigenomics datasets; uses CNNs to predict epigenomic signals (e.g., histone modifications) and refine enhancer predictions; supports in silico saturated mutagenesis for variant prioritization.	Enhancer prediction, variant prioritization.

Training Data

We constructed a dataset of positive and control enhancer sequences from four human

cell lines: HepG2, Hela, K562, and Neural Progenitor Cells (NPC). These cell lines

were selected for their relevance in genomic research and the availability of

comprehensive epigenomic data, such as DNase-Seq and histone modification profiles,

sourced from the ENCODE project using genome annotations for hg19 and hg38.

Positive enhancer sequences were defined through a two-step process. First, open

chromatin regions were identified using DNase-Seq data. These regions were then

intersected with H3K27ac histone modification signals, a hallmark of active enhancers.

To refine the dataset, we excluded non-enhancer regions such as exonic regions (coding

sequences), promoter regions (near transcription start sites), and low-confidence

regions listed in the ENCODE Blacklist (e.g., repetitive sequences).

Specific blacklisted peaks from UCSC browser tables

(wgEncodeDacMapabilityConsensusExcludable) were also removed.

Control sequences were selected using a similar filtering process but specifically

excluded regions overlapping both DNase-Seq peaks and H3K27ac signals to ensure

they represented non-active genomic regions. Both positive and control sequences

underwent identical exclusion criteria to eliminate confounding genomic features such

as exons and promoters.

All sequences were standardized to 1 kb in length to balance sufficient coverage of

regulatory regions with computational efficiency. This size is widely used in enhancer

studies as it captures relevant regulatory elements while avoiding extraneous genomic

features. The dataset includes an equal number of positive and control sequences for

each cell line: HepG2 (14,062 each), NPC (12,466 each), K562 (19,968 each), and Hela

(31,247 each). Metadata from the ENCODE project was used to retrieve biosample IDs

and associated DNase-Seq and histone modification data. Reference files provided

coordinates for exons, blacklist regions, and promoters, ensuring only valid enhancer

or control regions were included in the final dataset.

Validation Data

Datasets 1 and 2 were sourced from the study by [20], which utilized the Survey of

Regulatory Elements (SuRE) reporter assay to identify reporter assay quantitative trait

loci (raQTLs)—SNPs influencing regulatory element activity. For K562 cells, 19,237

raQTLs were identified with an average allelic fold change of 4.0-fold, while HepG2

cells yielded 14,183 raQTLs with a higher average fold change of 7.8-fold. Most

raQTLs were cell-type-specific, showing limited overlap between the two cell lines.

Dataset 3 was derived from [21] and focuses on a ~600 bp enhancer region at the

SORT1 locus (1p13). This region includes SNP rs12740374, which creates a C/EBP

binding site that influences SORT1 expression. The enhancer was tested in HepG2 cells

using a luciferase reporter assay. Data for flipped sequences were excluded due to the

orientation-independent nature of enhancers, and deletion datasets were omitted due to

computational model constraints, which focus on single-nucleotide variants.

Dataset 4, from [22] used MPRA to investigate 2,756 variants linked to red blood cell

traits in K562 cells. The study identified 84 highly significant SNPs with functional

effects, enriched in open chromatin regions shared between K562 cells and primary

human erythroid progenitors (HEPs). This dataset provides insights into erythroid-

specific regulatory activity.

Dataset 5, generated by [23], includes MPRA results for 284 SNPs tested in HepG2

cells. These SNPs, located within regions of high linkage disequilibrium identified

through eQTL analyses, offer a detailed map of allele-specific regulatory effects

relevant to liver-specific gene regulation and genotype-phenotype associations.

Dataset 6, from [24], includes lentiMPRA results for 14,042 single-nucleotide variants

fixed or nearly fixed in modern humans but absent in archaic humans (e.g.,

Neanderthals). Tested across three cell types—embryonic stem cells, neural progenitor

cells, and fetal osteoblasts—the study identified 1,791 sequences with regulatory

activity, including 407 with differential expression between modern and archaic

humans. This dataset focuses on NPC data.

Datasets 7, 8, and 9, from [25], include MPRA results for three coding exons—SORL1

exon 17, TRAF3IP2 exon 2, and PPARG exon 6—tested in HeLa cells. Respectively,

these datasets cover 1,962, 1,614, and 1,665 SNPs. The study explored enhancer

activity within these coding exons (eExons), finding that ~12% of mutations altered

enhancer activity by at least 1.2-fold, with ~1% causing changes of less than 2-fold.

This version eliminates repetition while maintaining all critical details about the

datasets and their relevance.

Acknowledgements

This work utilized the computational resources of the NIH HPC Biowulf cluster (http://hpc.nih.gov).

Funding

This research was supported by the Division of Intramural Research of the National Library of Medicine, National Institutes of Health; 1-ZIA-LM200881-12 (to I.O.).

References

1. Uffelmann E, Huang QQ, Munung NS, de Vries J, Okada Y, Martin AR, et al. Genome-wide association studies. Nat Rev Methods Primers. 2021 Dec;1(1):59.

2. Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, et al. 10 years of GWAS discovery: biology, function, and translation. Am J Hum Genet. 2017 Jul 6;101(1):5–22.

3. Knight JC. Approaches for establishing the function of regulatory genetic variants involved in disease. Genome Med. 2014 Oct 31;6(10):92.

4. Albert FW, Kruglyak L. The role of regulatory variation in complex traits and disease. Nat Rev Genet. 2015 Apr;16(4):197–212.

5. Zhou J, Troyanskaya OG. Predicting effects of noncoding variants with deep learning-based sequence model. Nat Methods. 2015 Oct;12(10):931–4.

6. Chen KM, Wong AK, Troyanskaya OG, Zhou J. A sequence-based global map of regulatory activity for deciphering human genetics. Nat Genet. 2022 Jul 11;54(7):940–9.

7. Hudaiberdiev S, Taylor DL, Song W, Narisu N, Bhuiyan RM, Taylor HJ, et al. Modeling islet enhancers using deep learning identifies candidate causal variants at loci associated with T2D and glycemic traits. Proc Natl Acad Sci USA. 2023 Aug 29;120(35):e2206612120.

8.   Ji Y, Zhou Z, Liu H, Davuluri RV. DNABERT: pre-trained Bidirectional Encoder
Representations from Transformers model for DNA-language in genome.
Bioinformatics. 2021 Aug 9;37(15):2112–20.

9.   Zhou Z, Ji Y, Li W, Dutta P, Davuluri R, Liu H. DNABERT-2: Efficient Foundation
Model and Benchmark For Multi-Species Genome. arXiv. 2023;

10. Dalla-Torre H, Gonzalez L, Mendoza Revilla J, Lopez Carranza N, Henryk
Grywaczewski A, Oteri F, et al. The nucleotide transformer: building and evaluating
robust foundation models for human genomics. BioRxiv. 2023 Jan 15;

11. de Almeida BP, Dalla-Torre H, Richard G, Blum C, Hexemer L, Gelard M, et al.
SegmentNT: annotating the genome at single-nucleotide resolution with DNA
foundation models. BioRxiv. 2024 Mar 15;

12. Avsec Ž, Agarwal V, Visentin D, Ledsam JR, Grabska-Barwinska A, Taylor KR, et al.
Effective gene expression prediction from sequence by integrating long-range
interactions. Nat Methods. 2021 Oct 4;18(10):1196–203.

13. Theodoris CV, Xiao L, Chopra A, Chaffin MD, Al Sayed ZR, Hill MC, et al. Transfer
learning enables predictions in network biology. Nature. 2023 Jun;618(7965):616–24.

14. Consens ME, Dufault C, Wainberg M, Forster D, Karimzadeh M, Goodarzi H, et al. To
Transformers and Beyond: Large Language Models for the Genome. arXiv. 2023;

15. Alharbi WS, Rashid M. A review of deep learning applications in human genomics
using next-generation sequencing data. Hum Genomics. 2022 Jul 25;16(1):26.

16. Yue T, Wang Y, Zhang L, Gu C, Xue H, Wang W, et al. Deep learning for genomics:
from early neural nets to modern large language models. Int J Mol Sci. 2023 Nov
1;24(21).

17. Fishman V, Kuratov Y, Petrov M, Shmelev A, Shepelin D, Chekanov N, et al. GENA-
LM: A Family of Open-Source Foundational Models for Long DNA Sequences.
BioRxiv. 2023 Jun 13;

18. Nguyen E, Poli M, Faizi M, Thomas A, Birch-Sykes C, Wornow M, et al. HyenaDNA:
Long-Range Genomic Sequence Modeling at Single Nucleotide Resolution. arXiv.
2023 Nov 14;

19. Linder J, Srivastava D, Yuan H, Agarwal V, Kelley DR. Predicting RNA-seq coverage
from DNA sequence as a unifying model of gene regulation. BioRxiv. 2023 Sep 1;

20. van Arensbergen J, Pagie L, FitzPatrick VD, de Haas M, Baltissen MP, Comoglio F, et
al. High-throughput identification of human SNPs affecting regulatory element activity.
Nat Genet. 2019 Jul;51(7):1160–9.

21. Kircher M, Xiong C, Martin B, Schubach M, Inoue F, Bell RJA, et al. Saturation
mutagenesis of twenty disease-associated regulatory elements at single base-pair
resolution. Nat Commun. 2019 Aug 8;10(1):3583.

22.   Ulirsch JC, Nandakumar SK, Wang L, Giani FC, Zhang X, Rogov P, et al. Systematic
621   functional dissection of common genetic variation affecting red blood cell traits. Cell.
622   2016 Jun 2;165(6):1530–45.

623   23.   Vockley CM, Guo C, Majoros WH, Nodzenski M, Scholtens DM, Hayes MG, et al.
624   Massively parallel quantification of the regulatory effects of noncoding genetic
625   variation in a human cohort. Genome Res. 2015 Aug;25(8):1206–14.

626   24.   Weiss CV, Harshman L, Inoue F, Fraser HB, Petrov DA, Ahituv N, et al. The cis-
627   regulatory effects of modern human-specific variants. eLife. 2021 Apr 22;10.

628   25.   Birnbaum RY, Patwardhan RP, Kim MJ, Findlay GM, Martin B, Zhao J, et al.
629   Systematic dissection of coding exons at single nucleotide resolution supports an
630   additional role in cell-specific transcriptional regulation. PLoS Genet. 2014 Oct
631   23;10(10):e1004592.