A periplasmic metallochaperone of the Cation Diffusion Facilitator YiiP is

required for Zn^{2+} sensing in *Pseudomonas aeruginosa*

Running title: An anchored membrane protein couples Zn^{2+} efflux to Zn^{2+} sensing in

Pseudomonas aeruginosa

Paula Mihelj[1], Tomás Moreyra[1], María Elena Carrizo[2], Daniel Raimunda[1]

[1]Instituto de Investigaciones Médicas Mercedes y Martín Ferreyra, INIMEC.

[2]Centro de Investigaciones en Química Biológica de Córdoba CIQUIBIC.

[1, 2] CONICET-Universidad Nacional de Córdoba. Argentina.

Abstract

Pseudomonas aeruginosa thrive to survive in harsh conditions imposed by the host. We have previously described two Zn^{2+}-transporting members of the Cation diffusion facilitator family, YiiP and CzcD, confering resistance to imipenem by decreasing the expression of the outer membrane porin OprD (A. Salusso and D. Raimunda, Frontiers in cellular and infection microbiology 7:84, 2017, https://doi.org/10.3389/fcimb.2017.00084). Here we provide evidence that a protein encoded in *yiiP*´s operon, PA3962, is fundamental to the coupling of this process. Immunodetection assay indicates that PA3962 locates in membranes. Supporting a role in *oprD* regulation, an insertion mutant has increased imipenem sensitivity, which is suppressed by gene complementation but not in presence of Zn^{2+}, as opposed to the YiiP mutant. We identified 2 pairs of conserved acidic residues in a hydrophobic juxtamembrane domain. Metal binding specificity and stoichiometry was explored in wild type and mutant versions of these. Zn^{2+} is the cognate metal being D40, D47 and D65 required for Zn^{2+} coordination. As the periplasmic Zn^{2+}-sensor CzcS regulates *oprD* expression, the interaction with it was analyzed *in vitro*. Interaction was Zn^{2+}-dependent, and mutations of D47A or D65A abolished it. We propose a role for PA3962, hereafter periplasmic metallochaperone of YiiP (PmcY), in the context of Zn^{2+} signaling pathways in *P. aeruginosa*. The rely YiiP–PmcY, supplies Zn^{2+} and activates CzcS/CzcR, down-regulating the transcription of the imipenem-permeable OprD. This mechanism would allow *P. aeruginosa* to put a brake on unspecific mechanisms for micronutrient uptake, with potential xenobiotic entry, while the cytosolic Zn^{2+} quota is still sufficient.

Introduction

Pseudomonas aeruginosa is a facultative anaerobic gram-negative bacteria and a prevalent opportunistic pathogen in respiratory infections of patients diagnosed with cystic fibrosis (1). It is considered an oligotrophic organism with a highly dynamic capacity for antibiotic resistance, metabolic adaptation, or phenotypical switching, which sometimes translates to fitness costs (2). Metabolic adaptations and antibiotic resistance involve optimization of Zn^{2+} usage in niches with low bioavailability (3). In the first case, the Zn^{2+}-dependent (C+) to Zn^{2+}-independent (C-) isoform switching of ribosomal subunits, increases the intracellular Zn^{2+} pool in this condition (4). In the second, Zn^{2+} induces carbapenem resistance by repressing the transcription, and then the antibiotic influx, of the outer membrane porin Occ1, also known as OprD (5). As carbapenem-resistant OprD mutants shows enhanced fitness *in vivo* (6) we hypothesize that this "master of survival", could benefit from a last resort-type of intracellular Zn^{2+} availability by finely tuning Zn^{2+}-sensing mechanisms. In this way this opportunistic pathogen could adapt metabolically and become resistant to carbapenems, avoiding the pay-offs of a fitness cost imposed by nutritional immunity(7).

We have previously shown that the Zn^{2+} exporters of the Cation diffusion facilitator family, YiiP and CzcD regulate levels of OprD (8). Single insertion mutants of these genes showed increased sensitivity to imipenem, and increased Zn^{2+} sensitivity when introduced into double mutant backgrounds where other Zn^{2+} resistance mechanisms were also affected. Considering studies by others showing that transcription of *czcD* responds to Zn^{2+} levels in the growth media with a slow

temporal activation (9) a role in Zn^{2+} detoxification might be considered. However,

the fact that *yiiP* transcripts are not upregulated by Zn^{2+}, opens new possibilities for

the role of this Zn^{2+}-transporter in Zn^{2+} mediated signaling.

Zn^{2+} sensing mechanisms in *P. aeruginosa* are well known. At the cytoplasmic level,

CadR, a member of the MerR transcription factor family, controls the expression of

CadA, also known as ZntA, a Zn^{2+}-transporting P_{IB}-ATPase (8, 10). Interestingly, it

has been proposed that under high Zn^{2+} conditions, CadA is required for timely

expression of CzcCBA, also known as HME-RND, a tripartite Zn^{2+}-export system

that promotes efflux from the periplasm to the extracellular space (10). This implies

that the specific CadA Zn^{2+}-export is coupled to Zn^{2+} sensing in the periplasm. In line

with this, extracellular Zn^{2+} bioavailability in *P. aeruginosa* is primarily detected in the

periplasm via the two-component system (TCS) CzcS-CzcR. CzcS senses

periplasmic Zn^{2+} by Zn^{2+} coordination in the periplasmic domain (11), which is

transduced to cytosolic domains promoting autophosphorylation of the histidine

kinase domain and finally, the transphosphorylation to CzcR. Phosphorylated CzcR

(P-CzcR) has dual effects on transcription of genes in the CzcS-CzcR regulon (12).

For example, P-CzcR up-regulates transcription of HME-RND, thus counteracting a

potential Zn^{2+} overload (9, 10) in the periplasm. However, P-CzcR affects negatively

the transcription of genes involved in pyocianin synthesis and of the outer membrane

porin *oprD* (11, 12).

Here we present evidence suggesting that a periplasmic metallochaperone of YiiP

(PmcY/PA3962) is required for OprD repression and thus, for Zn^{2+} sensing in the

periplasm. Our data shows that PmcY is a Zn^{2+}-metallochaperone that interacts with

CzcS, its putative acceptor or target protein. We also propose that the intracellular

Zn^{2+} efflux to the periplasm mediated by YiiP-PmcY has coevolved in

Pseudomonadales as a mechanism for limiting the fitness cost of adapting to

common environmental or host-imposed scenarios with low Zn^{2+} bioavailability, by

repressing an unspecific outer membrane porin.

Materials and methods

Bacterial strains, plasmids and growth media conditions

Pseudomonas aeruginosa MPAO1 wild-type (WT), and the insertion mutants

yiiP::dTn5 and *pmcY*/PA3962*::dTn5* were obtained from the *P. aeruginosa* Mutant

Library at the University of Washington ((13); Table 1). The *pmcY*/PA3962*::dTn5*

complemented strain was obtained by cloning the PA3962 locus plus the 250bp

upstream in pSEVA621 (14). The sequence for *gfp* was introduced at the 3'-end as

a fusion in order to promote protein stability. The primers used in this and other

procedures are listed in Table 1. Electrocompetent *P. aeruginosa* strains were

transformed by electroporation as described (15) and selected on Luria–Bertani (LB)

agar plates supplemented with gentamicin (30µg/ml). Cells were grown at 37°C in

LB or Müller-Hinton media as indicated, supplemented with tetracycline (30–

60µg/ml), and gentamicin (30µg/ml) as required.

Plasmid constructions and protein expression and purification

In a first attempt to yield a whole version of PmcY by heterologous expression in *E.*

coli PA3962 locus was cloned from *P. aeruginosa* MPAO1 gDNA and inserted by

ligation in the expression plasmids pET23b(+) (Novagen) or pGEX4T-2 using

primers described in Table 1. These constructs contained at C-terminal a

hexahistidine ($(His)_6$) tag preceded by Tobacco Etch Virus (TEV) protease

recognition site. Due to low yield of the protein and guided by bioinformatics analyses

we expressed the juxtamembrane domain of PmcY ($PmcY_j$) as a GST fusion protein

by deletion of N- and C-termini and $(His)_6$ tag of PmcY by the one-step site-directed

deletion PCR protocol (16). Site directed mutagenesis of conserved residues were

obtained using the same procedure. Similarly, the soluble domain (SD) of CzcS

(amino acids 40-166, $CzcS_{SD}$) was cloned by PCR amplification from MPAO1 gDNA

and ligated in pET16bStrep (Table 1) with a twin-strep tag at N-terminal. All

constructs and mutants were confirmed by DNA sequencing.

For protein expression and purification *E. coli* BL21(DE3)pLysS were transformed

with one of the above constructs. After selection cells were grown in LB overnight at

37°C on a shaker at 200 rpm. Next day 1-2 L of LB media with antibiotic was

inoculated with cells to reach $OD_{600}= 0.1$ and grown in the same conditions till early

log-phase. At this point cultures were induced with 0.15 mM IPTG for 16 hours at

20°C. For purification of GST and GST fusion versions of $PmcY_j$ cells were

resuspendend in 50 ml of Tris-HCl 20 mM pH 7.5, NaCl 500 mM, TCEP 0.25 mM,

0.1% Triton X-100 (Buffer A) plus PMSF 1 mM, and lysed in a cell disruptor

EmulsiFlex C3 (Avestin). After ultracentrifugation at 100,000 *g* the supernatant was

incubated with 1 ml of Glutathion Sepharose 4B or Fast Flow (GE Healthcare) for 1

hour at room temperature. Resin was packed by gravity in a polypropylene column

(Bio-Rad) and washed with 30 resin volumes of buffer A. The fusion protein was

eluted with buffer A containing 40 mM glutathione. The soluble domain for CzcS was

expressed and purified as described earlier (11) with minor modifications. After cell

growth and induction cells were lysed as above and ultracentrifuged at 100,000 *g*.

The supernatant was incubated for 1 hour at room temperature with 0.5 ml of Strep-

Tactin resin (IBA Lifesciences) equilibrated with buffer A for 1h and eluted after

washing with 30 resin volumes. Bound protein was eluted with 10 resin volumes of

buffer A containing 50 mM of D-biotin. In both cases elution fractions were analyzed

by SDS-PAGE 12%, Coomassie brilliant blue staining and Western blot using $(His)_6$-

tag antibody (pAb, Rabbit, Genscript), GST-tag antibody (pAb, Rabbit, Genscript) or

StrepMAB-Classic antibody (IBA Lifesciences) as primary antibodies. Those

fractions containing protein were pooled for dialysis in buffer A using a dialysis tubing

(SnakeSkin, Thermo Scientific) with 3.5 kDa cut-off. After addition of 10% glycerol,

protein concentration was estimated by Bradford (Bio-Rad) and 280nm lectures in

NanoDrop (Thermo Scientific). Protein samples were frozen and maintained at -

80°C for later use. For in-column interaction assays the bait protein, GST-PmcY$_i$ WT

and mutant versions, were not eluted from the resin and kept at 4°C in the dark for

later use and store for no more than 2 weeks. In this case protein concentrations

were determined by densitometry. Denatured resin samples were run in SDS-PAGE

12%, stained with Coomassie brilliant blue (CBB) and images analyzed with ImageJ

(17). For calibration, bovine serum albumin (Sigma-Aldrich) standards were run in

the same gel.

Imipenem sensitivity assays

Susceptibility to imipenem (Sigma-Aldrich) was assessed by the agar dilution or the

inhibition halo methods in Müller–Hinton (MH) medium as described before (8). For

agar dilution overnight grown cells were serially diluted from an initial $OD_{600}= 1$ and

then 10 µl were spotted on MH agar plates. For inhibition halos, a suspension of

cells at $OD_{600}=0.05$ in soft agar were layered on top of MH agar plates. After

solidifying, 7-mm diameter filter paper discs embedded with the indicated amount of

imipenem were located on top of the soft agar. In all cases growth or inhibition was

evaluated after overnight incubation at 37°C.

Metal binding assays

Transition metal (TM) binding capacity of GST fusion proteins was evaluated using

the chromogenic dye 4-(2-Pyridylazo) resorcinol (PAR, Tokyo Chemical Industry).

For incubation of the proteins with TM we followed described protocols for in-solution

(18) or in-column (19) assays with some modifications. To assess binding to proteins

10-20 µM samples (or the mass bound to resin to yield this concentration before

adding PAR) were incubated for 30 min with 5 molar excess of TM in buffer A. For

in-solution binding assays unbound TM were separated by exclusion

chromatography with Sephadex G25-80 (Sigma-Aldrich) and for the in-column type

the resin was washed with 10 volumes of buffer A to remove unbound TM. Before

TM quantification protein concentrations were estimated as described earlier and

samples containing proteins in solution or bound to resin were treated with 0.4-0.8

mg/ml of proteinase K. For TM quantification, PAR was added to a final

concentration of 100 µM and read in microplates at 500 nm. Calibration curves were

done in parallel for each TM and the molar ratio of protein to TM calculated. In both

conformations GST alone or GST fusions were run in parallel to detect TM bound in

proteins as purified. These values represented no more than 10% of those found in

TM incubated samples and were subtracted in all cases. For the quantification of Mn^{2+} the buffer pH was modified from 7.5 to 11 as the colorimetric reaction of this TM is more sensitive. Therefore, after the digestion reaction with proteinase K, the samples were evaporated completely and resuspended in 200 mM CAPS (3-(Cyclohexylamino)-1-propanesulfonic acid) buffer pH 11.

Protein localization

The *E. coli* strain BL21(DE3)pLysS transformed with the plasmid pET-23b(+)-PA3962, which expresses the PA3962 protein fused to a 6-histidine tag, was grown in 12.5 ml of LB medium at 37°C for 16 h. This culture was placed in two flasks with 500 ml of LB medium and grown with shaking (200 rpm) at 37°C for 1 hour. One of the cultures was induced with 0.15 mM of IPTG and the other was used as a control (basal). After 3 hours of growth the cells were harvested by centrifugation at 4,000 *g*, 4°C for 10 minutes. Cell pellets were washed in ice-cold 30 ml of 50 mM Tris-HCl pH 7.5 and 0.3 M sorbitol (buffer S) 3 times, with addition of 10 µM EDTA in the second wash. Cells were pelleted by centrifugation at 9,000 *g*, 4°C for 10 minutes. Finally, cells were resuspended in 30 ml of ultrapure water and incubated for 5 minutes at room temperature. They were then centrifuged at 11,000 *g*, 4°C for 30 minutes. The supernatants were collected and centrifuged again to eliminate any possible contamination. After this they were considered the periplasmic fractions. For cytosolic and total membrane fractions cell pellets were resuspended in 7 ml of 10 mM Tris-HCl pH 7.5 and disrupted at 4°C using a tip sonicator (Vibra Cell, Sonics & Materials, Inc.) with 6 cycles of 20 seconds on and 60 seconds off pulses. The lysate was centrifuged at 6,000 *g* for 10 minutes at 4°C to remove undisrupted cells

and centrifuged again at 100,000 *g* for 30 minutes at and 4°C. The supernatants

were recovered and classified as cytoplasmic fractions. The resultant pellets were

resuspended in 5 ml of 10 mM Tris-HCl pH 7.5 and classified as total membrane

fractions. The protein concentration of all fractions was quantified by Bradford assay,

and those with a value less than 1 mg/ml were concentrated using centrifuge filtration

tubes (Centricon, Millipore). Samples of all fractions were prepared at concentrations

of 1 mg/ml and 5X Laemmli sample buffer was added. A final total protein mass of

10 µg was loaded per lane on 14% SDS-PAGE gels in duplicate, one for CBB

staining and the other for Western blot. $(His)_6$-tag antibody (pAb, Rabbit, Genscript)

and anti-rabbit antibody labeled with horseradish peroxidase were used to detect

expression in fractions. Images were taken using Amersham Image Quant 800 Fluor

(Cytiva).

Protein-protein interaction assay

To evaluate the interaction between PmcY and CzcS, the recombinant proteins

GST-PmcY$_j$ (wild-type and mutants D40A, D47A, and D65A) and CzcS$_{SD}$ were used.

A 100±20 mg sample of purified WT or mutant versions of GST-PmcY$_j$ were treated

with EDTA-Tris. For this, Glutathion Fast Flow (GE Healthcare) resin with bound

protein was incubated with 1 ml of buffer A plus 1 mM EDTA-Tris in a 10 ml

polypropylene chromatography column (Bio-Rad). 3 ml of buffer A were used to

remove residual EDTA from column. The interaction was tested in two conditions,

with GST-PmcY$_j$ loaded with ($+Zn^{2+}$) and without ($-Zn^{2+}$) ZnCl$_2$. For this resin was

incubated for 5 min with 500 µl of 100 µM ZnCl$_2$ in buffer A or just buffer A,

respectively. The CzcS$_{SD}$ protein was used in solution and previously treated with

0.5 mM EDTA-Tris for 30 minutes in ice. EDTA-Tris was removed with PD-10 column

equilibrated in buffer A. In-column GST-PmcY$_j$ and mutant versions, in its two

conditions (+Zn^{2+}) and (-Zn^{2+}), were incubated with 25 µM CzcS$_{SD}$, added in 250 µl,

during two passes through the column (Flow through, FT) and 4 washes (W1-4) were

performed with 250 µl of buffer A. Finally, the resin was resuspended in 250 µl of

buffer A and samples prepared with 5X Laemmli loading buffer for SDS-PAGE 12%

analyses. After CBB staining images were taken using Amersham Image Quant 800

Fluor (Cytiva).

Bioinformatics analyses

To analyze the conservation of residues in a putative PmcY family, homologs were

searched in a local database of all protein sequences of RefSeq reference and

representative genomes protein sequences (Refseq_select_prot) (January 2024) by

the reciprocal best hit (RBH) method (20). 407 unique organisms having PmcY-like

members with unique accession numbers were found. Multiple sequence alignment

(MSA) was done in Clustal Omega (21) using a similar number of sequences and

the same MSA was used in ChimerX (22) to create the amino acids conservation

map in the putative 3D AF structure of PA3962 (AF-Q9HX54-F1).

This number of organisms were used later to compare and create a list of non-

redundant bacterial organisms having, or not, the *pmcY-yiiP* architecture. The

search of the *pmcY-yiiP* operon was done using Multigenblast (23). Similar RBH

searches were performed for OprD and CzcS homologs in the same data base. After

removal of duplicated organism names, a 4-group diagram was prepared with these

organisms (https://bioinformatics.psb.ugent.be/webtools/Venn/) and a phylogenetic

tree was constructed for visualization of coevolution in iTOL (24).

Results

PA3962 is a membrane protein required for intrinsic imipenem resistance

PA3962 is part of a putative operon with YiiP (PA3963). Previous bioinformatics

studies of identified *P. aeruginosa* secreted proteins pointed that the N-terminal

sequence in PA3962 is a type II signal peptide (25). These results were supported

with a plasmid library screening in *E. coli* where several PAO1 genomic DNA

fragments, including PA3962, fused to truncated alkaline phosphatase (PhoA) drove

its periplasmic localization. Signal peptides are recognized by different secretion

systems in bacteria and classified in different groups. The type II are related to

proteins anchored to the membrane, also called lipoproteins. A consensus sequence

called "lipobox" is found in the type II signal peptide, which allows for molecular

recognition by the machinery involved in processing and finally anchoring the protein

to the inner or outer membrane. For this, first a conserved cysteine residue in the

lipobox is diacylated with diacylgicerol. Second, the signal peptide is removed by a

specific protease leaving free and reactive the amino group of the cysteine for a final

third acylation (26). Figure 1, shows the predicted structure of PA3962 by

AlphaFold2 (AF2) (27) showcasing the colour–coded hydrophobic profile generated

in ChimeraX and position of Cys23 in the lipobox. The ribbon structures depict the

putative position of the protein facing the periplasmic space after the anchoring

occurs. In order to confirm the membrane localization of PA3962 we expressed the

whole protein with a C-terminal $(His)_6$ tag in *E. coli* in basal and induced conditions.

After fractioning cells, cytosolic, total membranes, and periplasmic samples were

analyzed by SDS-PAGE and Western blot using anti-$(His)_6$ as primary antibody (Fig.

2B). Protein detection was evident in most of the fractions and signal was increased

under basal conditions, pointing to cytotoxicity when the full length protein is

overexpressed. Most importantly the detection in total membrane fraction was

observed. As the mass of protein loaded in all lanes was similar, and assuming an

equal contribution of each compartment in cell mass, a qualitative conclusion is that

PA3962 is a membrane protein. PhoA detection in *E. coli* requires folding in the

periplasm, thus this orientation is the most likely scenario in *P. aeruginosa* and we

renamed PA3962 as Periplasmic metallochaperone of YiiP, PmcY.

As previous results indicated a role for YiiP in imipenem resistance (8) we next

analyzed the minimal inhibitory concentration (MIC) for the antibiotic in Tn5-

generated mutant strains (Fig. 1C). YiiP and PmcY mutants are sensitive to

imipenem *vs.* WT and complementation of PmcY strain was achieved by

transformation of mutant cells with the low copy plasmid pSEVA621 (14) carrying

PA3962 locus under transcriptional regulation of 250 bp upstream (Fig. 1D). These

results indicated that the molecular mechanism leading to the loss of imipenem

resistance involves the lack of PmcY and YiiP interaction. As YiiP is involved in Zn^{2+}

transport, and imipenem resistance is induced by periplasmic Zn^{2+} sensing we

hypothesized that both YiiP and PmcY mutants would remain unresponsive to Zn^{2+}

content in the growth media. Only PmcY mutant strain showed the same antibiotic

resistance in presence of high levels of Zn^{2+}, indicating that it is required for Zn^{2+}

sensing in the periplasm (Fig. 1E). As part of a sensing mechanism leading to

intrinsic imipenem resistance, PmcY should bind Zn^{2+}, interact with a sensor, and

likely transfer the metal to it.

PmcY binds Zn^{2+} specifically through conserved aspartate residues

We hypothesized that PmcY plays a role as metallochaperone. These function in

metalation of proteins or in detoxification pathways, and in both cases the

commonality is their capacity to bind a specific transition metal through conserved

residues. A multiple sequence alignment of PmcY homologs obtained by a reciprocal

best hit search in complete bacterial genomes allowed us to identified conserved

residues (Fig. 2A, maroon). Most of these would be located in a periplasmic domain

juxtaposed to the membrane. Conserved amino acids are all acidic, with oxygen (O)

ligands of lateral chains of aspartates and glutamates. After several failed trials of

full length PmcY expression and purification to get enough protein for biochemical

characterization, we opted to fuse this domain (amino acids 30 to 93), which we

called PmcY juxtamembrane ($PmcY_j$), to GST. The fusion protein was purified and

its capacity and specificity to bind transition metals analyzed. For this, five molar

excess of different metals were incubated with the fusion protein (Fig. 2B-C) and

after unbound metal was separated by exclusion chromatography (Fig. 2B) or

washed (Fig. 2C), protein was quantified and metal bound to protein detected as

described in Materials and methods. This allowed us to confirmed that Zn^{2+} is the

cognate metal for PmcY and that the stoichiometry of binding is 2 Zn^{2+} per PmcY

molecule. To identified possible binding sites, we analyzed the Zn^{2+} binding of the

single mutants of highly conserved positions with O ligands; i.e., D40A, D47A and

D65A. All these mutations lowered the binding stoichiometry to 1, indicating their

likely participation in two Zn^{2+} binding sites. A closer look to the distances between

O atoms in lateral chains of the PmcY structure, underscores residues D47-D65 as

candidates for one Zn^{2+} binding with the archetypical tetrahedral geometry (Fig. 2D).

PmcY interacts with the periplasmic Zn^{2+} sensor CzcS

Another property which defines a metallochaperone is the interaction with its target/s

protein/s. This is key in order to promote the ligand transfer reaction, and the metal

transfer, between the donor, the metallochaperone, and the acceptor, the target

protein. Thus an interaction between PmcY and CzcS could promote Zn^{2+} sensing

by means of a ligand transfer reaction between partners. In order to test this, we

expressed and purified the periplasmic soluble domain (SD) of CzcS (amino acids

40 to 166, named CzcS$_{SD}$) as described by Wang et al. (11) with a twin Strep-tag at

the N-terminal end. The protein preparation was treated with EDTA to eliminate Zn^{2+}

carried-over during purification, and then incubated with GST-PmcYj, in two

conditions, EDTA-treated or loaded with Zn^{2+}. After collecting flow through and

washes we detected the presence of CzcS in the final resin (Rf) only when GST-

PmcY$_j$ was loaded with Zn^{2+} (Fig. 3A, black asterisk). In order to study how Zn^{2+}

occupancy affected this interaction, GST-PmcY$_j$ mutants, D40A, D47A and D65A

were analyzed similarly. The Zn^{2+}-dependent interaction was lost only in D47A and

D65A mutants. This supports our previous proposition in which these acidic residues

might coordinate Zn^{2+} in one binding site. Mutation D40A did not preclude the

interaction in presence of Zn^{2+}.

PmcY-YiiP operon coevolved with OprD and CzcS in Pseuodomonadales

The data presented suggest that YiiP, PmcY, CzcS/CzcR participate in a sensing

mechanism coupling Zn^{2+} efflux via YiiP, to imipenem resistance (as *oprD*

expression is downregulated). If this mechanism is conserved among gram-negative

bacteria, we hypothesize that PmcY homologs found in other organisms would

present the same operon architecture as in *P. aeruginosa* MPAO1. A multigene

BLAST search did not support this. However, organisms presenting the operon

architecture were mostly of the Pseudomonadales order. Most importantly,

contrasted with organisms with non-*pmcY-yiiP* operon detected, *oprD* and *czcS* were

mostly distributed in organisms with the operon *pmcY-yiiP*; i.e., the

Pseudomonadales order. The lack of CzcS in a high number of members in this

clade is surprising. However, the periplasmic Cu^+ sensor CopS also has been

proposed to downregulate *oprD* transcription (28). Altogether these data support a

likely coevolution of these genes. We discuss in the next section what could be the

selection pressure of the coevolution and the physiological role in a context of Zn^{2+}

bioavailability.

Discussion

The association between the Zn^{2+} induced repression of *oprD* transcription, and the

Zn^{2+} induced resistance to imipenem is long recognized (29). Moreover, several

works have used this response to dissect important molecular and mechanistic

aspects of the CzcS-CzcR signaling pathway (11, 30) in the periplasmic space. Here

we present evidence pointing to the participation of YiiP and PmcY in periplasmic

Zn^{2+}-sensing mechanisms. In particular, we propose that both players affect *oprD*

transcription, and likely other genes, by promoting Zn^{2+}-sensing; i.e., the metalation

of the Zn^{2+} sensor CzcS. We provide evidences supporting our model, showing that

PmcY binds Zn^{2+} specifically, with high specificity and with a molar stoichiometry of

2 Zn^{2+} per molecule. Metal coordination is given by mean of conserved Asp residues.

Phenotypical analyses showed that different to WT and YiiP mutant, the mutant

strain of PmcY lacks the Zn^{2+}-dependent response to imipenem resistance likely by

a lack or reduced Zn^{2+}-sensing capacity of CzcS. These results lead us to propose

PmcY as a Zn^{2+}-metalochaperone.

Many of the players involved in orchestrating Zn^{2+} homeostasis in *P. aeruginosa*

have been described (31). High Zn^{2+} bioavailability is sensed by periplasmic and

cytosolic sensors, CzcS and CadR, respectively which lead to transcription of high

capacity Zn^{2+} transport systems, such as HME-RND and P_{IB}-type ATPases (9, 10).

The integration of these compartmentalized sensing mechanisms can be interpreted

considering electrochemical Zn^{2+} gradient. As Zn^{2+} entries and accumulates first in

the periplasm transcription of HME-RND are first expected. However, free Zn^{2+} is

almost negligible in cells (32) and a role for cytosolic Zn^{2+} exporters have been

proposed in periplasmic Zn^{2+} sensing mechanisms (10). This is important as it

suggests the participation of Zn^{2+}-transport systems allowing cross-talk between

cellular compartments to promote intracellular Zn^{2+} exit even when extracellular Zn^{2+}

could turn into lower levels.

Could a similar scenario, involving Zn^{2+} transport systems, participate in cross-talk

between compartments when Zn^{2+} bioavailability is low? Considering the risk of

losing Zn^{2+} in such scenario, the process should be accompanied of a "favourable"

fitness cost. We proposed that Zn^{2+}-export by YiiP is linked to Zn^{2+}-sensing by CzcS

with the final outcome of delaying *oprD* expression, which role allows entry of

nutrients, but also xenobiotic molecules such as antibiotics. Thus this

hypersensitivity to sense intracellular Zn^{2+}, when scarce but still available, would be

beneficial for this opportunistic pathogen as *oprD* induction is still repressed.

HME-RND also is part of the CzcS-CzcR regulon (30) and Zn^{2+}-binding to CzcS

upregulates the transcripts levels. Even when transcription could be triggered by

YiiP-PmcY activity we speculate that low Zn^{2+} availability is not sufficient to promote

HME-RND activity as this transport system would have similar biochemical

properties to CadA; i.e.: high transport capacity but low metal affinity. Incidental Zn^{2+}-

entry via OprD has been suggested (33). Although counterintuitive, the up-regulation

of HME-RND by CzcS activation, would prepare the cell for a potential Zn^{2+} overflow

in the periplasm with minimal basal activity of HME-RND until Zn^{2+} bioavailability

increases. However, evidence to support this is still missing and the lack of Zn^{2+}

resistance shown by OprD mutants contradicts this role (30).

Lastly, and considering our working model, although it is tempting to hypothesize

that a Zn^{2+}-basic amino acid abduct could be the physiological substrate transported

by OprD, the metal-chelating molecule EDTA was not able to outcompete

arginine/lysine transport by OprD in *in vitro* studies (5). Nonetheless, considering the

Zn^{2+} binding capacity of several xenobiotics, including antibiotics, and the evident

lack of specificity of *oprD* for carboxylated substrates, its overexpression could be

viewed as a "razors-edge"-like situation promoting Zn^{2+} uptake linked to the

xenobiotc molecule carrying the ion. Thus, braking the system could be beneficial

until Zn^{2+} reservoirs are downsized to the limits of a "go-hunting" for survival

condition.

Conflict of interest statement

404 All authors declare that research was conducted in the absence of any commercial

405 or financial relationships.

Author contributions

408 Conceived and designed the experiments: D.R., P.M.

409 Analyzed the data: D.R., P.M., T.M., M.E.C

410 Contributed reagents/samples/analysis/tools: D.R., M.E.C

411 Wrote or edited the manuscript: D.R., P.M.

412 All authors read and approved the final manuscript.

Funding

This work was supported by the Agencia Nacional de Promoción Científica y

Tecnológica (ANPCyT) PICT-2018-03139 to D.R., and PICT-2021-0147 to M.E.C.

D.R. and M.E.C. are Consejo Nacional de Investigaciones Científicas y

Tecnológicas (CONICET) investigators. P.M. is recipient of a CONICET doctoral

fellowship. T.M. is recipient of a doctoral fellowship from ANPCyT (PICT-2018-

03139).

Abbreviations

TM, transition metal; CDF, cation diffusion facilitator; IPTG, Isopropyl β-D-1-thiogalactopyranoside; PAR, 4-(2-Pyridylazo) resorcinol; HME-RND, heavy metal exporter of the resistance-nodulation-division family.

Acknowledgments

We thank Camila Semeniuk, Dr. Laura Montroull and Dr. Andrea Pellegrini for their invaluable help during this work.

References

1. Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JAJ, Sommer LM, Molin S, Johansen HK. 2021. *Pseudomonas aeruginosa* adaptation and evolution in patients with cystic fibrosis. Nature Reviews Microbiology 19:331-342.
2. Zhu M, Dai X. 2024. Shaping of microbial phenotypes by trade-offs. Nat Commun 15:4238.
3. Panina EM, Mironov AA, Gelfand MS. 2003. Comparative genomics of bacterial zinc regulons: enhanced ion transport, pathogenesis, and rearrangement of ribosomal proteins. Proc Natl Acad Sci U S A 100:9912-7.
4. Pederick VG, Eijkelkamp BA, Begg SL, Ween MP, McAllister LJ, Paton JC, McDevitt CA. 2015. ZnuA and zinc homeostasis in *Pseudomonas aeruginosa*. Sci Rep 5:13139.
5. Eren E, Vijayaraghavan J, Liu J, Cheneke BR, Touw DS, Lepore BW, Indic M, Movileanu L, van den Berg B. 2012. Substrate specificity within a family of outer membrane carboxylate channels. PLoS Biol 10:e1001242.
6. Skurnik D, Roux D, Cattoir V, Danilchanka O, Lu X, Yoder-Himes DR, Han K, Guillard T, Jiang D, Gaultier C, Guerin F, Aschard H, Leclercq R, Mekalanos JJ, Lory S, Pier GB. 2013. Enhanced in vivo fitness of carbapenem-resistant *oprD* mutants of *Pseudomonas aeruginosa* revealed through high-throughput sequencing. Proc Natl Acad Sci U S A 110:20747-52.
7. Kehl-Fie TE, Skaar EP. 2010. Nutritional immunity beyond iron: a role for manganese and zinc. Curr Opin Chem Biol 14:218-24.
8. Salusso A, Raimunda D. 2017. Defining the Roles of the Cation Diffusion Facilitators in Fe^{2+}/Zn^{2+} Homeostasis and Establishment of Their Participation in Virulence in *Pseudomonas aeruginosa*. Front Cell Infect Microbiol 7:84.
9. Ducret V, Abdou M, Goncalves Milho C, Leoni S, Martin-Pelaud O, Sandoz A, Segovia Campos I, Tercier-Waeber ML, Valentini M, Perron K. 2021.

Global Analysis of the Zinc Homeostasis Network in *Pseudomonas aeruginosa* and Its Gene Expression Dynamics. Front Microbiol 12:739988.

10. Ducret V, Gonzalez MR, Leoni S, Valentini M, Perron K. 2020. The CzcCBA Efflux System Requires the CadA P-Type ATPase for Timely Expression Upon Zinc Excess in *Pseudomonas aeruginosa*. Front Microbiol 11:911.

11. Wang D, Chen W, Huang S, He Y, Liu X, Hu Q, Wei T, Sang H, Gan J, Chen H. 2017. Structural basis of Zn(II) induced metal detoxification and antibiotic resistance by histidine kinase CzcS in *Pseudomonas aeruginosa*. PLoS Pathog 13:e1006533.

12. Dieppois G, Ducret V, Caille O, Perron K. 2012. The transcriptional regulator CzcR modulates antibiotic resistance and quorum sensing in *Pseudomonas aeruginosa*. PLoS One 7:e38148.

13. Held K, Ramage E, Jacobs M, Gallagher L, Manoil C. 2012. Sequence-verified two-allele transposon mutant library for *Pseudomonas aeruginosa* PAO1. J Bacteriol 194:6387-9.

14. Martinez-Garcia E, Fraile S, Algar E, Aparicio T, Velazquez E, Calles B, Tas H, Blazquez B, Martin B, Prieto C, Sanchez-Sampedro L, Norholm MHH, Volke DC, Wirth NT, Dvorak P, Alejaldre L, Grozinger L, Crowther M, Goni-Moreno A, Nikel PI, Nogales J, de Lorenzo V. 2023. SEVA 4.0: an update of the Standard European Vector Architecture database for advanced analysis and programming of bacterial phenotypes. Nucleic Acids Res 51:D1558-D1567.

15. Choi KH, Schweizer HP. 2006. mini-Tn7 insertion in bacteria with single *att*Tn7 sites: example *Pseudomonas aeruginosa*. Nat Protoc 1:153-61.

16. Liu H, Naismith JH. 2008. An efficient one-step site-directed deletion, insertion, single and multiple-site plasmid mutagenesis protocol. BMC Biotechnol 8:91.

17. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9:671-675.

18. Raimunda D, Subramanian P, Stemmler T, Argüello JM. 2012. A tetrahedral coordination of Zinc during transmembrane transport by P-type Zn^{2+}-ATPases. Biochim Biophys Acta 1818:1374-7.

19. Modrof J, Becker S, Muhlberger E. 2003. Ebola virus transcription activator VP30 is a zinc-binding protein. J Virol 77:3334-8.

20. Mihelj P, Abreu I, Moreyra T, Gonzalez-Guerrero M, Raimunda D. 2023. Functional Characterization of the Co^{2+} Transporter AitP in *Sinorhizobium meliloti*: A New Player in Fe^{2+} Homeostasis. Appl Environ Microbiol 89:e0190122.

21. Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. Nucleic acids research 52:W521-W525.

22. Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023. UCSF ChimeraX: Tools for structure building and analysis. Protein Sci 32:e4792.

23. Medema MH, Takano E, Breitling R. 2013. Detecting sequence homology at the gene cluster level with MultiGeneBlast. Mol Biol Evol 30:1218-23.

502   24.   Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to
503   the phylogenetic tree display and annotation tool. Nucleic Acids Res 52:W78-
504   W82.
505   25.   Lewenza S, Gardy JL, Brinkman FS, Hancock RE. 2005. Genome-wide
506   identification of *Pseudomonas aeruginosa* exported proteins using a
507   consensus computational strategy combined with a laboratory-based PhoA
508   fusion screen. Genome Res 15:321-9.
509   26.   El Rayes J, Rodriguez-Alonso R, Collet JF. 2021. Lipoproteins in Gram-
510   negative bacteria: new insights into their biogenesis, subcellular targeting and
511   functional roles. Curr Opin Microbiol 61:25-34.
512   27.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,
513   Tunyasuvunakool K, Bates R, Zidek A, Potapenko A, Bridgland A, Meyer C,
514   Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler
515   J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M,
516   Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,
517   Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure
518   prediction with AlphaFold. Nature 596:583-589.
519   28.   Caille O, Rossier C, Perron K. 2007. A copper-activated two-component
520   system interacts with zinc and imipenem resistance in *Pseudomonas*
521   *aeruginosa*. J Bacteriol 189:4561-8.
522   29.   Conejo MC, Garcia I, Martinez-Martinez L, Picabea L, Pascual A. 2003. Zinc
523   eluted from siliconized latex urinary catheters decreases OprD expression,
524   causing carbapenem resistance in *Pseudomonas aeruginosa*. Antimicrob
525   Agents Chemother 47:2313-5.
526   30.   Perron K, Caille O, Rossier C, Van Delden C, Dumas JL, Kohler T. 2004.
527   CzcR-CzcS, a two-component system involved in heavy metal and
528   carbapenem resistance in *Pseudomonas aeruginosa*. J Biol Chem 279:8761-
529   8.
530   31.   Ducret V, Gonzalez D, Perron K. 2023. Zinc homeostasis in *Pseudomonas*.
531   Biometals 36:729-744.
532   32.   Maret W. 2013. Zinc biochemistry: from a single zinc enzyme to a key element
533   of life. Adv Nutr 4:82-91.
534   33.   Wang S, Cheng J, Niu Y, Li P, Zhang X, Lin J. 2021. Strategies for Zinc
535   Uptake in *Pseudomonas aeruginosa* at the Host-Pathogen Interface. Front
536   Microbiol 12:741873.

 Tables

 Table 1. List of Strains, plasmids and primers used in this study.

Strain/plasmid/primer	Relevant characteristics/sequence	Reference/Use
Pseudomonas aeruginosa		
MPAO1	Wild-type	Held *et al.* (2012)
yiiP::dTn5	Tn5 derivative ISphoA/hah insertion in *yiiP* (PA3963), Tc^r	Held *et al.* (2012)
pmcY::dTn5	Tn5 derivative ISphoA/hah insertion in *pmcY* (PA3962), Tc^r	Held *et al.* (2012)
C-pmcY	Tn5 derivative ISphoA/hah insertion in *pmcY*, pSEVA621-P$_{pmcY}$-*pmcY-gfp*, Tc^r Gm^r	This work
Escherichia coli		
BL21(DE3) pLysS	*F- ompT hsdSB* (rB-mB-) *gal dcm* (DE3) *pLysS* (Cam^r)	Invitrogen, Carlsbad, CA,USA
DH5α	*supE44 ∆lacU169* (φ80 *lacZ* ∆M15*) hsdR17 recA1 endA1 gyrA96 thi1 relA1*	Invitrogen, Carlsbad, CA,USA
Plasmids		
pET23b(+)	T7 expression plasmid, Ap^r	Novagen, Darmstadt, Germany
pGEX-4-T2	T7*lac* expression plasmid*,* Ap^r	Cytiva, Marlborough, MA, USA
pSEVA621	RK2 cloning plasmid, Gm^r	Martínez-García *et al.* 2023
pET16bStrep	T7 expression plasmid, Ap^r	Luis Rubio´s laboratory
Primers		
F- pET PA3962	CAGTCATATGCGCCTTGCCACCCGCG	Cloning PA3962 in pET23b(+) + NdeI at 5'-end
R- pET PA3962	ACTCGAGGGACTGAAAATACAGGTTTTCGCCGCCG CCGCGCTTCAGGCC	Cloning PA3962 in pET23b(+) +

F-pGEX PA3962	GGATCCATGCGCCTTGCCACCCGCG	TEV site + XhoI at 3'- end Cloning PA3962 in pGEX4T-2 + BamHI at 5'-end
R-pGEX PA3962	CCCGGGTCAGTGGTGGTGGTGGTGGTGCTCGAGG GACTGAAAATACAGGTTTTCGCCGGCGGCCGGCTT CAGGCC	Cloning in pGEX4T-2 + TEV site + (His)$_6$ tag + SmaI at 3'-end
F-pGEX-$_{1-29}$PA3962	CGTGGATCCGAGCGACGCGGCCAGATCGACGGTC G	Deletion N-terminal residues (1-29) of PmcY
R-pGEX-$_{1-29}$PA3962	TCGCTCGGATCCACGCGGAACCAGATCCGATTTTG	Deletion N-terminal residues (1-29) of PmcY
F-pGEX-$_{94-138}$PA3962	GCGGACGGCGGCGAAAACCTGTATTTTCAGTCCCT	Deletion C-terminal residues (94-138) of PmcY
R-pGEX-$_{94-138}$PA3962	TTCGCCGCCGTCCGCGCCGACGGCATCCTTGAGCA	Deletion C-terminal residues (94-138) of PmcY
F-pGEX$_{30-93}$PA3962His*6	CGGACGGCTGATCTAGACTAGGGGTCGACTCGAGC GGC	Deletion of (His)$_6$ tag
R-pGEX$_{30-93}$PA3962His*6	TCTAGATCAGCCGTCCGCGCCGACGGCATCCTTGA GCAC	Deletion of (His)$_6$ tag
F-D40xA– PA3962	TCGCATCGCCCCGGCCATCGTCGCCTTCGACGCCA TCGGCC	D40A mutation in PmcY$_j$
R-D40xA– PA3962	GCCGGGGCGATGCGACCGTCGATCTGGCCGCGTC GCTC	D40A mutation in PmcY$_j$
F-D47xA– PA3962	CGCCTTCGCCGCCATCGGCCTGCTGTTCTATATCAT CCCCGGCCTG	D47A mutation in PmcY$_j$
R-D47xA– PA3962	GATGGCGGCGAAGGCGACGATGGCCGGATCGATG CGACCG	D47A mutation in PmcY$_j$
F-D65xA– PA3962	CGGCGTCGCCTTCGCCACCGGCGCCATCTACCTGC CCGAC	D65A mutation in PmcY$_j$
R-D65xA– PA3962	GCGAAGGCGACGCCGAAGGCGATCAGGCCGGGGA TGATATAGAACAGC	D65A mutation in PmcY$_j$

F-pSEVA PA3962	CTAGTGAATTCCCCGGGTGGTCTTGCCGGC	Cloning *pmcY* plus 250pb up stream + EcoRI at 5'-end
R-pSEVA PA3962	ACTGGTACCGGCGGCCGGCTTCA	Cloning *pmcY* plus 250pb up stream + KpnI at 3'-end
F-GFP	AGGGTACCGCTAGCAAAGGAGAA	Cloning *gfp* + KpnI at 5'-end
R-GFP	CCTTCTAGATTAGCAGCCGGATCCT	Cloning *gfp* + KpnI at 3'-end
F-CzcS	ATCCATATGGCCCGCGAGCGGGA	Cloning *czcS* soluble domain + NdeI at 5'-end
R-CzcS	AGTGGATCCTCATTTTCGAACTGCGGGTGGCTCCAG CGCTCGTCGGCGT	Cloning *czcS* soluble domain plus StrepII + BamHI at 3'-end

Figure legends

Figure 1. PmcY is a membrane protein required for imipenem resistance. (A)

Left, hydrophobic surface profile of the AF2 predicted structure for PmcY/PA3962.

Position of Cys23 is indicated. Center and right, ribbon diagram of PmcY with lipobox

amino acids indicated in green. The thiol group of Cys23 would promote acylation

anchoring the protein, right. Notice that independent of the position at the inner or

the outer membrane, PmcY faces the periplasmic space. P= periplasm; M=

membrane; I/O= inside/outside the cell. (B) Analysis of heterologously expressed

PmcY-(His)$_6$ in *E. coli* compartments (cytosolic= C; total membranes= TM;

periplasm= P) by cell fractioning. 10 μg of protein from each fraction obtained from

cells grown in LB media (basal= B; induced = I) were analyzed by SDS-PAGE (left)

and Western blot (right). PmcY-(His)$_6$ expression is apparently increased in basal

conditions and mostly found in the membrane fraction. (C) Serial dilutions (10 μl) of

P. aeruginosa PAO1 (WT), *pmcY::dTn5* and *yiiP::dTn5* cells initially at OD$_{600nm}$=1.0

were spotted on MH-agar supplemented with the indicated imipenem concentration

and incubated for 16 h at 37°C. (D) Complementation of *pmcY::dTn5* restores

imipenem resistance. PA3962 plus 250bp upstream bases were cloned in the low

copy plasmid pSEVA621 containing GFP. This construct was introduced by

electroporation into *pmcY::dTn5*. (E) Lack of Zn^{2+} induced resistance in *pmcY::dTn5*.

Bacterial cells suspensions in LB soft-agar plates were layered on top of the same

media plates containing Zn^{2+} (0.5 mM) or basal labels (0 mM). A paper disc

containing 10 μg of imipenem was located on top of the surface and the inhibition

halos were registered upon 16 hours of incubation at 37 °C. Notice the increased

resistance of WT, and *yiiP::dTn5* and the loss of the effect with *pmcY::dTn5*.

oprD::dTn5 was used as a control.

Figure 2. Conserved residues in the juxtamembrane region/domain (PmcYs)

have a role in Zn^{2+} binding. (A) Left, conserved amino acids profile depicted in the

structure for PmcY/PA3962. Positions colored with maroon indicates high

conservation while cyan points to low conservation. Right, lateral chains of putative

amino acids with metal binding capacity are shown. (B) Up, representation of the

fusion protein GST-PmcYs. Down, metal (MT^{2+}) binding capacity and specificity of

the fusion product. Metal bound to GST was subtracted in each case and

represented 10% of the total metal present in GST-PmcYs. (C) Up, representation

of the fusion protein with conserved amino acids positions indicated with blue lines.

Down, binding capacity of mutants D40A, D47A and D65A. Notice that Zn^{2+} binding

decrease to a half vs. WT. (D) Representation of putative Zn^{2+} binding sites in PmcY.

Zn^{2+} atoms are indicated in red and putative bonds by O-ligands with dotted lines.

In all cases, data are the mean ± SE of three independent experiments. *$P<0.01$; [#],

P, 0.05 from a two-tailed Student's *t* test versus Zn^{2+} (B) or WT (C).

Figure 3. PmcY interacts with CzcS. On-column Zn^{2+}-dependent protein-protein

interaction assay. (A) Upper and lower panel, 100 mg (200 µl) of resin bound with

10 nmole of GST-PmcYj (PmcY$_{30-93}$) were washed with 10 bed volume of Buffer B

(Buffer A + 1 mM EDTA), (Upper panel, lane REDTA$_t$), or Buffer C (Buffer A + 0.1

mM Zn^{2+}) (Lower panel, lane RZn^{2+}$_t$) and incubated with 5 nmole CzcS$_{40-166}$ (lane

+I). The flow through was collected (FT) and the resin washed 4X with 2 bed volumes

of Buffer A (lanes W1-4). Finally, the resin was resuspended at the initial volume

(lane Rf) and 25 µl of each fraction analyzed by SDS-PAGE 12% and CBB staining.

Notice that PmcY-CzcS interaction requires Zn^{2+} bound to PmcY (black asterisk).

(B) Protein interaction analyses of PmcY mutants. EDTA treated (upper panel) or

Zn^{2+}-loaded (lower panel) GST-PmcY mutants (5 nmole D40A, 1 nmole D47A, 2.5

nmole D65A) were incubated with 5 nmole CzcS. 25 µl of fractions from resin treated

(lanes REDTA$_t$ and RZn^{2+}$_t$), incubated with CzcS (lanes +I) and final (lanes Rf)

samples were analyzed as in (A). Black asterisk denotes the Zn^{2+}-dependent

interaction and white, the lack of it.

Figure 4. PmcY-YiiP-like operons coevolved with OprD and likely CzcS in

Pseudomonadales. Bioinformatic analyses. (A) Two operon configurations

involving PmcY homologs were searched in complete bacterial genomes by

multigene Blast using protein sequences; *pmcY-yiiP*-like operon (blue) and non-

pmcY-yiiP-like operon (white). Parallel searches by RBH were done with protein

sequences coded by *czcS* (green) and *oprD* (yellow). (B) Venn diagram showing the

distribution of organisms with detected CzcS (green) and OprD (yellow) between the

two configurations of PmcY (*pmcY-yiiP*-like operon, in blue vs. *non-pmcY-yiiP*-like

operon, in yellow). Notice the lack of association between *non-pmcY-yiiP*-like operon

organisms (226) with CzcS and OprD (1 and 13 organisms among 226,

respectively). (C) Phylogenetic analysis of the organisms with PmcY homolog

detected in the genome. Most species with *pmcY-yiiP*-like operon (blue strip)

belongs to the order of Pseudomonadales (ocher). Coevolution of OprD and CzcS

is depicted with yellow and green strips labels, respectively.

Figures

Figure 1

Figure 2

Figure 3

Figure 4