A periplasmic metallochaperone (PmcY) couples Zn^{2+}-transport to sensing in

Pseudomonas aeruginosa

Running title: An anchored membrane protein couples YiiP mediated Zn^{2+} efflux to

Zn^{2+} sensing in *Pseudomonas aeruginosa*

Paula Mihelj[1], Tomás Moreyra[1], Monserrat Olea-Flores[2], María Elena Carrizo[3],

Teresita Padilla-Benavides[2], Daniel Raimunda[1]

[1]Instituto de Investigaciones Médicas Mercedes y Martín Ferreyra, INIMEC.

[2]Department of Molecular Biology and Biochemistry, Wesleyan University,

Middletown CT, 06459, USA.

[3]Centro de Investigaciones en Química Biológica de Córdoba CIQUIBIC.

[1,3] CONICET-Universidad Nacional de Córdoba. Argentina.

Abstract

Pseudomonas aeruginosa thrive to survive in harsh conditions imposed by the host. We have previously described two Zn^{2+}-transporting members of the Cation Diffusion Facilitator family, YiiP and CzcD, inhibiting susceptibility to imipenem by decreasing the expression of the outer membrane porin OprD (A. Salusso and D. Raimunda, Frontiers in cellular and infection microbiology 7:84, 2017, https://doi.org/10.3389/fcimb.2017.00084). Here we provide evidence that a protein encoded in *yiiP´*'s operon, PA3962, is fundamental to the coupling of these processes. Immunodetection assay indicates that PA3962 locates in membranes. Supporting a role in *oprD* regulation, a PA3962 insertional mutant has a significant increase of *oprD* expression levels and imipenem sensitivity, which is suppressed by gene complementation but not in presence of Zn^{2+}, as opposed to the YiiP mutant. We identified 2 pairs of conserved acidic residues in a hydrophobic juxtamembrane domain. Metal binding specificity and stoichiometry was explored in wild-type and mutant versions of these. Zn^{2+} appears as the cognate metal of PA3962, with residues D40, D47 and D65 required for its coordination. As the periplasmic Zn^{2+}-sensor CzcS regulates *oprD* expression, the interaction with it was analyzed *in vitro*. Interaction was Zn^{2+}-dependent, and mutations of D47A or D65A abolished it. We propose a role for PA3962, hereafter periplasmic metallochaperone of YiiP (PmcY), in the context of Zn^{2+} signaling pathways in *P. aeruginosa*. The relay YiiP–PmcY, supplies Zn^{2+} and activates CzcS/CzcR, down-regulating the transcription of the imipenem-permeable OprD. This mechanism would allow *P. aeruginosa* to put a

brake on unspecific mechanisms for micronutrient uptake, with potential xenobiotic

entry, while the cytosolic Zn^{2+} quota is still sufficient.

Introduction

Pseudomonas aeruginosa is a facultative anaerobic gram-negative bacteria and a

prevalent opportunistic pathogen in respiratory infections of patients diagnosed with

cystic fibrosis (1). It is considered an oligotrophic organism with a highly dynamic

capacity for antibiotic resistance, metabolic adaptation, or phenotypical switching,

which sometimes translates to fitness costs (2). Metabolic adaptations and antibiotic

resistance involve optimization of Zn^{2+} usage in niches with low bioavailability (3). In

the first case, the Zn^{2+}-dependent (C+) to Zn^{2+}-independent (C-) isoform switching

of ribosomal subunits, increases the intracellular Zn^{2+} pool in this condition (4). In the

second, Zn^{2+} induces carbapenem resistance by repressing the transcription, and

then the antibiotic influx, of the outer membrane porin Occ1, also known as OprD

(5). As carbapenem-resistant OprD mutants show enhanced fitness *in vivo* (6) we

hypothesize that this "master of survival", could benefit from a last resort-type of

intracellular Zn^{2+} availability by finely tuning Zn^{2+}-sensing mechanisms. In this way

this opportunistic pathogen could adapt metabolically and become resistant to

carbapenems, avoiding the pay-offs of a fitness cost imposed by nutritional

immunity(7).

We have previously shown that the Zn^{2+} exporters of the Cation Diffusion Facilitator

family, YiiP and CzcD regulate levels of OprD (8). Single insertion mutants of these

genes showed increased sensitivity to imipenem, and increased Zn^{2+} sensitivity

when introduced into double mutant backgrounds where other Zn^{2+} resistance mechanisms were also affected. Considering studies by others showing that transcription of *czcD* responds to Zn^{2+} levels in the growth media with a slow temporal activation (9) a role in Zn^{2+} detoxification might be considered. However, the fact that *yiiP* transcripts are not upregulated by Zn^{2+}, opens new possibilities for the role of this Zn^{2+}-transporter in Zn^{2+} mediated signaling.

Zn^{2+} sensing mechanisms in *P. aeruginosa* are well known. At the cytoplasmic level, CadR, a member of the MerR transcription factor family, controls the expression of CadA, also known as ZntA, a Zn^{2+}-transporting P_{IB}-ATPase (8, 10). Interestingly, it has been proposed that under high Zn^{2+} conditions, CadA is required for timely expression of CzcCBA, also known as HME-RND, a tripartite Zn^{2+}-export system that promotes efflux from the periplasm to the extracellular space (10). This implies that the specific CadA Zn^{2+}-export is coupled to Zn^{2+} sensing in the periplasm. In line with this, extracellular Zn^{2+} bioavailability in *P. aeruginosa* is primarily detected in the periplasm via the two-component system (TCS) CzcS-CzcR. CzcS senses periplasmic Zn^{2+} by coordinating the metal in the periplasmic domain (11), which is transduced to cytosolic domains promoting autophosphorylation of the histidine kinase domain and finally, the transphosphorylation to CzcR. Phosphorylated CzcR (P-CzcR) has dual effects on transcription of genes in the CzcS-CzcR regulon (12). For example, P-CzcR up-regulates transcription of HME-RND, thus counteracting a potential Zn^{2+} overload (9, 10) in the periplasm. However, P-CzcR affects negatively the transcription of genes involved in pyocyanin synthesis and of the outer membrane porin *oprD* (11, 12).

Here we present evidence suggesting that a periplas<u>mic</u> metallochaperone of <u>Y</u>iiP

(PmcY/PA3962) is required for *oprD* repression and thus, for Zn^{2+} sensing in the

periplasm. Our data shows that PmcY is a Zn^{2+}-metallochaperone that interacts with

CzcS, its putative acceptor or target protein. We also propose that the intracellular

Zn^{2+} efflux to the periplasm mediated by YiiP-PmcY has coevolved in

Pseudomonadales as a mechanism for limiting the fitness cost of adapting to

common environmental or host-imposed scenarios with low Zn^{2+} bioavailability, by

repressing an unspecific outer membrane porin.

Materials and methods

Bacterial strains, plasmids and growth media conditions

Pseudomonas aeruginosa MPAO1 wild-type (WT), and the insertion mutants

yiiP::dTn5 and *pmcY*/PA3962*::dTn5* were obtained from the *P. aeruginosa* Mutant

Library at the University of Washington ((13); Table 1). For complementation of the

pmcY/PA3962*::dTn5* strain PA3962 locus plus the 250bp upstream bases was PCR

amplified and cloned in pSEVA621 using EcoRI and KpnI sites (14). The sequence

for *gfp* was introduced at the 3'-end as a fusion in order to promote protein stability

using KpnI and XbaI sites. Electrocompetent cells were transformed by

electroporation as described (15) and selected on Luria–Bertani (LB) agar plates

supplemented with gentamicin (30 µg/ml). Double mutant *oprD-pmcY* (DM3004)

was obtained via double recombination events following methods previously

described (16). The up and down-stream 500 bp fragments of PA3962 were

amplified and stiched by PCR. After purification and digestions with EcoRI and

HindIII restriction enzymes the fragment was ligated into the suicide plasmid pEX18-

Gm. Electrocompetents *oprD::dTn5* cells were transformed with this plasmid and

recombinants were selected on LB-agar supplemented with 60 µg/ml gentamicin.

Several colonies were grown in LB supplemented with sucrose 5%. The PA3962

locus deletion was confirmed by PCR using genomic DNA of isolated colonies grown

in LB + sucrose 5%, supplemented with tetracycline (30 µg/ml). The primers used in

these and other procedures are listed in Table 1.

Plasmid constructions and protein expression and purification

In a first attempt to yield a whole version of PmcY by heterologous expression in *E.*

coli PA3962 locus was cloned from *P. aeruginosa* MPAO1 gDNA and inserted by

ligation in the expression plasmids pET23b(+) (Novagen) or pGEX4T-2 using

primers described in Table 1. These constructs contained at C-terminal a

hexahistidine ($(His)_6$) tag preceded by Tobacco Etch Virus (TEV) protease

recognition site. Due to low yield of the protein and guided by bioinformatics analyses

we expressed the juxtamembrane domain of PmcY ($PmcY_j$) as a GST fusion protein

by deletion of N- and C-termini and $(His)_6$ tag of PmcY by the one-step site-directed

deletion PCR protocol (17). Site directed mutagenesis of conserved residues were

obtained using the same procedure. Similarly, the soluble domain (SD) of CzcS

(amino acids 40-166, $CzcS_{SD}$) was cloned by PCR amplification from MPAO1 gDNA

and ligated in pET16bStrep (Table 1) with a twin-strep tag at N-terminal. All

constructs and mutants were confirmed by DNA sequencing.

For protein expression and purification *E. coli* BL21(DE3)pLysS were transformed

with one of the above constructs. After selection cells were grown in LB overnight at

37°C on a shaker at 200 rpm. Next day 1-2 L of LB media with antibiotic was

inoculated with cells to reach OD_{600}= 0.1 and grown in the same conditions till early

log-phase. At this point cultures were induced with 0.15 mM IPTG for 16 h at 20°C.

For purification of GST and GST fusion versions of $PmcY_j$ cells were resuspendend

in 50 ml of Tris-HCl 20 mM pH 7.5, NaCl 500 mM, TCEP 0.25 mM, 0.1% Triton X-

100 (Buffer A) plus PMSF 1 mM, and lysed in a cell disruptor EmulsiFlex C3

(Avestin). After ultracentrifugation at 100,000 *g* the supernatant was incubated with

1 ml of Glutathione Sepharose 4B or Fast Flow (GE Healthcare) for 1 h at room

temperature. Resin was packed by gravity in a polypropylene column (Bio-Rad) and

washed with 30 resin volumes of buffer A. The fusion protein was eluted with buffer

A containing 40 mM glutathione. The soluble domain for CzcS was expressed and

purified as described earlier (11) with minor modifications. After growth and

induction cells were lysed as above and ultracentrifuged at 100,000 *g*. The

supernatant was incubated for 1 h at room temperature with 0.5 ml of Strep-Tactin

resin (IBA Lifesciences) equilibrated with buffer A for 1 h and eluted after washing

with 30 resin volumes. Bound protein was eluted with 10 resin volumes of buffer A

containing 50 mM of D-biotin. In both cases elution fractions were analyzed by SDS-

PAGE 12%, Coomassie brilliant blue staining and Western blot using $(His)_6$-tag

antibody (pAb, Rabbit, Genscript), GST-tag antibody (pAb, Rabbit, Genscript) or

StrepMAB-Classic antibody (IBA Lifesciences) as primary antibodies. Those

fractions containing protein were pooled for dialysis in buffer A using a dialysis tubing

(SnakeSkin, Thermo Scientific) with 3.5 kDa cut-off. After addition of 10% glycerol,

protein concentration was estimated by Bradford (Bio-Rad) and 280 nm lectures in

NanoDrop (Thermo Scientific). Protein samples were frozen and maintained at -

80°C for later use. For in-column interaction assays the bait protein, GST-PmcY$_j$ WT

and mutant versions, were not eluted from the resin and kept at 4°C in the dark for

later use and store for no more than 2 weeks. In this case protein concentrations

were determined by densitometry. Denatured resin samples were run in SDS-PAGE

12%, stained with Coomassie brilliant blue (CBB) and images analyzed with ImageJ

(18). For calibration, bovine serum albumin (Sigma-Aldrich) standards were run in

the same gel.

Antibiotic sensitivity assays

Susceptibility to imipenem (Sigma-Aldrich) and ciprofloxacin was assessed by the

agar dilution or the inhibition halo methods in Müller–Hinton (MH) medium as

described before (8). For agar dilution overnight grown cells were serially diluted

from an initial OD$_{600}$= 1 and then 10 µl were spotted on MH agar plates with different

antibiotic concentrations. For inhibition halos, a suspension of cells at OD$_{600}$=0.05 in

soft agar were layered on top of MH agar plates. After solidifying, 7-mm diameter

filter paper discs embedded with the indicated amount of imipenem were located on

top of the soft agar. In all cases growth or inhibition was evaluated after overnight

incubation at 37°C. Analyses of the imipenem MIC values found by agar dilution

method (ADM) *vs.* our previously data published using Epsilon-test method

(Biomerieux Diagnostics) yielded a lineal correlation between MICs following

y=8.082 x with and R² = 0.797, where *y* equals the apparent ADMs MICs, and *x* the

Epsilon-test ones. Thus, apparent imipenem concentrations are informed in this

work.

Quantitative RT-PCR analysis

Overnight cultures from *P. aeruginosa* WT, *yiiP::dTn5* and *pmcY::dTn5* were diluted

to an OD_{600} of 0.05 in LB and incubated until they reached an OD_{600} of 0.6-0.8. Total

RNA was extracted with TRIzol™ Reagent (Invitrogen) according to the supplier's

instructions and treated with RNase-free DNase I (TransGen Biotech) for 2 h at 37°C

to remove residual DNA. After phenol/chloroform extraction, RNAs were precipitated

with ethanol and the pellets resuspended in RNase-free water. For cDNA synthesis,

200-300 ng of RNA was reverse transcribed using random primers and *EasyScript*®

Reverse Transcriptase (TransGen Biotech) according to the manufacturer's

instructions. Quantitative PCR was performed using FastStart Universal SYBR

Green Mastermix (Rox) (Roche), according to the supplier's instructions. Fold

change expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method as previously

described (19). Results were normalized with the *rpsL* gene (PA4268). Primers used

for qPCR studies are listed in Table 1.

Metal binding assays

Transition metal (TM) binding capacity of GST fusion proteins was evaluated using

the chromogenic dye 4-(2-Pyridylazo) resorcinol (PAR, Tokyo Chemical Industry).

For incubation of the proteins with TM we followed described protocols for in-solution

(20) or in-column (21) assays with some modifications. To assess binding to proteins

10-20 µM samples (or the mass bound to resin to yield this concentration before

adding PAR) were incubated for 30 min with 5 molar excess of TM in buffer A. For

in-solution binding assays unbound TM were separated by exclusion

chromatography with Sephadex G25-80 (Sigma-Aldrich) and for the in-column type

the resin was washed with 10 volumes of buffer A to remove unbound TM. Before

TM quantification protein concentrations were estimated as described earlier and

samples containing proteins in solution or bound to resin were treated with 0.4-0.8

mg/ml of proteinase K. For TM quantification, PAR was added to a final

concentration of 100 µM and read in microplates at 500 nm. Calibration curves were

done in parallel for each TM and the molar ratio of protein to TM calculated. In both

conformations GST alone or GST fusions were run in parallel to detect TM bound in

proteins as purified. These values represented no more than 10% of those found in

TM incubated samples and were subtracted in all cases. For the quantification of

Mn^{2+} the buffer pH was modified from 7.5 to 11 as the colorimetric reaction of this

TM is more sensitive. Therefore, after the digestion reaction with proteinase K, the

samples were evaporated completely and resuspended in 200 mM CAPS (3-

(Cyclohexylamino)-1-propanesulfonic acid) buffer pH 11.

Protein localization

The *E. coli* strain BL21(DE3)pLysS transformed with the plasmid pET-23b(+)-

PA3962, which expresses the PA3962 protein fused to a 6-histidine tag, was grown

in 12.5 ml of LB medium at 37°C for 16 h. This culture was placed in two flasks with

500 ml of LB medium and grown with shaking (200 rpm) at 37°C for 1 h. One of the

cultures was induced with 0.15 mM of IPTG and the other was used as a control

(basal). After 3 h of growth the cells were harvested by centrifugation at 4,000 *g*, 4°C

for 10 min. Cell pellets were washed in ice-cold 30 ml of 50 mM Tris-HCl pH 7.5 and

0.3 M sorbitol (buffer S) 3 times, with addition of 10 µM EDTA in the second wash.

Cells were pelleted by centrifugation at 9,000 *g*, 4°C for 10 min. Finally, cells were

resuspended in 30 ml of ultrapure water and incubated for 5 min at room

temperature. They were then centrifuged at 11,000 g, 4°C for 30 min. The

supernatants were collected and centrifuged again to eliminate any possible

contamination. After this they were considered the periplasmic fractions. For

cytosolic and total membrane fractions cell pellets were resuspended in 7 ml of 10

mM Tris-HCl pH 7.5 and disrupted at 4°C using a tip sonicator (Vibra Cell, Sonics &

Materials, Inc.) with 6 cycles of 20 seconds on and 60 seconds off pulses. The lysate

was centrifuged at 6,000 g for 10 min at 4°C to remove undisrupted cells and

centrifuged again at 100,000 g for 30 min at 4°C. The supernatants were recovered

and classified as cytoplasmic fractions. The resultant pellets were resuspended in 5

ml of 10 mM Tris-HCl pH 7.5 and classified as total membrane fractions. The protein

concentration of all fractions was quantified by Bradford assay, and those with a

value less than 1 mg/ml were concentrated using centrifuge filtration tubes

(Centricon, Millipore). Samples of all fractions were prepared at concentrations of 1

mg/ml and 5X Laemmli sample buffer was added. A final total protein mass of 10 μg

was loaded per lane on 14% SDS-PAGE gels in duplicate, one for CBB staining and

the other for Western blot. (His)$_6$-tag antibody (pAb, Rabbit, Genscript) and anti-

rabbit antibody labeled with horseradish peroxidase were used to detect expression

in fractions. Images were taken using Amersham Image Quant 800 Fluor (Cytiva).

Protein-protein interaction assay

To evaluate the interaction between PmcY and CzcS, the recombinant proteins

GST-PmcY$_j$ (WT and mutants D40A, D47A, and D65A) and CzcS$_{SD}$ were used. A

100±20 mg sample of purified WT or mutant versions of GST-PmcY$_j$ were treated

with EDTA-Tris. For this, Glutathione Fast Flow (GE Healthcare) resin with bound

protein was incubated with 1 ml of buffer A plus 1 mM EDTA-Tris in a 10 ml

polypropylene chromatography column (Bio-Rad). Three ml of buffer A were used to

remove residual EDTA from column. The interaction was tested in two conditions,

with GST-PmcY$_j$ loaded with (+Zn^{2+}) and without (-Zn^{2+}) ZnCl$_2$. For this, resin was

incubated for 5 min with 500 µl of 100 µM ZnCl$_2$ in buffer A or just buffer A,

respectively. The CzcS$_{SD}$ protein was used in solution and previously treated with

0.5 mM EDTA-Tris for 30 min in ice. EDTA-Tris was removed with PD-10 column

equilibrated in buffer A. In-column GST-PmcY$_j$ and mutant versions, in its two

conditions (+Zn^{2+}) and (-Zn^{2+}), were incubated with 25 µM CzcS$_{SD}$, added in 250 µl,

during two passes through the column (Flow through, FT) and 4 washes (W1-4) were

performed with 250 µl of buffer A. Finally, the resin was resuspended in 250 µl of

buffer A and samples prepared with 5X Laemmli loading buffer for SDS-PAGE 12%

analyses. After CBB staining images were taken using Amersham Image Quant 800

Fluor (Cytiva).

Bioinformatics analyses

To analyze the conservation of residues in a putative PmcY family, homologs were

searched in a local database of all protein sequences of RefSeq reference and

representative genomes protein sequences (Refseq_select_prot) (January 2024) by

the reciprocal best hit (RBH) method (19). Four hundred and sevenunique organisms

having PmcY-like members with unique accession numbers were found. Multiple

sequence alignment (MSA) was done in Clustal Omega (22) using a similar number

of sequences and the same MSA was used in ChimeraX (23) to create the amino

acids conservation map in the putative 3D AF structure of PA3962 (AF-Q9HX54-F1).

This number of organisms were used later to compare and create a list of non-

redundant bacterial organisms having, or not, the *pmcY-yiiP* architecture. The

search of the *pmcY-yiiP* operon was done using Multigenblast (24). Similar RBH

searches were performed for OprD and CzcS homologs in the same data base. After

removal of duplicated organism names, a 4-group diagram was prepared with these

organisms (https://bioinformatics.psb.ugent.be/webtools/Venn/) and a phylogenetic

tree was constructed for visualization of coevolution in iTOL (25).

Results

PA3962 is a membrane protein

PA3962 is part of a putative operon with YiiP (PA3963). Previous bioinformatics

studies of identified *P. aeruginosa* secreted proteins pointed that the N-terminal

sequence in PA3962 is a type II signal peptide (26). These results were supported

with a plasmid library screening in *E. coli* where several PAO1 genomic DNA

fragments, including PA3962, fused to truncated alkaline phosphatase (PhoA) drove

its periplasmic localization. Signal peptides are recognized by different secretion

systems in bacteria and classified in different groups. The type II are related to

proteins anchored to the membrane, also called lipoproteins. A consensus sequence

called "lipobox" is found in the type II signal peptide, which allows for molecular

recognition by the machinery involved in processing and finally anchoring the protein

to the inner or outer membrane. For this, first a conserved cysteine residue in the

lipobox is diacylated with diacylglycerol. Second, the signal peptide is removed by a

specific protease leaving free and reactive the amino group of the cysteine for a final

third acylation (27). Figure 1, shows the predicted structure of PA3962 by

AlphaFold2 (AF2) (28) showcasing the colour–coded hydrophobic profile generated

in ChimeraX and position of Cys23 in the lipobox. The ribbon structures depict the

putative position of the protein facing the periplasmic space after the anchoring

occurs. In order to confirm the membrane localization of PA3962 we expressed the

whole protein with a C-terminal (His)$_6$ tag in *E. coli* in basal and induced conditions.

After fractioning cells, cytosolic, total membranes, and periplasmic samples were

analyzed by SDS-PAGE and Western blot using anti-(His)$_6$ as primary antibody (Fig.

1B). Protein detection was evident in most of the fractions and signal was increased

under basal conditions, pointing to an inhibitory effect of the translation in presence

of the inducer when the full length protein is overexpressed. Most importantly the

detection in total membrane fraction was observed. As the mass of protein loaded in

all lanes was similar, and assuming an equal contribution of each compartment in

cell mass, a qualitative conclusion is that PA3962 is a membrane protein. PhoA

detection in *E. coli* requires folding in the periplasm, thus this orientation is the most

likely scenario in *P. aeruginosa* and we renamed PA3962 as <u>P</u>eriplas<u>m</u>ic

metallochaperone of <u>Y</u>iiP, PmcY.

309   PmcY is involved in *oprD* mediated imipenem susceptibility

310   As previous results indicated a role for YiiP in Zn^{2+}-dependent antibiotic susceptibility

311   involving OprD and altered membrane permeability (8) we next analyzed if absence

312   of PmcY yielded similar effects. The minimal inhibitory concentration (MIC) for the

antibiotic in Tn5-generated mutant strains (Fig. 1C) YiiP and PmcY mutants showed

that both are sensitive to imipenem *vs.* WT. Complementation of PmcY strain was

achieved by transformation of mutant cells with the low copy plasmid pSEVA621

(14) carrying PA3962 locus under transcriptional regulation of 250 bp upstream

sequence (Fig. 1D). We have previously demonstrated that lack of YiiP induces

expression of OprD (8). If PmcY is involved in imipenem susceptibility the expression

of *oprD* should be altered compared to the WT strain. The results of *oprD* locus

expression between PmcY and YiiP mutant strains (*vs.* WT) correlates to the

imipenem susceptible phenotype of both (Fig 1E). These results indicated that the

molecular mechanism leading to the loss of imipenem susceptibility involves the lack

of Zn^{2+} sensing capacity. This loss would involve specifically Zn^{2+} transport from

cytosol to the periplasm. Thus, we hypothesized that both, YiiP and PmcY, mutants

would remain sensitive to imipenem at high Zn^{2+} content (0.5 mM) in the growth

media. However, only PmcY mutant strain showed the same imipenem susceptibility

in presence of high levels of Zn^{2+} indicating a key role for PmcY. The apparent

dispense of YiiP to sense Zn^{2+} at high extracellular Zn^{2+} conditions, correlates with

a save-for-later strategy to sustain steady-state levels of the intracellular quota of

this vital element. To test the Zn^{2+}-specificity of the mechanism we tested also

ciprofloxacin sensitivity (Fig. 1F right). The lack of Zn^{2+} complementation to this

antibiotic susceptibility observed in both mutants, points to the involvement of

another mechanism likely related to membrane permeability of ciprofloxacin and

other more lipophilic antibiotics, as suggested previously (8). To discard any

agonistic effects between OprD and PmcY related to membrane permeability, we

generated the double mutant *oprD-pmcY* (DM3004) and evaluated its imipenem

susceptibility. Figure 1G shows that both strains, *oprD::Tn5* and DM3004, behave

similarly. These results indicate that PmcY is required for Zn^{2+} sensing in the

periplasm and that *oprD* is regulated by PmcY activity. Thus we focused on the

pivotal role of PmcY as part of a sensing mechanism leading to decrease intrinsic

imipenem susceptibility, and hypothesize that PmcY should bind Zn^{2+} and interact

with a sensor to likely transfer the metal to it.

PmcY binds Zn^{2+} specifically through conserved aspartate residues

We hypothesized that PmcY plays a role as metallochaperone. These function in

metalation of proteins or in detoxification pathways, and in both cases the

commonality is their capacity to bind a specific transition metal through conserved

residues. A multiple sequence alignment of PmcY homologs obtained by a reciprocal

best hit search in complete bacterial genomes allowed us to identified conserved

residues (Fig. 2A, maroon). Most of these would be located in a periplasmic domain

juxtaposed to the membrane. Conserved amino acids are all acidic, with oxygen (O)

ligands of lateral chains of aspartates and glutamates. After several failed trials of

full length PmcY expression and purification to get enough protein for biochemical

characterization, we opted to fuse this domain (amino acids 30 to 93), which we

called PmcY juxtamembrane (PmcY$_j$), to GST. The fusion protein was purified and

its capacity and specificity to bind transition metals analyzed. For this, five molar

excess of different metals were incubated with the fusion protein (Fig. 2B-C) and

after unbound metal was separated by exclusion chromatography (Fig. 2B) or

washed (Fig. 2C), protein was quantified and metal bound to protein detected as

described. This strategy allowed us to confirm that Zn^{2+} is the cognate metal for

PmcY and that the stoichiometry of binding is 2 Zn^{2+} per PmcY molecule. To identify

possible binding sites, we analyzed the Zn^{2+} binding of the single mutants of highly

conserved positions with O ligands; *i.e.*, D40A, D47A and D65A. All these mutations

lowered the binding stoichiometry to 1, indicating their likely participation in two Zn^{2+}

binding sites. A closer look to the distances between O atoms in lateral chains of the

PmcY structure, underscores residues D47-D65 as candidates for one Zn^{2+} binding

with the archetypical tetrahedral geometry (Fig. 2D).

PmcY interacts with the periplasmic Zn^{2+} sensor CzcS

Another property which defines a metallochaperone is the interaction with its

target(s) protein(s). This is key in order to promote the ligand transfer reaction, and

the metal transfer, between the donor, the metallochaperone, and the acceptor, the

target protein. Thus an interaction between PmcY and CzcS could promote Zn^{2+}

sensing by means of a ligand transfer reaction between partners. In order to test

this, we expressed and purified the periplasmic soluble domain (SD) of CzcS (amino

acids 40 to 166, named CzcS$_{SD}$) as described by Wang et al. (11) with a twin Strep-

tag at the N-terminal end. The protein preparation was treated with EDTA to

eliminate Zn^{2+} carried-over during purification, and then incubated with GST-PmcYj,

in two conditions, EDTA-treated or loaded with Zn^{2+}. After collecting flow through

and washes we detected the presence of CzcS in the final resin (Rf) only when GST-

PmcY$_j$ was loaded with Zn^{2+} (Fig. 3A, black asterisk). In order to assess how Zn^{2+}

occupancy affected this interaction, GST-PmcY$_j$ mutants, D40A, D47A and D65A

were analyzed similarly. The Zn^{2+}-dependent interaction was lost only in D47A and

D65A mutants. This data supports the hypothesis in which these acidic residues

might coordinate Zn^{2+} in one binding site. Mutation D40A did not preclude the

interaction in presence of Zn^{2+}.

PmcY-YiiP operon coevolved with OprD and CzcS in Pseuodomonadales

The data presented suggest that YiiP, PmcY, CzcS/CzcR participate in a sensing

mechanism coupling Zn^{2+} efflux via YiiP, to decrease imipenem suceptibility (as *oprD*

expression is downregulated). If this mechanism is conserved among gram-negative

bacteria, we hypothesized that PmcY homologs found in other organisms would

present the same operon architecture as in *P. aeruginosa* MPAO1. A multigene

BLAST search did not support this. However, organisms presenting the operon

architecture were mostly of the Pseudomonadales order. Most importantly,

contrasted with organisms with non-*pmcY-yiiP* operon detected, *oprD* and *czcS* were

mostly distributed in organisms with the operon *pmcY-yiiP*; *i.e.*, the

Pseudomonadales order. The lack of CzcS in a high number of members in this

clade is surprising. However, the periplasmic Cu^{+} sensor CopS has also been

proposed to downregulate *oprD* transcription (29). Altogether these data support a

likely coevolution of these genes. We discuss in the next section what could be the

selection pressure of the coevolution and the physiological role in a context of Zn^{2+}

bioavailability.

Discussion

The association between the Zn^{2+} induced repression of *oprD* transcription, and the

Zn^{2+} induced resistance to imipenem is long recognized (30). Moreover, several

studies have used this response to dissect important molecular and mechanistic

aspects of the CzcS-CzcR signaling pathway (11, 31) in the periplasmic space. Here we present evidence pointing to the participation of YiiP and PmcY in periplasmic Zn^{2+}-sensing mechanisms. In particular, we propose that both players affect *oprD* transcription, and likely other genes, by promoting Zn^{2+}-sensing; *i.e.*, the metalation of the Zn^{2+} sensor CzcS. We provide evidences supporting our model, showing that PmcY binds Zn^{2+}, with high specificity and with a molar stoichiometry of 2 Zn^{2+} per molecule. Metal coordination is given by mean of conserved Asp residues. Phenotypical analyses showed that different to WT and YiiP mutant, the mutant strain of PmcY lacks the Zn^{2+}-dependent response to imipenem susceptibility likely by a lack or reduced Zn^{2+}-sensing capacity of CzcS. These results lead us to propose PmcY as a Zn^{2+}-metalochaperone.

Many of the players involved in orchestrating Zn^{2+} homeostasis in *P. aeruginosa* have been described (32). High Zn^{2+} bioavailability is sensed by periplasmic and cytosolic sensors, CzcS and CadR, respectively which lead to transcription of high capacity Zn^{2+} transport systems, such as HME-RND and P_{IB}-type ATPases (9, 10). The integration of these compartmentalized sensing mechanisms can be interpreted considering electrochemical Zn^{2+} gradient. As Zn^{2+} entries and accumulates first in the periplasm transcription of HME-RND are first expected. However, free Zn^{2+} is almost negligible in cells (33) and a role for cytosolic Zn^{2+} exporters have been proposed in periplasmic Zn^{2+} sensing mechanisms (10). This is important as it suggests the participation of Zn^{2+}-transport systems allowing cross-talk between cellular compartments to promote intracellular Zn^{2+} exit even when extracellular Zn^{2+} could turn into lower levels.

Could a similar scenario, involving Zn^{2+} transport systems, participate in cross-talk

between compartments when Zn^{2+} bioavailability is low? Considering the risk of

losing Zn^{2+} in such scenario, the process should be accompanied of a "favourable"

fitness cost. We proposed that Zn^{2+}-export by YiiP is linked to Zn^{2+}-sensing by CzcS

with the final outcome of delaying *oprD* expression, which role allows entry of

nutrients, but also xenobiotic molecules such as antibiotics. Thus this

hypersensitivity to sense intracellular Zn^{2+}, when scarce but still available, would be

beneficial for this opportunistic pathogen as *oprD* induction is still repressed.

HME-RND also is part of the CzcS-CzcR regulon (31) and Zn^{2+}-binding to CzcS

upregulates the transcripts levels. Even when transcription could be triggered by

YiiP-PmcY activity we speculate that low Zn^{2+} availability is not sufficient to promote

HME-RND activity as this transport system would have similar biochemical

properties to CadA; i.e.: high transport capacity but low metal affinity. Incidental Zn^{2+}-

entry via OprD has been suggested (34). Although counterintuitive, the up-regulation

of HME-RND by CzcS activation, would prepare the cell for a potential Zn^{2+} overflow

in the periplasm with minimal basal activity of HME-RND until Zn^{2+} bioavailability

increases. However, evidence to support this is still missing and the lack of Zn^{2+}

resistance shown by OprD mutants contradicts this role (31).

Lastly, and considering our working model, although it is tempting to hypothesize

that a Zn^{2+}-basic amino acid abduct could be the physiological substrate transported

by OprD, the metal-chelating molecule EDTA was not able to outcompete

arginine/lysine transport by OprD in *in vitro* studies (5). Nonetheless, considering the

Zn^{2+} binding capacity of several xenobiotics, including antibiotics, and the evident

lack of specificity of *oprD* for carboxylated substrates, its overexpression could be

viewed as a "razors-edge"-like situation promoting Zn^{2+} uptake linked to the

xenobiotc molecule carrying the ion. Thus, braking the system could be beneficial

until Zn^{2+} reservoirs are downsized to the limits of a "go-hunting" for survival

condition.

Conflict of interest statement

All authors declare that research was conducted in the absence of any commercial

or financial relationships.

Author contributions

Conceived and designed the experiments: D.R., P.M.

Analyzed the data: D.R., P.M., T.M., M.E.C.

Contributed reagents/samples/analysis/tools: D.R., M.E.C., M.O.-F., T.P.-B.

Wrote or edited the manuscript: D.R., P.M.

All authors read and approved the final manuscript.

Funding

This work was supported by the Agencia Nacional de Promoción Científica y

Tecnológica (ANPCyT) PICT-2018-03139 to D.R., and PICT-2021-0147 to M.E.C.

D.R. and M.E.C. are Consejo Nacional de Investigaciones Científicas y

Tecnológicas (CONICET) investigators. P.M. is recipient of a CONICET doctoral

fellowship. T.M. is recipient of a doctoral fellowship from ANPCyT (PICT-2018-

03139). T.P.-B. was supported by Wesleyan University institutional funds.

Abbreviations

478 TM, transition metal; CDF, cation diffusion facilitator; IPTG, Isopropyl β-D-1-

479 thiogalactopyranoside; PAR, 4-(2-Pyridylazo) resorcinol; HME-RND, heavy metal

480 exporter of the resistance-nodulation-division family.

Acknowledgments

483 We thank Camila Semeniuk, Dr. Laura Montroull and Dr. Andrea Pellegrini for their

484 invaluable help during this work.

References

1. Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JAJ, Sommer LM, Molin S, Johansen HK.
2021. Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis.
Nature Reviews Microbiology 19:331-342.
2. Zhu M, Dai X. 2024. Shaping of microbial phenotypes by trade-offs. Nat Commun 15:4238.
3. Panina EM, Mironov AA, Gelfand MS. 2003. Comparative genomics of bacterial zinc
regulons: enhanced ion transport, pathogenesis, and rearrangement of ribosomal proteins.
Proc Natl Acad Sci U S A 100:9912-7.
4. Pederick VG, Eijkelkamp BA, Begg SL, Ween MP, McAllister LJ, Paton JC, McDevitt CA. 2015.
ZnuA and zinc homeostasis in Pseudomonas aeruginosa. Sci Rep 5:13139.
5. Eren E, Vijayaraghavan J, Liu J, Cheneke BR, Touw DS, Lepore BW, Indic M, Movileanu L, van
den Berg B. 2012. Substrate specificity within a family of outer membrane carboxylate
channels. PLoS Biol 10:e1001242.
6. Skurnik D, Roux D, Cattoir V, Danilchanka O, Lu X, Yoder-Himes DR, Han K, Guillard T, Jiang
D, Gaultier C, Guerin F, Aschard H, Leclercq R, Mekalanos JJ, Lory S, Pier GB. 2013. Enhanced
in vivo fitness of carbapenem-resistant oprD mutants of Pseudomonas aeruginosa revealed
through high-throughput sequencing. Proc Natl Acad Sci U S A 110:20747-52.
7. Kehl-Fie TE, Skaar EP. 2010. Nutritional immunity beyond iron: a role for manganese and
zinc. Curr Opin Chem Biol 14:218-24.

8. Salusso A, Raimunda D. 2017. Defining the Roles of the Cation Diffusion Facilitators in
Fe(2+)/Zn(2+) Homeostasis and Establishment of Their Participation in Virulence in
Pseudomonas aeruginosa. Front Cell Infect Microbiol 7:84.
9. Ducret V, Abdou M, Goncalves Milho C, Leoni S, Martin-Pelaud O, Sandoz A, Segovia Campos
I, Tercier-Waeber ML, Valentini M, Perron K. 2021. Global Analysis of the Zinc Homeostasis
Network in Pseudomonas aeruginosa and Its Gene Expression Dynamics. Front Microbiol
12:739988.
10. Ducret V, Gonzalez MR, Leoni S, Valentini M, Perron K. 2020. The CzcCBA Efflux System
Requires the CadA P-Type ATPase for Timely Expression Upon Zinc Excess in Pseudomonas
aeruginosa. Front Microbiol 11:911.
11. Wang D, Chen W, Huang S, He Y, Liu X, Hu Q, Wei T, Sang H, Gan J, Chen H. 2017. Structural
basis of Zn(II) induced metal detoxification and antibiotic resistance by histidine kinase CzcS
in Pseudomonas aeruginosa. PLoS Pathog 13:e1006533.
12. Dieppois G, Ducret V, Caille O, Perron K. 2012. The transcriptional regulator CzcR modulates
antibiotic resistance and quorum sensing in Pseudomonas aeruginosa. PLoS One 7:e38148.
13. Held K, Ramage E, Jacobs M, Gallagher L, Manoil C. 2012. Sequence-verified two-allele
transposon mutant library for Pseudomonas aeruginosa PAO1. J Bacteriol 194:6387-9.
14. Martinez-Garcia E, Fraile S, Algar E, Aparicio T, Velazquez E, Calles B, Tas H, Blazquez B,
Martin B, Prieto C, Sanchez-Sampedro L, Norholm MHH, Volke DC, Wirth NT, Dvorak P,
Alejaldre L, Grozinger L, Crowther M, Goni-Moreno A, Nikel PI, Nogales J, de Lorenzo V.
2023. SEVA 4.0: an update of the Standard European Vector Architecture database for
advanced analysis and programming of bacterial phenotypes. Nucleic Acids Res 51:D1558-
D1567.
15. Choi KH, Schweizer HP. 2006. mini-Tn7 insertion in bacteria with single attTn7 sites: example
Pseudomonas aeruginosa. Nat Protoc 1:153-61.
16. Huang W, Wilks A. 2017. A rapid seamless method for gene knockout in Pseudomonas
aeruginosa. BMC Microbiol 17:199.
17. Liu H, Naismith JH. 2008. An efficient one-step site-directed deletion, insertion, single and
multiple-site plasmid mutagenesis protocol. BMC Biotechnol 8:91.
18. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
analysis. Nature Methods 9:671-675.
19. Mihelj P, Abreu I, Moreyra T, Gonzalez-Guerrero M, Raimunda D. 2023. Functional
Characterization of the Co(2+) Transporter AitP in Sinorhizobium meliloti: A New Player in
Fe(2+) Homeostasis. Appl Environ Microbiol 89:e0190122.
20. Raimunda D, Subramanian P, Stemmler T, Arguello JM. 2012. A tetrahedral coordination of
Zinc during transmembrane transport by P-type Zn(2+)-ATPases. Biochim Biophys Acta
1818:1374-7.
21. Modrof J, Becker S, Muhlberger E. 2003. Ebola virus transcription activator VP30 is a zinc-
binding protein. J Virol 77:3334-8.
22. Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S.
2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. Nucleic
acids research 52:W521-W525.
23. Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023. UCSF
ChimeraX: Tools for structure building and analysis. Protein Sci 32:e4792.
24. Medema MH, Takano E, Breitling R. 2013. Detecting sequence homology at the gene cluster
level with MultiGeneBlast. Mol Biol Evol 30:1218-23.
25. Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. Nucleic Acids Res 52:W78-W82.

26.   Lewenza S, Gardy JL, Brinkman FS, Hancock RE. 2005. Genome-wide identification of
553   Pseudomonas aeruginosa exported proteins using a consensus computational strategy
554   combined with a laboratory-based PhoA fusion screen. Genome Res 15:321-9.
555   27.   El Rayes J, Rodriguez-Alonso R, Collet JF. 2021. Lipoproteins in Gram-negative bacteria: new
556   insights into their biogenesis, subcellular targeting and functional roles. Curr Opin Microbiol
557   61:25-34.
558   28.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
559   R, Zidek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-
560   Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M,
561   Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,
562   Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
563   AlphaFold. Nature 596:583-589.
564   29.   Caille O, Rossier C, Perron K. 2007. A copper-activated two-component system interacts
565   with zinc and imipenem resistance in Pseudomonas aeruginosa. J Bacteriol 189:4561-8.
566   30.   Conejo MC, Garcia I, Martinez-Martinez L, Picabea L, Pascual A. 2003. Zinc eluted from
567   siliconized latex urinary catheters decreases OprD expression, causing carbapenem
568   resistance in Pseudomonas aeruginosa. Antimicrob Agents Chemother 47:2313-5.
569   31.   Perron K, Caille O, Rossier C, Van Delden C, Dumas JL, Kohler T. 2004. CzcR-CzcS, a two-
570   component system involved in heavy metal and carbapenem resistance in Pseudomonas
571   aeruginosa. J Biol Chem 279:8761-8.
572   32.   Ducret V, Gonzalez D, Perron K. 2023. Zinc homeostasis in Pseudomonas. Biometals 36:729-
573   744.
574   33.   Maret W. 2013. Zinc biochemistry: from a single zinc enzyme to a key element of life. Adv
575   Nutr 4:82-91.
576   34.   Wang S, Cheng J, Niu Y, Li P, Zhang X, Lin J. 2021. Strategies for Zinc Uptake in Pseudomonas
577   aeruginosa at the Host-Pathogen Interface. Front Microbiol 12:741873.

Tables

 Table 1. List of Strains, plasmids and primers used in this study.

Strain/plasmid/primer	Relevant characteristics/sequence	Reference/Use
Pseudomonas aeruginosa		
MPAO1	Wild-type	Held *et al.* (2012)
yiiP::dTn5	Tn5 derivative ISphoA/hah insertion in *yiiP* (PA3963), Tc^r	Held *et al.* (2012)
pmcY::dTn5	Tn5 derivative ISphoA/hah insertion in *pmcY* (PA3962), Tc^r	Held *et al.* (2012)
C-pmcY	Tn5 derivative ISphoA/hah insertion in *pmcY*, pSEVA621-P$_{pmcY}$-*pmcY-gfp*, Tc^r Gm^r	This work
oprD::dTn5	Tn5 derivative ISphoA/hah insertion in *pmcY* (PA0958), Tc^r	Held *et al.* (2012)
DM3004	Tn5 derivative ISphoA/hah insertion in *pmcY* (PA0958), ΔPA3962,Tc^r	This work
Escherichia coli		
BL21(DE3) pLysS	*F- ompT hsdSB* (rB-mB-) *gal dcm* (DE3) *pLysS* (Cam^r)	Invitrogen, Carlsbad, CA,USA
DH5α	*supE44 ΔlacU169* (φ80 *lacZ* ΔM15*) hsdR17 recA1 endA1 gyrA96 thi1 relA1*	Invitrogen, Carlsbad, CA,USA
Plasmids		
pET23b(+)	T7 expression plasmid, Ap^r	Novagen, Darmstadt, Germany
pGEX-4-T2	T7*lac* expression plasmid*,* Ap^r	Cytiva, Marlborough, MA, USA
pSEVA621	RK2 cloning plasmid, Gm^r	Martínez-García *et al.* 2023
pET16bStrep	T7 expression plasmid, Ap^r	Luis Rubio´s laboratory
pEX18-Gm	*oriT$^+$ sacB$^+$*, gene replacement vector with MCS from pUC18, Gm^r	Huang & Wilks (2017)
Primers		

F- pET PA3962	CAGTCATATGCGCCTTGCCACCCGCG	Cloning PA3962 in pET23b(+) + NdeI at 5'-end
R- pET PA3962	ACTCGAGGGACTGAAAATACAGGTTTTCGCCGCCGCCGCGCTTCAGGCC	Cloning PA3962 in pET23b(+) + TEV site + XhoI at 3'- end
F-pGEX PA3962	GGATCCATGCGCCTTGCCACCCGCG	Cloning PA3962 in pGEX4T-2 + BamHI at 5'-end
R-pGEX PA3962	CCCGGGTCAGTGGTGGTGGTGGTGGTGCTCGAGGGACTGAAAATACAGGTTTTCGCCGGCGGCCGGCTTCAGGCC	Cloning in pGEX4T-2 + TEV site + $(His)_6$ tag + SmaI at 3'-end
F-pGEX-$_{1-29}$PA3962	CGTGGATCCGAGCGACGCGGCCAGATCGACGGTCG	Deletion N-terminal residues (1-29) of PmcY
R-pGEX-$_{1-29}$PA3962	TCGCTCGGATCCACGCGGAACCAGATCCGATTTTG	Deletion N-terminal residues (1-29) of PmcY
F-pGEX-$_{94-138}$PA3962	GCGGACGGCGGCGAAAACCTGTATTTTCAGTCCCT	Deletion C-terminal residues (94-138) of PmcY
R-pGEX-$_{94-138}$PA3962	TTCGCCGCCGTCCGCGCCGACGGCATCCTTGAGCA	Deletion C-terminal residues (94-138) of PmcY
F-pGEX$_{30-93}$PA3962His*6	CGGACGGCTGATCTAGACTAGGGGTCGACTCGAGCGGC	Deletion of $(His)_6$ tag
R-pGEX$_{30-93}$PA3962His*6	TCTAGATCAGCCGTCCGCGCCGACGGCATCCTTGAGCAC	Deletion of $(His)_6$ tag
F-D40xA– PA3962	TCGCATCGCCCCGGCCATCGTCGCCTTCGACGCCATCGGCC	D40A mutation in PmcY$_j$
R-D40xA– PA3962	GCCGGGGCGATGCGACCGTCGATCTGGCCGCGTCGCTC	D40A mutation in PmcY$_j$
F-D47xA– PA3962	CGCCTTCGCCGCCATCGGCCTGCTGTTCTATATCATCCCCGGCCTG	D47A mutation in PmcY$_j$

R-D47xA– PA3962	GATGGCGGCGAAGGCGACGATGGCCGGATCGATG CGACCG	D47A mutation in PmcY$_j$
F-D65xA– PA3962	CGGCGTCGCCTTCGCCACCGGCGCCATCTACCTGC CCGAC	D65A mutation in PmcY$_j$
R-D65xA– PA3962	GCGAAGGCGACGCCGAAGGCGATCAGGCCGGGGA TGATATAGAACAGC	D65A mutation in PmcY$_j$
F-pSEVA PA3962	CTAGTGAATTCCCCGGGTGGTCTTGCCGGC	Cloning *pmcY* plus 250pb up stream + EcoRI at 5'-end
R-pSEVA PA3962	ACTGGTACCGGCGGCCGGCTTCA	Cloning *pmcY* plus 250pb up stream + KpnI at 3'-end
F-GFP	AGGGTACCGCTAGCAAAGGAGAA	Cloning *gfp* + KpnI at 5'-end
R-GFP	CCTTCTAGATTAGCAGCCGGATCCT	Cloning *gfp* + XbaI at 3'-end
F-CzcS	ATCCATATGGCCCGCGAGCGGGA	Cloning *czcS* soluble domain + NdeI at 5'-end
R-CzcS	AGTGGATCCTCATTTTCGAACTGCGGGTGGCTCCAG CGCTCGTCGGCGT	Cloning *czcS* soluble domain plus StrepII + BamHI at 3'-end
F-RpsL	TCGGCACTGCGTAAGGTATG	qRT-PCR
R-RpsL	CTTACGGTCTTTGACACCCG	qRT-PCR
F-OprD	GCACAAACGATGAAGGCAAG	qRT-PCR
R-OprD	GTAGGCCAAGGTGAAAGTGT	qRT-PCR
F-up500bp	ACGTAAGCTTAGGCCGACACCGTCCAGCGC	*pmcY* deletion in *P. aeruginosa*
R-up500bp	GGTCGCGCTCGGAGGGCGGAGTATGACCTCCGGG TGAACATGACG	*pmcY* deletion in *P. aeruginosa*
F-down500bp	CGTCATGTTCACCCGGAGGTCATACTCCGCCCTCC GAGCGCGACC	*pmcY* deletion in *P. aeruginosa*

R-down500bp	ACGTGAATTCTCTGGAAGGCCAGGAGAAGG	*pmcY* deletion in *P. aeruginosa*

Figure 1. PmcY is a membrane protein that counteracts imipenem susceptibility. (A) Left, hydrophobic surface profile of the AF2 predicted structure for PmcY/PA3962. Position of Cys23 is indicated. Center and right, ribbon diagram of PmcY with lipobox amino acids indicated in green. The thiol group of Cys23 would promote acylation anchoring the protein, right. Notice that independent of the position at the inner or the outer membrane, PmcY faces the periplasmic space. P= periplasm; M= membrane; I/O= inside/outside the cell. (B) Analysis of heterologously expressed PmcY-(His)$_6$ in *E. coli* compartments (cytosolic= C; total membranes= TM; periplasm= P) by cell fractioning. Ten µg of protein from each fraction obtained from cells grown in LB media (basal= B; induced = I) were analyzed by SDS-PAGE (left) and Western blot (right). PmcY-(His)$_6$ expression is apparently increased in basal conditions and mostly found in the membrane fraction. (C) Serial dilutions (10 µl) of *P. aeruginosa* PAO1 (WT), *pmcY::dTn5* and *yiiP::dTn5* cells initially at OD$_{600nm}$=1.0 were spotted on MH-agar supplemented with the indicated imipenem concentration and incubated for 16 h at 37°C. (D) Complementation of *pmcY::dTn5* restores imipenem susceptibility. PA3962 plus 250bp upstream bases were cloned in the low copy plasmid pSEVA621 as a fusion to GFP. This construct was introduced by electroporation into *pmcY::dTn5*. No complementation was observed in *pmcY::dTn5* cells transformed with empty pSEVA621 nor with pSEVA621-GFP (not shown) (E) Fold change expression levels ($2^{-\Delta\Delta Ct}$ with *rpsL* as reference gene) of *oprD* transcripts *vs.* WT. Cells were grown until mid-log phase at 37°C in LB before RNA extraction. Data are the mean ± SE of three independent experiments. *,

P<0.05 from a two-tailed Student's t test of averaged ΔCt (*vs.* WT). (F) Zn^{2+} induced

lack of susceptibility response is specific to imipenem and is PmcY dependent.

Notice that *pmcY::dTn5* is unresponsive to Zn^{2+} and the lack of response to

ciprofloxacin in presence of the TM in both mutant strains. Agar dilution method was

used to determine MICs. (G) Double mutant *oprD-pmcY* (DM3004) shows similar

sensitivity to imipenem as the single mutant *oprD::dTn5*. Inhibition halos were

photographed after 16 h incubation at 37°C. Similar sensitivities between mutant

strains were obtained with increasing imipenem masses in the filter discs (not

shown).

Figure 2. Conserved residues in the juxtamembrane region/domain (PmcYs)

have a role in Zn^{2+} binding. (A) Left, conserved amino acids profile depicted in the

structure for PmcY/PA3962. Positions colored with maroon indicates high

conservation while cyan points to low conservation. Right, lateral chains of putative

amino acids with metal binding capacity are shown. (B) Up, representation of the

fusion protein GST-PmcYs. Down, metal (TM^{2+}) binding capacity and specificity of

the fusion product. Metal bound to GST was subtracted in each case and

represented 10% of the total metal present in GST-PmcYs. (C) Up, representation

of the fusion protein with conserved amino acids positions indicated with blue lines.

Down, binding capacity of mutants D40A, D47A and D65A. Notice that Zn^{2+} binding

decrease to a half vs. WT. (D) Representation of putative Zn^{2+} binding sites in PmcY.

Zn^{2+} atoms are indicated in red and putative bonds by O-ligands with dotted lines.

In all cases, data are the mean ± SE of three independent experiments. *P<0.01; [#],

P, 0.05 from a two-tailed Student's *t* test versus Zn^{2+} (B) or WT (C).

Figure 3. PmcY interacts with CzcS. On-column Zn^{2+}-dependent protein-protein

interaction assay. (A) Upper and lower panel, 100 mg (200 µl) of resin bound with

10 nmole of GST-PmcYj (PmcY$_{30-93}$) were washed with 10 bed volume of Buffer B

(Buffer A + 1 mM EDTA), (Upper panel, lane REDTA$_t$), or Buffer C (Buffer A + 0.1

mM Zn^{2+}) (Lower panel, lane RZn^{2+}$_t$) and incubated with 5 nmole CzcS$_{40-166}$ (lane

+I). The flow through was collected (FT) and the resin washed 4X with 2 bed volumes

of Buffer A (lanes W1-4). Finally, the resin was resuspended at the initial volume

(lane Rf) and 25 µl of each fraction analyzed by SDS-PAGE 12% and CBB staining.

Note that PmcY-CzcS interaction requires Zn^{2+} bound to PmcY (black asterisk). (B)

Protein interaction analyses of PmcY mutants. EDTA treated (upper panel) or Zn^{2+}-

loaded (lower panel) GST-PmcY mutants (5 nmole D40A, 1 nmole D47A, 2.5 nmole

D65A) were incubated with 5 nmole CzcS. Twenty five µl of fractions from resin

treated (lanes REDTA$_t$ and RZn^{2+}$_t$), incubated with CzcS (lanes +I) and final (lanes

Rf) samples were analyzed as in (A). Black asterisk denotes the Zn^{2+}-dependent

interaction and white, the lack of it.

Figure 4. PmcY-YiiP-like operons coevolved with OprD and likely CzcS in

Pseudomonadales. Bioinformatic analyses. (A) Two operon configurations

involving PmcY homologs were searched in complete bacterial genomes by

multigene Blast using protein sequences; *pmcY-yiiP*-like operon (blue) and non-

pmcY-yiiP-like operon (white). Parallel searches by RBH were done with protein

sequences coded by *czcS* (green) and *oprD* (yellow). (B) Venn diagram showing the

distribution of organisms with detected CzcS (green) and OprD (yellow) between the

two configurations of PmcY (*pmcY-yiiP*-like operon, in blue vs. *non-pmcY-yiiP*-like

operon, in yellow). Note the lack of association between *non-pmcY-yiiP*-like operon

organisms (226) with CzcS and OprD (1 and 13 organisms among 226,

respectively). (C) Phylogenetic analysis of the organisms with PmcY homolog

detected in the genome. Most species with *pmcY-yiiP*-like operon (blue strip)

belongs to the order of Pseudomonadales (ocher). Coevolution of OprD and CzcS

is depicted with yellow and green strips labels, respectively.

Figures

Figure 1

Figure 2

Figure 3

Figure 4