Temperature and genetic background drive mobilization of diverse transposable elements in a global human fungal pathogen

Anna I. Mackey[1], Vesper Fraunfelter[1], Samantha Shaltz[1], John McCormick[2], Callan Schroeder[1], John R. Perfect[3], Cedric Feschotte[2], Paul M. Magwene[4], Asiya Gusa[1]

[1]Department of Molecular Genetics and Microbiology, Duke University, Durham, North Carolina, United States of America

[2]Department of Molecular Biology and Genetics, Cornell University, Ithaca, New York, United States of America

[3]Department of Medicine, Duke University, Durham, North Carolina, United States of America

[4]Department of Biology, Duke University, Durham, North Carolina, United States of America

* **Corresponding author**

Email: asiya.gusa@duke.edu

ABBREVIATIONS

5mC – 5-methylcytosine

CMC – CACTA, Mirage, Chapaev

Indel – Insertion-deletion

KDZ – Kyakuja, Dileera, Zisupton

LTR – Long terminal repeat

Rap+FK506 – Rapamycin and FK506

RNAi – RNA interference

SNP – Single nucleotide polymorphism

TSD – Terminal site duplication

TE – Transposable element

TIR – Terminal inverted repeat

ABSTRACT

Transposable elements (TEs) are key agents of genome evolution across all domains of

life. These mobile genetic elements can cause mutations through transposition or by promoting

structural rearrangements. Stress conditions can amplify TE mobility, either by impairing TE

suppression mechanisms or through stress-induced interactions between transcription factors

and TE sequences, offering a route for rapid genetic change. As such, TEs represent an

important source of adaptability within populations. To investigate the interplay between

environmental stress and eukaryotic TE dynamics relevant to infectious disease, we examined

how heat stress and host-mimicking medium (RPMI) affect TE mobility in the global human

fungal pathogen *Cryptococcus neoformans*, using a collection of clinical and environmental

isolates. Using a selection-based screen, we captured the mobilization of seven distinct mobile

element families, encompassing diverse retrotransposons and DNA transposons, whose

insertions conferred antifungal resistance. This includes a novel element, CNEST, which

belongs to the CACTA, Mirage, Chapaev (CMC) supergroup. Heat stress at human body

temperature (37°C) significantly increased the mobilization of a subset of these TEs, leading to

higher rates of acquired antifungal resistance. Whole-genome assemblies revealed that,

compared to retrotransposons, DNA transposons were hypomethylated and approximately

uniformly distributed throughout the genome, features that may contribute to their frequent

mobilization. We further assessed TE-driven genomic changes within hosts using serial isolates

from patients with recurrent cryptococcal infections and from isolates passaged through mice.

While we observed evidence of TE copy number changes near chromosome ends, we found no

indication of TE-mediated alterations near gene-coding regions across any of the serial isolates.

Finally, TE mobility was isolate- and strain-dependent, with significant variation even among

clonally related strains collected from the same patient, emphasizing the role of genetic

background in shaping TE activity. Together, these findings reveal a complex and dynamic

relationship between environmental stress, genetic background, TE type-specific epigenetic

regulation, and TE mobility, with important implications for adaptation and acquired antifungal

resistance in *C. neoformans*.

AUTHOR SUMMARY

Transposable elements (TEs) are mobile segments of DNA that can "jump" from one place to another in a genome, causing genetic changes that can impact fitness and evolution. In this study, we investigate how different environmental conditions influence TE mobility in the pathogenic yeast *Cryptococcus neoformans*, which is globally distributed in the environment and causes life-threatening infections. We compared genetically diverse strains and found that growth at higher temperatures, specifically human body temperature (37°C), increases the mobility of some TEs. In contrast, other TEs were more active at lower temperatures (30°C), indicating distinct pathways of temperature regulation for different TE groups. By examining *C. neoformans* isolates from patients with recurrent infections, we observed genetic changes by one TE group that likely occurred during infection. We also found that TE groups are distributed differently across the genome and show varying levels of DNA methylation, a modification that can suppress their activity and control TE mobility in the cell. Overall, this work shows that environmental stresses, such as increased temperature, can accelerate genetic change by promoting TE mobilization. These changes may contribute to how *C. neoformans* adapts during infection or in the environment.

INTRODUCTION

Transposable elements (TEs) are powerful drivers of genomic change and provide a mechanism for adaptive evolution in fungal pathogens [1–9]. TEs are mobile genetic elements that fall into two broad categories: retrotransposons, which mobilize via an RNA intermediate through a "copy-and-paste" mechanism, and DNA transposons, which mobilize via a DNA intermediate, typically through a "cut-and-paste" mechanism [10]. TE activity can lead to genomic instability through unchecked transposition or by providing homologous sequences throughout the genome that promote rearrangements [11]. However, TEs can also promote beneficial genetic changes, especially under changing or stressful environmental conditions, by modulating the expression of nearby genes, disrupting endogenous gene functions, or providing beneficial cargo [11–15]. Notably, various stressors can increase TE expression and mobilization rates in fungi [1,5,15–18]. TE-mediated genetic changes have been linked to multiple adaptive phenotypes, including increased fitness under heavy metal stress in *Schizosaccharomyces pombe* and nitrogen-limited conditions in *Saccharomyces cerevisiae* [1–

90   5]. However, despite the demonstrated influence of TEs on adaptative genomic change,

91   spontaneous transposition events are rarely observed in *S. pombe* and *S. cerevisiae* under

92   many environmental conditions [19–22]. Therefore, the extent to which TEs contribute to the

93   rapid adaptation of yeast to environmental stressors remains an open question.

94   Diverse TEs are present in *Cryptococcus neoformans*, a pathogenic yeast which

primarily causes disease in immunocompromised individuals such as cancer patients, organ

transplant recipients, and people living with HIV/AIDS [23–25]. *C. neoformans* is responsible for

approximately 147,000 deaths annually and accounts for up to 20% of AIDS-related mortality

[26,27]. It is a ubiquitous environmental yeast, and inhalation of spores or desiccated yeast cells

can lead to pulmonary infection that may disseminate across the blood-brain barrier, causing

cryptococcal meningitis [23]. Approximately 10 – 20% of patients diagnosed with cryptococcal

meningitis experience recurrent disease, which can persist for years despite antifungal

treatment [28]. In-host microevolution can result in adaptive genetic changes that enhance

antifungal resistance, alter virulence traits, or increase overall pathogenicity, contributing to

persistent infections and treatment failure [29–32]. Although in-host microevolution influences

disease outcomes, the genetic basis of altered virulence traits and antifungal resistance often

remains unknown, potentially because short-read sequencing and variant calling, the primary

detection methods, are less suited for identifying structural rearrangements and TE

polymorphisms [33].

109   During infection, *C. neoformans* encounters a variety of stressors during the

environment-to-host transition. These include shifts in nutrient availability, pH, exposure to

reactive oxygen species, and elevated temperature (37°C) [34]. Previous studies in the less

pathogenic sister species *Cryptococcus deneoformans* have shown that heat stress significantly

increases TE mobilization rates, exceeding those of single nucleotide polymorphisms (SNPs)

and insertions/deletions (indels) by more than threefold per generation during in vitro passaging

at 37°C [35,36]. These findings suggest that in stressful environments, TEs can be the primary

drivers of genomic change. Indeed, TE mobilization in *C. deneoformans* has been observed

during murine host infection and, in both *C. neoformans* and *C. deneoformans*, has been linked

to the development of resistance to 5-fluorocytosine (5-FC), a first-line antifungal treatment [35–

37]. A prior study in *C. neoformans* assessed whether heat stress increases mutation rates in

the reference strain H99 and two additional clinical isolates, finding no association between heat

stress and elevated mutation rates [38]. However, given the small number of isolates examined,

it remains unclear whether heat stress can increase TE mobilization rates in *C. neoformans* and

how it may affect mutation rates across additional isolates and lineages. Furthermore, it is

unknown whether TEs contribute to genetic changes accumulated during persistent human

infection in *C. neoformans*, which causes approximately 95% of cryptococcal infections [39].

C. neoformans is known to harbor a variety of TEs, including DNA transposons as well

as long terminal repeat (LTR) and non-LTR retrotransposons [24,25]. Due to significant diversity

within TE sequences, structures, and mobilization mechanisms, TEs are further subdivided

using a hierarchical classification system [40]. TE orders (e.g., LTR) are subdivided into

superfamilies (e.g., Ty1/Copia) based on defining structural or sequence features. For instance,

DNA transposons that mobilize via a cut-and-paste mechanism typically contain a conserved

DDE or DDD catalytic domain within the transposase, which can be used for identification [41].

To date, only a subset of TEs in *C. neoformans* have been classified to the superfamily level,

including DNA transposons belonging to the Kyakuja/Dileera/Zisupton (KDZ) and HARBINGER

superfamilies, and LTR retrotransposons from the Ty1/Copia and Ty3/Gypsy superfamilies

[37,42,43]. Refining TE classification can provide insights into the mechanisms by which stress

influences TE mobilization and genome evolution.

Because uncontrolled TE activity can cause genome instability, hosts have evolved

mechanisms to suppress their mobilization [44–47]. The best-characterized TE suppression

system in *C. neoformans* is RNA interference (RNAi). Through this pathway, transcripts targeted

for silencing are processed into small interfering RNAs, which are loaded into the RNA-induced

silencing complex to target homologous transcripts for degradation [43,48–50]. RNAi has been

shown to silence double-stranded RNAs, transcripts with sub-optimal splicing signals, repetitive

transgenes, and TEs in *C. neoformans* [38,43,48,50]. Unlike in the yeast *S. pombe*, RNAi is

strictly involved in post-transcriptional silencing in *C. neoformans* and does not lead to

transcriptional silencing by facilitating the assembly or maintenance of heterochromatin [43].

Prior studies have shown that loss-of-function mutations in key RNAi genes can lead to a

hypermutator phenotype, driven by rampant TE mobilization, but only in strains with high TE

burden [37,38]. TE mobilization may also be shaped by epigenetic context. LTR and non-LTR

elements are enriched in centromeric and subtelomeric regions [38,42], where they co-localize

with epigenetic features of silent chromatin, such as 5-methylcytosine DNA methylation and

repressive histone modifications, including H3K27me3 and H3K9me2 [38,42,51–53]. These

marks are also enriched in centromeric and subtelomeric regions, and are rarely detected in

gene-rich chromosomes arms [38,42,51–53]. Despite this association, the regulatory effects of

DNA methylation and heterochromatin on TEs in *C. neoformans* remain poorly understood,

particularly regarding their influence on the mobilization of active TE families.

In this study, we investigated how infection-associated stressors, including elevated

temperature and exposure to a host-mimicking medium (RPMI), influence TE mobilization in *C.*

neoformans. Using a reporter gene-based approach, we examined TE mobility in genetically

diverse clinical and environmental isolates, including representatives from three of the four

major lineages (VNI, VNII, and VNBII) [54]. We found diverse, active TEs in *C. neoformans*,

including several whose identity or mobility were previously unknown. TE mobility was

temperature-dependent for all elements assessed, with heat stress significantly increasing

transposition rates for a subset of elements. De novo telomere-to-telomere genome assemblies

were used to examine TE-driven genetic changes in serially collected isolates from patients with

recurrent cryptococcal meningitis. Using these assemblies, we annotated TEs to assess the

genomic distribution and DNA methylation status of various transposable element types. DNA

transposons were approximately uniformly distributed throughout the genome and had

significantly reduced levels of DNA methylation compared to retrotransposons. Lastly, we found

that TE copy number and mobility were highly isolate- and strain-dependent, suggesting that

genetic background is a major determinant of TE-mediated genomic impacts. Together, these

results highlight the critical roles of environmental factors, particularly temperature, in driving

stress-induced genomic changes and adaptive evolution via transposition in the human fungal

pathogen *C. neoformans*.

RESULTS

Diverse transposable elements mobilize in a subset of *C. neoformans* clinical isolates

To investigate transposable element (TE) mobilization dynamics in *Cryptococcus*, we

obtained a set of 33 clinical isolates from 18 South African patients with recurrent cryptococcal

meningitis, collected as part of the GERMS-SA surveillance study (Table S1) [55]. Isolates were

collected by plating cerebrospinal fluid on non-selective medium and selecting a single colony at

the time of diagnosis (incident) and after subsequent disease relapse(s), which occurred 57 –

238 days later, depending on the patient (Fig 1A, Table S1). Using short read sequencing, a

prior study confirmed that isolates collected from the same patient differed by few single

nucleotide polymorphisms (SNPs) and insertions/deletions (indels), and that these isolates

formed monophyletic clades in phylogenetic analyses, suggesting that disease relapses were

clonal and not due to reinfection [29]. This set of clinical isolates is genetically diverse, including

Cryptococcus neoformans isolates belonging to three of the four major clades, VNI, VNII, and

VNBII, and *Cryptococcus gattii* isolates belonging to clades VGI and VGIV (Table S1) [54,56].

Prior work from our group showed that growth at 37°C increased TE mobilization rates in the

less pathogenic species *Cryptococcus deneoformans* [35,36]. To investigate TE mobility under

heat stress in *C. neoformans* and *C. gattii*, both recognized as priority fungal pathogens by the

World Health Organization [57], we conducted a preliminary screen using a "transposon trap"

assay that detects mutations in the ~934 bp *FRR1* (CNAG_03682) reporter gene, a system

previously used by our group and others [35,37,38]. Only mutations in *FRR1*, which encodes

the shared drug target of the antifungals rapamycin and FK506 (rap+FK506), can confer

resistance to both antifungals (Fig 1B) [58]. While mutations in other genes can confer

resistance to rapamycin or FK506 individually, rap+FK506 resistance during a short growth

period is most likely caused by a mutation in *FRR1*. This selection-based assay facilitates

assessment of TE activity by requiring mutation analysis at only one locus. For the preliminary

screen, incident and relapse isolates from the 18 South African cases, as well as the reference

strain H99, were cultured in non-selective, nutrient-rich YPAD medium at 37°C and plated on

medium containing rap+FK506 (Fig 1B). From a total of 37 isolates, ten independent rap+FK506

resistant mutants were collected per isolate, yielding 370 independent mutants. The *FRR1* gene

was PCR amplified for each independent mutant to determine the causative mutation of drug

resistance. Large DNA insertions, approximately 150 bp – 11 kb, detected by PCR amplification

and indicative of TE insertions were confirmed by sequencing. Isolates from three of the 18

cases screened contained TE insertions disrupting the *FRR1* locus at 37°C: cases 7, 15, and

77, which belong to the VNBII, VNI, and VNII clades, respectively (Table 1). Isolates from these

three cases were selected to assess the impact of temperature on drug resistance rates and TE

mobility. TE mobilization was detected exclusively in *C. neoformans*; of the three *C. gattii* cases

screened, only point mutations were identified in *FRR1*. Additionally, no TE insertions were

identified in H99, consistent with prior studies [38]. Subsequent screening of rap+FK506-

resistant mutants from cases 7 and 77 following growth at 30°C (see below) revealed additional

mobile element families.

Table 1. Cases used for in-depth profiling of the impact of temperature on TE mobility,

whole-genome assembly, and TE mapping.

Case	Collection Point	Clade	Days After Incident Isolate Collection	TEs identified inserting in *FRR1* in YPAD
7	Incident	VNBII	-	TCN3
7	Relapse	VNBII	137	TCN3, CNL1, KDZ2, KDZ3
15	Incident	VNI	-	CNIRT6
15	Relapse	VNI	166	CNIRT6
15	Relapse 2	VNI	223	CNIRT6
77	Incident	VNII	-	CNEST, KDZ1
77	Relapse	VNII	182	CNEST, KDZ1
77	Relapse 2	VNII	290	CNEST

In addition to the 370 independent rap+FK506 resistant mutants collected as part of the
preliminary screen, we collected additional transposon trap mutants from cases 7, 15, and 77,
along with an environmental isolate, for in-depth profiling of the effects of temperature (30°C
and 37°C) and RPMI medium on TE dynamics. In total, we examined over 1,000 independent
rap+FK506-resistant mutants and identified over 350 independent TE insertion events in *FRR1*.
Elements were assigned to known families based on sequence similarity to *C. neoformans* TE
family consensus sequences in the Repbase TE database, or to GenBank submissions when
family-level consensus sequences were unavailable. Family membership was assigned based
on the 80-80-80 rule: nucleotide similarity had to exceed 80% over at least 80 base pairs and
cover at least 80% of the recovered sequence length [40]. An exception to this was the non-LTR
retrotransposon CNL1, which shares only ~60% nucleotide similarity with the CNL1 consensus
sequence in Repbase. However, it exhibits high (> 99%) nucleotide similarity to an active mobile
element previously classified as CNL1 in *C. neoformans* by Priest et al. (2022), so we have
retained this name. In total, we identified seven diverse TE families mobilizing in *C. neoformans*
clinical isolates (Fig 1C, Table 1). This includes an element belonging to a family that, to our
knowledge, has not been identified in previous studies (CNEST), as well as elements not
previously reported to be mobile (CNIRT6, TCN3, KDZ3) (Fig 1C, Table S2). These elements
included DNA transposons, as well as long terminal repeat (LTR) and non-LTR
retrotransposons.

CNIRT6 transposition drives increased drug resistance rates at 37°C in case 15

To assess how heat stress impacts rap+FK506 drug resistance rates, fluctuation assays
were performed on case 15 incident, relapse, and relapse 2 isolates. For all fluctuation assays,
cultures were grown to saturation at 30°C or 37°C and the average cell population for each
isolate and temperature condition was determined to ensure that differences in growth rates did
not affect mutation rate estimates. All case 15 isolates displayed significant increases in
rap+FK506 resistance rates during growth at 37°C compared to 30°C in YPAD medium, as
indicated by non-overlapping 95% confidence intervals (21-, 12-, and 4-fold, respectively) (Fig
2A). The spectrum of mutations in *FRR1* was obtained by sequencing the *FRR1* locus of
independent mutants, followed by mapping to the wild-type *FRR1* sequence for each isolate
(Fig S1A). In cases where sequencing across the locus proved unsuccessful, junction PCR
using TE-specific primers was used to determine the mutation type. Strikingly, many of the
mutations in *FRR1*, up to 91% in the incident isolate at 37°C, were caused by insertions of a
large (~7.5 kb) DNA transposon with 99.7% nucleotide similarity to a putative element named
CNIRT6 (GenBank: JQ277267.1) (Fig 2B). CNIRT6 was originally identified in *C. neoformans*
strain JF289 through sequence analysis, however, this transposon has not been described in
the literature and is absent from TE databases. Mapping of a subset of these insertions
revealed CNIRT6 transposition into *FRR1* introns and exons, resulting in 3 bp terminal site
duplications (TSDs) with variable sequences (Fig S2A, B). CNIRT6 insertion rates were
significantly higher at 37°C compared to 30°C, increasing by 28-, 20-, and 16-fold in the
incident, relapse, and relapse 2 isolates, respectively (Fig 2C). Notably, drug-resistance rates at
37°C and CNIRT6 mobility at both temperatures decreased progressively in relapse isolates
collected 166 and 223 days after the incident isolate.

Given the significant differences in CNIRT6 mobilization rates among case 15 isolates, we
hypothesized that inter-isolate variability may be driven by differences in copy number, genomic
position, or epigenetic regulation. To investigate these factors, and to assess all inter-isolate
polymorphisms caused by mobile elements, we performed long-read sequencing to generate
contiguous, telomere-to-telomere genome assemblies for the wild-type case 15 incident,
relapse, and relapse 2 isolates, which were polished with Illumina reads (Table S3). 5-
methylcytosine (5mC) DNA methylation was identified using Nanopore signal data. Given that
all CNIRT6 insertions into *FRR1* appeared to be the same length by gel electrophoresis and
shared high nucleotide similarity to one another (> 99%), the full-length CNIRT6 sequence,
obtained from an insertion into *FRR1*, was used as the query in BLAST-like alignment tool
(BLAT) [59] homology searches against case 15 genome assemblies to identify full-length
copies (Fig S1A). For all BLAT homology searches predetermined cutoffs of nucleotide identity ≥

90% and coverage ≥ 95% were used to call full-length copies. This analysis revealed two full-
length CNIRT6 copies that are positionally conserved between incident and relapse isolates:
one is an exact match to the CNIRT6 sequence identified inserting into *FRR1*, and the second
differs by a single nucleotide substitution (Fig 2D). The nucleotide sequences of both CNIRT6
copies are identical between the incident, relapse, and relapse 2 isolates. No DNA methylation
was detected in any full-length CNIRT6 copies across all case 15 isolates, as indicated by
methylation frequencies < 0.75 at each CpG site, meaning fewer than 75% of long-read signal
data supported a methylated call (Table S4). Therefore, differences in genomic copy number,
chromosomal location, or DNA methylation do not appear to underlie differences in CNIRT6
mobilization rates between serial isolates.

To further characterize the active DNA transposon CNIRT6, we performed sequence
analysis and found that this element was ~7.5 kb in length and contained 24 bp terminal
inverted repeats beginning with 'CACTG' (Fig 1C, Table S2). Though no CNIRT6 movement has
been reported in the H99 reference strain, a homology search revealed two approximately full-
length copies in H99 (identity ≥ 90% and coverage ≥ 95%), containing a ~6.9 kb gene
annotation (CNAG_01367). Repeat masking of the predicted protein sequence to assess
similarity to known TEs using Repbase CENSOR revealed hits to transposases from TEs in the
CACTA (also known as EnSpm) superfamily, which contain a 'CACTA' motif at their distal ends,
resembling the 'CACTG' motif found in CNIRT6 [60]. To confirm that CNAG_01367 encodes a
transposase and refine this classification, a multiple sequence alignment was performed using
the amino acid sequences from CNAG_01367 and transposases from several cut-and-paste
DNA transposons from the CMC (CACTA, Mirage, Chapaev) supergroup [41]. This alignment
revealed a conserved DDE motif, which catalyzes the cut-and-paste mechanism of
transposition, indicating that CNIRT6 represents a full-length element predicted to encode a
transposase of the CMC supergroup (Fig S3A). To examine the relationship of CNIRT6 to other
CMC transposons, an unrooted tree was constructed using transposase amino acid sequences
representative of CMC families from animals, plants, and fungi. CNIRT6 formed a statistically
supported clade (Bootstrap value 79%) with CACTA elements previously identified in other
fungi, as well as plants and animals (Fig S3B). We conclude that CNIRT6 is a member of the
CACTA superfamily within the CMC supergroup.

To comprehensively assess all genomic changes between case 15 isolates that may have
been mediated by TEs other than CNIRT6, we performed synteny and read depth analyses, and
used Synteny and Rearrangement Identifier (SyRI) [61], to define structural rearrangements and

identify insertions, deletions, and duplications. We repeat masked insertions, deletions, and duplications, which revealed no evidence of transposition events between serial isolates, as indicated by the absence of sequence similarity to contiguous full-length TEs and transposition signatures such as TSDs. Relative to the genome reference strain H99, all three case 15 isolates contain a segmental duplication of ~614 kb of the left arm of chromosome 5 and a translocation between chromosomes 4 and 14 where the breakpoint lies in the predicted centromeres (Fig 2D). When comparing the case 15 isolates to one another, we identified an inversion of ~1.26 Mb on chromosome 3 present in both the relapse and relapse 2 isolates, and a translocation between chromosomes 1 and 9 present only in the relapse 2 isolate (Fig 2D). Breakpoints for both inter-isolate rearrangements were outside of the predicted centromeres. No TE annotations were observed near rearrangement breakpoints, suggesting that another source of sequence homology or homology-independent DNA repair mechanisms mediated these rearrangements.

Lastly, given the robust activity of CNIRT6 in case 15 isolates grown in YPAD, we hypothesized that CNIRT6 may have mobilized during recurrent infections associated with other cases. To test this, we performed Southern blots on a subset of cases to identify putative transposition events between incident and relapse isolates. We generated a 551 bp CNIRT6-specific probe and utilized a restriction enzyme that cuts once within CNIRT6, outside of the probe site, and produces a range of fragment lengths separable by gel electrophoresis (Fig S4A). Consistent with whole-genome assembly comparisons, there was no difference in the hybridization band pattern for CNIRT6 among case 15 isolates (Fig S4B). However, several other patient cases displayed differences in CNIRT6 hybridization band patterns among serial isolates (Fig S4B). Case 1 was of particular interest, as it exhibited changes in the molecular weight of three bands, suggesting that these copies may occupy different genomic locations and indicating putative CNIRT6 transposition events between the incident and relapse isolates.

To investigate putative CNIRT6 transposition events in case 1 isolates, we generated whole-genome assemblies of the incident and relapse isolates and performed homology searches using full-length CNIRT6 as the query sequence. No full-length copies of CNIRT6 were detected in either genome; an identical number of truncated copies were present at the same genomic locations in both case 1 isolates (Fig S4C). Additionally, repeat masking of insertions, deletions, and duplications revealed no evidence of TE mobilization events between the isolates. To assess whether mutations that altered recognition sites of the restriction enzyme BspEI (which cleaves at the palindromic sequence 'TCCGGA') might explain differences in Southern blot

banding patterns, we identified the closest BspEI sites upstream and downstream of CNIRT6

copies in the genomes. We found three instances where changes in BspEI sites led to the

predicted production of CNIRT6-containing fragments of different lengths, which correspond to

the observed band pattern differences, highlighting a limitation of this approach in detecting

transposition events. Lastly, in the case 1 relapse isolate a translocation was detected between

chromosomes 3 and 10, with the breakpoint located outside of the predicted centromere

regions. There were no TEs near the breakpoint, suggesting that this rearrangement was not

mediated by TE-derived sequence homology.

CNEST transposition drives increased drug resistance rates at 37°C in case 77

For case 77, the relapse, and relapse 2 isolates displayed significant increases in

rap+FK506 drug resistance rates during growth at 37°C vs 30°C (Fig 3A). Of mutations in *FRR1*

observed after growth at 37°C, 29 – 82% were caused by insertion of a novel DNA transposon,

which we named CNEST (*Cryptococcus neoformans* EnSpm-like Transposon; naming rationale

described below) (Fig 3B). CNEST insertions appeared identical in length by gel electrophoresis

and mapped to introns and exons in *FRR1* (Fig S1B, S5A). Insertions resulted in 2 bp 'TA' TSDs

(Fig S5B). CNEST insertion rates were significantly higher for all isolates at 37°C; 22-, 83-, and

36-fold in the incident, relapse, and relapse 2, respectively (Fig 3C). The full-length CNEST

sequence was ~7.2 kb in length, with 33 bp TIRs beginning with 'CACGG' (Fig 1C, Table S2).

Interestingly, we also identified insertions of the DNA transposon KDZ1, recently

characterized by Huang et al. 2024, which occurred at a lower frequency and only during growth

at 30°C (Fig 3B) [37]. KDZ1 is an ~11 kb element with 241 bp TIRs, and its insertion into *FRR1*

resulted in 8 bp TSDs (Fig S5B, Table S2). The KDZ1 putative transposase contains a catalytic

core affiliated with the Kyakuja/Dileera/Zisupton (KDZ) superfamily and a predicted Zn-chelating

CxC1 domain, both of which have been previously described in fungi (Fig 1C) [62].

Assemblies of the case 77 relapse and relapse 2 isolates revealed conserved copy number

and position of CNEST and KDZ1 by homology searches (identity ≥ 90% and coverage ≥ 95%)

with 4 full-length copies of CNEST and 5 full-length copies of KDZ1 (Fig 3D). No DNA

methylation was detected in any CNEST copies, as indicated by methylation frequencies < 0.75

at each CpG site from long-read signal data (Table S4). For KDZ1, a single copy contained one

methylated CpG in the relapse (Table S4). In the relapse 2, two KDZ1 copies contained one

methylated CpG. The additional methylated CpG in the relapse 2 isolate is ~900 bp upstream of

the predicted transposase start codon; however, a single methylated CpG seems unlikely to

impact transcription of the transposase.

To further characterize CNEST, which has not previously been identified as mobilizing in the

H99 reference strain, we performed a homology search, which revealed two approximately full-

length copies in H99 (≥ 90% identity and ≥ 95% coverage), containing a ~4.4 kb gene

annotation (CNAG_05910). Repeat masking of CNEST and the predicted amino acid sequence

of CNAG_05910 revealed no hits to TEs or transposases in the Repbase TE database.

However, given the identification of CNIRT6 as a member of the CMC supergroup, we predicted

that CNEST may also encode a CMC transposase. Multiple sequence alignment of the

CNAG_05910 amino acid sequence with transposases of the CMC supergroup revealed a well-

conserved DDE catalytic core, confirming CNAG_05910 as a CMC-type transposase encoded

within full-length CNEST (Fig S3A). Phylogenetic analysis places CNEST in a distinct clade from

CNIRT6, forming part of a strongly supported clade (Bootstrap value 100%) containing

unclassified transposases from diverse animals, including a transposase annotated as a CACTA

(also known as EnSpm) element in the Pacific oyster (*Crassostrea virginica*) (Fig S3B). This

clade is sister to clades of elements containing Chapaev and Mirage transposases from

animals, forming a well-supported monophyletic group (Bootstrap value 86%) (Fig S3B). Thus,

CNEST encodes a transposase defining a new superfamily within the CMC supergroup that is

distinct but most closely related to Chapaev and Mirage superfamilies.

Lastly, no major genomic rearrangements were identified between the case 77 relapse and

relapse 2 assemblies and no other TE-mediated genomic changes were detected by repeat

masking, as all insertions, deletions, and duplications identified showed no significant sequence

similarity to known TEs. Although mutation rate and spectra data are provided for the case 77

incident isolate, there was some variability in the growth of independent colonies and some

inconsistency in mutation rate observed between independent fluctuation assays. The likelihood

of genome instability in this isolate led to its exclusion from nanopore sequencing and genome

assembly. In summary, two elements mobilized to mediate drug resistance in case 77: KDZ1,

which was detected only at 30°C, and CNEST, which mobilized significantly more frequently at

37°C and is now characterized as a novel active TE in *C. neoformans* belonging to the CMC

supergroup. Note that the KDZ superfamily, to which KDZ1 belongs, is completely distinct from

the CMC supergroup, which includes CNEST and CNIRT6.

Diverse TEs drive temperature and strain-specific TE mobilization dynamics in case 7

The case 7 incident isolate displayed a significantly higher rap+FK506 resistance rate

following growth at 37°C compared to 30°C, while the relapse isolate showed no significant

difference in drug resistance rates (Fig 4A). For the incident and relapse isolates at 37°C, ~30%

of mutations in *FRR1* were caused by insertions of the LTR retrotransposon TCN3, with both

full-length and solo LTR insertions detected (Fig 4B, Fig S1C). TCN3 is ~6.3 kb in length with

606 bp LTRs, and insertion into *FRR1* occurred in introns and exons, resulting in 5 bp TSDs of

variable sequence (Fig 1C, Fig S6A, B, Table S2). TCN3 is a member of the Ty3/gypsy

superfamily that has been previously characterized [63]. The TCN3 insertion we identified

contains a ~4.7 kb open reading frame with apparently intact Pol domains (reverse

transcriptase, RNase H, protease, and integrase) (Fig 1C). Although we did not identify a Gag

domain, the first 306 amino acids lack recognizable functional domains and correspond to the

region that typically encodes Gag in Ty3/gypsy elements [63]. Moreover, the Gag gene is the

most variable component of LTR retrotransposons, making it particularly difficult to detect [64].

These observations suggest that TCN3 likely represents an autonomous LTR retrotransposon

(Fig 1C, Table S2). We recovered no TCN3 insertions at 30°C out of a total of 68 independent

mutants screened, suggesting that the element may exclusively mobilize at higher temperatures

(Fig 4B).

Surprisingly, over 60% of *FRR1* mutations in the case 7 relapse isolate at 30°C were caused

by insertions of diverse TEs, including the non-LTR retrotransposon CNL1 and the DNA

transposons KDZ2 and KDZ3 (Fig 4B, Fig S1C). In contrast, mutations in the incident isolate at

30°C were exclusively SNPs/indels (Fig 4B). The longest CNL1 insertion we identified in *FRR1*

was ~3.5 kb in length and contained a ~2.5 kb open reading frame with an apparently intact

reverse transcriptase domain (Fig 1C, Fig S1C). This maximum insertion length matches

previous reports of CNL1 transposition [38]. CNL1 insertions identified in *FRR1* ranged from 180

bp to ~3.5 kb likely due to premature termination of reverse transcription which can occur during

target primed reverse transcription (TPRT) [65]. CNL1 insertions occurred in the *FRR1* 5′

untranslated region (UTR) and in an exon, and were associated with TSDs of variable length (0

– 13 bp), consistent with findings from prior studies (Fig S6A, B) [38]. For KDZ2 (~9.2 kb with

139 bp TIRs) and KDZ3 (~10 kb with 169 bp TIRs), insertion into *FRR1* occurred in introns and

exons and resulted in 4 – 8 bp TSDs (Fig 1C, Fig S6A, B, Table S2). The internal sequences of

KDZ2 and KDZ3 encode putative transposases with KDZ and CxC1 domains (Fig 1C). Although

TCN3 and KDZ3 were previously annotated, this study is the first to demonstrate their mobility.

Homology searches of case 7 genome assemblies using full-length TE sequences that
inserted into *FRR1* as queries revealed conserved copy number and position of full-length
TCN3 (2 copies), KDZ2 (2 copies), and KDZ3 (3 copies) between the incident and relapse
isolates (Fig 4C). This indicated that the observed KDZ2 and KDZ3 activity in the relapse isolate
was not due to an increase in copy number or change in genomic position. Additionally, no DNA
methylation was detected on any full-length KDZ2 or KDZ3 copies in case 7 isolates, and no TE
sequence polymorphisms were observed between serial isolates (Table S4). We did, however,
identify an expansion of CNL1 copies in the relapse isolate, with the number increasing from 16
to 39 full-length copies in arrays in the subtelomeres (Fig 4C). CNL1 copies exhibited varying
levels of DNA methylation (Fig 4C, Table S4). The expansion in CNL1 copy number may explain
the observed CNL1 mobility in the relapse isolate, which was undetected in the incident isolate
(Fig 4B). No genomic rearrangements or other TE-mediated changes, including alterations in
copy number or position of TCN3 solo LTRs, were identified between the case 7 isolates.

Interestingly, we recovered 13 total CNL1 insertions in *FRR1*; however, all full-length CNL1
copies in case 7 genome assemblies are located at the ends of incident and relapse
chromosome arms, flanked externally by telomeric repeats ("TAACCCCC," or slight variations
thereof), consistent with previous reports (Fig 4C) [24,66]. To further characterize CNL1, we
performed a phylogenetic analysis of the reverse transcriptase domain from a full-length CNL1
insertion in *FRR1*, and the CNL1 consensus sequence from the Repbase TE database. This
analysis revealed that CNL1 is more closely related to non-LTR elements of the R2 superfamily
(Fig S3C), which are characterized by a single open reading frame and often insert into specific
target sequences [67,68], in contrast to other non-LTR retrotransposons such as LINE1
elements which have weak target site specificity [69,70]. CNL1 was found to be most closely
related to MoTeR, a sequence-specific telomeric-targeting element described in the fungal plant
pathogen *Magnaporthe oryzae* (Bootstrap value 100%) [71]. Additionally, we observed similarity
in the TSDs between independent CNL1 insertions in *FRR1*, suggesting potential sequence-
specific target site preferences for this element (Fig S6A, B).

Centromeres in *C. neoformans* are typically enriched for LTR retrotransposons with > 95%
of LTR retrotransposons, including TCN3, restricted to centromeric regions in the H99 genome
reference strain, though the mechanism or selective pressure driving centromere targeting of
insertions remains unclear [42]. Although endogenous TCN3 copies in case 7 isolates are
restricted to centromeres, 21 independent TCN3 insertions were detected in *FRR1* (Fig 4B).

Mapped insertions occurred at variable sequences and positions within *FRR1*, indicating a lack

of strong sequence-specific insertion preference (Fig S6A, B).

C. neoformans centromeres are also enriched for DNA methylation, which is thought to

contribute to repression of TEs present in these regions [51,52]. Therefore, we assessed

whether the centromeric copies of TCN3 were methylated. If methylated, demethylation and

subsequent derepression of TCN3 under heat stress in vitro might explain the observed TCN3

mobility at 37°C in case 7 isolates [72]. We found that the full-length copies of TCN3 in the case

7 isolate genomes reside in hypomethylated regions within the predicted centromere region of

Chr 3 for the incident and relapse (data for the incident isolate shown) (Fig 4D). While five to

seven 5mC DNA methylation marks were observed in full-length TCN3 copies, they are located

at the end of the 3' LTR (Fig 4D, Table S4). This region is not typically transcribed, making the

observed DNA methylation unlikely to impact TCN3 transposition [73]. This suggests that loss of

DNA methylation at full-length TCN3 copies is unlikely to explain the differential activity of TCN3

observed in vitro, where transposition occurred only at 37°C.

Our group previously identified the LTR-retrotransposon TCN12 in *Cryptococcus*

deneoformans as heat stress-activated, mobilizing almost exclusively at 37°C [35,36]. In *C.*

neoformans, heat shock binding elements (HSEs) in the promoter region of heat-responsive

genes have been shown to be bound by the transcriptional activator heat shock factor 1 (Hsf1)

during heat stress [74,75]. A putative HSE was previously identified in the TCN12 LTR [36], and

upon scanning the TCN3 LTR, we identified two putative HSEs (Fig 4E). This suggests that the

elevated mobilization rates of TCN3 and TCN12 at 37°C may result from Hsf1 binding and

subsequent upregulation.

TE mobility is altered in RPMI medium

To more closely mimic the environment *C. neoformans* experiences in-host and assess

impacts on transposition, we compiled the spectra of mutations in the *FRR1* reporter gene for

cases 7, 15, and 77 following growth in chemically defined RPMI tissue culture medium at 37°C

[76]. We additionally tested the impact of RPMI medium at 30°C for case 7 isolates given the

activity of CNL1, KDZ2, and KDZ3 observed in the relapse isolate at 30°C in YPAD medium (Fig

4B). RPMI medium was supplemented with less glucose (0.2%) compared to YPAD (2%) and

had a defined starting pH of 7.0. For four of the six isolates, the same TEs were identified as

mobile in both YPAD and RPMI medium (Fig S7A, B, C). In case 77 and case 7 incident

isolates, CNL1 and KDZ3, respectively, were observed inserting into *FRR1*, events not

previously detected in YPAD medium (Fig. S7B, S7C).

We next assessed differences in the frequency of TE insertion events between RPMI

and YPAD medium. In the case 15 relapse isolate, there was a significant difference in TE

insertions into *FRR1*, with fewer insertions occurring in RPMI compared to YPAD at 37°C (two-

tailed Fisher's Exact test, *p* = 5.62e-6) (Fig S7A, Fig 2B). Interestingly, the use of RPMI medium

altered TE dynamics in an isolate- and temperature-specific manner in case 7 isolates (Fig S7C,

Fig 4B). Differences between RPMI and YPAD medium include: (1) a significant decrease in

TCN3 insertions in the relapse at 37°C in RPMI, with no insertions detected (two-tailed Fisher's

Exact test, *p* = 0.00905), (2) a significant increase in KDZ3 insertions in the incident at 30°C in

RPMI, with no insertions occurring in YPAD (two-tailed Fisher's Exact test, *p* = 2.56e-4), and (3)

a significant increase in CNL1 insertions in the relapse at 30°C in RPMI (two-tailed Fisher's

Exact test, *p* = 0.0261), highlighting the complex relationship between environment, genetic

background, and TE mobility in case 7 isolates. Across all cases assessed, growth in RPMI

medium altered either the TE families observed mobilizing or the frequency of insertion events

for at least one isolate.

CNEST heat stress-activation is conserved in a geographically and temporally distinct

environmental isolate

Homology searches revealed full-length CNEST and KDZ1 copies in a previously

published *C. neoformans* VNII genome, the same lineage as the case 77 isolates, sequenced

from an environmental sample collected from cockatoo droppings in Boston in 2000 (designated

cockatoo guano isolate) [31,77]. While the cockatoo guano and case 77 isolates both belong to

the VNII clade, they are geographically and temporally distinct, as the case 77 isolates were

collected eight years later in South Africa. However, a pairwise whole-genome comparison

revealed that the isolates are highly genetically similar, differing by 1,610 SNPs and indels

across the genome, corresponding to approximately 0.0085% divergence at the nucleotide

level. To better understand mobility of CNEST and KDZ1 in a different strain background, we

assessed rap+FK506 drug resistance rates and the spectra of mutations in *FRR1*. Indeed,

cockatoo guano isolate drug resistance rates were significantly elevated during growth at 37°C

compared to 30°C (Fig 5A). Approximately 20% of mutations in *FRR1* at 37°C were CNEST

insertions, while no CNEST insertions were observed at 30°C, strongly supporting that CNEST

transposition is induced by heat stress in the cockatoo guano isolate (Fig 5B). To assess the

genomic context of CNEST and KDZ1, we performed synteny analysis, which demonstrated

that both the copy number and genomic position of CNEST and KDZ1 were conserved between

the cockatoo guano and case 77 isolates (Fig 5C). This may suggest that mobilization of

CNEST and KDZ1 in the VNII lineage is infrequent in situ, that mobilization events are typically

detrimental to fitness, or that there is selective pressure to maintain both the copy number and

genomic context of these elements, all possibilities that will require further investigation.

Copy number analysis of other mobile TEs identified in this study revealed two full-length

copies of KDZ3 in the case 77 genomes and one full-length copy in the cockatoo guano isolate,

located at a conserved site shared with one of the case 77 copies (Fig S8A). The second KDZ3

copy in the case 77 relapse isolate, located on Chr 12, appears to result from a transposition

event, as synteny flanking this copy is maintained between the two isolates and 8 bp TSDs were

identified (Fig S8A, B). Repeat masking of insertions and deletions also revealed evidence of an

additional TE family, Gypsy-1_CrGa, originally identified in *C. gattii* [78], whose copy number

differs between the case 77 relapse and cockatoo guano isolates (Fig S8A). Two Gypsy-1_CrGa

transposition events were detected in the cockatoo guano isolate and one in the case 77

relapse genome (Fig S8A). All insertions of this element occurred within predicted centromeres

and generated 6 bp TSDs of variable sequence (Fig S8B). Other insertions and deletions, which

commonly occurred in the centromeres, contained TE sequences but were likely not mediated

by transposition, as they involved multiple or fragmented TE copies and lacked transposition

signatures such as TSDs. Thus, while KDZ1 and CNEST copy number and position are

conserved between these isolates, transposition events involving other TE families have

occurred.

A recent study passaged the cockatoo guano isolate twice through A/J mice via retro-

orbital intravenous injection, collecting a total of ten isolates from the brains and lungs after the

second passage [31]. The cockatoo guano isolate was also the point-source isolate for a case

of cryptococcal meningitis, from which a single isolate was collected from the patient (PU

isolate) [77]. To assess whether transposition of the two highest copy number TEs located

outside of the predicted centromeres, CNEST and KDZ1, occurred during murine passage or

human infection, we performed Southern analysis using 551 bp CNEST- and 567 bp KDZ1-

specific probes on all ten isolates that had undergone two rounds of murine passage, as well as

the cockatoo guano and PU isolates. Restriction enzymes that cut once within the TE sequence,

outside of the probe site, were used to produce a range of fragment lengths (Fig S9A). The

same hybridization band patterns were observed in the original cockatoo guano isolate and in

all isolates recovered following infection (Fig S9B, C), indicating that no CNEST or KDZ1

mobilization occurred during infection, despite detection of CNEST transposition in vitro at 37°C

(Fig 5B). Thus, while heat-stress activation of CNEST appears conserved between

environmental and clinical isolates, no transposition events were detected in isolates recovered

from human or murine infections.

567   Full-length copies of TEs are present across multiple isolates and lineages

Because TE mobility was highly isolate- and strain-dependent in the case 15, 77, and 7

isolates, where the TEs captured mobilizing were unique to each case, we hypothesized that

this may be due to the absence of full-length copies in certain isolate backgrounds. To evaluate

the prevalence of full-length TE copies across *C. neoformans* isolates, we conducted homology

searches in assembled genomes using the active TE sequences that were identified inserting

into the *FRR1* gene as queries. Full-length copies were defined as hits with ≥ 90% nucleotide

identity and ≥ 95% coverage of the query sequence. We identified full-length copies of CNIRT6,

CNEST, TCN3, KDZ1, and KDZ3 in several genomes of isolates where transposition was not

detected using the transposon trap assay (Fig S10A). For example, at least one full-length copy

of CNEST was found in all genomes assessed, including strains from the VNI, VNII, and VNBII

lineages (Fig S10A). Additionally, while Gypsy-1_CaGr was not observed mobilizing in vitro, at

least one full-length copy was present in all genomes, with up to 12 copies detected in the case

7 isolates (Fig S10A). While it appears that there may be a link between increased genomic

copy number and TE mobility, we do not have sufficient data to formally assess this.

To assess whether full-length copies may be present in the 15 *C. neoformans* cases

initially screened without available whole-genome assemblies, we performed TE copy number

estimation by aligning short reads to representative full-length TE sequences obtained from an

isolate in which the TE family was identified as active. Copy numbers were estimated based on

normalized read depth, defined as the depth at which ≥ 95% of the TE sequence is covered

[37]. This analysis revealed that full-length copies of potentially active elements may be present

even in isolates in which no TE mobilization was detected in the preliminary screen, including

cases where estimated copy numbers exceeded those in isolates with observed mobilization

(Fig S10B).

591   DNA methylation status may explain differential mobility of TCN3

Although most TEs identified in this study were detected mobilizing exclusively in a

single isolate background, additional full-length copies were found in other assemblies or

inferred through copy number estimation (Fig S10A, B). This suggests that TE mobility may be

differentially regulated across isolate backgrounds. For example, elevated TE mobility in some

isolates may be explained by a loss of RNAi functionality or the absence of DNA methylation on

TE sequences.

To explore potential regulatory mechanisms, we first assessed the levels of DNA

methylation in full-length TE copies in which mobilization was either observed or not observed

using the transposon trap assay. Full-length TCN3 copies were highly methylated in cases 15

and 1, whereas hypomethylated copies were observed in case 7 isolates (Table S4). In contrast,

DNA methylation was generally absent from full-length DNA transposons in strains where no

mobilization was observed. For example, in whole-genome assemblies, full-length KDZ1, KDZ3,

and CNEST elements were unmethylated in isolates where transposition was not observed in

FRR1, with the exception of CNEST in case 1 (Table S4). These findings suggest that while

DNA methylation may play a role in regulating TCN3 mobility at the centromeres, the

methylation status of full-length DNA transposons (which are primarily unmethylated) does not

explain isolate- or strain-specific differences in mobility.

Next, we examined whether genes required for RNAi function were intact in cases where

TE mobility was observed. In cases 7, 15, 77, and the cockatoo guano isolate, we first screened

for predicted loss-of-function mutations defined as predicted protein sequences < 85% of the

expected length using a previously published list of genes essential for RNAi [37]. No genes

required for RNAi function exhibited truncated protein sequences. Second, we performed

variant analysis and identified zero nonsense or frameshift mutations across the entire coding

sequence for RNAi genes in all isolates, providing no strong evidence for loss-of-function

mutations.

DNA transposons are predominantly unmethylated and commonly found in gene-rich

regions

We observed that all full-length, active DNA transposons mapped in the case 15, 77, and

7 genomes were located in gene-rich regions along the chromosome arms (Fig 2D, 3D, 4C).

This pattern contrasts with the distribution of retrotransposons in *Cryptococcus*, as previous

studies have shown that LTR elements predominantly localized in centromeres [42], and non-

LTR elements are enriched at chromosome ends [38]. To further investigate the genomic

distribution of DNA transposons in *C. neoformans*, which has not previously been described, we

took a comprehensive, unbiased approach to assess genome-wide TE content and distribution.

De novo TE annotation was performed using the Earl Grey package [79] for cases 1, 7, 15, and

77, as well as the genome of the reference strain H99, in which TE mobilization has not been

observed unless RNAi is disrupted [38,49]. Results from the first assemblies of serially collected

isolates are shown, as assemblies of later relapse isolates from the same patients yielded

similar results. TEs occupied approximately 3 – 5% of the genomes analyzed (Fig 6A). While

the proportion of LTR retrotransposons remained relatively consistent (2.2 – 2.6%), greater

(more than tenfold) variation was observed in the proportion of DNA transposons (0.08 –

0.85%), and non-LTR retrotransposons (0.06 – 2.06%) (Fig 6A). Notably, the *C. neoformans*

reference strain H99 had the lowest proportion of DNA transposons among all isolates

analyzed.

To visualize TE distributions, chromosomes were split at the midpoint of the predicted

centromere. For each chromosome arm, the length, TE positions, and protein-coding gene

annotation positions were scaled from 0 to 1, allowing data from all chromosome arms to be

plotted on a single track. In agreement with previously published data, LTR and non-LTR

elements cluster in the predicted centromeres and subtelomeres, respectively; these are

regions with a lower density of protein-coding genes (Fig 6B). By contrast, DNA transposons

appear to be more evenly distributed across chromosome arms (Fig 6B). We used quantile-

quantile (QQ) plots to assess whether the positions of DNA transposons along chromosome

arms follow a uniform distribution. For each genome, we generated theoretical quantiles as

sorted random numbers between 0 and 1; the total number of theoretical quantiles matched the

total number of DNA transposon annotations. The plotted points fell approximately along the

diagonal of the QQ plots, indicating that DNA transposons were distributed roughly uniformly

across the genome (Fig S11).

To assess how frequently DNA transposons were methylated, given their presence in

gene-rich regions that are not typically methylated in *C. neoformans* [52,53], we compared their

5mC DNA methylation levels to those of LTR and non-LTR elements. First, we assessed

average methylation density per 100 bp. DNA transposons exhibited significantly lower average

5mC densities compared to both non-LTR and LTR retrotransposons (pairwise Wilcoxon rank-

sum test with continuity correction; adjusted *p* values < 2e-16 for both comparisons) (Fig 6C).

Additionally, LTR elements exhibited significantly higher DNA methylation densities compared to

non-LTR elements (pairwise Wilcoxon rank-sum test with continuity correction; adjusted *p* value

< 2e-16). Second, we assessed the proportion of TE annotations per genome containing at least

one high-confidence 5mC call (methylated frequency from nanopore long-read signal data >

0.75). According to this analysis, we found that LTR elements are by far the most consistently

methylated (55.6 – 71.2% of LTR retrotransposons contained at least one predicted 5mC),

followed by non-LTR elements (7.7 – 53.4%), and DNA transposons (5.4 – 15.5%) (Fig S12A,

B). The number of methylated LTR and non-LTR retrotransposons was significantly enriched

compared to DNA transposons (Fisher's method combined *p* values: 3.2e-64 for LTR elements

and 9.2e-19 for non-LTR elements).

To estimate background levels of DNA methylation in protein-coding genes, we lifted

over gene annotations from the H99 reference genome. The percentage of protein-coding

genes with at least one high-confidence 5mC call ranged from 0.5 – 4.5%, indicating that DNA

TEs remain significantly enriched for DNA methylation relative to endogenous protein-coding

genes (Fisher's method combined *p* value 3.9e-10), albeit at a lower frequency than LTR and

non-LTR elements. Thus, the majority of DNA transposons across all genomes analyzed are

unmethylated, and those that are methylated exhibit a lower methylation density compared to

non-LTR and LTR retrotransposons (Fig 6C, Fig S12A, B). This suggests that DNA methylation

is unlikely to play a major role in modulating the activity of DNA transposons in *C. neoformans*.

DISCUSSION

Understanding the interplay between mobile genetic elements, environmental stressors,

epigenetics, and genetic background is critical for uncovering factors that influence evolutionary

rates and the capacity for adaptive change, particularly in the context of fungal pathogens. In

this study, we screened a collection of clinical *Cryptococcus* isolates for transposable element

(TE) insertions in the *FRR1* reporter gene and profiled rates of rapamycin and FK506

(rap+FK506) resistance, as well as mutation spectra, at both 30°C and the heat-stress condition

of 37°C. We also generated telomere-to-telomere genome assemblies to assess TE copy

number, genomic position, and DNA methylation status in isolates from patients with recurrent

cryptococcal infections.

C. neoformans **hosts diverse active transposable elements**

We identified seven TE families mobilizing to confer rap+FK506 resistance, including

both retrotransposons (LTR and non-LTR) and DNA transposons (Fig 1). Several of these

elements (TCN3, CNIRT6, KDZ3) were not previously recognized as active. In addition, we

discovered a novel, active element, which we named CNEST. The diversity of active TEs in *C.*

neoformans stands in striking contrast to other yeasts commonly used to study TEs, such as

Saccharomyces cerevisiae and *Schizosaccharomyces pombe*, which harbor only LTR
retrotransposons [80,81]. In these species, TE mobilization studies typically rely on TE
constructs driven by inducible heterologous promoters, due in part to the low de novo
transposition rates of endogenous copies [82–85]. By contrast, in *C. neoformans*, selecting for
mutations at a single genomic locus frequently revealed insertions of endogenous TE copies
with diverse transposition mechanisms (Figs 2 – 5). The ability to readily capture multiple active
TE families mobilizing within a single *C. neoformans* strain underscores the value of this
genetically tractable model for studying TE mobilization strategies and regulation in a
basidiomycete.

Phylogenetic analysis of transposases from CNIRT6 and CNEST revealed that they
belong to the CMC supergroup first described by Yuan and Wessler (2011) (Fig S3). However,
CNIRT6 and CNEST are only distantly related and fall within two monophyletic clades among
CMC elements: one containing CNIRT6 and transposases from plants, animals, and fungi, and
a second containing CNEST and transposases from animals and fungi. Thus, *C. neoformans*
hosts active members of two different CMC superfamilies, a notable finding given that this group
of DNA transposons remains largely understudied in fungi and animals.

The use of a single reporter gene to capture TE insertions may have limited our potential
for active mobile element discovery, since some TEs have preferences for specific insertion
sequences or genetic compartments [11]. However, the *FRR1* reporter system recovered
insertions of TCN3 and CNL1, which appear to target centromeric and subtelomeric regions,
respectively. Mapping TCN3 insertions in *FRR1* suggested weak sequence specificity (Fig S6).
Alternatively, TCN3 insertions may be guided by non-sequence-based features, such as
centromere-specific histone variants [86] or heterochromatin-associated proteins [87], or its
insertion pattern may be shaped by selective pressures. It remains possible that TCN3 insertion
into *FRR1* is relatively rare and that most de novo insertions occur within centromeric regions.

CNL1 copies are enriched in subtelomeric regions in case 7 isolates and other *C.*
neoformans genomes [38]; however, we also recovered insertions of CNL1 within *FRR1* (Fig 4).
Phylogenetic analysis revealed that CNL1 is most closely related to R2 elements, and
specifically to MoTeR, a telomere-targeting element characterized in the fungal plant pathogen
Magnaporthe oryzae [71]. Although prior studies have reported that CNL1 preferentially inserts
into copies of itself within arrays near telomeric repeats, its precise sequence specificity remains
undefined. Limited mapping of CNL1 insertions into *FRR1* reveled that TSDs contained the
sequence 'CACCCCT,' which resembles the *C. neoformans* telomere seed sequence

('AACCCCCT' or similar variants), although interpretation was complicated by premature

termination of target primed reverse transcription (TPRT) at different positions within the

element (Fig S6). Further work is needed to clarify CNL1's target site specificity, including the

extent to which it targets telomeric repeats like MoTeR, inserts into its own sequence, or both.

Along with TCN3, these findings highlight open questions about factors driving target site

selection and TE distribution across genomic compartments. Intriguingly, stress may be one

such factor as nutrient deprivation alters Ty5 phosphorylation, shifting its insertion profile from

heterochromatic regions to more broadly distributed sites across the genome [87].

Lastly, most observed TE mobilization events were driven by apparently full-length TE

copies containing detectable functional domains required for mobilization. Phylogenetic analysis

revealed full-length, seemingly intact transposase sequences within mobilized DNA

transposons, and functional domains required for mobilization were identified in mobilized TCN3

and CNL1. We found no evidence for the mobilization of clearly non-autonomous TE copies

(significantly shortened or lacking apparent mobilization machinery). However, we did detect

truncated CNL1 insertions, likely resulting from premature termination of TPRT, as well as

instances of solo TCN3 LTRs, which likely arose from non-allelic recombination between LTR

sequences following TCN3 insertion. This contrasts with other organisms, such as humans,

plants, and *Drosophila*, where shortened non-autonomous TEs frequently transpose [88–90].

Collectively, we identified seven TE families in *C. neoformans*: DNA transposons from

the KDZ superfamily and CMC supergroup, alongside a Ty3/Gypsy LTR element and an R2

non-LTR element actively transposing. These TEs employ distinct transposition mechanisms

and show preferences for different genomic compartments, revealing a rich "ecosystem" of

active TEs within *C. neoformans*.

Transposable element mobilization in *C. neoformans* is strongly influenced by

temperature

Previous work in *C. deneoformans* demonstrated that heat stress increases TE

mobilization [35,36]. While some mobile elements in *C. neoformans* followed this pattern,

mobilizing more frequently at elevated temperature, we also uncovered TEs that behave

differently. Several elements (CNL1, KDZ1, KDZ3) were found to mobilize exclusively at 30°C in

vitro. Others (CNIRT6, CNEST) mobilized at both temperatures, but with higher frequency at

37°C. Therefore, by limiting our initial screen to 37°C and examining only ten independent

mutants, we may have overlooked strains harboring mobile elements that are active only at

lower temperatures or mobilizing at low frequency. Interestingly, we found completely non-
overlapping sets of mobilizing TE families in *C. neoformans* and *C. deneoformans* [35,36]. The
discovery of TE families that mobilize more frequently at lower temperatures adds further
complexity to this system, suggesting the involvement of multiple TE- or host-specific
mechanisms in temperature-regulated mobility in *Cryptococcus*. One such mechanism may
involve TCN3 exploiting the host's heat shock response via heat shock elements in its LTR to
facilitate mobilization (Fig 1E). While heat shock-induced TE activation via host machinery has
been demonstrated in plants and animals [91–93], it remains undescribed in fungi.

Other temperature-dependent effects on TE mobility may result from (1) impacts on
functionality of TE-encoded machinery, (2) altered TE suppression, or (3) changes in chromatin
accessibility at *FRR1* facilitating transposition. First, Ty1 mobilization in *S. cerevisiae* is most
frequent at 22°C and absent above 34°C due to inactivity of the Ty1-encoded protease [94].
Similarly, non-functional TE machinery may explain, for instance, why elements from the KDZ
superfamily were only captured mobilizing at 30°C. Second, heat stress can disrupt TE
suppression: in *Drosophila*, Hsp70 displaces PIWI-interacting RNA (piRNA) biogenesis factors
to the lysosome for degradation, leading to transcriptional activation of TEs [95]. In *C.*
neoformans, a recent study found a link between the Calcineurin signaling pathway, which is
necessary for growth at 37°C, and RNAi-mediated silencing of a repetitive transgene,
suggesting a potential mechanism for altered RNAi function during heat stress [96]. Third, heat
stress may increase chromatin accessibility at *FRR1*, facilitating higher TE insertion rates
independent of any direct effects on TE suppression or machinery [97,98]. Elucidating the
mechanisms underlying differential TE mobilization under heat stress in *C. neoformans* will
advance our understanding of the role of TEs in adaptive evolution during infection and in other
heat-stress contexts. Additionally, it is likely that other environmental factors impact TE
mobilization, given that RPMI medium altered TE dynamics across most strains, although it
remains unclear whether this is due to changes in nutrient availability, pH, or other factors (Fig
S7). Moreover, although CNL1 mobilization was only observed in vitro at 30°C, it was the sole
TE implicated in genomic changes during recurrent cryptococcal infection, highlighting that
temperature is not the sole regulator.

DNA transposons are hypomethylated relative to retrotransposons, and are commonly
found in gene-rich regions

Given that *C. neoformans* lacks a de novo cytosine DNA methyltransferase, DNA
methylation as a TE suppression mechanism is relatively inflexible and unlikely to target newly

inserted or unmethylated TE sequences [52]. Despite this, a substantial proportion of TEs
exhibit DNA methylation: 55.6 – 71.2% of LTR elements and 7.7 – 53.4% of non-LTR elements
(Fig S12). In contrast, DNA transposons are less frequently methylated than retrotransposons,
with only 5.4 – 15.5% showing methylation. When methylated, DNA transposons exhibit
significantly lower average 5-methylcytosine (5mC) density compared to LTR and non-LTR
elements (Fig 6). Nonetheless, DNA TEs are still significantly enriched for DNA methylation
compared to protein-coding genes. This reduced level of methylation may contribute to the more
frequent in vitro mobilization of DNA transposons observed in this study, where five of the seven
mobile elements identified were DNA TEs. Furthermore, full-length DNA transposons are more
evenly distributed across the genome and are commonly found within gene-rich regions
compared to LTR and non-LTR retrotransposons (Fig 6). As a result, DNA TEs may be more
likely to mediate genetic changes that alter the expression or function of protein-coding genes.

Transposable elements provide a mechanism for rapid genetic change and facilitate
adaptation in *C. neoformans*

Growth at mammalian body temperature (37°C) significantly increased the rate of
acquired resistance to rap+FK506 in both clinical and environmental isolates (Figs 2–5). In all
cases, this effect was driven by elevated TE insertion rates, highlighting the role of TEs in
generating adaptive phenotypes such as antifungal resistance. These findings suggest that
stress may enhance evolvability within *C. neoformans* populations through bursts of rapid TE
mobilization.

RNA interference (RNAi) is the primary known mechanism for TE suppression in *C.*
neoformans, and loss of RNAi function is associated with hypermutator phenotypes driven by
uncontrolled TE activity [38]. However, recent work has shown that this is not always the case,
as some isolates with loss-of-function mutations in essential RNAi genes maintain low TE
burdens and mutation rates [37]. In this study, none of the clinical or environmental isolates
profiled harbored predicted loss-of-function mutations in RNAi pathway genes. Nevertheless,
isolates from cases 15 and 77 exhibited rap+FK506 resistance rates at 37°C comparable to
those observed in hypermutator strains, including the mismatch repair-deficient *msh2Δ* mutant
[37,38]. This suggests that, in certain strain backgrounds, heat stress alone can be sufficient to
induce a hypermutator-like phenotype through increased TE mobility, even in strains with a
relatively low TE burden (e.g., two full-length copies of CNIRT6 in case 15 isolates).

When assessing the impact of TEs on in-host microevolution, we observed an expansion
of CNL1 copy number in serial isolates collected from a patient with recurrent cryptococcal
meningitis, suggesting a role for TE activity in within-host evolution (Fig 4). However, no
transposition or TE-mediated genomic changes were detected in other longitudinally sampled
clinical isolates or in isolates collected following murine passaging. This may suggest that TE
mobilization events are rare in vivo, that TE-mediated changes did not confer a selective in-host
growth advantage during infection, or that TE-mediated changes occurred in subpopulations not
captured by sampling a single isolate. Previous studies in *C. deneoformans* found evidence of
transposition occurring during murine infection [35,36]. In these cases, however, only drug-
resistant isolates were selected, requiring that loss-of-function mutations at specific reporter loci
had occurred. In contrast, there was no selective pressure for evolved drug resistance or other
traits during the recovery of patient and mice isolates examined in this study.

Genetic background influences TE mobility

Telomere-to-telomere genome assemblies revealed that the presence of full-length TEs
does not reliably predict their mobilization in vitro. For instance, in clonally related case 7
isolates, we observed striking differences in KDZ2 and KDZ3 mobilization at 30°C, despite
these elements being present at identical copy number and genomic positions, with identical
sequences, and lacking DNA methylation in both isolates. Interestingly, of the two single
nucleotide polymorphisms (SNPs) previously identified between case 7 isolates, one was an
amino acid substitution in *SET5* (CNAG_03890), a predicted protein lysine methyltransferase
[29]. This suggests that alternative epigenetic regulation may be driving differential mobility.
Similarly, unmethylated full-length CNEST copies were identified in all assembled genomes
except for case 1; however, mobilization was observed only in the case 77 and cockatoo guano
isolates. Notably, these isolates had the highest CNEST copy numbers, suggesting that copy
number may be positively correlated with increased mobility in vitro. These observations
suggest that isolate- and strain-specific TE mobility may be regulated by TE copy number
polymorphisms, sequence variation within TEs, differential targeting by RNAi, or genetic
variation in host factors that control transposition.

Altogether, our findings highlight the complex interplay between genetic background,
epigenetics, and environmental conditions in governing TE mobilization, with temperature
emerging as a major determinant. Further studies should investigate the extent to which stress-
induced transposition generates genetic diversity and whether this diversification drives

adaptation to infection-relevant stressors and the repeated emergence of antifungal drug
resistance in *C. neoformans*.

METHODS

Strains and growth media

The strains used in this study are listed in Table S1. Strains were grown in YPAD
medium (1% yeast extract, 2% peptone, 2% dextrose, 0.025% adenine) with 2% agar added for
solid media. For host-mimicking conditions, strains were cultured on RPMI plates containing
3.45% MOPS, 0.2% glucose, 2% agar, and 1.04% RPMI 1640 powder (Sigma #R1383).
Mutants resistant to rapamycin and FK506 (rap+FK506) were selected for on YPAD plates
supplemented with 1 µg/mL rapamycin (LC Laboratories) and 1 µg/mL FK506 (Astellas).

Fluctuation assay and mutation spectra

Cultures were inoculated with independent colonies. Each colony was split into 1 mL
cultures of YPAD and grown rotating at 30°C or 37°C for 3 days, allowing all cultures to reach
saturation. Cells were pelleted, washed, and resuspended in water. Cells were appropriately
diluted, plated on solid YPAD medium, and incubated at 30°C for 3 days. Colony counts were
used to calculate colony forming units (CFUs) per culture. The remainder of the culture was
plated on two rap+FK506 plates and incubated at 37°C for four days before counting colonies.
Total CFUs and rap+FK506-resistant colony counts were used to independently calculate the
drug resistance rate for each isolate and temperature using the Fluctuation AnaLysis
CalculatOR (FALCOR), applying the Ma-Sandri-Sarkar (MSS) maximum likelihood method [99].
For all fluctuation assays a minimum of 23 cultures were used to calculate the drug resistance
rate. A single rap+FK506-resistant colony from each culture was selected as an independent
mutant for genomic DNA isolation, performed using either in-house reagents or a MasterPure
Yeast DNA Purification Kit (VWR #76081-694). A minimum of 24 independent drug-resistant
mutants were collected per isolate and temperature for the YPAD mutation spectra analysis.

The *FRR1* gene was amplified for Sanger (Azenta) or linear amplicon long-read
sequencing (Plasmidsaurus). Amplification of *FRR1* was carried out using a variety of
polymerases: MyTaq Mix (Thomas Scientific #C755G91), Apex Taq RED Master Mix (Genesee
Scientific #42-138), Platinum SuperFi II PCR Master Mix (ThermoFisher #12368010), Q5 High-
Fidelity DNA Polymerase (NEB #M0491S), and LongAmp Taq DNA Polymerase (NEB

#M0323S). We often observed "phantom" wild-type–sized secondary bands when amplifying

across TEs, which can occur as a PCR artifact when amplifying across long repetitive DNA

sequences [100]. In these instances, gel extraction using the Zymoclean Gel DNA Recovery Kit

(Zymo Research #D4007) was performed on the higher molecular weight bands, which were

then sent for sequencing. For TEs identified as inserting into *FRR1*, junction PCR with primers

targeting the internal TE sequence and primers flanking *FRR1* were used to confirm a subset of

insertion events, as PCR amplification across long repetitive TE sequences can be difficult

[100]. Primers used to amplify *FRR1* and for TE-specific junction PCR are listed in Table S5.

Sanger sequencing data were analyzed in SeqMan Ultra 17 or DNASTAR Navigator 17

by aligning reads to the wild-type *FRR1* sequence for each isolate. TEs were assigned to

families based on the 80-80-80 rule: nucleotide similarity greater than 80%, over more than 80

base pairs, covering at least 80% of the recovered sequence length [40]. The TE consensus

sequence for TCN3 was obtained from Repbase. For TEs lacking family-level consensus

sequences, GenBank submissions were used to assign family membership (GenBank

accessions: PQ181658 [KDZ1], PQ181660 [KDZ2], PQ181661 [KDZ3], and JQ277267.1

[CNIRT6]). CNL1 family membership was called based on sequence similarity to CNL1 mapped

by Priest et al. (2022) in Bt65 and Bt81 genome assemblies (BioProject accession

PRJNA749953). Domains were identified within full-length TE sequences using InterPro [101].

RPMI mutation spectra

To generate independent mutants during growth on RPMI medium, a prong-plating

procedure was carried out. Single colonies were inoculated in liquid YPAD and grown overnight

rotating at 30°C, then cells were washed and diluted in water. A custom tool containing 121 flat-

tipped pins was used to transfer approximately ~10^4 cells/pin onto non-selective RPMI plates

[102]. To confirm there were no pre-existing mutations, cells were also spotted on rap+FK506

plates. Confluent spots of cells on these plates would indicate mutations occurred during the

overnight growth in YPAD and prong plating would not be possible to generate independent

mutants. If less than 5 colonies appeared on these control plates, experiments were continued.

RPMI plates were incubated for 3 days at either 30°C or 37°C. RPMI plates were then replica

plated to rap+FK506 plates and incubated at 37°C for 3 days. A single colony was picked from

each spot for genomic DNA isolation, followed by *FRR1* amplification and either Sanger

sequencing (Azenta) or junction PCR. A minimum of 18 independent drug-resistant mutants

were collected per isolate and temperature for the RPMI mutation spectra analysis. *FRR1*

amplification, junction PCR, Sanger sequencing data analysis, and TE classification were
carried out as described above.

TE insertion rate

TE insertion rate was calculated using the proportion of independent mutants with a
given TE insertion multiplied by the drug resistance rate. The 95% confidence intervals (CIs) for
proportions were determined using the vassarstats.net proportion CI calculator including
continuity correction. 95% CI for TE insertion rates were determined by a method employing the
square root of the sum of the squares (using the component 95% CI for rates and proportions)
[103].

Transposase and reverse transcriptase protein sequence alignments and phylogenies

Transposase and reverse transcriptase protein sequences were obtained from Yuan and
Wessler (2011) for CMC (CACTA, Mirage, Chapaev) related sequences, Repbase consensus
sequences (https://www.girinst.org/), and from BLASTp (for translated mRNA sequences) or
tBLASTn (for translated genomic sequences) via NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi).
Protein sequences were aligned using MAFFT (https://www.ebi.ac.uk/jdispatcher/msa/mafft)
with the following parameters, protein matrix = BLOSUM62, gap opening penalty = 1.53, gap
extension penalty = 0.123, tree rebulding number = 2, max iterations = 2, and FFTS = none.
Aligned protein sequences were then used to construct maximum likelihood phylogenies with
100 bootstrap replicates using MEGA version 11 [104].

Nanopore sequencing

To extract high molecular weight DNA for library preparation, overnight cultures were
grown in 50 mL liquid YPAD shaking at 30°C. Cell were pelleted and lyophilized overnight.
Genomic DNA was extracted using a cetyltrimethylammonium bromide (CTAB) extraction
method with some modifications [105]. Briefly, cells were lysed by bead beating, and DNA
extraction buffer (100 mM Tris-HCl, pH 7.5; 1% w/v CTAB; 0.7 M NaCl; 10 mM EDTA; 1% v/v 2-
mercaptoethanol) was added and incubated at 65°C for 30 minutes. Chloroform purification was
then performed, and layers were separated by centrifugation. The aqueous layer was isolated,
and DNA was subsequently precipitated, washed, and treated with RNase A. Sodium acetate
(NaOAc) was then added to a final concentration of 0.6 M, and chloroform purification was
repeated. DNA was precipitated again, washed, and spooled to collect high molecular weight
DNA.

DNA quality was assessed using NanoDrop One A260/A280 and A260/A230 ratios

(Thermo Fisher Scientific) and quantified using Qubit dsDNA Broad Range Assay Kit (Invitrogen

#Q32850) and the Qubit 4 Fluorometer (Invitrogen). To assess DNA fragment length, clamped

homogeneous electric field (CHEF) gel electrophoresis was performed using the CHEF Mapper

(Bio-Rad). Libraries were prepared using the SQK-LSK110 or SQK-NBD114.24 and EXP-

NBD104 kits and sequenced using the MinION Mk1C following manufacturer's instructions

(Oxford Nanopore). Sequencing was performed using the MinKNOW software at the default

voltage for at least 48 hours. Basecalling was performed using the latest Guppy software

(v7.1.4, v6.3.8, or v6.4.8), or with Dorado (v0.8.1). The specific sequencing kit, flow cell, and

basecalling methodology used for each isolate is listed in Table S3.

Genome assembly

Demultiplexing was performed using Guppy (v6.5.7). The Canu (v2.2) genome

assembler was used to assemble long reads into contigs, small contigs likely representing

mtDNA or contaminants (< 200 kb) were discarded [106]. For the case 15 incident isolate only,

assembly was performed by combining outputs from Canu and Flye (v2.9.5) [107]. Genome

polishing was performed using long reads generated in this study and short reads available

under the SRA umbrella project PRJNA37160 [29]. Long reads were mapped to contigs using

Minimap2 (v2.24) followed by Medaka (v1.5.0 or v2.0.1) polishing using the appropriate model

[108]. Short reads were mapped to the draft assembly using Bowtie2 (v2.4.4) followed by Pilon

(v1.24) polishing, which was performed in five iterations, re-mapping the short reads to each

new draft assembly before the next iteration of polishing [109,110]. Canu-corrected Nanopore

reads were mapped to the assembly using Minimap2 (v2.24) and coverage was visually

inspected in the Integrative Genome Viewer (IGV) [111]. To confirm generation of telomere-to-

telomere assemblies, contig ends were inspected for the presence of telomere seed sequences

[112]. Assembly breaks at the rDNA locus were artificially merged and separated by 50 Ns.

Contigs were oriented to match the first serially collected isolate assembly to produce final

nuclear genome assemblies. For each isolate, Canu parameters, polishing specifications, and

assembly statistics are provided in Table S3.

DNA methylation detection

5-methylcytosine (5mC) DNA methylation at CpG sites was called using Nanopolish

(v0.14.0) or f5c (v1.5) (see Table S3 for isolate-specific details). Minimap2 (v2.24) was used to

map long reads to the final nuclear assembly for each isolate. For DNA methylation detection

using f5c, POD5 files were converted to BLOW5 using blue-crab (v0.2.0). The Nanopolish

helper script methylation_frequency.py was used to calculate the methylation frequency for CpG

sites.

Centromere mapping

982   Predicted centromere regions were determined using a synteny based approach using

983   the H99 reference genome (GCA_000149245.3) where centromere regions have been

984   experimentally identified [38,42,112]. The 5 kb regions flanking H99 centromeres were used as

985   BLASTn (v2.14.0+) queries against assembled genomes for centromere mapping [113,114].

Full-length TE mapping in assembled genomes

To estimate full-length TE copy numbers from assembled genomes BLAT (v35) and

BLASTn (v2.14.0+) were used, full-length TE sequences obtained from linear amplicon long-

read sequencing of TE insertions into *FRR1* were provided as queries [59]. Full-length copies

were defined as hits with identity ≥ 90% and alignment length ≥ 95% of the query length. For

BLAT the -minIdentity=90 specification was used, and pslReps (v469) filtered hits to find the

best alignment with a minimum coverage of 95% (-minCover=0.95). BLASTn hits meeting full-

length criteria were filtered in R (v4.3.3). In instances were BLAT and BLASTn reported different

full-length copy number estimates, the higher estimate was used. Only BLASTn was used to

detect copies of CNL1 because BLAT reported overlapping CNL1 copies leading to

overestimation, this was not observed for other TEs.

Identification of structural rearrangements and synteny analysis

998   Whole-genome alignment was performed using Minimap2 (v2.24). Dotplots to visualize

999   primary alignments between assemblies were generated in R (v4.3.3) using pafr (v0.0.2).

1000   Synteny and Rearrangement Identifier (SyRI) (v1.6.3) was used to define syntenic regions and

1001   structural rearrangements between serial isolates using default parameters [61]. Plotsr was

1002   used to visualize syntenic regions [115].

Southern analysis

Genomic DNA was prepared using a CTAB DNA extraction method (described above)

and digested with restriction enzymes BspEI, PvuII, or PspXI for CNIRT6, MLDT, and KDZ1,

respectively (New England Biolabs) [105]. Digests were run on 0.7% agarose gels at 4°C for

approximately 16 h. Fragments from the gel were transferred to a charged nylon membrane

(Roche). Target probes were prepared using chemiluminescent DIG-labeling (Roche), see Table

S5 for primers used to amplify probes. Hybridization of the DIG probes (DIG Easy Hyb

Granules, Roche) and detection by chemiluminescence using an anti-DIG antibody (Anti-

Digoxigenin-AP Fab Fragments, Roche) were performed according to the recommended

manufacturer's protocols and imaged on film (Amersham Hyperfilm ECL).

TE annotation and repeat masking

1014 Transposon annotation was performed using Earl Grey (v4.1.0) with default parameters

1015 [79]. Insertions, deletions, and duplications identified by SyRI were repeat masked using

1016 Genetic Information Research Institute (GIRI) Repbase CENSOR [78].

Gene annotation

1018 Gene annotations were lifted over from the H99 reference genome (GCA_000149245.3)

1019 by Liftoff (v1.6.3) with the -polish flag [116].

TE copy number estimation using short reads

TE copy number estimation using short reads was performed as previously described

[37]. Briefly, publicly available short reads (PRJNA37160) were mapped to the H99 reference

genome (GCA_000149245.3) using Minimap2 (v2.24), mosdepth (v0.3.10) was used to

calculate the median read depth in 500 bp non-overlapping windows, the median depth reported

was used for normalization [117]. Short reads were also mapped using Minimap2 to a multi-

fasta containing representative full-length TE sequences identified inserting into *FRR1*.

Mosdepth was then used to produce the cumulative read depth distribution. For each TE, read

depth at 0.95 coverage was normalized by the genome-wide median read depth to estimate full-

length copy number.

RNAi gene loss-of-function prediction

Genome feature annotation was "lifted over" from the H99 reference genome (FungiDB

R68) to each isolate-specific genome assembly using the software tool Liftoff (v1.6.3) [116]. The

`-polish` option of Liftoff was employed to re-align exons in cases where the lift-over procedure

resulted in start/stop codon loss or introduced an in-frame stop codon. Based on the polished

lift-over annotation, the AGAT GTF/GFF Toolkit software (https://github.com/NBISweden/AGAT)

was used to predict protein sequences for all annotated genes in each strain-specific assembly

using the `agat_sp_extract_sequences.pl` script. Where multiple protein isoforms are annotated

in the reference genome, predictions were generated for each isoform. For each RNAi-related

gene of interest, the predicted amino acid sequences for each isolate were compared to the

H99 reference genome. Candidate loss-of-function alleles were classified as those encoding

proteins whose predicted length is < 85% of the length in H99.

Variant analysis of RNAi genes was performed by aligning the coding sequence of each

RNAi gene from each isolate's polished genome assembly using the corresponding Liftoff gene

annotation to the H99 reference genome coding sequence using MAFFT (v7.526) [118]. Snp-

sites (v2.5.1) was then used to generate a variant call file, and the impact of variants was

assessed using SnpEff (v5.3a) [119,120].

1047

DATA AVAILABILITY

DATA AVAILABILITY

Raw nanopore sequencing reads and whole-genome assemblies generated from

isolates used in this study are available on NCBI's Sequence Read Archive (SRA) under

BioProject PRJNA1222431, accession numbers are listed in Table S3. Full-length TE

sequences identified inserting into *FRR1* were uploaded to GenBank, accession numbers are

listed in Table S2.

1054

ACKNOWLEDGEMENTS

ACKNOWLEDGEMENTS

We thank Dr. Marco Dias Coelho for training and assistance with de novo genome

assembly. We also thank Dr. Arturo Casadevall for providing the cockatoo guano isolate, as well

as the isolates collected after murine passage. We thank Drs. Joseph Heitman and Jun Huang

for helpful discussions and for sharing unpublished results on KDZ transposons in *C.

neoformans*. Finally, we thank Dr. Sue Jinks-Robertson for her advice and discussions during

project conceptualization. The isolates from patients with recurrent cryptococcal meningitis in

this study were collected through GERMS-SA surveillance, provided by Dr. N.P. Govender of the

National Institute for Communicable Diseases, South Africa, and shared by Dr. J. Perfect at

Duke University. We thank members of GERMS-SA, 2005 to 2008, including S. Vasaikar

(Eastern Cape); N. Janse van Rensberg, A. Möller, P. Smith, and A. M. Pretorius (Free State);

K. Ahmed, A. Hoosen, R. Lekalakala, P. P. Sein, C. Feldman, A. S. Karstaedt, O. Perovic, J.

Wadula, M. Dove, K. Lindeque, L. Meyer, and G. Weldhagen (Gauteng); S. Harvey and P.

Jooste (Northern Cape); D. Cilliers and A. Rampe (North West Province); W. Sturm, T. Vanmali,

P. Bhola, P. Moodley, S. Sithole, and H. Dawood (KwaZulu Natal); K. Hamese (Limpopo); K.

Bauer, G. Hoyland, J. Lebudi, and C. Mutanda (Mpumalanga); R. Hoffmann, S. Martin, L.

1078

COMPETING INTERESTS

1080 The authors have declared that no competing interests exist.

1081

REFERENCES

1.   Feng G, Leem Y-E, Levin HL. Transposon integration enhances expression of stress

response genes. Nucleic Acids Res. 2013 Jan;41(2):775–789.

2.   Esnault C, Lee M, Ham C, Levin HL. Transposable element insertions in fission yeast

drive adaptation to environmental stress. Genome Res. 2019 Jan;29(1):85–95.

3.   Urquhart AS, Chong NF, Yang Y, Idnurm A. A large transposable element mediates metal

resistance in the fungus *Paecilomyces variotii*. Curr Biol. 2022 Mar 14;32(5):937–950.e5.

4.   McDonald MC, Taranto AP, Hill E, Schwessinger B, Liu Z, Simpfendorfer S, et al.

Transposon-Mediated Horizontal Transfer of the Host-Specific Virulence Protein ToxA

between Three Fungal Wheat Pathogens. MBio. 2019 Sep 10;10(5).

5.   Hays M, Schwartz K, Schmidtke DT, Aggeli D, Sherlock G. Paths to adaptation under

fluctuating nitrogen starvation: The spectrum of adaptive mutations in *Saccharomyces*

cerevisiae is shaped by retrotransposons and microhomology-mediated recombination.

PLoS Genet. 2023 May 16;19(5):e1010747.

6.   Singh NK, Badet T, Abraham L, Croll D. Rapid sequence evolution driven by

transposable elements at a virulence locus in a fungal wheat pathogen. BMC Genomics.

2021 May 27;22(1):393.

7.   Oggenfuss U, Croll D. Recent transposable element bursts are associated with the
1100   proximity to genes in a fungal plant pathogen. PLoS Pathog. 2023 Feb
1101   14;19(2):e1011130.

1102   8.   Lorrain C, Feurtey A, Möller M, Haueisen J, Stukenbrock E. Dynamics of transposable
1103   elements in recently diverged fungal pathogens: lineage-specific transposable element
1104   content and efficiency of genome defenses. G3 (Bethesda). 2021 Apr 15;11(4).

1105   9.   Badet T, Tralamazza SM, Feurtey A, Croll D. Recent reactivation of a pathogenicity-
1106   associated transposable element is associated with major chromosomal rearrangements
1107   in a fungal wheat pathogen. Nucleic Acids Res. 2023 Dec 24;

1108   10.   Wells JN, Feschotte C. A field guide to eukaryotic transposable elements. Annu Rev
1109   Genet. 2020 Nov 23;54:539–561.

1110   11.   Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten
1111   things you should know about transposable elements. Genome Biol. 2018 Nov
1112   19;19(1):199.

1113   12.   Gluck-Thaler E, Ralston T, Konkel Z, Ocampos CG, Ganeshan VD, Dorrance AE, et al.
1114   Giant starship elements mobilize accessory genes in fungal genomes. Mol Biol Evol.
1115   2022 May 3;39(5).

1116   13.   Baduel P, Sammarco I, Barrett R, Coronado-Zamora M, Crespel A, Díez-Rodríguez B, et
1117   al. The evolutionary consequences of interactions between the epigenome, the genome
1118   and the environment. Evol Appl. 2024 Jul 23;17(7):e13730.

1119   14.   Horváth V, Merenciano M, González J. Revisiting the Relationship between
1120   Transposable Elements and the Eukaryotic Stress Response. Trends Genet. 2017
1121   Nov;33(11):832–841.

1122   15.   Fouché S, Badet T, Oggenfuss U, Plissonneau C, Francisco CS, Croll D. Stress-Driven
1123   Transposable Element De-repression Dynamics and Virulence Evolution in a Fungal
1124   Pathogen. Mol Biol Evol. 2020 Jan 1;37(1):221–239.

1125   16.   Sehgal A, Lee C-YS, Espenshade PJ. SREBP controls oxygen-dependent mobilization of
1126   retrotransposons in fission yeast. PLoS Genet. 2007 Aug;3(8):e131.

1127   17.   Paquin CE, Williamson VM. Temperature effects on the rate of ty transposition. Science.
1128   1984 Oct 5;226(4670):53–55.

1129   18.   Servant G, Pinson B, Tchalikian-Cosson A, Coulpier F, Lemoine S, Pennetier C, et al.
1130   Tye7 regulates yeast Ty1 retrotransposon sense and antisense transcription in response
1131   to adenylic nucleotides stress. Nucleic Acids Res. 2012 Jul;40(12):5271–5282.

1132   19.   Scholes DT, Banerjee M, Bowen B, Curcio MJ. Multiple regulators of Ty1 transposition in
1133   *Saccharomyces cerevisiae* have conserved roles in genome maintenance. Genetics.
1134   2001 Dec;159(4):1449–1465.

1135   20.   Hénault M, Marsit S, Charron G, Landry CR. The genomic landscape of transposable
1136   elements in yeast hybrids is shaped by structural variation and genotype-specific
1137   modulation of transposition rate. Elife. 2024 Feb 27;12.

1138   21.   Garfinkel DJ, Nyswaner KM, Stefanisko KM, Chang C, Moore SP. Ty1 copy number
1139   dynamics in *Saccharomyces*. Genetics. 2005 Apr;169(4):1845–1857.

1140   22.   Esnault C, Levin HL. The Long Terminal Repeat Retrotransposons Tf1 and Tf2 of
1141   *Schizosaccharomyces pombe*. Microbiol Spectr. 2015 Aug;3(4).

1142   23.   Lin X, Heitman J. The biology of the *Cryptococcus neoformans* species complex. Annu
1143   Rev Microbiol. 2006;60:69–105.

1144   24.   Goodwin TJ, Poulter RT. The diversity of retrotransposons in the yeast *Cryptococcus*
1145   *neoformans*. Yeast. 2001 Jun 30;18(9):865–880.

1146   25.   Goodwin TJD, Butler MI, Poulter RTM. Cryptons: a group of tyrosine-recombinase-
1147   encoding DNA transposons from pathogenic fungi. Microbiology (Reading, Engl). 2003
1148   Nov;149(Pt 11):3099–3109.

1149   26.   Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, et al. Global
1150   burden of disease of HIV-associated cryptococcal meningitis: an updated analysis.
1151   Lancet Infect Dis. 2017 Aug;17(8):873–881.

1152   27.   Denning DW. Global incidence and mortality of severe fungal disease. Lancet Infect Dis.
1153   2024 Jul;24(7):e428–e438.

1154   28.   Chang CC, Dorasamy AA, Gosnell BI, Elliott JH, Spelman T, Omarjee S, et al. Clinical
1155   and mycological predictors of cryptococcosis-associated immune reconstitution
1156   inflammatory syndrome. AIDS. 2013 Aug 24;27(13):2089–2099.

29.   Chen Y, Farrer RA, Giamberardino C, Sakthikumar S, Jones A, Yang T, et al.
Microevolution of Serial Clinical Isolates of *Cryptococcus neoformans* var. *grubii* and
C. gattii. MBio. 2017 Mar 7;8(2).

30.   Ormerod KL, Morrow CA, Chow EWL, Lee IR, Arras SDM, Schirra HJ, et al. Comparative
Genomics of Serial Isolates of *Cryptococcus neoformans* Reveals Gene Associated With
Carbon Utilization and Virulence. G3 (Bethesda). 2013 Apr 9;3(4):675–686.

31.   Sephton-Clark P, McConnell SA, Grossman N, Baker RP, Dragotakes Q, Fan Y, et al.
Similar evolutionary trajectories in an environmental *Cryptococcus neoformans* isolate
after human and murine infection. Proc Natl Acad Sci USA. 2023 Jan
10;120(2):e2217111120.

32.   Hu G, Chen SH, Qiu J, Bennett JE, Myers TG, Williamson PR. Microevolution during
serial mouse passage demonstrates *FRE3* as a virulence adaptation gene in
Cryptococcus neoformans. MBio. 2014 Apr 1;5(2):e00941–14.

33.   Ewing AD. Transposable element detection from whole genome sequence data. Mob
DNA. 2015 Dec 29;6:24.

34.   Brown SM, Campbell LT, Lodge JK. *Cryptococcus neoformans*, a fungus under stress.
Curr Opin Microbiol. 2007 Aug 17;10(4):320–325.

35.   Gusa A, Williams JD, Cho J-E, Averette AF, Sun S, Shouse EM, et al. Transposon
mobilization in the human fungal pathogen *Cryptococcus* is mutagenic during infection
and promotes drug resistance in vitro. Proc Natl Acad Sci USA. 2020 May
5;117(18):9973–9980.

36.   Gusa A, Yadav V, Roth C, Williams JD, Shouse EM, Magwene P, et al. Genome-wide
analysis of heat stress-stimulated transposon mobility in the human fungal pathogen
Cryptococcus deneoformans. Proc Natl Acad Sci USA. 2023 Jan
24;120(4):e2209831120.

37.   Huang J, Larmore CJ, Priest SJ, Xu Z, Dietrich FS, Yadav V, et al. Distinct evolutionary
trajectories following loss of RNA interference in *Cryptococcus neoformans*. Proc Natl
Acad Sci USA. 2024 Nov 19;121(47):e2416656121.

38. Priest SJ, Yadav V, Roth C, Dahlmann TA, Kück U, Magwene PM, et al. Uncontrolled
transposition following RNAi loss causes hypermutation and antifungal drug resistance in
clinical isolates of *Cryptococcus neoformans*. Nat Microbiol. 2022 Aug 2;7(8):1239–1251.

39. Maziarz EK, Perfect JR. Cryptococcosis. Infect Dis Clin North Am. 2016 Mar;30(1):179–
206.

40. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified
classification system for eukaryotic transposable elements. Nat Rev Genet. 2007
Dec;8(12):973–982.

41. Yuan Y-W, Wessler SR. The catalytic domain of all eukaryotic cut-and-paste transposase
superfamilies. Proc Natl Acad Sci USA. 2011 May 10;108(19):7884–7889.

42. Yadav V, Sun S, Billmyre RB, Thimmappa BC, Shea T, Lintner R, et al. RNAi is a critical
determinant of centromere evolution in closely related fungi. Proc Natl Acad Sci USA.
2018 Mar 20;115(12):3108–3113.

43. Wang X, Hsueh Y-P, Li W, Floyd A, Skalsky R, Heitman J. Sex-induced silencing defends
the genome of *Cryptococcus neoformans* via RNAi. Genes Dev. 2010 Nov
15;24(22):2566–2582.

44. Ilık İA, Yang X, Zhang ZZZ, Aktaş T. Transcriptional and post-transcriptional regulation of
transposable elements and their roles in development and disease. Nat Rev Mol Cell
Biol. 2025 Oct;26(10):759–775.

45. Lisch D. Epigenetic regulation of transposable elements in plants. Annu Rev Plant Biol.
2009;60:43–66.

46. Amselem J, Lebrun M-H, Quesneville H. Whole genome comparative analysis of
transposable elements provides new insight into mechanisms of their inactivation in
fungal genomes. BMC Genomics. 2015 Feb 28;16(1):141.

47. Nicolás FE, Ruiz-Vázquez RM. Functional diversity of RNAi-associated sRNAs in fungi.
Int J Mol Sci. 2013 Jul 24;14(8):15348–15360.

48. Dumesic PA, Natarajan P, Chen C, Drinnenberg IA, Schiller BJ, Thompson J, et al.
Stalled spliceosomes are a signal for RNAi-mediated genome defense. Cell. 2013 Feb
28;152(5):957–968.

49. Burke JE, Longhurst AD, Natarajan P, Rao B, Liu J, Sales-Lee J, et al. A Non-Dicer
RNase III and Four Other Novel Factors Required for RNAi-Mediated Transposon
Suppression in the Human Pathogenic Yeast *Cryptococcus neoformans*. G3 (Bethesda).
2019 Jul 9;9(7):2235–2244.

50. Janbon G, Maeng S, Yang D-H, Ko Y-J, Jung K-W, Moyrand F, et al. Characterizing the
role of RNA silencing components in *Cryptococcus neoformans*. Fungal Genet Biol. 2010
Dec;47(12):1070–1080.

51. Schotanus K, Yadav V, Heitman J. Epigenetic dynamics of centromeres and
neocentromeres in *Cryptococcus deuterogattii*. PLoS Genet. 2021 Aug
31;17(8):e1009743.

52. Catania S, Dumesic PA, Pimentel H, Nasif A, Stoddard CI, Burke JE, et al. Evolutionary
Persistence of DNA Methylation for Millions of Years after Ancient Loss of a De Novo
Methyltransferase. Cell. 2020 Jan 23;180(2):263–277.e20.

53. Dumesic PA, Homer CM, Moresco JJ, Pack LR, Shanle EK, Coyle SM, et al. Product
binding enforces the genomic specificity of a yeast polycomb repressive complex. Cell.
2015 Jan 15;160(1-2):204–218.

54. Desjardins CA, Giamberardino C, Sykes SM, Yu C-H, Tenor JL, Chen Y, et al. Population
genomics and the evolution of virulence in the fungal pathogen *Cryptococcus*
neoformans. Genome Res. 2017 Jul;27(7):1207–1219.

55. Van Wyk M, Govender NP, Mitchell TG, Litvintseva AP, GERMS-SA. Multilocus sequence
typing of serially collected isolates of *Cryptococcus* from HIV-infected patients in South
Africa. J Clin Microbiol. 2014 Jun;52(6):1921–1931.

56. Bovers M, Hagen F, Kuramae EE, Boekhout T. Six monophyletic lineages identified
within *Cryptococcus neoformans* and *Cryptococcus gattii* by multi-locus sequence typing.
Fungal Genet Biol. 2008 Apr;45(4):400–421.

57. Antimicrobial Resistance Division (AMR), Control of Neglected Tropical Diseases (NTD),
Global Coordination and Partnership (GCP). WHO fungal priority pathogens list to guide
research, development and public health action. World Health Organization (WHO); 2022
Oct.

58.   Cruz MC, Cavallo LM, Görlach JM, Cox G, Perfect JR, Cardenas ME, et al. Rapamycin
1244   antifungal action is mediated via conserved complexes with FKBP12 and TOR kinase
1245   homologs in *Cryptococcus neoformans*. Mol Cell Biol. 1999 Jun;19(6):4101–4112.

1246   59.   Kent WJ. BLAT--the BLAST-like alignment tool. Genome Res. 2002 Apr;12(4):656–664.

1247   60.   Fedoroff NV. Molecular genetics and epigenetics of CACTA elements. Methods Mol Biol.
1248   2013;1057:177–192.

1249   61.   Goel M, Sun H, Jiao W-B, Schneeberger K. SyRI: finding genomic rearrangements and
1250   local sequence differences from whole-genome assemblies. Genome Biol. 2019 Dec
1251   16;20(1):277.

1252   62.   Iyer LM, Zhang D, de Souza RF, Pukkila PJ, Rao A, Aravind L. Lineage-specific
1253   expansions of TET/JBP genes and a new class of DNA transposons shape fungal
1254   genomic and epigenetic landscapes. Proc Natl Acad Sci USA. 2014 Feb 4;111(5):1676–
1255   1683.

1256   63.   Coffin JM, Hughes SH, Varmus HE. Structural Classes of Retroelements and Replication
1257   Strategies. 1997;

1258   64.   Eickbush TH, Jamburuthugoda VK. The diversity of retrotransposons and the properties
1259   of their reverse transcriptases. Virus Res. 2008 Jun;134(1-2):221–234.

1260   65.   Cost GJ, Feng Q, Jacquier A, Boeke JD. Human L1 element target-primed reverse
1261   transcription in vitro. EMBO J. 2002 Nov 1;21(21):5899–5910.

1262   66.   Priest SJ, Coelho MA, Mixão V, Clancey SA, Xu Y, Sun S, et al. Factors enforcing the
1263   species boundary between the human pathogens *Cryptococcus neoformans* and
1264   *Cryptococcus deneoformans*. PLoS Genet. 2021 Jan 19;17(1):e1008871.

1265   67.   Yang J, Malik HS, Eickbush TH. Identification of the endonuclease domain encoded by
1266   R2 and other site-specific, non-long terminal repeat retrotransposable elements. Proc
1267   Natl Acad Sci USA. 1999 Jul 6;96(14):7847–7852.

1268   68.   Christensen SM, Ye J, Eickbush TH. RNA from the 5' end of the R2 retrotransposon
1269   controls R2 protein binding to and cleavage of its DNA target site. Proc Natl Acad Sci
1270   USA. 2006 Nov 21;103(47):17602–17607.

69. Fujiwara H. Site-specific non-LTR retrotransposons. Microbiol Spectr. 2015
Apr;3(2):MDNA3–0001.

70. Jurka J. Sequence patterns indicate an enzymatic involvement in integration of
mammalian retroposons. Proc Natl Acad Sci USA. 1997 Mar 4;94(5):1872–1877.

71. Starnes JH, Thornbury DW, Novikova OS, Rehmeyer CJ, Farman ML. Telomere-targeted
retrotransposons in the rice blast fungus *Magnaporthe oryzae*: agents of telomere
instability. Genetics. 2012 Jun;191(2):389–406.

72. Korotko U, Chwiałkowska K, Sańko-Sawczenko I, Kwasniewski M. DNA Demethylation in
Response to Heat Stress in *Arabidopsis thaliana*. Int J Mol Sci. 2021 Feb 4;22(4).

73. Zhang L, Yan L, Jiang J, Wang Y, Jiang Y, Yan T, et al. The structure and
retrotransposition mechanism of LTR-retrotransposons in the asexual yeast *Candida*
albicans. Virulence. 2014 Aug 15;5(6):655–664.

74. Yang D-H, Jung K-W, Bang S, Lee J-W, Song M-H, Floyd-Averette A, et al. Rewiring of
Signaling Networks Modulating Thermotolerance in the Human Pathogen *Cryptococcus*
neoformans. Genetics. 2017 Jan;205(1):201–219.

75. Gao X, Fu Y, Sun S, Gu T, Li Y, Sun T, et al. Cryptococcal Hsf3 controls
intramitochondrial ROS homeostasis by regulating the respiratory process. Nat Commun.
2022 Sep 15;13(1):5407.

76. Kang YS, Jung J, Brown HL, Mateusiak C, Doering TL, Brent MR. Leveraging a new data
resource to define the response of *Cryptococcus neoformans* to environmental signals.
Genetics. 2025 Jan 8;229(1):1–29.

77. Nosanchuk JD, Shoham S, Fries BC, Shapiro DS, Levitz SM, Casadevall A. Evidence of
zoonotic transmission of *Cryptococcus neoformans* from a pet cockatoo to an
immunocompromised patient. Ann Intern Med. 2000 Feb 1;132(3):205–208.

78. Kohany O, Gentles AJ, Hankus L, Jurka J. Annotation, submission and screening of
repetitive elements in Repbase: RepbaseSubmitter and Censor. BMC Bioinformatics.
2006 Oct 25;7:474.

79. Baril T, Galbraith J, Hayward A. Earl Grey: A Fully Automated User-Friendly Transposable
Element Annotation and Analysis Pipeline. Mol Biol Evol. 2024 Apr 2;41(4).

80. Kim JM, Vanguri S, Boeke JD, Gabriel A, Voytas DF. Transposable elements and
genome organization: a comprehensive survey of retrotransposons revealed by the
complete *Saccharomyces cerevisiae* genome sequence. Genome Res. 1998
May;8(5):464–478.

81. Levin HL, Weaver DC, Boeke JD. Two related families of retrotransposons from
Schizosaccharomyces pombe. Mol Cell Biol. 1990 Dec;10(12):6791–6798.

82. Curcio MJ, Lutz S, Lesage P. The Ty1 LTR-Retrotransposon of Budding Yeast,
Saccharomyces cerevisiae. Microbiol Spectr. 2015 Apr;3(2):MDNA3–0053.

83. Curcio MJ, Garfinkel DJ. Single-step selection for Ty1 element retrotransposition. Proc
Natl Acad Sci USA. 1991 Feb 1;88(3):936–940.

84. Behrens R, Hayles J, Nurse P. Fission yeast retrotransposon Tf1 integration is targeted
to 5' ends of open reading frames. Nucleic Acids Res. 2000 Dec 1;28(23):4709–4716.

85. Singleton TL, Levin HL. A long terminal repeat retrotransposon of fission yeast has strong
preferences for specific sites of insertion. Eukaryotic Cell. 2002 Feb;1(1):44–55.

86. Tsukahara S, Bousios A, Perez-Roman E, Yamaguchi S, Leduque B, Nakano A, et al.
Centrophilic retrotransposon integration via CENH3 chromatin in *Arabidopsis*. Nature.
2025 Jan 1;637(8046):744–748.

87. Dai J, Xie W, Brady TL, Gao J, Voytas DF. Phosphorylation regulates integration of the
yeast Ty5 retrotransposon into heterochromatin. Mol Cell. 2007 Jul 20;27(2):289–299.

88. Cordaux R, Batzer MA. The impact of retrotransposons on human genome evolution. Nat
Rev Genet. 2009 Oct;10(10):691–703.

89. Thomas J, Vadnagara K, Pritham EJ. DINE-1, the highest copy number repeats in
Drosophila melanogaster are non-autonomous endonuclease-encoding rolling-circle
transposable elements (Helentrons). Mob DNA. 2014 Jun 4;5:18.

90. Feschotte C, Jiang N, Wessler SR. Plant transposable elements: where genetics meets
genomics. Nat Rev Genet. 2002 May;3(5):329–341.

91. Cavrak VV, Lettner N, Jamge S, Kosarewicz A, Bayer LM, Mittelsten Scheid O. How a
retrotransposon exploits the plant's heat stress response for its activation. PLoS Genet.
2014 Jan 30;10(1):e1004115.

92. Pietzenuk B, Markus C, Gaubert H, Bagwan N, Merotto A, Bucher E, et al. Recurrent
evolution of heat-responsiveness in *Brassicaceae* COPIA elements. Genome Biol. 2016
Oct 11;17(1):209.

93. Garrigues JM, Tsu BV, Daugherty MD, Pasquinelli AE. Diversification of the
Caenorhabditis heat shock response by Helitron transposable elements. Elife. 2019 Dec
11;8.

94. Lawler JF, Haeusser DP, Dull A, Boeke JD, Keeney JB. Ty1 defect in proteolysis at high
temperature. J Virol. 2002 May;76(9):4233–4240.

95. Cappucci U, Noro F, Casale AM, Fanti L, Berloco M, Alagia AA, et al. The Hsp70
chaperone is a major player in stress-induced transposable element activation. Proc Natl
Acad Sci USA. 2019 Sep 3;116(36):17943–17950.

96. Yadav V, Mohan R, Sun S, Heitman J. Calcineurin contributes to RNAi-mediated
transgene silencing and small interfering RNA production in the human fungal pathogen
Cryptococcus neoformans. Genetics. 2024 Jan 27;

97. Kiełbowicz-Matuk A, Smaczniak C, Mikołajczak K, Kuczyńska A, Xu X, Braeuning C, et
al. Heat stress causes chromatin accessibility and related gene expression changes in
crown tissues of barley (*Hordeum vulgare*). Plant Mol Biol. 2024 Oct 22;114(6):115.

98. Huang Y, An J, Sircar S, Bergis C, Lopes CD, He X, et al. HSFA1a modulates plant heat
stress responses and alters the 3D chromatin organization of enhancer-promoter
interactions. Nat Commun. 2023 Jan 28;14(1):469.

99. Hall BM, Ma C-X, Liang P, Singh KK. Fluctuation analysis CalculatOR: a web tool for the
determination of mutation rate using Luria-Delbruck fluctuation analysis. Bioinformatics.
2009 Jun 15;25(12):1564–1565.

100. Hommelsheim CM, Frantzeskakis L, Huang M, Ülker B. PCR amplification of repetitive
DNA: a limitation to genome editing technologies and many other applications. Sci Rep.
2014 May 23;4:5052.

101. Blum M, Andreeva A, Florentino LC, Chuguransky SR, Grego T, Hobbs E, et al. InterPro:
the protein sequence classification resource in 2025. Nucleic Acids Res. 2025 Jan
6;53(D1):D444–D456.

102.   Hum YF, Jinks-Robertson S. DNA strand-exchange patterns associated with double-
strand break-induced and spontaneous mitotic crossovers in *Saccharomyces cerevisiae*.
PLoS Genet. 2018 Mar 26;14(3):e1007302.

103.   Moore A, Dominska M, Greenwell P, Aksenova AY, Mirkin S, Petes T. Genetic control of
genomic alterations induced in yeast by interstitial telomeric sequences. Genetics. 2018
Jun;209(2):425–438.

104.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
version 11. Mol Biol Evol. 2021 Jun 25;38(7):3022–3027.

105.   Pitkin JW, Panaccione DG, Walton JD. A putative cyclic peptide efflux pump encoded by
the *TOXA* gene of the plant-pathogenic fungus *Cochliobolus carbonum*. Microbiology
(Reading, Engl). 1996 Jun;142 (Pt 6):1557–1565.

106.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res. 2017 May;27(5):722–736.

107.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using
repeat graphs. Nat Biotechnol. 2019 May;37(5):540–546.

108.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018 Sep
15;34(18):3094–3100.

109.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012 Mar 4;9(4):357–359.

110.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. PLoS One. 2014 Nov 19;9(11):e112963.

111.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotechnol. 2011 Jan;29(1):24–26.

112.   Janbon G, Ormerod KL, Paulet D, Byrnes EJ, Yadav V, Chatterjee G, et al. Analysis of
the genome and transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex
RNA expression and microevolution leading to virulence attenuation. PLoS Genet. 2014
Apr 17;10(4):e1004261.

113. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.
J Mol Biol. 1990 Oct 5;215(3):403–410.

114. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2009 Dec 15;10:421.

115. Goel M, Schneeberger K. plotsr: visualizing structural similarities and rearrangements
between multiple genomes. Bioinformatics. 2022 May 13;38(10):2922–2926.

116. Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations. Bioinformatics.
2021 Jul 19;37(12):1639–1643.

117. Pedersen BS, Quinlan AR. Mosdepth: quick coverage calculation for genomes and
exomes. Bioinformatics. 2018 Mar 1;34(5):867–868.

118. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: A novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002 Jul
15;30(14):3059–3066.

119. Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, et al. SNP-sites: rapid
efficient extraction of SNPs from multi-FASTA alignments. Microb Genom. 2016 Apr
29;2(4):e000056.

120. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs
in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. Fly (Austin). 2012
Jun;6(2):80–92.

1407

Fig 1. Diverse transposable elements (TEs) captured mobilizing in clinical isolates of *C. neoformans*. (**A**) Schematic of clinical isolate collection from patients with recurrent cryptococcal meningitis at disease onset and after subsequent relapse(s). Single colonies were isolated from patient cerebrospinal fluid on non-selective medium [55]. (**B**) Diagram of the preliminary screen using the "transposon trap" assay. Isolates were cultured in non-selective liquid YPAD medium at 37°C and plated on medium containing rapamycin and FK506 (rap+FK506) (N = 10 per isolate). Point mutations or TE insertions in the ~934 bp *FRR1* (CNAG_03682) reporter gene confer resistance to rap+FK506. (**C**) Illustration of seven mobile TEs detected in *C. neoformans* isolates, including DNA transposons from the CMC (CACTA, Mirage, Chapaev) supergroup and KDZ (Kyakuja/Dileera/Zisupton) superfamily, and retrotransposons from the Ty3/gypsy and R2 superfamilies. Terminal inverted repeats (TIRs) and long terminal repeats (LTRs) are indicated by triangles and arrows, respectively. Genes within TEs that are predicted to facilitate mobilization are shown as white arrows. Functional domains are indicated, including KDZ (Kyakuja/Dileera/Zisupton transposase), CxC1 (predicted zinc chelating domain), AP (aspartic protease), RT (reverse transcriptase), RH (RNase H), and IN (integrase). TEs are drawn to scale. Created in BioRender. Gusa, A. (2025) https://BioRender.com/1psvbhx.

Fig 2. Mobilization of the DNA transposon CNIRT6 is induced by heat stress. (**A**) Rap+FK506 drug resistance rates per cell per generation following growth in non-selective YPAD medium at 30°C or 37°C (N = 23 – 36). (**B**) Mutation spectra in the *FRR1* gene for independent rap+FK506-resistant mutants. The mutation type "Unknown" indicates failure to amplify *FRR1* or absence of detectable mutations in its sequence (N = 24 – 55). (**C**) Insertion rate of CNIRT6 into *FRR1* per cell per generation. (**D**) Chromosome maps of the wild-type case 15 incident, relapse, and relapse 2 isolates. Full-length copies of CNIRT6 are shown as orange circles, and centromeres are shown as dark gray blocks. Chromosome synteny is indicated by chromosome color, with inversions and duplications outlined in dashed and dotted lines, respectively. All chromosomes with identified structural rearrangements are marked with an asterisk. Error bars in (**A**) and (**C**) represent 95% confidence intervals (CIs); rates are considered statistically different if error bars do not overlap.

Fig 3. Mobilization of the DNA transposon CNEST is induced by heat stress. (**A**) Rap+FK506 drug resistance rates per cell per generation following growth in non-selective YPAD medium at 30°C or 37°C (N = 24). (**B**) Mutation spectra in the *FRR1* gene for independent rap+FK506-resistant mutants. The mutation type "Unknown" indicates failure to amplify *FRR1* or absence of detectable mutations in its sequence (N = 40 – 65). (**C**) Insertion rate of CNEST into *FRR1* per cell per generation. (**D**) Chromosome maps of the wild-type case 77 relapse and relapse 2 isolates. Full-length copies of CNEST (gold) and KDZ1 (light blue) are shown as circles, and centromeres are shown as dark gray blocks. Chromosome synteny is indicated by chromosome color. Error bars in (**A**) and (**C**) represent 95% confidence intervals (CIs); rates are considered statistically different if error bars do not overlap.

Fig 4. Temperature- and strain-specific TE dynamics in case 7. (**A**) Rap+FK506 drug resistance rates per cell per generation following growth in non-selective YPAD medium at 30°C or 37°C (N = 25 – 26). (**B**) Mutation spectra in the *FRR1* gene for independent rap+FK506-resistant mutants. The mutation type "Unknown" indicates failure to amplify *FRR1* or absence of detectable mutations in its sequence (N = 33 – 35). (**C**) Chromosome maps of the wild-type case 7 incident and relapse isolates. Full-length copies of TCN3 (green), CNL1 (purple), KDZ3 (blue), and KDZ2 (dark blue) are shown as circles, and centromeres are shown as dark gray blocks. Chromosome synteny is indicated by chromosome color. Insets show the 45 kb regions on the left arms of chromosomes in the incident and relapse isolates. Purple arrows indicate all CNL1 BLASTn hits; grey lines represent DNA methylation frequency from long reads, with the y-axis scaled from 0 to 1. (**D**) DNA methylation frequency plotted for the Chr 3 centromere region ±10 kb of the incident isolate. The position and orientation of full-length TCN3 copies is shown. (**E**) Diagram of HSE sequence motifs (red) identified in TCN3 LTRs. Position is shown relative to the gag-pol open reading frame (ORF) translation start site. Created in BioRender. Gusa, A. (2025) https://BioRender.com/ucajfxd.

Fig 5. Heat stress-induced CNEST activation is conserved in an environmental isolate. (**A**) Rap+FK506 drug resistance rates per cell per generation following growth in non-selective YPAD medium at 30°C or 37°C (N = 23 – 24). Error bars represent 95% confidence intervals (CIs); rates are considered statistically different if error bars do not overlap. (**B**) Mutation spectra in the *FRR1* gene for independent rap+FK506-resistant mutants. The mutation type "Unknown" indicates failure to amplify *FRR1* or absence of detectable mutations in its sequence (N = 42 – 52). (**C**) Chromosome-level synteny between the cockatoo guano and case 77 relapse isolates. Full-length copies of CNEST (gold) and KDZ1 (light blue) are shown as circles.

Fig 6. Genomic distribution and DNA methylation differ among non-LTR, LTR, and DNA transposons. (A) The proportion of the genome composed of non-LTR, LTR, and DNA transposons, based on Earl Grey TE annotations. **(B)** Scaled positions of protein coding genes and non-LTR, LTR, and DNA transposons across chromosome arms. Chromosomes were split at the midpoint of the centromere region, and protein-coding gene and TE positions were scaled from 0 to 1 based on chromosome arm length. Data for all chromosome arms is plotted. **(C)** Mean 5mC DNA methylation density per 100 bp for DNA, non-LTR, and LTR TEs in isolates from cases 1, 7, 15, and 77. Violin plots are overlaid to illustrate the distribution. Significance was assessed using the pairwise Wilcoxon rank-sum test with continuity correction. *** denotes Bonferroni-adjusted $p < 2e\text{-}16$. The first genome assembly from serially collected clinical isolates was chosen as a representative for all cases in **(A)**, **(B)**, and **(C)**.

Fig S1. PCR amplification of *FRR1* in wild-type and independent rap+FK506-resistant mutants. *FRR1* was amplified from independent drug-resistant mutants from case 15 (**A**), case 77 (**B**), and case 7 (**C**); a subset of representative mutants are shown. Higher molecular weight bands indicative of TE insertion in *FRR1* were verified by Sanger or linear amplicon sequencing, or junction PCR. Colored arrows indicate the molecular weights of bands identified as TEs.

Fig S2. CNIRT6 insertion into *FRR1*. (**A**) A subset of CNIRT6 insertion sites (orange arrows) mapped in *FRR1*. The 5′ and 3′ untranslated regions (UTRs) of *FRR1* are underlined; introns are shown in lowercase, exons in capital letters, and the start codon is bolded. (**B**) TSDs detected at CNIRT6 insertion sites. Insertions are ordered based on their appearance in the *FRR1* sequence from beginning to end.

Fig S3. Classification of CNIRT6, CNEST, and CNL1. (**A**) Regions of interest in a multiple sequence alignment highlighting conserved residues in CMC supergroup elements, which include members of the CACTA (also known as EnSpm), Mirage, and Chapaev superfamilies. Highlighted residues include the DDE catalytic core (yellow), conserved cysteine residues (blue) and conserved histidine residues (green) [41]. (**B**) Unrooted amino acid phylogeny of diverse CMC transposases, demonstrating the relationship of CNIRT6 and CNEST with known CMC families and other related sequences. CNIRT6 is marked with an orange star, and CNEST with a gold star. (**C**) Unrooted amino acid phylogeny of reverse transcriptase domains, demonstrating the relationship between CNL1 and MoTeR1, and more distantly related R2 elements from flies and fish, as well as mammalian L1 elements. Branches in (**B**) and (**C**) are colored by kingdom: animals (blue), plants (green), and fungi (yellow). Bootstrap values greater than 50% are shown at nodes. Scale bars represent amino acid substitutions per site. Abbreviations: HS – *Homo sapiens*, DR – *Danio rerio*, DM – *Drosophila melanogaster*, CE – *Caenorhabditis elegans,* CB – *Caenorhabditis briggsae,* PM – *Petromyzon marinus*, BF – *Branchiostoma floridae*, Dpulex – *Daphnia pulex*, Cis – *Ciona intestinalis*, Aca – *Aplysia californica*, Sm – *Schistosoma mansoni*, AA – *Aedes aegypti*, ChOp – *Chionoecetes opilio*, Jewel wasp/NVi – *Nasonia vitripennis*, HM – *Hydra magnipapillata*, Ccinerea – *Coprinopsis cinerea*, RIr – *Rhizophagus irregularis*, Roryzae – *Rhizopus oryzae*, Mlaricis – *Melampsora laricis*, PGr – *Puccinia graminis*, PB – *Phycomyces blakesleeanus*, Mo – *Magnaporthe oryzae*, AT – *Arabidopsis thaliana*.

Fig S4. Putative CNIRT6 transposition events in serially collected clinical isolates. (**A**) Location of the restriction site (dotted line) and CNIRT6-specific probe used in Southern analysis. (**B**) Southern blot of genomic DNA digested with BspEI probed for CNIRT6 in incident (I) and relapse (R) isolates; case numbers are indicated as C#. Asterisks denote the loss of CNIRT6 fragments, while arrowheads indicate the emergence of CNIRT6 fragments at new genomic locations between incident and relapse isolates. (**C**) BLAT homology search hits identified in case 1 isolates using full-length CNIRT6 as the query. The orange dashed line indicates the length of full-length CNIRT6.

Fig S5. CNEST and KDZ1 insertion into *FRR1*. (**A**) A subset of CNEST (gold) and KDZ1 (light blue) insertion sites are mapped with arrows in *FRR1*. The 5′ and 3′ UTRs of *FRR1* are underlined; introns are shown in lowercase, exons in capital letters, and the start codon is bolded. (**B**) TSDs detected at CNEST and KDZ1 insertion sites. Insertions are ordered based on their appearance in the *FRR1* sequence from beginning to end.

Fig S6. KDZ2, KDZ3, CNL1, and TCN3 insertion into *FRR1*. (A) A subset of KDZ2 (dark blue), KDZ3 (blue), CNL1 (purple), and TCN3 (green) insertion sites are mapped with arrows in *FRR1*. The 5′ and 3′ UTRs of *FRR1* are underlined; introns are shown in lowercase, exons in capital letters, and the start codon is bolded. **(B)** TSDs detected at KDZ2, KDZ3, CNL1, and TCN3 insertion sites. Insertions are ordered based on their appearance in the *FRR1* sequence from beginning to end.

Fig S7. Growth in RPMI medium alters TE mobilization dynamics. Mutation spectra in the *FRR1* gene for independent rap+FK506-resistant mutants in case 15 (**A**), case 77 (**B**), and case 7 (**C**). The mutation type "Unknown" indicates failure to amplify *FRR1* or absence of detectable mutations in its sequence (N = 18 – 31).

Fig S8. Transposition events between the cockatoo guano and case 77 relapse isolates. (**A**) Chromosome-level synteny between the case 77 relapse and cockatoo guano isolates. Full-length copies of CNEST (gold), KDZ1 (light blue), KDZ3 (dark blue), and Gypsy-1_CrGa (dark grey) are shown as circles. Predicted centromere regions are indicated by black rectangles and detected transposition events are numbered. (**B**) TSDs detected at KDZ3 and Gypsy-1_CrGa insertion sites.

Fig S9. No predicted transposition of CNEST or KDZ1 in murine-passaged isolates. (**A**) Location of the restriction sites (dotted lines) and CNEST- and KDZ1-specific probes used in the Southern analysis. Southern blots of (**B**) PvuII-digested genomic DNA probed for CNEST, and (**C**) PspXI-digested genomic DNA probed for KDZ1. Shown are case 77 isolates, the cockatoo guano isolate, the patient isolate (PU) infected by the guano point-source strain, and cockatoo guano isolates passaged one to two times through mice by Sephton-Clark et al. (2023), recovered from brain or lung tissue (see Table S1 for strain details).

A

	CNIRT6	CNEST	KDZ1	KDZ2	KDZ3	CNL1	TCN3	Gypsy-1_CaGr
Case 1 Incident	0	1	0	0	0	0	1	1
Case 1 Relapse	0	1	0	0	0	0	1	2
Case 7 Incident	0	1	1	3	2	16	2	12
Case 7 Relapse	0	1	1	3	2	39	2	12
Case 15 Incident	2	1	0	0	0	0	1	1
Case 15 Relapse	2	1	0	0	0	0	1	1
Case 15 Relapse 2	2	1	0	0	0	0	1	1
Case 77 Relapse	0	4	5	0	2	0	0	11
Case 77 Relapse 2	0	4	5	0	2	0	0	8
Cockatoo guano	0	4	5	0	1	0	0	11
H99	2	2	1	0	0	0	1	3

B

Fig S10. Full-length TE copies are present in genomes where no in vitro mobilization was detected. (**A**) Full-length TE copy numbers identified in whole-genome assemblies using BLAT and BLASTn homology searches. TE copy numbers in isolates with observed mobilization events into *FRR1* in YPAD medium are highlighted in pink. Isolate names are colored by lineage: VNI (blue), VNII (green), and VNBII (orange). (**B**) Full-length TE copy number estimates from short-read sequencing for all 15 *C. neoformans* incident isolates screened, except for case 7, where both incident and relapse isolates are shown. Isolates with no insertions into *FRR1* in YPAD medium are shown in grey, and those with insertions are shown in pink.

Fig S11. DNA TEs are approximately uniformly distributed across the genome. Quantile-quantile (QQ) plots are shown for the H99 reference genome and the first genome assembly from serially collected clinical isolates for cases 1, 7, 15, and 77 (representative examples). Observed quantiles correspond to the scaled positions (0 to 1) of DNA TEs along all chromosome arms. Theoretical quantiles are uniformly distributed random values between 0 and 1, sorted in ascending order, with their total number matching the number of observed DNA TEs.

Fig S12. DNA transposons are less frequently methylated than LTR and non-LTR retrotransposons. (**A**) Number of TEs per type and the percentage containing at least one high-confidence DNA methylation call (methylation frequency > 0.75), with median values shown. (**B**) TE types (colors) per genome (shapes) and the percentage with at least one high-confidence DNA methylation call (methylation frequency > 0.75). The first genome assembly from serially collected clinical isolates was chosen as a representative.

Table S1. Strains used in this study.

Table S2. Transposable elements characterized in this study.

Table S3. Whole-genome sequencing and assemblies used in this study.

Table S4. DNA methylation analysis of full-length transposable element copies.

Table S5. Primers used in this study.