Dissecting fluctuating selection: A unified population and quantitative genetics

framework

Esdras Tuyishimire[1], Molly K. Burke[2], Elizabeth G. King[1]

[1] Division of Biology, University of Missouri, Columbia, Mo 65211 USA

[2] Department of Integrative Biology, Oregon State University, Jefferson St, Corvallis, OR 97331

Abstract

One of the longstanding debates in evolutionary biology is the effect of fluctuating selection on

genetic changes in populations. However, the extent to which these periodic forces influence

organisms at both genomic and phenotypic levels remains unclear. Furthermore, despite the

compelling evidence of fluctuating selection from recent studies, there is a disconnect between

empirical findings and theoretical models concerning the underlying mechanisms due to the

limited evidence regarding the scale and processes that generate stable genome-wide oscillations.

This study aims to elucidate the genetic and ecological factors driving fluctuating selection and to

identify the parameters that produce consistent oscillatory patterns in allele frequencies. To address

these longstanding challenges, we developed a modeling framework integrating quantitative and

population genetics to simulate a population under various selection regimes. Using SLiM, a

forward evolution simulator, we varied genetic (heritability and genomic architecture) and

ecological (selection pressure and season length) parameters. Unlike previous models focusing on

selection acting directly on loci, our approach evaluates individual fitness based on the shift in the

seasonal optimum relative to the mean phenotype. We also applied spectral analysis to detect

periodicity, indicating cyclical selective environments. Our simulations shed light on conditions

sustaining oscillations in allele frequencies over time. Spectral analysis successfully identifies the

periodic patterns from allele frequency, even under highly complex selection regimes. Not only

does our study clarify the conditions that yield persistent oscillatory behaviors, but these

parameters are also relatively easy to predict from natural population, providing a possibility of

empirically testing these models.

Introduction

Understanding the dynamic nature of evolution in populations is at the core of evolutionary biology. The direction and intensity of selection can vary significantly through time [1–4].Various factors can lead to these fluctuations, perhaps the most notable of these being seasonal variability. In addition, as climate change imposes unprecedented changes in conditions across the globe, fluctuations in selection pressure are predicted to intensify [5] . Thus, the ability to predict how a population will evolve in response to dynamic environmental conditions has implications for the future of biodiversity. However, while there is widespread appreciation for the relevance of fluctuating selection for populations [6–11], we still lack a clear understanding of the potential impact of fluctuating selection on genome-wide allele frequency dynamics.

Many empirical studies have shown that strong selective pressures can produce rapid phenotypic and genomic changes across diverse species, a phenomenon that has been vigorously documented not only in domesticated plants and animals [12, 13] but also in wild populations exposed to well-defined selective pressures [1, 3, 14] and through laboratory experiments [15–17]. Furthermore, the direction and intensity of selection have been shown to fluctuate over time [1–3, 18–21] due to various factors, including seasonal temperature variability, boom and bust cycles of predator populations, and other biotic and abiotic agents [8, 14, 22–25]. In natural populations, a recent observation is very rapid changes in allele frequencies, with some occurring repeatedly in a seasonal cycle, oscillating back and forth each year (e.g., [9, 26–28]), with the explanation for the pattern being fluctuating selection. For example, Behrman et al. suggested that changes in bacterial populations infecting fruit flies, *Drosophila melanogaster*, due to temperature fluctuations between seasons influence the seasonal cycling of immune genes. In addition, Pfenninger & Foucault, studied fluctuating environments using *Chironomus riparius* and confirmed earlier

results from Rudman et al. about the genomic and phenotypic changes brought by the seasonal

environment on natural populations [9, 28]. However, population genetic models that include

fluctuating selection pressure generally find only a narrow range of parameter space leading to

stable oscillations in allele frequencies [23, 29, 30]. This disconnect between theory and empirical

results has led to disagreement and uncertainty in whether the observed empirical patterns of

oscillating allele frequencies are due to fluctuating selection or some other mechanism [23, 31].

Previous theoretical models of fluctuating selection have typically fallen broadly into two classes:

1) population genetic models that focus on predicting allele frequency trajectories, often focusing

on a single or very few loci; and 2) quantitative genetic models that focus on predicting phenotypic

trajectories and typically assume a polygenic (often infinitesimal) genetic architecture defined by

emergent genetic properties (i.e. heritabilities and genetic correlations) and ignored the dynamics

of individual loci. Past computational limitations of tracking loci genome-wide motivated the

contrast in these two approaches, but while these limits have been diminishing rapidly over time,

most models still fall largely into one framework without fully bridging these approaches, a gap

in the field that has been noted by others (e.g., Barghi *et al.* 2020). Many previous quantitative

genetic models of fluctuating selection have produced robust predictions for resulting phenotypic

evolution [32, 33], the evolution of phenotypic plasticity and bet-hedging [34], and the

maintenance of genetic variation [35, 36]. While most previous population genetic approaches to

fluctuating selection focused on allele frequency dynamics have found only very limited

conditions that will lead to stable oscillations in allele frequencies, recently, some models have

considered more complex scenarios to define the parameters under which allele frequency

oscillations are possible. For example, Wittmann et al. found the maintenance of polymorphism

on hundreds of loci under a segregating lift mechanism, which technically involves the change in

dominance over seasons[30]. Furthermore, the extension of these models indicates that additional

parameters, including genetic and ecological factors, might as well explain the perceived temporal

oscillating polymorphisms[23, 31], even though validation is still required. An important detail of

most population genetic approaches is that they assign a specific selection coefficient to each locus

under consideration. However, in nature, selection acts on the phenotype and dynamics can be

quite different when selection is modeled at the phenotypic level with fluctuating optima, rather

than with a fluctuating selection coefficient acting directly at specific loci. Thus, there are

advantages to integrating quantitative genetic and population genetic approaches in exploring the

possible conditions that might lead to stable oscillations in allele frequencies.

Here, we develop a general modeling framework integrating population genetic and quantitative

genetic approaches, implemented in the powerful and flexible simulation software SLiM [37]. As

in other quantitative genetic models, we parameterize the model by assigning a trait heritability

and defining specific trait optima and fitness functions based on phenotypic values. Like

population genetic models, we specifically consider individual loci genome-wide, tracking

frequency changes over time due to the forces of genetic drift and selection. We sought to identify:

1) The parameter space in which fluctuating phenotypic optima lead to stable oscillating

allele frequencies

2) The circumstances in which patterns of fluctuating selection can be inferred from allele

frequency and phenotypic data

Acquiring knowledge about the above objectives will not only help us understand evolutionary

dynamics, but it will also help in implementing accurate predictive models of evolution, enhance

conservation efforts, and play a role in both disease and pest control, especially nowadays when

species are experiencing various environmental shifts due to climate change.

Methods

Genome size, population setup, and model assumptions

We simulated evolutionary events that use a Wright-Fisher model of a constant population with 10,000 individuals for fluctuating and constant optima and neutral selection. We set a genome size to be approximately 1.2MB for an autosomal chromosome, which varied with a few additional bases (< 1,000 bases) depending on the number of loci affecting the phenotype. Conversely, our study has also considered different genetic architectures: monogenic, oligogenic, and polygenic traits, as shown in the model parameters section. The model assumes known quantitative trait loci (QTL) with a locus effect drawn from exponential distribution. For polygenic and oligogenic models, we assume most loci have minor effects on the phenotype and a few have a large effect size [38]. We did not include a term for genotype by environment interaction in this model, meaning it is not possible for phenotypic plasticity to evolve in our models.

In addition, our simulations assume sexual reproduction with an equal sex ratio, no new mutations, and a recombination rate of 10^{-8}, which is a commonly used value in other SLiM models [37], and it reflects the average recombination rate for many mammals [39]. The initial allele frequency was set to vary among the loci. The probability for an individual to carry a given QTL was drawn from a uniform distribution, and the presence of the allele in individual's genome was determined using a binomial distribution, leading to different initial frequencies across the population. All models were simulated using SLiM V4.2.2 [37], and we tracked each position's frequency and mean population phenotype over generations.

Model parameters.

To better understand the underlying processes governing the changes observed in a population under fluctuating selection, we assessed a range of parameters for genome architecture, heritability, selection strength, and selection length (summarized in Table 1). These parameters are considered essential for affecting the response to selection [40–45]. Our models considered QTLs distributed along the genome (genome size ≈ 1.2MB) with equal neutral regions between loci. In addition, we hypothesize that different values of heritability (check how we defined h^2 in selection models section) might affect the evolvability of a population in different ways, especially when experiencing selection, therefore, we considered its various values in this study. While some previous studies have identified that most fertility-related traits are lowly heritable, in contrast, morphological traits are highly heritable (e.g., fertility vs. height) [44, 46, 47]. Therefore, we predefined the heritability at the initial generation and sought to account for values that encompass a wide range of trait (traits with low to high heritability).

Our study also considered selection strength as a parameter to explore its role in driving genomic and phenotypic diversity within a population. Previous studies show that different selection pressures drive adaptation by influencing evolutionary and population dynamics [48, 49]. Our models used a selection criterion predicated on an individual's fitness derived from the distance between the phenotype and the new optimum (Equation 1). The strength of selection depended on the distance between the population's mean phenotype and the optimum at the current generation, modulated by a constant value that controls the sensitivity of fitness to changes in this distance (Figure 1 & Equation 1). The optima values are defined from the initial generation as the sum or difference of mean phenotype and a standard deviation (referred to as distance to optimum, SD),

depending on which selection model is considered (Equations 2 for constant, 3 for instantaneous,

and 4 for gradual selections).

Moreover, we considered different genetic architectures to explore how they interact with other

parameters. Various theoretical studies have shown that genetic architecture can significantly affect

population evolution while experiencing environmental changes [50]. Remarkably, while single

locus and multiple loci models behaved in a similar manner, e.g., driving the population adaptation

and extinction, researchers show that models with intermediate loci may contain complex

evolutionary behaviors [50–52]. Consequently, our models include a single locus, oligogenic with

10, and polygenic 70, 100, and 300 traits models. The consideration of multiple polygenic models

was to assess if there are significant changes in dynamics as the number of loci for a polygenic

model increases.

The final set of parameters considered in our models is the number of generations over which the

selection event is likely to last (selection length). As described in the next section, this parameter

had different ranges depending on the model under consideration (Table 1). All values were chosen

arbitrarily to illustrate model behavior, and we employ a four-season framework purely as a

complexity system rather than to mimic any specific population. To clarify, in this framework,

each season may correspond to a year during which the population undergoes a particular number

of generations (which may differ from one season to the next, for four years), and then the cycle

repeats.

Figure 1. Change in fitness as the optimum moves away from the phenotype (vertical dashed line). The numbers colored numbers are the fitness values at given optimum and each point is at 0.5 interval.

Table 1. Simulation parameters. Note that each parameter value is considered at once except the generation interval in the four unequal seasons model, where all values are used in a single simulation.

Parameters	Size
Population	10,000
Heritability (h^2)	0.1, 0.5, 0.8
Distance to optimum (SD)	1, 2, 3, 4
Selection length (Two equal seasons)	10, 20, 30
Selection length (Four unequal seasons)	12, 22, 10, 16
Number of loci	1, 10, 70, 100, 300

Selection models

The current study primarily focuses on fluctuating selection resulting from instantaneous changes in optima and extends to both simple and complex gradual changes. Additionally, we explored neutral and constant selection (Figure 2.A) to assess the effect of drift and directional selection, respectively, and how these models differ from fluctuating selection models [53, 54] in maintaining polymorphism.

We implement the instantaneous model by considering two binary environments with two equal seasons, where the change in optimum phenotype is direct, as illustrated in Figure 2.B. Moreover, it is common to find a natural population experiencing gradual environmental changes [55, 56]; a good example is a temperature change over the seasons. Therefore, we extended our models to consider a gradual shift in optimum over generations. The first gradual model (Figure 2.C) helps to study relatively simple evolutionary scenarios, while the second model (Figure 2.D) helps understand complex evolutionary scenarios in which a population may experience a gradual change in optima with multiple seasons, each resulting in a different number of generations, a typical situation for multivoltine organisms that exhibit different generation numbers over the seasons [57–61]. For instance, when the optimum temperature is very high during a given season, it can hinder an organism's development, and also the reproduction rate. Conversely, at moderate temperatures, the development rate can increase to reduce the time to maturity, which can result in many generations [62–64]. Essentially, a population can experience many generations under favorable environmental conditions subjected to a given selective force. In contrast, fewer generations will experience that selective force in a harsh environment, potentially affecting the observed genetic and phenotypic patterns.

For all models, an individual's fitness determined their contribution to the next generation via

sexual reproduction. The fitness (W) for constant and fluctuating selection is a function of the

squared distance between the individual's phenotype (Z) and the optimum (Z_{opt}), the number of

standard deviations (distance to optimum) from the initial mean population phenotype (Equation

1).

$W = \exp\left[-\frac{1}{125} * \left(Z - Z_{opt}\right)^2\right]$ Eq.1

Where the constant 125 controls the steepness of the fitness function. We did not alter this scaling

factor to change the strength of selection, instead altering how far the new optimum was from the

population mean to achieve different strengths of selection.

The phenotype (Z) is calculated as $Z = \sum_{l=1}^{L}(L) + e$ where L represents the total number of

segregating sites affecting trait. Also, e is the environmental effect where $e \sim N(0, VE)$ and VE is

the environmental variance from the heritability equation ($h^2 = VA/(VA + VE)$; VA is the

additive variance) [65, 66]. The expected value of VA was calculated using the equation $VA =$

$\frac{2 * nloci}{3}$, where $nloci$ is the number of loci affecting the trait. This calculation was based on the

results from 100 iterative simulations. Therefore, the expected phenotypic standard deviation is

$sdp = sqrt(\frac{VA}{h^2})$, and the expected mean phenotype $mup = nloci.$ We finally scaled Z to be

$Z_{(adj)} = \frac{Z - mup}{sdp}$ so that the population mean reflects the number of loci contributing to the

phenotype with an expected mean of 0 and standard deviation of 1 to keep the phenotypes on a

similar scale regardless of the number of contributing loci.

The optimum Z_{opt} for each mode of selection was obtained as:

$Z_{opt(constant)} = \bar{Z} + n\sigma_Z$ Eq.2

$Z_{opt(instantaneous)} = \bar{Z} \pm n\sigma_Z$ Eq.3

$Z_{opt\,(gradual)} = \dfrac{\left[\bar{Z} \pm n\sigma_Z * sin\left(2\pi * \frac{T}{i}\right)\right]}{S}$ Eq.4

$\bar{Z}$ is the mean phenotype, n is an integer number $(0 < n < 5)$, σ_Z is the phenotype's standard

deviation for a given generation, T is the current generation, i is the number of generations the

current season (environmental condition) takes, S is scaling factor equals 1 for gradual two equal

seasons and 4 for gradual four unequal seasons (Figure 2).

Figure 2. Illustration of simulation in SLiM featuring positions with known effect and how
optimum changes for each selection model: A reflects the constant model, B shows the instant
change in optimum, C shows the sinusoidal change for two equal seasons, and D is for sinusoidal
change for four unequal seasons. The lines show the change in optimum over generations.
228

Identifying signals of fluctuating selection from allele frequency and mean phenotype

We used ggplot2 [67] to visualize changes in allele frequency and the population mean phenotype

over time. Furthermore, spectral analysis as a mathematical tool that decomposes a time-domain

signal into its frequency-domain representation, it reveals how variance in a time series is

distributed across different frequencies [68]. Consequently, we used spectral analysis to

decompose the allele frequency signal and identify seasonality (the selection period), which

enabled us to detect fluctuating selection in our dataset. Specifically, we first averaged the allele

frequencies across all loci to obtain a single mean frequency per generation. We then organized

these mean values into time-series, with each generation corresponding to a time point, to illustrate

changes in allele frequency over time. Finally, we estimated the spectral density of each time series

using the spectrum function from base R [69] and applied a smoothing span of 2 to reduce noise

and reveal the dominant periodic components. This approach enabled us to detect and characterize

cyclical patterns in allele frequencies across generations.

Results

In our models, the key parameters were (1) selection strength, (2) heritability, (3) the number of

generations per selection season (season length), and (4) genetic architecture. We considered these

parameters for several different selection models, including instantaneous selection (Supplemental

1), gradual two equal seasons (Supplemental 2), gradual with four unequal seasons (Supplemental

3), neutral selection (Supplemental 4), and constant selection (Supplemental 5), with subsections

for parameters' interactions.

Instantaneous selection for polygenic traits

The instantaneous selection model focuses on sudden shifts in selection pressure after a time;

therefore, it provides a direct way to analyze evolutionary dynamics while accounting for multiple

genetic and ecological parameters. We first only consider the model with 100 loci and explore the

effects of the other parameters. Following these results based on a 100 loci model, we vary the

genetic architecture and discuss its effects in the next section. By applying the instantaneous

section model to various combinations of parameters under polygenic traits, we observed

meaningful effects on dynamics of allele frequency and population mean phenotype (Figure 3).

Higher parameter values (SD, h^2, and Gen) often produced more apparent amplitude oscillations

(Figure 3.A), especially in mean phenotypes (Figure 3.B). Also, spectral analysis confirmed these

findings by showing more pronounced peaks under high heritability and strong selection pressure

(Figure 3.C), while it reveals broader peaks when the selection period increased (Supplemental

Figure 1.2.3 - 1.2.4).

Among these parameters, heritability (h^2) had a particularly notable influence on the magnitude of

oscillations (h^2 = 0.8 > 0.5 > 0.1). Highly heritable traits amplified selection effects with

pronounced fluctuations in allele frequencies and phenotypes. Conversely, low heritability masked

changes in allele frequencies and reduced shifts in mean phenotypes, making ongoing selection

less obvious. Despite the allele frequency and phenotypic observations on the effect of heritability,

the spectral analysis showed a more prominent, distinct peak at all heritability levels

(Supplemental Figures 1.2. Heritability increases from top to bottom).

Besides heritability, selection pressure - defined by the distance between the phenotype and the

new optimum - also shaped how allele frequencies and phenotypes evolved over time. More

substantial selection (for example, SD = 4 vs. 1) led to greater fluctuation and variability in allele

frequencies (Figure 3.A), alongside more pronounced oscillations in phenotypes. The spectral

analysis supported these observations, indicating that strong selection reduced variation in spectral

density among replicates, and each unit of selection intensity still produced a clear peak (Figure

3.C).

In addition to heritability and selection pressure, the selection season (number of generations per

season, Gen) further influenced how much fluctuating selection could affect dynamics in the

population. Longer periods (e.g., 30 generations) allowed more time for selection to act, producing

more visible oscillations in allele frequencies and mean phenotypes than shorter periods (e.g., 10

generations). Although these differences sometimes appeared subtle in raw allele frequency data,

spectral analysis detected seasonal patterns under shorter selection periods, albeit with narrower

peaks (Supplemental Figure 1.2.4: from short (SI 1.2.4.1) to long (SI 1.2.4.3) period).

As these parameters interact, different combinations affect the persistence and intensity of

oscillations across generations. For instance, low selection pressure (SD = 1), low heritability, (h^2

= 0.1), and a short season (10 generations) yielded relatively stable frequencies with minimal

fluctuations. Increasing any of these factors, particularly selection pressure or heritability,

intensified oscillation amplitude. Notably, combining high selection pressure (SD = 4), high

heritability, (h^2= 0.8), and a long season (30 generations) produced persistent oscillations across

multiple loci (Figure 3.A, lower right). However, higher fluctuations tend to be intensified for loci

at intermediate initial frequencies (between 0.25 and 0.75). Additionally, phenotypic trends

paralleled these allele frequency changes and became more evident under stronger selection or

higher heritability. Unlike allele frequency, phenotypic change is very apparent when there is an

increase in either one of the parameters (heritability, selection period, and intensity), a condition

that is hard to quickly notice for heritability and selection season by simply visualizing allele

frequencies.

Spectral analysis under the instantaneous selection model further clarifies how those fluctuations

persist across parameter combinations. It consistently shows a peak for seasonal selection,

regardless of parameter size and combinations. However, it is possible to deduce the effect of each

parameter through comparisons, especially with multiple replicates. Smaller parameters cause

significant variability in spectral density, while increases in parameters like heritability and

selection strength reduce this variability (see Figure 3.C). Overall, high selection intensity, a longer

selection period, and substantial additive genetic variance maintain genome-wide variation over

generations, allowing observed allele frequency oscillations to persist for many generations.

*Figure 3. Evolutionary dynamics for a population with polygenic trait (loci = 100) experiencing an instantaneous fluctuating selection. **A**. represents allele frequency dynamics and shows parameter combinations from a single replicate for 1000 generations. Each line in the plot represents the frequency trajectory of a locus among 15 loci randomly sampled from 100 loci, and the magenta line highlights dynamics of single locus. **B**. Distribution of population mean phenotype dynamics representing 30 replicates over 1000 generations with red line representing the mean across all replicates. **C**. The spectral density from allele frequency for all 2000 generations; each colored line represents a single replicate. The large peak in the spectral plots marks a complete cycle of selection season. The variables are heritability (h^2), selection strength (SD: number of standard deviations also known as distance to optimum), and the selection length in terms of number of generations (Gen).*

Genetic architecture: Polygenic, Monogenic, and Oligogenic Traits

In addition to evaluating the parameters that influence an instantaneous selection on polygenic trait, we also examined how genetic architecture affects observed evolutionary dynamics. We hypothesized that the number of loci influencing the trait could interact with other parameters, heritability, selection strength, and selection duration. Overall, the patterns identified in our polygenic model extend to monogenic and oligogenic traits, although fewer positions affecting the trait tend to produce more pronounced oscillations in allele frequencies and phenotypes. That makes it easier to distinguish the influence of each parameter, mainly heritability and selection duration, which exert more subtle effects in polygenic contexts.

Overall, monogenic traits display the most conspicuous cyclic fluctuations in allele frequencies and allow seasonality detection (Supplemental figure 1.1.1). Oligogenic traits also exhibit clear, though somewhat weaker, oscillations than monogenic traits (Figure 4) due to some loci demonstrating no great changes in frequency. In monogenic and oligogenic models, higher heritability and intense selection accelerate and amplify allele-frequency shifts, whereas lower parameter values yield weaker oscillations (Supplemental Figures 1.1.1 and 1.1.2). Although polygenic traits seem to distribute selection effects across many loci, there are fewer dramatic changes per single locus as seen from figure 4. Subsequently, analyzing all loci collectively can reliably reveal cyclical patterns in such a complex genomic architecture while using allele frequency and phenotypic trajectories. These differences are driven by the fact that the heritability of the trait is held constant in each case. Thus, for a given heritability, having fewer causative loci means that each of those loci will contribute a larger share to the percentage of genetic variation for the trait and thus have a larger genetic effect. Thus, the observation that individual loci show

greater fluctuations when there are fewer loci is driven by this effect, rather than it being a feature

of the genetic architecture itself.

Moreover, spectral analysis effectively identifies seasonality/oscillations under all genetic

architectures but is less effective at distinguishing differences among these architectures for a

single replicate (Although not shown). Monogenic and oligogenic models often show distinct

peaks with high spectral densities, while polygenic traits typically have lower spectral densities

(Figure 4.C). Nevertheless, the greater variation in spectral peaks across replicates in monogenic

traits may demonstrate the susceptibility to genetic drift. Oligogenic traits occupy an intermediate

position with more remarkable changes in a few key loci, particularly under high heritability and

intense selection. Polygenic traits, however, require higher heritability and selection strength to

display cyclical changes visibly across many loci, but the spectral analysis clearly shows that it

more reliably detects fluctuations in the polygenic model. As a result, even though the shifts are

more subtle when there are more loci sharing the effect, the analysis across all of them gives a

well-defined signal. Ultimately, despite the differences, spectral analysis reliably detects

seasonality across all genetic architectures, particularly when considering aggregate behavior

across loci.

Finally, in addition to the described means of differentiating behaviors per genomic architecture,

we calculated the percentage of fixed loci, as they may as well reflect the variability in spectral

peak across replicates. The results show that when relatively few loci (monogenic and oligogenic)

affect the trait, the proportion of fixed alleles is substantially higher than in polygenic systems

(Figure 5). In polygenic traits, increasing the number of causative loci does not notably alter

fixation rates, and other parameters (e.g., heritability or selection duration) have little impact on

this outcome (Figure 5). The fixation rates are percentages that were calculated as the ratio of the

total number of loci fixed at one and lost to the total number of loci from all replicates, multiplied

by 100.

Figure 4. *Comparison of genetic architectures (monogenic and oligogenic traits) under*
instantaneous fluctuating selection. ***A.*** *represents allele frequency dynamics for a single replicate*
for small (Min combo: $h^2 = 0.1$, SD = 1, Gen = 10), medium (Med combo: $h^2 = 0.5$, SD = 2, Gen
= 20), and large (High combo: $h^2 = 0.8$, SD = 4, Gen = 30) parameter combinations over 1000
generations. Each line represents the frequency trajectory of a locus, with the magenta line
highlighting the dynamics of a single locus. ***B.*** *Shows distribution of population mean phenotype*
dynamics for 30 replicates over 1000 generations, with the red line representing the overall mean
for both small and high parameter combinations in each architecture. ***C.*** *displays spectral density*
of allele frequency over 2000 generations, where each colored line represents a single replicate

for both small and large parameter combinations for each architecture. The large peak in the
spectral plot marks a complete cycle of the selection season.

Figure 5. *Percentage of fixed loci across different genetic architectures based on parameter*
interactions. The monogenic trait exhibits the highest percentage of fixed loci among the 30
replicates, followed by the oligogenic model, while the polygenic model shows a consistent fixation
rate.

Beyond instantaneous selection: gradual selection models

Apart from the instantaneous model, we explored two gradual optima-change models, one with

two equal seasons and another with four unequal seasons, to capture more complex scenarios of

fluctuating selection [70]. These gradual-change models introduce selection more slowly, with

smooth transitions between seasonal optima; hence, they reflect natural environments where

conditions do not shift abruptly.

Unlike the instantaneous model, the gradual two-season model tends to obscure clear seasonal

selection events, particularly in polygenic traits (Figures 6 & 7). Although gradual two-season

behaves similarly to the instantaneous model, its fluctuations are less pronounced as the slower

transition reduces the overall selection impact. When selection strength is relatively low,

increasing heritability or extending the selection season does not produce the same prominent

allele frequency shifts seen under instantaneous selection (Figure 5, see Supplemental 2.1).

Observing cyclical allele frequency changes requires high selection intensity, especially in

polygenic traits. Monogenic and oligogenic traits (Supplemental Figure 2.1.1 and 2.1.2), although

fluctuations are weaker than in the instantaneous model, remain detectable. Likewise, spectral

analysis can identify periodicity effectively in these data, showing no major differences between

the gradual two-season and instantaneous models (Supplemental Figures 2.2).

The four-season gradual model is more complex than others because it involves different selection

season lengths (12, 22, 10, and 16 generations) in a single cycle. Thus, we do not isolate the effect

of season length but instead focus on heritability and selection intensity. This setup reflects

environments with uneven selection pressures throughout the year or across life stages,

complicating evolutionary dynamics and blurring distinctions between selection and neutral drift

(see later section). Even under high parameter combinations, observing periodic shifts in allele

frequencies for polygenic traits is challenging (Figure 7.A) under this type of selection model. In

contrast, monogenic and oligogenic traits can still exhibit noticeable oscillations at certain loci

under strong selection or high heritability (Thus, additional scenarios are needed to explore the

effect of specific season lengths).

Moreover, phenotypic changes generally follow the same trends as allele frequencies and can

sometimes reveal more distinct oscillations (7.A and 7.B). However, these oscillations are still less

prominent than in two-season or instantaneous models. Hence, as the selection model becomes

more complex, higher selection intensity is required for parameters such as heritability and season

length to effectively uncover seasonality in allele frequencies or phenotypes, as both gradual

models demonstrate. While the gradual two-season model can display phenotypic fluctuations at

lower heritability and shorter selection periods, the more complex four-season model struggles to

produce visible allele oscillations under low parameters. Both gradual models still reveal

distinctive phenotypic changes in monogenic and oligogenic traits, but polygenic traits under four-

season fluctuations show the least pronounced patterns, indicating that as patterns of fluctuation

in selection become more complex, it may be more challenging to detect fluctuating selection in

the genome.

Although clear spectral peaks are often difficult to detect in the four-seasons model, spectral

analysis remains more effective than direct inspection of allele frequency trajectories for

identifying periodic signals (Figure 7.C, Supplemental Figures 3.2.1 and 3.2.2). Under low

selection intensity, the spectral density peak among replicates is not well-defined, regardless of

heritability; however, highly heritable traits present less noise, thus increasing the chance of

recognizing the periodic patterns more readily. The results emphasize that genetic architecture and

selection model complexity significantly influence how populations respond to fluctuating

environments. Also, spectral analysis remains a powerful tool for confirming periodic selection

signals across various evolutionary scenarios while visualizing allele frequencies and phenotypes

can help deduce initial conclusions.

__Figure 6__: Allele frequencies from 15 sampled loci out 100 to compare all fluctuating selections models with instantaneous selection showing clearer seasonal oscillations, the gradual two-season, and finally gradual four unequal-season model with a $h^2 = 0.5$, SD =2, and Gen (for two season models) = 20.

*Figure 7: Comparison of gradual selection models. **A**. Allele frequency trajectories for a single replicate of monogenic, oligogenic, and polygenic traits are shown. In the polygenic model, only 15 of 100 loci are displayed. The highlighted loci illustrate that the dynamics are compared within the same genetic architecture (i.e., the highlighted positions are not identical across different architectures). **B**. Phenotypic dynamics for 30 replicates are presented, with the overall mean per generation highlighted. **C**. Spectral analysis for 30 replicates is displayed, with each color representing a different replicate. For all models, the parameters are $h^2 = 0.5$, SD = 0.2, and Gen = 20. Thus, the spectral peaks in both the instantaneous and gradual two-season models occur at 40, representing a full seasonal cycle. The four-season models exhibit four peaks, corresponding to the four-generation intervals, with a complete cycle occurring at twice the generation interval per season.*

Difference between fluctuating selection models to neutral and constant selection models

Comparing fluctuating optima models with neutral and constant selection models is indispensable to fully understanding how fluctuating selection affects allele frequencies and phenotypes. Under both neutral and constant selection scenarios, neither allele frequencies nor phenotype trajectories display clear seasonal oscillations, as confirmed by spectral density analyses (see supplemental file for more details: Sections 4 and 5 for neutral and constant selections, respectively).

In neutral models, there are random fluctuations with no discernible pattern mainly because allele frequencies change solely via genetic drift. That is, all the genetic architectures show similar allele frequency dynamics, and loci with terminal initial frequency fix (Supplemental figure SI 4.1). By contrast, constant selection models exert steady selective pressures, leading to predictable shifts in allele frequencies and phenotypes until they plateau, either when certain alleles fix, are lost, or when the population reaches its optimum. The plateauing behavior is particularly pronounced in polygenic models (Supplemental Figures, sections 5.1 and 5.2 for allele frequencies and phenotypes, respectively, under constant selection). Monogenic traits under strong selection often go to fixation more quickly, while polygenic and oligogenic traits tend to remain within specific

frequency ranges over many generations. Such behavior does not appear under neutrality, where

frequencies drift randomly.

Despite these distinctions, spectral analyses of neutral and constant selection models reveal no

evident periodicity in either regime (see Supplemental file, section 4.3 and 5.3 for neutral and

constant selections, respectively). Consequently, it is difficult to pinpoint out the effects of

individual parameters via spectral analysis in these two models.

In contrast, fluctuating selection models, both instantaneous and gradual, introduce seasonally or

temporally varying selective pressures that can drive noticeable oscillations in allele frequencies

and phenotypes. These oscillations are especially marked in monogenic and oligogenic traits, as

well as in polygenic traits with high heritability under strong selection. However, when selection

intensities or heritability are low, the behavior of polygenic traits under fluctuating selection may

appear similar to that under neutral or constant selection (Figure 8). Insufficient selection pressure

or limited heritability prevents large frequency shifts and cause some alleles to remain in stable

ranges (mimicking constant selection) or to drift randomly (mimicking neutrality). As selection

intensity and heritability increase, however, the oscillatory patterns become more pronounced,

clearly setting fluctuating selection apart from either purely directional (constant) or random

(neutral) processes.

The gradual two-season model often exhibits easily detectable seasonal oscillations in allele

frequencies, much like the instantaneous model. In more complex scenarios, such as the four

unequal season model, these oscillations may be harder to distinguish from neutrality, particularly

for polygenic traits (Figure 8.A and 8.B). At lower parameter values, the peaks in spectral analyses

may be weak, resembling neutral drift. Nonetheless, increasing certain parameters can amplify the

oscillatory signal, making it possible to differentiate fluctuating selection from neutrality in some

cases.

***Figure 8**: A represents a neutral model and B representing the four unequal-season model for each genomic structure with low parameter combinations. The oligogenic and polygenic trait models for both selection regimes are very similar although the gradual model represents less variation among replicates, and the monogenic trait shows the spectral peak representing the involved seasons for gradual model.*

Discussion

Parameters leading to persistent allele frequency oscillations

A topic of recent debate has been the conditions under which fluctuating selection will lead to stable oscillations that persist across time and whether the recently observed empirical results [6] are genuinely due to fluctuating environmental conditions. Previous models have produced mixed results, with some showing restrictive parameter spaces leading to stable oscillations [31, 36] and others suggest numerous loci may remain polymorphic under certain seasonal dominance assumptions [30] and protection from selection [23]. However, it remains challenging to empirically verify those conditions, partly because dominance relationships and other relevant parameters, for instance, degree of protection, are often difficult to quantify in natural populations, as noted by the authors. In contrast, our study examines relatively easy-to-measure parameters in natural populations and demonstrates that cyclic allele frequencies can persist at many positions even under low selective pressure, particularly when selection acts instantaneously. However, the magnitude of these oscillations scales directly with selection strength; thus, it is the primary determinant of how pronounced the cyclical patterns are, and the combination of selection strength with other parameters leads to numerous positions with stable oscillations.

When both heritability and selection intensity were high, the population exhibited pronounced oscillations, a finding consistent with classical quantitative genetic theory [44, 71, 72]. Moreover, with intermediate to high selection pressure and heritability, we can visually see those fluctuations at high amplitude, contradicting the claim that not all positions will experience strong selection [73]. Under these conditions, we observed marked fluctuations and, at times, near-fixation of alleles, consistent with other reports suggesting that increased selection intensity may lead to allele fixation [41, 53, 74–77]. However, complete fixation is often prevented because the environment

fluctuates, thereby maintaining numerous alleles at intermediate frequencies. Furthermore, longer

seasons grant sufficient time for selection to shift allele frequencies, as seen in the evolutionary

experiment study by [15] and [78], and mean phenotypes toward the new optimum, accentuating

oscillations. Conversely, shorter selection periods can diminish these effects because the

environment changes again before the population fully adapts, as seen in classic field studies such

as Grant and Grant's finch research [1] and theoretical predictions [79]. Also, the fact that we

cannot observe the oscillation directly from allele frequencies does not mean that the fluctuating

selection is not happening and is involved in maintaining polymorphism [80]. Our results illustrate

that extending the selection season amplifies the visible shifts in allele frequencies, especially

under high selection pressure, thereby producing more pronounced cyclical patterns. Oscillations

are very challenging to observe for individual loci in polygenic traits with small parameter

combinations, however, spectral analysis readily recovers the spectral peak, especially in the

instantaneous model and the two gradual season models. Therefore, the identification of oscillatory

behavior will be predicated on the parameter combination and the complexity of the selection

regimes. That is, trait's heritability, length of selection season and its strength will determine the

level of oscillation among alleles.

Genomic architecture effect on oscillation and fixation

Previous theoretical models investigating the effects of fluctuating selection on genome dynamics

often used the approach of selection acting directly on the causative loci, generally demonstrating

that polymorphism could be maintained under specific conditions, notably if heterozygote fitness

exceeded that of homozygotes [81, 82]. They also indicated that single-locus models could lead to

fixation over extended periods of fluctuating selection, thus limiting long-term polymorphism [32,

35, 82, 83]. In contrast to those models that focused on a single locus selection coefficient, our

study employs a phenotype-based selection approach [84, 85], where changes in environmental

conditions shift the optimal phenotype and shifts in selection on the causative loci are emergent.

The results support temporary genetic diversity by showing that monogenic models exhibit larger

fluctuations, mainly because a single genetic locus drives the entire phenotypic shift for a given

heritability. In some cases, these fluctuations persist until fixation occurs, while in others they

endure for an extended period. Although monogenic models tend to show a high rate of fixation,

alleles do not always fix, and the degree of fixation depends on the specific parameter combination.

Extending our analysis from single-locus models to oligogenic and polygenic frameworks reveals

that selection acts simultaneously across multiple loci, leading to oscillations in allele frequencies

and phenotypes under appropriate conditions. In contrast to single-locus models, where parameter

changes strongly influence allele fixation, polygenic models tend to mask selection effects at

individual loci, with the cumulative impact spreading across a large genomic architecture [86–89].

Therefore, that diffused selection effect can explain the need for intense selection on highly

heritable polygenic traits to sustain observable oscillations in allele frequencies across many loci.

In addition, our current results indicate that alleles with intermediate initial frequencies fluctuate

more than others. This pattern could reflect heterozygote advantage [32] or an antagonistic

pleiotropy, where one allele is favored in one season and disfavored in the next [30, 90]. If this

interpretation holds, then the parameters in our study would more effectively explain the observed

oscillations, and they might be easier to predict than the reversal dominance model proposed in

previous research. Therefore, further investigation into potential overdominance effects, including

elevated genetic load, is warranted, as discussed by Wittmann et al. (2017).

Fluctuations in complex selection regimes

Our study contrasts two principal types of fluctuating selection models: (1) instantaneous (where

the environment switches sharply between two states) and (2) gradual (where the environment

transitions more smoothly through two equal or four unequal seasons). The observed similarities

in instantaneous and gradual two equal season models likely arise from having constant season

lengths and selection peaks. In contrast, when there are four unequal seasons with different peaks

strength (highest optimum for each season), the population struggles to adapt to each peak; thus,

the complex oscillatory dynamics. Nevertheless, gradual models better represent complex natural

environments where multiple selection regimes occur over time. These more complex scenarios,

especially those with four unequal seasons, create subtle patterns that can resemble neutral or

constant selection when parameters (e.g., selection strength, heritability) are low [30, 91, 92].

Indeed, a particularly long or short season under these gradual models can either strengthen or

weaken selection signals, respectively, making detection more difficult [19, 93, 94] for the latter.

We employed spectral analysis to tackle the challenge of identifying subtle oscillatory signals.

This approach successfully discerned cyclical patterns in allele frequency across most parameter

combinations, particularly in simpler models (i.e., the instantaneous and two equal seasons

models). However, spectral analysis struggled to detect periodicity when parameter values were

low (e.g., low heritability or weak selection) or when the model included multiple complex

environments (as in the four-unequal-season model). These difficulties mirror empirical findings

[11] and may be explained by the short duration of each season, which hinders the population's

ability to adapt to new conditions. A similar challenge in identifying seasonality was observed in

the statistical analysis of experimental data from Bergland et al. (2014), who reported seasonal

allele-frequency oscillations in thousands of SNPs. However, a subsequent reanalysis [95] failed

to replicate this result, likely due to the complex selection pressures experienced by the population

or the timing of sample collection (i.e., the generation intervals used by Bergland et al. in their

2014 data set). These results highlight and define the limits for detecting fluctuating selection from

allele frequency data under complex scenarios.

Ideally, sequencing multiple sequential generations for a long time would help identify seasonality

from genomic data using the currently developed statistical tools. However, its feasibility is almost

impossible due to the associated cost and labor. Although Phillips et al. 2020 [96] did not examine

fluctuating selection, they argued that taking more samples over time can dramatically improve

our ability to understand the complexity into evolutionary dynamics. Therefore, researchers

studying populations under complex fluctuating conditions should also aim to estimate the optimal

sampling points required to capture such dynamics. Furthermore, new computational methods,

including machine learning and generative models, may reduce empirical demands or detect

cyclical selection signals even in sparser datasets [97].

Limitations and Future Directions

Our simulations assume non-overlapping generations, constant population sizes, no new

mutations, and fixed recombination rates, all of which may limit the extent to which these results

can generalize. In addition, our models did not allow for any genotype by environment interaction

at any loci. Previous quantitative genetic models of fluctuating selection have frequently identified

the conditions under which fluctuating selection is expected to lead to the evolution of phenotypic

plasticity driven by genotype by environment interactions [98–100]. Our models did not allow for

the evolution of plasticity in this way, removing a major potential way for populations to adapt to

fluctuating selection pressures. Thus, our results shed light on whether it is ever possible for

fluctuating selection to drive recurrent, stable, allele frequency oscillations in a population, given

these simplifying assumptions. Future models incorporating additional factors, including allowing

for mutation, genotype by environment interactions, and incorporating dynamic population sizes

are necessary to capture the complexity of natural populations [45, 101–103]. Future research

should also test these observations through evolve-and-resequence experiments and refine

methods for parameter estimation from time-series genomic data. This work offers a foundation

for understanding how traits evolve and persist under dynamic natural conditions by revealing how

seasonal and environmental changes modulate selective pressures over time.

Concluding remarks

This study highlights the intricate interplay among heritability, genomic architecture, selection

intensity, and season length in shaping oscillatory allele-frequency dynamics under fluctuating

environments. High heritability and strong selection produce pronounced and persistent frequency

oscillations, and these effects become more evident when selection acts over longer seasons.

Meanwhile, the broad genomic architecture of polygenic traits can mask individual locus

fluctuations, yet substantial oscillations still emerge when selection is sufficiently strong. Spectral

analysis is valuable for detecting cyclical patterns, but its effectiveness diminishes in highly

complex or weakly selective environments. Therefore, improved methods are needed to identify

seasonality from very complex models.

Overall, our results underscore the importance of the discussed ecological and genetic parameters

in understanding the evolutionary fate of populations under fluctuating selection. Accurately

modeling additional parameters and empirically testing these dynamics will be key to predicting

evolutionary outcomes and identifying conditions that promote sustained genetic diversity as

environmental variability increases.

Reference:

1. Grant PR, Grant BR (2002) Unpredictable evolution in a 30-year study of Darwin's
finches. Science (1979) 296:707–711
2. Bell G (2010) Fluctuating selection: the perpetual renewal of adaptation in variable
environments. Philosophical Transactions of the Royal Society B: Biological Sciences
365:87–97
3. Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards S V.
(2017) Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the
green anole lizard. Science (1979) 357:495–498
4. Endler JA (2020) Natural Selection in the Wild. (MPB-21), Volume 21.
https://doi.org/10.2307/j.ctvx5w9v9
5. Gienapp P, Reed TE, Visser ME (2014) Why climate change will invariably alter
selection pressures on phenology. Proceedings of the Royal Society B: Biological
Sciences 281:20141611
6. Bergland AO, Behrman EL, 'brien O, Schmidt KR, Petrov PS (2014) Genomic Evidence
of Rapid and Stable Adaptive Oscillations over Seasonal Time Scales in Drosophila.
PLoS Genet 10:1004775
7. Yair S, Coop G (2022) Population differentiation of polygenic score predictions under
stabilizing selection. Philosophical Transactions of the Royal Society B: Biological
Sciences. https://doi.org/10.1098/rstb.2020.0416
8. Van Buskirk J, Smith DC (2021) Ecological causes of fluctuating natural selection on
habitat choice in an amphibian. Evolution (N Y) 75:1862–1877
9. Pfenninger M, Foucault Q (2022) Population Genomic Time Series Data of a Natural
Population Suggests Adaptive Tracking of Fluctuating Environmental Changes. Integr
Comp Biol 62:1812–1826
10. Thomas E Reed, Daniel E Schindler, Robin S Waples (2011) Interacting Effects of
Phenotypic Plasticity and Evolution on Population Persistence in a Changing Climate.
Conservation Biology 25:56–63
11. Bitter MC, Berardi S, Oken H, Huynh A, Lappo E, Schmidt P, Petrov DA (2024)
Continuously fluctuating selection reveals fine granularity of adaptation. Nature 2024
634:8033 634:389–396
12. Trut L, Oskina I, Kharlamova A (2009) Animal evolution during domestication: the
domesticated fox as a model. BioEssays 31:349–360
13. Doebley JF, Gaut BS, Smith BD (2006) The Molecular Genetics of Crop Domestication.
Cell 127:1309–1321
14. Waldvogel A, Wieser A, Schell T, Patel S, Schmidt H, Hankeln T, Feldmeyer B,
Pfenninger M (2018) The genomic footprint of climate adaptation in *Chironomus*
riparius. Mol Ecol 27:1439–1456
15. Burke MK, Dunham JP, Shahrestani P, Thornton KR, Rose MR, Long AD (2010) Genome-
wide analysis of a long-term evolution experiment with Drosophila. Nature 467:587–
590
16. Tenaillon O, Barrick JE, Ribeck N, et al (2016) Tempo and mode of genome evolution in
a 50,000-generation experiment. Nature 536:165–170

17.   Bersaglieri T, Sabeti PC, Patterson N, Vanderploeg T, Schaffner SF, Drake JA, Rhodes
680   M, Reich DE, Hirschhorn JN (2004) Genetic signatures of strong recent positive
681   selection at the lactase gene. Am J Hum Genet 74:1111–1120

682   18.   Roberts Kingman GA, Vyas DN, Jones FC, et al (2021) Predicting future from past: The
683   genomic basis of recurrent and rapid stickleback evolution. Sci Adv 7:5285–5303

684   19.   Abdul-Rahman F, Tranchina D, Gresham D (2021) Fluctuating environments maintain
685   genetic diversity through neutral fitness effects and balancing selection. Mol Biol Evol
686   38:4362–4375

687   20.   Kelly JK (2022) The genomic scale of fluctuating selection in a natural plant population.
688   Evol Lett 6:506–521

689   21.   Gossmann TI, Waxman D, Eyre-Walker A (2014) Fluctuating selection models and
690   Mcdonald-Kreitman type analyses. PLoS One.
691   https://doi.org/10.1371/journal.pone.0084540

692   22.   Rusuwa BB, Chung H, Allen SL, Frentiu FD, Chenoweth SF (2022) Natural variation at
693   a single gene generates sexual antagonism across fitness components in Drosophila.
694   Current Biology 32:3161-3169.e7

695   23.   Bertram J, Masel J (2019) Different mechanisms drive the maintenance of
696   polymorphism at loci subject to strong versus weak fluctuating selection. Evolution (N
697   Y) 73:883–896

698   24.   McAdam AG, Boutin S, Dantzer B, Lane JE (2019) Seed Masting Causes Fluctuations
699   in Optimum Litter Size and Lag Load in a Seed Predator. Am Nat 194:574–589

700   25.   Louthan AM, Kay KM (2011) Comparing the adaptive landscape across trait types:
701   larger QTL effect size in traits under biotic selection. BMC Evol Biol 11:60

702   26.   Behrman EL, Howick VM, Kapun M, Staubach F, Bergland AO, Petrov DA, Lazzaro BP,
703   Schmidt PS (2018) Rapid seasonal evolution in innate immunity of wild Drosophila
704   melanogaster. Proceedings of the Royal Society B: Biological Sciences.
705   https://doi.org/10.1098/rspb.2017.2599

706   27.   Machado HE, Bergland AO, Taylor R, et al (2021) Broad geographic sampling reveals
707   the shared basis and environmental correlates of seasonal adaptation in Drosophila.
708   Elife. https://doi.org/10.7554/eLife.67577

709   28.   Rudman SM, Greenblum SI, Rajpurohit S, Betancourt NJ, Hanna J, Tilk S, Yokoyama T,
710   Petrov DA, Schmidt P (2022) Direct observation of adaptive tracking on ecological time
711   scales in Drosophila. Science (1979). https://doi.org/10.1126/science.abj7484

712   29.   Hoekstra RF, Bijlsma R, Dolman AJ (1985) Polymorphism from environmental
713   heterogeneity: models are only robust if the heterozygote is close in fitness to the
714   favoured homozygote in each environment. Genet Res (Camb) 45:299–314

715   30.   Wittmann MJ, Bergland AO, Feldman MW, Schmidt PS, Petrov DA (2017) Seasonally
716   fluctuating selection can maintain polymorphism at many loci via segregation lift.
717   Proceedings of the National Academy of Sciences 114:E9932–E9941

718   31.   Glaser-Schmitt A, Wittmann MJ, Ramnarine TJS, Parsch J (2021) Sexual Antagonism,
719   Temporally Fluctuating Selection, and Variable Dominance Affect a Regulatory
720   Polymorphism in *Drosophila melanogaster*. Mol Biol Evol 38:4891–4907

721   32.   Gillespie J (1973) Polymorphism in random environments. Theor Popul Biol 4:193–195

722   33.   Gillespie JH (1977) Natural Selection for Variances in Offspring Numbers: A New
723   Evolutionary Principle. Source: The American Naturalist 111:1010–1014
724   34.   Roff DA (1997) Phenotypic Plasticity and Reaction Norms. Evolutionary Quantitative
725   Genetics 196–240
726   35.   Bürger R, Ghnelfarb A (1999) Genetic Variation Maintained in Multilocus Models of
727   Additive Quantitative Traits Under Stabilizing Selection. Genetics 152:807–820
728   36.   Ellner S, Hairston Jnr NG (1994) Role of overlapping generations in maintaining genetic
729   variation in a fluctuating environment. Am Nat 143:403–417
730   37.   Haller BC, Messer PW (2023) SLiM 4: Multispecies Eco-Evolutionary Modeling.
731   American Naturalist. https://doi.org/10.1086/723601
732   38.   Barton NH, Etheridge AM, Véber A (2017) The infinitesimal model: Definition,
733   derivation, and implications. Theor Popul Biol 118:50–73
734   39.   Dumont BL, Payseur BA (2008) Evolution of the gnomic rate of recombination in
735   mammals. Evolution (N Y) 62:276–294
736   40.   Álvarez-Castro JM, Carlborg Ö (2007) A unified model for functional and statistical
737   epistasis and its application in quantitative trait loci analysis. Genetics 176:1151–
738   1167
739   41.   Wientjes YCJ, Bijma P, Van Den Heuvel J, Zwaan BJ, Vitezica ZG, Calus MPL (2023) The
740   long-term effects of genomic selection: 2. Changes in allele frequencies of causal loci
741   and new mutations. Genetics. https://doi.org/10.1093/GENETICS/IYAD141
742   42.   Barton NH, Turelli M (2004) Effects of genetic drift on variance components under a
743   general model of epistasis. Evolution (N Y) 58:2111–2132
744   43.   Beniwal BK, Hastings IM, Thompson R, Hill WG (1992) Estimation of changes in genetic
745   parameters in selected lines of mice using REML with an animal model. 2. Body
746   weight, body composition and litter size. Heredity (Edinb) 69:361–371
747   44.   Bijma P (2011) A general definition of the heritable variation that determines the
748   potential of a population to respond to selection. Genetics 189:1347–1359
749   45.   Holstad A, Voje KL, Opedal ØH, Bolstad GH, Bourg S, Hansen TF, Pélabon C (2024)
750   Evolvability predicts macroevolution under fluctuating selection. Science (1979)
751   384:688–693
752   46.   Visscher PM, Hill WG, Wray NR (2008) Heritability in the genomics era — concepts and
753   misconceptions. Nat Rev Genet 9:255–266
754   47.   Kruuk LEB (2004) Estimating genetic parameters in natural populations using the
755   'animal model.' Philos Trans R Soc Lond B Biol Sci 359:873–890
756   48.   Cisneros-Mayoral S, Graña-Miraglia L, Pérez-Morales D, Peña-Miller R, Fuentes-
757   Hernández A (2022) Evolutionary History and Strength of Selection Determine the
758   Rate   of   Antibiotic   Resistance   Adaptation.   Mol   Biol   Evol.
759   https://doi.org/10.1093/molbev/msac185
760   49.   Phillips MA, Briar RK, Scaffo M, Zhou S, Sandoval-Powers M, Burke MK (2023) Strength
761   of selection potentiates distinct adaptive responses in an evolution experiment with
762   outcrossing yeast. bioRxiv 2022.05.19.492575
763   50.   Kardos M, Luikart G (2021) The genetic architecture of fitness drives population
764   viability during rapid environmental change. American Naturalist 197:511–525

51. Orr HA, Unckless RL (2008) Population extinction and the genetics of adaptation. American Naturalist 172:160–169

52. Yamamichi M (2022) How does genetic architecture affect eco-evolutionary dynamics? A theoretical perspective. Philosophical Transactions of the Royal Society B. https://doi.org/10.1098/RSTB.2020.0504

53. Stephan W (2019) Selective Sweeps. Genetics 211:5–13

54. Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C (2005) Genomic scans for selective sweeps using SNP data. Genome Res 15:1566–1575

55. Gregory TR (2009) Understanding Natural Selection: Essential Concepts and Common Misconceptions. Evolution: Education and Outreach 2:156–175

56. Oz T, Guvenek A, Yildiz S, et al (2014) Strength of Selection Pressure Is an Important Parameter Contributing to the Complexity of Antibiotic Resistance Evolution. Mol Biol Evol 31:2387–2401

57. Bjørnstad ON, Nelson WA, Tobin PC (2016) Developmental synchrony in multivoltine insects: generation separation versus smearing. Popul Ecol 58:479–491

58. Mendonça, Jr. M de S, Romanowski HP (2012) Population ecology of the multivoltine Neotropical gall midge Eugeniamyia dispar (Diptera, Cecidomyiidae). Iheringia Ser Zool 102:170–176

59. Harvey JA, Tougeron K, Gols R, et al (2023) Scientists' warning on climate change and insects. Ecol Monogr. https://doi.org/10.1002/ecm.1553

60. Subedi B, Poudel A, Aryal S (2023) The impact of climate change on insect pest biology and ecology: Implications for pest management strategies, crop production, and food security. J Agric Food Res 14:100733

61. Buckley LB, Arakaki AJ, Cannistra AF, Kharouba HM, Kingsolver JG (2017) Insect development, thermal plasticity and fitness implications in changing, seasonal environments. Integr Comp Biol 57:988–998

62. Gaytán Á, Gotthard K, Tack AJM (2022) Strong impact of temperature and resource specialisation on patterns of voltinism within an oak-associated insect community. Ecol Entomol 47:544–552

63. Allen AP, Gillooly JF, Savage VM, Brown JH (2006) Kinetic effects of temperature on rates of genetic divergence and speciation. Proc Natl Acad Sci U S A 103:9130–9135

64. Sampaio F, Krechemer FS, Marchioro CA (2021) The hotter the better? Climate change and voltinism of Spodoptera eridania estimated with different methods. J Therm Biol 98:102946

65. Sella G, Barton NH (2019) Thinking About the Evolution of Complex Traits in the Era of Genome-Wide Association Studies. Annu Rev Genomics Hum Genet 20:461–493

66. Hayward LK, Sella G (2022) Polygenic adaptation after a sudden change in environment. Elife. https://doi.org/10.7554/eLife.66697

67. Wickham H (2016) ggplot2. https://doi.org/10.1007/978-3-319-24277-4

68. Jenkins GM (1965) A Survey of Spectral Analysis. Appl Stat 14:2

69. R Core Team (2022) R: A language and environment for statistical computing. https://www.r-project.org/. Accessed 14 Feb 2025

70. Reznick DN, Ghalambor CK (2005) Selection in Nature: Experimental Manipulations of Natural Populations. Integr Comp Biol 45:456–462

71. Barton NH, Turelli M (1989) Evolutionary quantitative genetics: How little do we know?
Annu Rev Genet 23:337–370

72. Hill WG, Mackay TFC (2004) D. S. Falconer and Introduction to Quantitative Genetics.
Genetics 167:1529–1536

73. Nuzhdin S V, Turner TL (2013) Promises and limitations of hitchhiking mapping. Curr
Opin Genet Dev 23:694–699

74. Wilson BA, Petrov DA, Messer PW (2014) Soft Selective Sweeps in Complex
Demographic Scenarios. Genetics 198:669–684

75. Chevin LM (2019) Selective sweep at a QTL in a randomly fluctuating environment.
Genetics 213:987–1005

76. Wientjes YCJ, Bijma P, Calus MPL, Zwaan BJ, Vitezica ZG, van den Heuvel J (2022) The
long-term effects of genomic selection: 1. Response to selection, additive genetic
variance, and genetic architecture. Genetics Selection Evolution 54:19

77. Franssen SU, Kofler R, Schlötterer C (2017) Uncovering the genetic signature of
quantitative trait evolution with replicated time series data. Heredity (Edinb) 118:42–
51

78. Turner TL, Stewart AD, Fields AT, Rice WR, Tarone AM (2011) Population-Based
Resequencing of Experimentally Evolved Populations Reveals the Genetic Basis of
Body Size Variation in Drosophila melanogaster. PLoS Genet 7:e1001336

79. Kopp M, Hermisson J (2009) The Genetic Basis of Phenotypic Adaptation II: The
Distribution of Adaptive Substitutions in the Moving Optimum Model. Genetics
183:1453–1476

80. Lynch M, Wei W, Ye Z, Pfrender M (2024) The genome-wide signature of short-term
temporal selection. Proceedings of the National Academy of Sciences
121:e2307107121

81. Haldane JBS, Jayakar SD (1963) Polymorphism due to selection of varying direction. J
Genet 58:237–242

82. Bürger Reinhard, Gimelfarb A (2002) Fluctuating environments and the role of
mutation in maintaining quantitative genetic variation. Genet Res 80:31–46

83. Turelli M, Barton NH (2004) Polygenic Variation Maintained by Balancing Selection:
Pleiotropy, Sex-Dependent Allelic Effects and $G \times E$ Interactions. Genetics 166:1053–
1079

84. Lande R (2007) Expected relative fitness and the adaptive topography of fluctuating
selection. Evolution (N Y) 61:1835–1846

85. Lande R (2009) Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. J Evol Biol 22:1435–1446

86. Höllinger I, Wölfl B, Hermisson J (2023) A theory of oligogenic adaptation of a
quantitative trait. Genetics. https://doi.org/10.1093/genetics/iyad139

87. Bertram J (2021) Allele frequency divergence reveals ubiquitous influence of positive
selection in Drosophila. PLoS Genet 17:e1009833

88. Barghi N, Hermisson J, Schlötterer C (2020) Polygenic adaptation: a unifying
framework to understand positive selection. Nature Reviews Genetics 2020 21:12
21:769–781

89. Sibly RM, Curnow RN (2023) Allele frequencies and selection coefficients in locally
adapted populations. J Theor Biol 565:111463

90. Hedrick PW (1976) Genetic variation in a heterogeneous environment. II. Temporal
heterogeneity and directional selection. Genetics 84:145–157

91. de Vladar HP, Barton N (2014) Stability and Response of Polygenic Traits to Stabilizing
Selection and Mutation. Genetics 197:749–767

92. Chevin LM, Gompert Z, Nosil P (2022) Frequency dependence and the predictability
of evolution in a changing environment. Evol Lett 6:21–33

93. Cvijović I, Good BH, Jerison ER, Desai MM (2015) Fate of a mutation in a fluctuating
environment. Proceedings of the National Academy of Sciences 112:E5021–E5028

94. West R, Mobilia M (2020) Fixation properties of rock-paper-scissors games in
fluctuating populations. J Theor Biol 491:110135

95. Buffalo V, Coop G (2020) Estimating the genome-wide contribution of selection to
temporal allele frequency change. Proc Natl Acad Sci U S A 117:20672–20680

96. Phillips MA, Kutch IC, Long AD, Burke MK (2020) Increased time sampling in an evolve-
and-resequence experiment with outcrossing Saccharomyces cerevisiae reveals
multiple paths of adaptive change. Mol Ecol 29:4898–4912

97. Greener JG, Kandathil SM, Moffat L, Jones DT (2021) A guide to machine learning for
biologists. Nature Reviews Molecular Cell Biology 2021 23:1 23:40–55

98. Via S, Lande R (1985) Genotype-environment interaction and the evolution of
phenotypic plasticity. Evolution (N Y) 39:505–522

99. Lande R (2014) Evolution of phenotypic plasticity and environmental tolerance of a
labile quantitative character in a fluctuating environment. J Evol Biol 27:866–875

100. Oostra V, Saastamoinen M, Zwaan BJ, Wheat CW (2018) Strong phenotypic plasticity
limits potential for evolutionary responses to climate change. Nature
Communications 2018 9:1 9:1–11

101. Gloss AD, Whiteman NK (2016) Balancing Selection: Walking a Tightrope. Current
Biology 26:R73–R76

102. Barghi N, Tobler R, Nolte V, Jakšić AM, Mallard F, Otte KA, Dolezal M, Taus T, Kofler R,
Schlötterer C (2019) Genetic redundancy fuels polygenic adaptation in Drosophila.
PLoS Biol. https://doi.org/10.1371/journal.pbio.3000128

103. Agashe D, Sane M, Singhal S (2023) Revisiting the Role of Genetic Variation in
Adaptation. Am Nat 202:486–502