Genetic background influences MAFA^{S64F}-mediated diabetes penetrance in male mice

Zachary Loyd[1,*], Dongsoo Lee[1,*], Mallory Maurer[1], Lizabeth Buzzelli[1,2], Jin-Hua Liu[2], Garrett Reynolds[2], Mark P. Keller[3], Jean-Philippe Cartailler[2,4], Mark A. Magnuson[2,4], Roland Stein[2], Jeeyeon Cha[1,2,5,§]

[1]Division of Diabetes, Endocrinology, and Metabolism, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

[2]Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, USA

[3]Department of Biochemistry, University of Wisconsin-Madison, Madison, WI, USA

[4]Creative Data Solutions, Vanderbilt Center for Stem Cell Biology, Vanderbilt University, Nashville, TN, USA

[5]Tennessee Valley Veterans Affairs Health Care System, Geriatric Research Education and Clinical Center (GRECC) and Medicine Section, Nashville, TN, USA

*Co-first authors

§Corresponding author: jeeyeon.cha@vumc.org

Keywords: Islet, beta cell, MAFA, MODY diabetes

ABSTRACT

Pancreatic β-cells require the coordinated expression of transcription factors such as MAFA to dynamically secrete insulin to maintain euglycemia. A naturally occurring mutation in MAFA (MAFA[S64F]) produces a long-lived variant protein which predisposes carriers to dichotomous conditions of either maturity (adult)-onset diabetes of the young (MODY) or hypoglycemia in a sex-dependent manner. Here we show that genetic background modulates disease penetrance in MafA[S64F/+] male mice. Specifically, MafA[S64F/+] males backcrossed to a C57/Bl6J ("C57") background prevented dysglycemia, while C57 MafA[S64F/+] males bred one generation to an SJL/J background ("mixed") manifested overt diabetes, impaired insulin secretion, and accelerated β-cell senescence. MafA protein levels, phosphorylation status, and target gene expression in C57 MafA[S64F/+] male islets were more comparable to wildtype males. RNA sequencing of C57 MafA[S64F/+] male islets revealed fewer differentially expressed genes than mixed background male islets, including transcriptional signatures of β-cell senescence. In addition, retinoic acid signaling, another signature of cellular aging, was uniquely downregulated in C57 MafA[S64F/+] male islets. Indeed, core retinoic acid signaling receptor RARα and known target genes were downregulated in C57 MafA[S64F/+] male islets. CUT&RUN mapping revealed that MafA directly impacted retinoic acid receptor alpha *Rara* expression. In sum, these data show that genetic factors can impact unique pathways to modulate disease penetrance in mice modeling MAFA-MODY.

Article Highlights:

- Genetic background in mice or ancestry in humans can profoundly influence diabetes susceptibility.

- We asked whether diabetes penetrance in our robust mouse model of MAFA-MODY on a mixed genetic background was impacted when backcrossed to C57/Bl6J ("C57").

- Here we show that C57 MafA$^{S64F/+}$ males have wildtype glucose-sensing properties, compared to overt diabetes which could be reproducibly re-created on a mixed background. Further analysis disclosed more muted gene expression changes in C57 MafA$^{S64F/+}$ male islets including unique downregulation of senescence and retinoic acid signaling.

- These results illustrate the influence of genetic landscape on β-cell function in response to a pathogenic MODY variant.

INTRODUCTION

MafA, the large V-Maf avian <u>m</u>usculo<u>a</u>poneurotic <u>f</u>ibrosarcoma transcription factor (TF), is a pancreatic islet β-cell-enriched protein that is essential for activating transcriptional programs promoting maturation (*1, 2*). Compromised levels of mouse MafA (*3-8*) or human MAFA (*9, 10*) also diminishes β-cell function. For example, genetic knockout models of pancreatic *MafA* manifest impaired glucose tolerance in male and female mice (*8*), while human *MAFA* expression is reduced in islets recovered from male and female donors with diabetes (*9*). Conversely, inducing *MafA* expression in MafA^{Low}, non-glucose responsive neonatal rat islets increases glucose-stimulated insulin secretion (GSIS) (*11*), while acute introduction of human *MAFA* alone in mouse and human stem cells directly promotes insulin production and β-cell maturity markers (*11-13*). Collectively, these results demonstrate a fundamental role for MafA in β-cell function and identity.

Pathogenic mutations in pancreatic islet-enriched TFs including MAFA can cause heritable, monogenic forms of islet dysfunction in adults (*14*). A *MAFA* missense mutation (p.Ser64Phe, c.191C>T) in male carriers develop sex-dependent diabetes, while female carriers develop persistent hyperinsulinemic hypoglycemia due to the production of non-syndromic, insulin-secreting neuroendocrine tumors (insulinomatosis) (*15*). The resultant MAFA^{Ser64Phe} (termed MAFA^{S64F}) variant protein impairs protein phosphorylation, which impacts both transcriptional activity and protein stability (i.e., t$_{1/2}$ of MAFA is normally only ~30 minutes, compared to many hours in MAFA^{S64F}) (*15*). A mouse model harboring this clinically relevant mutation in the endogenous *MafA* gene (termed MafA$^{S64F/+}$) on a mixed genetic background (C57/Bl6J and SJL/J) mimicked aspects of the heterogeneous, sex-dependent phenotypes: female MafA$^{S64F/+}$ mice were persistently hypoglycemic, while male MafA$^{S64F/+}$ mice showed impaired glucose tolerance due to widespread, premature β-cell aging and senescence (*16*).

Introduction of this protein in a male human β-cell line also drove accelerated cellular aging (*16*), implying that the regulatory mechanisms are conserved between mice and humans.

Genetic background in mice (*17-20*) or ancestry in humans (*21-23*) can profoundly influence diabetes susceptibility. Common genetic variants have been identified to modify disease risk and clinical presentation in monogenic diseases including diabetes (i.e., MODY) (*24-27*). These observations prompted our current study to determine if MafA[S64F/+] diabetes penetrance was impacted when backcrossed to the C57/Bl6J ("C57") background. C57 was chosen since these mice are susceptible to glucose intolerance under conditions of stress (*17-20*). Here we show that C57 MafA[S64F/+] males have essentially wildtype (WT) glucose-sensing properties by 6 weeks of age, compared to overt diabetes which could be re-created on a mixed background. Further analysis of C57 and mixed MafA[S64F/+] male islets disclosed more muted gene expression changes in the C57 background as well as downregulation of β-cell senescence and retinoic acid signaling pathways associated with aging. These results illustrate the influence of genetic landscape on β-cell function in response to a pathogenic MODY variant.

METHODS

Mice

MafA$^{S64F/+}$ heterozygous mice were originally generated using CRISPR/Cas9 targeting by the University of Michigan Transgenic Core on a mixed SJL/J and C57/Bl6J mouse strain (*16*). Mice were backcrossed eight generations with C57/Bl6J mouse strain (Jackson Lab) to generate C57 MafA$^{S64F/+}$ mice (Transnetyx Genetic Monitoring, >95% C57Bl/6J). These C57 MafA$^{S64F/+}$ mice were bred one time with SJL/J mice (Taconic) to produce the mixed genetic background strain (F1 generation, roughly 50:50 contribution of SJL/J and C57). Mice of both sexes were used in this study. All animal studies were reviewed and approved by the Vanderbilt University Institutional Animal Care and Use Committee. Mice were housed and cared for according to the Vanderbilt University Department of Animal Care and the Institutional Animal Care and Use Committee of Animal Welfare Assurance Standards and Guidelines.

Intraperitoneal glucose tolerance testing and serum hormone measurements

Glucose tolerance testing was performed on male and female MafA^{WT} and MafA$^{S64F/+}$ littermate mice (n = 6-16) after afternoon intraperitoneal injection of D-glucose (2 mg/g body weight) prepared in sterile PBS (20% w/v) preceded by a 5-hour fast (i.e. 8AM to 1PM). Afternoon insulin tolerance tests were conducted by intraperitoneal injection of 0.5 IU/kg body weight insulin (Novolin, regular human insulin, recombinant DNA origin) into mice (n = 3-5) fasted for 6 hours. Blood glucose was measured using a FreeStyle glucometer (Abbott Diabetes Care) before (0 minutes) and at 15, 30, 60, and 90 minutes following injection. Serum insulin and C-peptide was measured by Ultrasensitive Insulin ELISA (Mercodia) and C-peptide ELISA (Alpco), respectively.

Mouse islet isolation, RNA isolation, and gene expression analysis

Mouse islets were isolated using collagenase P (Sigma) injected into the pancreatic duct, followed by a Histopaque 1077 (Sigma-Aldrich) gradient. Handpicked islets were isolated in standard RPMI-1640 medium (Thermo Fisher Scientific) supplemented with 10% FBS, L-glutamine, and penicillin-streptomycin, then stored frozen. Islet RNA was isolated using the RNAqueous Micro Total RNA Isolation Kit (Invitrogen). cDNA was generated using Superscript III reverse transcriptase (Invitrogen) by the oligo(dT) priming method (*16*). Real-time PCR assays were performed using the LightCycler FastStart DNA Master PLUS SYBR Green kit (Roche) and a LightCycler PCR instrument (Roche). The real-time PCR primers are listed in **Supplemental Table 1**. Gapdh or β-Actin was used to normalize the real-time PCR data which were analyzed using the ΔΔCt method (*8*).

Cell culture and RNA isolation

Monolayer cultures of MIN6 mouse β-cells and HeLa cells were grown under conditions described previously (*4*). Transfection with scrambled control, MAFA^{WT} or MAFA^{S64F} constructs of confirmed sequences (*28*) was performed using Lipofectamine 2000 (Invitrogen) (average transfection efficiency 80%) before protein or RNA isolation. RNA was isolated 4 days posttransfection using the Trizol reagent (Fisher Scientific), while protein was isolated in RIPA buffer (*8*). cDNA synthesis and real-time PCR assays were performed as above (*16*).

Western blotting and gel shift assay

MAFA protein levels were normalized to endogenous β-actin by immunoblotting with antibodies against MAFA (Cell Signaling Technology, 79737) or β-actin (MilliporeSigma, MAB1501). Horseradish peroxidase–conjugated anti-rabbit (31460, Thermo Fisher Scientific) or anti-goat (31402, Thermo Fisher Scientific) secondary antibodies were used at 1:5000 dilution. Immunoblots were quantified with ImageJ (NIH). Gel shift assays were performed using 10 µg of nuclear protein extract and 200 fmol of biotin-labeled double-stranded human *INS* MAFA

binding site probe alone or mixed with WT unlabeled competitor DNA, mutant unlabeled

competitor DNA, or MAFA antibody (Bethyl Laboratories, Product # A700–067) in a total of

20 µL reactions (Lightshift Chemiluminescent EMSA Kit, Thermo Fisher Scientific) containing 1X

binding buffer, 2.5% glycerol, 5 mM MgCl, 50 ng/µL poly(dI-dC) (Thermo Fisher Scientific), and

0.05% NP-40 (Sigma-Aldrich). These reactions were electrophoresed through a 6% precast

DNA retardation gel in 0.5% Tris-borate-EDTA buffer (TBE, Thermo Fisher Scientific) at 100 V

for 1.5 h.

Immunohistochemical analyses

Mouse pancreata were fixed overnight at 4°C in 4% paraformaldehyde (Electron Microscopy

Services) in PBS, and pancreata were embedded in either Tissue-Plus OCT (Thermo Fisher

Scientific) or paraffin wax. Sections of rodent pancreata were cut at 6µm thickness. The paraffin

sections were deparaffinized and rehydrated before citrate buffer–based antigen retrieval.

Sections were then made permeable by 0.5% Triton in PBS treatment for 10 minutes. Following

blocking with 0.5% BSA in PBS for 120 minutes, the primary antibodies were applied overnight

at 4°C. Primary antibodies are listed in **Supplemental Table 1**. Species-matched antibodies

conjugated with the Cy2, Cy3, or Cy5 fluorophores were used for secondary detection (1:1000;

Jackson ImmunoResearch). DAPI was used for nuclear staining (Southern Biotech).

Immunofluorescence images were obtained using the Zeiss Axio Imager M2 widefield

microscope with ApoTome.

Senescence-associated β-galactosidase (SA-β-gal) staining

Pancreata were snap frozen in O.C.T. and cryosections were prepared at 16µm thickness. SA-

β-gal activity staining was performed at pH 6.0 (*16*) (Cell Signaling). To compare the intensity of

SA-β-gal staining, sections from different genotypes were processed on the same slide. Staining

reactions were developed for 18 hours at 37°C, then quenched by 3x PBS washes (pH 7.4).

These slides were subject to immunostaining for insulin by fixing in 4% paraformaldehyde for 45

minutes, permeabilized with Tris-buffered saline with 0.2% Triton in PBS for 15 minutes,

blocked in 2% normal donkey serum/1% BSA in PBS and incubated overnight with guinea pig

anti-insulin (1:500, Fitzgerald 20-IP35) at 4°C. HRP-conjugated secondary antibodies were

incubated on slides for 1 hour and detected with the DAB+ chromogen kit (DAKO). After

washing, slides were mounted and imaged by brightfield microscopy.

RNAscope and quantification

The RNAscope® Multiplex Fluorescent Reagent Kit V2 (ACD) was used following

manufacturer's instructions and modified for mouse pancreata as previously described (*29*).

RNAscope images from 6 week old mice were generated using a Leica DMi8 inverted

microscope at 40X objective and analyzed using QuPath (*30*). Regions of interest (ROI) around

islets were defined based on insulin-positive staining. Stardist (*31*) , a deep-learning-based

nuclei detection model, was used to detect nuclei and outline the cell. Max entropy or triangle

auto-threshold methods and image down-sampling (factor of 2) was applied to determine the

threshold for each channel with an RNA probe. RNA speckles were detected using the

subcellular detection feature in QuPath with the calculated thresholds. Percentage of β-cells

positive for *Cdkn1a* was calculated in QuPath and exported to Prism for statistical analysis.

Bulk RNA-sequencing analysis of mouse islet cells

Bulk RNA-sequencing was performed on mouse islets isolated at 6 weeks of age. RNAqueous

Micro Total RNA Isolation Kit (Invitrogen) was used to isolate total RNA, and RNA quality was

analyzed on an Agilent 2100 Bioanalyzer. Samples with RIN >8 were used for sequencing. The

cDNA libraries were constructed, and paired-end sequencing was performed on an Illumina

NovaSeq6000 (150-nucleotide reads). The generated FASTQ files were processed and

interpreted using the Genialis visual informatics platform (https://www.genialis.com) (*16*).

DESeq2 was used for differential gene expression analyses and statistical comparison (*32*).

Poorly expressed genes (i.e., expression count summed over all samples below 10) were

filtered out from the differential expression analysis input matrix. Sequencing data is available in

the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO)

database (accession no. GSE299343).

Position Weight Matrix (PWM) Motif Search Protocol

Putative MAFA responsive elements (MAREs) were identified in the *Rara* gene promoter and

other upstream regulatory regions by scanning genomic sequences for Position Weight Matrix

(PWM) motifs. The MAFA PWM was obtained from the JASPAR database (matrix ID

MA1521.2), representing the 13 bp core MARE consensus derived from ChIP-seq binding data

(*33*). The position frequency matrix was converted to a log-odds PWM with pseudocounts of 0.8

using TFBSTools (*34*). Genomic sequences were scanned on both strands using

TFBSTools::searchSeq(), and matches scoring at or above 80% of the maximum possible PWM

score were retained as candidate binding sites. The transcription start site (TSS) was defined as

the 5'-most position of a representative transcript from the UCSC Gene annotation (mm39

assembly); one TSS per gene symbol was used. Three non-overlapping search regions were

defined relative to each TSS: proximal promoter (TSS -200bp to +1000bp), distal promoter (TSS

+1kb to +5kb), and enhancer (TSS +5kb to +100kb). PWM relative scores were tiered as

follows: 80-85% (weak/marginal candidate), 85-90% (moderately strong candidate), and 90%+

(strong candidate with high confidence match to consensus sequence).

CUT&RUN assay and analysis

CUT&RUN was performed on 500,000 dispersed mouse MIN6 cells per condition using the

CUTANA ChIC/CUT&RUN protocol v3.1 (Epicypher). Nuclei were extracted with nuclear

extraction buffer (20 mM HEPES-KOH [pH 7.9]; 10 mM KCl; 0.1% Triton X-100; 20% glycerol; 1 mM MnCl2; 0.5 mM spermidine; 1uM protease inhibitor; Thermo Fisher Scientific) for 10 minutes on ice and immobilized onto Concanavalin-A beads (EpiCypher). After blocking and washes, samples were incubated with 0.5 µg of rabbit anti-MAFA (Cell Signaling 79737) or rabbit anti-IgG (EpiCypher 13-0042) antibodies overnight at 4°C. pAG-MNase (EpiCypher) was added to nuclei (1:20) and incubated at room temperature for 10 minutes. Targeted chromatin digestion was induced by adding 100 mM $CaCl_2$ and nutating for 2 hours at 4°C. DNA fragments were purified using the CUTANA ChIC/CUT&RUN kit. DNA was resuspended in 0.1M~ Tris-EDTA buffer solution and used for library preparation with the CUTANA CUT&RUN Library Prep Kit (EpiCypher, 14-1001), according to the version 1 (v1) manual. Libraries were sequenced as PE150 reads on the NovaSeq platform. All libraries had > 15 million reads and were processed using the nf-core/cutandrun workflow, v3.1 (*35*). The pipeline performs adapter trimming, alignment to the GRCm38 reference genome, filtering against the mm10 blacklist regions, spike-in normalization, and peak calling using MACS2. The workflow also generates IGV sessions and all track data, which allows for interactive visualization and exploration. Quality was confirmed by fragment length distribution and fingerprint plots. Nearest gene analysis was performed with peakScout (*36*). Coverage plots were created using ggcoverage (*37*). Sequencing data were deposited in the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) database (accession no. GSE298664).

Statistical Analysis

Data are expressed as the mean ± SEM. Statistical analysis was performed using GraphPad Prism 9.5.0 (GraphPad Software Inc.). The differences between groups were analyzed by unpaired 2-tailed Student's t test or 2-way ANOVA, as indicated. Differences were considered to be statistically significant at $P < 0.05$.

RESULTS

Glucose tolerance is largely unaffected in C57 MafA$^{S64F/+}$ males at 6 weeks of age, while those on a mixed background manifest overt diabetes and impaired insulin secretion.

As seen clinically, MafA$^{S64F/+}$ mice on a mixed genetic background (C57/Bl6J and SJL/J) were previously found to have distinct metabolic phenotypes (*16*). Here we find that impaired glucose tolerance of MafA$^{S64F/+}$ males was essentially eliminated by 6 weeks of age when backcrossed eight generations to the C57/Bl6J (C57) genetic background (**Fig. 1A**, C57). In contrast, the F1 generation of C57 MafA$^{S64F/+}$ males bred once with parental SJL/J mice were both glucose intolerant and hyperglycemic (**Fig. 1B**, mixed) due to impaired insulin and C-peptide secretion (**Fig. 1C-D**). The newly generated F1 mixed background mice were glucose intolerant, like the earlier MafA$^{S64F/+}$ males (**Fig. 1B**) (*16*) and showed transient hypoglycemia preceding impaired glucose tolerance at 5 weeks of age even on an extended wean protocol (**Supp Fig. 1A-B**). Moreover, this mixed MafA$^{S64F/+}$ mouse line had comparable weight gain as well as intact peripheral insulin tolerance in relation to littermate wild type (WT) mice (**Supp Fig. 1C-E**), all of which were properties of the previously reported mixed line (*16*). In contrast, MafA$^{S64F/+}$ females on both mixed and C57 backgrounds were hypoglycemic as reported previously (*16*). Consequently, we principally focused our efforts on determining how the C57 background prevented glucose intolerance in MafA$^{S64F/+}$ males.

MafA^{WT} protein status and target gene expression is augmented in C57 MafA$^{S64F/+}$ male islets compared to mixed background males.

MafA function is dependent upon posttranslational phosphorylation events which couple two distinct regulatory processes: increased transactivation activity *and* ubiquitin-mediated

degradation (*38*). *In vitro* analysis of MafA[S64F] found that impaired phosphorylation at neighboring Ser65 (S65) prevented the ability of glycogen synthase kinase 3 (GSK3) to phosphorylate S61, T57, T53, and S49, which results in a protein with faster SDS-PAGE mobility, greater stability, and altered transactivation capacity (*38*). Here we show that the MafA[WT] protein was visible in C57, but not mixed, background MafA[S64F/+] male islets (**Fig. 2A**). Interestingly, the quantity of MafA[S64F] protein did not appear to differ by genetic background (**Fig. 2B**). The expression of MafA and known MafA target genes were either less impacted (i.e., *Ins1*, *Ins2*) or unaffected (i.e., *Gck, Slc2a3*, *Slc30a8, MafA*) in C57 MafA[S64F/+] male islets (**Fig. 2C**), presumably reflecting MafA[WT] protein activity in this background. The relatively low level of total islet MafA (WT+S64F) protein in relation to WT mice reflects the ability of MafA[S64F] to reduce endogenous mRNA production (**Fig. 2C**) (*16*). Along these lines, low MafA protein levels were also evident in human MAFA[S64F] carriers (*15*).

Unbiased RNA sequencing illustrates how differently MafA[S64F] regulates gene expression in C57 and mixed MafA[S64F/+] male islets.

Bulk RNA sequencing was performed on MafA[WT] and MafA[S64F/+] male islets at 6 weeks of age to assess dominant gene expression patterns between these groups. Notably, the 6 week time point in MafA[S64F/+] males represents when distinct differences in male phenotypes between backgrounds is observed (**Fig. 1A-B**). Male C57 MafA[S64F/+] islets had 2,097 differentially expressed genes (DEGs, 616 up and 1481 down) compared to male WT islets, while a greater number of DEGs were found on the mixed background MafA[S64F/+] mice (3,366: 1505 up and 1861 down) (**Fig. 3A**). Of these, only 386 upregulated and 621 downregulated DEGs were similarly impacted between the genetic backgrounds. Unbiased clustering identified that the molecular signatures between MafA[WT] and MafA[S64F/+] males on a C57 background were more similar than those on a mixed background (**Fig. 3B**). In addition, GO pathway analysis showed

that the up- and down-regulated DEGs separated into distinct GO pathways in C57 and mixed MafA[S64F/+] male islets (**Fig. 3C**).

Senescence gene signatures are abrogated in male C57 MafA[S64F/+] islets.

GO analysis and STRING protein association network analysis (*39*) of the DEGs of MafA[S64F/+] male islets defined a distinct set of genetic background-specific pathways (**Fig. 3C**). As expected, cellular aging pathways were found in mixed MafA[S64F/+] male islets, which represent earlier reported signatures (*16*). However, these pathways were attenuated in the C57 MafA[S64F/+] males (e.g., see pancreatic secretion, protein digestion and absorption, ECM receptor interactions, cell adhesion, and calcium signaling pathways). In line with this, senescence gene signatures enriched in the mixed background MafA[S64F/+] male islets (*16*) were largely muted in the male C57 MafA[S64F/+] islets (**Fig. 3D-F**). As previously reported (*16*), islet senescence signatures were not seen in female MafA[S64F/+] islets (**Supp Fig. 2**). Furthermore, hallmark molecular changes associated with cellular senescence such as enrichment of senescence associated-β-galactosidase (SA-β-gal) staining, DNA damage response such as 53BP1, and impaired nuclear integrity such as by downregulation of LaminB1, found in mixed background MafA[S64F/+] males, were not detected or unchanged in C57 MafA[S64F/+] male islets (**Figs. 3E-F, 4A-B**). Collectively, these results revealed that glucose intolerance (**Fig. 1**) and accelerated islet aging was prevented in C57 MafA[S64F/+] male islets.

C57 MafA[S64F/+] male islets show unique downregulation of core retinoic acid signaling components.

The activity of mixed male MafA[S64F/+] islets is presumably mediated by unique GO pathways associated with cellular aging and MODY enriched in this sex and background (**Fig. 3C**).

Circadian rhythm was another downregulated pathway in the mixed background males. Indeed, oscillatory insulin secretion is synchronized with nutritional intake (*40-43*), thus the circadian dysregulation in β-cells observed in this context could be additive to accelerated aging to impair dynamic insulin secretion and promote dysglycemia (*42-44*). In addition, downregulation of nuclear receptors and retinol metabolism genes was observed by GO analysis in C57 MafA[S64F/+] islets (**Fig. 3C**). In line with this, persistent retinoic acid signaling through retinoic receptors RAR or RXR drives cell cycle arrest and senescence (*45-47*), and exogenous retinoic acid induces senescence programming via upregulation of p21 and p16 cell cycle inhibitors (*48-50*).

Importantly, the core retinoic acid signaling components *Retinoic acid receptor (Rar) a, Rarb*, and *Retinol binding protein (Rbp) 7* were only downregulated in C57 MafA[S64F/+] male islets (**Fig. 5A**)*. MafA CUT&RUN analysis was performed in mouse Min6 β-cells to determine whether retinoic acid signaling components could be gene targets of endogenous MafA. We expect that this will also reveal MafA[S64F] binding in β-cells, as this mutation resides in the transactivation, and not DNA binding, domain. Gel shift assay showed that MafA[WT] and MafA[S64F] have similar *cis*-element binding ability when transfected in HeLa cell nuclear extracts (**Supp Fig. 3**) (*8*).

Gene-annotated MafA CUT&RUN peaks (n=33,013 peaks) were overlaid with the DEGs uniquely identified in MafA[S64F/+] males of each genetic background (**Fig. 5B**). Mixed background MafA[S64F/+] male islets yielded 1210 putative targets (**Fig. 5B**). Importantly, these included genes associated with β-cell identity and islet hormones such as urocortin 3 (Ucn3), glucagon (Gcg), somatostatin (Sst), cholecystokinin (Cck), insulin (Ins), and gastrin (Gast); islet-enriched transcription factors and receptors such as Glis1, Glp1r, Nkx6.2, Nkx6.3, and Smarca1; and circadian regulators Period1/2 (Per1/2), Crytochrome2 (Cry2), RAR-related orphan receptor α (Rora), D-site albumin promotor binding protein (Dbp), basic helix-loop-helix family member (Bhlhe40/Dec1), and nuclear receptor subfamily 1 group D member 1 (Rev-erba and Rev-erbb), which sets up multiple highly effective feedback loops that promotes dynamic insulin secretion

(*18, 37, 38, 51, 56-59*). Collectively, these results suggest several avenues of dysregulation in mixed background MafA$^{S64F/+}$ male islets.

C57 MafA$^{S64F/+}$ male islets yielded 250 unique putative targets of MafA (**Fig. 5B**). Notably, these targets included factors in the senescence-associated prosurvival Bcl2 family (ie., Bcl2l14) and retinoic acid receptor (RARα and RARβ) signaling. Consistent with these findings, *Rara* houses a moderately strong candidate MARE (Maf responsive binding element) in its distal (+1000 to +5000bp) promoter region. Immunostaining also revealed downregulation of RARα protein in C57 MafA$^{S64F/+}$ male islets (**Fig. 5C**), and gene expression studies showed downward trends of classical RARα signaling targets *Rbp1* and *Cyp26a1* (**Fig. 5A**). Furthermore, a significant reduction in RARα target *Cdkn1a* (encoding p21) was seen in C57 MafA$^{S64F/+}$ islet β-cells compared to mixed background MafA$^{S64F/+}$ β-cells (**Fig. 5D**). Thus, we conclude that MafA can directly regulate *Rara* expression in C57 MafA$^{S64F/+}$ β-cells to attenuate *Cdkn1a* induction and β-cell senescence in MafA$^{S64F/+}$ islets.

DISCUSSION

Genetic ancestry is widely accepted to influence disease penetrance (*21-23*). Even in purported monogenic disease, there is increasing recognition that genetic factors can play modulating roles beyond the causative mutation (*25-27*), and a recent study has shown that common genetic variants can influence monogenic HNF-MODY diabetes risk to range from 11% to 81% (*24*). Genetic background influences can more easily be mechanistically addressed in monogenic diabetes than in polygenic type 1 (T1D) and type 2 (T2D) diabetes. As a result, here we investigated the influence that genetic background could have on the MAFA-MODY variant and found a profound impact on the male diabetic phenotype in mice.

Patient carriers of the MAFA^{S64F} missense mutation (*15*) manifest sex-dependent glycemic disorders: men show adult-onset diabetes while women primarily manifest

hypoglycemia with non-syndromic insulinomatosis (*15*). Since this original report, a second

human MAFA missense mutation in the transactivation domain (i.e., MAFA^{T57R}) has been

identified to block the actions of GSK3 and, as expected, promotes MAFA protein stability and

cosegregates with insulinomatosis and MODY-like diabetes (*51*). A mouse model harboring the

S64F mutation in the endogenous *Mafa* gene, which prevented phosphorylation of the priming

kinase at MAFA^{S65} and prevents GSK3 actions, recapitulated the sex-dependent responses

seen in patients (*16*), providing mechanistic insights into the impact of this MafA variant *in vivo*,

including a male sex-dependent acceleration of islet senescence that impaired β-cell function. In

fact, accelerated cellular aging is a common feature of islets from T1D and T2D donors, and is

seen in mouse models of diabetes such as those with insulin resistance modeling T2D (*52*),

NOD mice modeling T1D (*53*), and MafA$^{S64F/+}$ males modeling MAFA-MODY (*16*).

Given increasing appreciation for genetic background effects in diabetes penetrance,

efforts to backcross MafA$^{S64F/+}$ mice on the C57 background were examined. Here we show that

the C57 genetic background interacts with this MafA mutation to influence the penetrance of the

male dysfunction by: 1) preventing glucose intolerance and β-cell senescence; 2) allowing

increased MafA^{WT} protein production, 3) yielding gene expression pattern similar to MafA^{WT}

islets; and 4) downregulating retinoic acid signaling, an important mediator of β-cell maturity.

Spatiotemporally regulated retinoic acid signaling is a requisite for cellular differentiation

of many cell types including islet β-cells. Specifically, a necessary role of retinoic acid signaling

has been found in β-cell differentiation and function in mouse and human islets (*54, 55*), and it is

also a critical component for stem-cell derived maturation of β-like cells for cell-based therapies

(*56, 57*). However, persistent retinoic acid signaling and exogenous retinoic acid can activate

cell cycle arrest and senescence programs (*45, 46*) via core cell cycle inhibitors (e.g., p21)

across cell types (*48-50*). In fact, excess signaling through endogenous pancreas retinoids are

reported to contribute to glucose intolerance (*58*). Here we show that downregulation of this

accessory maturity pathway in C57 MafA$^{S64F/+}$ mice appears to dampen the expression of the common aging driver p21.

Conversely, in the mixed background, we uncovered an impact of MafA^{S64F} on islet circadian regulators, identifying a novel link between β-cell maturity driver MafA, cellular aging, and circadian entrainment. Interestingly, we find that male MafA$^{S64F/+}$ islets on the mixed background show marked downregulation of not only direct circadian regulators (e.g. *Per1*, *Per2*, *Cry2)* but also amplification factors in complementary circadian circuits such as *Dbp, Rora, Bhlhe40/Dec1, Rev-erb* and *Nfil3* (*42-44*). Interestingly, the expression of core circadian activators Bmal and Clock was unchanged between genotypes and genetic background, affirming that the genetic-background related circadian dysregulation engaged regulators rather than the core Clock:Bmal1 components themselves. A recent study in human stem-cell derived β-cells deficient in *Bhlhe40/Dec1* showed impaired MafA periodicity, suggesting a novel circadian islet feedback loop (*59*), but confirmation of this relationship in *MafA$^{S64F/+}$* islets requires further study. Notably, several senescence mediators such as *Cdkn1a*, *Bcl2a1d*, *Ankrd1*, or *Icam1* enriched in mixed background MafA^{S64F} male islets, were not identified as MafA targets by CUT&RUN, suggesting that their regulation is through the influence of MafA^{S64F} on intermediate pathways such as circadian regulators or retinoic acid signaling.

MafB is another islet-enriched Maf family member with distinct temporal and cell-type expression from MafA (*4*). In mouse islets, MafB is the dominant family member expressed developmentally before MafA is expressed postnatally, while human MAFB is postnatally expressed alongside MAFA in β-cells. The analogous mutation in human MAFB (MAFB^{Ser70Ala}, or MAFB^{S70A}) in the transactivation domain also impairs phosphorylation and increases protein stability (*28*). Notably, it is one of several mutations which drives the clinical syndrome Multicentric Carpotarsal Osteolysis characterized by bony deformities and renal failure (*60*). The clinical impact of MAFB^{S70A} reflect the broader expression pattern of MAFB compared to MAFA (*1*). While the effect of this mutation on islet cell function is unclear, MAFB is essential for the

development of human embryonic stem cell derived β-like Insulin$^+$ cells (*61*), thus future studies comparing MAFB^{S70A} to MAFA^{S64F} would provide insights into its pathogenicity to adult β-cells.

A crucial next step is to investigate MAFA^{S64F} or MAFB^{S70A} in genetically modified human islets. However, our results raise the possibility that studies in male human islets may not show consistent phenotypes due to donor genetic background. In such a case, further investigations on the male phenotype may require models controlling for genetic background, such as stem cell derived β-cells, or a return to mouse models such as ours to investigate genetic background influences. In mice, genome-wide differential single nucleotide polymorphism (SNP) analysis between inbred genetic backgrounds can be pursued, and prioritization of high impact variants in genes involved in retinoic acid signaling, cellular aging, and circadian dysregulation would be interesting. Furthermore, identification of SNPs in genes which can impact MafA turnover such as kinases, ubiquitination mediators or proteasome components would also be interesting given the relative integrity of MafA^{WT} protein in C57 MafA$^{S64F/+}$ mice. An unbiased forward genetic screen can mechanistically delineate specific genetic loci which can contribute to the dramatic differences in MafA expression and activity in MafA$^{S64F/+}$ males between genetic backgrounds.

Nevertheless, our work here demonstrates the combinatorial impact of genetic background and biologic sex on molecular and functional β-cell responses to produce heterogenous responses (i.e., cellular senescence driving dysfunction vs. hypoglycemia) to a clinically relevant, stable variant of MafA. Our data strongly suggest that additional genetic regulators can modulate cellular responses to a pathogenic mutation and ultimately will improve strategies for diabetes risk stratification and therapies to optimize β-cell function in people at risk of diabetes.

Acknowledgements. This research was performed using resources and/or funding provided in parts by NIH grants to JC (K08 DK132507); RS (R01 DK090570); Alan Attie and MPK (R01DK101573, R01DK102948, and RC2DK125961); and the Vanderbilt Diabetes Research and Training Center (DK20593). JC was also supported by a Doris Duke Physician Scientist Fellowship, Burroughs Wellcome Fund Career Award for Medical Scientists, and generous start-up funding from Vanderbilt University Medical Center. MPK is also supported by the University of Wisconsin-Madison Department of Biochemistry and Office of the Vice Chancellor for Research and Graduate Education with funding from the Wisconsin Alumni Research Foundation. The authors thank former and current members of the Cha lab for feedback on this manuscript.

Guarantor Data Access and Responsibility Statement. JC is the guarantor of this work, and as such, had full access to all data in the study and is responsible for the integrity of the data and accuracy of data analysis.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions.

ZL, DL, MAM, RS, and JC designed the study.

ZL, DL, MM, LB, JL, GR, JPC, and JC performed experiments.

ZL, DL, MM, MPK, JPC, and JC performed bioinformatic analysis.

ZL, DL, MM, LB, JL, GR, MPK, JPC, MAM, RS, and JC analyzed data.

ZL, DL, JPC, RS and JC wrote the manuscript.

ZL and DL are co-first authors due to their contributions in most of the experimental studies.

MAIN FIGURE LEGENDS

Figure 1. Glucose tolerance is largely unaffected in C57 MafA$^{S64F/+}$ males, while those on a mixed background manifest frank diabetes and impaired insulin secretion

A. Fasted male animals underwent intraperitoneal glucose tolerance tests (GTT) at time points indicated. Male MafA$^{S64F/+}$ heterozygous (Het) mice on a C57 background had no changes to glucose tolerance through adulthood compared to MafA$^{+/+}$ (WT) male littermates.

B. Het males on a mixed background had impaired glucose tolerance by 5 weeks of age which then progressed to overt diabetes by 7 weeks of age. **$p < 0.01$; ***$p < 0.001$.

C-D. Stimulated in vivo C-peptide (C) and insulin (D) measurements showed reduction in Het males at 6 weeks of age on a mixed background. Two-tailed Student t test; *$p < 0.05$; **$p < 0.01$.

E. Stimulated in vivo C-peptide measurements showed no significant changes between WT and Het males at 6 weeks of age on a C57 background.

Figure 2. MafA protein status and activity are augmented in islets of C57 MafA$^{S64F/+}$ male islets compared to those from a mixed background.

A. Left lanes (1-2), representative western blotting on MIN6 nuclear extract transfected to express either MafA^{WT} or MafA^{S64F} shows faster migration in mutant MAFA due to impaired posttranslational modification by phosphorylation. Right lanes (3-6), Isolated mouse islets from each genotype and background (150 islets per lane) showed detectable levels of phosphorylated MafA^{WT} in C57 background, but relative uniformity of MafA species with faster migration and impaired phosphorylation (MafA^{S64F}) in the mixed background.

B. Quantification of western blotting bands by line scan analysis shows greater proportion of phosphorylated MafA^{WT} species (gray) in Het male islets from the C57 background compared to the mixed background.

C. qPCR of known MafA target genes in male islets between backgrounds. Blue letters, comparison between C57 WT and C57 Het islets; Black letters, comparison between mixed WT and mixed Het. *a,* p< 0.05; *b,* p < 0.01; *c,* p < 0.001; **nd,** no difference. Mean ± SEM. n=3-4 mice per group.

Figure 3. Male C57 MafA$^{S64F/+}$ islet gene expression is more similar to MafA^{WT} islets.

A. Venn diagram illustrates the total number of RNA-seq-identified DEGs up- or down-regulated between 6-week-old WT and Het male islets of each background.

B. Unbiased hierarchical clustering showed congruent organization within genotypes and genetic background.

C. Gene Ontology (GO) molecular function analysis (p < 0.05) of the genes uniquely upregulated in each genetic background in male Het islets.

D. Heatmaps showing aging/senescence enrichment in mixed, but not C57, Het male islets. Each column represents islets from a unique animal.

E. Left, qPCR of senescence signature genes are enriched in Het islets from a mixed background. Right, senescence associated beta gal staining (SA-β-gal) is enriched in male Het islets from the mixed background. *p < 0.05, **p < 0.01. Scale bar, 20μm.

F. Left, qPCR of senescence signature genes are not enriched. Right, SA-β-gal staining not detectable in male Het islets from the C57 background at 6 weeks of age.

Figure 4. Islet senescence is abrogated in MafA^{S64F/+} male islets on a C57 background.

A-B. Immunostaining of senescence markers show enrichment of 53BP1 (A) and relative loss of nuclear LaminB1 (B) in the mixed, but not C57, background. Scale bar, 50μm. Arrowhead, positive stain for 53BP1.

Figure 5. MafA directly modulates downregulation of retinol signaling in C57 MafA^{S64F/+} male islets.

A. Expression of retinoic acid receptors signaling components and known target genes Rbp1 and Cyp26a are altered specifically in C57 MafA^{S64F/+} (Het) male islets. TPM±SEM. *p < 0.05, **p < 0.01. ***p < 0.001. n=5-6 animals/group.

B. Top, DEGs identified uniquely in Male Het islets from a mixed background (1238 + 1114 = 2352 genes) versus those uniquely from a C57 background (228 + 855 = 1083 genes) were then overlayed with peaks identified by endogenous MafA CUT&RUN in mouse MIN6 cells (n=33,013 peaks). Of these, 250 DEGs uniquely enriched in a C57 background overlapped with a MafA CUT&RUN peak, while 1210 DEGs were uniquely enriched in a mixed background overlapped with a MafA CUT&RUN peak.

Bottom, UCSC Genome Browser tracks showing genomic regions associated with endogenous MafA CUT&RUN peaks near known targets *Ins1* and *Ins2* (positive controls), and candidate genes *Rara* and *Rarb*. MafA CUT&RUN enriched peaks are highlighted in dashed boxes, and regulated genes are depicted below IgG control tracks.

C. Immunostaining of RARα show downregulation in the C57 but not mixed background MafA^{S64F/+} males. Scale bar, 20μm.

D. Top, RNAScope for *Cdkn1a* encoding p21 shows enrichment in Het islets on a mixed background in β-cells. Bottom, Quantification of *Cdkn1a*$^+$Ins$^+$ cells. *p < 0.05.

SUPPLEMENTAL FIGURE AND TABLE LEGENDS

Supplemental Figure 1. Mixed background male and female MafA$^{S64F/+}$ **show hypoglycemia while nursing, extended wean does not impact hyperglycemia in male Hets, and weight gain and insulin tolerance are similar between the genetic backgrounds**

A. Random blood glucose monitoring at 2-3 weeks of age while nursing shows similar mild, transient hypoglycemia in male and female Het mice on a mixed background. *p < 0.05; ****p < 0.0001.

B. Random glucose level of suckling male Het mice on a mixed background at 6 weeks of age during an extended wean still reveals hyperglycemia, suggesting that weaning does not precipitate dysglycemia. **p < 0.01.

C-D. Weight gain as a proxy for overall health across genotypes, biologic sex, and age does not appear significantly altered between groups.

E. Insulin tolerance testing showing comparable peripheral insulin sensitivity in male and female WT and Het mice on a mixed background, similar to results in the previously published mixed background.

Supplemental Figure 2. Senescent signatures are not evident in MafA$^{S64F/+}$ **female islets from C57 and mixed genetic backgrounds .**

qPCR showing senescence signature genes are not enriched in mixed (left) and C57 (right) background female Het islets at 6 weeks of age.

Supplemental Figure 3. Binding ability of MAFA^{WT} and mutant MAFA^{S64F} are comparable.

Gel shift assay of HeLa nuclear extract (NE) produced MAFA^{WT} and mutant MAFA^{S64F} bound to *INS* enhancer (known MAFA target), showing similar binding ability. WT comp, wild-type competitor. Mut comp, mutant competitor. Specificity of MAFA binding showed by super shift (SS) with MafA antibody.

Supplemental Table 1. Primer sequences and antibody sources

REFERENCES

1. Y. Hang, R. Stein, MafA and MafB activity in pancreatic β cells. *Trends Endocrinol Metab* **22**, 364–373 (2011).
2. I. El Khattabi, A. Sharma, Proper activation of MafA is required for optimal differentiation and maturation of pancreatic beta-cells. *Best Pract Res Clin Endocrinol Metab* **29**, 821–831 (2015).
3. I. Artner, Y. Hang *et al.*, MafA and MafB regulate genes critical to β cells in a unique temporal manner. *Diabetes* **59**, 2530–2539 (2010).
4. H. A. Cyphert, E. M. Walker *et al.*, Examining how the MAFB transcription factor affects islet β cell function postnatally. *Diabetes* **68**, 337–348 (2019).
5. Y. Hang, T. Yamamoto *et al.*, The MafA transcription factor becomes essential to islet β cells soon after birth. *Diabetes* **63**, 1994–2005 (2014).
6. C. Zhang, T. Moriguchi *et al.*, MafA is a key regulator of glucose-stimulated insulin secretion. *Mol Cell Biol* **25**, 4969–4976 (2005).
7. C. Luan, Y. Ye *et al.*, The calcium channel subunit gamma-4 is regulated by MafA and necessary for pancreatic β cell specification. *Commun Biol* **2**, 106 (2019).
8. J. Cha, X. Tong *et al.*, Species-specific roles for the MAFA and MAFB transcription factors in regulating islet beta cell identity. *JCI Insight* **8**, (2023).
9. S. Guo, C. Dai *et al.*, Inactivation of specific β cell transcription factors in type 2 diabetes. *J Clin Invest* **123**, 3305–3316 (2013).
10. L. R. Cataldo, T. Singh *et al.*, MAFA and MAFB regulate exocytosis-related genes in human beta-cells. *Acta Physiol (Oxf)* **234**, e13761 (2022).
11. C. Aguayo-Mazzucato, A. Koh *et al.*, MafA expression enhances glucose-responsive insulin secretion in neonatal rat β cells. *Diabetologia* **54**, 583–593 (2011).
12. L. Wang, Y. Huang *et al.*, Differentiation of iPSCs into insulin-producing cells via adenoviral transfection of PDX-1, NeuroD1 and MafA. *Diabetes Res Clin Pract* **104**, 383–392 (2014).
13. A. Jeyagaran, M. Urbanczyk *et al.*, Forward programming of hiPSCs towards beta-like cells using Ngn3, Pdx1, and MafA. *Sci Rep* **14**, 13608 (2024).
14. A. Bonnefond, R. Unnikrishnan *et al.*, Monogenic diabetes. *Nat Rev Dis Primers* **9**, 12 (2023).
15. D. Iacovazzo, S. E. Flanagan *et al.*, MAFA missense mutation causes familial insulinomatosis and diabetes mellitus. *Proc Natl Acad Sci U S A* **115**, 1027–1032 (2018).
16. E. M. Walker, J. Cha *et al.*, Sex-biased islet beta cell dysfunction is caused by the MODY MAFA S64F variant by inducing premature aging and senescence in males. *Cell Rep* **37**, 109813 (2021).
17. D. A. Fontaine, D. B. Davis, Attention to Background Strain Is Essential for Metabolic Research: C57BL/6 and the International Knockout Mouse Consortium. *Diabetes* **65**, 25–33 (2016).
18. M. P. Keller, M. E. Rabaglia *et al.*, Gene loci associated with insulin secretion in islets from non-diabetic mice. *J Clin Invest* **129**, 4419–4432 (2019).
19. S. Andrikopoulos, C. M. Massa *et al.*, Differential effect of inbred mouse strain (C57BL/6, DBA/2, 129T2) on insulin secretory function in response to a high fat diet. *J Endocrinol* **187**, 45–53 (2005).
20. R. L. Hull, J. R. Willard *et al.*, High fat feeding unmasks variable insulin responses in male C57BL/6 mouse substrains. *J Endocrinol* **233**, 53–64 (2017).
21. H. I. Ortega, M. S. Udler *et al.*, Diabetes mellitus polygenic risk scores: heterogeneity and clinical translation. *Nat Rev Endocrinol* **21**, 530–545 (2025).

22. P. Bowman, S. E. Flanagan, A. T. Hattersley, Future Roadmaps for Precision Medicine Applied to Diabetes: Rising to the Challenge of Heterogeneity. *J Diabetes Res* **2018**, 3061620 (2018).

23. S. Misra, Ethnic diversity in precision medicine: a reality or an aspiration? *Diabetologia* **68**, 2449–2464 (2025).

24. J. Murray Leech, R. N. Beaumont *et al.*, Common genetic variants modify disease risk and clinical presentation in monogenic diabetes. *Nat Metab* **7**, 1819–1829 (2025).

25. R. Kingdom, C. F. Wright, Incomplete Penetrance and Variable Expressivity: From Clinical Studies to Population Cohorts. *Front Genet* **13**, 920390 (2022).

26. R. Kingdom, R. N. Beaumont *et al.*, Genetic modifiers of rare variants in monogenic developmental disorder loci. *Nat Genet* **56**, 861–868 (2024).

27. A. C. Fahed, M. Wang *et al.*, Polygenic background modifies penetrance of monogenic variants for tier 1 genomic conditions. *Nat Commun* **11**, 3635 (2020).

28. J. Cha, X. Tong *et al.*, Defining unique structural features in the MAFA and MAFB transcription factors that control Insulin gene activity. *J Biol Chem*, 107938 (2024).

29. C. Dos Santos, A. Cambraia *et al.*, Calorie restriction increases insulin sensitivity to promote beta cell homeostasis and longevity in mice. *Nat Commun* **15**, 9063 (2024).

30. P. Bankhead, M. B. Loughrey *et al.*, QuPath: Open source software for digital pathology image analysis. *Sci Rep* **7**, 16878 (2017).

31. U. Schmidt, Weigert, M., Broaddus, C., Myers, G., Cell detection with star-convex polygons. *International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, (2018).

32. C. Benner, T. van der Meulen *et al.*, The transcriptional landscape of mouse beta cells compared to human beta cells reveals notable species differences in long non-coding RNA and protein-coding gene expression. *BMC Genomics* **15**, 620 (2014).

33. O. Fornes, J. A. Castro-Mondragon *et al.*, JASPAR 2020: update of the open-access database of transcription factor binding profiles. *Nucleic Acids Res* **48**, D87–D92 (2020).

34. G. Tan, B. Lenhard, TFBSTools: an R/bioconductor package for transcription factor binding site analysis. *Bioinformatics* **32**, 1555–1556 (2016).

35. M. P. Meers, T. D. Bryson *et al.*, Improved CUT&RUN chromatin profiling tools. *Elife* **8**, (2019).

36. A. L. Lin, J.P. Cartailler, in *peakScout – a user-friendly and reversible peak-to-gene translator for genomic peak calling results:* *https://github.com/vandydata/peakScout*. (2025).

37. Y. Song, in *ggcoverage - Visualize and annotate omics coverage with ggplot2:* *https://showteeth.github.io/ggcoverage/*. (2025).

38. N. Rocques, N. Abou Zeid *et al.*, GSK-3-mediated phosphorylation enhances Maf-transforming activity. *Mol Cell* **28**, 584–597 (2007).

39. D. Szklarczyk, R. Kirsch *et al.*, The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Res* **51**, D638–D646 (2023).

40. H. K. Hong, E. Maury *et al.*, Requirement for NF-kappaB in maintenance of molecular and behavioral circadian rhythms in mice. *Genes Dev* **32**, 1367–1379 (2018).

41. B. J. Weidemann, B. Marcheva *et al.*, Repression of latent NF-kappaB enhancers by PDX1 regulates beta cell functional heterogeneity. *Cell Metab* **36**, 90–102 e107 (2024).

42. J. Bass, M. A. Lazar, Circadian time signatures of fitness and disease. *Science* **354**, 994–999 (2016).

43. K. Rakshit, J. Qian *et al.*, The islet circadian clock: entrainment mechanisms, function and role in glucose homeostasis. *Diabetes Obes Metab* **17 Suppl 1**, 115–122 (2015).

44. N. Seshadri, C. A. Doucette, Circadian Regulation of the Pancreatic Beta Cell. *Endocrinology* **162**, (2021).

45. M. Colucci, S. Zumerle *et al.*, Retinoic acid receptor activation reprograms senescence response and enhances anti-tumor activity of natural killer cells. *Cancer Cell* **42**, 646–661 e649 (2024).
46. M. Mushtaq, J. Liano-Pons *et al.*, EZH2 inhibition sensitizes retinoic acid-driven senescence in synovial sarcoma. *Cell Death Dis* **15**, 836 (2024).
47. R. E. Watson, J. Arjuna Ratnayaka *et al.*, Retinoic acid receptor alpha expression and cutaneous ageing. *Mech Ageing Dev* **125**, 465–473 (2004).
48. S. Naderi, H. K. Blomhoff, Retinoic acid prevents phosphorylation of pRB in normal human B lymphocytes: regulation of cyclin E, cyclin A, and p21(Cip1). *Blood* **94**, 1348–1358 (1999).
49. S. H. Park, J. S. Lim, K. L. Jang, All-trans retinoic acid induces cellular senescence via upregulation of p16, p21, and p27. *Cancer Lett* **310**, 232–239 (2011).
50. G. C. Bao, J. G. Wang, A. Jong, Increased p21 expression and complex formation with cyclin E/CDK2 in retinoid-induced pre-B lymphoma cell apoptosis. *FEBS Lett* **580**, 3687–3693 (2006).
51. C. Fottner, S. Sollfrank *et al.*, Second MAFA Variant Causing a Phosphorylation Defect in the Transactivation Domain and Familial Insulinomatosis. *Cancers (Basel)* **14**, (2022).
52. C. Aguayo-Mazzucato, J. Andle *et al.*, Acceleration of β cell aging determines diabetes and senolysis improves disease outcomes. *Cell Metab* **30**, 129–142 e124 (2019).
53. P. J. Thompson, A. Shah *et al.*, Targeted elimination of senescent β cells prevents type 1 diabetes. *Cell Metab* **29**, 1045–1060 e1010 (2019).
54. M. Bandral, L. Sussel, D. S. Lorberbaum, Retinoid signaling in pancreas development, islet function, and disease. *Curr Top Dev Biol* **161**, 297–318 (2025).
55. P. J. Brun, A. Grijalva *et al.*, Retinoic acid receptor signaling is required to maintain glucose-stimulated insulin secretion and beta-cell mass. *FASEB J* **29**, 671–683 (2015).
56. N. J. Hogrebe, K. G. Maxwell *et al.*, Generation of insulin-producing pancreatic beta cells from multiple human stem cell lines. *Nat Protoc* **16**, 4109–4143 (2021).
57. D. S. Lorberbaum, S. Kishore *et al.*, Retinoic acid signaling within pancreatic endocrine progenitors regulates mouse and human beta cell specification. *Development* **147**, (2020).
58. M. A. Kane, A. E. Folias *et al.*, Identification of 9-cis-retinoic acid as a pancreas-specific autacoid that attenuates glucose-stimulated insulin secretion. *Proc Natl Acad Sci U S A* **107**, 21884–21889 (2010).
59. S. Preza, B. Zheng *et al.*, DEC1 regulates human beta cell functional maturation and circadian rhythm. *Cell Rep* **44**, 116666 (2025).
60. A. Zankl, E. L. Duncan *et al.*, Multicentric carpotarsal osteolysis is caused by mutations clustering in the amino-terminal transcriptional activation domain of MAFB. *Am J Hum Genet* **90**, 494–501 (2012).
61. R. Russell, P. P. Carnese *et al.*, Loss of the transcription factor MAFB limits β cell derivation from human PSCs. *Nat Commun* **11**, 2742 (2020).

Main figure 1

A

MIN6   Mouse islets

MAFA^WT   MAFA^S64F   C57 WT   C57 Het   Mixed WT   Mixed Het

50kDa

37kDa

MafA^WT
MafA^S64F

Actin

Lane: 1   2   3   4   5   6

B

Line scan analysis

MAFA^WT
MAFA^S64F
C57 WT
C57 Het
Mixed WT
Mixed Het

MafA^WT   MafA^S64F

Quantification of bands

MafA^WT
MafA^S64F

Band intensity normalized to Loading control (A.U.)

MAFA^WT (C57)   MAFA^S64F (C57)   MAFA^WT (Mixed)   MAFA^S64F (Mixed)

C

c b   *b a*   *c a* nd   *a*   *c*   *b*

Fold change over WT
(expression normalized to GAPDH)

Ins1   Ins2   Cacng4   Nkx6.1   Glp1r   Gck   Slc2a2   Slc30a8   MafA

Mixed WT
Mixed Het
C57 WT
C57 Het

Main figure 2

A Differentially expressed genes

C57 Down: 1481
Mixed Down: 1861

1119

1240

C57 Up: 616
Mixed Up: 1505

B Unbiased clustering

Mixed Het Male 1
Mixed Het Male 2
Mixed Het Male 3
Mixed Het Male 4
Mixed Het Male 5
Mixed Het Male 6
C57 Het Male 1
C57 Het Male 2
C57 Het Male 3
C57 Het Male 4
C57 Het Male 5
C57 Het Male 6
C57 WT Male 1
C57 WT Male 2
C57 WT Male 3
C57 WT Male 4
C57 WT Male 5
Mixed WT Male 1
Mixed WT Male 2
Mixed WT Male 3
Mixed WT Male 4
Mixed WT Male 5

C

GO Pathways: Mixed Upregulated Genes (1114)
Protein digestion and absorption
ECM-receptor interactions
Calcium signaling
Cell adhesion molecules
Neuroactive ligand-receptor interactions
$-\log_{10}(P)$

GO Pathways: Mixed Downregulated Genes (1238)
Circadian rhythm
Maturity onset diabetes of the young
Ala, Asp, Glu metabolism
Toll-like receptor signaling pathway
Phosphonate and phosphonate metabolism
$-\log_{10}(P)$

GO Pathways: C57 Upregulated Genes (228)
Pancreatic secretion
One carbon pool by folate
Protein digestion and absorption
Calcium signaling
PI3K-Akt signaling pathway
$-\log_{10}(P)$

GO Pathways: C57 Downregulated Genes (855)
Nuclear receptors in Lipid metabolism
Retinol Metabolism
Burn Wound Healing
Nuclear Receptors
Spinal cord injury
$-\log_{10}(P)$

D

Aging
Expressions score
Cdkn1a
Eng
Icam1
Id2
Map2k1
Mapk14
Mif
Prkcd
Serbp1
Tbx2
Tbx3
Zmiz1
C57 WT   C57 Het   Mixed WT   Mixed Het

Senescence Composite
Expressions score
Ankrd1
Bcl2a1d
Ccl11
Cd68
Cdkn1a
Cdkn2a
Gdf15
Glb1
Icam1
Igfbp3
Igfbp5
Lmnb1
C57 WT   C57 Het   Mixed WT   Mixed Het

E Mixed background

Fold change over WT (expression normalized to β-Actin)
WT
Het
Cdkn1a ** Lmnb1 Bcl2a1a Cd68 * Ankrd1 **

Insulin   SA-β-gal+
$MafA^{WT}$
$MafA^{S64F/+}$

F C57 background

Fold change over WT (expression normalized to β-Actin)
WT
Het
Cdkn1a Lmnb1 Bcl2a1d Cd68 Ankrd1

Insulin   SA-β-gal+
$MafA^{WT}$
$MafA^{S64F/+}$

Main figure 3

Main figure 5