Roost abandonment and behavioural shifts following human disturbance of vampire bats in complex landscapes

Rita Ribeiro[1,2], Carlos M. Zariquiey[3,4], Wendi Chavez[4], Kristhie Pillaca Rodriguez[4], Joaquin Clavijo Manuttupa[4], Marco Antonio Risco Huiman[4,5], Roselvira Zuniga Villafuerte[4,6], Walter Campana Quintanilla[4], Noelia Medrano-Uchuya[4], Hollie French[1,2], Jakub Czyzewski[7], Carlos Shiva[8], Nestor Falcon[8], William Valderrama[4,8], Daniel P. Walsh[9], Tonie E. Rocke[10], Jason Matthiopoulos[1#], Daniel G. Streicker[1,2#]

Joint Senior Authors

[1]School of Biodiversity, One Health and Veterinary Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

[2]Medical Research Council, Centre for Virus Research, University of Glasgow, Glasgow, UK

[3]Facultad de Salud Pública y Administración, Universidad Peruana Cayetano Heredia, Lima, Perú

[4]ILLARIY (Asociación para el Desarrollo y Conservación de los Recursos Naturales), Lima, Perú

[5]Escuela Profesional de Ciencias Biológicas. Facultad de Ciencias, Universidad Nacional de Piura, Piura, Perú

[6]Centro de Investigación Vertebrate, Universidad Nacional San Antonio Abad del Cusco, Cusco, Perú

[7]School of Biodiversity, One Health and Veterinary Medicine, BioElectronics Unit,

University of Glasgow, Glasgow, UK

[8]Facultad de Medicina Veterinaria y Zootecnia, Universidad Peruana Cayetano

Heredia, Lima, Perú

[9]U.S. Geological Survey, Montana Cooperative Wildlife Research Unit, Wildlife

Biology Program, University of Montana, Missoula, Montana, USA

[10]U.S. Geological Survey, National Wildlife Health Centre, Madison, Wisconsin, USA

Correspondence: Rita Ribeiro, SBOHVM, College of Medical, Veterinary & Life

Sciences, Graham Kerr Building, University of Glasgow, Glasgow G12 8QQ

ritaclaudia.cardosoribeiro@glasgow.ac.uk

Abstract

Background Geomorphologically complex landscapes constrain animal movement

and, therefore, the spatial spread of host-associated pathogens. Monitoring animal

movement in such landscapes is required to understand these processes but

remains challenging because structural barriers to animal movement (forests,

mountains) also disrupt the visibility and signal quality of satellites used by animal-

borne Global Positioning System (GPS) tags. In Latin America, rabies transmitted by

vampire bats (*Desmodus rotundus*) epitomises this challenge: viral spread is strongly

constrained by mountains and may be exacerbated by bat dispersal responses to

human disturbance.

Methods We used GPS tags to track the post-disturbance (i.e., capture, handling

and tagging) movement of 93 vampire bats from five roosts in the highly complex

landscape of inter-Andean valleys in Peru, where signal degradation reduced usable

data by 89%. To overcome this data sparsity, the information gap needed to be filled

by modelling and data imputation. Therefore, we developed a probabilistic Bayesian

state-space model to infer the movement dynamics of vampire bats and to identify

correlates of departure time and distance from the roost, with a focus on disturbance

effects.

Results Immediately following the disturbance of capture, 43% of bats, particularly

males and stronger flyers, disappeared from their roosts for the remainder of the

study period, suggesting an acute disturbance response. For the 55 remaining GPS-

tagged vampire bats, we observed longer flight distances and high individual

heterogeneity on the night of disturbance, with bats gradually remaining closer to the

roost over the 9-day study period.

Conclusions Roost abandonment and expanded foraging areas are expected to

exacerbate rabies spread, demonstrating how human disturbances that alter animal

movement may have unintended consequences for disease transmission. More

generally, our Bayesian model accounting for GPS uncertainty and missing data

extends the utility of animal-borne tracking to species living in complex landscapes

with poor satellite visibility.

Keywords Bayesian integrated model, disease control, elevation, Lyssavirus,

perturbation effect, maximum distances, topography, wildlife management, satellite

tracking, zoonoses

Background

The composition and geomorphological structure of landscapes are fundamental determinants of how animals move within their environment and therefore of how host-associated infectious diseases spread. For example, large urban areas shape the movement of bobcats (*Lynx rufus*) and their Feline Immunodeficiency Virus (Retroviridae: Lentivirus), while in less urban areas, the same virus in puma (*Puma concolor*) is structured by mountain ranges and other factors that determine host-relatedness and spatial proximity (1,2). Pathogen spread within or across complex landscapes may also influence management decisions. For example, the fungal pathogen *Batrachochytrium dendrobatidis* has spread through complex mountainous landscapes in Latin America, causing major amphibian declines; the capacity to forecast spatial and temporal spread would be valuable for invasion monitoring or prevention (3). The adoption of smaller, lighter and longer-lasting animal-borne Global Positioning System (GPS) tags has the potential to revolutionise our understanding of pathogen spread by providing a mechanistic understanding of the underlying animal movements (4). While declining size and capacity to remotely download data without recapture means such tags can now be applied to a larger number of wildlife species than ever before, satellite obstructions remain a limiting factor, particularly for animal species that live in complex habitats such as dense forests or mountains that block GPS signals. Currently, inferring animal movement in such habitats is frustrated by imprecise or entirely missing data points.

In Latin America, rabies transmitted by vampire bats (*Desmodus rotundus*) represents a key example of how uncertainty in animal movement through complex landscapes impedes control of an important disease. In this region, vampire bats are

the main reservoir of the rabies virus (Rhabdoviridae, Lyssavirus), which causes

sporadic lethal outbreaks in humans and a sustained agricultural burden through

livestock mortality (5). Outbreaks can spread as epizootic waves into historically

rabies-free areas. In mountainous regions, such as the Andes of Peru, spread is

highly constrained by terrain and varies considerably in speed, from about 9-20 km

per year (6,7). Moreover, invasions appear to be accelerated by efforts to control bat

population sizes through sanctioned use of anticoagulant poisons ('vampiricides') or

unsanctioned killing of captured bats and destruction, sealing or burning of roosts

(8–10). Consistent with this idea, evaluation of a culling campaign in Peru showed

that culling in response to livestock rabies cases increased viral spatial spread, an

effect that was hypothesised to arise from increased movement of rabies-incubating

bats (9). While these observations suggest that culling affects bat movement with

downstream repercussions for rabies virus transmission, the behavioural responses

of vampire bats to disturbance remain speculative.

In principle, GPS tracking of vampire bats could test how land use, landscape

geomorphology or human disturbance affect bat movement, connecting colonies and

allowing the rabies virus to spread to new areas. Indeed, analogous approaches

combining experimental perturbation and GPS tracking in topographically simpler

landscapes (i.e. flat farmland with forest patches) revealed increased ranging and

dispersal behaviours in badgers (*Meles meles*) that survived culls, likely explaining

increasing tuberculosis (*Mycobacterium bovis*) risk in cattle outside of culled areas in

the United Kingdom (UK) (11). Although recent technological advances, including

smaller size and remote data recovery, allow GPS monitoring of small animals such

as bats, reliance on satellite visibility introduces challenges in complex landscapes

where vampire bats occur and where the rabies problem can be most acute (6,12).

Indeed, to our knowledge, no GPS study on vampire bats has been reported.

Alternative techniques to directly measure fine-scale behaviour in vampire bats are

precluded by technical and ecological challenges. In mark-recapture studies, low

between-roost recapture rates limit inference on movement (13). Very high

frequency (VHF)-radio tracking has also been applied, but is labour-intensive, error-

prone and only records animal presence near antennas (14). Use of these methods

in vampire bats has thus been limited to coarse-scale studies at foraging sites and

identification of unknown roosts (15,16). Overcoming the limitations of GPS satellite

visibility and signal quality would therefore represent a major opportunity to study the

relationships between landscape composition, human activity and infectious disease

spread in geomorphologically complex areas.

We used lightweight GPS tags, enabling remote data recovery, to assess fine-

scale movement of vampire bats in response to human disturbance (i.e., from

capture, handling, and tagging). We focused on the Apurimac, Ayacucho and Cusco

(AAC) departments in the Peruvian Andes, where bats live within a network of

interconnected inter-Andean valleys. The AAC is a hotspot of vampire bat rabies

(VBR) in Peru, accounting for 60.2% of all reported livestock cases between 2003

and 2019. This region is subjected to regular bat culls and has experienced wave-

like invasions of rabies into historically rabies-free areas (9). To address the

challenges imposed by the obstructive local topography, we developed a new

Bayesian state-space method to reconstruct times and distances from sparse and

error-prone GPS data (17). This model characterised the trips of individual bats,

addressing data sparsity by 1) simultaneously reconstructing and identifying

ecological correlates for two key movement metrics, nightly departure time and

distance from the roost every 15 minutes of a night; and 2) fitting the model to all

bats at once, which allowed the full dataset to be used to inform and fill in missing

data. This approach leveraged all available information to statistically impute missing

data and propagate the uncertainty, resulting in more robust inference from sparse

datasets than conventional interpolation methods. We first investigated the causes of

immediate roost switching or abandonment after disturbance (hereafter, roost

abandonment, defined as the non-recapture of bats and non-recovery of GPS data

after disturbance during the study period). Then, for the bats that remained in the

roost, using the Bayesian model, we tested the hypothesis that disturbance would

induce longer flights.

Methods

Study area and design

The study was conducted during the Peruvian dry season, between April-

October 2022 and May-June 2023, in five vampire bat roosts (Table S1), each

located >50 km apart in southern Peru. The study area is characterised by rugged

topography, with altitudes ranging from 1200 to 6000 meters, with vampire bat

presence and rabies virus spread constrained to elevations below ca. 3600 meters

(6) (Fig. 1a).

At each roost, two capture and tagging sessions were carried out, with a

minimum interval of three months. Roost 2 (Fig. 1c) had a single tagging session due

to data recovery challenges. Each session aimed to tag 10 adult vampire bats (five

males, five females). All capture sessions began after dusk, except the first capture

in April 2022, when, due to logistical constraints, captures began in the morning.

Short-term follow-up recapture sessions (one to two nights) were conducted at the

end of the GPS-tracking study, whereas long-term sessions occurred during

subsequent tagging rounds or in later years to monitor colony dynamics (Table S1).

Capture and handling of vampire bats

Vampire bats were captured using mist and butterfly nets. Captured bats were

fitted with a 3.5 mm wing ring (Porzana, UK), and weight (g), sex, reproductive status

(non-reproductive, scrotal male, pregnant, or lactating female), age (adult, subadult,

or juvenile), and forearm length (mm) were recorded. We applied GPS tags (2.8 g, 8

cm length [including a 5 cm antenna]; nanoFix®, PathTrack, UK) to bats that met our

inclusion criteria (see below). After light sedation using intramuscular (pectoral)

injection (29-gauge needle, 1.0 mL syringe) of Ketamine (20–25 mg/kg) diluted in

saline (NaCl) at a 50/50 volume ratio or Midazolam (0.25 mg/kg, used in 2023 to

allow faster recovery times), GPS tags were attached to the skin of the dorsal side of

each bat using an absorbable suture (Monosyn®, B Braun), threaded through

custom tag loops (18). According to the manufacturer, the absorbable sutures lose

50% of their strength after 14 days, allowing natural tag detachment. Before suturing,

hair was trimmed, and a small amount of cyanoacrylate-based glue was used to

secure the tags. Sedation and tag suturing were performed by a veterinarian,

ensuring it did not impair flight or walking. GPS tags were painted with UV

fluorescent paint to aid recovery for reuse (Starglow®, Glowtec, UK).

GPS tagging was limited to healthy adult bats, excluding pregnant and

lactating females. The average weight of tagged bats was 43.4 g ($\pm$ 3.4 g, N=93),

44.1 g ($\pm$ 3.3 g, N=43) for females and 42.7 g ($\pm$ 3.4 g, N=50) for males. For both

sexes, the tags represented a maximum of 7% of the bat's weight (mean = 6.4 $\pm$

0.47% for females and mean = 6.6 $\pm$ 0.47% for males). Recommendations for bat

tracking studies indicate that attached devices should not exceed 10% of body mass,

with a best practice of limiting to 5%. However, heavier tags are considered

acceptable for short-term use in species with high physiological tolerance for weight

change (19,20). Our previous records of wild adult vampire bat mass (N=3,429

individuals) showed that they can gain over 9 g of weight after feeding (pre-feeding:

41.9 g ± 8.44, N=2,751; post-feeding: 51.5 g ± 9.1, N=678), implying daily weight

gains of >20% of body mass. Additionally, during pregnancy and after birth, females

can weigh over 85 g and continue foraging with their pups attached. These weight

fluctuations and their ability to carry significant additional mass strongly indicate that

the tags we used should be tolerable for short-term use.

During initial deployment in April 2022, GPS tags were set to record locations

every 5 minutes (10 tags) or 10 minutes (five tags) for three and six nights,

respectively. To extend the duration of tracking for up to 9 nights in subsequent

sessions, later deployments used a lower temporal resolution of 15 minutes (see

later sections for how data with different time intervals were integrated). Tags

operated nightly from 6 PM to 6 AM, with data stored onboard and downloaded

remotely via one or two solar-powered UHF base stations (PathTrack, UK) placed

near the roost.

Ethical approval and permits for the capture, handling, and GPS tagging of

vampire bats were granted by the Research Ethics Committee of the University of

Glasgow (Reference EA28a/21) and by Peruvian national authorities, including

SERFOR (Servicio Nacional Forestal y de Fauna Silvestre, Nos. D000024-2022,

D000107-2022, D000103-2024 and D000188-2024-MIDAGRI-SERFOR-

DGGSPFFS-DGSPFS) and SERNANP (Servicio Nacional de Áreas Naturales

Protegidas por el Estado, N° 021-2023-SERNANP-SHM/J).

Data cleaning and management

According to the manufacturer, the nanoFix GPS accuracy depends on signal

quality and the number of satellites tracked by the tag. Specifically, tags require at

least four visible satellites to record a position ('fix') such that fixes with fewer than 4

satellites are automatically not available and not recoverable. Due to the complex

topography of our study area, 89% of the GPS points were not recoverable, resulting

in a sparse and irregular dataset. Prior to developing the modelling framework

outlined below to reconstruct movement from such data, we carried out manual data

cleaning in R version 4.4.0 (21). Specifically, GPS fixes recorded before tag

application and 1 hour after capture were removed to account for the tag attachment

process. Data from the first night were otherwise included because quantifying the

disturbance immediately after capture was one of our objectives. Cleaning of the

remaining data included one inaccurate fix flagged by the software, all fixes from one

bat with a malfunctioning tag (distances up to 6 km recorded before attachment), all

data from three bats with less than three fixes across the study period, and all data

from an unrealistically sessile bat (suggesting tag loss or death). As detailed in the

supporting information (S1), we also removed GPS fix jumps and outliers of

estimated flight speed (22). The speed (km/hour) of each segment between

subsequent fixes was calculated, along with the Euclidean distance from the roost

for each fix, using base R and the sf package (v1.0-19) (23). Our state-space model

was designed to be robust to GPS errors, so we could have omitted data filtering

based on speed entirely. However, we anticipated that simultaneously inferring and

filtering speed within the model could hinder convergence.

To manage the three sampling rates (5, 10 and 15 minutes) and provide a

uniform input for the Bayesian model, each GPS fix was assigned to a 15-minute

block during the night. The model's input matrix was constructed from these blocks

(i.e., 48 15-minute blocks between 6 PM and 6 AM of the following day), with each

block recorded as either containing an observed distance from the roost or marked

as missing (NA).

Variable definition and processing

We investigated predictors of bats' nightly departure time and distance

travelled from the roost. Vampire bats are thought to be lunar phobic, departing

before moonrise or after moonset (24). We therefore modelled departure time as a

function of the number of hours until complete darkness, which depended on the

interplay of moonrise, moonset, astronomical night start (occurs after sunset and

dusk) and end (occurs before dawn). For each roost and night, we calculated the

hours from sunset until darkness using the *suncalc* package (v0.5.1) (25). We

predicted that longer periods until darkness would delay departure times. We also

tested whether sex influences departure time.

We modelled distance from the roost as a function of variables related to roost

location, individual bat characteristics, and human disturbance. Roost-related

variables within 5 km radii of roosts included proportion of permanent water,

predicted vampire bat roost density (26) and cattle density (27) (Table S2),

calculated using the packages *sf* (v1.0-19) and *terra* (v1.7-71) (23,28). We predicted

that the availability of alternative roosts and preferred food sources (cattle) near the

roost would reduce the distances travelled. Since cattle graze near water and

vampire bats can use rivers as corridors, we hypothesised that permanent water

sources near roosts could aid prey searching, decreasing distances from the roost

(29). Individual characteristics included bat sex and forearm length. We

hypothesised that males would fly further distances and would be more likely to

abandon their roosts after capture due to reduced offspring care (30). We also

expected that individuals with longer forearms may travel farther from the roost or

would be more likely to switch roosts after capture, since forearm length scales with

wingspan and area, providing a proxy for wing loading and flight speed (31,32), and

is positively associated with migration distance in bats (33).

Human disturbance was defined as the effects of capture, handling, and

tagging. Key variables included the night of capture ('night 0') versus subsequent

nights, and the time post-disturbance, calculated as the difference in hours between

the start of the tracking session and each GPS fix date and time stamp. We

averaged the time post-disturbance for each bat and each night to obtain a single

value per bat per night. Averaging reduced the influence of irregular GPS fix intervals

and temporal autocorrelation, and provided a covariate structure consistent with our

model, which included effects of individual, roost and night. We anticipated that

disturbance would increase flight distance following capture, with distances

decreasing over time as behaviour normalised. Finally, because vampire bats in our

study area are deliberately or accidentally disturbed by humans (e.g., capture,

culling, roost destruction or shifts in agricultural practices), we included the number

of rural human settlements within 5 km of each roost as a proxy for historical

disturbance levels, allowing us to isolate the effects of the immediate disturbance

caused by our capture, handling and tagging from longer term behavioural effects

arising from historical persecution (Table S2).

Drivers of roost abandonment

By placing UHF base stations at roost sites, we were able to monitor whether GPS-tagged bats returned to the roosts where they were first tagged during the 9 nights of GPS tracking. An absence of recorded GPS data from base stations across the study period, therefore, indicated roost switching or, at a minimum, short-term roost abandonment. We fitted binomial generalised linear models (GLM) to examine whether roost abandonment was associated with individual bat traits (i.e., sex and forearm length). Vampire bat weight was not included because it fluctuates considerably within a night, depending on feeding and time since feeding. Nevertheless, with this caveat, for each bat, the percentage of body mass represented by the tag was treated as a fixed covariate to evaluate the potential influence of tag burden on roost abandonment behaviour. We assessed multicollinearity between variables by calculating variance inflation factors, adopting a threshold of 2.5 as an indicator of high multicollinearity (34). Model performance was compared across 14 models (single effect models and their interaction terms) using the Akaike information criterion (AIC) and pseudo-R^2. Nested ANOVAs were used to test the statistical significance of interaction terms and added variables.

Modelling framework to infer bat movement metrics

We developed a Bayesian state-space model with two latent variables: the bats' departure time (t, in minutes since the daily activation of the GPS at 6 PM) and its distance from the roost (ψ, in km). The model reconstructed distances from the roost every 15 minutes and estimated the duration of activity and return time to the roost, accounting for both process error (i.e., the natural stochasticity in the biological processes) and observation error (i.e., GPS measurement error). To reconstruct

nightly trips for each bat and distances from the roost, the model needs to have a

known starting and ending point. Hence, we assumed that each trip begins and ends

at the main roost. Although this assumption could be challenged by the potential use

of multiple roosts, a preliminary analysis showed that at least 56.6% of bats-nights

returned to the roost, as indicated by GPS fixes near the roost the following morning.

This estimate is likely conservative due to data limitations, such as missed fixes

during rapid returns and signal loss at the roost, which may underestimate actual

return rates. We also assumed that most bats would depart before 1 AM, as 95% of

our experiments were conducted on nights when the moon set before 1 AM.

For each bat i and night j, $t_{i,j}$ is the departure time in minutes after 6 PM, with

a maximum value of 420 minutes (1 AM). The expected and realised departure

instances, $\bar{\tau}_{i,j}$ and $\tau_{i,j}$, are expressed as a proportion of the interval between 6 PM

and 1 AM,

(1) $\tau_{i,j} = \frac{t_{i,j}}{420}$,

(2) $t_{i,j} = 420\tau_{i,j}$.

The expected departure instance $\bar{\tau}_{i,j}$ was modelled as a logit function of the number

of hours until darkness and bat sex,

(3) $logit(\bar{\tau}_{i,j}) = \theta_0 + \theta_1 hours\_dark_j + \theta_2 sex_i$.

$\tau_{i,j}$ was modelled using a Beta distribution with mean $\bar{\tau}_{i,j}$ and a process error

quantified as the standard deviation of departure instances σ. This error represents

variability in trip start times and was assigned a wide prior (estimated by the model,

table S3), to capture pooled inter-trip and inter-individual variation,

(4) $\tau_{i,j} \sim Beta(a_{i,j}, b_{i,j})$,

(5) $a_{i,j} = \bar{\tau}_{i,j}(\frac{\bar{\tau}_{i,j}(1-\bar{\tau}_{i,j})}{\sigma^2} - 1)$,

(6) $b_{i,j} = \left(1 - \bar{\tau}_{i,j}\right)\left(\frac{\bar{\tau}_{i,j}(1-\bar{\tau}_{i,j})}{\sigma^2} - 1\right).$

The duration of nightly activity after departure was modelled with a uniform

distribution, assuming that once a bat departed, it remained away for at least 20

minutes. The maximum duration of a trip was the time left after departing and the

end of GPS activation (720 minutes),

(7) $\Delta t_{i,j} \sim U\left(20, 720 - t_{i,j}\right).$

To estimate distances travelled from the roost, we defined a parametric model

of distance against time, for each bat i, night j, and each block k of 15 minutes after

6 PM,

(8) $\bar{\psi}_{i,j,k} = exp\left(\beta_0 - \frac{\left(\left(k15 - t_{i,j}\right) - \Delta t_{i,j}\tau_{mid,i,j}\right)^2}{\left(\Delta t_{i,j}\sigma_{mid,i,j}\right)^2} + \beta_i night_j + \sum_{n=1}^{N} B_n X_{i,j,n} + \zeta_i\right).$

This equation assumes that only one trip is made by each bat every night (i.e., the

bat leaves the roost, travels to a destination and then comes back), but is flexible

regarding specific details of the trip, such as when it starts, ends, and maximum

distance from the roost. It also assumes that the net velocity is away from the roost

until the maximum distance is reached, and then toward the roost thereafter, for 15-

minute intervals. The exponential of the intercept β_0, corresponds to the maximum

distance from the roost. β_i corresponds to the terms for the effect of night of

disturbance ($night_j$) for each bat, and B_n is the parameter for the n^{th} covariate effect

(forearm length, sex, roost density, number of human rural settlements, cattle

density, permanent water and time post-disturbance). We added an individual-level

random effect ζ_i with mean 0 and standard deviation σ_1 ($\zeta_i = N(0, \sigma_1)$), to capture

the influence of individual variation (Table S3). In both departure time and distance

models, we considered random effects of individual, roost, calendar night, and

session of capture. Only the random effect of the individual on the distance model

was relevant, hence it was kept.

The mid-point of the trip $\tau_{mid,i,j}$ was defined as the proportion of time between

departure and return until reaching maximum distance, with a parameter $\sigma_{mid,i,j}$,

which made sure that the distance from the roost climbed smoothly from the start of

the trip to the midpoint and then declined to the trip's endpoint, preventing unrealistic

jumps in movement. $\tau_{mid,i,j}$ was given a beta prior distribution and was set to be

close to the middle of the trip duration (mean 0.5 and variance 0.001). $\sigma_{mid,i,j}$ was

fixed to a sufficiently large value to ensure that the bats did not immediately reach

their maximum distance after departure or instantaneously returned to the roost,

$$(9)\ \sigma_{mid,i,j} = \left(\tau_{mid,i,j}, 1 - \tau_{mid,i,j}\right) / 2.$$

The realisation of distance giving rise to observations, $\psi_{i,j,k}$, was modelled using a

normal distribution, with mean $\bar{\psi}_{i,j,k}$ and variance $O_{i,j,k}$,

$$(10)\ \psi_{i,j,k} \sim N(\bar{\psi}_{i,j,k},\ O_{i,j,k}),\ \text{with}\ \psi_{i,j,k} \geq 0\ \text{km}.$$

$O_{i,j,k}$ is a combined error that incorporates observation error $\xi_{i,j,k}$, the standard

deviation of GPS error, and process error, in the form of an overdispersion

parameter φ, to account for further uncertainty that cannot be explained by the

observation model (i.e., in equation 11, 0.6 km is the GPS error for four satellites, the

minimum number of visible satellites required),

$$(11)\ O_{i,j,k} = \xi_{i,j,k}^2 + 0.6\varphi^2.$$

The overdispersion parameter φ was estimated by the model, but we encouraged it

to be close to zero (Table S3) to prioritise the identification of biological effects. The

nanoFix manufacturer indicated using the number of visible satellites to estimate

GPS error. Using data from a GPS tag placed at a single location in our study area,

we fitted a regression of the standard deviation of distances against the number of

satellites and estimated the standard deviation of GPS error for each satellite count

(standard deviation of 0.3 km when four satellites were visible; deviance explained =

0.986). This regression produced the two parameters γ_0 and γ_1 which, in

combination with the number of visible satellites that generated each positional fix,

$N_{i,j,k}$, informed $\xi_{i,j,k}$

(12) $\xi_{i,j,k} = exp\ (\gamma_1 + \gamma_2 N_{i,j,k})$.

We used Bayesian Stochastic Search Variable Selection (SSVS) with spike

and slab priors to estimate model coefficients conditional on a variable being

retained in the model (35). If a variable was participating in the model, the coefficient

value was given a normal distribution, with mean 0 and standard deviation τ_1, with τ_1

being estimated by the model, except for the coefficient for hours until darkness,

which was given a gamma prior distribution to preserve its positivity (mean 0.01 and

variance 0.0001). The posterior for the probability of retention ρ was modelled with a

Bernoulli distribution (variable included or excluded), with the parameter ρ being

estimated by the model with a constrained Beta prior (i.e., a variable has a 25%

probability of being included), meaning that variables need strong evidence to be

included.

We assessed goodness of fit by calculating the R^2, comparing observed

distances from the roost to the posterior predicted distance for every single GPS fix,

both for the entire model and for each bat separately. The models were fitted in

JAGS 4.3.2 using rjags (v4-15) and runjags (v2.2.2-4) (36,37). The model ran with

four independent chains, for 10,000 samples per chain after discarding the first

500,000 iterations as burn-in, with thinning of 200 (retaining every 200[th] iteration).

Convergence across chains was assessed through visual inspections of the

posterior and the Gelman-Rubin diagnostic test (using coda (v0.19-4.1) (38)). Table

S3 lists the final model parameter priors and posteriors.

Results

Between April 2022 and June 2023, 93 adult vampire bats were GPS tagged

across five roosts in southern Peru (Table S1). Twenty-four percent of tags (20 out of

83 tags used) were recovered after use, most of them found inside roosts, allowing

partial redeployment. Among the 60 bats with recovered GPS data, 89% of possible

observations (14,142 out of 15,842) were not recorded due to having less than four

visible satellites, leaving a total of 1700 GPS fixes. GPS fix availability was highest

on the night of capture ('night 0', 24.7%), followed by 'night 1' after capture (18.2%),

and generally declined each subsequent night to a minimum of 0.6% on 'night 8'

after capture (Fig. S1). Data cleaning excluded a further 22.3% of the fixes, leaving

1321 fixes from 55 bats (Fig. S2). Visualisation of the GPS data showed that vampire

bats occasionally flew to distances higher than 5 km from the roost but spent most

time near roosts (Fig. 1b-f; Fig. S3). Individual bats from the same roost foraged in

different areas, did not visit the same locations every night, and used a variety of

flight paths, highlighting behavioural heterogeneity among and within individuals (Fig.

1b, d, e, f; Fig. S4d, e).

Filtering out three bats which were confirmed dead, we recovered no GPS

data from 30 bats, and nine additional bats were not detected after 'night 0',

indicating the possibility that 43% (39/90) of bats left our study areas following our

capture and tagging effort. While we cannot rule out tag loss or death of these

missing animals in unobservable locations within our study area, recapture data

point to short- or medium-term roost abandonment rather than death or tag loss.

Specifically, of the 39 bats which disappeared, we recaptured only one (short-term

recapture rate = 2.6%) during capture sessions immediately following each tracking

period. In contrast, of the 51 bats that we were able to GPS track, eight were

recaptured in the sessions immediately after the GPS tracking period (short-term

recapture rate = 15.7%). As missing bats would be expected to have similar short-

term recapture rates if their apparent disappearance from the study area was an

artefact of GPS tag loss or failure, we would have expected at least six recaptures.

Moreover, during surveys conducted 4-32 months after GPS tagging, six of the

missing bats were recaptured (long-term recapture rate=15.4%), while six of the bats

that had remained were also recaptured (long-term recapture rate=11.8%). These

results suggest that disappearance immediately after tagging was due to roost

abandonment or switching rather than bat mortality.

A binomial GLM testing for the effects of bat forearm length and sex, and tag

burden, revealed that bats with longer forearms (-0.96 [SE = 0.35], P<0.01) and male

bats (-1.11 [SE = 0.66], P=0.09) were less likely to return to the roost during the 9-

day study period, although there was only weak evidence for the latter effect (Fig. 2;

Table S4; Fig. S5). Non-random disappearance of bats following patterns that might

be expected from bat behaviour and physiology provides further support for the idea

that disappearances reflected short- to medium-term roost abandonment by vampire

bats following disturbance. Across all models tested, including those with interaction

terms, we found no evidence that the tag burden influenced the probability of roost

abandonment.

We next explored correlates of departure time and maximum nightly distance

from the roost among the bats that remained sufficiently close to the roost to

contribute GPS data. A Bayesian state-space model demonstrated a strong overall

fit to the GPS data (R^2 = 0.80). Because the number of observed GPS fixes varied

across bats, we also calculated individual-level R^2 values to assess model fit

consistency. The model fitted well for most bats, with a median individual R^2 of 0.71.

Only 10.9% of bats were poorly fit (R^2 <0.20) (Fig. S6). Predicted maximum

distances were larger than observed maximum distances per bat and night,

reflecting the property of our model to fill gaps due to missing observations (Fig. 3c-

d). Our model estimated patterns and correlates of the time that bats departed from

the roost. On average, bats departed ~2 hours after the GPS started at 6 PM

(posterior mean of 124 minutes, 95% credible interval (CI) [58.3; 220.5]) and were

out of the roost for >5 hours (posterior mean of 345 minutes, 95% CI [190.0; 497.4])

(Fig. S7). Although the number of hours until darkness had a low posterior inclusion

probability, its posterior distribution indicated that bats left the roost as soon as there

was no sunlight or moonlight (Fig. 3a; Table S5).

Among drivers of the nightly distances that bats moved from the roost, time

post-disturbance was the only variable with strong posterior support for inclusion

(probability = 0.95, Fig. 3a, d; Table S5; Figs. S8-S9). This negative effect implied

that bats initially flew farther from the roost following disturbance, then shifted to

flying shorter distances, presumably returning to a more typical flight distance.

Specifically, the maximum nightly distance decreased by 0.17 km per night (Fig. S8)

and stabilised approximately four nights after tagging (Fig. 3d). Deeper examination

of model outputs also showed an interaction between individual bat identity and night

(night of disturbance versus all the other nights), whereby on the night of the

disturbance, some bats stayed unusually close to the roost, and others undertook

the longest movements recorded in the study period (Figs. 3b, 4; Table S6; Figs.

S10-S11). This putative individual heterogeneity in the response to disturbance

occurred across all roosts except roost 2, which had data from only one bat (Fig.

S12). Taken together, these results suggest that while on average bats flew further

from the roost on the night of disturbance, this result was dampened by marked

individual heterogeneity, where some bats had the opposite response of reducing

movement.

Discussion

Fine-scale studies of wildlife movement are currently restricted to relatively

geomorphologically simple habitats, limiting our ability to understand how natural or

human-induced behavioural variation affects infectious disease transmission in

complex habitats. By developing a Bayesian state-space model to reconstruct

movement metrics from sparse and error-prone GPS data from 93 vampire bats in a

rabies hotspot in the Peruvian Andes, we identified three behavioural responses to

human disturbance which could exacerbate rabies transmission and dilute the

effectiveness of bat population control efforts.

First, 43% of the GPS-tagged bats were not detected in the study area for the

duration of the tracking sessions that followed capture and tagging. Because any

return to the roost area would have triggered GPS data upload to our receivers,

these findings suggest that a surprisingly large fraction of bats abandoned the focal

roosts for at least nine nights (length of the study) because of the disturbance arising

from capture and tagging. While we cannot completely rule out the possibility that

tags detached in areas with poor satellite coverage, the scarcity of stationary tags

and lack of GPS fixes before presumed detachment make this explanation unlikely.

Moreover, missing bats had lower short-term recapture rates, but higher long-term

recapture rates compared to bats that remained in the study area during the GPS

tracking study, arguing against mortality as a main cause. Previous research has

shown that vampire bats switch roosts frequently (39,40), so some disappearances

in our data could reflect normal switching rather than disturbance-induced

abandonment. Unfortunately, since our measure of roost abandonment (GPS data

upload to a receiver at the roost site) required GPS tagging, a non-disturbance

control was not feasible to rule out this explanation. However, given that vampire bat

dispersal is generally male-biased and that forearm length is linked to flight ability,

the results of our GLM that males and stronger flyers more frequently disappeared

after tagging further support transient roost abandonment (7,30). Future studies

might use less invasive monitoring of roost fidelity, for example, automated

monitoring systems with radio frequency identification (RFID) tags, to estimate

baseline roost switching rates.

We observed two other behavioural responses among bats that remained

detectable: a general tendency for longer distance movements after disturbance and

elevated individual heterogeneity on the night of capture. As with the roost

abandonment analysis, the necessity of GPS tagging meant baseline flight distances

in the absence of tagging could not be inferred. While we interpret the negative effect

of time post-disturbance on flight distances as an initial increase with gradual return

to shorter distances, we cannot entirely rule out that data from the initial nights after

capture and tagging represented normal behaviour and the shorter distances

observed at the end of the observation period reflected shorter than typical flight

distances, perhaps because of fatigue from tags. However, we argue this

interpretation is unlikely since high individual heterogeneity on the night of capture

(Fig. 3b) and the putative roost abandonment effect (Fig. 2), are more consistent with

an immediate disturbance effect (41).

Collectively, these three perturbations could have unintended consequences

for both the transmission and management of VBR. Roost abandonment following

the application of vampiricide poison would dilute local population reduction and may

disperse bats to lower-density roosts or roosts where bats have fewer social

connections, reducing the efficiency of poison transfer. Disturbance-induced roost

abandonment could also have complex effects on rabies transmission. On the one

hand, roost abandonment could decelerate rabies spread if rabies-infected bats

immigrate into areas where bats have been dispersed into many small roosts, as this

would reduce the likelihood of the virus successfully establishing in new areas. On

the other hand, if disturbance occurs in areas where rabies already circulates, roost

abandonment or greater travel distances by rabies-infected bats are predicted to

raise rabies incidence in bats, through overlap of foraging areas and colony mixing,

and promote spread to nearby roosts. As such, our results potentially explain

findings that reactive culls accelerated viral spatial spread while preventive culls

slowed spread (8,9). Since viral spatial spread is believed to be strongly influenced

by male bat movement, the tendency for male bats to abandon roosts after capture

would be particularly problematic (7). In contrast, disturbance-induced movement

could benefit the recently proposed application of vaccines with the capacity for self-

dissemination since roost switching would 'seed' vaccine transmission to a larger

number of roosts than previously appreciated (42).

A caveat when drawing conclusions about culling is that our study measured

the effects of a relatively mild disturbance. While our disturbance shares similarities

with culling (capture, handling, the application of a foreign object, and release), there

are key differences. Vampire bats have strong cooperative networks, characterised

by blood meal sharing and intense allogrooming, which could be disrupted by the

sickness behaviours and mortality associated with poisoning (43,44). As such, culls

may lead to stronger effects on movement than the relatively mild disturbance we

induced. Capture locations may also vary between our study and real-world

management. While we carried out captures in roosts for greater tractability,

captures can also occur at properties where bats attack livestock (9,45). Additional

field experiments that separate the effects of human presence at roost sites, capture,

handling and poisoning, could clarify vampire bat population management.

High-resolution GPS tracking also revealed that geomorphological constraints

on bat movement and VBR spread may be less restrictive than implied by previous

observational or radiotelemetry data. For example, rather than consistently foraging

downstream of their roosts, following valleys or rivers, we observed vampire bats

travelling over mountains and foraging in both directions along rivers (46,47).

Furthermore, similar to other colonial species such as seabirds, vampire bats

exhibited high movement behavioural heterogeneity within the colony (48). Our data

also challenge the notion that vampire bats return to the same foraging location and

even individual animals for blood feeding every night (49). Overall, heterogeneity

across individual bats and within bats over short periods helps explain the

sometimes-sporadic evidence of vampire bat bites on livestock and humans (50) and

implies that pathogen-infected bats are likely to expose multiple animals at different

locations. Incorporating these behavioural ecological insights on vampire bats into

models of rabies spread could enhance our understanding of its dynamics.

The Bayesian framework we developed incorporated all available data on

departure times and distances from the roost, jointly estimating movement

parameters across individuals. By partially pooling information across bats, the

model leveraged well-sampled individuals to inform estimates for those with sparse

data, statistically imputing missing values and propagating their uncertainty.

Importantly, our objective was not to infer exact animal locations throughout the

night, but rather to estimate how individual, landscape, disturbance and temporal

factors affected key characteristics of bat movement while accounting for

uncertainty. Furthermore, this approach enabled us to estimate the true distances

bats can travel (Fig. 3c-d), which are critical for disease management, but are often

missed in sparse datasets, which tend to underrepresent maximum distances (51). A

potential limitation of our modelling approach is the assumption that vampire bats do

a single trip and return to the roost every night. This assumption was broadly

supported by overall model fit and parameter convergence. However, a subset of

bat-nights (13%) had bimodal posterior distributions that were more consistent with

multiple trips within a night (Fig. S10). If some long-distance movements instead

reflect roost switching, our results may overestimate true foraging distances;

however, we note that eventual return to the roost was required for data recovery, so

either behaviour reflects short-term movements after disturbance and will not affect

our estimates of maximum distances travelled.

Conclusions

Our study demonstrates how advances in GPS technology, despite persistent

data challenges in complex habitats, can be utilised through analytical frameworks to

extend wildlife tracking to species and environments where conventional analyses

may fail, or be strongly biased by satellite visibility constraints. Moreover, our

findings demonstrate that human disturbance can have significant repercussions for

VBR management, revealing movement dynamics that would otherwise remain

unknown in complex landscapes and highlighting how altered bat dispersal and

colony mixing may facilitate livestock rabies emergence into new areas in Latin

America.

List of abbreviations

AAC – Ayacucho, Apurímac, Cusco

AIC - Akaike Information Criterion

ANOVA – Analysis of Variance

CI – Credible interval

GLM – Generalised linear model

GPS – Global Positioning System

NA – Not available

RFID - radio frequency identification

SE – Standard Error

SSVS - Bayesian Stochastic Search Variable Selection

VBR – vampire bat rabies

VHF – Very high frequency

Declarations

Ethics approval

Ethical approval and permits for the capture, handling, and GPS tagging of vampire

bats were granted by the Research Ethics Committee of the University of Glasgow

(Reference EA28a/21) and by Peruvian national authorities, including SERFOR

(Servicio Nacional Forestal y de Fauna Silvestre, Nos. D000024-2022, D000107-

2022, D000103-2024 and D000188-2024-MIDAGRI-SERFOR-DGGSPFFS-

DGSPFS) and SERNANP (Servicio Nacional de Áreas Naturales Protegidas por el

Estado, N° 021-2023-SERNANP-SHM/J).

Consent for publication

Not applicable.

Availability of data and materials

Data and code for this study are deposited in the Dryad Digital Repository, and will

be available after publication of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

The project was funded by the NSF/BBSRC Ecology and Evolution of Infectious

Diseases Program (DEB 2011069, BB/V003798/1). D.G.S. was funded by a

Wellcome Trust Senior Research Fellowship (217221/Z/19/Z). This research

received support from the UK Medical Research Council through core funding of the

MRC-University of Glasgow Centre for Virus Research (Virus Cross Species

Transmission Programme: MC_UU_00034/3).

Authors' contributions

Conceptualisation: R.R., J.M., D.G.S. Methodology: R.R., J.C., J.M., D.G.S. Analysis

and visualisation: R.R., J.M. Investigation: R.R., C.M.Z., W.C., K.P.R., J.C.M.,

M.A.R.H., R.Z.V., W.C.Q., N.M-U., H.F., D.G.S. Resources and management:

C.M.Z., C.S., N.F., W.V. Data Curation: R.R. Project administration: D.G.S., Funding

acquisition: D.P.W., T.E.R., J.M., D.G.S. Supervision: R.R., J.M., D.G.S. Writing

(original draft): R.R., J.M., D.G.S. All authors contributed critically to the drafts and

gave final approval for publication.

Acknowledgements

We are grateful to Elsa Cardenas Canales, Haris Malik and Carlos Tello for field

assistance; Lynda Dunbar and Marieke Pingen for help with the suturing technique;

Fred Touzalin and Pathtrack for insights and assistance with the GPS. We thank

Holly Niven for helpful discussions about the analysis, and Nardus Mollentze, Travis

Mcdevitt-Galles, and our colleagues from the Streicker group for peer review and

feedback on earlier versions of the manuscript. We thank SENASA (National and

Regional offices) for feedback and support.

References

1. Fountain-Jones NM, Craft ME, Funk WC, Kozakiewicz C, Trumbo DR,

Boydston EE, et al. Urban landscapes can change virus gene flow and

evolution in a fragmentation-sensitive carnivore. Mol Ecol. 2017

Nov;26(22):6487–98.

2. Fountain-jones NM, Kraberger S, Gagne RB, Trumbo DR, Salerno PE, Funk

WC, et al. Host relatedness and landscape connectivity shape pathogen

spread in the puma, a large secretive carnivore. Commun Biol. 2021;1–9.

Available from: http://dx.doi.org/10.1038/s42003-020-01548-2

3.   Cheng TL, Rovito SM, Wake DB, Vredenburg VT. Coincident mass extirpation

of neotropical amphibians with the emergence of the infectious fungal

pathogen *Batrachochytrium dendrobatidis*. PNAS. 2011;108(23):9502–7.

4.   Nathan R, Monk CT, Arlinghaus R, Adam T, Alós J, Assaf M, et al. Big-data

approaches lead to an increased understanding of the ecology of animal

movement. Science (1979). 2022;375(6582).

5.   Schneider MC, Romijn PC, Uieda W, Tamayo H, da Silva DF, Belotto A, et al.

Rabies transmitted by vampire bats to humans: An emerging zoonotic disease

in Latin America? Rev Panam Salud Publica. 2009 Mar;25(3):260–9.

6.   Benavides JA, Valderrama W, Streicker DG. Spatial expansions and travelling

waves of rabies in vampire bats. Proceedings of the Royal Society B:

Biological Sciences. 2016;283(1832):1–9.

7.   Streicker DG, Winternitz JC, Satterfield DA, Condori-Condori RE, Broos A,

Tello C, et al. Host-pathogen evolutionary signatures reveal dynamics and

future invasions of vampire bat rabies. Proc Natl Acad Sci U S A.

2016;113(39):10926–31.

8.   Blackwood JC, Streicker DG, Altizer S, Rohani P. Resolving the roles of

immunity, pathogenesis, and immigration for rabies persistence in vampire

bats. Proc Natl Acad Sci U S A. 2013 Dec;110(51):20837–42.

9.   Viana M, Benavides JA, Broos A, Loayza DI, Niño R, Bone J, et al. Effects of

culling vampire bats on the spatial spread and spillover of rabies virus. Sci

Adv. 2023;9(10).

10.   O'Shea TJ, Cryan PM, Hayman DTS, Plowright RK, Streicker DG. Multiple

mortality events in bats: A global review. Mamm Rev. 2016;46(3):175–90.

11.   Ham C, Donnelly CA, Astley KL, Jackson SYB, Woodroffe R. Effect of culling

on individual badger *Meles meles* behaviour: Potential implications for bovine

tuberculosis transmission. Journal of Applied Ecology. 2019;56(11):2390–9.

12.   Kays R, Crofoot MC, Jetz W, Wikelski M. Terrestrial animal tracking as an eye

on life and planet. Science (1979). 2015;348(6240):aaa2478.

13.   Britzke ER, Gumbert MW, Hohmann MG. Behavioral response of bats to

passive integrated transponder tag reader arrays placed at cave entrances. J

Fish Wildl Manag. 2014;5(1):146–50.

14.   Thomas B, Holland JD, Minot EO. Wildlife tracking technology options and

cost considerations. Wildlife Research. 2011;38(8):653–63.

15.   Streicker DG, Recuenco S, Valderrama W, Benavides JG, Vargas I, Pacheco

V, et al. Ecological and anthropogenic drivers of rabies exposure in vampire

bats: Implications for transmission and control. Proceedings of the Royal

Society B: Biological Sciences. 2012;279(1742):3384–92.

16.   Rocha F, Ulloa-Stanojlovic FM, Rabaquim VCV, Fadil P, Pompei JC, Brandão

PE, et al. Relations between topography, feeding sites, and foraging behavior

of the vampire bat, *Desmodus rotundus*. J Mammal. 2020;101(1):164–71.

17.   Patterson TA, Thomas L, Wilcox C, Ovaskainen O, Matthiopoulos J. State –

space models of individual animal movement. Trends in Ecology and

Evolution. 2008;(2)23:87-94.

18.   Castle KT, Weller TJ, Cryan PM, Hein CD, Schirmacher MR. Using sutures to

attach miniature tracking tags to small bats for multimonth movement and

behavioral studies. Ecol Evol. 2015;5(14):2980–9.

19.   Aldridge HDJN, Brigham RM. Load Carrying and Maneuverability in an

734   Insectivorous Bat: a Test of the 5% "Rule" of Radio-Telemetry. J Mammal

735   1988 May 20;69(2):379–82. Available from: https://doi.org/10.2307/1381393

736   20.   O'Mara MT, Amorim F, Scacco M, McCracken GF, Safi K, Mata V, et al. Bats

737   use topography and nocturnal updrafts to fly high and fast. Current Biology.

738   2021;31(6):1311-1316.e4.

739   21.   R Core Team. R: A Language and Environment for Statistical Computing.

740   Vienna, Austria: R Foundation for Statistical Computing, Vienna, Austria; 2024.

741   Available from: https://www.r-project.org/

742   22.   Gupte PR, Beardsworth CE, Spiegel O, Lourie E, Toledo S, Nathan R, et al. A

743   guide to pre-processing high-throughput animal tracking data. Journal of

744   Animal Ecology. 2022;91(2):287–307.

745   23.   Pebesma E. Simple Features for R: Standardized Support for Spatial Vector

746   Data. R J. 2018;10(1):439–46.

747   24.   Crespo RF, Linhart SB, Burns RJ, Mitchell GC. Foraging Behavior of the

748   Common Vampire Bat Related to Moonlight. J Mammal. 1972;53(2):366–8.

749   25.   Thieurmel B, Elmarhraoui A. Package "suncalc": Compute Sun Position,

750   Sunlight Phases, Moon Position and Lunar Phase. 2022. Report No.: R

751   package version 0.5.1. Available from: https://cran.r-

752   project.org/package=suncalc

753   26.   Ribeiro R, Matthiopoulos J, Lindgren F, Tello C, Zariquiey CM, Valderrama W,

754   et al. Incorporating environmental heterogeneity and observation effort to

755   predict host distribution and viral spillover from a bat reservoir. Proceedings of

756   the Royal Society B: Biological Sciences. 2023 Jan 1;290(20231739):11.

757   27.   Greenhall AM, Joermann G, Schmidt U. *Desmodus rotundus*. Mammalian

758   Species. 1983;202:1–6.

759   28.   Hijmans R. terra: Spatial Data Analysis. 2024. Available from: https://cran.r-

760   project.org/package=terra

761   29.   Mialhe P. Characterization of *Desmodus rotundus* (E. Geoffroy, 1810)

762   (Chiroptera, Phyllostomidae) shelters in the Municipality of São Pedro - SP.

763   Brazilian Journal of Biology. 2013;73(3):521–6.

764   30.   Delpietro HA, Russo RG, Carter GG, Lord RD, Delpietro GL. Reproductive

765   seasonality, sex ratio and philopatry in Argentina's common vampire bats. R

766   Soc Open Sci. 2017;4(4):1–14.

767   31.   Norberg UM, Rayner JM V. Ecological Morphology and Flight in Bats

768   (Mammalia; Chiroptera): Wing Adaptations, Flight Performance, Foraging

769   Strategy and Echolocation. Philos Trans R Soc Lond B Biol Sci. 1987 Nov

770   20;316(1179):335–427. Available from: http://www.jstor.org/stable/2396486

771   32.   Findley JS, Studier EH, Wilson DE. Morphologic Properties of Bat Wings. J

772   Mammal. 1972 Sep 27;53(3):429–44. Available from:

773   https://doi.org/10.2307/1379035

774   33.   Lehnert LS, Kramer-schadt S, Teige T, Hoffmeister U, Popa-lisseanu A,

775   Bontadina F, et al. Variability and repeatability of noctule bat migration in

776   Central Europe: evidence for partial and differential migration. Proceedings of

777   the Royal Society B: Biological Sciences. 2018;(285).

778   34.   Johnston R, Jones K, Manley D. Confounding and collinearity in regression

779   analysis: A cautionary tale and an alternative procedure, illustrated by studies

780   of British voting behaviour. Qual Quant. 2018;52(4):1957–76.

35. O'Hara RB, Sillanpää MJ. A review of bayesian variable selection methods:

What, how and which. Bayesian Anal. 2009;4(1):85–118.

36. Plummer M. rjags: Bayesian graphical models using MCMC. 2023. Available

from: https://cran.r-project.org/package=rjags

37. Denwood M, Plummer M. runjags: Interface Utilities, Model Templates, Parallel

Computing Methods and Additional Distributions for MCMC Models in JAGS. R

package version 2.2.2-4; 2024. p. 61. Available from: https://github.com/ku-

awdc/runjags

38. Plummer M, Best N, Cowles K, Vines K. CODA: Convergence Diagnosis and

Output Analysis for MCMC. R News. 2006;6:7–11.

39. Wilkinson GS. The social organization of the common vampire bat. Behav Ecol

Sociobiol. 1985;17(2):111–21.

40. Hartman CAR, Wilkinson GS, Razik I, Hamilton IM, Hobson EA, Carter GG.

Hierarchically embedded scales of movement shape the social networks of

vampire bats. Proceedings of the Royal Society B: Biological Sciences.

2024;291(2021):1–12.

41. Streicker DG, Franka R, Jackson FR, Rupprecht CE. Anthropogenic roost

switching and rabies virus dynamics in house-roosting big brown bats. Vector-

Borne and Zoonotic Diseases. 2013;13(7):498–504.

42. Griffiths ME, Broos A, Bergner LM, Meza DK, Suarez NM, Da Silva Filipe A, et

al. Longitudinal deep sequencing informs vector selection and future

deployment strategies for transmissible vaccines. PLoS Biol. 2022;20(4):1–19.

Available from: http://dx.doi.org/10.1371/journal.pbio.3001580

43. Carter G, Leffer L. Social grooming in bats: Are vampire bats exceptional?

PLoS One. 2015;10(10):1–11.

44.   Ripperger SP, Carter GG. Social foraging in vampire bats is predicted by long-

term cooperative relationships. PLoS Biol. 2021;19(9):1–17.

45.   Vargas LÁ. Anticoagulants for the Control of the Common Vampire Bat

(*Desmodus rotundus*). Zoonoses and Public Health. 2025;101–16.

46.   Ávila-Flores R, Bolaina-Badal AL, Gallegos-Ruiz A, Sánchez-Gómez WS. Use

of linear features by the common vampire bat (*Desmodus rotundus*) in a

tropical cattle-ranching landscape. Therya. 2019;10(3):229–34.

47.   Trajano E. Movements of cave bats in southeastern Brazil, with emphasis on

the population ecology of the common vampire bat, *Desmodus rotundus*

(Chiroptera). Biotropica. 1996;28(1):121–9.

48.   Cleasby IR, Owen E, Butler A, Baer J, Blackburn J, Bogdanova MI, et al.

Assessing the importance of individual- and colony-level variation when using

seabird foraging ranges as impact assessment and conservation tools. Ibis.

2023;166(3):871–95.

49.   Greenhall AM. Feeding behavior. In: Schmidt AM, Greenhall RK, editors.

Natural History of Vampire Bats. 1st Editio. Florida: CRC Press Inc, Boca

Raton; 1988. p. 7–17.

50.   Streicker DG, Allgeier JE. Foraging choices of vampire bats in diverse

landscapes: potential implications for land-use change and disease

transmission. Journal of Applied Ecology. 2016;53(4):1280–8.

51.   Hooten MB, Johnson DS, McClintock BT, Morales JM. Animal Movement:

Statistical Models For Telemetry Data. 1st Editio. Florida: CRS Press, Boca

Raton; 2017. 320 p.

52. Kahle D, Wickham H. ggmap: Spatial visualization with ggplot2. R Journal.

2013;5(1):144–61. Available from: http://journal.r-project.org/archive/2013-

1/kahle-wickham.pdf

Figure 1. Study area and GPS data from 55 vampire bats (*Desmodus rotundus*)

from five roosts. (a) Elevation map with the five study roosts (numbered triangles)

in AAC, with an inset map of Peru highlighting the study region. (b) roost 1, (c) roost

2, (d) roost 3, (e) roost 4 and (f) roost 5. In panels b-f, each colour represents a bat

tracked for up to nine nights. Due to the varying scales of movement across the

roosts, this figure does not use a standardised scale, and only 95% of the fixes

closest to the roost were plotted to avoid outliers. Maps used ggmap (52). Base map

tiles show terrain; in panel (c) the map is zoomed in. Data © OpenStreetMap

Figure 2. Predicted probabilities of vampire bats (*Desmodus rotundus*)

returning to the roost following disturbance. Results from a binomial GLM: (a)

Shows sex-specific estimates with 95% confidence intervals; (b) shows the

relationship of probability of return with forearm length (mm), with the 95%

confidence interval shaded.

Figure 3. Predictors of vampire bat (*Desmodus rotundus*) departure time and

distance from the roost. (a) Standardised effect sizes on the log-odds scale

(departure time, logit model) or log scale (distance, exponential model), coloured by

SSVS retention probability. (b) Individual variation in the distance from the roost on

the disturbance night (bat-night interaction). (c) Observed maximum distances (km)

from the roost per bat and night. (d) Predicted maximum distances (km) per bat and

night. In (c) and (d), night 0 is the disturbance night; squares indicate means, and

black dots are outliers.

Figure 4. Reconstructed movement curves for nine vampire bats (*Desmodus rotundus*) across nine nights. First row: bats that travelled longer distances on night 0; middle row: bats that travelled similar distances across different nights; bottom row: bats that left the roost after night 0 and did not return. The shaded area indicates the credible intervals, and the dots the observed values. Purple lines indicate night zero, with progressively lighter colours indicating later nights. Due to the varying scales of movement, the y-axis is not standardised.