This information product has been peer reviewed and approved for publication as a

preprint by the U.S. Geological Survey.

Diverse behavioural responses of vampire bats to human disturbance revealed

by state-space modelling of sparse GPS data.

Rita Ribeiro[1,2], Carlos M. Zariquiey[3,4], Wendi Chavez[4], Kristhie Pillaca Rodriguez[4],

Joaquin Clavijo Manuttupa[4], Marco Antonio Risco Huiman[4,5], Roselvira Zuniga

Villafuerte[4,6], Walter Campana Quintanilla[4], Noelia Medrano-Uchuya[4], Hollie

French[1,2], Jakub Czyzewski[7], Carlos Shiva[8], Nestor Falcon[8], William Valderrama[4,8],

Daniel P. Walsh[9], Tonie E. Rocke[10], Jason Matthiopoulos[1#], Daniel G. Streicker[1,2#]

Joint Senior Authors

[1]School of Biodiversity, One Health and Veterinary Medicine, College of Medical,

Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

[2]Medical Research Council, Centre for Virus Research, University of Glasgow,

Glasgow, UK

[3]Facultad de Salud Pública y Administración, Universidad Peruana Cayetano

Heredia, Lima, Perú

[4]ILLARIY (Asociación para el Desarrollo y Conservación de los Recursos Naturales),

Lima, Perú

[5]Escuela Profesional de Ciencias Biológicas. Facultad de Ciencias, Universidad

Nacional de Piura, Piura, Perú

[6]Centro de Investigación Vertebrate, Universidad Nacional San Antonio Abad del

Cusco, Cusco, Perú

[7]School of Biodiversity, One Health and Veterinary Medicine, BioElectronics Unit,

University of Glasgow, Glasgow, UK

[8]Facultad de Medicina Veterinaria y Zootecnia, Universidad Peruana Cayetano

Heredia, Lima, Perú

[9]U.S. Geological Survey, Montana Cooperative Wildlife Research Unit, Wildlife

Biology Program, University of Montana, Missoula, Montana, USA

[10]U.S. Geological Survey, National Wildlife Health Center, Madison, Wisconsin, USA

Correspondence: Rita Ribeiro, SBOHVM, College of Medical, Veterinary & Life

Sciences, Graham Kerr Building, University of Glasgow, Glasgow G12 8QQ

ritaclaudia.cardosoribeiro@glasgow.ac.uk

Acknowledgements:

We are grateful to Elsa Cardenas Canales, Haris Malik and Carlos Tello for field

assistance; Lynda Dunbar and Marieke Pingen for help with the suturing technique;

Fred Touzalin and Pathtrack for insights and assistance with the GPS. We thank

Holly Niven for helpful discussions about the analysis, and Nardus Mollentze, Travis

Mcdevitt-Galles, and our colleagues from the Streicker group for peer review and

feedback on earlier versions of the manuscript. We thank SENASA (National and

Regional offices) for feedback and support. The project was funded by the

NSF/BBSRC Ecology and Evolution of Infectious Diseases Program (DEB 2011069,

BB/V003798/1). D.G.S. was funded by a Wellcome Trust Senior Research

Fellowship (217221/Z/19/Z). This research received support from the UK Medical

Research Council through core funding of the MRC-University of Glasgow Centre for

Virus Research (Virus Cross Species Transmission Programme: MC_UU_00034/3).

Any use of trade, firm, or product names is for descriptive purposes only and does

not imply endorsement by the U.S. Government.

Author Contributions:

Conceptualization: R.R., J.M., D.G.S. Methodology: R.R., J.C., J.M., D.G.S. Analysis

and visualisation: R.R., J.M. Investigation: R.R., C.M.Z., W.C., K.P.R., J.C.M.,

M.A.R.H., R.Z.V., W.C.Q., N.M-U., H.F., D.G.S. Resources and management:

C.M.Z., C.S., N.F., W.V. Data Curation: R.R. Project administration: D.G.S., Funding

acquisition: D.P.W., T.E.R., J.M., D.G.S. Supervision: R.R., J.M., D.G.S. Writing

(original draft): R.R., J.M., D.G.S. All authors contributed critically to the drafts and

gave final approval for publication.

Statement on inclusion: Our study involved scientists from multiple countries,

including those based in the study location, fostering diversity and inclusion from the

project's outset. Co-authors were actively involved in project design, infrastructure,

management, investigation, and manuscript preparation.

Data availability: Data and code for this study are deposited in the Dryad Digital

Repository and will be made publicly available upon publication, at which point the

corresponding DOI will be shared.

ABSTRACT

1. Human activities can alter animal movement, having unexpected consequences for disease transmission. In Latin America, culls of common vampire bats (*Desmodus rotundus*) for rabies control have been indicated to raise viral spatial spread by increasing long-range movement and inter-colony mixing among surviving bats, but behavioural responses to disturbance have not been quantified.

2. We used Global Positioning System (GPS) tags to track the movements of 93 vampire bats from 5 roosts in the Peruvian Andes over 9 nights after human disturbance (capture, handling and tag attachment). Responses to disturbance were modelled under a Bayesian state-space framework that estimated bat departure time, activity duration, and travel distances while accounting for the sparse GPS data obtainable in a topographically complex landscape where vampire bats occur.

3. A large fraction of bats (43%, 39/90) abandoned the study area on the night of disturbance, with more abandonment observed in groups expected to have greater dispersal capacity (males and bats with longer forearms). Among the remaining bats, we observed high individual heterogeneity on the night of disturbance, with some bats staying atypically close to the roost and others travelling unusually long distances. With increasing time post-disturbance, bats tended to remain closer to the roost.

4. Analysis of inferred flight histories showed that bats travelled >5 km on most nights, with high variability both within individuals across nights and between individuals in the same roost.

5. By inferring movement from irregular and error-prone data, our model extends the utility of animal-borne GPS tracking to landscapes with poor satellite visibility,

revealing greater complexity in vampire bat movement than previously

recognised. Vampire bat disturbance was associated with three distinct

behavioural responses: roost abandonment, increased flight distances and

extreme individual heterogeneity on the night of disturbance. In the context of

vampire bat management, such perturbations would be expected to exacerbate

the spatial spread of rabies and potentially undermine the efficacy of culling

activities which are currently practiced by most Latin American governments.

Key words: Bayesian integrated model, disease control measures, JAGS,

perturbation effect, travel distance, wildlife management, wildlife telemetry, zoonoses

1| INTRODUCTION

Human-induced changes in animal movement can influence disease dynamics

and increase spillover risk to humans or domestic animals by forcing the mixing of

infected and uninfected populations, potentially increasing the incidence and spatial

extent of pathogens (Blackwood et al., 2013). For example, in Brazil, urbanization

and habitat loss forced non-human primates into cities potentially increasing human

exposure to yellow fever virus (Couto-Lima et al., 2017). In Europe, badger culling

for managing bovine tuberculosis increased survivors' ranging and dispersal

behaviours, likely explaining increased tuberculosis risk in cattle outside of culled

areas (Ham et al., 2019). Independently, disturbance may induce chronic stressors

that diminish immunity and increase infection rate and intensity (Becker et al., 2018;

Seltmann et al., 2017).

Bats are important reservoirs of zoonotic viruses including Nipah virus

(Paramyxoviridae, *Henipavirus nipahense*), Hendra virus (Paramyxoviridae,

Henipavirus hendraense), Marburg virus (Filoviridae, Orthomarburgvirus), and

various Lyssaviruses (Calisher et al., 2006). The high mobility of bats arising from

their capacity for flight, together with their high sensitivity to human

disturbance - including occasional, deliberate human persecution - may

inadvertently increase human disease risk (O'Shea et al., 2016). Indeed, bat

disturbance has been linked to increases in the spread or incidence of both

rabies virus and Marburg virus (Amman et al., 2014; Streicker et al., 2012).

Despite the potential links between zoonotic risk and bat movement, few

studies have quantified how human disturbance affects bat movement, likely

due to challenges posed by bats' cryptic lifestyles and small size, which

restricts available tracking technologies (O'Mara et al., 2014). One study using

radiotelemetry revealed that disturbance from removal of big brown bats

(*Eptesicus fuscus*) from houses induced roost switching, potentially

exacerbating rabies virus transmission (Streicker et al., 2013). Another study

using Global Positioning System (GPS) tags in straw-coloured fruit bats

(*Eidolon helvum*) showed that disturbance from human capture led to

increased daytime movements (Costa et al., 2020). While these studies

suggest that human disturbance can affect bat behaviour, reactions and

intensity of responses may vary across species and types of disturbance.

In Latin America, *Desmodus rotundus* (the common vampire bat) is the

main reservoir of rabies virus (Rhabdoviridae, Lyssavirus), which causes

sporadic lethal outbreaks in humans and a sustained agricultural burden

through livestock mortality (Schneider et al., 2009). This species is routinely

disturbed during campaigns aimed at controlling vampire bat-transmitted

rabies (VBR). Sanctioned population control (culling) uses anticoagulant

poisons ('vampiricides') that can be topically applied to bats or injected into livestock

for later consumption by bats during blood feeding (World Health Organization,

2018). However, unsanctioned efforts to reduce vampire bat populations are

common and include direct killing of captured bats and destruction, sealing or

burning of roosts (O'Shea et al., 2016). Despite decades of culling, high VBR

burdens persist in enzootic areas and spatiotemporal analyses show viral

expansions into historically rabies-free regions of Latin America (Benavides et al.,

2016; Rojas-Sereno et al., 2022). Epidemiological models explaining the limited

efficacy of culls indicated that increased bat mixing after disturbance could raise

VBR incidence and local spillovers (Blackwood et al., 2013). Consistent with this

idea, evaluation of a large culling campaign in Peru showed that culling did not limit

rabies spillover to livestock, and culling in response to livestock rabies cases

(reactive culling) increased viral spatial spread, an effect that was hypothesized to

arise from increased movement of rabies-incubating bats (Viana et al., 2023). While

these observational studies suggest that culling affects bat movement with

downstream repercussions for rabies transmission, the behavioural responses of

vampire bats to disturbance remains unexplored.

Directly measuring the fine-scale behavioural response of vampire bats to

human disturbances is hampered by technical and ecological challenges. In mark-

recapture studies, low between-roost recapture rates limit inference on movement

(Britzke et al., 2014). Very High Frequency (VHF)-radio tracking has also been

applied, but it is labour-intensive, error-prone and only detects animal presence in

close proximity to antennas (Thomas et al., 2011). Use in vampire bats has thus

been limited to coarse-scale studies at foraging sites and identification of unknown

roosts (Rocha et al., 2020; Streicker et al., 2012). Recent advances in GPS allow for

the monitoring of animal movements at high spatial and temporal resolutions

with lightweight tags equipped for remote data recovery, making them suitable

for small animals such as bats (Kays et al., 2015). However, the reliance of

GPS on satellite visibility introduces challenges in complex landscapes,

including highly mountainous or densely forested areas. Indeed, to our

knowledge, no GPS study on vampire bats has been reported.

Here, we used lightweight GPS tags to assess fine-scale movement of

vampire bats in the Peruvian Andes in response to human disturbance,

developing a new method to reconstruct movement despite the sparse and

error-prone data expected from the obstructive local topography. We jointly

reconstructed and identified ecological correlates of two fundamental metrics

of movement: nightly departure time and distance from the roost at different

times of the night. We hypothesised that bats would leave the roost under

conditions of darkness at night, and that disturbance from capture, handling,

and tagging, would induce roost abandonment and longer flights.

2| MATERIALS AND METHODS

2.1| Study area and design

The study was conducted between April 2022 and June 2023 in 5

vampire bat roosts, located >51.6 km apart (range 51.6-186 km). Roosts are

in a region in southern Peru (departments of Ayacucho, Apurimac, and

Cusco, AAC), that is characterised by rugged topography, with altitudes

ranging from 1200 to 6000 meters (Fig. 1a). AAC is a hotspot of rabies in

Peru, accounting for 60.2% of all reported livestock cases between 2003 and

2019, and this region is subjected to regular bat culls (Viana et al., 2023).

Vampire bats live below 3600 meters, roosting predominantly in natural caves,

abandoned houses, mines, and irrigation canals (Benavides et al., 2016, 2020).

2.2| Data collection

Vampire bats were captured and handled under approved ethics protocols

(University of Glasgow, EA28a/21) and permits from local authorities, including

SERFOR (Servicio Nacional Forestal y de Fauna Silvestre, Nos. D000024-2022,

D000107-2022, D000103-2024 and D000188-2024-MIDAGRI-SERFOR-

DGGSPFFS-DGSPFS) and SERNANP (Servicio Nacional de Áreas Naturales

Protegidas por el Estado, N° 021-2023-SERNANP-SHM/J).

Two capture and tagging sessions per roost were carried out from April to

October (the Peruvian dry season), with at least three months between sessions.

Roost 2 (Fig. 1c) had only one tagging session due to data recovery challenges.

Each session aimed to tag 10 adult vampire bats (5 males, 5 females). All capture

sessions began after dusk, except the first capture in April 2022, when due to

logistical constraints, captures began in the morning. Vampire bats were captured

using mist and butterfly nets. Captured bats were fit with a 3.5 mm wing ring

(Porzana, UK), and weight (g), sex, reproductive status (non-reproductive, scrotal

male, pregnant, or lactating female), age (adult, subadult, or juvenile), and forearm

length (mm) were recorded. We applied lightweight GPS tags (2.8 g, 8 cm length

[including a 5 cm antenna]; nanoFix®, PathTrack, UK) to bats that met specific

inclusion criteria (see below). After light sedation using intramuscular (pectoral)

injection (29-gauge needle, 1.0 mL syringe) of Ketamine (20–25 mg/kg) diluted in

saline (NaCl) at a 50/50 volume ratio or Midazolam (0.25 mg/kg, used in 2023 to

allow faster recovery times), GPS tags were attached to the skin of the dorsal side of

each bat using an absorbable suture (Monosyn®, B Braun), threaded through

custom tag loops (Castle et al., 2015). According to the manufacturer, the absorbable

sutures lose 50% of their strength after 14 days, allowing natural tag

detachment. Prior to suturing, hair was trimmed, and a small amount of

cyanoacrylate-based glue was used to secure the tags. Sedation and tag

suturing were carried out by a veterinarian, ensuring it didn't impair flight or

walking. GPS tags were painted with UV fluorescent paint to aid in recovery

for reuse (Starglow®, Glowtec, UK).

GPS tagging was limited to healthy adult bats, excluding pregnant and

lactating females. The average weight of tagged bats was 43.4 g ($\pm$ 3.4 g,

N=93), so the tags represented a maximum of 7% of the bat's weight (mean =

6.5 $\pm$ 0.48%, N=93). Recommendations for bat tracking studies indicate that

attached devices should not exceed 10% of body mass, with a best practice of

limiting to 5%. However, heavier tags are acceptable for short-term use in

some species (Aldridge & Brigham, 1988; O'Mara et al., 2014). From our prior

fieldwork, adult vampire bats (N=3,429) weigh, on average 41.9 g (± 8.44,

N=2,751) and 51.5 g (± 9.1, N=678) after feeding, indicating nightly weight

gains of over 9 g. Additionally, during pregnancy and after birth, females can

weigh over 85 g and continue foraging. These observations indicate that the

tags would be tolerable for short-term use.

During the initial deployment in April 2022, GPS tags were programmed

to record locations every 5 minutes (10 tags) or 10 minutes (5 tags) for 3 and

6 nights, respectively. To extend tracking duration, later deployments used a

15-minute interval for 9 nights. Our model (see below) considered this

difference between deployments and reconstructed distances from the roost

every 15 minutes. Tags operated nightly from 6 PM to 6 AM, with data stored

onboard and downloaded remotely via one or two solar-powered UHF base

stations (PathTrack, UK) placed near the roost.

2.3| Data cleaning

Data cleaning was performed in R version 4.4.0 (R Core Team, 2024). The

nanoFix GPS tags require at least 4 visible satellites to record a position ('fix'). The

number of visible satellites, along with an accuracy indicator for high satellite counts,

determines the positional accuracy. Due to the complex topography of the Peruvian

Andes, most GPS points were missing since <4 satellites were in range at the

attempted connection time, resulting in a sparse and irregular dataset that required

the model framework outlined below to reconstruct movement.

GPS fixes recorded before tag application and 1 hour after capture were

removed to account for the tag attachment process. Data from the first night were

otherwise included because quantifying the disturbance immediately after capture

was one of our objectives. Cleaning of the remained data excluded: one inaccurate

fix (6 satellites connected, but accuracy indicator >30, considered unreliable), all

fixes from one bat with a malfunctioning tag (distances up to 6 km recorded before

attachment), all data from 3 bats with <3 fixes across the study period, and all data

from an unrealistically sessile bat (suggesting tag loss or death). As detailed in the

supplementary information, we also removed GPS fix jumps, and outliers of

estimated flight speed (Gupte et al., 2022). The speed (km/hour) of each segment

between subsequent fixes was calculated, along with the Euclidean distance from

the roost for each fix, using base R and the sf package (v1.0-19) (Pebesma, 2018).

Our state-space model was designed to be robust to GPS errors, so we could have

omitted data filtering based on speed entirely. However, we anticipated that

simultaneously inferring and filtering speed within the model could hinder

convergence.

2.3.1| Variable definition and processing

We investigated predictors of bats' nightly departure time and distance

travelled from the roost. Vampire bats are thought to be lunar phobic,

departing before moonrise or after moonset (Crespo et al., 1972). We

therefore modelled departure time as a function of the number of hours until

complete darkness, which depended on the interplay of moonrise, moonset,

astronomical night start (occurs after sunset and dusk) and end (occurs before

dawn). For each roost and night, we calculated the hours from sunset until

darkness using the *suncalc* package (v0.5.1) (Thieurmel & Elmarhraoui,

2022). We predicted that longer periods until darkness delay departure times.

We also hypothesised that sex influences departure time, as female vampire

bats, which maintain cooperative relationships that include behaviours such as

visiting other roosts, might leave earlier in the night (Ripperger & Carter, 2021;

Wilkinson, 1985).

Distance from the roost was modelled with variables related to roost

location, individual bat characteristics, and human disturbance (Table S1).

Roost-related variables within 5 km radii of roosts included proportion of

permanent water, predicted vampire bat roost density (Ribeiro et al., 2023)

and cattle density (Greenhall et al., 1983), calculated using the packages *sf*

(v1.0-19) and *terra* (v1.7-71) (Hijmans, 2024; Pebesma, 2018). We predicted

that the availability of alternative roosts and preferred food sources (cattle)

near the roost would reduce the distance bats travel. Since cattle graze near water

and vampire bats can use rivers as corridors, we hypothesized that permanent water

sources near roosts could aid prey searching, decreasing distances from the roost

(Mialhe, 2013). Individual characteristics included bat sex and forearm length. We

hypothesised that males would fly further distances due to reduced offspring care

(Delpietro et al., 2017). As flight speed is correlated with forearm length, we

expected a positive effect of bat forearm length on distance from the roost (Hopkins

et al., 2003).

Human disturbance was defined as the effects of capture, handling, and

tagging, with key variables being the night of capture ('night 0') versus subsequent

nights, and the time (in hours) post-disturbance, calculated as the difference

between the start of the tracking session and each GPS fix. We averaged time post-

disturbance per bat and night to summarise nightly patterns, while reducing the

effect of the irregular GPS fix intervals and potential autocorrelation. This also aligns

with our model structure, which accounts for individual, roost and night effects. We

anticipated that disturbance would increase flight distance following capture, with

distances decreasing over time as behaviour normalised. Finally, because vampire

bats in our study area are deliberately or accidentally disturbed by humans (e.g.,

capture, culling, roost destruction or shifts in agricultural practices), we also included

a variable describing the number of rural human settlements within 5 km of each

roost as a proxy for historical disturbance levels.

2.3.2| Modelling framework

We developed a Bayesian state-space model with two latent variables: the

bats' departure time (t, in minutes since the daily activation of the GPS at 6 PM) and

its distance from the roost (ψ, in km). The model reconstructed distances from

the roost every 15 minutes and estimated the duration of activity and return

time to the roost, accounting for both process error (i.e., the natural

stochasticity in the biological processes) and observation error (i.e., GPS

measurement error). The model assumed nightly returns to the roost.

Although our data indicated that returns occurred on at least 56.6% of bats

and nights, this is likely a conservative estimate due to data limitations –

including missed fixes during 15-minute intervals when bats departed or

returned quickly, and signal loss at the roost – potentially leading to

underestimation of return rates. We also assumed that most bats would depart

before 1 AM, as our experiments were conducted on nights when the moon

set before 1 AM in 95% of cases.

For each individual bat i and night j, $t_{i,j}$ is the departure time in minutes

after 6 PM, with a maximum value of 420 minutes (1 AM). The expected and

realised departure instances, $\bar{\tau}_{i,j}$ and $\tau_{i,j}$, are expressed as a proportion of the

interval between 6 PM and 1 AM,

(1) $\tau_{i,j} = \dfrac{t_{i,j}}{420}$,

(2) $t_{i,j} = 420\tau_{i,j}$.

The expected departure instance $\bar{\tau}_{i,j}$ was modelled as a logit function of the number

of hours until darkness and bat sex,

(3) $logit(\bar{\tau}_{i,j}) = \theta_0 + \theta_1 hours\_dark_j + \theta_2 sex_i$.

$\tau_{i,j}$ was modelled by a Beta distribution, with mean $\bar{\tau}_{i,j}$ and a process error,

quantified by the standard deviation of departure instance, σ (Table S2), capturing

pooled inter-trip and inter-individual variation,

(4) $\tau_{i,j} \sim Beta(a_{i,j}, b_{i,j})$,

(5) $a_{i,j} = \bar{\tau}_{i,j}(\frac{\bar{\tau}_{i,j}(1-\bar{\tau}_{i,j})}{\sigma^2} - 1)$,

(6) $b_{i,j} = (1 - \bar{\tau}_{i,j})(\frac{\bar{\tau}_{i,j}(1-\bar{\tau}_{i,j})}{\sigma^2} - 1)$.

The duration of nightly activity after departure was modelled with a uniform

distribution, assuming that once a bat departed, it remained away for at least 20

minutes. The maximum duration of a trip was the time left after departing and the

end of GPS activation (720 minutes),

(7) $\Delta t_{i,j} \sim U(20, 720 - t_{i,j})$.

To estimate distances travelled from the roost, we defined a parametric model

of distance against time, for each individual bat i, night j, and each block k of 15

minutes after 6 PM,

(8) $\bar{\psi}_{i,j,k} = \exp(\beta_0 - \frac{((k15-t_{i,j})-\Delta t_{i,j}\tau_{mid,i,j})^2}{(\Delta t_{i,j}\sigma_{mid,i,j})^2} + \beta_i night_j + \sum_{n=1}^{N} B_n X_{i,j,n} + \zeta_i)$.

This equation assumes that only one trip is made per night and is flexible regarding

the point of start, end, and maximum distance. It also assumes that the net velocity

is away from the roost until the maximum distance is reached, and then toward the

roost thereafter, for 15-minute intervals. The exponential of the intercept β_0,

corresponds to the maximum distance from the roost. β_i corresponds to the terms

for the effect of night of disturbance ($night_j$) for each bat, and B_n is the parameter for

the n^{th} covariate effect (forearm length, sex, roost density, number of human rural

settlements, cattle density, permanent water and time post-disturbance). We added

an individual-level random effect ζ_i with mean 0 and standard deviation σ_1 ($\zeta_i =$

$\mathcal{N}(0, \sigma_1)$), to capture the influence of individual variation (Table S2). In both

departure time and distance models, we considered random effects of individual,

roost, calendar night, and session of capture. Only the random effect of individual on

the distance model was relevant, hence it was kept.

The mid-point of the trip, $\tau_{mid,i,j}$, was defined as the proportion of time

between departure and return until reaching maximum distance, with a

parameter $\sigma_{mid,i,j}$, which made sure that the distance from the roost climbed

smoothly from the start of the trip to the midpoint and then declined to the

trip's endpoint, preventing unrealistic jumps in movement. $\tau_{mid,i,j}$ was given a

beta prior distribution and was set to be close to the middle of the trip duration

(mean 0.5 and variance 0.001). $\sigma_{mid,i,j}$ was fixed to a sufficiently large value to

ensure that the bats did not immediately reach their maximum distance after

departure or instantaneously returned to the roost,

(9) $\sigma_{mid,i,j} = \max(\tau_{mid,i,j}, 1 - \tau_{mid,i,j})/2.$

The realisation of distance giving rise to observations, $\psi_{i,j,k}$, was modelled using a

normal distribution, with mean $\bar{\psi}_{i,j,k}$ and variance $O_{i,j,k}$,

(10) $\psi_{i,j,k} \sim \mathcal{N}(\bar{\psi}_{i,j,k}, O_{i,j,k})$, with $\psi_{i,j,k} \geq 0$ km.

$O_{i,j,k}$ is a combined error, that incorporates observation error $\xi_{i,j,k}$, the standard

deviation of GPS error, and process error, in the form of an overdispersion

parameter φ, to account for further uncertainty that cannot be explained by the

observation model (i.e., 0.6 km is the GPS error for 4 satellites),

(11) $O_{i,j,k} = \xi_{i,j,k}^2 + 0.6\varphi^2.$

The overdispersion parameter φ was estimated by the model, but we encouraged it

to be close to 0 (Table S2) to prioritize the identification of biological effects. Using

data from a GPS tag placed at a single location in our study area, we fitted a

regression of the standard deviation of distances against the number of satellites and

estimated the standard deviation of GPS error for each satellite count (standard

deviation of 0.3 km when 4 satellites were visible; deviance explained = 0.986). This

regression produced the two parameters γ_0 and γ_1 which, in combination with the

number of visible satellites that generated each positional fix, $N_{i,j,k}$, informed $\xi_{i,j,k}$,

(12) $\xi_{i,j,k} = \exp(\gamma_1 + \gamma_2 N_{i,j,k})$.

We used Bayesian Stochastic Search Variable Selection (SSVS) with spike

and slab priors to estimate model coefficients conditional to a variable being retained

in the model (O'Hara & Sillanpää, 2009). If a variable was participating in the model,

the coefficient value was given a normal distribution, with mean 0 and standard

deviation τ_1, with τ_1 being estimated by the model, except for the coefficient for

hours until darkness, which was given a gamma prior distribution to preserve its

positivity (mean 0.01 and variance 0.0001). The posterior for the probability of

retention ρ was modelled with a Bernoulli distribution (variable included or excluded),

with the parameter ρ being estimated by the model with a constrained Beta prior

(i.e., a variable has a 25% probability of being included), meaning that variables

need strong evidence to be included.

To assess goodness of fit, we calculated the R^2 by comparing the observed

and predicted distances from the roost. The models were fitted in JAGS 4.3.2 using

rjags (v4-15) and runjags (v2.2.2-4) (Denwood & Plummer, 2024; Plummer, 2023).

The model ran with four independent chains, for 10,000 samples per chain after

discarding the first 500,000 interactions as burn-in, with thinning of 200 (retaining

every 200th iteration). Convergence across chains was assessed through visual

inspections of the posterior and the Gelman-Rubin diagnostic test (using coda

(v0.19-4.1) (Plummer et al., 2006)). Table S2 lists the final model parameter priors

and posteriors.

2.3.2.1| Drivers of roost abandonment

We fitted binomial generalised linear models (GLM) to examine whether roost abandonment was associated with bat individual traits (sex, forearm length, and weight), hypothesising that stronger bats (i.e., males, heavier, and bats with longer forearms) would be less likely to return. Model performance was compared across six models using the Akaike information criterion (AIC) and pseudo-R^2.

3| RESULTS

Between April 2022 and June 2023, 93 adult vampire bats were GPS tagged across 5 roosts in southern Peru (Table S3). Twenty-four percent of tags (20/83) were recovered after use, most of them found inside roosts, allowing partial redeployment. Among the 60 bats with recovered GPS data, 89% of possible observations (14,142 out of 15,842) were not recorded due to having <4 visible satellites, leaving a total of 1700 GPS fixes. The night of capture ('night 0') had the most fixes (24.7%), followed by 'night 1' after capture (18.2%), 'night 2' (17.9%), 'night 3' (14.1%), 'night 4' (11.1%), 'night 5' (7.7%), 'night 6' (4.5%), 'night 7' (1.2%) and 'night 8' (0.6%). Data cleaning excluded a further 22.3% of the fixes, leaving 1321 fixes from 55 bats (Fig. S1). Visualization of the GPS data showed that vampire bats occasionally flew to distances >5km from the roost but spent most time near roosts (Fig. 1b-f; Fig.S2). Individual bats from the same roost foraged in different areas, did not visit the same locations every night, and used a variety of flight paths,

highlighting great behavioural variation among and within individuals (Fig. 1b, d, e, f;

Fig. S3d, e).

Filtering out 3 bats which were confirmed dead, we recovered no GPS data

from 30 bats and 9 additional bats were not detected after night 0, indicating the

possibility that 43% (39/90) of bats abandoned our study areas following our capture

and tagging effort. While we cannot rule out tag loss or death of these missing

animals in unobservable locations within our study area, recapture data point to long-

or medium-term roost abandonment. Of the 51 bats which returned to the roost

according to GPS records, 15.7% were recaptured during capture sessions

immediately following the tracking period. In contrast, of the 39 missing bats, we

recaptured only 1 bat (2.6%) from roost 2 (where we faced challenges in recovering

data), while none from other roosts were recaptured. As missing bats would be

expected to have similar recapture rates if their apparent disappearance from the

study area was an artefact of GPS tag loss or failure, we would have expected at

least 6 recaptures. Moreover, during surveys conducted between 4-32 months after

GPS tagging, 5 of the missing bats (12.8%) were recaptured. Comparable long-term

recapture rates between bats that disappeared from the study area and those that

remained argue against disappearance being due to mortality.

A binomial GLM revealed that roost abandonment was non-random and

consistent with expectations of vampire bat dispersal biology. Bats with longer

forearms (-0.96 [SE = 0.35], P<0.01) and male bats (-1.11 [SE = 0.66], P=0.09) were

less likely to return to the roost, although there was only weak evidence for the latter

effect (Fig. 2; Table S4; Fig. S4). These findings strongly support considerable short-

term roost abandonment by vampire bats following disturbance. Complementing this,

the GPS data also revealed high variation in movement on the night of

disturbance across all roosts except roost 2, which had data from only one bat

(Fig. S5).

By fitting a Bayesian state-space model to the GPS data, we explored

correlates of departure time and maximum nightly distance from the roost. The

model gave a good fit to the data with an R^2 of 0.80 (Fig. S6), estimating that

bats typically departed ~2 hours after the GPS started at 6 PM (posterior

mean of 124 minutes, 95% credible interval (CI) [58.3; 220.5]) and were out of

the roost for <6 hours (345 minutes, 95% CI [190.0; 497.4]) (Fig. S7). The

number of hours until darkness had a small but significant effect on departure

time, with bats leaving the roost as soon as there was no sunlight or moonlight

(Fig. 3a; Table S5). Our model, which reconstructs the movement curve for

each bat, filling gaps due to missing observations, predicted larger distances

from the roost compared to observed data (Fig 3c-d). Within the distance

component of the model, time post-disturbance had a negative effect on

maximum nightly distance travelled from the roost (retention probability

[P]=0.95) (Fig. 3a, d; Table S5; Figs. S8-S9). This result indicates that bats

tended to fly further away after disturbance but then returned to their baseline

distances, with a decrease in maximum nightly distance of 0.17 km per night

(Fig. S8). The interaction between bat and night highlighted high heterogeneity

among individuals, particularly on the night of the disturbance, when some

bats stayed unusually close to the roost and others undertook the longest

movements recorded in the study period (Figs. 3b, 4; Table S6; Figs. S10-

S11).

4| DISCUSSION

Behavioural responses of wildlife to human disturbances can undermine

efforts to prevent disease spillover but fine-scale studies are currently restricted to

relatively large-bodied species and in simple habitats. By developing a Bayesian

state-space model to reconstruct movement histories from sparse and error-prone

GPS data from 93 vampire bats in the Peruvian Andes, we identified three

behavioural responses to human disturbance which could exacerbate rabies

transmission and dilute the effectiveness of bat population control efforts seeking to

prevent bat bites and rabies spillover.

First, we found that 43% of the GPS-tagged bats appeared to abandon the

roost, an effect that was unlikely to be explained by bat mortality or GPS failure.

Since return to the vicinity of the roost at any point in the study would have triggered

data upload to our receivers, we believe this indicates that bats left the study area, at

least for the 9-day duration of the study, though their destination is unknown. Given

that vampire bat dispersal is generally male-biased and that forearm length is linked

to flight ability, the finding of putatively stronger flyers more frequently disappearing

after tagging further supports transient roost abandonment (Delpietro et al., 2017;

Streicker et al., 2016). Although we cannot rule out the possibility that tags detached

in areas with poor satellite coverage, the small number of stationary tags detected,

and the absence of GPS fixes prior to this putative detachment makes this

explanation unlikely. We observed two other behavioural responses among bats that

remained detectable: a general tendency for longer distance movements after

disturbance and elevated individual heterogeneity on the night of capture.

Collectively, these three perturbations could have unintended consequences

for both the transmission and management of VBR. Roost switching following the

application of vampiricide poison would dilute local population reduction and

may potentially disperse bats to lower density roosts or roosts where bats

have fewer social connections, reducing the efficiency of poison transfer.

Similarly, increased dispersal of rabies-positive vampire bats following

disturbance is predicted to raise rabies incidence in bats from nearby roosts,

potentially explaining findings why reactive culls increased livestock rabies risk

in adjacent areas (Blackwood et al., 2013; Viana et al., 2023). Since viral

spatial spread is believed to be strongly influenced by male bat movement, the

tendency for male bats to abandon roosts after capture would be particularly

problematic (Streicker et al., 2016). For future research, the effects of

disturbance that we report are consistent with those from other mammals,

indicating that studies seeking to understand the natural, post-disturbance

behaviour of vampire bats using GPS tags should delay data recording for 4

nights (Stiegler et al., 2024).

A caveat when drawing conclusions about culling is that our study

measured the effects of a relatively mild disturbance. While our disturbance

shares similarities with culling - capture, handling, the application of a foreign

object, and release - there are key differences. Vampire bats have strong

cooperative networks, characterised by blood meal sharing and intense

allogrooming, which could be disrupted by the sickness behaviours and

mortality associated with poisoning (Carter & Leffer, 2015; Ripperger & Carter,

2021). As such, culls may lead to stronger effects on movement than the

relatively mild disturbance we induced. Capture locations may also vary

between our study and real-world management. While we carried out captures

in roosts for greater tractability, this form of capture accounts for only 26% of

culling events, with captures at properties where bats attack livestock being more

common (Viana et al., 2023). Additional field experiments that separate effects of

human presence at roost sites, capture, handling and poisoning, could clarify

vampire bat population management.

High resolution GPS tracking also revealed that constraints on bat movement

and VBR spread may be less restrictive than implied by previous observational or

radiotelemetry data. For example, rather than consistently foraging downstream of

their roosts, following valleys or rivers we observed vampire bats travelling over

mountains and foraging in both directions along rivers (Ávila-Flores et al., 2019;

Trajano, 1996). Furthermore, similar to other colonial species such as seabirds,

vampire bats exhibited high movement behavioural heterogeneity within the colony

(Cleasby et al., 2023). Our data also challenge the notion that vampire bats return to

the same foraging location and even individual animals for blood feeding every night

(Greenhall, 1988). Overall, heterogeneity across individual bats and within bats over

short time periods helps explain the sometimes-sporadic evidence of vampire bat

bites on livestock and humans (Streicker & Allgeier, 2016) and implies that

pathogen-infected bats are likely to expose multiple animals at different locations.

Incorporating these behavioural ecological insights on vampire bats into models of

rabies spread could enhance our understanding of its dynamics.

Our study fills an analytical gap that has arisen between the declining size and

increasing battery life of GPS tags - which have expanded their applicability in

wildlife research - and the problematic data generated when such studies are

conducted in challenging habitats (D'Eon et al., 2002). The Bayesian framework,

which jointly estimated departure times and distances from a fixed starting point (the

roost), enabled inferences from sparse and irregular GPS data. While our objective

was not intended to infer exact animal locations throughout the night, it did

allow us to estimate how individual, landscape, disturbance and temporal

factors affected key characteristics of bat movement while accounting for

uncertainty. Furthermore, this approach enabled us to estimate the true

distances bats can travel - that are critical for disease management – but that

are often missed in sparse datasets, which tend to underrepresent maximum

distances (Hooten et al., 2017). By enabling inference from such a challenging

dataset, our approach extends the use of GPS tracking to other species and

environmental conditions where conventional analyses may fail or be strongly

biased by satellite visibility constraints.

A potential limitation of our approach are the assumptions that vampire

bats do a single trip and return to the roost every night. If some long-distance

movements instead reflect roost switching, our results may overestimate true

foraging distances; however, we note that eventual return to the roost was

required for data recovery so either behaviour reflects short term movements

after disturbance. The assumption of a single trip per night was broadly

supported by overall model fit and parameter convergence. However, a subset

of bat-nights (13%) showed non-convergence, resulting in a bimodal

distribution more consistent with multiple trips within a night (Fig. S10).

However, our model can identify such misspecifications and highlight that

vampire bats may undertake multiple trips per night. The significance of

multiple trips from a rabies perspective will depend on whether trips represent

scouting activities that inform foraging activities (e.g. evaluating lunar

brightness) or multiple bouts of foraging which would increase rabies

exposures in prey. Empirical data on vampire bat activity, roost-switching and

abandonment rates through automated radio frequency identification (RFID) systems

would be useful to validate our interpretation.

Understanding animal movement in response to disturbance is important to

design effective strategies for disease management but conventionally has been

frustrated in some species and environments due to sparse data and error in

movement data. This study provides evidence to support a change in the

management of VBR. By enabling simultaneous inference and predictors of nightly

movement curves for individual vampire bats, we show three responses to

disturbance (increase in maximum distance travelled, individual heterogeneity in

movement, and roost abandonment) which are compatible with the hypothesis that

disturbing vampire bats by culling may trigger them to abandon the roost or to fly

further, potentially promoting viral spread. Increased movement and roost

abandonment immediately following disturbance from culling would also be expected

to dilute the distribution of poison across roosts, potentially reducing efficacy. In

contrast, disturbance induced movement could benefit the recently proposed

application of vaccines with the capacity for self-dissemination since roost switching

would 'seed' vaccine transmission to a larger number of roosts than previously

appreciated (Griffiths et al., 2022). Most generally, our study highlights the value and

analytical challenges of incorporating new data streams provided by increasingly

accessible field-deployable technologies into wildlife ecology and management.

REFERENCES

Aldridge, H. D. J. N., & Brigham, R. M. (1988). Load Carrying and Maneuverability in

an Insectivorous Bat: a Test of the 5% "Rule" of Radio-Telemetry. *Journal of*

Mammalogy, *69*(2), 379–382. https://doi.org/10.2307/1381393

Amman, B. R., Nyakarahuka, L., McElroy, A. K., Dodd, K. A., Sealy, T. K., Schuh, A.

J., Shoemaker, T. R., Balinandi, S., Atimnedi, P., Kaboyo, W., Nichol, S. T., &

Towner, J. S. (2014). Marburgvirus resurgence in Kitaka mine bat population

after extermination attempts, Uganda. *Emerging Infectious Diseases*, *20*(10),

1761–1764. https://doi.org/10.3201/eid2010.140696

Ávila-Flores, R., Bolaina-Badal, A. L., Gallegos-Ruiz, A., & Sánchez-Gómez, W. S.

(2019). Use of linear features by the common vampire bat (Desmodus rotundus)

in a tropical cattle-ranching landscape. *Therya*, *10*(3), 229–234.

https://doi.org/10.12933/therya-19-890

Becker, D. J., Hall, R. J., Forbes, K. M., Plowright, R. K., & Altizer, S. (2018).

Anthropogenic resource subsidies and host –parasite dynamics in wildlife.

Philosophical Transactions of the Royal Society B: Biological Sciences,

373(1745). https://doi.org/10.1098/rstb.2017.0086

Benavides, J. A., Valderrama, W., Recuenco, S., Uieda, W., Suzán, G., Avila-Flores,

R., Velasco-Villa, A., Almeida, M., de Andrade, F. A. G., Molina-Flores, B.,

Vigilato, M. A. N., Pompei, J. C. A., Tizzani, P., Carrera, J. E., Ibanez, D., &

Streicker, D. G. (2020). Defining New Pathways to Manage the Ongoing

Emergence of Bat Rabies in Latin America. *Viruses*, *12*(9), 1–13.

https://doi.org/10.3390/v12091002

Benavides, J. A., Valderrama, W., & Streicker, D. G. (2016). Spatial expansions and

travelling waves of rabies in vampire bats. *Proceedings of the Royal Society B:*

Biological Sciences, *283*(1832), 1–9. https://doi.org/10.1098/rspb.2016.0328

Blackwood, J. C., Streicker, D. G., Altizer, S., & Rohani, P. (2013). Resolving the

roles of immunity, pathogenesis, and immigration for rabies persistence in

vampire bats. *Proceedings of the National Academy of Sciences of the United*

Amman, B. R., Nyakarahuka, L., McElroy, A. K., Dodd, K. A., Sealy, T. K., Schuh, A.

J., Shoemaker, T. R., Balinandi, S., Atimnedi, P., Kaboyo, W., Nichol, S. T., &

Towner, J. S. (2014). Marburgvirus resurgence in Kitaka mine bat population

after extermination attempts, Uganda. *Emerging Infectious Diseases*, *20*(10),

1761–1764. https://doi.org/10.3201/eid2010.140696

Ávila-Flores, R., Bolaina-Badal, A. L., Gallegos-Ruiz, A., & Sánchez-Gómez, W. S.

(2019). Use of linear features by the common vampire bat (Desmodus rotundus)

in a tropical cattle-ranching landscape. *Therya*, *10*(3), 229–234.

https://doi.org/10.12933/therya-19-890

Becker, D. J., Hall, R. J., Forbes, K. M., Plowright, R. K., & Altizer, S. (2018).

Anthropogenic resource subsidies and host –parasite dynamics in wildlife.

Philosophical Transactions of the Royal Society B: Biological Sciences,

373(1745). https://doi.org/10.1098/rstb.2017.0086

Benavides, J. A., Valderrama, W., Recuenco, S., Uieda, W., Suzán, G., Avila-Flores,

R., Velasco-Villa, A., Almeida, M., de Andrade, F. A. G., Molina-Flores, B.,

Vigilato, M. A. N., Pompei, J. C. A., Tizzani, P., Carrera, J. E., Ibanez, D., &

Streicker, D. G. (2020). Defining New Pathways to Manage the Ongoing

Emergence of Bat Rabies in Latin America. *Viruses*, *12*(9), 1–13.

https://doi.org/10.3390/v12091002

Benavides, J. A., Valderrama, W., & Streicker, D. G. (2016). Spatial expansions and

travelling waves of rabies in vampire bats. *Proceedings of the Royal Society B:*

Biological Sciences, *283*(1832), 1–9. https://doi.org/10.1098/rspb.2016.0328

Blackwood, J. C., Streicker, D. G., Altizer, S., & Rohani, P. (2013). Resolving the

roles of immunity, pathogenesis, and immigration for rabies persistence in

vampire bats. *Proceedings of the National Academy of Sciences of the United*

States of America, *110*(51), 20837–20842.

https://doi.org/10.1073/pnas.1308817110

Britzke, E. R., Gumbert, M. W., & Hohmann, M. G. (2014). Behavioral response of

bats to passive integrated transponder tag reader arrays placed at cave

entrances. *Journal of Fish and Wildlife Management*, *5*(1), 146–150.

https://doi.org/10.3996/082012-JFWM-065

Calisher, C. H., Childs, J. E., Field, H. E., Holmes, K. V., & Schountz, T. (2006).

Bats: Important reservoir hosts of emerging viruses. *Clinical Microbiology*

Reviews, *19*(3), 531–545. https://doi.org/10.1128/CMR.00017-06

Carter, G., & Leffer, L. (2015). Social grooming in bats: Are vampire bats

exceptional? *PLoS ONE*, *10*(10), 1–11.

https://doi.org/10.1371/journal.pone.0138430

Castle, K. T., Weller, T. J., Cryan, P. M., Hein, C. D., & Schirmacher, M. R. (2015).

Using sutures to attach miniature tracking tags to small bats for multimonth

movement and behavioral studies. *Ecology and Evolution*, *5*(14), 2980–2989.

https://doi.org/10.1002/ece3.1584

Cleasby, I. R., Owen, E., Butler, A., Baer, J., Blackburn, J., Bogdanova, M. I.,

Coledale, T., Daunt, F., Dodd, S., Evans, J. C., Green, J. A., Guilford, T., Harris,

M. P., Hughes, R., Newell, M. A., Newton, S. F., Robertson, G. S., Ruffino, L.,

Shoji, A., … Bolton, M. (2023). Assessing the importance of individual- and

colony-level variation when using seabird foraging ranges as impact

assessment and conservation tools. *Ibis*, *166*(3), 871–895.

https://doi.org/10.1111/ibi.13284

Costa, T. D., Santos, C. D., Rainho, A., Abedi- Lartey, M., Fahr, J., Wikelski, M., &

Dechmann, D. K. N. (2020). Assessing roost disturbance of strawcoloured fruit

bats (Eidolon helvum) through tri-axial acceleration. *PLoS ONE*, *15*(11

November), 1–12. https://doi.org/10.1371/journal.pone.0242662

Couto-Lima, Di., Madec, Y., Bersot, M. I., Campos, S. S., Motta, M. D. A., Dos

Santos, F. B., Vazeille, M., Da Costa Vasconcelos, P. F., Lourenço-De-Oliveira,

R., & Failloux, A. B. (2017). Potential risk of re-emergence of urban transmission

of Yellow Fever virus in Brazil facilitated by competent Aedes populations.

Scientific Reports, *7*(1), 1–12. https://doi.org/10.1038/s41598-017-05186-3

Crespo, R. F., Linhart, S. B., Burns, R. J., & Mitchell, G. C. (1972). Foraging

Behavior of the Common Vampire Bat Related to Moonlight. *Journal of*

Mammalogy, *53*(2), 366–368. https://doi.org/10.2307/1379175

D'Eon, R. G., Serrouya, R., Smith, G., & Kochanny, C. O. (2002). GPS

radiotelemetry error and bias in mountainous terrain. *Wildlife Society Bulletin*,

30(2), 430–439.

Delpietro, H. A., Russo, R. G., Carter, G. G., Lord, R. D., & Delpietro, G. L. (2017).

Reproductive seasonality, sex ratio and philopatry in Argentina's common

vampire bats. *Royal Society Open Science*, *4*(4), 1–14.

https://doi.org/10.1098/rsos.160959

Denwood, M., & Plummer, M. (2024). *runjags: Interface Utilities, Model Templates,*

Parallel Computing Methods and Additional Distributions for MCMC Models in

JAGS (R package version 2.2.2-4; p. 61). R package version 2.2.2-4.

https://github.com/ku-awdc/runjags

Greenhall, A. M. (1988). Feeding behavior. In A. M. Schmidt & R. K. Greenhall

(Eds.), *Natural History of Vampire Bats* (1st Editio, pp. 7–17). CRC Press Inc,

Boca Raton.

Greenhall, A. M., Joermann, G., & Schmidt, U. (1983). Desmodus rotundus.

694 *Mammalian Species*, *202*, 1–6. https://doi.org/10.2307/3503895

695 Griffiths, M. E., Broos, A., Bergner, L. M., Meza, D. K., Suarez, N. M., Da Silva Filipe,

696  A., Tello, C., J. Becker, D., & Streicker, D. G. (2022). Longitudinal deep

697  sequencing informs vector selection and future deployment strategies for

698  transmissible vaccines. *PLoS Biology*, *20*(4), 1–19.

699  https://doi.org/10.1371/journal.pbio.3001580

700 Gupte, P. R., Beardsworth, C. E., Spiegel, O., Lourie, E., Toledo, S., Nathan, R., &

701  Bijleveld, A. I. (2022). A guide to pre-processing high-throughput animal tracking

702  data. *Journal of Animal Ecology*, *91*(2), 287–307. https://doi.org/10.1111/1365-

703  2656.13610

704 Ham, C., Donnelly, C. A., Astley, K. L., Jackson, S. Y. B., & Woodroffe, R. (2019).

705  Effect of culling on individual badger Meles meles behaviour: Potential

706  implications for bovine tuberculosis transmission. *Journal of Applied Ecology*,

707  *56*(11), 2390–2399. https://doi.org/10.1111/1365-2664.13512

708 Hijmans, R. (2024). *terra: Spatial Data Analysis* (R package version 1.7-71).

709  https://cran.r-project.org/package=terra

710 Hooten, M. B., Johnson, D. S., McClintock, B. T., & Morales, J. M. (2017). *Animal*

711  *Movement: Statistical Models For Telemetry Data.* (1st Editio). CRS Press, Boca

712  Raton. https://doi.org/10.1201/9781315117744

713 Hopkins, H. L., Sánchez-Hernández, C., Romero-Almaraz, M. D. L., Gilley, L. M.,

714  Schnell, G. D., & Kennedy, M. L. (2003). Flight speeds of four species of

715  Neotropical bats. *Southwestern Naturalist*, *48*(4), 711–714.

716  https://doi.org/10.1894/0038-4909(2003)048<0711:FSOFSO>2.0.CO;2

717 Kahle, D., & Wickham, H. (2013). ggmap: Spatial visualization with ggplot2. *The R*

718  *Journal*, *5*(1), 144–161. http://journal.r-project.org/archive/2013-1/kahle-

wickham.pdf

Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking

as an eye on life and planet. *Science*, *348*(6240), aaa2478.

https://doi.org/10.1126/science.aaa2478

Mialhe, P. (2013). Characterization of Desmodus rotundus (E. Geoffroy, 1810)

(Chiroptera, Phyllostomidae) shelters in the Municipality of São Pedro - SP.

Brazilian Journal of Biology, *73*(3), 521–526.

O'Hara, R. B., & Sillanpää, M. J. (2009). A review of bayesian variable selection

methods: What, how and which. *Bayesian Analysis*, *4*(1), 85–118.

https://doi.org/10.1214/09-BA403

O'Mara, T. M., Wikelski, M., & Dechmann, D. K. N. (2014). 50 years of bat tracking:

Device attachment and future directions. *Methods in Ecology and Evolution*,

5(4), 311–319. https://doi.org/10.1111/2041-210X.12172

O'Shea, T. J., Cryan, P. M., Hayman, D. T. S., Plowright, R. K., & Streicker, D. G.

(2016). Multiple mortality events in bats: A global review. *Mammal Review*,

46(3), 175–190. https://doi.org/10.1111/mam.12064

Pebesma, E. (2018). Simple Features for R: Standardized Support for Spatial Vector

Data. *The R Journal*, *10*(1), 439–446. https://doi.org/10.32614/RJ-2018-009

Plummer, M. (2023). *rjags: Bayesian graphical models using MCMC* (R package

version 4-15). https://cran.r-project.org/package=rjags

Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). CODA: Convergence

Diagnosis and Output Analysis for MCMC. *R News*, *6*, 7–11.

R Core Team. (2024). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/

Ribeiro, R., Matthiopoulos, J., Lindgren, F., Tello, C., Zariquiey, C. M., Valderrama,

W., Rocke, T. E., & Streicker, D. G. (2023). Incorporating environmental

heterogeneity and observation effort to predict host distribution and viral

spillover from a bat reservoir. *Proceedings of the Royal Society B: Biological*

Sciences, *290*(20231739), 11. https://doi.org/10.1101/2023.04.04.535562

Ripperger, S. P., & Carter, G. G. (2021). Social foraging in vampire bats is predicted

by long-term cooperative relationships. *PLoS Biology*, *19*(9), 1–17.

https://doi.org/10.1371/journal.pbio.3001366

Rocha, F., Ulloa-Stanojlovic, F. M., Rabaquim, V. C. V., Fadil, P., Pompei, J. C.,

Brandão, P. E., Dias, R. A., & Ortega, J. (2020). Relations between topography,

feeding sites, and foraging behavior of the vampire bat, Desmodus rotundus.

Journal of Mammalogy, *101*(1), 164–171.

https://doi.org/10.1093/jmammal/gyz177

Rojas-Sereno, Z. E., Streicker, D. G., Medina-Rodríguez, A. T., & Benavides, J. A.

(2022). Drivers of spatial expansions of vampire bat rabies in Colombia. *Viruses*,

14(11), 1–12. https://doi.org/10.3390/v14112318

Schneider, M. C., Romijn, P. C., Uieda, W., Tamayo, H., da Silva, D. F., Belotto, A.,

da Silva, J. B., & Leanes, L. F. (2009). Rabies transmitted by vampire bats to

humans: An emerging zoonotic disease in Latin America? *Revista*

Panamericana de Salud Publica = Pan American Journal of Public Health,

25(3), 260–269. https://doi.org/10.1590/s1020-49892009000300010

Seltmann, A., Czirják, G., Courtiol, A., Bernard, H., Struebig, M. J., & Voigt, C. C.

(2017). Habitat disturbance results in chronic stress and impaired health status

in forest-dwelling paleotropical bats. *Conservation Physiology*, *5*(1), 1–14.

https://doi.org/10.1093/conphys/cox020

Stiegler, J., Gallagher, C. A., Hering, R., Müller, T., Tucker, M., Apollonio, M., Arnold,

J., Barker, N. A., Barthel, L., Bassano, B., Beest, F. M. van, Belant, J. L.,

Berger, A., Beyer Jr, D. E., Bidner, L. R., Blake, S., Börner, K., Brivio, F., Brogi,

R., … Blaum, N. (2024). Mammals show faster recovery from capture and

tagging in human-disturbed landscapes. *Nature Communications*, *15*(1), 8079.

https://doi.org/10.1038/s41467-024-52381-8

Streicker, D. G., & Allgeier, J. E. (2016). Foraging choices of vampire bats in diverse

landscapes: potential implications for land-use change and disease

transmission. *Journal of Applied Ecology*, *53*(4), 1280–1288.

https://doi.org/10.1111/1365-2664.12690

Streicker, D. G., Franka, R., Jackson, F. R., & Rupprecht, C. E. (2013).

Anthropogenic roost switching and rabies virus dynamics in house-roosting big

brown bats. *Vector-Borne and Zoonotic Diseases*, *13*(7), 498–504.

https://doi.org/10.1089/vbz.2012.1113

Streicker, D. G., Recuenco, S., Valderrama, W., Benavides, J. G., Vargas, I.,

Pacheco, V., Condori Condori, R. E., Montgomery, J., Rupprecht, C. E., Rohani,

P., & Altizer, S. (2012). Ecological and anthropogenic drivers of rabies exposure

in vampire bats: Implications for transmission and control. *Proceedings of the*

Royal Society B: Biological Sciences, *279*(1742), 3384–3392.

https://doi.org/10.1098/rspb.2012.0538

Streicker, D. G., Winternitz, J. C., Satterfield, D. A., Condori-Condori, R. E., Broos,

A., Tello, C., Recuenco, S., Velasco-Villa, A., Altizer, S., & Valderrama, W.

(2016). Host-pathogen evolutionary signatures reveal dynamics and future

invasions of vampire bat rabies. *Proceedings of the National Academy of*

Sciences of the United States of America, *113*(39), 10926–10931.

https://doi.org/10.1073/pnas.1606587113

Thieurmel, B., & Elmarhraoui, A. (2022). *Package "suncalc" Title Compute Sun*

Position, Sunlight Phases, Moon Position and Lunar Phase (R package version

0.5.1). https://cran.r-project.org/package=suncalc

Thomas, B., Holland, J. D., & Minot, E. O. (2011). Wildlife tracking technology

options and cost considerations. *Wildlife Research*, *38*(8), 653–663.

https://doi.org/10.1071/WR10211

Trajano, E. (1996). Movements of cave bats in southeastern Brazil, with emphasis

on the population ecology of the common vampire bat, Desmodus rotundus

(Chiroptera). *Biotropica*, *28*(1), 121–129. https://doi.org/10.2307/2388777

Viana, M., Benavides, J. A., Broos, A., Loayza, D. I., Niño, R., Bone, J., da Silva

Filipe, A., Orton, R., Bazan, W. V., Matthiopoulos, J., & Streicker, D. G. (2023).

Effects of culling vampire bats on the spatial spread and spillover of rabies virus.

Science Advances, *9*(10). https://doi.org/10.1126/sciadv.add7437

Wilkinson, G. S. (1985). The social organization of the common vampire bat. *Behav*

Ecol Sociobiol, *17*, 111–121.

World Health Organization. (2018). WHO Expert Consultation on rabies, third report.

In *WHO technical report series Licence: CC BY-NC-SA 3.0 IGO.* (Vol. 1012).

Figure 1: Study area and GPS data from 55 bats (1 colour, 1 bat) captured from 5

roosts (numbered triangles) and tracked for up to 9 nights. (a) Elevation map with the

5 study roosts in AAC, with an inset map of Peru highlighting the study region. (b)

roost 1, (c) roost 2, (d) roost 3, (e) roost 4 and (f) roost 5. Due to the varying scales

of movement across the roosts, this figure does not use a standardised scale, and

only 95% of the fixes closest to the roost were plotted to avoid outliers. Maps used

ggmap (Kahle & Wickham, 2013).

Figure 2: Predicted probabilities of vampire bats returning to the roost following

disturbance, from a binomial GLM. (a) Shows sex-specific estimates with 95%

confidence intervals; (b) shows the relationship of probability of return with forearm

length (mm), with the 95% confidence interval shaded.

Figure 3: Predictors of vampire bat departure time and distance from the roost. (a)

Standardised effect sizes on the log-odds scale (departure time, logit model) or log

scale (distance, exponential model), coloured by SSVS retention probability. (b)

Individual variation in the distance from the roost on the disturbance night (bat-night

interaction). (c) Observed maximum distances (km) from the roost per bat and night.

(d) Predicted maximum distances (km) per bat and night. In (c) and (d), night 0 is the

disturbance night; squares indicate means, and black dots are outliers.

Figure 4: Reconstructed movement curves for 9 bats across 9 nights. First row: bats

that travelled longer distances on night 0; middle row: bats that travelled similar

distances across different nights; bottom row: bats that left the roost after night 0 and

did not return. The shaded area indicates the credible intervals. Purple lines indicate

night zero, with progressively lighter colours indicating later nights. Due to the

varying scales of movement, the y-axis is not standardised.