Substrate heterogeneity promotes cancer cell dissemination through interface roughening

Zuzana Dunajova[1*], Saren Tasciyan[1*], Juraj Majek[1], Jack Merrin[1], Erik Sahai[2], Michael Sixt[1#], Edouard Hannezo[1#]

[1] Institute of Science and Technology Austria (IST Austria), am Campus 1, 3400 Klosterneuburg, Austria

[2] Tumour Cell Biology Laboratory, The Francis Crick Institute, London, UK

[*]Contributed equally, [#]contributed equally

[#]Correspondence: michael.sixt@ist.ac.at, edouard.hannezo@ist.ac.at

Abstract

While tumor malignancy has been extensively studied under the prism of genetic and epigenetic heterogeneity, tumor cell states also critically depend on reciprocal interactions with the microenvironment. This raises the hitherto untested possibility that heterogeneity of the untransformed tumor stroma can actively fuel malignant progression. As biological heterogeneity is inherently difficult to control, we adopted a reductionist approach and let tumor cells invade micro-engineered environments harboring obstacles with precision-controlled geometry. We find that not only the presence of obstacles, but more surprisingly their spatial disorder, causes a drastic shift from a collective to a single-cell mode of invasion – comparable in strength to cadherin loss. Combining live-imaging and perturbation experiments with minimal biophysical modeling, we demonstrate that cell detachments result both from local geometrical constraints and a global integration of spatial disorder over time. We show that different types of microenvironments map onto different universality classes of invasion dynamics - homogeneous substrates follow Kardar–Parisi–Zhang (KPZ) scaling, while disordered ones exhibit exponents consistent with KPZ with quenched disorder (KPZq). Our findings highlight generic physical principles for how the mode of cancer cell invasion depends on environmental heterogeneity, with potential implications to understand tumor evolution *in vivo*.

Introduction

Cellular heterogeneity is at the core of tumor evolution. The canonical view suggests that new genetic variants produce new cancer cell phenotypes that invade extra-tumoral micromilieus. When such variants experience favorable conditions, they clonally expand, amplifying genetic heterogeneity in an iterative process that ultimately leads to a fatal state of systemic dissemination[1–3]. While heritable genetic variations have long been recognized to play a crucial role, phenotypic variation can also arise from epigenetic regulation or transient signaling states[4,5]. Importantly, variation and selection of cancer cells are not autonomous to the tumor, but occur through reciprocal crosstalk with the untransformed microenvironment [6–8]. Remarkably, in some tumor models, invasiveness is not an intrinsic property of tumor cells but rather a transient response of rare cancer cells to signals released from a benign stroma[9–11]. This raises the possibility that heterogeneous extra-tumoral microenvironments may act as drivers of malignancy[12], although this remains largely untested due to the experimental difficulty of controlling substrate heterogeneity.

In recent years, reductionist approaches have gained increasing attention, showing that cell behaviors can emerge from physical principles without requiring genetic or epigenetic changes. For instance, bottom-up reconstitution of microenvironments has been successfully used to model cellular responses[13] and cancer dissemination in response to specific extracellular matrix (ECM) topographies[14,15]. Cellular collectives can also undergo fluid-solid transitions as a function of cell-cell and cell-ECM adhesion, cell density or active migratory forces[16–22], with fluidized states favoring invasion[23–27]. Such fluidization can create large mechanical stresses on nuclei, triggering mechano-adaptive responses that feed back to further promote malignancy[28]. In parallel, the interfaces of spreading tumor colonies display fractal-like features, that have been proposed to follow generic models for growing interfaces, such as the Kardar-Parisi-Zhang (KPZ) equation[29,30] or the Molecular Beam Epitaxy (MBE) equation[31], validated in diverse non-living systems[32,33]. However, prior studies have almost exclusively examined migration in homogeneous or ordered environments, leaving the impact of extracellular heterogeneity unexplored.

Here, we combine reductionist microengineering pillar arrays, live cell imaging and minimal particle-based simulations to dissect how environmental disorder shapes collective invasion. We chose to specifically manipulate environmental geometry, while leaving all other aspects, like cell-cell and cell-matrix adhesions, constant. This minimal change demonstrated complex impacts on interfacial growth and tumor dispersion. In particular, we found that spatial disorder robustly enhances single-cell detachments from the invasive front, a phenomenon that emerges both from local features of substrate heterogeneity, as well as a much more global ef-

fect of temporal integration of the heterogeneity causing the roughening of the tumor front. In-
deed, substrate heterogeneity leads to a transition to a different universality class of invasion
dynamics, which we characterize experimentally and theoretically. This led us to derive a quan-
titative description of the determinants of cellular dispersion, which we tested by experiments in
different environments with structured geometries. Our results link microenvironmental hetero-
geneity to invasion dynamics through universal physical principles, providing a mechanistic ba-
sis for how disorder promotes cellular dispersion and, potentially, malignant progression.
75
Results
Environmental mechanics and heterogeneity are sufficient to drive dispersion *in vitro*
As a cellular model for tumor invasion, we employed A431 human carcinoma cells[34].
This cell line is derived from a primary tumor of epithelial origin and forms stable cell-cell adhe-
sions due to the expression of E- and P-Cadherin[35]. On fibronectin-coated 2D surfaces, A431
cells spread radially as a cohesive front, without cellular detachments (Supplementary Fig. 1a).
To test invasion into geometrically precisely controlled environments, we engineered microfluid-
ic devices where a centrally applied pool of cells spread into a fibronectin-coated confined
space of 9µm height. The two confining surfaces were intersected with arbitrarily positionable
9µm diameter pillars, creating a 3D chamber that forces cells to invade into the "pillar forest" as
a monolayer. This 9µm pillar spacing was chosen both so that cells could pass through by only
moderately deforming their ~12-18µm diameter nuclei, consistent with dimensions of 3D ECM
pores *in vivo* [36,37]. The setup allowed free medium exchange and long-term life imaging
(>1week) using genetically encoded fluorescent reporters for cytoplasm (mCherry) and nuclei
(GFP).
Each microfluidic chip was divided into four sectors, allowing parallel testing of different
pillar patterns under identical conditions. The ordered patterns consisted of grid-like arrays of
obstacles with constant spacing (Fig. 1a), whereas in disordered patterns the average spacing
was preserved, but individual pillar positions randomly shifted according to a Laplace distribu-
tion (Fig. 1b). We define the level of spatial disorder by the average displacement of pillars rela-
tive to an ordered array (see Methods). Tracking invasion over 8 days showed that cells invaded
ordered patterns with occasional detachments. Strikingly, in disordered patterns cells advanced
with enhanced single cell detachments at the invasion front (Mann–Whitney U test, p = 0.0075,
Fig. 1c,e, Supplementary Movie 1). This was in stark contrast to experiments on obstacle-free
2D surfaces, where A431 cells remained cohesive with no detachments (Fig. 1d, Supplemen-
tary Movie 2).

To further asses the magnitude of the dispersion effect in disordered patterns, we com-
pared wild-type (WT) A431 cells to cells that were genetically engineered to lack E- and P- cad-
herin (EPCad$^{-/-}$). As expected, EPCad$^{-/-}$ cells invaded primarily as single cells, irrespective of
environmental geometry (Supplementary Fig. 1b-c, Supplementary Movie 3). Notably, WT cells
in disordered environments showed detachment frequencies comparable to EPCad$^{-/-}$ cells
(Supplementary Fig. 1d). Together, these data demonstrate that substrate disorder alone can
drive a drastic shift from collective to single cell invasion, with an effect comparable in magni-
tude to genetically inflicted loss of cell-cell adhesion.

Several mechanisms might explain why environmental heterogeneity promotes single
cell detachments, as different geometrical features change in ordered vs. disordered patterns.
Firstly, we reasoned that disordered pillar patterns contain regions of very narrow pore sizes,
that mechanically restrict cell-cell contacts. Such constrictions could also compress the nucleus,
potentially triggering an active cellular response that might propel the cell forward and cause
detachment[38–43]. To test this 'pore size' hypothesis, we repeated the assay in ordered pillar pat-
terns with smaller spacing (6µm), forcing nuclei to severely squeeze between obstacles. We
found that cells nonetheless invaded collectively without increased detachment, arguing against
pore size alone as the cause (Supplementary Fig. 1d).

Next, we hypothesized that gradients in pore size, present only in disordered environ-
ments, could promote detachments. In this "funnel" scenario, cells at the leading edge might
pass through a narrow constriction into more open space, while the bulk of the tissue remains
stalled. To test this, we engineered two ordered but graded pillar arrays, in which pore size var-
ied only along the invasion axis; one gradual (sinusoidal design) and one abrupt (stepwise de-
sign) (see Supplementary Fig. 1e). Importantly, neither design triggered a significant increase in
detachments (Mann–Whitney U test: $p = 0{,}61$ and $p = 0.88$, respectively; Supplementary Fig.
1f). Together, these data show that favorable local obstacle geometry alone is not sufficient to
increase cellular detachments. This led us to explore other, more collective mechanisms, where
repeated transitions through heterogeneous configurations accumulate to drive detachment in
disordered environments.

Fig. 1: Substrate heterogeneity promotes tumor cell dispersal *in vitro*. a. Experimental setup of the microfluidic invasion assay. Devices are coated with fibronectin and filled with cell culture media before A431 cells are seeded at the center **b**. Illustration of the ordered and disordered pillar patterns used to modulate substrate geometry. Cells invade outward from the central seeding region. **c.** A431 cell invasion imaged with time-lapse fluorescence microscopy over the days 3, 4 and 5.5 from left to right. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green). Right panel shows segmented nuclei at day 4. **d.** Scratch assay in homogeneous (pillar-free) environment, serving as a control for collective migration. **e.** Number of detached cells normalized by the radial interface length in the respective pillar patterns for A431 WT. Statistical comparison (Mann–Whitney U): Ordered vs. Disordered, p = 0.0075. Scale bars: 100 μm, box and whiskers plot display median, 1st and 3rd quartiles (hinges) and bars display 1.5 interquartile range from the hinges, n=16 experiments for ordered and n=18 per disordered environment,

followed over 8 days. These data show that substrate disorder significantly increases the frequency of
single-cell detachment events, despite identical average pore size and cell type, indicating a direct role for
environmental geometry in driving dispersal.

Minimal biophysical simulations recapitulate the effect of environmental heterogeneity

To further test the role of disorder, we turned to *in silico* simulations of invasion in com-
plex environments. We deliberately adopted a minimal modeling approach, asking whether spa-
tial disorder alone, in the absence of complex cell-intrinsic features, was sufficient to promote
cell detachment. Cells were modeled as soft, adhesive, and polar self-propelled particles[44–47],
interacting repulsively with static pillars arranged in different geometries (Fig. 2a). To mirror our
analysis of experimental data, detached cells were defined in the simulations as cells separated
by distance 1.55 *L* from the bulk, i.e. when the cell-cell contact force becomes negligible (see
Methods for details). Before quantitatively constraining the model parameters, we first per-
formed a broad parameter screen. Despite the model's simplicity, it robustly recapitulated key
qualitative features of the experiments: detachments increased strongly with the presence of
pillars, and even more so with pillar heterogeneity (Supplementary Fig. 2a). These features
were robust to model variations, for example, adding transient cell-pillar adhesion did not
change the phenomenology (Supplementary Fig. 2b).

We then constrained model parameters more precisely, to understand quantitatively
what causes detachments in the simulations. As detailed in the Methods (see also Supplemen-
tary Table 1 for a summary of the parameter fitting), parameters governing single-cell migration
(e.g. migration speed and bias, persistence time) were inferred from the measured behavior of
single detached cells in pillar patterns. Cell-cell adhesion strength was constrained based on the
lack of detachments in homogeneous environment and infrequent cell-cell rearrangements,
consistent with a glass-like rheology (Supplementary Fig. 2c, Supplementary Movie 4). Simula-
tions within this parameter regime confirmed that pillar heterogeneity consistently increased de-
tachments (OLS regression, R^2 = 0.98 and R^2 = 0.82 for pore sizes 0.92 and 1.06 *L*, respective-
ly – see Fig. 2b, Supplementary Movie 5). Furthermore, reducing adhesion strength not only
increased detachments overall, but also diminished the additional effect of pillar heterogeneity
(T-test: p=0.47; Supplementary Fig. 2d,e, Supplementary Movie 6), in good agreement with ex-
perimental results from EPCad[-/-] cells (Supplementary Fig. 1d).

With the constrained parametrization, we next ran many repetitions of simulations with a
specific disordered pattern and calculated the probability distribution of detachment as a func-
tion of position within the pillar pattern. This analysis, based on clustering instantaneous cell po-

sitions and tracking both transient and long-term detachments, revealed that detachment prob-
ability was most strikingly correlated with time, rather than a specific local position. Indeed, the
frequency of detachments consistently increased (OLS regression: $R^2 = 0.9$, $p < 10^{-4}$) as the in-
vasion progressed, with much steeper increase in disordered patterns (Fig. 2c).

Fig. 2: Minimal particle-based simulations reproduce heterogeneity driven detachments and reveal
differences in dynamics of interfacial growth. a. Schematic of the simulation setup. Cells are modeled
as soft, adhesive particles (gray) invading arrays with pillar obstacles (black). Relevant forces and interac-
tion potentials are illustrated. **b.** Simulated number of detached cells (at day 6) as a function of environ-
mental disorder. Disorder is controlled via the scale parameter of a Laplace distribution (see Supplemen-
tary Table 1). Bars show mean values; error bars indicate standard error. Increase of detachment fre-
quency was approximated by ordinary least squares (OLS) regression model: 3.31 ± 0.26 ($R^2 = 0.98$, $p <$
10^{-4}) and 2.19 ± 0.51 ($R^2 = 0.82$, $p = 0.013$) for pore sizes of $0.92\,L$ and $1.06\,L$, respectively. **c.** Rate of
detachment events increases over time in simulations, with steeper increase in disordered patterns. OLS

regression slopes are 2.28 ± 0.19 (R^2 = 0.89, p < 10^{-4}), and 4.29 ± 0.32 (R^2 = 0.91, p < 10^{-4}) for ordered

and disordered, respectively. **d.** Snapshots of simulated invasion fronts at different time points (magenta),

alongside experimental cell interface segmentations (see Fig. 1c,d). Interface roughness increases over

time and with environmental disorder. Detached cells are shown in red. **e.** Quantification of interfacial

roughness over time in simulations. Power-law fits yield growth exponents of β_{hom}= 0.33 ± 0.02, β_{ord}= 0.46

± 0.01, β_{dis}= 0.53 ± 0.02 (R^2 > 0.92, p < 10^{-4}), consistent with increasing roughening in disordered envi-

ronments. Markers indicate means, shaded regions show standard error and solid lines the linear fits. **f.**

Quantification of interfacial roughness in experiments yields growth exponents β_{hom}= 0.25 ± 0.01, β_{ord} =

0.54 ± 0.04, β_{dis}= 0.63 ± 0.04 (weighted least squares regression, R^2 > 0.92, p < 10^{-4}), showing a good

agreement with the theoretical predictions. **g.** Across all experiments, interface roughness at day N

strongly correlates with the number of detachments observed on day N+1 (n_{ord}= 16; n_{dis}=18, n_{hom}= 6,

Pearson r = 0.66, p < 10^{-4}). Together, these results show that increasing substrate disorder leads to

stronger interfacial roughening and elevated detachment rates, both in simulations and experiments.

Substrate heterogeneity drives KPZq-type roughening of tumor invasion fronts

From a physical perspective, a classical result on the invasion dynamics of a fluid mate-

rial such as a tissue is that its interface becomes progressively rougher over time, following a

highly universal trend[29,48–51], which depends only on a few key characteristics of the system. We

hypothesized that cell detachments could be connected to these interface properties, with "time-

integrated" heterogeneity causing cumulative surface roughening, as each pillar perturbs the

advancing front.

To test this, we computed in simulations both the overall interface perimeter (which in-

creased continuously during cell invasion, Supplementary Fig. 3a), as well as the surface

roughness (i.e. deviations in interface height from its mean, Fig. 2d,e). Interestingly, the inter-

face roughness always increased in time, but the growth exponent depended strongly on geom-

etry. Power-law fits yielded β = 0.33 ± 0.02 in homogeneous environments – consistent with

KPZ dynamics, a highly universal model of interfacial growth. Strikingly, disordered patterns

showed markedly faster roughening with β = 0.5-0.6. This is close to predictions for the KPZq

universality class[52–55], which describes interfaces in the presence of quenched noise (Fig. 2e).

Given that pillars are static in our simulations, they can indeed be considered as quenched

noise sources for the migrating interface.

Computing β across a wide parameter space (Supplementary Table 2) confirmed that

these exponents were robust despite the additional features absent from continuum KPZ mod-

els (e.g. adhesion and self-propulsion). To further test identification of the universality classes,

we independently analyzed the height–height correlation function $C(r) = \langle [h(x + r) - $

$h(x)]^2\rangle \sim r^{2\alpha}$ from the simulations at late times, to extract the roughness exponent α[56,57]. In ho-
mogeneous environments, $\alpha = 0.54 \pm 0.03$, again consistent with KPZ scaling ($\alpha \approx 0.5$, Supple-
mentary Fig. 3b). In disordered environments α increased to 0.71 ± 0.01, which is in agreement
with the KPZq universality class[57]. Finally, we asked whether the measured exponents jointly
satisfy the Family–Vicsek scaling relation[48], which predicts that data from different lateral sys-
tem sizes collapse onto a universal curve when roughness and time are rescaled as w/L^α and
t/L^z, respectively. Using the independently extracted exponents ($\alpha \approx 0.55$, $\beta \approx 0.33$ in homoge-
neous and $\alpha \approx 0.70$, $\beta \approx 0.55$ in disordered environments), we observed a very good collapse
across system sizes (Supplementary Fig. 3c). This stringent scaling test[58] provides strong evi-
dence that environmental disorder fundamentally alters the universality class of collective cell
invasion.

Strikingly, computing surface roughness and the growth exponent β in the experimental
data revealed strong agreement with the simulations ($\beta=0.25 \pm 0.01$ in homogeneous and
$\beta=0.63 \pm 0.04$ in disordered environments, Fig. 2f). In simulations of ordered pillar patterns, we
found that the predicted growth exponent depended both on the average pore size and the ori-
entation of the pattern relative to invasion direction (Supplementary Table 2, Supplementary Fig.
3d,e). Typically, ordered environments yielded intermediate exponents, with systematically low-
er interface width than disordered environments. This trend was confirmed experimentally: or-
dered patterns exhibited $\beta=0.54 \pm 0.04$, and the average interface width was 25% lower com-
pared to disordered environments (Mann-Whitney U test, p=0.0004).

Having rationalized the dynamics of interfacial growth, we next asked whether this could
explain the different trends of temporal increase in detachments across conditions. To test this,
we correlated interface roughness at day N with the number of detachments at day N+1 for
each experimental replicate. We found a strong correlation (Pearson r = 0.66, $p < 10^{-4}$) across
all conditions, arguing that there might be a causal link between interfacial roughening and cell
detachments (Fig. 2g), suggesting a direct mechanistic link between roughening and detach-
ment. To test further this idea, we revisited geometries that did not produce detachments, such
as ordered patterns with small pores and sinusoid/stepwise designs (Supplementary Fig. 3f,
Supplementary Movie 7). Simulations run under the same parameter regime but with these pillar
geometries reproduced the experimental outcome: both interfacial roughness and number of
detachments remained lower than in disordered environments (Supplementary Fig. 3g). To-
gether, these results indicate that substrate disorder promotes detachments by enhancing inter-
face roughening, motivating us to next ask where along the interface detachments preferentially
occur and which local geometrical features make them more likely.

Fig. 3: Local interface geometry predicts cell detachment across conditions. **a.** Interface shape (magenta) analyzed in simulations. Cells are shown as grey and pillar as black beads, tips of interface protrusions are highlighted in orange, detached cells in red. **b.** Schematic of how the position of detaching cells is measured relative to interface shape in both simulations and experiments. Detachment positions are normalized between the base (0) and tip (1) of the nearest protrusion. The last time frame prior to detachment is used. Cell bodies express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green). Scale bar, 100μm. **c**. Distribution of detachment positions relative to the interface shape. Detachments predominantly occur near protrusion tips across all conditions. Middle line, upper and lower lines indicate the median, first and third quartiles, respectively (collection of n=100 detaching cells from n=7 experiments and n>700 from n=40 simulations for each condition). **d.** Local curvature distributions of the interface in homogeneous, ordered and disordered environments, compared with curvature at detachment points. Pillar patterns, especially disordered, show a higher frequency of large-curvature regions. Data from 6 (homogeneous), 16 (ordered), and 18 (disordered) experiments, and 40 simulations per condition. **e.** Aspect ratio of interface protrusions in experiments, defined by the angle between the tip and the adjacent local

minima ("shoulders"). Disordered environments produce more elongated, narrow protrusions (Ordered vs.
Disordered distribution, $p < 10^{-4}$). Same dataset was used as in d). **f-h.** Spatial distribution of interfacial
protrusions (h) and detachments (g) across n=10 simulations with identical disordered pillar pattern (f),
confirming that the location of interface protrusions is highly predictive of detachments (Pearson r = 0.6, p
$< 10^{-4}$). These results show that local geometric features - especially sharp, convex interface protrusions -
strongly predict detachment events in both experiments and simulations.

Quantitative local and global determinants of cellular detachments

To follow up on this finding, we asked how a global property such as interface roughness
connects to the local occurrence of cell detachments. We hypothesized that rougher fronts gen-
erate interface regions with higher curvature and fewer cell contacts, which might facilitate de-
tachments[59–61]. Indeed, analysis of individual detachment events in both simulation and experi-
ment showed that the cells were most likely to detach from the tip of convex, "finger-like" protru-
sions (Fig. 3a-c). We next analyzed the local curvature of the interface immediately before de-
tachment, which is inversely related to the number of neighbors (Fig. 3d). Most detachments
occurred from interface positions above a threshold curvature value. By contrast, cells migrating
in pillar-free environments never reached such high interfacial curvature values, quantitatively
explaining the absence of detachment in this condition. Although the shape of curvature distri-
butions of ordered and disordered patterns appear similar in experiments, likely due to the chal-
lenges of segmenting very rough interfaces (Supplementary Fig. S4a), we found that the disor-
dered patterns yield 50 % more sharper protrusions than ordered ones (finger angle $< 50°$, p <
10^{-4}), especially at late times (Fig. 3e, Supplementary Fig. 4b). Our theory was further supported
by a strong spatial relationship between detachments and interfacial protrusions created by the
heterogeneous environment (Pearson r = 0.6, $p < 10^{-4}$; Fig. 3f-h, Supplementary Fig. 4c). To-
gether, these findings support the idea that cell dispersion in pillar patterns is a cumulative phe-
nomenon. Cell front invades with gradual increase of interfacial roughness, with geometry-
dependent scaling exponents. The existence of a critical roughness which is permissive for de-
tachments explains that detachments do not occur at early times in the pillar patterns and never
in the pillars-free conditions (within our experimental window).

To test this mechanism further via independent experiments, we designed new pillar ge-
ometries consisting of sinusoidal patterns oriented perpendicular to the invasion axis. By making
the sinusoidal modulation relatively mild, we ensured invasion across the entire pillar pattern
while systematically controlling curvature modulation (Fig. 4a, Supplementary Movie 8). As pre-
dicted by simulations, this geometry promoted formation of periodic finger-like protrusions at re-

gions of maximal pore width (Fig. 4b), and to a higher detachment probability when the frequen-

cy of sinusoidal modulation increased, producing sharper fingers (Supplementary Fig. 4d). In

both simulations and experiments, the detachment probability scaled with the local interface

shape, confirming that curvature modulation alone is sufficient to drive detachments (Fig. 4b-c,

e-f).

Finally, we asked whether additional local geometrical features predict detachment be-

yond global integrated heterogeneity. Indeed, as discussed above, repeated simulations on the

same disordered geometry revealed "hot-spots" where more detachments occurred. Given that

sinusoidal patterns aligned with the invasion axis did not yield detachments (Supplementary Fig.

1d-e, 3c-d), whereas perpendicular sinusoidal did (Fig. 4a-c), we quantified metrics of local pillar

anisotropy relative to the invasion direction (front, back, sides; Fig. 4g, Supplementary Fig. 4e-f).

In simulations, regions with locally anisotropic free space (i.e. fewer pillars parallel to invasion

compared to perpendicular) displayed a small but significant correlation with detachments

(R^2=0.1, $p < 10^{-4}$). When combined with the long-term trend of roughness-induced increases in

detachment, predictive power increased substantially (R^2=0.64, $p < 10^{-4}$, Supplementary Fig.

4e). Consistently, individual simulated cell trajectories showed increased pillar anisotropy specif-

ically at detachment events (Fig. 4h). To test this experimentally, we quantified local pillar ani-

sotropy across n=68 detaching cell trajectories. We found indeed significantly more pillars ori-

ented perpendicular than parallel to the invasion direction at the time of detachment (Fig. 4i).

Together, these results demonstrate that disordered environments promote detachments pri-

marily via globally increased interfacial roughening and finger-like protrusions, but that local

substrate anisotropy further modulates the likelihood of cell detachment.

Fig. 4: Predicting the statistics of detachments as a function of global and local heterogeneity. **a.** Comparison of simulations (top) and experiments (bottom) using a longitudinal sinusoidal pillar pattern that introduces periodic variation in lateral pore size. Cells invade with forming regularly spaced finger-like protrusions. Simulated cells are shown in grey, pillar in black, protrusion tips in orange, cell interface in magenta, detached cells in red. In experiment, cell bodies express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green). **b.** Simulated interface profile averaged over all timepoints and sinusoidal periods (n=10 simulations), showing alignment of protrusions with regions of lower pillar density. Shaded area indicates standard error. **c.** Spatial distribution of detachment events in simulations across all timepoints and periods of sinusoid profile showing a strong preference for detachment at low-density regions. **d.** Interface profile in experiment in sinusoidal geometry, averaged over all timepoints and plotted in spherical coordinates (n=4 experiments). **e-f.** Average interface profile (e, purple) and corresponding positions of

recently detached cells (f, red bars), showing strong agreement with simulation predictions. **g.** Schematic

of the method used to quantify local anisotropy in pillar geometry. At each spatial location, surrounding

pillars are grouped into 4 quadrants – front, back, left and right. **h.** In simulations, cells approaching de-

tachment exhibit a decrease in front-facing pillar density and an increase in side-facing density, leading to

detachments in locally anisotropic regions (time=0 at detachment, n>10000 detachment events from 40

simulations, plot shows mean ± SEM). **i.** Experimental measurements of local pillar density at the time of

detachment (n=68 detaching cells from 7 experiments). Cells detach more often in regions with increased

side confinement, as simulations predicted. Dot plot shows mean ± SEM. Statistical comparisons (Wil-

coxon test): Back vs. Left/Right, p = 0.0045; Front vs. Left/Right, p = 0.018; Back vs. Front, p = 0.74. To-

gether, these results show that both global pattern-induced interface shaping and local anisotropic con-

finement jointly predict where cells detach from the invading front.

Discussion

Our results demonstrate that environmental heterogeneity is a key driver of cancer cell

dissemination that the geometry of the collective invasion front critically determines detachment

events. We find that substrate heterogeneity drives detachments from the bulk through two

complementary mechanisms. First, as cells invade disordered environments, the interface grad-

ually roughens, effectively integrating heterogeneity over time. This appears to be a generic ef-

fect, since the roughening dynamics map onto the KPZq universality class of growing interfaces.

After a critical level of roughness is reached, cells at the tips of protrusions have few enough

neighbors to detach from the collective. Second, this global permissive effect is compounded by

local substrate geometry, where anisotropic pillar arrangements selectively increase detach-

ment probability at specific sites. Thus, disorder influences invasion both globally and locally,

providing a robust physical framework that may generalize to other collective cell systems be-

yond cancer.

Although our study was conducted in a controlled *in vitro* setting, where environmental

geometry could be finely tuned, the next challenge is to relate our findings to more realistic *in*

vivo situations[62]. For instance, different organs exhibit different extracellular matrix architec-

tures[63], which often fall between fully ordered and fully disordered states, although this spectrum

has not been systematically quantified. Generalizing our simulations to 3D fibrous environments

could provide new insight into how tissue organization shapes invasion through environment-

tumor cross-talk. In line with that idea, quantitative imaging of human tumor samples revealed

that spatially disordered or tilted stromal architectures can be negative prognostic factors for

clinical outcome[64,65]. Another natural extension of the model would be to incorporate remodeling

of the environment due to mechanical forces exerted by cell migration[66], which could create

feedbacks between cells and their microenvironment that further influence invasion dynamics. We anticipate that combining microengineered models, minimal simulations, and quantitative pathology could yield a predictive framework for how stromal heterogeneity impacts tumor progression.

Using an experimentally controlled bottom-up approach, we introduce stromal heterogeneity as a potential driver of malignant progression. In analogy to organismal evolution, where heterogenous habitats drive genotypic diversity, a heterogenous tumor stroma might shift tumor cells into novel phenotypic states (in our case, the single cell rather than the collective state). Although these states are transient phenotypes without heritable traits, they may later become consolidated[28] by epigenetic reprogramming and ultimately genetic accommodation. Our findings therefore highlight a potential route by which non-genetic environmental disorder can generate phenotypic diversity that later becomes fixed during tumor evolution.

Acknowledgments

The authors thank all the members of the Hannezo and Sixt groups for fruitful discussions and Takuya Kato for advices on the experimental set-up. We are also thankful for support from the Scientific Service Units (SSU) of IST Austria through resources provided by the Imaging and Optics Facility (IOF) and the Lab Support Facility (LSF). Z.D. has received funding from the Doctoral Programme of the Austrian Academy of Sciences (OeAW): grant agreement 26360. This work was supported by the European Research Council (grant ERC-SyG 101071793 to M.S).

Author contributions

S.T., Z.D., M.S., and E.H. designed the research. S.T. performed the experiments: S.T., Z. D. and Ju.M. performed data and image analysis; Z.D. and Ju. M. performed simulations; Ja.M. supported microfluidic engineering; E.S. provided cellular model systems. M.S. and E.H. supervised the study; S. T., Z.D., M.S. and E.H. wrote the manuscript, with input from all authors.

References

1. Almendro, V., Marusyk, A. & Polyak, K. Cellular heterogeneity and molecular evolution in cancer. *Annu. Rev. Pathol.* **8**, 277–302 (2013).

2. Altschuler, S. J. & Wu, L. F. Cellular Heterogeneity: Do Differences Make a Difference? *Cell*

141, 559–563 (2010).

3. McGranahan, N. & Swanton, C. Clonal Heterogeneity and Tumor Evolution: Past, Present,

and the Future. *Cell* **168**, 613–628 (2017).

4. Cheung, K. J. & Ewald, A. J. A collective route to metastasis: Seeding by tumor cell clusters.

Science **352**, 167–169 (2016).

5. Yang, R. *et al.* Distinct epigenetic features of tumor-reactive CD8+ T cells in colorectal can-

cer patients revealed by genome-wide DNA methylation analysis. *Genome Biol.* **21**, 2

(2019).

6. Ferguson, L. P., Diaz, E. & Reya, T. The Role of the Microenvironment and Immune System

in Regulating Stem Cell Fate in Cancer. *Trends Cancer* **7**, 624–634 (2021).

7. Friedl, P. & Alexander, S. Cancer Invasion and the Microenvironment: Plasticity and Recip-

rocity. *Cell* **147**, 992–1009 (2011).

8. Spill, F., Reynolds, D. S., Kamm, R. D. & Zaman, M. H. Impact of the physical microenvi-

ronment on tumor progression and metastasis. *Curr. Opin. Biotechnol.* **40**, 41–48 (2016).

9. Giampieri, S. *et al.* Localized and reversible TGFβ signalling switches breast cancer cells

from cohesive to single cell motility. *Nat. Cell Biol.* **11**, 1287–1296 (2009).

10. Brabletz, T. EMT and MET in Metastasis: Where Are the Cancer Stem Cells? *Cancer Cell*

22, 699–701 (2012).

11. Padmanaban, V. *et al.* E-cadherin is required for metastasis in multiple models of breast

cancer. *Nature* 1–6 (2019) doi:10.1038/s41586-019-1526-3.

12. Rezk, R. *et al.* Spatial heterogeneity of cell-matrix adhesive forces predicts human glioblas-

toma migration. *Neuro-Oncol. Adv.* **2**, vdaa081 (2020).

13. Flommersfeld, J., Stöberl, S., Shah, O., Rädler, J. O. & Broedersz, C. P. Geometry-

Sensitive Protrusion Growth Directs Confined Cell Migration. *Phys. Rev. Lett.* **132**, 098401

(2024).

14. Jouybar, M., Sleeboom, J. J. F., Vaezzadeh, E., Sahlgren, C. M. & den Toonder, J. M. J. An

in vitro model of cancer invasion with heterogeneous ECM created with droplet microflu-

idics. *Front. Bioeng. Biotechnol.* **11**, (2023).

15. Wang, W., Law, R. A., Perez Ipiña, E., Konstantopoulos, K. & Camley, B. A. Confinement,

Jamming, and Adhesion in Cancer Cells Dissociating from a Collectively Invading Strand.

PRX Life **3**, 013012 (2025).

16. Bocanegra-Moreno, L., Singh, A., Hannezo, E., Zagorski, M. & Kicheva, A. Cell cycle dy-

namics controls fluidity of the developing mouse neuroepithelium. 2022.01.20.477048 Pre-

print at https://doi.org/10.1101/2022.01.20.477048 (2022).

17. Mongera, A. *et al.* A fluid-to-solid jamming transition underlies vertebrate body axis elonga-

tion. *Nature* (2018) doi:10.1038/s41586-018-0479-2.

18. Petridou, N. I., Corominas-Murtra, B., Heisenberg, C. P. & Hannezo, E. Rigidity percolation

uncovers a structural basis for embryonic tissue phase transitions. *Cell* **184**, 1914-1928.e19

(2021).

19. Pinheiro, D., Kardos, R., Hannezo, É. & Heisenberg, C.-P. Morphogen gradient orchestrates

pattern-preserving tissue morphogenesis via motility-driven unjamming. *Nat. Phys.* **18**,

1482–1493 (2022).

20. Bi, D., Lopez, J. H., Schwarz, J. M. & Manning, M. L. A density-independent rigidity transi-

tion in biological tissues. *Nat. Phys.* **11**, 1074–1079 (2015).

21. Mitchel, J. A. *et al.* In primary airway epithelial cells, the unjamming transition is distinct from

the epithelial-to-mesenchymal transition. *Nat. Commun.* **11**, 1–14 (2020).

22. Kato, T. *et al.* Interplay of adherens junctions and matrix proteolysis determines the invasive

pattern and growth of squamous cell carcinoma. *eLife* **12**, e76520 (2023).

23. Grosser, S. *et al.* Cell and Nucleus Shape as an Indicator of Tissue Fluidity in Carcinoma.

Phys. Rev. X **11**, (2021).

24. Ilina, O. *et al.* Cell–cell adhesion and 3D matrix confinement determine jamming transitions

in breast cancer invasion. *Nat. Cell Biol.* **22**, 1103–1115 (2020).

25. Malinverno, C. *et al.* Endocytic reawakening of motility in jammed epithelia. *Nat. Mater.* **16**,

587–596 (2017).

26. Oswald, L., Grosser, S., Smith, D. M. & Käs, J. A. Jamming transitions in cancer. *J. Phys.*

Appl. Phys. **50**, 483001 (2017).

27. Fuhs, T. *et al.* Rigid tumours contain soft cancer cells. *Nat. Phys.* **18**, 1510–1519 (2022).

28. Frittoli, E. *et al.* Tissue fluidification promotes a cGAS–STING cytosolic DNA response in

invasive breast cancer. *Nat. Mater.* (2022) doi:10.1038/s41563-022-01431-x.

29. Kardar, M., Parisi, G. & Zhang, Y.-C. Dynamic Scaling of Growing Interfaces. *Phys. Rev.*

Lett. **56**, 889–892 (1986).

30. Muzzio, N. E., Pasquale, M. A., González, P. H. & Arvia, A. J. Influence of individual cell

motility on the 2D front roughness dynamics of tumour cell colonies. *J. Biol. Phys.* **40**, 285–

308 (2014).

31. Brú, A., Albertos, S., Luis Subiza, J., García-Asenjo, J. L. & Brú, I. The Universal Dynamics

of Tumor Growth. *Biophys. J.* **85**, 2948–2961 (2003).

32. Maunuksela, J. *et al.* Kinetic Roughening in Slow Combustion of Paper. *Phys. Rev. Lett.* **79**,

1515–1518 (1997).

33. Takeuchi, K. A. & Sano, M. Evidence for Geometry-Dependent Universal Fluctuations of the

Kardar-Parisi-Zhang Interfaces in Liquid-Crystal Turbulence. *J. Stat. Phys.* **147**, 853–890

(2012).

34. Giard, D. J. *et al.* In vitro cultivation of human tumors: establishment of cell lines derived

from a series of solid tumors. *J. Natl. Cancer Inst.* **51**, 1417–1423 (1973).

35. Macpherson, I. R. *et al.* p120-catenin is required for the collective invasion of squamous cell

carcinoma cells via a phosphorylation-independent mechanism. *Oncogene* **26**, 1210334

(2007).

36. Wolf, K. *et al.* Physical limits of cell migration: Control by ECM space and nuclear deforma-

tion and tuning by proteolysis and traction force. *J. Cell Biol.* **201**, 1069–1084 (2013).

37. Wolf, K. *et al.* Collagen-based cell migration models in vitro and in vivo. *Semin. Cell Dev.*

Biol. **20**, 931–941 (2009).

38. Bertillot, F., Miroshnikova, Y. A. & Wickström, S. A. SnapShot: Mechanotransduction in the

nucleus. *Cell* **185**, 3638-3638.e1 (2022).

39. Kalukula, Y., Stephens, A. D., Lammerding, J. & Gabriele, S. Mechanics and functional con-

sequences of nuclear deformations. *Nat. Rev. Mol. Cell Biol.* **23**, 583–602 (2022).

40. Venturini, V. *et al.* The nucleus measures shape changes for cellular proprioception to con-

trol dynamic cell behavior. *Science* **370**, eaba2644 (2020).

41. Venturini, V. *et al.* The nucleus as a mechanosensitive controller of cell behavior. *Biophys.*

J. **121**, 490a (2022).

42. Lomakin, A. J. *et al.* The nucleus acts as a ruler tailoring cell responses to spatial con-

straints. *Science* **370**, eaba2894 (2020).

43. Kidiyoor, G. R. *et al.* ATR is essential for preservation of cell mechanics and nuclear integ-

rity during interstitial migration. *Nat. Commun.* **11**, 4828 (2020).

44. Alert, R. & Trepat, X. Physical Models of Collective Cell Migration. *Annu. Rev. Condens.*

Matter Phys. **11**, 77–101 (2020).

45. Basan, M., Elgeti, J., Hannezo, E., Rappel, W.-J. & Levine, H. Alignment of cellular motility

forces with tissue flow as a mechanism for efficient wound healing. *Proc. Natl. Acad. Sci.*

110, 2452–2459 (2013).

46. Garcia, S. *et al.* Physics of active jamming during collective cellular motion in a monolayer.

Proc. Natl. Acad. Sci. **112**, 15314–15319 (2015).

47. Henkes, S., Fily, Y. & Marchetti, M. C. Active jamming: Self-propelled soft particles at high

density. *Phys. Rev. E* **84**, 040301 (2011).

48. Family, F. & Vicsek, T. Scaling of the active zone in the Eden process on percolation net-

works and the ballistic deposition model. *J. Phys. Math. Gen.* **18**, L75 (1985).

49. Chepizhko, O. *et al.* Bursts of activity in collective cell migration. *Proc. Natl. Acad. Sci.* **113**,

11408–11413 (2016).

50. Podewitz, N., Jülicher, F., Gompper, G. & Elgeti, J. Interface dynamics of competing tissues.

New J. Phys. **18**, 083020 (2016).

51. Rapin, G. *et al.* Roughness and dynamics of proliferating cell fronts as a probe of cell–cell

interactions. *Sci. Rep.* **11**, 8869 (2021).

52. Olami, Z., Procaccia, I. & Zeitak, R. Interface roughening in systems with quenched disor-

der. *Phys. Rev. E* **52**, 3402–3414 (1995).

53. Dias, C. S., Yunker, P. J., Yodh, A. G., Araújo, N. a. M. & Gama, M. M. T. da. Interaction

anisotropy and the KPZ to KPZQ transition in particle deposition at the edges of drying

drops. *Soft Matter* **14**, 1903–1907 (2018).

54. Csahók, Z., Honda, K., Somfai, E., Vicsek, M. & Vicsek, T. Dynamics of surface roughening

in disordered media. *Phys. Stat. Mech. Its Appl.* **200**, 136–154 (1993).

55. Muzzio, N. E. *et al.* Spatio-temporal morphology changes in and quenching effects on the

2D spreading dynamics of cell colonies in both plain and methylcellulose-containing culture

media. *J. Biol. Phys.* **42**, 477–502 (2016).

56. Barabási, A.-L. & Stanley, H. E. *Fractal Concepts in Surface Growth.* (Cambridge University

Press, Cambridge, 1995). doi:10.1017/CBO9780511599798.

57. Takeuchi, K. A. & Sano, M. Universal Fluctuations of Growing Interfaces: Evidence in Tur-

bulent Liquid Crystals. *Phys. Rev. Lett.* **104**, 230601 (2010).

58. Yamazaki, Y., Saito, K., Kobayashi, N., Ozawa, T. & Matsushita, M. Reconstruction and Ex-

tension of the Family–Vicsek Scaling Hypothesis for Growing Rough Interfaces. *J. Phys.*

Soc. Jpn. **76**, 104002 (2007).

59. Mark, S. *et al.* Physical Model of the Dynamic Instability in an Expanding Cell Culture. *Bio-*

phys. J. **98**, 361–370 (2010).

60. Tarle, V., Ravasio, A., Hakim, V. & Gov, N. S. Modeling the finger instability in an expanding

cell monolayer. *Integr. Biol.* **7**, 1218–1227 (2015).

61. Büscher, T., Diez, A. L., Gompper, G. & Elgeti, J. Instability and fingering of interfaces in

growing tissue. *New J. Phys.* **22**, 083005 (2020).

62. Dai, W. *et al.* Tissue topography steers migrating Drosophila border cells. *Science* **370**,

987–990 (2020).

63. Park, D. *et al.* Extracellular matrix anisotropy is determined by TFAP2C-dependent regula-

tion of cell collisions. *Nat. Mater.* 1–12 (2019) doi:10.1038/s41563-019-0504-3.

64. Conklin, M. W. *et al.* Aligned Collagen Is a Prognostic Signature for Survival in Human

Breast Carcinoma. *Am. J. Pathol.* **178**, 1221–1232 (2011).

65. Mukherjee, A., Jana, A., Koons, B. & Nain, A. Design of Fiber Networks for Studying Metas-

tatic Invasion. in *Biomechanics in Oncology* (eds Dong, C., Zahir, N. & Konstantopoulos, K.)

289–318 (Springer International Publishing, Cham, 2018). doi:10.1007/978-3-319-95294-

9_14.

66. Tsingos, E., Bakker, B. H., Keijzer, K. A. E., Hupkes, H. J. & Merks, R. M. H. Hybrid cellular

Potts and bead-spring modeling of cells in fibrous extracellular matrix. *Biophys. J.* **122**,

2609–2622 (2023).

67. Labernadie, A. *et al.* A mechanically active heterotypic E-cadherin/N-cadherin adhesion en-

ables fibroblasts to drive cancer cell invasion. *Nat. Cell Biol.* **19**, 224–237 (2017).

68. Leithner, A. *et al.* Diversified actin protrusions promote environmental exploration but are

dispensable for locomotion of leukocytes. *Nat. Cell Biol.* **18**, 1253–1259 (2016).

69. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*

9, 676–682 (2012).

70. Tinevez, J.-Y. *et al.* TrackMate: An open and extensible platform for single-particle tracking.

Methods **115**, 80–90 (2017).

71. Linkert, M. *et al.* Metadata matters: access to image data in the real world. *J. Cell Biol.* **189**,

777–782 (2010).

72. Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat. Methods* **16**,

1226–1232 (2019).

73. Agostinelli, C. & Lund, U. *R Package \textttcircular: Circular Statistics (Version 0.4-95).* (CA:

Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University,

Venice, Italy. UL: Department of Statistics, California Polytechnic State University, San Luis

Obispo, California, USA, 2022).

74. Pewsey, A., Neuhäuser, M. & Ruxton, G. D. *Circular Statistics in R.* (OUP Oxford, 2013).

75. Howse, J. R. *et al.* Self-Motile Colloidal Particles: From Directed Propulsion to Random

Walk. *Phys. Rev. Lett.* **99**, 048102 (2007).

76. Shaebani, M. R., Wysocki, A., Winkler, R. G., Gompper, G. & Rieger, H. Computational

models for active matter. *Nat. Rev. Phys.* **2**, 181–199 (2020).

77. Anderson, J. A., Glaser, J. & Glotzer, S. C. HOOMD-blue: A Python package for high-

performance molecular dynamics and hard particle Monte Carlo simulations. *Comput. Ma-

ter. Sci.* **173**, 109363 (2020).

78. Mandal, R., Bhuyan, P. J., Chaudhuri, P., Dasgupta, C. & Rao, M. Extreme active matter at

high densities. *Nat. Commun.* **11**, 2581 (2020).

79. Angelini, T. E. *et al.* Glass-like dynamics of collective cell migration. *Proc. Natl. Acad. Sci.*

108, 4714–4719 (2011).

80. Kang, W. *et al.* A novel jamming phase diagram links tumor invasion to non-equilibrium

phase separation. *iScience* **24**, 103252 (2021).

81. Narinder, N., F. Bos, M., Abaurrea-Velasco, C., Graaf, J. de & Bechinger, C. Understanding enhanced rotational dynamics of active probes in rod suspensions. *Soft Matter* **18**, 6246–6253 (2022).

82. Yamashita, H. *et al.* Lysophosphatidic Acid Upregulates Laminin-332 Expression during A431 Cell Colony Dispersal. *J. Oncol.* **2010**, 107075 (2010).

83. Tweedy, L. *et al.* Seeing around corners: Cells solve mazes and respond at a distance using attractant breakdown. *Science* **369**, eaay9792 (2020).

84. Zhou, W. & Yan, H. Alpha shape and Delaunay triangulation in studies of protein-related interactions. *Brief. Bioinform.* **15**, 54–64 (2014).

85. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

Methods

Experiments

Cell Culture

A431 cells expressing mCherry were grown in DMEM (Gibco, 10569010) containing phenol red,
high glucose, GlutaMAX, pyruvate, 10% fetal calf serum (FBS, Gibco, 10270106), non-essential
aminoacids (Gibco, 11140050), penicillin (50U/ml) and streptomycin (50µg/ml, Gibco 15070063)
in humidified incubators at 37°C with 5% CO_2. EPCad$^{-/-}$ A431 cells were made as described in
Labernadie et al. 2017[67]. Cells were regularly tested for mycoplasma contamination and pre-
served in growth medium containing 5% (v/v) DMSO in cryogenic conditions for longer periods.
Cells were transfected with lentivirus containing H2B-GFP construct and selected for blasticidin
resistance and then later sorted for GFP signal, regularly.

Device Fabrication

Patterns were generated with Python script in SVG format. Designs of photomasks were com-
pleted in (Inkscape, v0.92) in layers for each confinement. SVG files were converted to Auto-
CAD DXF R14 files, further converted to Gerber files (LinkCAD) and finally printed on chrome
photomasks at a resolution of 1 µm (JD Photo Data & Photo Tools). For high-resolution struc-
tures, wafers were first coated with a sub-micrometre layer of SU-8 as a sticky substrate for lat-
er SU-8 structures. We used GM-1040 SU-8 photoresist (Gersteltec) at 5,000 rpm for 2 min with
flood UV exposure. The feature layer was then applied with SU8-2005 (Microchem). According
to the datasheet, a UV-absorbing glass plate and higher exposure were used to make sharp
features. PL-360-LP from Omega Optical and 550 mJ/cm^2 with an EVG mercury lampbased
mask aligner gave good results in the 3–9-µm-thickness range. The thick (25 µm) loading chan-
nels were made with Microchem SU8-3025 using the EVG mask aligner according to the
Microchem data sheet without the PL-360-LP plate. The master wafers underwent vapour
silanization for 1 h with (1H, 1H, 2H, 2H-perfluorooctyl) trichlorosilane. The final results were
single hydrophobic master wafers containing several aligned layers. Polydimethylsiloxane
(PDMS) devices were made with Sylgard 184 Elastomer Kit (Dow Corning) using a ratio of 10:1.
The PDMS was mixed in a Thinky mixing machine and then poured on the wafer in a 3D printed
tray and further degassed several times in a vacuum chamber. The devices were baked at 80
°C overnight. The hardened devices were carefully peeled off the mould and cut into single de-

vices using a razor blade. Holes were punched with a custom arbor press or with skin biopsy

puncture tools (1 mm) for the central hole, and sticky tape was used to remove debris. Glass

coverslips were sonicated for 5 min in 100% isopropanol and blown dry. Devices were bonded

to the glass slides by oxygen plasma then sealed together at 85 °C for 1 h to make the bond

permanent.

Invasion Assays

Microfabricated devices were inserted into 12-well cell culture plates from the bottom side and

glued with molten paraffin. Devices were filled with fibronectin (F1141, Sigma) solution (diluted

in 1:5 from 1mg/ml stock solution in PBS) using a syringe pump, sealed and incubated for 3h in

cell culture growth conditions. Devices were later flushed with 100 µl cell culture growth medium

with syringe pumps. Debris were removed from the center hole, either spheroids of cancer cells

(generated with hanging drops) were dropped or a pipette tip filled with medium was inserted

into the hole, where 1.5 µl of the cell suspension was filled. Cells were incubated until they sank

into the hole and imaged at later days. Wells were checked for contamination under micro-

scope.

2D Spread Assay

A431 cells were mixed with cytoplasmic mCherry reporter expressing A431 cells (10:1) and as-

sembled into 1000 cell spheroids in hanging drops overnight. Glass cover slips were coated with

fibronectin (F1141, Sigma, diluted in 1:40 from 1mg/ml stock solution in PBS) for 1h and then

washed with PBS. Subsequently, chambers were filled with cell culture media and spheroids

were placed to adhere on the substrate. Cells were observed with fluorescence and DIC mi-

croscopy.

Scratch Assay

In order to observe invasion on non-confined 2D surface with an initial straight interface, cells in

suspension were placed and grown to full confluence on fibronectin coated glass substrates

(see 2D spread assay above). Then, approximately at the center of the cover slip, cells were

scratched with a rubber cell scraped and detached cells were aspirated and washed away with

cell culture media. Cells were observed with fluorescence and DIC microscopy.

Imaging

Cells were either imaged once a day at room conditions or in cell culture conditions for time-lapse imaging (10 min interval) for several days (typically 3-4 days). Invasion was recorded with an inverted wide-field Nikon Eclipse Ti-2 microscope equipped with a Lumencor light source (475 nm for GFP and 594 nm for mCherry) and an incubation chamber, heated stage and CO_2 mixer (Ibidi). Generally, the fluorescence imaging was accompanied with light microscopy (DIC) imaging. For long term imaging, phenol red color was checked and the temperature was monitored with a temperature sensor. Custom made CNC milled box and a pair 3D printed plugs were used to prevent CO_2 and humidity leak.

Collagen Assay

A collagen mixture was created with a final collagen concentration of 1.7 mg/ml by mixing bovine collagen (PureCol, Advanced BioMatrix) in 1x minimum essential Eagle (Sigma) and 0.4% sodium bicarbonate solution (Sigma Aldrich). Mixture was checked for a pale pink color to ensure for correct pH for polymerization. 20 µl of the collagen mixture was mixed with 9 µl of the growth medium and mixed gently. Subsequently, 1 µl of the spheroid containing growth medium was added to the mixture and mixed by stirring with the tip. Final mixture was gently placed on a glass coverslip, which was previously glued under a 60 mm dish with molten paraffin. The dish was flipped to let the collagen drop hang from the glass coverslip, whereas the plastic lid was filled with droplets of PBS to keep the dish humidified inside the incubator. Collagen drop was let to polymerize in the incubator for 45-60 min under cell culture conditions. Before imaging, PBS was removed and the dish was flipped again and medium was added to the chamber with the collagen drops. Invasion into the collagen was imaged with light microscopy (DIC) over several days under cell culture conditions.

Lentivirus Production

Lentivirus production was performed as described previously[68]. In brief, LX-293 HEK cells (Clontech) were co-transfected with LentiCRISPRv1 or LentiCRISPRv2, packaging-ps-PAX2 (Addgene no. 12260) and pCMV-VSV-G envelope plasmids using Lipofectamine 2000 (ThermoFisher Scientific) as recommended by the manufacturer, and cells were resuspended in the growth medium 1 day before transfection. Supernatant was collected after 72 h and stored at –80 °C.

Image processing and experimental analysis

Image Analysis

FiJi was used for image processing, including using the TrackMate plugin for manual cell tracking[69,70]. Proprietary ND2 files from Nikon NIS Elements were converted using Bio-Formats plugin[71] and the physically calibrated (pixel size and imaging interval). Images were aligned and processed into TIFF and HDF files. Most of the analysis processes were automated with ImageJ macros. Cell nuclei (H2B-GFP, green) and the remaining cell body (mCherry, blue) and nuclear H2B-GFP (green) were initially segmented using Ilastik and then particle analysis feature was used to quantify invasion[72]. Cells were tracked semi-automatically with TrackMate, tracks were corrected manually. Dividing and dying cells were excluded, all gaps were filled with interpolation. Final images were processed with automated scripts for display and exported into PNG. In images, cells were considered detached, when their cell bodies were not considered touching the bulk. Finger measurements were performed manually by measuring the distance from the tip to the finger baseline. Detachments were selected over a wide range of recorded experiments. The pore size of the detaching cell during detachment vs. the pore size in front of the cell, which is being detached from, was measured manually with an interactive macro.

Experimental data analysis

The cell interface in experiment was obtained by identifying the nuclei positions of cells at the interface, for better comparison with simulations, which do not model the cell bodies (Supplementary Fig. 4a). Identification of cells at interface was done by first running Delaunay triangulation of all cell nuclei in each timestep and subsequently removing the vertices, which cross through the segmented interface of bulk cell bodies (blue line in Supplementary Fig. 4a). This way we could efficiently find all cells, which are in contact with the very rough and irregular bulk interface. Before using the interface positions for analyzing the roughness exponents, we had to remove the overall tilts in the interface present in some experiments (see left panel in Fig. 1d, Supplementary Fig. 5a). This was done by removing long-wavelength Fourier modes (> 500 µm). The analysis of the local pillar geometry around the detaching cells (Fig. 4i) was performed by identifying all close pillar in a given radius around a the x,y position of the detaching cells and dividing them into 4 quadrants (front, back, left right, see Fig. 4g, the center of the front quadrant being the orientation of the detaching cell). Multiple radii were tested, all resulting in qualitatively similar result (final value: r_{max}= 27 µm, i.e three times the pore size). The identified pillars in

each quadrant were subsequently weighted by their distance from the reference position (r_{max}-

r)/r_{max} and summed.

Statistics and Data Visualization

All statistics for experimental data were performed in *R (4.0.3)* and plotted using *ggplot2* pack-

age (R Core Team 2020; Wickham 2016:2). Tracking data was analyzed with *TraXpert*, an in-

house developed particle tracking analysis application. Directionality data was analyzed with

circular package and circular statistics[73,74]. Minor cosmetic adjustments were performed in

Inkscape, without affecting the data points or the interpretation of the plots. Figures were as-

sembled in *Inkscape* and exported into high resolution lossless PNG images.

Numerical model

We implemented two dimensional Brownian dynamics simulations as a minimal model of collec-

tive cell migration[44–47] through complex environments (Fig. 2a). In this framework, each cell is

represented as a soft self-propelled particle (approximating its nucleus[41]) moving on a substrate.

The overdamped equation of motion for each particle is given by:

$$\gamma \frac{d\vec{r_i}}{dt} = \frac{dV_{pair}}{dr_{i,j}} + \vec{F_p}$$

731   where γ is the friction coefficient, $\vec{r_i}$ the particle coordinates, V_{pair} the pairwise interaction poten-

732   tial, and $r_{i,j}$ the distance between particles. The active self-propulsion force $\vec{F_p}$ is given by

$$\vec{F_p} = \gamma v_0 \vec{p_i}$$

733   Where v_0 is the intrinsic propulsion speed and $\hat{p}_i = (cos\theta_i, sin\theta_i)$ defines the orientation of parti-

734   cle i. In the absence of interactions, this force yields an average cell speed of v_0.

735   The polarity of migrating cells is subject to rotational diffusion and an external biasing torque

736   τ_{bias}, which directs cell migration away from the bulk (see below for discussion of its origin). The

737   orientation dynamics are given by

$$\frac{\delta\theta_i}{\delta t} = \tau_{bias} + \sqrt{2D_r}\eta = -k_{bias}sin(\theta_i) + \sqrt{2D_r}\eta$$

where D_r is the rotational diffusion coefficient, η a unit-variance Gaussian random variable, and

k_{bias} controls the strength of the bias torque. We define $\theta_i = 0$ as the invasion direction (to the

right in all simulations shown). To validate the numerical implementation, we computed the

mean square displacement of a single active Brownian particle without bias torque[75]. The ob-

served crossover from ballistic to diffusive motion was consistent with the used value of D_r.

To describe cell-cell interactions, we assume a simple pairwise potential consisting of short-

range repulsion and intermediate-range attraction, as previously described[44,45,76]. The potential

takes the form

$$V_{pair} = \sum_i \sum_{j>i} v_{pair}(r_{i,j})$$

$$v_{pair}(r_{i,j}) = \begin{cases} 4\epsilon\left[\left(\dfrac{\sigma}{r_{i,j}}\right)^4 - \left(\dfrac{\sigma}{r_{i,j}}\right)^2\right] + \Delta v_{pair}(r_{i,j}) \,, & r_{i,j} < r_{cut} \\ 0, & r_{i,j} \geq r_{cut} \end{cases}$$

where $r_{i,j}$ is the distance between particles *i* and *j,* $\square$ is the depth of the potential well, σ is the

characteristic interaction distance, and r_{cut} is the cutoff distance. The potential is shifted by a lin-

ear correction term Δv_{pair}, ensuring that both the potential and its derivative smoothly vanish at

the cutoff $\dfrac{dv_{pair}}{dr}(r_{cut}) = 0$.

To model the structured environments (pillar geometries), we introduced stationary particles into

the simulation box, which contribute to the total pair potential. The cell-pillar interaction was

truncated at its minimum ($r_{cut} = \sqrt{2}\sigma$) so that only repulsive interactions remained, consistent

with the absence of strong experimental evidence for stable cell-pillar adhesion. Importantly, a

central assumption of our minimal modeling approach is that intrinsic cellular parameters (e.g.,

motility, adhesion strength) are held constant across conditions; thus the observed differences

in detachment dynamics arise solely from variations in substrate heterogeneity.

Details of the simulation setup

We used the HOOMD-blue v3 library to perform numerical simulations[77]. The equations of mo-

tion were integrated using the Euler-Maruyama scheme with a timestep of *dt* =0.001τ. All simu-

lations were conducted in two dimensions.

Simulation parameters and results are reported in dimensionless units. For convenient compari-

son with experiments, lengths are expressed in units of cell-cell equilibrium distance *L* (defined

as the interaction minimum $1L = \sqrt{2}\sigma \approx 16$ um) and time in units of $\tau = 10^3$ s. Simulation output

was saved every 600 frames, corresponding to the smallest experimental timestep (10 minutes).

Unless stated otherwise, the simulation parameters used in the model are summarized in Sup-
plementary Table 1:

Supplementary Table 1: Summary of the simulation parameters.

Parameter	Notation	Chosen value (simulation units)	Parameter fitting
Friction coefficient	γ	1	Fixed parameter, non-dimensionalizes timescales
Attraction cutoff	r_{cut}	1.55 L (1 L for cell-pillar interaction)	Fixed parameter (on the order of cell size to avoid next-neighbor interactions)
Self-propulsion speed	v_0	0.43 L/τ	Fitted to correspond to mean single-cell speed $6.9*10^{-3}$ μm/s inside the pillar pattern (Supplementary Fig. 5a,b)
Rotational diffusion	D_r	0.7 τ^{-1}	Fitted according to MSAD of experimental trajectories of detached cells (Supplementary Fig. 5c)
Bias torque	k_{bias}	0.75 rad/τ	Fitted according to the drift of experimental trajectories of detached cells (Suplementary Fig. 5d)
Cell fluidity number	$\gamma v_0/\varepsilon$	0.52	Constrained by cell confluency while allowing occasional bulk cell rearrangements (Supplementary Fig. 2b, Supplementary Movie 4, Supplementary Table 3)
Pore size	L_p	0.88 -1.3 L	Free parameter (on the order of cell size; lower bound ensures cells can pass through pores individually)
Pore heterogeneity	$\mathscr{H}$	$\mu = b = 0.1*L_p$ - $0.5*L_p$	Defined by parameters of Laplace distribution; upper bound taken from disordered pattern in experiments

Before doing a systematic parameter search, we first constrained parameters to reasonable
ranges and conducted non-dimensional analysis to identify the key dimensionless ratios. The

fluidity of a tissue in the absence of obstacles is governed by a dimensionless number relating

self-propulsion to cell-cell interaction strength $(v_0\gamma/\epsilon)$[46,78,79]. If the ratio is too low, cells behave

as a solid, with very few neighbor rearrangements. Conversely, above a certain threshold, the

tissue fluidizes, as motility-driven rearrangements become possible. However, when considering

not only bulk tissue but also a cell interfaces, very large values of this ratio can lead to sponta-

neous cell detachments in pillar-free environments, resembling a gas[80]. Importantly, experi-

ments with monolayers invading a homogeneous environment support an intermediate value of

this ratio, as we observed virtually no detachments, while cells remained able to move relative

to one another (Supplementary Fig. 1b, Supplementary Movie 4, Supplementary Table 3).

Furthermore, we leveraged the presence of single detached cells in pillar experiments to con-

strain several parameters related to single-cell migratory properties. In particular, by calculating

instantaneous (x,y) displacements and the mean squared angular displacement (MSAD)[81] of

detached cells in both simulations and experiments, we could strongly constrain the values of

self-propulsion speed (v_0) and particle persistence length (v_0/D_r) (Supplementary Fig. 5b-d). The

value of external bias (k_{bias}) was fixed by analyzing the radial drift in trajectories of detached

cells in experiment and comparing it to simulations (Supplementary Fig. 5e). Notably, we de-

tected strong biased migration even in single cells far from the leading edge, indicating that this

bias does not simply arise from pushing forces due to bulk proliferation. Given previous reports

of self-generated gradients guiding long-range migration[82,83], such mechanisms are compatible

with this bias, although we chose to remain agnostic about its origin in the simulations. Finally,

we performed an additional sanity check by comparing the mean squared displacement (MSD)

of single detached cells in simulations and experiments, which allowed us to validate the chosen

combination of parameters D_r, v_0, and k_{bias}. The final parameter set reproduced the experimental

MSD curve most closely, yielding the highest coefficient of determination (pseudo-R^2 = 0.89),

particularly at long times (Supplementary Fig. 5f).

The simulations of migrating cells used for comparative statistics across various pillar geome-

tries and parameters (Fig. 2b, Supplementary Fig. 2ab, Supplementary Fig. 5, Supplementary

Table 2) were initialized in a rectangular box of size $Lx=Ly=425$ L. The left portion of the box

was filled with cells in hexagonal packing, while the right side contained the pillar geometry.

Simulations used for close temporal comparison with experiments (Fig. 2c-e, Fig. 3-4, Supple-

mentary Fig. 3-4) were initiated in a smaller box (Lx=425 L, Ly=106 L) with new cells added on

the left side during the simulation to maintain a stable bulk pressure. Each parameter combina-

tion was run with at least 10 repeats. The ordered pillar geometry was constructed as a square

lattice, with pillar spacing being twice the effective pillar diameter. Reported results for ordered

pattern simulations (e. g. Fig. 2b,c,e and Fig. 3d) are averaged over four orientations of the

square lattice (0°, 15°, 30° and 45°, Supplementary Fig. 3b). The disordered pattern was gener-

ated from the ordered patterns of all four tilts by shifting each pillar by a random angle and dis-

tance following a Laplace distribution from its original position. The fully disordered pattern cor-

responded to Laplace distribution parameters $\mu = b = 0.5 * L_p$.

Simulation analysis

The initial time required for cells to accumulate at the front of the pillar pattern and enter it was

excluded from the analysis. Unless stated otherwise, simulation frames were analyzed at a fre-

quency of $6\,\tau$ after the start of the invasion. Simulations were run until bulk cells reached the

end of the pillar pattern. All simulation analyses were performed in Python.

The overall number of detached cells (Fig. 2b,d, Fig. 3a, Supplementary Fig. 2a,b,d, Supple-

mentary Fig. 3d) was determined by distance-based clustering of cells, using r_{cut} as the natural

clustering cutoff from a physical perspective, and summing the number of cells within the pat-

tern that were not connected to the bulk at a given timepoint. The identification of pattern loca-

tions with high detachment probability and instantaneous detachments (Fig. 2c, Fig. 3c-d,f-g,

Fig. 4c,h, Supplementary Fig. 2e, Supplementary Fig. 3g, Supplementary Fig. 4c-e) was per-

formed in two steps. First, the same clustering analysis was applied to each simulation frame.

Second, a cell was considered detached if it remained separated from the bulk (largest cluster)

for at least 5 consecutive frames. For each detachment event, we recorded the x,y coordinates,

cell ID, and timestep for further analyses, such as relating detaching events to interface proper-

ties (Fig. 3c,d) or local pillar positions (Fig. 4d).

The shape of the cellular interface (eg. Fig. 2d,e, Fig. 3a, Fig. 4a) was analyzed by first compu-

ting the Delaunay triangulation (convex hull) of cellular positions in each simulation frame. The

Delaunay triangulation was then filtered using the alpha shape algorithm[84] (Python alphashape

library) to determine the concave hull of the cellular interface (alpha = 0.6). The positions of in-

terface cells were subsequently used in the roughness scaling analysis (Fig. 2e, Supplementary

Fig. 3c,e,g) and in the analysis of interface shape, perimeter and curvature (Fig. 3a,c,d,h, Fig.

4b, Supplementary Fig. 3a). The local maxima and minima of the cell interface were identified

using the argrelextrema function in SciPy library[85]. Transient protrusions on the order of cell size

(protrusion length = average distance to next local minimum < 1 L) were filtered out.

The interface roughness (Fig. 2e, Supplementary Fig. S3e,g) was quantified using the height

function $h(y)$,which represents the level of invasion of cells at different points along the interface

(polar coordinates were used for pillar pattern experiments). We analyzed the temporal increase

in roughness by computing the interface width $w(l,t)$, which measures the root-mean-square de-

viation of the interface from its mean value, defined as:

$$w(l, t) = \left\{ \frac{1}{N} \sum_{i-1}^{N} [(h(y, t) - \bar{h}(t)]^2 \right\}_{L_{int}}^{1/2}$$

850   where L_{int}=1000 µm is the length of the interface contour. These fluctuations grow over time fol-

851   lowing a power law scaling, $w(l, t) \sim t^{\beta}$, with a characteristic critical growth exponent β[31].

853   To quantify the spatial roughness of the invasion front (Supplementary Fig. S3b), we computed

854   the height–height correlation function of the interface profile in simulations. The function is de-

855   fined as:

$$C(r) = \langle [h(x + r) - h(x)]^2 \rangle$$

856   Where $h(x)$ denotes the height of the interface at lateral position x, and $\langle \cdot \rangle$ represents an aver-

857   age over all positions x and multiple simulation runs. For self-affine interfaces, this function ex-

858   hibits power-law scaling with distance r:

$$C(r) \sim r^{2\alpha}$$

We computed $C(r)$ from late-time snapshots of the invasion front, after roughness had reached a

statistically stationary regime. The roughness exponent α was extracted by fitting the log–log

plot of $C(r)$ versus r over the intermediate scaling range, avoiding short-range discretization arti-

facts and long-range saturation effects. We repeated this analysis for simulations conducted in

homogeneous and disordered environments. The resulting values of α provide an independent

estimate of the interface roughness exponent, complementary to the growth exponent β ob-

tained from time-dependent roughness evolution. To test for Family–Vicsek scaling (Supple-

mentary Fig. S3c), we performed simulations for multiple lateral system sizes L. For each L, we

ran and averaged ~20 independent simulations; in homogeneous environments these differed

only by stochastic cell dynamics, whereas in disordered environments they additionally differed

by independently generated random pillar patterns. We then rescaled roughness and time as

w/L^{α} and t/L^{z}, with independently measured exponents (α = 0.55, β = 0.33 in homogeneous; α =

0.70, $\beta = 0.55$ in disordered environments) and the dynamic exponent defined as $z = \alpha/\beta$. Col-

lapse of the rescaled curves across system sizes was taken as evidence that the observed dy-

namics fall into the corresponding universality class.

The analysis of local pillar heterogeneity (Fig. 4h, Supplementary Fig. 4e,f) was performed by

identifying all neighboring pillars within a radius r_{max}= 3.18 L (three times average pore size)

around a given (x,y) position in the disordered pattern. These pillars were then classified into

four quadrants (front, back, left right; Fig. 4g). The pillars in each quadrant were weighted by

their distance from the reference position using the factor $(r_{max} - r)/r_{max}$ and the weights were

then summed.

Supplementary Figures

882
Supplementary Fig. 1: Invasion dynamics of cells in different environments a. 2D invastion assay
of WT A431 cells with and without mCherry reporter with 1:10 mix respectively. **b.** Collagen invasion as-
say of WT and E-,P-Cad[-/-] spheroids imaged with brightfield microscopy. **c.** Detachments of the WT and
E-,P-Cad[-/-] cells invading into ordered and disordered pillar patterns. White arrows indicate individual de-
tached cells or cell clusters. **d.** Detached cell area in the respective pillar patterns for A431 WT and E-,P-
CadKO cells. Scale bars: 100 µm, box and whiskers plot displays median, 1[st] and 3[rd] quartiles (hinges)
and bars display 1.5 interquartile range from the hinges, n=3 experiments per genotype and pattern fol-
lowed over 8 days. **e-f.** Cell invasion assays into smaller (6 µm) sized ordered pillar pattern (d) and step-
wise or sinusoidal pattern (e). **g.** Quantified number of detachments for each pillar patterns, where box
and whiskers plot displays median, 1[st] and 3[rd] quartiles (hinges) and bars display 1.5 interquartile range

Supplementary Figure 2: Phase diagram of invasion dynamics for different model parameters. a.
Temporal increase in detachment rate is a robust phenomenon present both in ordered and disordered
pillar patterns, as well as for different simulation parameters. The simulation snapshots show a repre-
sentative state of the systems at day 6 (cells are shown as grey and pillar as black beads, detached cells
are highlighted in red), with the best-fit set of simulation parameters used in the main text (left column,
see also Supplementary Table 1). Each subsequent column shows the effect of changing a single simula-
tion parameter ($D_r = 1\ \tau^{-1}$, $D_r = 0.5\ \tau^{-1}$, $k_{bias} = 0.5\ rad/\tau$, $k_{bias} = 1\ rad/\tau$, $v_0 = 0.24\ L/\tau$ respectively). **b.**
Snapshots of simulation od cells in ordered and disordered environments at day 6 (cells are shown as
grey and pillar as black beads, detached cells are highlighted in red), for different values of cell-pillar ad-
hesion (r_{cut}=1.55 L). **c.** Snapshots of simulation of cells in homogeneous environments (grey beads
shown with smaller diameter for clarity, day 6), for different values of the fluidity number. High fluidity
number (left) leads to numerous detachments, something not observed in wild-type experiments, while
low fluidity (right) leads to a solid-like rheology without any re-arrangements on the time scale of the simu-
lations (also not observed). **d.** Snapshots from simulations of cells with lower cell-cell adhesion, corre-
sponding to E-,P-Cad$^{-/-}$ cells (fluidity 0.87, cells are shown as grey and pillar as black beads, detached
cells are highlighted in red). This leads to detachments even in the absence of any pillars, i.e. to a much
weaker sensitivity to the environment, as seen experimentally. **e.** Quantification for the increase of de-
tachments when lowering the cell-cell adhesion, for either ordered or disordered pillar patterns, showing
that the difference between ordered and disordered pattern disappears (T-test: p=0.47) compared to in-
termediate adhesion/fluidity.
918
919

Supplementary Figure 3: Universality class identification and sensitivity analysis of simulation outputs for different pillar patterns. a. Temporal evolution of the interface perimeter, quantified in simulations (n=40 simulations, the shaded intervals indicate standard error). **b.** Height-height correlation function of the invasion front in simulations in steady-state snapshots of the interface (day 25) to estimate the roughness exponent . Shaded intervals represent standard error across the analyzed snapshots. Fits were performed over the intermediate scaling range (1 < r < 12). **c** Family–Vicsek scaling collapse of interface roughness for homogeneous (left) and disordered (right) environments. When the original curves for different *L* (insets) are rescaled according to the Family–Vicsek relation / vs. / , with independently measured exponents, they collapse onto a single universal function, confirming assignment to the KPZ and KPZq universality classes, respectively. **d.** Differences in invasion behavior with different orientations of regular pattern (snapshots of day 4, cells are shown as grey and pillar as black beads,

magenta line represents cell interface). We used the average results of the 4 shown orientations for com-

paring regular pattern with other geometries, given that invasion is radial in experiments. **e.** Interfacial

roughness and roughness exponents in 4 orientations of regular patterns (corresponding to b) (n=10 sim-

ulations each, the shaded intervals correspond to standard error). **f.** Simulation snapshots of invasion in

sinusoid and stepwise pillar patterns, exhibiting lower or comparable detachments, as in ordered pattern,

as seen experimentally in Supplementary Fig 1e-g. Cells are represented as grey and pillars as black

beads, detached cells are marked red, cell interface in magenta, day 4. **g.** Quantifications of interface

width and detachment numbers in simulations of panel d, showing that it follows a stepwise trend corre-

sponding to the pillar pattern, and that some cells tend to reattach to the bulk resulting in lower overall

number of detachments.

Supplementary Figure 4: Detachments statistics in data and simulations. a. Analysis of interface roughness in experiments. The blue line represents the segmented outline of cell bodies and magenta line connects the cell nuclei along the interface, which are used for subsequent roughness and interface curvature analysis (see Methods). **b.** Difference between the finger angle (defined as the angle between the tip of finger and neighboring local minima, with small angles thus corresponding to sharp fingers, see Fig. 3e) of the protrusions in ordered and disordered patterns, which becomes more pronounced in later timepoints (n=16 experiments for regular and n=18 for disordered pattern). **c.** Distribution of detachment events in simulations with increasing disorder in pillar positions (color coded in red for increasing probability). **d.** Temporal increase of detachments in perpendicular sinusoid simulations with smaller wavelength (see Fig. 4a). **e.** Heat map of detachment positions compared to the local anisotropy of pillar positions in disordered pattern (Metric: Left + Right – Front, $R^2 = 0.1$) and local anisotropy of pillar positions multiplied by the mask, which takes into account the increase of detachments in the direction of invasion ($R^2=0.63$). detachments analyzed from n=10 simulations. **f.** Example of a different metric calculated using the same analysis (Back-Front), which exhibited lower spatial correlation with detachment positions.

Supplementary Figure 5: Fitting procedure for different model parameters. a) Correction of tilts in
the detected experimental interface (blue line) by removing long Fourier modes (orange line). **b-c)** Fitting
the active force v_0 based on displacements of single detached cells in a pillar patterns, respectively (ex-
perimental data shown in the first two columns of panel b, orange line in panel c represents the median of
distribution of detached cells, n=91 trajectories from n=6 experiments). The displacements do not depend
on D_r and grow linearly with v_0 (further columns in panel b show n=3 simulations of n=100 sparsely added
single cells in ordered pillar patterns with pore size 1.06 L and different combinations of parameters). By
comparing the data to different values of v_0, we constrained v_0=0.43 $L/$ as the best-fit parameter used in
all simulations unless explicitly mentioned **d)** Fitting of the rotational diffusion constant based on MSAD

of single detached cells in a regular pillar pattern (color coded according to value of v_0, black dashed line
represents MSAD of detached cells in experiment). As above, we compared the experimental results from
trajectories of detached cells to simulations with different rotational diffusion coefficients D_r. We found that
the best-fit occurred for 0.7 τ^{-1} (linear fit to simulation data, R^2=0.93), which we use for all simulations
unless explicitly mentioned. **e)** Fitting of the k_{bias} based on the radial bias in trajectories of single detached
cells in a regular pillar pattern (same procedure as above). We found that k_{bias}=0.75 *rad/τ* reproduced
best the experimental data (Kolmogorov-Smirnoff test, p=0.46), which we use for all simulations unless
explicitly mentioned. **f)** MSD comparing the detached cells in pillar patterns in experiment and different
combinations of simulation parameters D_r, v_0, and k_{bias}. The final set of parameters chosen according to
b-e exhibited highest pseudo-R^2 related to the experimental MSD curve.

Supplementary Table 2. Interface roughness of cells invading in ordered and disordered patterns in different parameter combinations (green – exponents from 0.25-0.44, blue – larger than 0.55).

Pore size	D_r [τ^{-1}]	Ordered pattern fluidity				Disordered pattern fluidity			
		0.44	0.52	0.60	0.76	0.44	0.52	0.60	0.76
0.88 *L*	0.5	0.27	0.33	0.41	0.50	0.55	0.70	0.70	0.57
	1	0.19	0.36	0.43	0.44	0.45	0.65	0.72	0.71
	2	0.37	0.25	0.23	0.37	0.44	0.36	0.39	0.49
1.06 *L*	0.5	0.48	0.57	0.62	0.59	0.54	0.62	0.78	0.57
	1	0.30	0.45	0.52	0.55	0.44	0.69	0.64	0.59
	2	0.34	0.23	0.28	0.42	0.35	0.32	0.40	0.43
1.24 *L*	0.5	0.32	0.32	0.38	0.38	0.57	0.66	0.80	0.49
	1	0.37	0.46	0.49	0.50	0.58	0.69	0.64	0.59
	2	0.33	0.27	0.31	0.41	0.56	0.33	0.40	0.43

Supplementary Table 3: Behavior of the cell bulk in homogeneous environment in simulations with different parameters (green – KPZ scaling, yellow – KPZ scaling with holes and detachments, red – jammed, collective movement; brown – jammed, no movement

Supplementary Movie Legends

Supplementary Movie 1: Invasion of A431 human skin carcinoma cells in ordered vs. disordered environments. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green).

Supplementary Movie 2: Invasion of A431 human skin carcinoma cells in free environments. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green).

Supplementary Movie 3: Invasion of EPCad[-/-] A431 human skin carcinoma cells in ordered vs. disordered environments. Cells express cytoplasmic mCherry (blue) and nuclear H2B-GFP (green).

Supplementary Movie 4: Simulations of cells with different fluidity number (i.e. with varying cell-cell in-
teraction strength) in homogeneous environment showing different level of cell rearrangements. Cells are
represented as grey beads.

Supplementary Movie 5: Simulations of cells migrating in different environments (homogeneous, or-
dered and disordered. Cells are represented as grey and pillars as black beads.

Supplementary Movie 6: Simulations of EPCad$^{-/-}$ cells (lower cell-cell adhesion) invading different types
of environments. Simulations of cells migrating in different environments (homogeneous, ordered and
disordered. Cells are represented as grey and pillars as black beads.

Supplementary Movie 7: Simulations of cell invasion in spatially structured environments which do not
increase detachment rate (regular with smaller pore, sinusoid, stepwise). Cells are represented as grey
and pillars as black beads.

Supplementary Movie 8: Simulation and experiment of cell invasion in sinusoid perpendicular pattern,
which creates localized interface protrusions and detachments. Cells in experiment express cytoplasmic
mCherry (blue) and nuclear H2B-GFP (green). Cells in simulations are represented as grey and pillars as
black beads.

1010
1011